A-level

ENVIRONMENTAL SCIENCE

Questions by topic: Agroecosystems: abiotic & biotic factors - Set 1

Mark scheme

AQA spec 3.5.1.1; Biological resources; textbook pp. 306-320

Marking guidance

• Each part opens with a line in bold saying how its marks are shared out, followed by one sentence on how to mark it.
• The figure in bold at the start of a marking point is the mark that point earns. Lines indented beneath it, starting with a dash, are the detail that belongs to the same mark; a line starting **OR** is another route to it.
• **Also accept** lists other wordings and answers that earn the mark - wording never has to match these notes, only the science. **Do not credit** lists the answers that look right and earn nothing.
• Where a question asks for a fixed number of points (eg *two* reasons), credit only the first points given.
• For levels-of-response questions, determine the level from the descriptors first, then position the mark within the level using the indicative content.
• Calculations: credit correct working (method marks) even if the final answer is wrong; incorrect answers with no working score 0.

Textbook references: every question below cites the printed page numbers in R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018) from which the content is drawn, together with the AQA 7447 specification reference.

Question	Marking guidance	Marks
01.1	**1 mark each** 1 mark for each of the 3 points below. **1** B: frost-free period (150 days) exceeds the 120 needed whereas A (110) is too short - late frosts would kill seedlings **1** B is warmer / longer thermal growing season because south-facing slopes receive more insolation – valley floors are frost pockets where cold dense air collects **1** But B has lower rainfall and a steeper slope (erosion / machinery difficulties) so irrigation / contour cultivation may be needed **Also accept** • Any other correct, relevant point of the same kind - wording never has to match these notes. • **insolation** → incoming solar radiation / energy arriving from the Sun	3

Do not write outside the box

Question	Marking guidance	Marks
01.2	**Any 3 of 4** 1 mark each for any 3 of the 4 points below, in any order, maximum 3. **1** Greenhouses / polytunnels heated by solar gain or burning gas / paraffin (ventilated to prevent overheating) **1** Woven plastic cloches / fleece over rows to retain warm air near the ground **1** Burning oil / gas or using large fans in orchards to prevent frost damage to blossom **1** Site selection (aspect, avoiding frost pockets) – heated buildings and shelter for livestock **Also accept** • Any other correct, relevant point of the same kind - wording never has to match these notes. **Do not credit** • Points after the first three - the question asks for three, so a list of extras cannot pick up the marks.	3
01.3	**1 mark each** 1 mark for each of the 2 points below. **1** Mammals and birds use food energy to maintain a constant body temperature (thermoregulation) **1** Reducing heat loss (shelter, heated buildings) means less food energy is used for heat and more is available for growth **Also accept** • Any other correct, relevant point of the same kind - wording never has to match these notes.	2
01.4	**1 mark each** 1 mark for each of the 2 points below. **1** Long days increase milk production and egg laying – poultry grow best with short days – sheep mate as days shorten in autumn **1** Artificial lighting extends the growing season for greenhouse crops or simulates autumn in spring to give sheep a second mating / lambing season **Also accept** • Any other correct, relevant point of the same kind - wording never has to match these notes.	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 306-309; AQA spec 3.5.1.1.	
02.1	**1 mark each** 1 mark for each of the 3 points below. **1** Plants close their stomata to reduce transpiration losses **1** Closed stomata stop the absorption of carbon dioxide so photosynthesis and growth stop **1** Less water for nutrient absorption / transport and cell turgidity – severe shortage dehydrates cells and inhibits biochemical reactions **Also accept** • Any other correct, relevant point of the same kind - wording never has to match these notes. • **transpiration** → water lost as vapour from the leaves	3

Do not write outside the box

Question	Marking guidance	Marks
02.2	**Marks shared** Up to 4 marks in total - share them between the 3 points below according to how completely each one is made. **up to 2** Anaerobic soil favours denitrifying bacteria and inhibits nitrifying bacteria so nitrate is lost faster than it forms – higher risk of fungal disease – root respiration inhibited **up to 2** Drainage ditches or pipes **up to 2** Deep ploughing – avoiding compaction by machinery / livestock – adding organic matter to encourage worms **Also accept** • Any other correct, relevant point of the same kind - wording never has to match these notes. • **waterlogged** → soil pores filled with water so there is no air • **denitrifying** → nitrate converted back to nitrogen gas • **nitrifying** → ammonium oxidised to nitrite and then nitrate	4
02.3	**1 mark each** 1 mark for each of the 2 points below. **1** Groundwater often contains dissolved salts – when the water evaporates from the soil the salts are left behind and accumulate **1** High soil salinity causes osmotic dehydration of roots / crop death – washing out with extra water uses more water and salinises rivers **Also accept** • Any other correct, relevant point of the same kind - wording never has to match these notes. • **salinisation** → salt building up in the soil / soil becoming salty	2
02.4	1 mark for the point below. **1** Add crushed lime (calcium carbonate / hydroxide) – low pH increases leaching, inhibits nutrient uptake and mobilises toxic aluminium and lead **Also accept** • Any other correct, relevant point of the same kind - wording never has to match these notes. • **leaching** → nutrients or ions washed down out of the soil / washed away by water	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 308-310, 313; AQA spec 3.5.1.1.	

Do not write outside the box

Question	Marking guidance	Marks
03.1	**Marks shared** Up to 6 marks in total - share them between the 4 points below according to how completely each one is made. **up to 2 Organic advantages:** often waste products, locally available and cheap – add humus improving structure and water retention – increase soil biota – slow release reduces leaching **up to 2 Organic disadvantages:** nutrient composition cannot be controlled – released slowly so needs long-term planning – bulky with high water content so expensive to transport – cannot be applied to a growing crop – odour / pathogens **up to 2 Inorganic advantages:** composition controlled to match crop needs – nutrients released rapidly – concentrated and easy to apply **up to 2 Inorganic disadvantages:** high energy (fossil fuel) manufacture (Haber process) – no organic matter so humus and biota decline – some toxic to worms – high solubility so leached causing eutrophication – raw materials (phosphate rock) finite **Also accept** • Any other correct, relevant point of the same kind - wording never has to match these notes. • **humus** → dark, decomposed organic matter in soil • **leaching** → nutrients or ions washed down out of the soil / washed away by water • **eutrophication** → nutrient enrichment of water leading to algal growth **Do not credit** • 'the algae use up all the oxygen' on its own - the oxygen is used by aerobic decomposers breaking down the dead algae	6
03.2	**Marks shared** Up to 4 marks in total - share them between the 3 points below according to how completely each one is made. **up to 2** Crops grown in a circulating nutrient solution (inert medium) in greenhouses – nutrient concentrations monitored and adjusted so nutrients are never limiting – abiotic factors controlled **up to 2** Roots are small (all in contact with nutrients) so more growth goes into the harvested crop – no soil-borne pathogens or weeds – crop stays fresh with roots **up to 2 Disadvantages:** high inputs of nutrients and energy (heating, lighting, pumping) – high level of technical knowledge – only economic for high-value crops **Also accept** • Any other correct, relevant point of the same kind - wording never has to match these notes. • **abiotic** → non-living / physical factors	4
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 310-312; AQA spec 3.5.1.1.	

Do not write outside the box

Question	Marking guidance	Marks
04.1	**Marks shared** Up to 4 marks in total - share them between the 3 points below according to how completely each one is made. **up to 2 Rotation:** each crop has its own pests – growing a different crop each year (4-5 year cycle) means pests left at the end of a season die before that crop returns, so populations cannot build up early **up to 2** Beetle banks / hedgerows provide habitat and overwintering sites for natural predators (ground beetles, ladybirds) that eat aphids **up to 2** Rotation also evens out nutrient demands and includes legumes **Also accept** • Any other correct, relevant point of the same kind - wording never has to match these notes.	4
04.2	**1 mark each** 1 mark for each of the 3 points below. **1** Integrated control gives almost the same yield as chemical control (18.1 vs 18.5) with far fewer applications (1-2 vs 6) **1** It retains most of the biodiversity of cultural control (26 vs 31 species) whereas chemical control alone leaves only 8 **1** Cultural control alone loses about 12% yield – integrated control combines cultural methods, resistant varieties and carefully timed specific non-persistent pesticides only when essential - best balance of yield, cost and impact **Also accept** • Any other correct, relevant point of the same kind - wording never has to match these notes. • **biodiversity** → the variety of life: habitat, species and genetic diversity • **persistent** → does not break down / stays in the environment for a long time **Do not credit** • pests 'getting used to' a pesticide - the few naturally resistant individuals survive and pass the trait on	3
04.3	**1 mark each** 1 mark for each of the 3 points below. **1** Large numbers of males are sterilised with gamma radiation and released **1** Females of many insects mate only once and store sperm, so mating with a sterile male produces no offspring – enough sterile males reduce fertile matings below replacement level (screw worm fly, Mediterranean fruit fly) **1** Only the target species' females mate with them so non-target species are unaffected – fails if sterilised males behave abnormally or re-colonisation occurs **Also accept** • Any other correct, relevant point of the same kind - wording never has to match these notes.	3
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 314-318; AQA spec 3.5.1.1.	

Do not write outside the box

Question	Marking guidance	Marks
05.1	**Marks shared** Up to 4 marks in total - share them between the 3 points below according to how completely each one is made. **up to 2 Success:** Cactoblastis moth introduced from South America controlled prickly pear cactus in Australia – Encarsia parasitic wasps control greenhouse whitefly **up to 2 Problems:** cane toads introduced to Australia to control sugar cane beetles did not control them but ate many other species and are toxic to predators – rosy wolfsnail introduced to Polynesia to control African land snails ate endemic Partulid snails (48 species extinct) – harlequin ladybirds introduced to Europe eat native ladybirds **up to 2 Lesson:** control agents should be specialist feeders on the pest **Also accept** • Any other correct, relevant point of the same kind - wording never has to match these notes. • **biological control** → using a natural predator, parasite or pathogen of the pest • **endemic** → found naturally only in that one place	4
05.2	**1 mark each** 1 mark for each of the 2 points below. **1** Release an artificial sex scent attracting males (or females) of the pest species **1** Used to monitor when the pest arrives so pesticide is timed correctly, or (in enclosed greenhouses) to trap enough of one sex to prevent breeding **Also accept** • Any other correct, relevant point of the same kind - wording never has to match these notes. **Do not credit** • pests 'getting used to' a pesticide - the few naturally resistant individuals survive and pass the trait on	2
05.3	**1 mark each** 1 mark for each of the 3 points below. **1** Regular low doses kill only the most sensitive bacteria so resistant strains survive and multiply **1** Agricultural use exceeds medical use, so resistance spreads **1** Resistant zoonotic bacteria (E. coli, Salmonella, Campylobacter) can transfer to humans and are harder to treat – hence reduced use of antibiotics is a sustainability goal **Also accept** • Any other correct, relevant point of the same kind - wording never has to match these notes. • **sustainability** → meeting present needs without stopping future generations meeting theirs **Do not credit** • 'sustainable means it lasts a long time' - it means use that can continue without reducing what is available in future	3

Do not write outside the box

Question	Marking guidance	Marks
05.4	1 mark for the point below. **1** Grow nectar-rich flowering plants alongside crops – restrict pesticides harmful to bees / spray when not in flower – introduce bee hives **Also accept** • Any other correct, relevant point of the same kind - wording never has to match these notes. **Do not credit** • pests 'getting used to' a pesticide - the few naturally resistant individuals survive and pass the trait on	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 316-320; AQA spec 3.5.1.1.	

Do not write outside the box