

A-level

ENVIRONMENTAL SCIENCE

Paper 1 - Generated practice paper Set 16

Mark scheme

120 marks: Questions 1-10 (95 marks) plus one essay from Question 11 (25 marks).

Marking guidance

- The number in brackets after each marking point is the mark it earns; the note above the points says how many are needed.
- Where a question asks for a fixed number of points (eg *two reasons*), credit only the first points given.
- Alternative correct wording and other relevant, accurate points are credited.
- For levels-of-response questions, first determine the level using the descriptors, then position the mark within the level using the indicative content.
- Calculations: credit correct working (method marks) even if the final answer is wrong; incorrect answers with no working score 0.

Textbook references: every question below cites the printed page numbers in R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018) from which the content is drawn, together with the AQA 7447 specification reference.

Question	Marking guidance	Marks
01.1	<i>1 mark for each point.</i> • Deaths rise as smoke concentration rises, peaking at about 930 per day on 8 Dec when smoke was highest (4.5 mg m^{-3}) (1) • Deaths fall as smoke falls but remain above the starting level for several days (time lag / delayed effect) (1)	2
01.2	<i>A correct final answer (with unit where asked) scores full marks. If the answer is wrong, award 1 mark for each correct step of working shown.</i> • $(930 - 250) / 250 \times 100$ (1) • = 272% (1)	2
01.3	<i>1 mark for each point.</i> • Anticyclonic conditions: clear skies allowed rapid radiation of heat from the ground at night so the ground and air near it cooled (1) • Low wind velocity so air layers did not mix; London lies in a valley / basin where cold dense air collected: a temperature inversion formed (1) • Smoke from coal fires was trapped below the inversion; moist air cooled to its dew point forming fog; smoke + fog = smog; the smog's high albedo prolonged the inversion (1)	3
01.4	<i>1 mark for each point.</i> • Smoke particles (PM10) are inhaled deeply and irritate the lungs / kill cilia so particles and bacteria are not cleared, causing bronchitis / infections (1) • These groups have weaker lungs / immune systems / less reserve so the extra stress (with sulfur dioxide acting synergistically) was fatal (1)	2
01.5	<i>1 mark.</i> • Clean Air Act (1956) (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 245-247; AQA spec 3.4.3.2.1.	

Question	Marking guidance	Marks
02.1	<i>A correct final answer (with unit where asked) scores full marks. If the answer is wrong, award 1 mark for each correct step of working shown.</i> • Single: $5.8 \times 20 \times 12 = 1392 \text{ W}$ (1) • Argon: $1.6 \times 20 \times 12 = 384 \text{ W}$ (1) • Reduction = 1008 W (accept 1010) (1)	3
02.2	<i>1 mark for each point.</i> • A wider gap of trapped gas reduces conduction because gas is a poor conductor (1) • If the gap is too wide convection currents start in the gas, carrying heat across; the optimum depends on the gas and window size (1)	2
02.3	<i>1 mark for each point.</i> • Low-e coating reflects long-wavelength infrared back into the room, reducing radiation losses (1) • Argon / krypton / xenon have lower thermal conductivity than air so less conduction across the gap (1)	2
02.4	<i>1 mark for each point.</i> • Panes must be held apart by small spacers which cannot be long, so the gap is only about 0.2 mm (1) • Used where there is not room for thick gas-filled multiple glazing (eg historic buildings) (1)	2
02.5	<i>1 mark.</i> • Each extra layer saves less energy (diminishing returns) while cost rises, so beyond the optimum the cost of insulation exceeds the value of energy saved (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 228-231; AQA spec 3.3.4.2.2.	
03.1	<i>A correct final answer (with unit where asked) scores full marks. If the answer is wrong, award 1 mark for each correct step of working shown.</i> • Human = $120 + 60 + 30 = 210$; total = 325 (1) • $210 / 325 \times 100 = 64.6\%$ (accept 65%) (1)	2
03.2	<i>1 mark for each point.</i> • Nitrogen and hydrogen react to form ammonia over an iron catalyst at high temperature and pressure; ammonia then converted to nitrates for fertiliser (1) • Uses large amounts of energy from fossil fuels (also the hydrogen source), releasing CO_2 and NO_x (1) • Biological fixation by bacteria occurs at low temperature using solar energy captured by plants (enzymes lower activation energy) (1)	3
03.3	<i>Any 3 from the points below - 1 mark each, maximum 3.</i> • Eutrophication of water bodies by leached nitrates / sewage (1) • Photochemical smogs and tropospheric ozone from NO_x (1) • Acid rain (nitric acid) (1) • Nitrous oxide as a greenhouse gas (1) • NO_x toxicity / respiratory disease (1) • Nitrate in drinking water (methaemoglobinaemia) (1)	3
03.4	<i>1 mark for each point.</i> • Carbon is lost as CO_2 (composting) or CH_4 (digestion) so the C:N ratio improves / more nitrogen-rich (1) • Bulk and odour reduced, pathogens killed, easier to apply; digestion also yields biogas (1)	2

Question	Marking guidance	Marks
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 154-155; AQA spec 3.2.4.3.	
04.1	<i>A correct final answer (with unit where asked) scores full marks. If the answer is wrong, award 1 mark for each correct step of working shown.</i> • $(4.8 - 1.4) / 4.8 \times 100$ (1) • = 70.8% (accept 71%) (1)	2
04.2	<i>1 mark for each point.</i> • A tributary joins the river, more than doubling the flow (15 to $36 \text{ m}^3 \text{ s}^{-1}$) (1) • So the pollutant is diluted by the extra water (dispersal), not broken down (1)	2
04.3	<i>1 mark for each point.</i> • The pollutant is persistent so it is not degraded / broken down as it travels (1) • Little extra water joins the river (flow 36 to 38) so little further dilution (1)	2
04.4	<i>Any 2 from the points below - 1 mark each, maximum 2.</i> • Lake has a long residence time / water does not flow away so the pollutant accumulates rather than being carried away (1) • Smaller volume of water / less dilution / less dispersal by currents (1) • Persistent pollutant gradually becomes more concentrated (eg by evaporation) (1)	2
04.5	<i>1 mark for each point.</i> • Marine discharge: large volume of water and currents dilute and disperse the pollutant to acceptably low concentrations (control of emission location) (1) • River: small volume so concentrations stay high / drinking water abstraction / sensitive wildlife downstream (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 236-241, 254-255; AQA spec 3.4.1, 3.4.2.1.	
05.1	<i>1 mark for each point.</i> • When there is a difference in level between the sea and the basin: on the ebb (about 6-8 hours) water flows out through the turbines (1) • Also possible on the flood when the sea is higher than the basin; no generation when levels are equal (slack water) (1)	2
05.2	<i>1 mark for each point.</i> • Two periods of slack water in each 12.5-hour cycle (twice in about 25 hours) give no flow; the timing does not match demand (1) • Tides depend on the positions of the Moon and Sun which are known, so output can be forecast years ahead and alternative supplies planned (1)	2
05.3	<i>1 mark for each point.</i> • Spring tides: Sun and Moon aligned so their gravitational effects combine giving a large tidal range; neap: effects oppose giving a small range (1) • Energy depends on the mass of water and height difference, so output is much greater at springs (1)	2
05.4	<i>1 mark for each point.</i> • When demand is low, surplus electricity pumps water into the lagoon above sea level (1) • The stored water is released through the turbines later when demand is high (pumped storage) (1)	2

Question	Marking guidance	Marks
05.5	<i>1 mark for each point.</i> • Large tidal range (eg Severn Estuary) (1) • Estuaries, bays or headlands / channels that focus the flow and increase velocity (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 214-215; AQA spec 3.3.4.1.	
06.1	<i>1 mark for each point.</i> • Surplus electricity (eg windy nights) electrolyses water to produce hydrogen which is stored (1) • Hydrogen has a high energy density so it can be transported / stored for long periods (1) • Released by combustion for heat, industrial processes or steam turbines; or by fuel cells combining hydrogen and oxygen electrochemically to produce electricity and water (1) • Can also be used as a vehicle fuel or converted to methane / ammonia (1)	4
06.2	<i>Any 3 from the points below - 1 mark each, maximum 3.</i> • Compressed gas (up to 700 times normal volume) - energy to run the compressor, strong tanks (1) • Liquefied at very low temperature - energy for refrigeration and compression (1) • Adsorbed on a metal matrix (metal hydride) at low pressure - tanks larger and heavier than a petrol tank storing the same energy (1) • Conversion to ammonia - some energy needed but less than the alternatives (1)	3
06.3	<i>1 mark for each point.</i> • Both convert chemical energy to electricity electrochemically, but a battery stores a fixed amount of chemicals which are reformed by recharging (1) • A fuel cell keeps producing electricity as long as fresh fuel (hydrogen / alcohol) and oxidant are supplied (1)	2
06.4	<i>1 mark.</i> • Using hydrogen made from surplus renewable electricity as the main store and carrier of energy, released when demand exceeds supply or intermittent sources fail, for heating, vehicles and electricity (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 219-221; AQA spec 3.3.4.1.	
07.1	<i>1 mark for each point.</i> • Natural pressure of gas above the oil (or dissolved in it) or water beneath forces oil up the production well; pump-jacks ('nodding donkeys') increase flow (1) • Pressure falls as oil is removed so the remaining oil no longer flows to the well (1)	2
07.2	<i>1 mark for each point.</i> • Water or natural gas is pumped down an injection well to maintain the pressure and flow of oil (recovery to about 40%) (1) • Captured CO₂ can be pumped down instead, increasing oil recovery while storing the CO₂ underground (1)	2

Question	Marking guidance	Marks
07.3	<i>Any 3 from the points below - 1 mark each, maximum 3.</i> • Steam injection (from combustion or solar parabolic concentrators) to heat the oil and reduce its viscosity (1) • Controlled underground combustion to heat the oil (1) • Detergents / solvents to reduce surface tension so oil flows more easily (1) • Bacteria that partially digest heavy oil into lighter oils and produce CO₂ to maintain pressure (1)	3
07.4	<i>1 mark for each point.</i> • Many wells can be drilled from a single platform, reducing land take / number of rigs (1) • Deposits under locations where rigs cannot be placed (urban areas, sensitive habitats) can be reached (1) • Drilling can follow softer strata (quicker) and target multiple small reservoirs up to 10 km away, increasing total recovery (1)	3
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 187-188; AQA spec 3.3.4.1.	
08.1	<i>1 mark for each point.</i> • Dissolved oxygen fluctuates daily / 24-hour cycle: highest in the afternoon (about 13-13.5 mg dm⁻³ at 16 and 40 h) and lowest at night / early morning (2.5-3.5 mg dm⁻³ at 4 and 28 h) (1) • The night-time minimum is lower on the second night (2.5 vs 3.5) (1)	2
08.2	<i>1 mark for each point.</i> • Nitrates (from fertiliser runoff) and phosphates (from sewage effluent) cause eutrophication: rapid growth of algae / cyanobacteria (algal bloom) (1) • During the day photosynthesis by the dense algae releases oxygen so concentrations rise (above the normal saturation level) (1) • At night there is no photosynthesis but respiration by algae / animals / bacteria continues, using oxygen (1) • Aerobic bacterial decomposition of dead algae / shaded macrophytes uses oxygen; warm summer water holds less oxygen and decay is rapid (1)	4
08.3	<i>1 mark for each point.</i> • At night no photosynthesis replaces the oxygen used in respiration / decomposition so concentrations fall below the minimum fish can tolerate (1) • At the end of summer large amounts of dead vegetation decay rapidly in warm water causing severe deoxygenation (1)	2
08.4	<i>Any 2 from the points below - 1 mark each, maximum 2.</i> • Cyanobacteria release toxins harmful to livestock, pets and people (1) • Floating algae shade submerged / rooted macrophytes which die, disrupting food webs (1) • Algae are an unreliable food source (die off rapidly) so food webs are poor (1) • Blocked filters / unpleasant taste and smell in water supplies (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 266-268; AQA spec 3.4.3.2.9.	
09.1	<i>Correct answer only.</i> • The readings are precise but may not be accurate (1)	1

Question	Marking guidance	Marks
09.2	1 mark for each point. • A result that differs from other results with which it was expected to be similar (1) • Repeat the study / measurement: if it recurs it is a 'real' result caused by another variable; if not it was an error and can be excluded (1)	2
09.3	1 mark for each point. • To test the reliability of the method / equipment (1) • To decide the sample size, number of samples, spacing or timing needed to detect the variation without collecting unnecessary data (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 390-391; AQA spec 3.7.1.	
10.1	Levels of response Level 3 (7-9 marks): Answer is comprehensive, relevant and logically structured. Shows detailed knowledge and understanding of the relevant scientific ideas and applies them accurately to the context. Evaluation/analysis is well supported by evidence and a clear judgement or conclusion is reached. Level 2 (4-6 marks): Answer is mostly relevant and shows some structure. Knowledge and understanding of relevant ideas is sound but application to the context or evaluation may be incomplete or lack detail. A judgement may be made but is not fully supported. Level 1 (1-3 marks): Answer is basic, with limited structure. Some relevant knowledge is shown but there are omissions or errors. Little application to the context and little or no evaluation. 0 marks: No relevant content. Indicative content (credit other relevant, accurate points) • Stages of scientific method illustrated by ozone: identification (Antarctic UV data), hypothesis (Rowland-Molina 1974), data collection / analysis (Dobson units, satellites, balloons confirming CIO), proposed solutions, enactment (Vienna 1985, Montreal 1987) • Evidence of success: ozone hole area and depth stabilised and beginning to shrink; CFC concentrations in the atmosphere falling; UV levels not rising further; global ozone loss limited to about 4% • Reasons for success: international recognition of serious consequences (skin cancer); agreement of nearly every country; legally binding targets with a fund for developing countries; relatively few producers / products; alternatives (HCFCs, HFCs, pump sprays, hydrocarbons) already available and cheap; no lifestyle change needed • Limits: persistence of CFCs means recovery takes decades; some ODSs still used; HFC replacements are greenhouse gases • Climate change contrast: greenhouse gases come from almost every human activity (energy, transport, agriculture, industry) in every country; fossil fuels underpin economies so control requires major lifestyle and economic change • Agreements (Kyoto, Paris) have weaker / non-binding targets, disputes between developed and developing nations; emissions still rising; natural fluctuations, time delays and tipping points make cause-effect less clear to the public • Judgement: ozone control was a technical substitution problem; climate change is a systemic one, so the ozone success is only partly transferable	9

Question	Marking guidance	Marks
10.2	1 mark. Vienna Convention for the Protection of the Ozone Layer - no legally binding goals (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 108-113, 105-106; AQA spec 3.2.1.3, 3.2.1.2.	
11.1	Soil is often described as a non-renewable resource. Discuss the causes and consequences of soil degradation and evaluate the strategies that can be used to manage soils sustainably. Levels of response Level 5 (21-25 marks): Comprehensive and detailed knowledge and understanding drawn from across the specification, applied accurately to the question. Argument is coherent, well-structured and supported by a wide range of relevant, well-chosen examples and quantitative detail where appropriate. Interconnections between topics are made explicitly and a clear, justified conclusion is reached. Level 4 (16-20 marks): Detailed knowledge and understanding with good application to the question. Argument is logically organised with relevant examples; some links between different areas of the specification are made. Conclusions are drawn but may be less fully justified. Level 3 (11-15 marks): Sound knowledge and understanding, mostly applied to the question. Some structure and relevant examples, but coverage may be uneven or descriptive rather than analytical. Limited linking of topics. Level 2 (6-10 marks): Basic knowledge and understanding with partial relevance. Answer is largely descriptive with few examples and limited structure; some errors or omissions. Level 1 (1-5 marks): Limited, fragmented knowledge with little relevance to the question. Little structure; examples absent or inappropriate. 0 marks: No relevant content. Indicative content (students are not expected to cover all of these; credit other relevant material) - Features of fertile soil: water, nutrients, air, organic matter, pH, biota, texture, structure, depth; slow rate of soil formation vs rapid erosion (dynamic equilibrium broken) - Causes of degradation: vegetation removal, ploughing vulnerable soils, overgrazing, reduced organic matter and biota, cultivating steep slopes, compaction, salinisation from irrigation, nutrient depletion, pollution - Types of erosion: rain splash, runoff, wind, slumping; consequences: reduced productivity, river and reservoir sedimentation, flooding, coastal sedimentation, particulates, desertification, landslides - USLE as a tool for estimating loss and testing management - Management: long-term crops, low / zero tillage, contour ploughing, tied ridging, terracing, stone rows, windbreaks, multicropping / strip cropping, mulching, organic matter, livestock control, drainage / pH / nutrient management, conservation of soil biota - Links to sustainable agriculture, carbon storage, hydrological cycle, circular economy - Evaluation and judgement on which strategies are most effective in different environments Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 158-169, 328-337; AQA spec 3.2.5, 3.5.1.3, 3.5.1.5.	25

Question	Marking guidance	Marks
11.2	Discuss the difficulties in monitoring and predicting global climate change and evaluate the methods used to overcome them. Levels of response Level 5 (21-25 marks): Comprehensive and detailed knowledge and understanding drawn from across the specification, applied accurately to the question. Argument is coherent, well-structured and supported by a wide range of relevant, well-chosen examples and quantitative detail where appropriate. Interconnections between topics are made explicitly and a clear, justified conclusion is reached. Level 4 (16-20 marks): Detailed knowledge and understanding with good application to the question. Argument is logically organised with relevant examples; some links between different areas of the specification are made. Conclusions are drawn but may be less fully justified. Level 3 (11-15 marks): Sound knowledge and understanding, mostly applied to the question. Some structure and relevant examples, but coverage may be uneven or descriptive rather than analytical. Limited linking of topics. Level 2 (6-10 marks): Basic knowledge and understanding with partial relevance. Answer is largely descriptive with few examples and limited structure; some errors or omissions. Level 1 (1-5 marks): Limited, fragmented knowledge with little relevance to the question. Little structure; examples absent or inappropriate. 0 marks: No relevant content. Indicative content (students are not expected to cover all of these; credit other relevant material) • Nature of the problem: interconnected atmosphere-hydrosphere-biosphere systems; changes over different time and spatial scales; natural fluctuations (solar output, orbit) masking or exaggerating anthropogenic change; time delay between cause and effect (ocean heat capacity) • Limitations of historic data: sparse global coverage, unsophisticated equipment, heat island bias, no atmospheric composition records • Proxy data and their reliability: dendrochronology, coral growth rings, pollen in dated sediments, ice cores (gas bubbles, isotope ratios, volcanic ash layers, compaction problems) • Modern monitoring: long-term data sets (Mauna Loa), electronic equipment, satellites (LEO vs geostationary), Argo floats, remote sensing of ice thickness / area / sea level • Feedback mechanisms and tipping points adding uncertainty; ecological uncertainty (species survival, distribution, fragmentation) • Computer models: how they are tested and refined; importance of accurate representative data; scenario-based predictions and ranges • Uncertainty over ENSO / Conveyor / jet stream responses • Evaluation and conclusion: confidence in the overall trend and human cause despite detailed uncertainty; role of peer review and international bodies (IPCC) Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 89-90, 100-104, 10; AQA spec 3.2.1.2, 3.1.1.3.	25