

A-level

ENVIRONMENTAL SCIENCE

Paper 1 - Generated practice paper Set 35

Mark scheme

120 marks: Questions 1-10 (95 marks) plus one essay from Question 11 (25 marks).

Marking guidance

- The number in brackets after each marking point is the mark it earns; the note above the points says how many are needed.
- Where a question asks for a fixed number of points (eg *two reasons*), credit only the first points given.
- Alternative correct wording and other relevant, accurate points are credited.
- For levels-of-response questions, first determine the level using the descriptors, then position the mark within the level using the indicative content.
- Calculations: credit correct working (method marks) even if the final answer is wrong; incorrect answers with no working score 0.

Textbook references: every question below cites the printed page numbers in R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018) from which the content is drawn, together with the AQA 7447 specification reference.

Question	Marking guidance	Marks
01.1	<i>Correct answer only.</i> • Oxygen (1)	1
01.2	<i>1 mark for each point.</i> • A change to one process causes other processes to alter as a direct result (chains of linked processes / feedback) (1) • There is still a lot that is not understood about how atmospheric processes work (1)	2
01.3	<i>1 mark for each point.</i> • The surface warmed in the tropics heats the air above; warm winds (south-westerlies from the Caribbean) carry heat energy to higher latitudes (1) • Winds carry water vapour (evaporated from warm oceans) to areas that would otherwise receive little precipitation (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 86-88; AQA spec 3.2.1.1.	
02.1	<i>1 mark for each point.</i> • A fluid in a buried ground loop absorbs low-temperature heat from the ground; at the expansion valve the fluid boils at low pressure absorbing this heat (1) • A compressor raises the pressure so the gas condenses, releasing heat at a higher temperature inside the building (1) • The electricity only moves heat rather than creating it, so up to four times as much heat is delivered as electricity used (less if the source is colder) (1)	3

Question	Marking guidance	Marks
02.2	<i>1 mark for each point.</i> - Parabolic reflectors focus sunlight onto tubes of oil, increasing the energy density and reaching high temperatures (1) - The heat is transferred to molten salt (potassium nitrate) stored in large insulated tanks at up to 550 °C (1) - The stored heat boils water to drive steam turbines whenever electricity is required, including at night (1)	3
02.3	<i>1 mark for each point.</i> - Reflectors can only focus parallel rays of direct sunlight; cloud, dust and smoke scatter light so it cannot be concentrated (1) - PV cells convert diffuse as well as direct light, though output is lower (1)	2
02.4	<i>Any 2 from the points below - 1 mark each, maximum 2.</i> - Manufacture: extraction / processing of silicon, metals and plastics; toxic wastes (silicon tetrachloride, cadmium) (1) - Land take by large solar farms competing with other uses / affecting desert flora and fauna (1) - Water for cleaning panels in arid areas (1) - Disposal of old panels (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 200-202; AQA spec 3.3.4.1.	
03.1	<i>1 mark for each point.</i> - There is an environmental gradient (distance from the hedge / changing conditions) so samples must be taken at regular intervals along it to detect the change (1) - Random sampling might leave gaps / miss the pattern; systematic sampling avoids bias while covering the gradient (1)	2
03.2	<i>1 mark for each point.</i> - Sample at close intervals (eg every 1 m) first and see at what spacing the variations are still detected without unnecessary data; wider intervals (eg 4 m) may miss peaks and troughs (1) - Take increasing numbers of samples at one site and calculate the running mean; the number needed is where the mean stops fluctuating / stabilises (1) - Calculate standard deviation to assess scatter: more variable data need more samples (1) - Test the method (extraction technique, timing) for reliability before the main study (1)	4
03.3	<i>Up to 4 marks in total; award marks in proportion to the completeness of each point.</i> - Remove surface vegetation; pour an irritant solution (dilute mustard / detergent in water) onto the area until saturated (and around it), keep adding as it infiltrates (up to 2) - Collect worms that come to the surface for a standardised time; count / identify / weigh (up to 2) - Limitations: solution may not percolate impermeable soils or reach deep worms; worms may move sideways or downwards; some may die or not respond; slow-moving worms missed (up to 2)	4
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 390-393; AQA spec 3.7.1.	

Question	Marking guidance	Marks
04.1	<i>Any 4 from the points below - 1 mark each, maximum 4.</i> • High persistence means DDT remains in the environment for years / decades, so it has time to disperse and be taken up (1) • High liposolubility means it dissolves in fat and is stored in body tissues rather than excreted (bioaccumulation) (1) • Concentration increases at each trophic level as predators eat many contaminated prey (biomagnification) (1) • Birds of prey are at the top of the food chain so receive the highest concentrations (1) • Sub-lethal doses caused thin eggshells that broke / reduced breeding success; higher doses caused sterility or death (1)	4
04.2	<i>1 mark for each point.</i> • Advantage: low persistence / low liposolubility so they do not bioaccumulate or biomagnify / break down quickly (1) • Disadvantage: high mammalian toxicity so farm workers / humans are at risk of acute poisoning (neurotoxin inhibiting acetylcholinesterase); chronic effects (memory loss, depression); suspected carcinogens (1)	2
04.3	<i>1 mark.</i> • Pyrethroids are highly toxic to fish (even though mammal toxicity is low) (1)	1
04.4	<i>Any 3 from the points below - 1 mark each, maximum 3.</i> • Neonicotinoids are water soluble so they are absorbed by the roots (from the soil) and translocated through the plant (systemic) (1) • They are moderately persistent so remain in the soil long enough to be absorbed (1) • Seed coating avoids spraying so no spray drift / no dust onto flowers or surrounding habitats (1) • So bees / pollinators are less exposed than when the crop is sprayed (1)	3
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 263-265; AQA spec 3.4.3.2.8.	
05.1	<i>1 mark for each point.</i> • Acidophilic bacteria (eg Acidithiobacillus) oxidise sulfide ores, producing sulfuric acid which dissolves the metal ions (1) • Leachate (rainwater / recirculated solution) percolating through the spoil heap is collected; recirculation increases the metal concentration (1) • Copper is recovered from solution by electrolysis or by displacement with scrap iron (iron is more reactive so copper is deposited as solid metal) (1) • Smelting needs chemical reduction at high temperatures; energy needed rises rapidly as ore grade falls, so low-grade ores are uneconomic (1) • Bioleaching works at low temperatures with low energy inputs, so it can exploit low-grade deposits and wastes (increasing reserves) - mimics natural low-temperature processes (1)	5
05.2	<i>1 mark for each point.</i> • Plants (hyperaccumulators) absorb metal ions from soil / water and concentrate them in their leaves (1) • The vegetation is harvested and incinerated (1) • The concentrated metal in the ash is dissolved in acid and separated by electrolysis; also decontaminates polluted sites (1)	3

Question	Marking guidance	Marks
05.3	<i>1 mark for each point.</i> - Uranium ions dissolved in seawater adsorb onto the surface of synthetic or natural polymers (lignin from wood, chitin from shrimp shells) (1) - The polymers are collected and the uranium recovered, avoiding mining / high-energy extraction (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 143-145; AQA spec 3.2.3.7.	
06.1	<i>2 marks for each point (partial credit for a partly correct point).</i> - Asbestos: fibres become airborne when it disintegrates and are inhaled, causing asbestosis (scar tissue thickens alveoli) and mesothelioma; disposal: double-wrapped in heavy-duty polythene in a specialised, recorded landfill (secure, permanent, sealed storage) (2) - Cyanide: very toxic enzyme inhibitor (aerobic respiration) if inhaled / ingested; disposal: incineration so carbon and nitrogen are separated and oxidised to CO₂ and NO_x (much smaller problem) (2) - High-level radioactive waste: intense ionising radiation, heat generation, long half-lives; disposal: vitrification - mixed with molten glass in sealed stainless steel containers in ventilated concrete stores (2)	6
06.2	<i>1 mark.</i> The fibres are bound / cannot become airborne so cannot be inhaled (1)	1
06.3	<i>1 mark for each point.</i> - Waste is mixed with a cement slurry and poured into containers of impermeable, unreactive material (1) - The solid cement immobilises the waste so it cannot leak / leach out (1)	2
06.4	<i>1 mark.</i> Even if the glass shatters the waste stays encapsulated in the fragments; glass is more stable / resistant to heat and leaching over very long timescales (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 284-285, 303; AQA spec 3.4.3.2.12.	
07.1	<i>1 mark for the correct answer.</i> 7×10^{11} tonnes (1)	1
07.2	<i>A correct final answer (with unit where asked) scores full marks. If the answer is wrong, award 1 mark for each correct step of working shown.</i> 10% of 10 000 = 1000 billion tonnes (1) 1000 / 700 x 100 = 143% (accept 142.9) (1)	2
07.3	<i>1 mark for each point.</i> - Carbon is transferred into and out of rocks extremely slowly (fossilisation / sedimentation, weathering, volcanic release) (1) - Residence time is millions of years so the flux to the atmosphere is tiny compared with respiration / photosynthesis / combustion (1)	2
07.4	<i>1 mark for each point.</i> - CO₂ dissolves in the sea (forming carbonic acid / hydrogen carbonate ions) and is absorbed by phytoplankton through photosynthesis (1) - Organisms such as molluscs, corals and planktonic organisms build calcium carbonate shells / exoskeletons (1) - When they die the shells sink and sediment on the seabed, becoming limestone / chalk over long periods (1)	3

Question	Marking guidance	Marks
07.5	<i>1 mark for each point.</i> • Aerobic respiration uses oxygen to break down organic compounds completely to CO₂ (and water), releasing much more energy (1) • Without oxygen, anaerobic organisms only partly break down organic compounds, releasing carbon as methane (CH₄), which is later oxidised to CO₂ in the atmosphere (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 148-149; AQA spec 3.2.4.2.	
08.1	<i>1 mark for each point.</i> • Soil loss increases as vegetation cover decreases: forest lowest (0.5), bare soil highest (60 t ha⁻¹ yr⁻¹) - over 100 times more (1) • Ploughing downslope loses nearly three times as much as contour ploughing for the same crop (1)	2
08.2	<i>Any 4 from the points below - 1 mark each, maximum 4.</i> • Leaves / leaf litter intercept raindrops so soil particles are not dislodged by rain splash (1) • Roots bind soil particles together (1) • Vegetation acts as a windbreak reducing wind velocity and the kinetic energy available to carry particles (1) • Plants increase infiltration (root channels, organic matter) so less surface runoff (1) • Organic matter / humus from vegetation binds particles into stable peds (1)	4
08.3	<i>1 mark for each point.</i> • Furrows running downslope act as channels / gulleys that increase the velocity (kinetic energy) of runoff (1) • Furrows along the contour trap the water so it loses energy, deposits particles and infiltrates (1)	2
08.4	<i>1 mark for each point.</i> • In the dry season the soil (especially low-clay soils) is loose and dry with little cohesion so wind blows it away (1) • In the wet season heavy rain causes rain splash and surface runoff when infiltration capacity is exceeded (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 162-164; AQA spec 3.2.5.2.	
09.1	<i>Any 3 from the points below - 1 mark each, maximum 3.</i> • Disposal of items containing mercury: batteries, thermometers, fluorescent lights (1) • Chemical plants producing chlorine using mercury electrodes (1) • Combustion of coal (1) • Gold mining (amalgamation) (1)	3
09.2	<i>Any 3 from the points below - 1 mark each, maximum 3.</i> • Replacement of mercury thermometers with alcohol / electronic thermometers (discontinued uses) (1) • Removal from effluents by reverse osmosis / activated carbon filters / ion exchange filters (1) • Disposal / storage of wastes at high pH to reduce solubility (1) • Encapsulation of mercury-containing hazardous waste in cement (1)	3

Question	Marking guidance	Marks
09.3	1 mark for each point. • Most heavy metals are more soluble at low pH / less soluble at high pH (1) • So at high pH they remain insoluble / immobile and cannot be leached into water or absorbed by organisms (1)	2
09.4	1 mark for each point. • The effects of two pollutants interact to produce a different (greater) effect than either alone (1) • Cadmium and zinc; ozone and sulfur dioxide (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 275-278; AQA spec 3.4.3.2.11.	
10.1	Levels of response Level 3 (7-9 marks): Answer is comprehensive, relevant and logically structured. Shows detailed knowledge and understanding of the relevant scientific ideas and applies them accurately to the context. Evaluation/analysis is well supported by evidence and a clear judgement or conclusion is reached. Level 2 (4-6 marks): Answer is mostly relevant and shows some structure. Knowledge and understanding of relevant ideas is sound but application to the context or evaluation may be incomplete or lack detail. A judgement may be made but is not fully supported. Level 1 (1-3 marks): Answer is basic, with limited structure. Some relevant knowledge is shown but there are omissions or errors. Little application to the context and little or no evaluation. 0 marks: No relevant content. Indicative content (credit other relevant, accurate points) • Stages of scientific method illustrated by ozone: identification (Antarctic UV data), hypothesis (Rowland-Molina 1974), data collection / analysis (Dobson units, satellites, balloons confirming CIO), proposed solutions, enactment (Vienna 1985, Montreal 1987) • Evidence of success: ozone hole area and depth stabilised and beginning to shrink; CFC concentrations in the atmosphere falling; UV levels not rising further; global ozone loss limited to about 4% • Reasons for success: international recognition of serious consequences (skin cancer); agreement of nearly every country; legally binding targets with a fund for developing countries; relatively few producers / products; alternatives (HCFCs, HFCs, pump sprays, hydrocarbons) already available and cheap; no lifestyle change needed • Limits: persistence of CFCs means recovery takes decades; some ODSs still used; HFC replacements are greenhouse gases • Climate change contrast: greenhouse gases come from almost every human activity (energy, transport, agriculture, industry) in every country; fossil fuels underpin economies so control requires major lifestyle and economic change • Agreements (Kyoto, Paris) have weaker / non-binding targets, disputes between developed and developing nations; emissions still rising; natural fluctuations, time delays and tipping points make cause-effect less clear to the public • Judgement: ozone control was a technical substitution problem; climate change is a systemic one, so the ozone success is only partly transferable	9

Question	Marking guidance	Marks
10.2	1 mark. Vienna Convention for the Protection of the Ozone Layer - no legally binding goals (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 108-113, 105-106; AQA spec 3.2.1.3, 3.2.1.2.	
11.1	Explain how feedback mechanisms and tipping points influence the rate of global climate change and discuss the implications for the strategies needed to control it. Levels of response Level 5 (21-25 marks): Comprehensive and detailed knowledge and understanding drawn from across the specification, applied accurately to the question. Argument is coherent, well-structured and supported by a wide range of relevant, well-chosen examples and quantitative detail where appropriate. Interconnections between topics are made explicitly and a clear, justified conclusion is reached. Level 4 (16-20 marks): Detailed knowledge and understanding with good application to the question. Argument is logically organised with relevant examples; some links between different areas of the specification are made. Conclusions are drawn but may be less fully justified. Level 3 (11-15 marks): Sound knowledge and understanding, mostly applied to the question. Some structure and relevant examples, but coverage may be uneven or descriptive rather than analytical. Limited linking of topics. Level 2 (6-10 marks): Basic knowledge and understanding with partial relevance. Answer is largely descriptive with few examples and limited structure; some errors or omissions. Level 1 (1-5 marks): Limited, fragmented knowledge with little relevance to the question. Little structure; examples absent or inappropriate. 0 marks: No relevant content. Indicative content (students are not expected to cover all of these; credit other relevant material) - Negative feedback: increased low-level cloud (albedo), increased photosynthesis and sequestration; hydrological cycle evaporation-precipitation - Positive feedback: melting permafrost (methane), reduced albedo of snow / ice / sea ice, methane hydrate release, ocean acidification and coral loss, forest and peat fires, soil decomposition, water vapour, cirrus clouds - Tipping points: self-sustaining processes; why control must precede them; examples - Implications: urgency and precautionary principle; need for deep cuts in greenhouse gases (energy, transport, buildings, agriculture, forestry, CCS) rather than only adaptation; geoengineering risks - Difficulty of prediction because feedbacks interact at different scales; models; international agreements and per capita footprints - Judgement on whether current strategies are adequate Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 103-107, 371-372; AQA spec 3.2.1.2, 3.6.1.	25

Question	Marking guidance	Marks
11.2	Discuss how the atmosphere supports life on Earth and evaluate the extent to which human activities threaten these life-support functions. Levels of response Level 5 (21-25 marks): Comprehensive and detailed knowledge and understanding drawn from across the specification, applied accurately to the question. Argument is coherent, well-structured and supported by a wide range of relevant, well-chosen examples and quantitative detail where appropriate. Interconnections between topics are made explicitly and a clear, justified conclusion is reached. Level 4 (16-20 marks): Detailed knowledge and understanding with good application to the question. Argument is logically organised with relevant examples; some links between different areas of the specification are made. Conclusions are drawn but may be less fully justified. Level 3 (11-15 marks): Sound knowledge and understanding, mostly applied to the question. Some structure and relevant examples, but coverage may be uneven or descriptive rather than analytical. Limited linking of topics. Level 2 (6-10 marks): Basic knowledge and understanding with partial relevance. Answer is largely descriptive with few examples and limited structure; some errors or omissions. Level 1 (1-5 marks): Limited, fragmented knowledge with little relevance to the question. Little structure; examples absent or inappropriate. 0 marks: No relevant content. Indicative content (students are not expected to cover all of these; credit other relevant material) • Composition and structure: N₂, O₂, Ar, CO₂, ozone; troposphere and stratosphere; pressure and liquid water; gases for biological molecules and industry • Energy processes: UV absorbed by O / O₂ / O₃ dynamic equilibrium; visible light absorbed at the surface; infrared absorbed by greenhouse gases (33 °C warming); heat distribution by winds and ocean currents; transport of water vapour; energy budget and albedo • Global climate change: anthropogenic greenhouse gases, sources, feedback and tipping points, consequences for climate, sea level, currents, species; difficulties of monitoring and prediction; control strategies and agreements • Ozone depletion: CFCs and halons, catalytic destruction, effects of UV, Montreal Protocol and replacements, recovery time • Other threats: aerosols, smoke, land-use change; interconnection of processes • Evaluation: which functions are most at risk; success of ozone controls vs slow progress on greenhouse gases; justified conclusion Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 86-113; AQA spec 3.2.1.	25