

A-level

ENVIRONMENTAL SCIENCE

Paper 2 - Generated practice paper Set 30

Mark scheme

120 marks: Questions 1-10 (95 marks) plus one essay from Question 11 (25 marks).

Marking guidance

- The number in brackets after each marking point is the mark it earns; the note above the points says how many are needed.
- Where a question asks for a fixed number of points (eg *two reasons*), credit only the first points given.
- Alternative correct wording and other relevant, accurate points are credited.
- For levels-of-response questions, first determine the level using the descriptors, then position the mark within the level using the indicative content.
- Calculations: credit correct working (method marks) even if the final answer is wrong; incorrect answers with no working score 0.

Textbook references: every question below cites the printed page numbers in R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018) from which the content is drawn, together with the AQA 7447 specification reference.

Question	Marking guidance	Marks
01.1	<i>1 mark for each point.</i> • Hydrosere (1) • Single-celled algae (and floating plants) - arrive on birds / insects / from soil (1)	2
01.2	<i>1 mark for each point.</i> • Pioneers and later plants grow and die; dead organic matter plus sediment washed in fill the pond so water becomes shallower (1) • As depth falls rooted plants (pondweeds then emergent reeds with leaves above water) can anchor and colonise; emergents shade submerged plants which die out (1) • Sediment / soil accumulates and becomes able to support larger plants; water-tolerant trees (willow, alder) colonise the waterlogged soil (1) • Transpiration by trees and further sediment dry the soil and make it aerobic so oak can colonise; the denser canopy shades out smaller plants (1) • Each community changes abiotic conditions (depth, light, nutrients, oxygen), making it suitable for the next colonisers which out-compete them; conditions become less extreme and biotic factors increasingly important (1)	5
01.3	<i>1 mark for each point.</i> • Starting conditions (bare rock vs open water) differ but each stage makes conditions less extreme and builds soil / organic matter (1) • The final community is controlled by the regional climate (temperature, water availability), which is the same for both - climatic climax (eg oak woodland in the UK) (1)	2

Question	Marking guidance	Marks
01.4	<i>1 mark.</i> Lichens and algae colonise bare rock (extreme temperatures, no soil, little water); rock fragments and dead organic matter accumulate; mosses, then grasses and ferns as thin soil forms; wind-dispersed trees (birch) then animal-dispersed (oak, beech) (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 73-76; AQA spec 3.1.3.3.	
02.1	<i>Correct answer only.</i> Environmental DNA (eDNA) analysis (1)	1
02.2	<i>1 mark for each point.</i> - Transmitters are too heavy for small birds / their batteries would not last for a long-term study (1) - Geolocator tags are small and only record time and light levels; latitude and longitude for each day are worked out when the bird is recaptured and the data downloaded (1)	2
02.3	<i>1 mark for each point.</i> - Each separate (regional) population has its own unique gene pool (1) - DNA from seized ivory can be compared with the database to identify where the elephant came from, so poaching hotspots / trade routes can be targeted (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 404-405; AQA spec 3.1.2.3.6.	
03.1	<i>1 mark for each point.</i> - Net held with its straight edge on the river bed facing the current (open to the flow) (1) - The bed in front is disturbed by kicking / scraping a boot for a standard number of kicks and time (eg 30 s) (1) - Same substrate type, water depth and flow rate at both sites; same net mesh; repeat samples; invertebrates identified with a key and counted (1)	3
03.2	<i>Any 3 from the points below - 1 mark each, maximum 3.</i> - Difficult to standardise the vigour / area kicked (1) - Buried organisms or those fixed to stones are not dislodged (1) - Some organisms swim and avoid the net (1) - Flow rate varies and affects capture (1) - Fragile organisms destroyed (1)	3
03.3	<i>1 mark for each point.</i> - A fixed area of bed is enclosed by a frame so the sampled area is standardised; side nets reduce escapes (1) - But only works in fast enough flow and is hard to set on rough substrates / deep water (1)	2
03.4	<i>1 mark for each point.</i> - Pollutant / oxygen levels fluctuate (eg with flow, time of day) so one reading may miss a pollution event (1) - Sensitive taxa (stonefly, mayfly larvae) disappear and tolerant ones (midge larvae, hoglouse) remain, integrating conditions over weeks / months (biotic index) (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 395-396, 415-416; AQA spec 3.7.2.1.	

Question	Marking guidance	Marks
04.1	<i>1 mark for each point.</i> • Area must be large enough to support viable populations (breeding numbers high enough, gene pool large enough to avoid inbreeding); species higher up food chains need larger areas (lower density) (1) • Large areas contain a full range of resources (eg different fruiting trees / feeding areas through the year) (1) • Shape: the perimeter has edge conditions (a mix of neighbouring habitats); a compact shape maximises core habitat for species needing core conditions (1) • Some species benefit from edges (higher light) or small isolated habitats (ponds too small for predatory fish suit amphibians) (1)	4
04.2	<i>1 mark for each point.</i> • Varied topography: banks, lagoons, islands, ditches created with machinery give a range of water depths (emergent, floating-leaved, submerged plant zones) (1) • Islands provide roosting / nesting sites safe from terrestrial predators (foxes) (1) • Control of salinity (eg intertidal conditions for opossum shrimps feeding dunlin and avocet); scrapes for skylarks; different water depths for waders and ducks (1)	3
04.3	<i>1 mark for each point.</i> • Mobile species (birds, flying insects) colonise easily (1) • Less mobile species (many plants, amphibians, invertebrates) may not reach an isolated site so must be introduced; keystone species especially (1)	2
04.4	<i>1 mark.</i> • Allows dispersal of surplus young, mixing of gene pools (less inbreeding) and recolonisation if a population dies out; avoids hazards (eg rope bridges for howler monkeys over roads) (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 40-42; AQA spec 3.1.2.3.4.	
05.1	<i>1 mark for each point.</i> • Overall decline from 62% to 22% (1) • Sharp falls in 1998 (to 35%) and 2016 (to 18%) coinciding with bleaching events, with partial recovery after each (1)	2
05.2	<i>1 mark for each point.</i> • The polyps expel their symbiotic zooxanthellae so the coral loses colour; bleached corals are alive but grow very slowly and may die if algae do not return (1) • Raised water temperature (climate change / El Niño) (1) • Pollution: sewage, pesticides, cyanide, sunscreen; low oxygen from zooplankton blooms after overfishing (1) • High light / solar activity; ocean acidification hinders skeleton formation (1)	4
05.3	<i>1 mark for each point.</i> • Coastal construction and dredging increase sedimentation / turbidity covering and killing polyps (1) • Sewage and fertiliser nutrients cause algal growth that shades corals and prevents egg release (1) • Physical damage from anchors, divers, souvenir collection; loss of mangroves / seagrass nurseries that trap sediment (1)	3

Question	Marking guidance	Marks
05.4	<i>1 mark.</i> Protected breeding populations grow larger and produce surplus young / adults that spill over into fished areas (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 60-63; AQA spec 3.1.2.3.4.	
06.1	<i>1 mark for each point.</i> - The new opportunity is overwhelmingly attractive (economic benefit) (1) - Lack of knowledge / understanding of the impacts (1) - The impacts occur elsewhere or much later, so are not evident to those who benefit (1)	3
06.2	<i>1 mark for each point.</i> - Cultural belief that resources should be exploited to satisfy needs only (1) - Problems of over-exploitation become obvious quickly as local supplies dwindle, so behaviour changes (1)	2
06.3	<i>1 mark for each point.</i> - Resources are brought in from other areas (1) - If one area is depleted, exploitation moves to another area (eg timber, fish, metals, energy) without the consumers noticing (1)	2
06.4	<i>1 mark for each point.</i> - Easier access to energy has made many new activities possible (1) - The scale of activities has greatly increased (1) - Toxic wastes are produced that cannot be decontaminated by natural processes (1)	3
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 370-371; AQA spec 3.6.1.	
07.1	<i>Any 3 from the points below - 1 mark each, maximum 3.</i> - Sinkers / weighted lines so hooks sink quickly below the reach of albatrosses (1) - Setting lines at night when birds are not feeding (1) - Bird-scaring streamers / decoys (1) - Curved hooks that catch tuna but not birds (1)	3
07.2	<i>1 mark for each point.</i> - Effort: limits on boat size, engine power, days at sea, number of licences; decommissioning payments to reduce fleet size where overfishing is unavoidable (1) - Methods: bans or length limits on drift nets; bans on demersal trawling over sensitive seabed (Rockall Bank, Hatton Bank deep-water reefs) (1) - Shorter trawl tows so turtles / by-catch survive (over 99% survive under 10 minutes) (1)	3
07.3	<i>1 mark for each point.</i> - Eggs / larvae are reared in captivity through the planktonic stage when most die (predation, failure to find habitat) (1) - Juveniles are released when survival chances are good, increasing recruitment to the adult population (1)	2
07.4	<i>1 mark for each point.</i> - Egg / larval survival varies greatly between years so adult numbers do not predict recruitment (1) - Several poor years would lower the MSY; egg surveys allow catches to be reduced in advance rather than after a sudden collapse (1)	2

Question	Marking guidance	Marks
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 344-348; AQA spec 3.5.2.2.	
08.1	<i>1 mark for each point.</i> - Domesticated crops were bred from a small number of original plants so they have a small gene pool / low genetic diversity (1) - Selective breeding for yield, appearance or shelf life may have lost other useful genes (eg taste, disease resistance) (1) - Genetically similar plants share the same susceptibilities so a new disease or environmental change can affect the whole crop (1)	3
08.2	<i>1 mark for each point.</i> - A region with a high concentration of the wild relatives of important crops - usually where the crop was first domesticated (eg wheat / barley in the Middle East; potato, tomato in South America; rice in India; maize in Central America) (1) - They hold the largest gene pools / most genetic variety for future crop breeding (disease resistance, drought and salt tolerance, yield) (1) - Many are in regions where environmental degradation / habitat loss threatens the wild plants (1)	3
08.3	<i>1 mark for each point.</i> - A large population may all descend from a few closely related individuals so they are genetically similar (1) - Similar adaptations mean all are susceptible to the same change (disease, climate) (1) - Greater risk of inbreeding so disadvantageous recessive genes are expressed (eg amelanism, poor muscle development in lion cubs); so species should be protected across their whole range to retain regional genes (1)	3
08.4	<i>1 mark.</i> Morama bean (drought resistant, soya-like protein); yeheb tree (drought resistant nuts, poor soils); potato bean (high protein); perennial maize / Kernza (no annual replanting, less ploughing / erosion); salt-tolerant spinach relatives (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 15-17; AQA spec 3.1.2.1.2.	
09.1	<i>Up to 6 marks in total; award marks in proportion to the completeness of each point.</i> - Non-indigenous species (Sitka spruce from North America) chosen for fast growth and pest resistance; indigenous wildlife's needs for food, pollination, seed dispersal and habitat are not met (up to 2) - Single species: easy planting, thinning and felling over large areas; but each tree species supports its own community so few species overall (up to 2) - Close planting makes trees compete for light so they grow tall and straight (good timber) but shades out ground flora and the species that depend on them (up to 2) - Simple age structure: uniform size makes management easy but gives little variety of light, temperature, wind; no old trees or dead wood for invertebrates and woodpeckers (up to 2) - Mixed woodland: varied niches, stratification, dead wood, indigenous trees each with their own dependent species (up to 2)	6

Question	Marking guidance	Marks
09.2	1 mark for each point. - Trees are felled when income will be greatest - approaching full size when the growth rate starts to decline (1) - Continuing to grow them ties up land and capital for little extra timber; natural forests keep old and dead trees (1)	2
09.3	1 mark for each point. - Industrial society replaced species-specific uses (baskets, fencing, charcoal, tool handles) with large-scale uses: chipboard, plywood, pulp (1) - Wood fuel replaced by fossil fuels / electricity; so traditional management declined and woods became uniform (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 365-367; AQA spec 3.5.3.2.	
10.1	Levels of response Level 3 (7-9 marks): Answer is comprehensive, relevant and logically structured. Shows detailed knowledge and understanding of the relevant scientific ideas and applies them accurately to the context. Evaluation/analysis is well supported by evidence and a clear judgement or conclusion is reached. Level 2 (4-6 marks): Answer is mostly relevant and shows some structure. Knowledge and understanding of relevant ideas is sound but application to the context or evaluation may be incomplete or lack detail. A judgement may be made but is not fully supported. Level 1 (1-3 marks): Answer is basic, with limited structure. Some relevant knowledge is shown but there are omissions or errors. Little application to the context and little or no evaluation. 0 marks: No relevant content. Indicative content (credit other relevant, accurate points) - Why needed: growing population (to 12 billion), land degradation since the fertile crescent, reliance on artificial fertilisers, pesticides and unsustainable irrigation; limits to new land and imports - Pest control: cultural methods (weeding, mulching, rotation, barrier / companion crops, predator habitats, biological control, sterile males, pheromones), resistant varieties, integrated control, reduced antibiotics - Nutrients: natural processes - nitrogen-fixing legumes, decomposition, recycling organic matter (manure, compost), crop rotation, permaculture, conservation of soil biota and mycorrhizae, reduced artificial fertiliser; phosphate recycling - Energy: natural processes instead of Haber-process fertiliser, low tillage, less machinery, renewable energy, low food miles, reduced methane (cattle diets, rice varieties) - Water: low water-use crops, drip irrigation, reservoirs and aquifer recharge, avoiding saline water; soil: organic matter maintenance, erosion control - Biodiversity: retaining hedgerows, ditches, ponds, woodlands; CWR conservation and seed banks; agri-environment schemes - Social: control of impacts on rural communities, Fairtrade, local food; consumer choices - Evaluation: yields vs sustainability, cost, need for policy support; judgement	9
10.2	1 mark. Agricultural systems designed to mimic natural ecosystems - perennial crops, mixed species, recycling of nutrients, minimal external inputs (1)	1

Question	Marking guidance	Marks
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 335-337; AQA spec 3.5.1.5.	
11.1	Natural systems have remained sustainable over very long periods. Discuss how the features of natural systems could be copied to make human systems more sustainable. Levels of response Level 5 (21-25 marks): Comprehensive and detailed knowledge and understanding drawn from across the specification, applied accurately to the question. Argument is coherent, well-structured and supported by a wide range of relevant, well-chosen examples and quantitative detail where appropriate. Interconnections between topics are made explicitly and a clear, justified conclusion is reached. Level 4 (16-20 marks): Detailed knowledge and understanding with good application to the question. Argument is logically organised with relevant examples; some links between different areas of the specification are made. Conclusions are drawn but may be less fully justified. Level 3 (11-15 marks): Sound knowledge and understanding, mostly applied to the question. Some structure and relevant examples, but coverage may be uneven or descriptive rather than analytical. Limited linking of topics. Level 2 (6-10 marks): Basic knowledge and understanding with partial relevance. Answer is largely descriptive with few examples and limited structure; some errors or omissions. Level 1 (1-5 marks): Limited, fragmented knowledge with little relevance to the question. Little structure; examples absent or inappropriate. 0 marks: No relevant content. Indicative content (students are not expected to cover all of these; credit other relevant material) • Dynamic equilibria and negative feedback in climate, hydrological cycle and populations; positive feedback and tipping points caused by humans • Energy: solar-driven, low energy density, low temperature (photosynthesis, enzymes, nitrogen fixation vs Haber process); renewable energy, heat pumps, bioleaching, fuel cells • Materials: cycles, few abundant elements, biodegradable non-toxic wastes vs fossil fuels and dispersed minerals; recycling, Cradle to Cradle, industrial symbiosis • Diversity and stability: rainforests / reefs vs agroecosystems; diverse energy mixes • Optimum vs maximum production; supporting ecosystems in agriculture, forestry, fisheries, urban design • Case studies and metrics; limits and judgement Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 370-379; AQA spec 3.6.1-3.6.4.	25

Question	Marking guidance	Marks
11.2	Discuss the methods that can be used to conserve biodiversity and evaluate their effectiveness. Levels of response Level 5 (21-25 marks): Comprehensive and detailed knowledge and understanding drawn from across the specification, applied accurately to the question. Argument is coherent, well-structured and supported by a wide range of relevant, well-chosen examples and quantitative detail where appropriate. Interconnections between topics are made explicitly and a clear, justified conclusion is reached. Level 4 (16-20 marks): Detailed knowledge and understanding with good application to the question. Argument is logically organised with relevant examples; some links between different areas of the specification are made. Conclusions are drawn but may be less fully justified. Level 3 (11-15 marks): Sound knowledge and understanding, mostly applied to the question. Some structure and relevant examples, but coverage may be uneven or descriptive rather than analytical. Limited linking of topics. Level 2 (6-10 marks): Basic knowledge and understanding with partial relevance. Answer is largely descriptive with few examples and limited structure; some errors or omissions. Level 1 (1-5 marks): Limited, fragmented knowledge with little relevance to the question. Little structure; examples absent or inappropriate. 0 marks: No relevant content. Indicative content (students are not expected to cover all of these; credit other relevant material) • Setting priorities: IUCN Red List criteria, EDGE, endemic, keystone, flagship species, population dispersal • Legislation and protocols: Wildlife and Countryside Act, designated areas (SSSI to MCZ), CITES Appendices, IWC, EU CFP, ITTO - successes and limits of enforcement • Captive breeding and release: selection, difficulties, techniques, release site criteria, hard / soft release, monitoring; seed banks • Habitat conservation: creation, design (area, shape, corridors, age structure, colonisation), abiotic and biotic management, plagioclimax maintenance, population control, reintroductions • Habitat case studies (woodland, rainforest, coral reefs, islands, mangroves, Antarctica) illustrating threats and conservation efforts • Monitoring and new technologies informing decisions • Evaluation: examples of success (Iberian lynx, Hawaiian goose, Scilly rats, Yellowstone) and failure (ITTO, tigers); cost, politics, climate change; judgement on the most effective combination Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 26-71; AQA spec 3.1.2.3.	25