

A-level

ENVIRONMENTAL SCIENCE

Paper 2 - Generated practice paper Set 33

Mark scheme

120 marks: Questions 1-10 (95 marks) plus one essay from Question 11 (25 marks).

Marking guidance

- The number in brackets after each marking point is the mark it earns; the note above the points says how many are needed.
- Where a question asks for a fixed number of points (eg *two reasons*), credit only the first points given.
- Alternative correct wording and other relevant, accurate points are credited.
- For levels-of-response questions, first determine the level using the descriptors, then position the mark within the level using the indicative content.
- Calculations: credit correct working (method marks) even if the final answer is wrong; incorrect answers with no working score 0.

Textbook references: every question below cites the printed page numbers in R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018) from which the content is drawn, together with the AQA 7447 specification reference.

Question	Marking guidance	Marks
01.1	<i>1 mark for each point.</i> • Each regional population has its own unique gene pool / DNA profile (1) • DNA from seized timber or ivory is compared with the database of regional populations (1) • This identifies where the material came from, showing whether it was legally harvested and where poaching occurs (1)	3
01.2	<i>1 mark for each point.</i> • Radio / GPS transmitters use radio waves - work in air and fresh water, giving positions (GPS stores or transmits location) (1) • Radio waves do not travel through seawater so acoustic transmitters (sound) are used for marine animals, giving real-time positions (1)	2
01.3	<i>Up to 3 marks in total; award marks in proportion to the completeness of each point.</i> • Drones / ROVs, AUVs (eg Autosub), balloons, aircraft, satellites, animals (tags) (up to 2) • Drones: cheap, low disturbance, high-resolution images of areas hard to reach (cliffs, canopy, ice), repeated surveys; infrared cameras detect animals at night (up to 2)	3
01.4	<i>1 mark for each point.</i> • Cameras record automatically (visible light by day, infrared at night) without researchers being present (1) • So animals behave naturally and nests are not deserted; monitoring continues 24 hours (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 404-406; AQA spec 3.7.2.3.	

Question	Marking guidance	Marks
02.1	<i>1 mark for each point.</i> • Silver carp eat phytoplankton, bighead carp zooplankton, grass carp and tilapia vegetation, common carp bottom-feed on molluscs / insects, catfish or perch eat small fish (1) • Species occupy different niches / trophic levels so they do not compete and all food sources are used (1) • Bottom feeders re-suspend nutrients boosting plankton / plant growth (1) • Predators control over-breeding species (tilapia) so remaining fish grow larger; total yield exceeds any single species (1)	4
02.2	<i>1 mark for each point.</i> • IMTA: fed species (salmon, shrimp) alongside inorganic extractive species (seaweeds absorbing ammonia and phosphate) and organic extractive species (filter-feeding mussels eating waste particles / plankton) (1) • Wastes of one species become nutrients for another so less pollution and extra harvests (1) • Aquaponics: aquaculture water rich in nutrients / organic matter feeds hydroponic salad crops (lettuce, herbs, watercress) (1) • Plants remove the nutrients that would cause deoxygenation and cleaned water returns to the fish (tilapia, carp, catfish); rice-fish paddies similar (1)	4
02.3	<i>1 mark for each point.</i> • Bluefin tuna do not breed well in captivity so young fish are caught from the wild and fattened in sea cages (1) • Production still depends on (and removes juveniles from) the wild population; carnivores need large fish inputs (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 357-358; AQA spec 3.5.2.3.	
03.1	<i>2 marks for each point (partial credit for a partly correct point).</i> • Normal diagonal mesh closes as the net fills / stretches so small fish cannot escape; square-mesh panels do not close so juveniles escape to grow and breed; TEDs let turtles out; larger mesh sizes generally (2) • Minimum size lets fish reach breeding age before capture (eg UK lobster minimum carapace above breeding size, cod 35 cm); maximum size / V-notching egg-carrying females protects large, highly fecund breeding adults (Hebridean lobsters over 145 mm, Nassau grouper) (2) • NTZs (Lundy, Lamlash Bay, Chagos, Great Barrier Reef) protect breeding populations and habitats; surplus young spill over and increase surrounding catches; damaged communities recover (2) • Closed seasons, ideally in the breeding season, let fish grow and breed; fishermen may oppose because shoaling breeders are easy to catch (2)	8
03.2	<i>1 mark for each point.</i> • Pelagic species form single-species shoals so fishing stops when the quota is reached (1) • In mixed fisheries many species are caught together; once one species' quota is reached it must be discarded (dead) while fishing for others continues (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 344-348; AQA spec 3.5.2.2.	
04.1	<i>1 mark.</i> • Development that meets the needs of the present generation without compromising the ability of future generations to meet their own needs (1)	1

Question	Marking guidance	Marks
04.2	<i>1 mark for each point.</i> - Impact depends on total population x per capita impact (1) - With few people, natural processes (forest regrowth, fish recruitment, aerobic decomposition / dilution) recover faster than the resource is used or the waste added (1) - Larger populations exceed the rate of natural recovery: forest and soil do not regenerate, catches exceed MSY, water deoxygenates (1)	3
04.3	<i>1 mark for each point.</i> - Urban areas are supported by resources brought in from elsewhere; when one area is depleted, exploitation simply moves (timber, fish, minerals, energy) so consumers see no shortage (1) - Rural communities exploit local resources directly so the effects of over-exploitation become obvious quickly, encouraging cultural limits to need (1)	2
04.4	<i>Up to 4 marks in total; award marks in proportion to the completeness of each point.</i> - Climate: warming increases evaporation, cloud and albedo; higher CO₂ increases photosynthesis and sequestration; hydrological cycle: more evaporation gives more precipitation; populations: density-dependent factors return numbers to carrying capacity (up to 2) - In agriculture soil degradation, water shortage and loss of predators would reduce yields (negative feedback limiting the activity) (up to 2) - Humans respond with fertilisers, irrigation and pesticides to maintain yields, further damaging the natural systems that support production (up to 2)	4
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 370-372; AQA spec 3.6.1.	
05.1	<i>2 marks for each point (partial credit for a partly correct point).</i> - Drainage makes soils aerobic and drier, increasing grass growth but wetland plants and the animals they support (waders, amphibians, invertebrates) cannot survive (2) - Fertiliser increases nutrient availability so vigorous species (grasses, nettles) grow taller and out-compete smaller wildflowers for light; diversity falls (2) - Hedgerow removal destroys habitat / corridors and nesting sites; loss of natural pest predator habitats increases pest damage; larger fields increase wind erosion; genetic contamination and introduced species also accepted (2)	6
05.2	<i>1 mark for each point.</i> - Long-continued traditional farming has created plagioclimax habitats colonised by many species: hay meadows, chalk grassland, grazed heath / moorland, hedgerows (1) - These communities depend on the farming continuing (1)	2
05.3	<i>1 mark for each point.</i> - CO₂ from fossil fuel use in machinery and embodied energy of fertiliser (Haber process) and machinery; ploughing increases soil respiration (1) - Methane from livestock intestines and rice paddies; nitrous oxide from manure and nitrogen fertilisers (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 328-330; AQA spec 3.5.1.3.	

Question	Marking guidance	Marks
06.1	<i>1 mark for each point.</i> • Zooxanthellae are photosynthetic algae living inside polyp cells (up to 30 000 per mm²) (1) • They use the polyp's CO₂ and excretory wastes (nitrogen, phosphorus) for photosynthesis and growth, so nutrients are recycled efficiently (1) • They supply the polyp with glucose and oxygen - over 95% of its food energy; the polyp provides a protected habitat and nutrients (1) • Polyps also trap plankton with cilia and stinging nematocysts; surplus CO₂ is used to build the calcium carbonate skeleton (1)	4
06.2	<i>1 mark for each point.</i> • High light (shallow, clear water) for the algae; warm stable temperatures (about 18-30 °C); low turbidity; constant salinity; well-oxygenated water (1) • Found only where all these coincide (about 1% of ocean area) (1) • Small changes (warming, sediment, freshwater runoff) move conditions outside the narrow range of tolerance (1)	3
06.3	<i>Any 3 from the points below - 1 mark each, maximum 3.</i> • Fisheries: support over a quarter of marine species including fish, lobsters, conch (1) • Erosion protection: complex structure dissipates wave energy protecting low coral islands; coral sand builds shores (1) • Medicinal discoveries: AZT and Ara-C from sponges, dolastatin from sea hares, bryostatin from Bugula, trabectedin from sea squirts, secosteroids, coral for bone grafts (1) • Climate control: carbonate skeletons sequester CO₂ (1) • Tourism (diving, snorkelling) (1)	3
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 57-60; AQA spec 3.1.2.3.4.	
07.1	<i>2 marks for each point (partial credit for a partly correct point).</i> • Light: selective felling to create clearings for plants needing high light while retaining dense canopy for shade species; clearings also give warm sand / soil for lizard eggs; leave dead wood; woodland floor plants (bluebells) grow before the canopy leaves (2) • Dissolved oxygen: narrowing a slow river increases flow and turbulence so more oxygen dissolves, suiting trout and mayfly larvae; shallow water warms quickly for egg development (2) • Nutrients: removing fertile topsoil / not fertilising allows plants of nutrient-poor soils (eg sundew on acidic nitrate-deficient soils) that cannot compete on fertile ground; dung patches add local nutrients (2)	6
07.2	<i>Any 4 from the points below - 1 mark each, maximum 4.</i> • Food: wildflower grasslands for seed-eating birds; providing food species (1) • Control of predation: island nest sites unreachable by foxes; trapping / culling / eradicating introduced rats (Scilly Isles - Manx shearwater and storm petrel breeding success increased) (1) • Control of competitors: removing rhododendron so wildflowers get light (1) • Pollination: a range of flowering plants with different flowering times so pollinators have food all season (1) • Seed dispersal species (hippos, elephants, monkeys) conserved; control of introduced pathogens (1)	4
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 43-46; AQA spec 3.1.2.3.4, 3.1.3.1.	

Question	Marking guidance	Marks
08.1	<i>Correct answer only.</i> • Waste products that are the raw materials for other processes (1)	1
08.2	<i>1 mark for each point.</i> • Bacteria fix nitrogen at low temperature using enzymes and energy ultimately from sunlight (renewable) (1) • The Haber process needs high temperature and pressure powered by non-renewable fossil fuels, releasing CO ₂ (1)	2
08.3	<i>1 mark for each point.</i> • Incineration uses high temperatures / fuel and releases pollutants; nutrients are not returned as usable raw materials (1) • Decomposition at low temperature recycles nutrients as the raw materials for other natural processes (material cycle) (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 373; AQA spec 3.6.2-3.6.3.	
09.1	<i>1 mark for each point.</i> • As sea otters declined (1960-1980) sea urchins increased (to 350) and kelp area fell (to 25) (1) • As otters recovered after 1990 urchins declined and kelp recovered (1) • Otter and kelp trends are positively correlated; urchins negatively correlated with both (1)	3
09.2	<i>1 mark for each point.</i> • Sea otters are predators of sea urchins so fewer otters allowed the urchin population to rise (1) • Sea urchins graze kelp so the kelp forest was over-grazed and destroyed (1) • Kelp forests are habitat / food for many species including otters (keystone species / trophic cascade); otter recovery restored the balance (1)	3
09.3	<i>Any 2 from the points below - 1 mark each, maximum 2.</i> • Over-fishing of sandeels reduced puffins (1) • Over-collection of turtle eggs reduced turtles so jellyfish (their food) increased; over-fishing of sardines / herring (competitors) also increased jellyfish (1) • Loss of pollinating bees (pesticides, wild flower loss) reduces plant reproduction; loss of elephants / large herbivores reduces seed dispersal of trees (1)	2
09.4	<i>1 mark for each point.</i> • They control abiotic features: beaver dams create ponds colonised by aquatic species; elephants create clearings and keep water holes open (1) • Species that depend on these features (light, water, minerals) cannot survive in their absence (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 23-25; AQA spec 3.1.2.2.	

Question	Marking guidance	Marks
10.1	Levels of response Level 3 (7-9 marks): Answer is comprehensive, relevant and logically structured. Shows detailed knowledge and understanding of the relevant scientific ideas and applies them accurately to the context. Evaluation/analysis is well supported by evidence and a clear judgement or conclusion is reached. Level 2 (4-6 marks): Answer is mostly relevant and shows some structure. Knowledge and understanding of relevant ideas is sound but application to the context or evaluation may be incomplete or lack detail. A judgement may be made but is not fully supported. Level 1 (1-3 marks): Answer is basic, with limited structure. Some relevant knowledge is shown but there are omissions or errors. Little application to the context and little or no evaluation. 0 marks: No relevant content. Indicative content (credit other relevant, accurate points) • Resources: wood, fibres (cotton, wool, silk), oils, fuels (wood, biofuels), food; new food species (morama bean, perennial cereals, eland, bison) better adapted to local conditions • Biomimetics: vehicle design, architecture, structures, adhesion, materials, ultrasound • Medicines from plants and animals (examples); most species unstudied - argument for habitat conservation • Physiological research using wildlife species; wildlife as pest control agents (ladybirds, Encarsia, Cactoblastis, Bt) • Genetic resources: crop wild relatives, Vavilov centres, gene pools • Ecosystem services: atmospheric composition, hydrological and biogeochemical cycles, soil maintenance, pollination, seed dispersal, habitat provision • Ethical / aesthetic / recreational arguments; the unknown value of species not yet investigated (precautionary principle) • Evaluation: which arguments are most persuasive to decision makers; economic valuation vs intrinsic value; judgement	9
10.2	<i>1 mark.</i> • The variety and abundance of life (species, genes, habitats) in an area / on Earth (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 10-19; AQA spec 3.1.2.1.	

Question	Marking guidance	Marks
11.1	Discuss the methods that can be used to conserve biodiversity and evaluate their effectiveness. Levels of response Level 5 (21-25 marks): Comprehensive and detailed knowledge and understanding drawn from across the specification, applied accurately to the question. Argument is coherent, well-structured and supported by a wide range of relevant, well-chosen examples and quantitative detail where appropriate. Interconnections between topics are made explicitly and a clear, justified conclusion is reached. Level 4 (16-20 marks): Detailed knowledge and understanding with good application to the question. Argument is logically organised with relevant examples; some links between different areas of the specification are made. Conclusions are drawn but may be less fully justified. Level 3 (11-15 marks): Sound knowledge and understanding, mostly applied to the question. Some structure and relevant examples, but coverage may be uneven or descriptive rather than analytical. Limited linking of topics. Level 2 (6-10 marks): Basic knowledge and understanding with partial relevance. Answer is largely descriptive with few examples and limited structure; some errors or omissions. Level 1 (1-5 marks): Limited, fragmented knowledge with little relevance to the question. Little structure; examples absent or inappropriate. 0 marks: No relevant content. Indicative content (students are not expected to cover all of these; credit other relevant material) • Setting priorities: IUCN Red List criteria, EDGE, endemic, keystone, flagship species, population dispersal • Legislation and protocols: Wildlife and Countryside Act, designated areas (SSSI to MCZ), CITES Appendices, IWC, EU CFP, ITTO - successes and limits of enforcement • Captive breeding and release: selection, difficulties, techniques, release site criteria, hard / soft release, monitoring; seed banks • Habitat conservation: creation, design (area, shape, corridors, age structure, colonisation), abiotic and biotic management, plagioclimax maintenance, population control, reintroductions • Habitat case studies (woodland, rainforest, coral reefs, islands, mangroves, Antarctica) illustrating threats and conservation efforts • Monitoring and new technologies informing decisions • Evaluation: examples of success (Iberian lynx, Hawaiian goose, Scilly rats, Yellowstone) and failure (ITTO, tigers); cost, politics, climate change; judgement on the most effective combination Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 26-71; AQA spec 3.1.2.3.	25

Question	Marking guidance	Marks
11.2	Discuss how an understanding of energy flow and the manipulation of food species can be used to increase food production sustainably. Levels of response Level 5 (21-25 marks): Comprehensive and detailed knowledge and understanding drawn from across the specification, applied accurately to the question. Argument is coherent, well-structured and supported by a wide range of relevant, well-chosen examples and quantitative detail where appropriate. Interconnections between topics are made explicitly and a clear, justified conclusion is reached. Level 4 (16-20 marks): Detailed knowledge and understanding with good application to the question. Argument is logically organised with relevant examples; some links between different areas of the specification are made. Conclusions are drawn but may be less fully justified. Level 3 (11-15 marks): Sound knowledge and understanding, mostly applied to the question. Some structure and relevant examples, but coverage may be uneven or descriptive rather than analytical. Limited linking of topics. Level 2 (6-10 marks): Basic knowledge and understanding with partial relevance. Answer is largely descriptive with few examples and limited structure; some errors or omissions. Level 1 (1-5 marks): Limited, fragmented knowledge with little relevance to the question. Little structure; examples absent or inappropriate. 0 marks: No relevant content. Indicative content (students are not expected to cover all of these; credit other relevant material) • Energy flow: autotrophs and heterotrophs; losses by reflection, transmission, respiration; 100 000 to 1 000 units; few food chains beyond four levels; plant food most productive; grazing animals where crops impossible; pigs on wastes • Energetics: intensive vs extensive, law of diminishing returns, energy subsidies (fertiliser, pesticides, irrigation, machinery, drying, processing, transport), energy ratios, food conversion ratios, dependence on fossil fuels • Control of genetics: asexual reproduction / cuttings / cloning, selective breeding and breeds, cross-breeding and hybrid vigour, AI and embryo transfer • GM: Roundup Ready soya, Bt crops, golden rice, omega-3 oilseed rape; advantages; risks (gene transfer, antibiotic markers, allergies, costs to LEDC farmers, loss of local diversity) • Sustainability links: energy dependence, trophic level of diet, feeding 12 billion people • Justified conclusion on the balance between productivity, efficiency and sustainability Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 321-328; AQA spec 3.5.1.2.	25