

A-level

ENVIRONMENTAL SCIENCE

Questions by topic: Agroecosystems: abiotic & biotic factors - Set 1

Mark scheme

AQA spec 3.5.1.1; Biological resources; textbook pp. 306-320

Marking guidance

- The number in brackets after each marking point is the mark it earns; the note above the points says how many are needed.
- Where a question asks for a fixed number of points (eg *two* reasons), credit only the first points given.
- Alternative correct wording and other relevant, accurate points are credited.
- For levels-of-response questions, first determine the level using the descriptors, then position the mark within the level using the indicative content.
- Calculations: credit correct working (method marks) even if the final answer is wrong; incorrect answers with no working score 0.

Textbook references: every question below cites the printed page numbers in R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018) from which the content is drawn, together with the AQA 7447 specification reference.

Question	Marking guidance	Marks
01.1	<i>1 mark for each point.</i> • B: frost-free period (150 days) exceeds the 120 needed whereas A (110) is too short - late frosts would kill seedlings (1) • B is warmer / longer thermal growing season because south-facing slopes receive more insolation; valley floors are frost pockets where cold dense air collects (1) • But B has lower rainfall and a steeper slope (erosion / machinery difficulties) so irrigation / contour cultivation may be needed (1)	3
01.2	<i>Any 3 from the points below - 1 mark each, maximum 3.</i> • Greenhouses / polytunnels heated by solar gain or burning gas / paraffin (ventilated to prevent overheating) (1) • Woven plastic cloches / fleece over rows to retain warm air near the ground (1) • Burning oil / gas or using large fans in orchards to prevent frost damage to blossom (1) • Site selection (aspect, avoiding frost pockets); heated buildings and shelter for livestock (1)	3
01.3	<i>1 mark for each point.</i> • Mammals and birds use food energy to maintain a constant body temperature (thermoregulation) (1) • Reducing heat loss (shelter, heated buildings) means less food energy is used for heat and more is available for growth (1)	2

Question	Marking guidance	Marks
01.4	<i>1 mark for each point.</i> • Long days increase milk production and egg laying; poultry grow best with short days; sheep mate as days shorten in autumn (1) • Artificial lighting extends the growing season for greenhouse crops or simulates autumn in spring to give sheep a second mating / lambing season (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 306-309; AQA spec 3.5.1.1.	
02.1	<i>1 mark for each point.</i> • Plants close their stomata to reduce transpiration losses (1) • Closed stomata stop the absorption of carbon dioxide so photosynthesis and growth stop (1) • Less water for nutrient absorption / transport and cell turgidity; severe shortage dehydrates cells and inhibits biochemical reactions (1)	3
02.2	<i>Up to 4 marks in total; award marks in proportion to the completeness of each point.</i> • Anaerobic soil favours denitrifying bacteria and inhibits nitrifying bacteria so nitrate is lost faster than it forms; higher risk of fungal disease; root respiration inhibited (up to 2) • Drainage ditches or pipes (up to 2) • Deep ploughing; avoiding compaction by machinery / livestock; adding organic matter to encourage worms (up to 2)	4
02.3	<i>1 mark for each point.</i> • Groundwater often contains dissolved salts; when the water evaporates from the soil the salts are left behind and accumulate (1) • High soil salinity causes osmotic dehydration of roots / crop death; washing out with extra water uses more water and salinises rivers (1)	2
02.4	<i>1 mark.</i> • Add crushed lime (calcium carbonate / hydroxide); low pH increases leaching, inhibits nutrient uptake and mobilises toxic aluminium and lead (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 308-310, 313; AQA spec 3.5.1.1.	
03.1	<i>Up to 6 marks in total; award marks in proportion to the completeness of each point.</i> • Organic advantages: often waste products, locally available and cheap; add humus improving structure and water retention; increase soil biota; slow release reduces leaching (up to 2) • Organic disadvantages: nutrient composition cannot be controlled; released slowly so needs long-term planning; bulky with high water content so expensive to transport; cannot be applied to a growing crop; odour / pathogens (up to 2) • Inorganic advantages: composition controlled to match crop needs; nutrients released rapidly; concentrated and easy to apply (up to 2) • Inorganic disadvantages: high energy (fossil fuel) manufacture (Haber process); no organic matter so humus and biota decline; some toxic to worms; high solubility so leached causing eutrophication; raw materials (phosphate rock) finite (up to 2)	6

Question	Marking guidance	Marks
03.2	<i>Up to 4 marks in total; award marks in proportion to the completeness of each point.</i> • Crops grown in a circulating nutrient solution (inert medium) in greenhouses; nutrient concentrations monitored and adjusted so nutrients are never limiting; abiotic factors controlled (up to 2) • Roots are small (all in contact with nutrients) so more growth goes into the harvested crop; no soil-borne pathogens or weeds; crop stays fresh with roots (up to 2) • Disadvantages: high inputs of nutrients and energy (heating, lighting, pumping); high level of technical knowledge; only economic for high-value crops (up to 2)	4
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 310-312; AQA spec 3.5.1.1.	
04.1	<i>Up to 4 marks in total; award marks in proportion to the completeness of each point.</i> • Rotation: each crop has its own pests; growing a different crop each year (4-5 year cycle) means pests left at the end of a season die before that crop returns, so populations cannot build up early (up to 2) • Beetle banks / hedgerows provide habitat and overwintering sites for natural predators (ground beetles, ladybirds) that eat aphids (up to 2) • Rotation also evens out nutrient demands and includes legumes (up to 2)	4
04.2	<i>1 mark for each point.</i> • Integrated control gives almost the same yield as chemical control (18.1 vs 18.5) with far fewer applications (1-2 vs 6) (1) • It retains most of the biodiversity of cultural control (26 vs 31 species) whereas chemical control alone leaves only 8 (1) • Cultural control alone loses about 12% yield; integrated control combines cultural methods, resistant varieties and carefully timed specific non-persistent pesticides only when essential - best balance of yield, cost and impact (1)	3
04.3	<i>1 mark for each point.</i> • Large numbers of males are sterilised with gamma radiation and released (1) • Females of many insects mate only once and store sperm, so mating with a sterile male produces no offspring; enough sterile males reduce fertile matings below replacement level (screw worm fly, Mediterranean fruit fly) (1) • Only the target species' females mate with them so non-target species are unaffected; fails if sterilised males behave abnormally or re-colonisation occurs (1)	3
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 314-318; AQA spec 3.5.1.1.	

Question	Marking guidance	Marks
05.1	<i>Up to 4 marks in total; award marks in proportion to the completeness of each point.</i> • Success: Cactoblastis moth introduced from South America controlled prickly pear cactus in Australia; Encarsia parasitic wasps control greenhouse whitefly (up to 2) • Problems: cane toads introduced to Australia to control sugar cane beetles did not control them but ate many other species and are toxic to predators; rosy wolfsnail introduced to Polynesia to control African land snails ate endemic Partulid snails (48 species extinct); harlequin ladybirds introduced to Europe eat native ladybirds (up to 2) • Lesson: control agents should be specialist feeders on the pest (up to 2)	4
05.2	<i>1 mark for each point.</i> • Release an artificial sex scent attracting males (or females) of the pest species (1) • Used to monitor when the pest arrives so pesticide is timed correctly, or (in enclosed greenhouses) to trap enough of one sex to prevent breeding (1)	2
05.3	<i>1 mark for each point.</i> • Regular low doses kill only the most sensitive bacteria so resistant strains survive and multiply (1) • Agricultural use exceeds medical use, so resistance spreads (1) • Resistant zoonotic bacteria (E. coli, Salmonella, Campylobacter) can transfer to humans and are harder to treat; hence reduced use of antibiotics is a sustainability goal (1)	3
05.4	<i>1 mark.</i> • Grow nectar-rich flowering plants alongside crops; restrict pesticides harmful to bees / spray when not in flower; introduce bee hives (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 316-320; AQA spec 3.5.1.1.	