

Optimal Rates for Generalization of Gradient Descent for Deep ReLU Classification

Yuanfan Li

School of Mathematical Sciences
Zhejiang University
22335070@zju.edu.cn

Yunwen Lei

Department of Mathematics
The University of Hong Kong
leiyw@hku.hk

Zheng-Chu Guo

School of Mathematical Sciences
Zhejiang University
guozc@zju.edu.cn

Yiming Ying *

School of mathematics and statistics
University of Sydney
yiming.ying@sydney.edu.au

Abstract

Recent advances have significantly improved our understanding of the generalization performance of gradient descent (GD) methods in deep neural networks. A natural and fundamental question is whether GD can achieve generalization rates comparable to the minimax optimal rates established in the kernel setting. Existing results either yield suboptimal rates of $O(1/\sqrt{n})$, or focus on networks with smooth activation functions, incurring exponential dependence on network depth L . In this work, we establish optimal generalization rates for GD with deep ReLU networks by carefully trading off optimization and generalization errors, achieving only polynomial dependence on depth. Specifically, under the assumption that the data are NTK separable from the margin γ , we prove an excess risk rate of $\tilde{O}(L^4(1 + \gamma L^2)/(n\gamma^2))$, which aligns with the optimal SVM-type rate $\tilde{O}(1/(n\gamma^2))$ up to depth-dependent factors. A key technical contribution is our novel control of activation patterns near a reference model, enabling a sharper Rademacher complexity bound for deep ReLU networks trained with gradient descent.

1 Introduction

Deep neural networks trained through first-order optimization methods have achieved remarkable empirical success in diverse domains [Krizhevsky et al., 2012]. Despite their widespread adoption, a rigorous theoretical understanding of their optimization dynamics and generalization behavior remains incomplete, particularly for ReLU networks. The inherent challenges arise from the non-smoothness and non-convexity of the loss landscape induced by ReLU activations and network architectures, which complicate the classical analysis. Intriguingly, empirical evidence demonstrates that over-parameterized models often achieve zero training error but still generalize well even in the absence of explicit regularization [Zhang et al., 2016]. This phenomenon has spurred significant theoretical research to understand its underlying mechanisms.

A prominent line of research uses the neural tangent kernel (NTK) framework to analyze neural network training [Jacot et al., 2018]. In the infinite-width limit, gradient descent (GD) dynamics can be characterized by functions in the NTK’s reproducing kernel Hilbert space (RKHS), with convergence guarantees established for both shallow and deep networks [Du et al., 2019b,a]. These

*Corresponding author

	Activation	Width	Training error	Generalization error	Network
Ji and Telgarsky	ReLU	$\tilde{\Omega}\left(\frac{1}{\gamma^8}\right)$	$\tilde{O}\left(\frac{1}{\gamma^{2T}}\right)$	$\tilde{O}\left(\frac{1}{\gamma^2\sqrt{n}}\right)$	Shallow
Chen et al.	ReLU	$\tilde{\Omega}\left(\frac{L^{22}}{\gamma^8}\right)$	$\tilde{O}\left(\frac{1}{\gamma^{2T}}\right)$	$\tilde{O}\left(\frac{e^{O(L)}}{\gamma^2}\sqrt{\frac{m}{n}}\right)$	Deep
Taheri et al.	Smooth	$\tilde{\Omega}\left(\frac{1}{\gamma^{6L+4}}\right)$	$\tilde{O}\left(\frac{1}{\gamma^{2T}}\right)$	$\tilde{O}\left(\frac{e^{O(L)}}{\gamma^2 n}\right)$	Deep
Ours	ReLU	$\tilde{\Omega}\left(\frac{L^{16}}{\gamma^8}\right)$	$\tilde{O}\left(\frac{1}{\gamma^{2T}}\right)$	$\tilde{O}\left(\frac{L^4(1+\gamma L^2)}{\gamma^2 n}\right)$	Deep

Table 1: Comparison of learning neural networks with GD on NTK separable data with prior works. Here m is the network width, L is the network depth, n is the sample size, γ is the NTK-margin, T is the number of iterations.

results demonstrate that GD converges to global minima within a local neighborhood of initialization, provided that the network width is sufficiently large. In particular, the appealing work [Arora et al., 2019a] showed that, if the network width $m = \tilde{O}(\text{poly}(n, 1/\lambda_0, L))$, then the generalization error is of the order $L\sqrt{\frac{\mathbf{y}^\top(\mathbf{H}^\infty)^{-1}\mathbf{y}}{n}}$, where $\mathbf{H}^\infty$ denotes the NTK gram matrix over the training data and $\lambda_0 = \lambda_{\min}(\mathbf{H}^\infty)$. However, the assumption $\lambda_0 > 0$ is a strong assumption because it tends to zero if the size of the training data tends to infinity as shown by Su and Yang [2019]. Ji and Telgarsky [2020] achieved logarithmic width requirements for NTK-separable data with a margin γ . They derived the risk bound of order $1/(\gamma^2\sqrt{n})$ for two-layer ReLU networks using Rademacher complexity. The recent work Chen et al. [2021] extended their results from shallow to deep neural networks, the authors considered the NTK feature learning and proved the bound of $\tilde{O}\left(\frac{e^{O(L)}}{\gamma^2}\sqrt{\frac{m}{n}}\right)$. However, all the above bounds explicit $\frac{1}{\sqrt{n}}$ dependence on the sample size n , which is suboptimal.

Complementing the NTK framework, another line of research employs algorithmic stability to analyze neural networks. In particular, Liu et al. [2020] demonstrated that the Hessian spectral norm scales with width m as $\tilde{O}\left(\frac{1}{\sqrt{m}}\right)$, providing the theoretical basis for subsequent studies on generalization in overparameterized networks [Richards and Kuzborskij, 2021, Lei et al., 2022, Taheri and Thrampoulidis, 2024, Taheri et al., 2025]. The work [Taheri and Thrampoulidis, 2024] achieves the bound of $\tilde{O}(1/n\gamma^2)$ for shallow neural networks, which is almost optimal, as illustrated by Shamir [2021], Schliserman and Koren [2023]. More recently, Taheri et al. [2025] extended this line of work to deep networks, obtaining a generalization bound of $\tilde{O}(e^{O(L)}/(n\gamma^2))$. However, this approach often assume smooth activation functions and can not apply to the non-smooth ReLU networks. In summary, these works either consider smooth neural networks or develop generalization bounds with exponential dependency on L . These limitations motivate two fundamental questions:

Can we develop optimal risk bounds of $1/(\gamma^2 n)$ for deep ReLU networks through refined Rademacher complexity analysis? Furthermore, is it possible to replace the exponential dependence on L with $\text{poly}(L)$ by using neural networks with width as a polylogarithmic function of n ?

In this paper, we provide affirmative answers to both questions. Our main contributions are listed below.

1. We prove that gradient decent with step size η and T iterations achieves the convergence rate of $F_S(\bar{\mathbf{W}})/(\eta T)$, where $F_S(\bar{\mathbf{W}}) = 3\eta T\mathcal{L}_S(\bar{\mathbf{W}}) + \|\bar{\mathbf{W}} - \mathbf{W}(0)\|_F^2$, $\bar{\mathbf{W}}$ is a reference model, $\mathbf{W}(0)$ is the initialization point and $\mathcal{L}_S(\cdot)$ is the training error. This indicates that all iterates remain within a neighborhood of $\bar{\mathbf{W}}$. By refining the analysis of ReLU activation patterns, we reduce the overparameterization requirement by a factor of L^6 as compared to Chen et al. [2021] (see Remark 1).
2. We establish a population risk bound of $\tilde{O}(L^4 F(\bar{\mathbf{W}})/n)$, where $F(\bar{\mathbf{W}})$ extends the empirical counterpart $F_S(\bar{\mathbf{W}})$ to the population setting. Our analysis introduces two key technical contributions. First, we derive sharper Rademacher complexity bounds for the hypothesis class encompassing all gradient descent iterates. A central innovation is our representation of the complexity via products of sparse matrices, whose norms are tightly controlled using optimization-

informed estimates (see Remark 2). Second, by leveraging the covering number techniques, we prove that ReLU networks remain $\tilde{O}(L^2)$ -Lipschitz continuous in a neighborhood around the initialization—a substantial improvement over previous exponential bounds [Xu and Zhu, 2024, Taheri et al., 2025] (see Remark 3).

3. For NTK separable data with a margin γ , we show that neural networks with $m = \tilde{O}(L^{16} \log(n/\delta)(\log n)^8/\gamma^8)$ can achieve $\tilde{O}(L^4(1 + \gamma L^2)/(\gamma^2 n))$ risk. This improves on Chen et al. [2021]’s $\tilde{O}(e^{O(L)}\gamma^{-2}\sqrt{m/n})$ by simultaneously (a) removing the exponential depth dependence, (b) eliminating the $\sqrt{m}$ width factor and (c) achieving the optimal dependence on sample complexity (see Table 1 for a comparison with existing work).

2 Related Works

2.1 Optimization

The foundational work of Jacot et al. [2018] introduced the Neural Tangent Kernel (NTK) framework, demonstrating that in the infinite-width limit, neural networks behave as linear models with a fixed tangent kernel [Liu et al., 2020, Lee et al., 2019]. This lazy training regime [Chizat et al., 2019], where parameters remain close to initialization, enables gradient descent to converge to global optima near initialization [Du et al., 2019a, Arora et al., 2019a]. These analyses showed that the training dynamics can be governed by the NTK Gram matrix, which leads to substantial overparameterization ($m \gtrsim n^6/\lambda_0^4$). This was later improved by Oymak and Soltanolkotabi [2020]. They showed that if the square-root of the number of the network parameters exceeds the size of the training data, randomly initialized gradient descent converges at a geometric rate to a nearby global optima. The work (Ji and Telgarsky [2020]) achieved polylogarithmic width requirements for logistic loss by leveraging the 1-homogeneity of two-layer ReLU networks. However, it should be mentioned that this special property does not hold for deep networks. The NTK framework was extended to deep architectures by Arora et al. [2019b] for CNNs and by Du et al. [2019b] for ResNets using the last-layer NTK. Xu and Zhu [2024] pointed out that such a characterization is loose, only capturing the contribution from the last layer. They further gave the uniform convergence of all layers as $m \rightarrow \infty$ and convergence guarantee for stochastic gradient descent (SGD) in streaming data setting. Allen-Zhu et al. [2019] showed that the optimization landscape is almost-convex and semi-smooth, based on which they proved that SGD can find global minima. Cao and Gu [2019] introduced the neural tangent random feature and showed the convergence of SGD under the overparameterized assumption $m \gtrsim n^7$.

2.2 Generalization

The NTK framework has yielded generalization bounds scaling as $\sqrt{\mathbf{y}^\top (\mathbf{H}^\infty)^{-1} \mathbf{y}/n}$ [Arora et al., 2019a, Cao and Gu, 2019]. This data-dependent complexity measure helps to distinguish between random labels and true labels. Li and Liang [2018] showed that SGD trained networks can achieve small test error on specific structured data. A very popular approach to studying the generalization of neural networks is via the uniform convergence, which analyzes generalization gaps in a hypothesis space using tools such as Rademacher complexity or covering numbers [Neyshabur et al., 2015, Bartlett et al., 2017, Golowich et al., 2018, Liu et al., 2024]. However, this could lead to vacuous generalization bound in some cases [Nagarajan and Kolter, 2019]. Moreover, these bounds typically exhibit exponential dependence on depth L , thus often leading to loose bounds [Chen et al., 2021]. This capacity-based method usually results in the generalization rate of the order $\tilde{O}(1/\sqrt{n})$. Recent work has also exploited stability arguments for generalization guarantees [Richards and Kuzborskij, 2021, Lei et al., 2022, Taheri and Thrampoulidis, 2024, Deora et al., 2023, Taheri et al., 2025]. The main idea of algorithmic stability is to study how the perturbation of training samples would affect the output of an algorithm [Rogers and Wagner, 1978]. The connection to generalization bound was established in Bousquet and Elisseeff [2002]. Hardt et al. [2016] gave the stability analysis of SGD for Lipschitz, smooth and convex problems. Lei and Ying [2020] further studied SGD under much wilder assumptions. Liu et al. [2020] identified weak convexity of neural networks, enabling stability analyses with polynomial width requirements for quadratic loss [Richards and Kuzborskij, 2021, Lei et al., 2022]. Moreover, Taheri and Thrampoulidis [2024], Taheri et al. [2025] obtained generalization bounds of order $\tilde{O}(1/n)$ by using a generalized local quasi-convexity property for

sufficiently parameterized networks. However, these methods depend on smooth activations, and whether similar or even better bound can be established for deep ReLU networks is still unknown.

3 Preliminaries

Notation Throughout the paper, we denote $a \lesssim b$ if there exists a constant $c > 0$ such that $a \leq cb$, and denote $a \asymp b$ if both $a \lesssim b$ and $b \lesssim a$ hold. We use the standard notation $O(\cdot), \Omega(\cdot)$ and use $\tilde{O}(\cdot), \tilde{\Omega}(\cdot)$ to hide polylogarithmic factors. Denote by $\mathbb{I}\{\cdot\}$ the indicator function (i.e., taking the value 1 if the argument holds true, and 0 otherwise). We use $\mathcal{N}(\mu, \sigma^2)$ to denote the Gaussian distribution of mean μ and variance σ^2 . For a positive integer n , we denote $[n] := \{1, \dots, n\}$. For a vector $x \in \mathbb{R}^d$, we use $\|x\|_2$ to denote its Euclidean norm. For a matrix $A \in \mathbb{R}^{m \times n}$, we denote $\|A\|_2$ and $\|A\|_F$ the corresponding spectral norm and Frobenius norm respectively. The $(2, 1)$ -norm of A is defined as $\|A\|_{2,1} = \sum_{j=1}^n \|A_{:,j}\|_2$. Let $\langle \cdot, \cdot \rangle$ be the inner product of a vector or a matrix, i.e., for any matrices $A, B \in \mathbb{R}^{m \times n}$, we have $\|A\|_F^2 = \text{tr}(A^\top A)$ and $\langle A, B \rangle = \text{tr}(A^\top B)$. Let $L \in \mathbb{N}$, $\mathbf{A} = (A_1, \dots, A_L)$ and $\mathbf{B} = (B_1, \dots, B_L)$ be two collections of arbitrary matrices such that A_i and B_i have the same size for all $i \in [L]$. We define $\langle \mathbf{A}, \mathbf{B} \rangle = \sum_{i=1}^L \text{tr}(A_i^\top B_i)$. For a matrix $\mathbf{W}$, we define $(\mathbf{w}_r)^\top$ the r -th row of $\mathbf{W}$. Denote $\|\cdot\|_0$ the l^0 -norm which is the number of nonzero entries of a matrix or a vector. We denote $C \geq 1$ as an absolute value, which may differ from line to line.

Let $\mathcal{X} \subseteq \mathbb{R}^d$ be the input space, $\mathcal{Y} = \{1, -1\}$ be the output space, and $\mathcal{Z} = \mathcal{X} \times \mathcal{Y}$. Let ρ be a probability measure defined on $\mathcal{Z}$. Let $S = \{z_i = (\mathbf{x}_i, y_i) : i = 1, \dots, n\}$ be a training dataset drawn from ρ . Let $\mathcal{W} := \mathbb{R}^{m \times d} \times (\mathbb{R}^{m \times m})^{L-1}$ be the parameter space. $\mathbf{W}^1 \in \mathbb{R}^{m \times d}$ and $\mathbf{W}^l \in \mathbb{R}^{m \times m}$ for $l = 2, \dots, L$ is the weight of the l -th hidden layer. $\mathbf{W} = (\mathbf{W}^1, \dots, \mathbf{W}^L) \in \mathcal{W}$ denotes the collection of weight matrices for all layers. Let $\mathbf{a} = (a_1, \dots, a_m)^\top \in \mathbb{R}^m$ be the weight vector of the output layer and $\sigma(\cdot) = \max\{\cdot, 0\}$ denote the ReLU activation function. For $\mathbf{x} \in \mathcal{X}$, we consider the L -layer deep ReLU neural networks with width m as follows,

$$f_{\mathbf{W}}(\mathbf{x}) = \mathbf{a}^\top \sqrt{\frac{2}{m}} \sigma \left(\mathbf{W}^L \dots \sqrt{\frac{2}{m}} \sigma(\mathbf{W}^1 \mathbf{x}) \right). \quad (1)$$

Given an input $\mathbf{x} \in \mathcal{X}$ and parameter matrix $\mathbf{W} = (\mathbf{W}^1, \dots, \mathbf{W}^L)$ of an L -layer ReLU network $f_{\mathbf{W}}(\mathbf{x})$. We denote the output of the l -th layer by $h^l(\mathbf{x}) = \sqrt{\frac{2}{m}} \sigma(\mathbf{W}^l h^{l-1}(\mathbf{x}))$ with $h^0(\mathbf{x}) = \mathbf{x}$. Then $f_{\mathbf{W}}(\mathbf{x}) = \mathbf{a}^\top h^L(\mathbf{x})$. We define $\mathcal{B}_R(\mathbf{W}) = \{\tilde{\mathbf{W}} \in \mathcal{W} : \|\mathbf{W}^l - \tilde{\mathbf{W}}^l\|_F \leq R, l \in [L]\}$. The performance of the network $f_{\mathbf{W}}(\mathbf{x})$ is measured by the following empirical risk $\mathcal{L}_S(\mathbf{W})$ and population risk $\mathcal{L}(\mathbf{W})$, respectively:

$$\mathcal{L}_S(\mathbf{W}) = \frac{1}{n} \sum_{i=1}^n \ell(y_i f_{\mathbf{W}}(\mathbf{x}_i)) \quad \text{and} \quad \mathcal{L}(\mathbf{W}) = \mathbb{E}_z \ell(y f_{\mathbf{W}}(\mathbf{x})),$$

where we use logistic loss $\ell(z) := \log(1 + \exp(-z))$ throughout this paper. We further assume the following symmetric initialization [Nitanda and Suzuki, 2020, Kuzborskij and Szepesvári, 2023, Xu and Zhu, 2024]:

Assumption 1 (Symmetric initialization). *Without loss of generality, we assume the network width m is even, and $a_{r+\frac{m}{2}} = -a_r \in \{-1, +1\}$ for $1 \leq r \leq m/2$. $\mathbf{W}(0) \in \mathcal{W}$ satisfies*

$$\begin{aligned} \mathbf{w}_r^1(0) &\sim \mathcal{N}(0, \mathbf{I}_d), \mathbf{w}_r^l(0) \sim \mathcal{N}(0, \mathbf{I}_m) \quad 2 \leq l \leq L-1 \text{ and } r \in [m], \\ \mathbf{w}_r^L(0) &\sim \mathcal{N}(0, \mathbf{I}_m) \text{ for } r = \{1, \dots, \frac{m}{2}\}, \text{ and } \mathbf{w}_{r+\frac{m}{2}}^L(0) = \mathbf{w}_r^L(0). \end{aligned} \quad (2)$$

We remark that this initialization is for theoretical simplicity, using general initialization techniques will not affect the main results. We fix the output weights $\{a_r\}$ and use Gradient Descent (GD) to train the weight matrix $\mathbf{W}$ [Ji and Telgarsky, 2020, Arora et al., 2019a, Zou et al., 2018].

Definition 1 (Gradient Descent). *GD updates $\{\mathbf{W}(k)\}$ by*

$$\mathbf{W}^l(k+1) = \mathbf{W}^l(k) - \eta \frac{\partial \mathcal{L}_S(\mathbf{W}(k))}{\partial \mathbf{W}^l(k)} \text{ for all } l \in [L], \quad (3)$$

where $\eta > 0$ is the step size.

Note that in each layer we employ $\sqrt{2/m}$ as the regularization factor instead of the conventional $\sqrt{1/m}$ [Ji and Telgarsky, 2020], which is due to $\mathbb{E}_{x \sim \mathcal{N}(0,1)} \sigma^2(x) = 1/2$ for our activation function $\sigma(\cdot)$. This scaling matches both the theoretical framework of Du et al. [2019a] and the initialization scheme of [He et al., 2015] (where weights $\mathbf{w}_r^l \sim \mathcal{N}(0, 2/m)$). As will be shown later (Appendix A), this regularization ensures stable gradient propagation and maintains consistent variance across layers.

The following assumption is standard in the literature [Cao and Gu, 2019, Ji and Telgarsky, 2020, Chen et al., 2021].

Assumption 2. We assume $\mathcal{X} = S^{d-1}$ be the sphere.

Throughout the paper, we assume that Assumptions 1 and 2 always hold true.

Error decomposition In this work, we analyze the performance of gradient descent through the lens of population risk. To facilitate this analysis, we decompose the population risk $\mathcal{L}(\mathbf{W})$ as follows

$$\mathcal{L}(\mathbf{W}) = (\mathcal{L}(\mathbf{W}) - \mathcal{L}_S(\mathbf{W})) + \mathcal{L}_S(\mathbf{W}),$$

where the first term captures the generalization error, quantifying the network’s performance on unseen data. The second term represents the optimization error, which reflects GD’s ability to find global minima. We will use tools in the optimization theory to study the optimization error [Ji and Telgarsky, 2020, Schliserman and Koren, 2022], and Rademacher complexity to control the generalization error [Mohri, 2018].

4 Main Results

In this section, we present the main results. In Section 4.1, we show the optimization analysis of gradient descent. In Section 4.2, we use Rademacher complexity to control the generalization error. In Section 4.3, we apply our generalization results to NTK-separable data with a margin γ .

4.1 Optimization Analysis

We introduce the following notations for a reference model $\bar{\mathbf{W}}$

$$F_S(\bar{\mathbf{W}}) := 3\eta T \mathcal{L}_S(\bar{\mathbf{W}}) + \|\mathbf{W}(0) - \bar{\mathbf{W}}\|_F^2, \quad \tilde{F}_S(\bar{\mathbf{W}}) = \frac{1}{n} \sum_{i=1}^n |\ell'(y_i f_{\bar{\mathbf{W}}}(\mathbf{x}_i))|.$$

Without loss of generality, we assume $F_S(\bar{\mathbf{W}}) \geq 1$.

Theorem 1. Let Assumptions 1, 2 hold. If $m \gtrsim L^{16}(\log m)^4 \log(nL/\delta) F_S^4(\bar{\mathbf{W}})$, $\eta \leq \min\{4/(5L), 1/(20L\tilde{F}_S(\bar{\mathbf{W}}))\}$, then with probability at least $1 - \delta$, for all $t \leq T$ we have

$$\|\mathbf{W}(t) - \bar{\mathbf{W}}\|_F^2 \leq F_S(\bar{\mathbf{W}}) \quad \text{and} \quad \eta \sum_{k=0}^{t-1} \mathcal{L}_S(\mathbf{W}(k)) \leq F_S(\bar{\mathbf{W}}).$$

Remark 1. Our theorem shows that the convergence rate is bounded by the optimization error of a reference model, implying that any low-loss reference point guarantees good convergence. While prior works relied on NTK-induced solutions [Richards and Kuzborskij, 2021, Arora et al., 2019a], we prove that there exists a reference model near initialization under the milder Assumption 3. Furthermore, our analysis implies that all training iterates remain within a neighborhood of the reference point, and thus near initialization, aligned with NTK-based observations but without studying the kernel or the corresponding Gram matrix [Du et al., 2019a,b].

Here we provide the proof sketch and compare it with previous works. The starting point is the following bound given in Lemma 19

$$f_{\mathbf{W}}(\mathbf{x}_i) - f_{\bar{\mathbf{W}}}(\mathbf{x}_i) - \left\langle \frac{\partial f_{\mathbf{W}}(\mathbf{x}_i)}{\partial \mathbf{W}}, \mathbf{W} - \bar{\mathbf{W}} \right\rangle = \tilde{O}\left(\frac{L^{8/3} R^{4/3}}{m^{1/6}}\right) \quad (4)$$

for $\mathbf{W}, \bar{\mathbf{W}} \in \mathcal{B}_R(\mathbf{W}(0))$.

Then we can show the following inequality holds (Lemma 21),

$$\|\mathbf{W}(t+1) - \bar{\mathbf{W}}\|_F^2 \leq \|\mathbf{W}(t) - \bar{\mathbf{W}}\|_F^2 - \eta \mathcal{L}_S(\mathbf{W}(t)) + 3\eta \mathcal{L}_S(\bar{\mathbf{W}}). \quad (5)$$

Telescoping gives the theorem. Chen et al. [2021] introduce the following neural tangent random feature (NTRF) function class:

$$\mathcal{F}(\mathbf{W}(0), R) = \left\{ F_{\mathbf{W}(0), \mathbf{W}}(\mathbf{x}) = f_{\mathbf{W}(0)}(\mathbf{x}) + \left\langle \frac{\partial f_{\mathbf{W}(0)}(\mathbf{x})}{\partial \mathbf{W}(0)}, \mathbf{W} - \mathbf{W}(0) \right\rangle : \mathbf{W} \in \mathcal{B}_R(\mathbf{W}(0)) \right\}.$$

They show that gradient descent achieves a training loss of at most $3\epsilon_{\text{NTRF}}$, where ϵ_{NTRF} denotes the minimal loss over the NTRF function class (see Theorem 3.3 therein). In contrast, our approach directly analyzes the GD iterates and shows that the existence of a nearby reference point with small training error is sufficient to ensure convergence—extending the analysis beyond the NTK regime. For a fair comparison, under Assumption 3, both analyses yield an optimization error of $\tilde{O}(1/T)$. However, our method significantly relaxes the overparameterization requirement, improving the width dependence by a factor of L^6 (see Remark 5).

4.2 Generalization Analysis

We use Rademacher complexity to study the generalization error, which measures the ability of a function class to correlate random noises.

Definition 2 (Rademacher complexity). *Let $\mathcal{F}$ be a class of real-valued functions over a space $\mathcal{X}$, $S_1 = \{\mathbf{x}_1, \dots, \mathbf{x}_n\} \subset \mathcal{X}$. We define the following empirical Rademacher complexity as*

$$\mathfrak{R}_{S_1}(\mathcal{F}) = \mathbb{E}_\epsilon \left[\sup_{f \in \mathcal{F}} \frac{1}{n} \sum_{i \in [n]} \epsilon_i f(\mathbf{x}_i) \right],$$

where $\epsilon = (\epsilon_i)_{i \in [n]} \sim \{\pm 1\}^n$ are independent Rademacher variables, i.e., taking values in $\{\pm 1\}$ with the same probability.

We further define the following worst-case Rademacher complexity,

$$\mathfrak{R}_{S_1, n}(\mathcal{F}) = \sup_{\tilde{S} \subset S_1: |\tilde{S}|=n} \mathfrak{R}_{\tilde{S}}(\mathcal{F}).$$

We define $G = \sup_z \ell(y f_{\bar{\mathbf{W}}}(\mathbf{x}))$, and

$$F(\bar{\mathbf{W}}) = 3\eta T \left(2\mathcal{L}(\bar{\mathbf{W}}) + \frac{7G \log(2/\delta)}{6n} \right) + \|\mathbf{W}(0) - \bar{\mathbf{W}}\|_F^2. \quad (6)$$

We consider the following function space

$$\mathcal{F} := \{\mathbf{x} \rightarrow f_{\mathbf{W}}(\mathbf{x}) : \mathbf{W} \in \mathcal{W}_1\}, \quad (7)$$

where the parameter space is defined as

$$\mathcal{W}_1 = \{\mathbf{W} \in \mathcal{W} : \|\mathbf{W} - \bar{\mathbf{W}}\|_F^2 \leq F(\bar{\mathbf{W}})\}. \quad (8)$$

As will be shown in Lemma 22, with high probability there holds

$$\mathfrak{R}_{S_1, n}(\mathcal{F}) = \tilde{O} \left(L^2 \sqrt{\frac{F(\bar{\mathbf{W}})}{n}} \right). \quad (9)$$

Remark 2 (Improved Rademacher complexity). In previous work [Chen et al., 2021], they derived the bound of $\tilde{O}(4^L L \sqrt{m F(\bar{\mathbf{W}})/n})$. Specifically, they use the generalization analysis in Bartlett et al. [2017], which requires to estimate the following term

$$\left[\prod_{l=1}^L \|\mathbf{W}^l\|_2 \right] \cdot \left[\sum_{l=1}^L \frac{\|(\mathbf{W}^l)^\top - (\bar{\mathbf{W}}^l)^\top\|_{2,1}^{2/3}}{\|\mathbf{W}^l\|_2^{2/3}} \right]^{3/2}.$$

Note that $\prod_{l=1}^L \|\mathbf{W}^l\|_2$ could lead to exponential dependence on the depth. Moreover, $\|(\mathbf{W}^l)^\top - (\bar{\mathbf{W}}^l)^\top\|_{2,1} \leq \sqrt{m}\|\mathbf{W}^l - \bar{\mathbf{W}}^l\|_F$, inducing an explicit $\sqrt{m}$ term. However, we reduce the dependence of L to polynomial (L^2). Furthermore, we sharpen the dependence on width from $\sqrt{m}$ to logarithmic terms. The key idea is to introduce the following expression

$$f_{\mathbf{W}}(\mathbf{x}_i) - f_{\mathbf{W}(0)}(\mathbf{x}_i) = \mathbf{a}^\top \sum_{l=1}^L \hat{\mathbf{G}}_{L,0}^l(\mathbf{x}_i)(\mathbf{W}^l - \mathbf{W}^l(0))h_0^{l-1}(\mathbf{x}_i),$$

where $\hat{\mathbf{G}}_{L,0}^l(\mathbf{x}_i)$ is a matrix defined in (36). We further show that $\hat{\mathbf{G}}_{L,0}^l(\mathbf{x}_i)$ is of the order $\tilde{O}(L/\sqrt{m})$ with high probability, from which we can derive $\mathfrak{R}_{S_1,n}(\mathcal{F}) = \tilde{O}(L^2/\sqrt{n})$. We will show that this improved Rademacher complexity is crucial to get almost optimal risk bounds in NTK separable data.

The following lemma shows that generalization bounds of smooth loss functions can be controlled by Rademacher complexity.

Lemma 1 (Srebro et al. [2010]). *Let $G' = \sup_{z, \mathbf{W} \in \mathcal{W}_1} \ell(yf_{\mathbf{W}}(\mathbf{x}))$. For any $0 < \delta < 1$, we have with probability at least $1 - \delta/2$ over S , for any $\mathbf{W} \in \mathcal{W}_1$,*

$$\begin{aligned} \mathcal{L}(\mathbf{W}) - \mathcal{L}_S(\mathbf{W}) &\lesssim \mathcal{L}_S^{1/2}(\mathbf{W}) \left(\frac{1}{2}(\log n)^{3/2} \mathfrak{R}_{S_1,n}(\mathcal{F}) + \left(\frac{G' \log(2/\delta)}{n} \right)^{1/2} \right) \\ &\quad + \frac{1}{4}(\log n)^3 \mathfrak{R}_{S_1,n}^2(\mathcal{F}) + \frac{G' \log(2/\delta)}{n}. \end{aligned}$$

Here we choose $G' = 2G + CL^4 \log m F(\bar{\mathbf{W}})$. Then the excess risk error is established as follows.

Theorem 2. *Let Assumptions 1, 2 hold. If $m \gtrsim L^{16} d(\log m)^5 \log(nL/\delta) F_S^4(\bar{\mathbf{W}})$, $\eta \leq \min\{4/(5L), 1/(20L\tilde{F}_S(\bar{\mathbf{W}}))\}$, $\eta T \asymp n$, then with probability at least $1 - \delta$, we have*

$$\frac{1}{T} \sum_{t=0}^{T-1} \mathcal{L}(\mathbf{W}(t)) = \tilde{O} \left(\frac{L^4 F(\bar{\mathbf{W}}) + G \log(2/\delta)}{n} \right).$$

Remark 3 (Analysis of Lipschitzness). To bound the term $G' = \sup_{z, \mathbf{W} \in \mathcal{W}_1} \ell(yf_{\mathbf{W}}(\mathbf{x}))$, we analyze the difference $f_{\mathbf{W}}(\mathbf{x}) - f_{\bar{\mathbf{W}}}(\mathbf{x})$. Since both $\mathbf{W}, \bar{\mathbf{W}} \in \mathcal{B}_R(\mathbf{W}(0))$ for some R , we only need to study the local variation $f_{\mathbf{W}}(\mathbf{x}) - f_{\mathbf{W}(0)}(\mathbf{x})$. This approach necessitates characterizing the uniform behavior of deep networks in $\mathcal{B}_R(\mathbf{W}(0))$, specifically establishing control over their Lipschitz constants near initialization. Existing works usually lead to an exponential dependence on L [Xu and Zhu, 2024, Taheri et al., 2025], thus resulting in a $e^{O(L)}$ term in the generalization bound. In particular, Lemma F.3 and F.5 in Liu et al. [2020] pointed out that $\|h^l(\mathbf{x})\| \leq C^L$, $\|\partial f_{\mathbf{W}}(\mathbf{x})/\partial h^l(\mathbf{x})\|_2 \leq C^{L-l+1}\sqrt{m}$. Based on these observations, Taheri et al. [2025] showed that

$$\left\| \frac{\partial f_{\mathbf{W}}(\mathbf{x})}{\partial \mathbf{W}^l} \right\|_2 \leq C^L.$$

On the other hand, to analyze the output difference near initialization, we observe that

$$f_{\mathbf{W}}(\mathbf{x}) - f_{\mathbf{W}(0)}(\mathbf{x}) = \mathbf{a}^\top (h^L(\mathbf{x}) - h_0^L(\mathbf{x})) \leq \sqrt{m} \|h^L(\mathbf{x}) - h_0^L(\mathbf{x})\|_2,$$

reducing our task to bounding the hidden layer perturbation. Previous approaches, including Xu and Zhu [2024] and Du et al. [2019b], employ a recursive estimation:

$$\begin{aligned} \|h^L(\mathbf{x}) - h_0^L(\mathbf{x})\|_2 &= \sqrt{\frac{2}{m}} \|\sigma(\mathbf{W}^L h^{L-1}(\mathbf{x})) - \sigma(\mathbf{W}^L(0) h_0^{L-1}(\mathbf{x}))\|_2 \\ &\leq \sqrt{\frac{2}{m}} (\|(\mathbf{W}^L - \mathbf{W}^L(0)) h^{L-1}(\mathbf{x})\|_2 + \|\mathbf{W}^L(0) (h^{L-1}(\mathbf{x}) - h_0^{L-1}(\mathbf{x}))\|_2) \\ &\lesssim \frac{R}{\sqrt{m}} (\|h^{L-1}(\mathbf{x}) - h_0^{L-1}(\mathbf{x})\|_2 + C^L) + \|h^{L-1}(\mathbf{x}) - h_0^{L-1}(\mathbf{x})\|_2 \leq \frac{C^L R}{\sqrt{m}}, \end{aligned}$$

where in the second inequality they used $\|h_0^l(\mathbf{x})\|_2 \leq C^L$ and $\|\mathbf{W}^l(0)\|_2 \lesssim \sqrt{m}$. Although this method provides a straightforward bound, it leads to an exponential dependence on depth L due to the recursive nature of the estimation.

In contrast to previous work, we develop the covering-number strategy to avoid the exponential dependence on depth. Specifically, we first show that for n training data, if $m = \tilde{\Omega}(L^{10} \log(n) R^2)$, then $\|h^l(\mathbf{x}_i) - h_0^l(\mathbf{x}_i)\|_2 = \tilde{O}\left(\frac{L^2 R}{\sqrt{m}}\right)$ holds for $i \in [n], l \in [L]$ (Lemma 15). We further take a $1/(C^L \sqrt{m})$ -covering $D = \{\mathbf{x}^j : j = 1, \dots, |D|\}$ of the input space. Recall that the input space $\mathcal{X} = S^{d-1}$, it is well known from Corollary 4.2.13 in Vershynin [2018] that the number of $1/(C^L \sqrt{m})$ -covering is given by $|D| \leq (1 + 2C^L \sqrt{m})^d$. Applying Lemma 15 to D derives that if $m = \tilde{\Omega}(L^{10} \log(|D|) R^2)$

$$\|h^l(\mathbf{x}^j) - h_0^l(\mathbf{x}^j)\|_2 = \tilde{O}\left(\frac{L^2 R}{\sqrt{m}}\right), \quad \mathbf{x}^j \in D, l \in [L].$$

Note that although the covering number could be exponential in L , we only require logarithm of it, thus leading to polynomial dependence! For any input $\mathbf{x}$, we use the closest cover point $\mathbf{x}^j \in D$ to approximate $\|h^l(\mathbf{x}) - h^l(\mathbf{x}^j)\|_2, \|h_0^l(\mathbf{x}) - h_0^l(\mathbf{x}^j)\|_2$. Combining these yields the key technical lemma (Lemma 16):

$$\sup_{\mathbf{x} \in \mathcal{X}} \|h^l(\mathbf{x}) - h_0^l(\mathbf{x})\|_2 = \tilde{O}\left(\frac{L^2 R}{\sqrt{m}}\right).$$

This directly implies that the network is $\tilde{O}(L^2)$ -Lipschitz near initialization. More details can be found in Lemma 16 and its proof.

From Theorem 1, we can prove that all iterates fall into $\mathcal{W}_1$ with high probability. Hence we can choose $R \asymp \sqrt{F(\overline{\mathbf{W}})}$. Combining the above results and the smoothness of ℓ yields

$$\ell(yf_{\mathbf{W}}(\mathbf{x})) - \ell(yf_{\overline{\mathbf{W}}}(\mathbf{x})) \lesssim L^4 \log m F(\overline{\mathbf{W}}),$$

which motivates our choice of G' . It then follows that

$$\begin{aligned} \mathcal{L}(\mathbf{W}(t)) - 2\mathcal{L}_S(\mathbf{W}(t)) &\lesssim (\log n)^3 \mathfrak{R}_{S,n}^2(\mathcal{F}) + \frac{G' \log(2/\delta)}{n} \\ &= \tilde{O}\left(\frac{L^4 F(\overline{\mathbf{W}}) + G \log(2/\delta)}{n}\right). \end{aligned}$$

Taking summarization over t and applying Theorem 1 complete the proof of Theorem 2. The full proofs are provided in Appendix C.

4.3 Optimal rates on NTK-separable data

In this section, we apply our general analysis to NTK-separable data [Ji and Telgarsky, 2020, Nitanda et al., 2020, Chen et al., 2021, Taheri and Thrampoulidis, 2024, Deora et al., 2023], and obtain the optimal rates.

Assumption 3. *There exists $\gamma > 0$ and a collection of matrices $\mathbf{W}_* = \{\mathbf{W}_*^1, \dots, \mathbf{W}_*^L\}$ satisfying $\sum_{l=1}^L \|\mathbf{W}_*^l\|_F^2 = 1$, such that*

$$y_i \left\langle \mathbf{W}_*, \frac{\partial f_{\mathbf{W}(0)}(\mathbf{x}_i)}{\partial \mathbf{W}(0)} \right\rangle \geq \gamma, \quad i \in [n].$$

This means that the dataset is separable by the NTK feature at initialization with a margin γ . Nitanda et al. [2020] pointed out that this assumption is weaker than positive eigenvalues of NTK Gram matrix, which has been widely used in the literature [Du et al., 2019b,a, Arora et al., 2019a]. With the above assumption, we have the following optimal risk bound on NTK separable data. The proof is given in Appendix D.

Theorem 3. *Let Assumptions 1, 2, 3 hold. If $m \gtrsim L^{16} d (\log m)^5 \log(nL/\delta) (\log T)^8 / \gamma^8, \eta \leq 4/(5L), \eta T \asymp n$, then with probability at least $1 - \delta$, we have*

$$\frac{1}{T} \sum_{t=0}^{T-1} \mathcal{L}(\mathbf{W}(t)) = \tilde{O}\left(\frac{L^4 (1 + \gamma L^2) (\log T)^2}{n \gamma^2}\right).$$

Remark 4 (Proof sketch). To apply the result in Theorem 2, we need to estimate $F(\bar{\mathbf{W}})$, for which it suffices to bound $\mathcal{L}(\bar{\mathbf{W}})$ and $G = \sup_{\mathbf{x}} \ell(y f_{\bar{\mathbf{W}}}(\mathbf{x}))$. For the first part, we control it by $\mathcal{L}_S(\bar{\mathbf{W}})$ using Bernstein inequality (Eq.(14)). Let $\bar{\mathbf{W}} = \mathbf{W}(0) + 2 \log T \mathbf{W}_* / \gamma$, plugging into (4) obtains $\ell(y_i f_{\bar{\mathbf{W}}}(\mathbf{x}_i)) \leq 1/T$ and further $\mathcal{L}_S(\bar{\mathbf{W}}) \leq 1/T$, implying $F_S(\bar{\mathbf{W}}) = \tilde{O}(1/\gamma^2)$. In order to control $\ell(y f_{\bar{\mathbf{W}}}(\mathbf{x}))$, we leverage the $\tilde{O}(L^2)$ -Lipschitzness of $f_{\mathbf{W}}(\mathbf{x})$. Indeed, for any $\mathbf{x} \in \mathcal{X}$, there holds

$$|f_{\bar{\mathbf{W}}}(\mathbf{x})| \leq |f_{\mathbf{W}(0)}(\mathbf{x})| + |f_{\bar{\mathbf{W}}}(\mathbf{x}) - f_{\mathbf{W}(0)}(\mathbf{x})| = \tilde{O}\left(\frac{L^2}{\gamma}\right).$$

It then follows that $G = \tilde{O}\left(\frac{L^2}{\gamma}\right)$ and $F(\bar{\mathbf{W}}) = \tilde{O}\left(\frac{(\log T)^2(1+\gamma L^2)}{\gamma^2}\right)$.

Remark 5 (Discussion on optimization error). Under Assumption 3, Theorem 3.3 and Proposition 4.2 of Chen et al. [2021] show that when the network width satisfies $m = \tilde{\Omega}(L^{22}/\gamma^8)$, the training error is of the order $\tilde{O}(1/T)$. We achieve the same guarantee under a significantly milder width condition of $m = \tilde{\Omega}(L^{16}/\gamma^8)$. This improvement is enabled by two key technical advances: a sharper bound for (4), and a tighter estimate of the iterate distance $\|\mathbf{W}(t) - \bar{\mathbf{W}}\|_F$. Specifically, we improve the bound in (4) by a factor of $L^{1/3}$, and show that $\|\mathbf{W}(t) - \bar{\mathbf{W}}\|_F = \tilde{O}(1/\gamma)$, improving upon the previous $\tilde{O}(\sqrt{L}/\gamma)$ bound. Together, these refinements reduce the required network width by a factor of L^6 .

Remark 6 (Comparison). Under similar assumptions, Ji and Telgarsky [2020] derived the bound $\tilde{O}\left(\frac{1}{\gamma^2 \sqrt{n}}\right)$ for shallow networks. For deep ReLU networks, Chen et al. [2021] developed the bound of the order $\tilde{O}\left(\frac{e^{O(L)}}{\gamma^2} \sqrt{\frac{m}{n}}\right)$ via Rademacher complexity [Bartlett et al., 2017], suffering from exponential depth dependence, explicit width requirement and suboptimal $\sqrt{1/n}$ scaling. Taheri et al. [2025] improved the result to $\tilde{O}\left(\frac{e^{O(L)}}{\gamma^2 n}\right)$ for deep networks. The dependence on n, γ is optimal up to a logarithmic factor [Shamir, 2021, Schliserman and Koren, 2023]. However, their results require smooth activations and exponential width in L . Our rate is almost-optimal and enjoys polynomial dependence over the network depth. Furthermore, our bound holds under the overparameterization $\tilde{\Omega}(1/\gamma^8)$, matching the requirement in Ji and Telgarsky [2020], Chen et al. [2021]. This is much better than $1/\gamma^{6L+4}$ in Taheri et al. [2025]. To the best of our knowledge, these are the best generalization bound and width condition for deep neural networks.

5 Conclusion and Future Work

In this paper, we present optimization and generalization analysis of gradient descent-trained deep ReLU networks for classification tasks. We explore the optimization error of $F_S(\bar{\mathbf{W}})/(\eta T)$ under a milder overparameterization requirement than before. We establish sharper bound of Rademacher complexity and Lipschitz constant for neural networks. This helps to derive generalization bound of order $\tilde{O}(F(\bar{\mathbf{W}})/n)$. For NTK-separable data with a margin γ , our methods lead to the optimal rate of $\tilde{O}(1/(n\gamma^2))$. We improve the existing analysis and require less overparameterization than previous works.

There remain several interesting questions for future works. First, it is an interesting question to extend our methods to SGD. Second, while we establish polynomial Lipschitz constants near initialization, investigating whether similar bounds hold far from initialization would deepen our theoretical understanding. Finally, we only consider fully-connected neural networks. It is interesting to study the generalization analysis of networks with other architectures, such as CNNs and Resnets [Du et al., 2019b].

References

Zeyuan Allen-Zhu, Yuanzhi Li, and Zhao Song. A convergence theory for deep learning via over-parameterization. In Kamalika Chaudhuri and Ruslan Salakhutdinov, editors, *Proceedings of the 36th International Conference on Machine Learning*, volume 97 of *Proceedings of Machine Learning Research*, pages 242–252. PMLR, 09–15 Jun 2019. URL <https://proceedings.mlr.press/v97/allen-zhu19a.html>.

- Sanjeev Arora, Simon Du, Wei Hu, Zhiyuan Li, and Ruosong Wang. Fine-grained analysis of optimization and generalization for overparameterized two-layer neural networks. In Kamalika Chaudhuri and Ruslan Salakhutdinov, editors, *Proceedings of the 36th International Conference on Machine Learning*, volume 97 of *Proceedings of Machine Learning Research*, pages 322–332. PMLR, 09–15 Jun 2019a. URL <https://proceedings.mlr.press/v97/arora19a.html>.
- Sanjeev Arora, Simon S Du, Wei Hu, Zhiyuan Li, Russ R Salakhutdinov, and Ruosong Wang. On exact computation with an infinitely wide neural net. *Advances in neural information processing systems*, 32, 2019b.
- Peter L Bartlett, Dylan J Foster, and Matus J Telgarsky. Spectrally-normalized margin bounds for neural networks. *Advances in neural information processing systems*, 30, 2017.
- Olivier Bousquet and André Elisseeff. Stability and generalization. *Journal of machine learning research*, 2 (Mar):499–526, 2002.
- Yuan Cao and Quanquan Gu. Generalization bounds of stochastic gradient descent for wide and deep neural networks. In H. Wallach, H. Larochelle, A. Beygelzimer, F. d’Alché-Buc, E. Fox, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 32. Curran Associates, Inc., 2019. URL https://proceedings.neurips.cc/paper_files/paper/2019/file/cf9dc5e4e194fc21f397b4cac9cc3ae9-Paper.pdf.
- Zixiang Chen, Yuan Cao, Difan Zou, and Quanquan Gu. How much over-parameterization is sufficient to learn deep relu networks?, 2021. URL <https://arxiv.org/abs/1911.12360>.
- Lenaïc Chizat, Edouard Oyallon, and Francis Bach. On lazy training in differentiable programming. *Advances in neural information processing systems*, 32, 2019.
- Puneesh Deora, Rouzbeh Ghaderi, Hossein Taheri, and Christos Thrampoulidis. On the optimization and generalization of multi-head attention. *arXiv preprint arXiv:2310.12680*, 2023.
- Simon Du, Jason Lee, Haochuan Li, Liwei Wang, and Xiyu Zhai. Gradient descent finds global minima of deep neural networks. In Kamalika Chaudhuri and Ruslan Salakhutdinov, editors, *Proceedings of the 36th International Conference on Machine Learning*, volume 97 of *Proceedings of Machine Learning Research*, pages 1675–1685. PMLR, 09–15 Jun 2019a. URL <https://proceedings.mlr.press/v97/du19c.html>.
- Simon S. Du, Xiyu Zhai, Barnabas Póczos, and Aarti Singh. Gradient descent provably optimizes over-parameterized neural networks, 2019b. URL <https://arxiv.org/abs/1810.02054>.
- Noah Golowich, Alexander Rakhlin, and Ohad Shamir. Size-independent sample complexity of neural networks. In *Conference On Learning Theory*, pages 297–299. PMLR, 2018.
- Moritz Hardt, Ben Recht, and Yoram Singer. Train faster, generalize better: Stability of stochastic gradient descent. In *International conference on machine learning*, pages 1225–1234. PMLR, 2016.
- Kaiming He, Xiangyu Zhang, Shaoqing Ren, and Jian Sun. Delving deep into rectifiers: Surpassing human-level performance on imagenet classification. In *Proceedings of the IEEE international conference on computer vision*, pages 1026–1034, 2015.
- Arthur Jacot, Franck Gabriel, and Clément Hongler. Neural tangent kernel: Convergence and generalization in neural networks. *Advances in neural information processing systems*, 31, 2018.
- Ziwei Ji and Matus Telgarsky. Polylogarithmic width suffices for gradient descent to achieve arbitrarily small test error with shallow relu networks, 2020. URL <https://arxiv.org/abs/1909.12292>.
- Alex Krizhevsky, Ilya Sutskever, and Geoffrey E Hinton. Imagenet classification with deep convolutional neural networks. In F. Pereira, C.J. Burges, L. Bottou, and K.Q. Weinberger, editors, *Advances in Neural Information Processing Systems*, volume 25. Curran Associates, Inc., 2012. URL https://proceedings.neurips.cc/paper_files/paper/2012/file/c399862d3b9d6b76c8436e924a68c45b-Paper.pdf.
- Ilja Kuzborskij and Csaba Szepesvári. Learning lipschitz functions by gd-trained shallow overparameterized relu neural networks, 2023. URL <https://arxiv.org/abs/2212.13848>.
- Jaehoon Lee, Lechao Xiao, Samuel Schoenholz, Yasaman Bahri, Roman Novak, Jascha Sohl-Dickstein, and Jeffrey Pennington. Wide neural networks of any depth evolve as linear models under gradient descent. *Advances in neural information processing systems*, 32, 2019.
- Yunwen Lei and Yiming Ying. Fine-grained analysis of stability and generalization for stochastic gradient descent. In *International Conference on Machine Learning*, pages 5809–5819. PMLR, 2020.

- Yunwen Lei, Rong Jin, and Yiming Ying. Stability and generalization analysis of gradient methods for shallow neural networks. In S. Koyejo, S. Mohamed, A. Agarwal, D. Belgrave, K. Cho, and A. Oh, editors, *Advances in Neural Information Processing Systems*, volume 35, pages 38557–38570. Curran Associates, Inc., 2022. URL https://proceedings.neurips.cc/paper_files/paper/2022/file/bf8fe6b79288f3d83696a5d276f4fc9d-Paper-Conference.pdf.
- Yuanzhi Li and Yingyu Liang. Learning overparameterized neural networks via stochastic gradient descent on structured data. *Advances in neural information processing systems*, 31, 2018.
- Chaoyue Liu, Libin Zhu, and Misha Belkin. On the linearity of large non-linear models: when and why the tangent kernel is constant. In H. Larochelle, M. Ranzato, R. Hadsell, M.F. Balcan, and H. Lin, editors, *Advances in Neural Information Processing Systems*, volume 33, pages 15954–15964. Curran Associates, Inc., 2020. URL https://proceedings.neurips.cc/paper_files/paper/2020/file/b7ae8fecf15b8b6c3c69eceae636d203-Paper.pdf.
- Fanghui Liu, Leello Dadi, and Volkan Cevher. Learning with norm constrained, over-parameterized, two-layer neural networks. *Journal of Machine Learning Research*, 25(138):1–42, 2024. URL <http://jmlr.org/papers/v25/22-1250.html>.
- Mehryar Mohri. Foundations of machine learning, 2018.
- Vaishnavh Nagarajan and J Zico Kolter. Uniform convergence may be unable to explain generalization in deep learning. *Advances in Neural Information Processing Systems*, 32, 2019.
- Behnam Neyshabur, Ryota Tomioka, and Nathan Srebro. Norm-based capacity control in neural networks. In *Conference on learning theory*, pages 1376–1401. PMLR, 2015.
- Atsushi Nitanda and Taiji Suzuki. Optimal rates for averaged stochastic gradient descent under neural tangent kernel regime. *arXiv preprint arXiv:2006.12297*, 2020.
- Atsushi Nitanda, Geoffrey Chinot, and Taiji Suzuki. Gradient descent can learn less over-parameterized two-layer neural networks on classification problems, 2020. URL <https://arxiv.org/abs/1905.09870>.
- Samet Oymak and Mahdi Soltanolkotabi. Toward moderate overparameterization: Global convergence guarantees for training shallow neural networks. *IEEE Journal on Selected Areas in Information Theory*, 1(1): 84–105, 2020.
- Dominic Richards and Ilja Kuzborskij. Stability & generalisation of gradient descent for shallow neural networks without the neural tangent kernel. *Advances in neural information processing systems*, 34:8609–8621, 2021.
- William H Rogers and Terry J Wagner. A finite sample distribution-free performance bound for local discrimination rules. *The Annals of Statistics*, pages 506–514, 1978.
- Matan Schliserman and Tomer Koren. Stability vs implicit bias of gradient methods on separable data and beyond. In *Conference on Learning Theory*, pages 3380–3394. PMLR, 2022.
- Matan Schliserman and Tomer Koren. Tight risk bounds for gradient descent on separable data. In A. Oh, T. Naumann, A. Globerson, K. Saenko, M. Hardt, and S. Levine, editors, *Advances in Neural Information Processing Systems*, volume 36, pages 68749–68759. Curran Associates, Inc., 2023. URL https://proceedings.neurips.cc/paper_files/paper/2023/file/d8ca28a32c05cd3b9b0940e43720f31b-Paper-Conference.pdf.
- Ohad Shamir. Gradient methods never overfit on separable data. *Journal of Machine Learning Research*, 22(85): 1–20, 2021. URL <http://jmlr.org/papers/v22/20-997.html>.
- Nathan Srebro, Karthik Sridharan, and Ambuj Tewari. Smoothness, low noise and fast rates. *Advances in neural information processing systems*, 23, 2010.
- Lili Su and Pengkun Yang. On learning over-parameterized neural networks: A functional approximation perspective. *Advances in Neural Information Processing Systems*, 32, 2019.
- Hossein Taheri and Christos Thrampoulidis. Generalization and stability of interpolating neural networks with minimal width. *Journal of Machine Learning Research*, 25(156):1–41, 2024. URL <http://jmlr.org/papers/v25/23-0422.html>.
- Hossein Taheri, Christos Thrampoulidis, and Arya Mazumdar. Sharper guarantees for learning neural network classifiers with gradient methods. In *The Thirteenth International Conference on Learning Representations*, 2025. URL <https://openreview.net/forum?id=h7GAgbLSmC>.

- Roman Vershynin. *High-dimensional probability: An introduction with applications in data science*, volume 47. Cambridge university press, 2018.
- Jiaming Xu and Hanjing Zhu. Overparametrized multi-layer neural networks: Uniform concentration of neural tangent kernel and convergence of stochastic gradient descent. *Journal of Machine Learning Research*, 25(94):1–83, 2024. URL <http://jmlr.org/papers/v25/23-0740.html>.
- Chiyuan Zhang, Samy Bengio, Moritz Hardt, Benjamin Recht, and Oriol Vinyals. Understanding deep learning requires rethinking generalization. *arXiv preprint arXiv:1611.03530*, 2016.
- Difan Zou, Yuan Cao, Dongruo Zhou, and Quanquan Gu. Stochastic gradient descent optimizes over-parameterized deep relu networks, 2018. URL <https://arxiv.org/abs/1811.08888>.

A Technical Lemmas

We define the diagonal sign matrix $\Sigma^l(\mathbf{x})$ with $l \in [L]$ by

$$\Sigma^l(\mathbf{x}) = \text{diag}\{\mathbb{I}\{\langle \mathbf{w}_r^l, h^{l-1}(\mathbf{x}) \rangle \geq 0\}\} \in \mathbb{R}^{m \times m}. \quad (10)$$

Then the deep ReLU network has the following matrix product representation:

$$f_{\mathbf{W}}(\mathbf{x}) = \mathbf{a}^\top \sqrt{\frac{2}{m}} \Sigma^L(\mathbf{x}) \mathbf{W}^L \cdots \sqrt{\frac{2}{m}} \Sigma^1(\mathbf{x}) \mathbf{W}^1 \mathbf{x}, \quad (11)$$

together with the presentation of $h^l(\mathbf{x})$:

$$h^l(\mathbf{x}) = \sqrt{\frac{2}{m}} \Sigma^l(\mathbf{x}) \mathbf{W}^l \cdots \sqrt{\frac{2}{m}} \Sigma^1(\mathbf{x}) \mathbf{W}^1 \mathbf{x}, \quad l \in [L]. \quad (12)$$

We further define $\mathbf{G}_l^l(\mathbf{x}) = \sqrt{2/m} \Sigma^l(\mathbf{x})$ and

$$\mathbf{G}_b^a(\mathbf{x}) = \sqrt{\frac{2}{m}} \Sigma^b(\mathbf{x}) \mathbf{W}^b \cdots \sqrt{\frac{2}{m}} \Sigma^a(\mathbf{x}), \quad 1 \leq a \leq b \leq L, \quad (13)$$

from which we can rewrite $f_{\mathbf{W}}(\mathbf{x})$ as

$$f_{\mathbf{W}}(\mathbf{x}) = \mathbf{a}^\top \mathbf{G}_L^L(\mathbf{x}) \mathbf{W}^L h^{L-1}(\mathbf{x}) = \langle (\mathbf{G}_L^L(\mathbf{x}))^\top \mathbf{a} (h^{L-1}(\mathbf{x}))^\top, \mathbf{W}^L \rangle.$$

Hence, for $l \in [L]$, we have

$$\frac{\partial f_{\mathbf{W}}(\mathbf{x})}{\partial \mathbf{W}^l} = (\mathbf{G}_L^l(\mathbf{x}))^\top \mathbf{a} (h^{l-1}(\mathbf{x}))^\top.$$

Similarly, we define

$$\mathbf{H}_b^a(\mathbf{x}) = \sqrt{\frac{2}{m}} \Sigma^b(\mathbf{x}) \mathbf{W}^b \cdots \sqrt{\frac{2}{m}} \Sigma^a(\mathbf{x}) \mathbf{W}^a, \quad 2 \leq a \leq b \leq L. \quad (14)$$

We denote $\Sigma_0^l(\mathbf{x})$, $h_0^l(\mathbf{x})$, $\mathbf{G}_{b,0}^a(\mathbf{x})$, $\mathbf{H}_{b,0}^a(\mathbf{x})$ as (10), (12), (13) and (14) with $\mathbf{W} = \mathbf{W}(0)$.

A.1 Properties of the Initialization

Lemma 2 (Theorem 4.4.5 in Vershynin [2018]). *With probability at least $1 - L \exp(-Cm)$ over the random choice of $\mathbf{W}(0)$, there exists an absolute constant $c_0 > 1$ such that for any $l \in [L]$, there holds*

$$\|\mathbf{W}^l(0)\|_2 \leq c_0 \sqrt{m}. \quad (15)$$

For a sub-exponential random variable X , its sub-exponential norm is defined as follows:

$$\|X\|_{\phi_1} = \inf\{t > 0 : \mathbb{E} \exp(|X|/t) \leq 2\}.$$

$X - \mathbb{E}X$ is sub-exponential too, satisfying

$$\|X - \mathbb{E}X\|_{\phi_1} \leq 2\|X\|_{\phi_1}. \quad (16)$$

If Y is a sub-gaussian random variable, we define the sub-gaussian norm of Y by

$$\|Y\|_{\phi_2} = \inf\{t > 0 : \mathbb{E} \exp(Y^2/t^2) \leq 2\}.$$

Suppose $Y \sim \mathcal{N}(0, r^2)$, then $\sigma(Y)$ is also sub-gaussian and we have

$$\|\sigma(Y)\|_{\phi_2} \leq \|Y\|_{\phi_2} \leq Cr. \quad (17)$$

We have the following lemma:

Lemma 3 (Lemma 2.7.6 in Vershynin [2018]). *A random variable X is sub-gaussian if and only if X^2 is sub-exponential. Moreover,*

$$\|X\|_{\phi_2}^2 = \|X^2\|_{\phi_1}.$$

Now we introduce Bernstein inequality with respect to $\|\cdot\|_{\phi_1}$,

Lemma 4 (Theorem 2.8.2 in Vershynin [2018]). *Let $X_1, \dots, X_m$ be independent, mean zero, sub-exponential random variables, and $d = (d_1, \dots, d_m) \in \mathbb{R}^m$, $K \geq \max_r \|X_r\|_{\phi_1}$. Then for every $t \geq 0$, we have*

$$\mathbb{P}\left(\left|\sum_{r=1}^m d_r X_r\right| \geq t\right) \leq 2 \exp\left[-c \min\left(\frac{t^2}{K^2 \|d\|_2^2}, \frac{t}{K \|d\|_\infty}\right)\right].$$

for some absolute constant c .

We introduce the following technical lemma related to the conditional expectation of Gaussian indicator function.

Lemma 5. *Suppose $\mathbf{w}$ is a m -dim Gaussian random vector with distribution $\mathcal{N}(0, \mathbf{I})$. Let $\mathbf{c} \neq 0, \mathbf{b}$ be two given vectors of m -dim. Then we have the following property*

$$\mathbb{E}[\mathbb{I}\{\langle \mathbf{w}, \mathbf{c} \rangle \geq 0\} \langle \mathbf{w}, \mathbf{b} \rangle^2] = \frac{\|\mathbf{b}\|_2^2}{2}.$$

Proof. Let $\mathbf{u} = \langle \mathbf{w}, \mathbf{c} \rangle, \mathbf{v} = \langle \mathbf{w}, \mathbf{b} \rangle$. Then $\mathbf{u} \sim \mathcal{N}(0, \|\mathbf{c}\|_2^2), \mathbf{v} \sim \mathcal{N}(0, \|\mathbf{b}\|_2^2)$. We decompose $\mathbf{v}$ into a component dependent on $\mathbf{u}$ and an independent residual $\mathbf{z}$:

$$\mathbf{v} = \frac{\text{Cov}(\mathbf{u}, \mathbf{v})}{\text{Var}(\mathbf{u})} \mathbf{u} + \mathbf{z} = \frac{\langle \mathbf{c}, \mathbf{b} \rangle}{\|\mathbf{c}\|_2^2} \mathbf{u} + \mathbf{z},$$

where $\mathbf{z} \sim \mathcal{N}\left(0, \|\mathbf{b}\|_2^2 - \frac{\langle \mathbf{c}, \mathbf{b} \rangle^2}{\|\mathbf{c}\|_2^2}\right)$ is independent of $\mathbf{u}$. Hence, we have

$$\begin{aligned} \mathbb{E}[\mathbb{I}\{\langle \mathbf{w}, \mathbf{c} \rangle \geq 0\} \langle \mathbf{w}, \mathbf{b} \rangle^2] &= \mathbb{E}[\mathbb{I}\{\mathbf{u} \geq 0\} \mathbf{v}^2] \\ &= \frac{\langle \mathbf{c}, \mathbf{b} \rangle^2}{\|\mathbf{c}\|_2^4} \mathbb{E}[\mathbb{I}\{\mathbf{u} \geq 0\} \mathbf{u}^2] + \frac{2\langle \mathbf{c}, \mathbf{b} \rangle}{\|\mathbf{c}\|_2^2} \mathbb{E}[\mathbb{I}\{\mathbf{u} \geq 0\} \mathbf{u} \mathbf{z}] + \mathbb{E}[\mathbb{I}\{\mathbf{u} \geq 0\} \mathbf{z}^2] \\ &= \frac{\langle \mathbf{c}, \mathbf{b} \rangle^2}{2\|\mathbf{c}\|_2^2} + 0 + \frac{1}{2} \mathbb{E} \mathbf{z}^2 \\ &= \frac{\langle \mathbf{c}, \mathbf{b} \rangle^2}{2\|\mathbf{c}\|_2^2} + \frac{1}{2} \cdot \left(\|\mathbf{b}\|_2^2 - \frac{\langle \mathbf{c}, \mathbf{b} \rangle^2}{\|\mathbf{c}\|_2^2}\right) = \frac{\|\mathbf{b}\|_2^2}{2}, \end{aligned}$$

where the second equality is due to $\mathbb{E}[\mathbb{I}\{\mathbf{u} \geq 0\} \mathbf{u}^2] = \frac{\|\mathbf{c}\|_2^2}{2}$ and the independence of $\mathbf{u}, \mathbf{z}$. The third equality follows from the distribution of $\mathbf{z}$. The proof is completed. $\square$

The following lemma studies the output of initialization at each layer.

Lemma 6. *For any $\delta > 0$, if $m \gtrsim L^2 \log(nL/\delta)$, then with probability at least $1 - \delta$,*

$$\|h_0^l(\mathbf{x}_i)\|_2 \in \left[\sqrt{\frac{2}{3}}, \sqrt{\frac{4}{3}}\right], \quad i = 1, \dots, n \quad \text{and} \quad l = 1, \dots, L.$$

Proof. This result directly follows Corollary A.2 in Zou et al. [2018], and we give the proof here for completeness. Note that for $1 \leq i \leq n, 1 \leq l \leq L$,

$$\|h_0^l(\mathbf{x}_i)\|_2^2 = \left\| \sqrt{\frac{2}{m}} \sigma(\mathbf{W}^l(0) h_0^{l-1}(\mathbf{x}_i)) \right\|_2^2 = \frac{1}{m} \sum_{r=1}^m 2\sigma^2(\langle \mathbf{w}_r^l(0), h_0^{l-1}(\mathbf{x}_i) \rangle).$$

Condition on $h_0^{l-1}(\mathbf{x}_i)$, we have $\langle \mathbf{w}_j^l(0), h_0^{l-1}(\mathbf{x}_i) \rangle \sim \mathcal{N}(0, \|h_0^{l-1}(\mathbf{x}_i)\|_2^2)$, hence

$$\mathbb{E} 2\sigma^2(\langle \mathbf{w}_j^l(0), h_0^{l-1}(\mathbf{x}_i) \rangle) = 2 \|h_0^{l-1}(\mathbf{x}_i)\|_2^2 \mathbb{E}_{u \sim \mathcal{N}(0,1)} \sigma^2(u) = \|h_0^{l-1}(\mathbf{x}_i)\|_2^2 \quad (18)$$

By (16) and Lemma 3, we have

$$\begin{aligned} &\|2\sigma^2(\langle \mathbf{w}_j^l(0), h_0^{l-1}(\mathbf{x}_i) \rangle) - \|h_0^{l-1}(\mathbf{x}_i)\|_2^2\|_{\phi_1} \\ &\leq 2\|2\sigma^2(\langle \mathbf{w}_j^l(0), h_0^{l-1}(\mathbf{x}_i) \rangle)\|_{\phi_1} \leq C\|\sigma(\langle \mathbf{w}_j^l(0), h_0^{l-1}(\mathbf{x}_i) \rangle)\|_{\phi_2}^2 \leq C\|h_0^{l-1}(\mathbf{x}_i)\|_2^2, \end{aligned}$$

where the last inequality is due to (17). Let $X_r = 2\sigma^2(\langle \mathbf{w}_r^l(0), h_0^{l-1}(\mathbf{x}_i) \rangle) - \|h_0^{l-1}(\mathbf{x}_i)\|_2^2$, $d_r = 1/m$, $K = C \|h_0^{l-1}(\mathbf{x}_i)\|_2^2$ and apply Lemma 4. We have for any $0 \leq t \leq 1$,

$$\begin{aligned} & \mathbb{P}(\|h_0^l(\mathbf{x}_i)\|_2^2 - \|h_0^{l-1}(\mathbf{x}_i)\|_2^2 \leq Ct \|h_0^{l-1}(\mathbf{x}_i)\|_2^2 | h_0^{l-1}(\mathbf{x}_i)) \\ &= \mathbb{P}\left(\left|\frac{1}{m} \sum_{r=1}^m (\mathbb{E} 2\sigma^2(\langle \mathbf{w}_r^l(0), h_0^{l-1}(\mathbf{x}_i) \rangle) - \|h_0^{l-1}(\mathbf{x}_i)\|_2^2)\right| \leq Ct \|h_0^{l-1}(\mathbf{x}_i)\|_2^2 | h_0^{l-1}(\mathbf{x}_i)\right) \\ &\geq 1 - 2 \exp(-Cm \min\{t^2, t\}) = 1 - 2 \exp(-Cmt^2). \end{aligned}$$

Taking union bounds over i, l , there holds for any $1 \leq i \leq n$ and $1 \leq l \leq L$,

$$\mathbb{P}(\|h_0^l(\mathbf{x}_i)\|_2^2 - \|h_0^{l-1}(\mathbf{x}_i)\|_2^2 \leq Ct \|h_0^{l-1}(\mathbf{x}_i)\|_2^2) \geq 1 - 2nL \exp(-Cmt^2).$$

Since $m \gtrsim L^2 \log(nL/\delta)$, let $t = \sqrt{\log(nL/\delta)/m}$, we have with probability at least $1 - \delta$, there holds

$$\|h_0^l(\mathbf{x}_i)\|_2^2 - \|h_0^{l-1}(\mathbf{x}_i)\|_2^2 \leq C \sqrt{\frac{\log(nL/\delta)}{m}} \|h_0^{l-1}(\mathbf{x}_i)\|_2^2, \quad 1 \leq i \leq n, 1 \leq l \leq L. \quad (19)$$

Now we show the following inequality holds with probability at least $1 - \delta$,

$$\|h_0^l(\mathbf{x}_i)\|_2^2 - 1 \leq \frac{4lC}{3} \sqrt{\frac{\log(nL/\delta)}{m}} \leq \frac{1}{3}. \quad (20)$$

When $l = 0$, it is true. If (20) holds for $l \in [L-1]$, then $\|h_0^l(\mathbf{x}_i)\|_2^2 \leq 4/3$. Combined with (19), we have with probability at least $1 - \delta$,

$$\begin{aligned} \||h_0^{l+1}(\mathbf{x}_i)\|_2^2 - 1| &\leq \||h_0^l(\mathbf{x}_i)\|_2^2 - 1| + \||h_0^{l+1}(\mathbf{x}_i)\|_2^2 - \|h_0^l(\mathbf{x}_i)\|_2^2| \\ &\leq \frac{4lC}{3} \sqrt{\frac{\log(nL/\delta)}{m}} + C \sqrt{\frac{\log(nL/\delta)}{m}} \|h_0^l(\mathbf{x}_i)\|_2^2 \\ &\leq \frac{4(l+1)C}{3} \sqrt{\frac{\log(nL/\delta)}{m}} \leq \frac{1}{3}. \end{aligned}$$

Hence, (20) holds for all $i \in [n], l \in [L]$, which implies the lemma. $\square$

Lemma 7. For any $\mathbf{x} \in \mathcal{X}$, we have

$$\mathbf{a}^\top \Sigma_0^L(\mathbf{x}) \mathbf{W}^L(0) = 0 \text{ and } \frac{\partial f_{\mathbf{W}(0)}(\mathbf{x})}{\partial \mathbf{W}^l(0)} = 0, \quad l \in [L-1].$$

Proof. Note the r -th row of $\Sigma_0^L(\mathbf{x}) \mathbf{W}^L(0)$ is $\mathbb{I}\{\langle \mathbf{w}_r^L(0), h_0^{L-1}(\mathbf{x}) \rangle \geq 0\} \mathbf{w}_r^L(0)$. Since $a_r = -a_{r+\frac{m}{2}}$ and $\mathbf{w}_r^L(0) = \mathbf{w}_{r+\frac{m}{2}}^L(0)$ for all $r \in [\frac{m}{2}]$, we have

$$\begin{aligned} & \mathbf{a}^\top \Sigma_0^L(\mathbf{x}) \mathbf{W}^L(0) \\ &= \sum_{r=1}^m a_r \mathbb{I}\{\langle \mathbf{w}_r^L(0), h_0^{L-1}(\mathbf{x}) \rangle \geq 0\} \mathbf{w}_r^L(0) \\ &= \sum_{r=1}^{\frac{m}{2}} a_r \mathbb{I}\{\langle \mathbf{w}_r^L(0), h_0^{L-1}(\mathbf{x}) \rangle \geq 0\} \mathbf{w}_r^L(0) + \sum_{r=1}^{\frac{m}{2}} a_{r+\frac{m}{2}} \mathbb{I}\{\langle \mathbf{w}_{r+\frac{m}{2}}^L(0), h_0^{L-1}(\mathbf{x}) \rangle \geq 0\} \mathbf{w}_r^L(0) \\ &= \sum_{r=1}^{\frac{m}{2}} (a_r - a_r) \mathbb{I}\{\langle \mathbf{w}_r^L(0), h_0^{L-1}(\mathbf{x}) \rangle \geq 0\} \mathbf{w}_r^L(0) = 0. \end{aligned}$$

It then follows that for all $l \in [L-1]$,

$$\left(\frac{\partial f_{\mathbf{W}(0)}(\mathbf{x})}{\partial \mathbf{W}^l(0)}\right)^\top = h_0^{l-1}(\mathbf{x}) \mathbf{a}^\top (\mathbf{G}_{L,0}^l(\mathbf{x}))^\top = h_0^{l-1}(\mathbf{x}) \mathbf{a}^\top \sqrt{\frac{2}{m}} \Sigma_0^L(\mathbf{x}) \mathbf{W}^L(0) \cdots \sqrt{\frac{2}{m}} \Sigma_0^l(\mathbf{x}) = 0.$$

Hence, the proof is completed. $\square$

Lemma 8. Suppose $m \gtrsim L^2 \log(nL/\delta)$, then with probability at least $1 - \delta$, for all $i \in [n]$,

$$\left\| \frac{\partial f_{\mathbf{W}(0)}(\mathbf{x}_i)}{\partial \mathbf{W}^L(0)} \right\|_F \leq \sqrt{2}.$$

Proof. By Hoeffding inequality, condition on $h_0^{L-1}(\mathbf{x}_i)$, with probability at least $1 - \delta$, there holds

$$\frac{1}{m} \mathbf{a}^\top \Sigma_0^L(\mathbf{x}_i) \Sigma_0^L(\mathbf{x}_i) \mathbf{a} = \frac{1}{m} \sum_{j=1}^m \mathbb{I}\{\langle \mathbf{w}_j^L(0), h_0^{L-1}(\mathbf{x}_i) \rangle \geq 0\} \leq \frac{1}{2} + C \sqrt{\frac{\log(n/\delta)}{m}} \leq \frac{3}{4}.$$

Combined with Lemma 6, we have

$$\left\| \frac{\partial f_{\mathbf{W}(0)}(\mathbf{x}_i)}{\partial \mathbf{W}^L(0)} \right\|_F^2 = \frac{2}{m} \|h_0^{L-1}(\mathbf{x}_i)\|_2^2 \mathbf{a}^\top \Sigma_0^L(\mathbf{x}_i) \Sigma_0^L(\mathbf{x}_i) \mathbf{a} \leq 2 \cdot \frac{4}{3} \cdot \frac{3}{4} = 2.$$

The proof is completed. $\square$

Let $\Sigma_1, \Sigma_2 \in \mathbb{R}^{m \times m}$ be two diagonal matrices with entries in $\{0, 1\}$.

Lemma 9. Suppose $m \gtrsim L^2 \log(nL/\delta)$, $s \lesssim m/(L^2 \log m)$, then with probability at least $1 - \delta$ we have for all $i \in [n]$, $2 \leq a \leq b \leq L$,

$$\sup_{\|\Sigma_1\|_0 \leq s} \|\mathbf{H}_{b,0}^a(\mathbf{x}_i) \Sigma_1\|_2 \lesssim 1. \quad (21)$$

Proof. We need to prove that for any $v \in S^{m-1}$, there holds

$$\|\mathbf{H}_{b,0}^a(\mathbf{x}_i) \Sigma_1 v\|_2 \lesssim 1. \quad (22)$$

Note that $\|\Sigma_1 v\|_0 \leq s$ and $\|\Sigma_1 v\|_2 \leq 1$. Let $\mathcal{P} = \{v \in S^{m-1} : \|v\|_0 \leq s\}$. We only need to prove that the following inequality holds with probability at least $1 - \delta$:

$$\sup_{v \in \mathcal{P}} \|\mathbf{H}_{b,0}^a(\mathbf{x}_i) v\|_2 \lesssim 1. \quad (23)$$

Let S be a subspace of S^{m-1} that has at most s non-zero coordinates. For such a subspace, we choose a $1/2$ -cover of it and denote this cover by $\mathcal{Q}$. By Lemma 4.2.13 in Vershynin [2018],

$$|\mathcal{Q}| \leq 5^s.$$

The number of such subspaces is $M = \binom{m}{s}$. We denote all subspaces by $\mathcal{S}_1, \dots, \mathcal{S}_M$, and the corresponding covers $\mathcal{Q}_1, \dots, \mathcal{Q}_M$. Let $\bigcup \mathcal{Q} = \{v_1, \dots, v_{M'}\}$ with $M' \leq \binom{m}{s} 5^s$. We first prove that (23) is true for all v_j , then it holds simultaneously for all elements in $\mathcal{P}$.

For a unit vector v , we define

$$v^l(\mathbf{x}_i) = \mathbf{H}_{l,0}^a(\mathbf{x}_i) v, \quad a \leq l \leq b.$$

and $v^{a-1}(\mathbf{x}_i) = v$. Note that condition on $h_0^{L-1}(\mathbf{x}_i), v^{l-1}(\mathbf{x}_i)$, we take expectation over $\mathbf{w}_r^L(0)$, applying Lemma 5 implies that

$$\mathbb{E}[2\mathbb{I}\{\langle \mathbf{w}_r^L(0), h_0^{L-1}(\mathbf{x}_i) \rangle \geq 0\} (\langle \mathbf{w}_r^L(0), v^{l-1}(\mathbf{x}_i) \rangle)^2] = \|v^{l-1}(\mathbf{x}_i)\|_2^2.$$

It then follows that

$$\begin{aligned} \|v^l(\mathbf{x}_i)\|_2^2 &= \|\mathbf{H}_{l,0}^a(\mathbf{x}_i) v\|_2^2 = \left\| \sqrt{\frac{2}{m}} \Sigma_0^L(\mathbf{x}_i) \mathbf{W}^L(0) v^{l-1}(\mathbf{x}_i) \right\|_2^2 \\ &= \frac{1}{m} \sum_{r=1}^m \mathbb{E} 2\mathbb{I}\{\langle \mathbf{w}_r^L(0), h_0^{L-1}(\mathbf{x}_i) \rangle \geq 0\} (\langle \mathbf{w}_r^L(0), v^{l-1}(\mathbf{x}_i) \rangle)^2 = \|v^{l-1}(\mathbf{x}_i)\|_2^2. \end{aligned}$$

Similar to the proof of Lemma 6, for every v_j , $1 \leq j \leq M'$, we apply Lemma 4 to get

$$\mathbb{P} \left(\left| \|v_j^l(\mathbf{x}_i)\|_2^2 - \|v_j^{l-1}(\mathbf{x}_i)\|_2^2 \right| \leq \frac{C \|v_j^{l-1}(\mathbf{x}_i)\|_2^2}{L} |v_j^{l-1}(\mathbf{x}_i) \right) \geq 1 - 2 \exp \left(-Cm \min \left\{ \frac{1}{L^2}, \frac{1}{L} \right\} \right).$$

Taking the union bounds for all j, l, i yields

$$\begin{aligned}
& \mathbb{P} \left(\left| \|v_j^l(\mathbf{x}_i)\|_2^2 - \|v_j^{l-1}(\mathbf{x}_i)\|_2^2 \right| \leq \frac{C\|v_j^{l-1}(\mathbf{x}_i)\|_2^2}{L} \right) \\
& \geq 1 - 2 \cdot \binom{m}{s} 5^s nL \exp \left(\frac{-Cm}{L^2} \right) \\
& \geq 1 - 2 \cdot (5em)^s nL \exp \left(\frac{-Cm}{L^2} \right) \\
& = 1 - \exp \left[\log \delta + s \log m + s \log(5e) + \log \left(\frac{2nL}{\delta} \right) - \frac{C}{L^2} m \right] \geq 1 - \delta,
\end{aligned}$$

where we have used $\binom{m}{s} \leq (em/s)^s \leq (em)^s$ in the second inequality and the last inequality is due to $m \gtrsim L^2 \log(nL/\delta)$, $s \lesssim m/(L^2 \log m)$. Hence, we have with probability at least $1 - \delta$,

$$\|v_j^l(\mathbf{x}_i)\|_2^2 \leq \left(1 + \frac{C}{L}\right) \|v_j^{l-1}(\mathbf{x}_i)\|_2^2 \leq \left(1 + \frac{C}{L}\right)^L \|v_j\|_2^2 \lesssim 1.$$

For any unit vector v with $\|v\|_0 \leq s$, consider the subspace S containing it and the corresponding $1/2$ -cover $\mathcal{Q}$. There exists a unit vector $v_j, j \in [M']$, $\|v - v_j\|_0 \leq s$ and $\|v - v_j\|_2 \leq 1/2$. Thus,

$$\begin{aligned}
& \|\mathbf{H}_{b,0}^a(\mathbf{x}_i)v\|_2 \\
& \leq \|\mathbf{H}_{b,0}^a(\mathbf{x}_i)v_j\|_2 + \|\mathbf{H}_{b,0}^a(\mathbf{x}_i)(v - v_j)\|_2 \\
& = \|v_j^b(\mathbf{x}_i)\|_2 + \|v - v_j\|_2 \left\| \mathbf{H}_{b,0}^a(\mathbf{x}_i) \frac{v - v_j}{\|v - v_j\|_2} \right\|_2 \\
& \lesssim 1 + \|v - v_j\|_2 \sup_{v \in \mathcal{P}} \|\mathbf{H}_{b,0}^a(\mathbf{x}_i)v\|_2 \leq 1 + \frac{1}{2} \sup_{v \in \mathcal{P}} \|\mathbf{H}_{b,0}^a(\mathbf{x}_i)v\|_2.
\end{aligned}$$

Taking sup to the both sides yields

$$\sup_{v \in \mathcal{P}} \|\mathbf{H}_{b,0}^a(\mathbf{x}_i)v\|_2 \lesssim 1 + \frac{1}{2} \sup_{v \in \mathcal{P}} \|\mathbf{H}_{b,0}^a(\mathbf{x}_i)v\|_2,$$

which implies $\sup_{v \in \mathcal{P}} \|\mathbf{H}_{b,0}^a(\mathbf{x}_i)v\|_2 \lesssim 1$. Hence (23) holds and the proof is completed. $\square$

Remark 7. Our proofs are inspired by Lemma A.9 in Zou et al. [2018], which establishes the estimates under the condition $s \gtrsim \log(nL/\delta)$. However, we eliminate this assumption via a more refined analysis.

Lemma 10. Suppose $m \gtrsim L^2 \log(nL/\delta)$, then with probability at least $1 - \delta$, we have for all $i \in [n]$, $2 \leq a \leq b \leq L$,

$$\|\mathbf{H}_{b,0}^a(\mathbf{x}_i)\|_2 \lesssim L\sqrt{\log m}.$$

Although the left-hand side of above inequality is the production of L terms, it is bounded by $\tilde{O}(L)$. This lemma shows that the introduction of ReLU activation can avoid exponential explosion.

Proof. For any unit vector v , we decompose it as $v = v_1 + \dots + v_q$, where $v_j, j \in [q]$ are all s -sparse vectors on different coordinates. Therefore,

$$\|v\|_2^2 = \sum_{j=1}^q \|v_j\|_2^2.$$

Here we choose $s \asymp m/(L^2 \log m)$, then $q \lesssim m/s \lesssim L^2 \log m$. Applying Lemma 9, we have

$$\begin{aligned}
\|\mathbf{H}_{b,0}^a(\mathbf{x}_i)v\|_2 & = \left\| \sum_{j=1}^q \mathbf{H}_{b,0}^a(\mathbf{x}_i)v_j \right\|_2 \leq \sum_{j=1}^q \|\mathbf{H}_{b,0}^a(\mathbf{x}_i)v_j\|_2 \leq \sqrt{q \sum_{j=1}^q \|\mathbf{H}_{b,0}^a(\mathbf{x}_i)v_j\|_2^2} \\
& \lesssim \sqrt{q \sum_{j=1}^q \|v_j\|_2^2} = \sqrt{q\|v\|_2^2} = \sqrt{q} \lesssim L\sqrt{\log m}.
\end{aligned}$$

where we have used Cauchy-Schwartz's inequality in the second inequality. Hence,

$$\|\mathbf{H}_{b,0}^a(\mathbf{x}_i)\|_2 \lesssim L\sqrt{\log m}.$$

The proof is completed. $\square$

From the above lemma, we know that if $m \gtrsim L^2 \log(nL/\delta) \log m$, then

$$\|\mathbf{G}_{b,0}^a(x_i)\|_2 \lesssim L\sqrt{\frac{\log m}{m}}, \quad i \in [n], 1 \leq a \leq b \leq L. \quad (24)$$

Remark 8. In Lemma 10, we introduce a useful technique that decomposes the unit vector into sparse components. This approach reduces the covering number from 5^m to $5^s \binom{m}{s}$, making it easier for high-probability bounds to hold. A related method appears in Lemma 7.3 of Allen-Zhu et al. [2019], but their width exhibits polynomial dependence on n . In contrast, our analysis achieves polylogarithmic width, substantially relaxing the overparameterization requirement.

Using similar techniques we can obtain the following lemma:

Lemma 11. *Suppose $m \gtrsim L^2 \log(nL/\delta)$, then with probability at least $1 - \delta$, for all $i \in [n]$, $2 \leq a \leq b \leq L$, $\|\Sigma_1\|_0, \|\Sigma_2\|_0 \leq s \lesssim m/(L^2 \log m)$,*

$$\left\| \Sigma_1 \sqrt{\frac{2}{m}} \mathbf{W}^b(0) \mathbf{H}_{b-1,0}^a(\mathbf{x}_i) \Sigma_2 \right\|_2 \lesssim \frac{1}{L}.$$

Proof. If $s = 0$, the above inequality becomes $0 \lesssim 1/L$, which holds true. Now we assume $s \geq 1$. Similar to Lemma 9, we only need to prove that for any s -sparse unit vector v there holds

$$\left\| \Sigma_1 \sqrt{\frac{2}{m}} \mathbf{W}^b(0) \mathbf{H}_{b-1,0}^a(\mathbf{x}_i) v \right\|_2 \lesssim \frac{1}{L}. \quad (25)$$

We use the same notation as in Lemma 9, it then follows that for all $j \in [M']$, $a \leq l \leq b, i \in [n]$,

$$\mathbb{P} \left(\left| \|v_j^l(\mathbf{x}_i)\|_2^2 - \|v_j^{l-1}(\mathbf{x}_i)\|_2^2 \right| \leq \frac{C \|v_j^{l-1}(\mathbf{x}_i)\|_2^2}{L} |v_j^{l-1}(\mathbf{x}_i)| \right) \geq 1 - 2 \exp \left(-Cm \min \left\{ \frac{1}{L^2}, \frac{1}{L} \right\} \right).$$

For a fixed Σ_1 , we assume $\Sigma_1 = \text{diag}\{d_1, \dots, d_m\}$ with $d_r \in \{0, 1\}$, $\sum_r d_r \leq s, r \in [m]$. We have

$$\left\| \Sigma_1 \sqrt{\frac{2}{m}} \mathbf{W}^b(0) v_j^{b-1}(\mathbf{x}_i) \right\|_2^2 = \sum_{r=1}^m d_r^2 \frac{2}{m} (\langle \mathbf{w}_r^b, v_j^{b-1}(\mathbf{x}_i) \rangle)^2. \quad (26)$$

Condition on $v_j^{b-1}(\mathbf{x}_i)$, there holds

$$\mathbb{E} \frac{2}{m} (\langle \mathbf{w}_r^b(0), v_j^{b-1}(\mathbf{x}_i) \rangle)^2 = \frac{2}{m} \|v_j^{b-1}(\mathbf{x}_i)\|_2^2.$$

Let $X_r = \frac{2}{m} (\langle \mathbf{w}_r^b(0), v_j^{b-1}(\mathbf{x}_i) \rangle)^2 - \frac{2}{m} \|v_j^{b-1}(\mathbf{x}_i)\|_2^2$ and $d = (d_1, \dots, d_m)$. Then X_r are mean-zero sub-exponential random variables, following similar discussions in Lemma 6, we have $\|X_r\|_{\phi_1} \leq \frac{C}{m} \|v_j^{b-1}(\mathbf{x}_i)\|_2^2$. Moreover, $\|d\|_2^2 = \sum_{r=1}^m d_r^2 = \sum_{r=1}^m d_r \leq s, \|d\|_\infty = 1$. Applying Lemma 4, we have

$$\begin{aligned} & \mathbb{P} \left(\left| \sum_{r=1}^m d_r \left(\frac{2}{m} (\langle \mathbf{w}_r^b(0), v_j^{b-1}(\mathbf{x}_i) \rangle)^2 - \frac{2}{m} \|v_j^{b-1}(\mathbf{x}_i)\|_2^2 \right) \right| \geq t \|v_j^{b-1}(\mathbf{x}_i)\|_2^2 |v_j^{b-1}(\mathbf{x}_i)| \right) \\ & \leq 2 \exp \left[-C \min \left(\frac{t^2 m^2}{s}, tm \right) \right]. \end{aligned} \quad (27)$$

Choosing $t = 1/L^2$ and note that $s \leq m/(L^2 \log m)$, we have

$$\begin{aligned} & \mathbb{P} \left(\left\| \Sigma_1 \sqrt{\frac{2}{m}} \mathbf{W}^b(0) v_j^{b-1}(\mathbf{x}_i) \right\|_2^2 \leq \left(\frac{2s}{m} + \frac{1}{L^2} \right) \|v_j^{b-1}(\mathbf{x}_i)\|_2^2 | v_j^{b-1}(\mathbf{x}_i) \right) \\ & \geq \mathbb{P} \left(\sum_{r=1}^m d_r^2 \frac{2}{m} (\langle \mathbf{w}_r^b(0), v_j^{b-1}(\mathbf{x}_i) \rangle)^2 \leq \sum_{r=1}^m \frac{2d_r}{m} \|v_j^{b-1}(\mathbf{x}_i)\|_2^2 + \frac{1}{L^2} \|v_j^{b-1}(\mathbf{x}_i)\|_2^2 | v_j^{b-1}(\mathbf{x}_i) \right) \\ & \geq 1 - 2 \exp \left(-\frac{Cm}{L^2} \right), \end{aligned}$$

where the first inequality is due to (26) and $\sum_{r=1}^m d_r \leq s$, the last inequality results from (27).

Taking union bounds over all $\mathbf{x}_i, \Sigma_1, v_j, l$, and note that

$$2nL \binom{m}{s} 5^s \binom{m}{s} \exp \left(-\frac{Cm}{L^2} \right) \leq \frac{\delta}{2}.$$

We have with probability at least $1 - \delta$,

$$\begin{aligned} & \left\| \Sigma_1 \sqrt{\frac{2}{m}} \mathbf{W}^b(0) v_j^{b-1}(\mathbf{x}_i) \right\|_2^2 \leq \left(\frac{2s}{m} + \frac{1}{L^2} \right) \|v_j^{b-1}(\mathbf{x}_i)\|_2^2 \\ & \leq \left(\frac{2s}{m} + \frac{1}{L^2} \right) \left(1 + \frac{C}{L} \right) \|v_j^{b-2}(\mathbf{x}_i)\|_2^2 \leq \left(\frac{2s}{m} + \frac{1}{L^2} \right) \left(1 + \frac{C}{L} \right)^L \|v_j\|_2^2 \lesssim \frac{1}{L^2}. \end{aligned}$$

Hence, we have

$$\left\| \Sigma_1 \sqrt{\frac{2}{m}} \mathbf{W}^b(0) \mathbf{H}_{b-1,0}^a(\mathbf{x}_i) v_j \right\|_2 \lesssim \frac{1}{L}.$$

Following same techniques of using 1/2-cover in the proof of Lemma 9, we can prove (25), and then complete the proof of the lemma. $\square$

Additionally, apply similar methods in Lemma 10 by decomposing the unit vector into s -sparse vectors, we have

$$\left\| \Sigma_1 \sqrt{\frac{2}{m}} \mathbf{W}_0^b(\mathbf{x}_i) \mathbf{H}_{b-1,0}^a(\mathbf{x}_i) \right\|_2 \lesssim \sqrt{\log m}. \quad (28)$$

Remark 9. To analyze the influence of the sparse matrix Σ_1 on (25), we propose a key technical improvement: instead of resorting to covering number arguments as in Zou et al. [2018], Allen-Zhu et al. [2019], we leverage a weighted Bernstein inequality. Particularly, existing methods require taking another union bound over both all s -sparse subspaces and their covers (of size $\sim \binom{m}{s} 9^s$), whereas our analysis only needs to union over the sparse subspaces themselves (of cardinality $\binom{m}{s}$). Our method directly demonstrates that sparsity inherently lowers computational costs by avoiding the need for dense covers. The simplicity of our technique also underscores the intrinsic benefits of sparse structures in optimization.

A.2 Properties of Perturbation Terms

Recall that

$$\mathbf{G}_{b,0}^a(\mathbf{x}) = \sqrt{\frac{2}{m}} \Sigma_0^b(\mathbf{x}) \mathbf{W}^b(0) \cdots \sqrt{\frac{2}{m}} \Sigma_0^{a+1}(\mathbf{x}) \mathbf{W}^{a+1}(0) \sqrt{\frac{2}{m}} \Sigma_0^a(\mathbf{x}).$$

For any $l \in [L]$, let $\widehat{\mathbf{W}}^l$ and the diagonal matrix $\widehat{\Sigma}^l(\mathbf{x})$ be the matrices with the same size of $\mathbf{W}^l(0)$ and $\Sigma_0^l(\mathbf{x})$, respectively. Define

$$\begin{aligned} \widehat{\mathbf{G}}_b^a(\mathbf{x}) &= \sqrt{\frac{2}{m}} (\Sigma_0^b(\mathbf{x}) + \widehat{\Sigma}^b(\mathbf{x})) (\mathbf{W}^b(0) + \widehat{\mathbf{W}}^b) \cdots \sqrt{\frac{2}{m}} (\Sigma_0^{a+1}(\mathbf{x}) + \widehat{\Sigma}^{a+1}(\mathbf{x})) \\ &\quad \times (\mathbf{W}^{a+1}(0) + \widehat{\mathbf{W}}^{a+1}) \sqrt{\frac{2}{m}} (\Sigma_0^a(\mathbf{x}) + \widehat{\Sigma}^a(\mathbf{x})), \quad 1 \leq a \leq b \leq L \end{aligned} \quad (29)$$

and $\widehat{\mathbf{G}}_l^l(\mathbf{x}) = \sqrt{\frac{2}{m}} (\Sigma_0^l(\mathbf{x}) + \widehat{\Sigma}^l(\mathbf{x}))$ for all $l \in [L]$.

Lemma 12. Let $\widehat{\mathbf{G}}_b^a(\mathbf{x})$ with $1 \leq a \leq b \leq L$ be the matrix defined in (29). Assume $\max_{l \in [L]} \|\widehat{\mathbf{W}}^l\|_2 \leq R \lesssim \sqrt{m}/(L^2 \sqrt{\log m})$, $m \gtrsim L^2 \log(nL/\delta)$ and $\widehat{\Sigma}^l(\mathbf{x}_i) \in [-1, 1]^{m \times m}$, $\|\widehat{\Sigma}^l(\mathbf{x}_i)\|_0 \leq s \lesssim m/(L^2 \log m)$ for all $i \in [n], l \in [L]$. Then, with probability at least $1 - \delta$ for all $1 \leq a \leq b \leq L, i \in [n]$, there holds

$$\|\widehat{\mathbf{G}}_b^a(\mathbf{x}_i)\|_2 \lesssim L \sqrt{\frac{\log m}{m}}.$$

Proof. The proof is similar to that of Lemma 8.6 in Allen-Zhu et al. [2019], we give the proof here for the sake of completeness. We first prove that for any $1 \leq a \leq b \leq L$,

$$\left\| \sqrt{\frac{2}{m}}(\Sigma_0^b(\mathbf{x}_i) + \widehat{\Sigma}^b(\mathbf{x}_i))\mathbf{W}^b(0) \cdots \sqrt{\frac{2}{m}}(\Sigma_0^{a+1}(\mathbf{x}_i) + \widehat{\Sigma}^{a+1}(\mathbf{x}_i))\mathbf{W}^{a+1}(0) \right\|_2 \lesssim L \sqrt{\log m}. \quad (30)$$

We define a diagonal matrix $(\widehat{\Sigma}_1^l(\mathbf{x}_i))_{k,k} = \mathbb{I}\{\widehat{\Sigma}^l(\mathbf{x}_i)_{k,k} \neq 0\}$, and $\|\widehat{\Sigma}_1^l(\mathbf{x}_i)\|_0 \leq s$. Therefore, $\widehat{\Sigma}^l(\mathbf{x}_i) = \widehat{\Sigma}_1^l(\mathbf{x}_i)\widehat{\Sigma}^l(\mathbf{x}_i)\widehat{\Sigma}_1^l(\mathbf{x}_i)$. We decompose the left term of (30) into 2^{b-a} terms and control them respectively. Each matrix can be written as (ignoring the superscripts and $\mathbf{x}_i$).

$$\begin{aligned} & \left(\Sigma_0 \sqrt{\frac{2}{m}} \mathbf{W}(0) \cdots \widehat{\Sigma}_1 \right) \widehat{\Sigma} \left(\widehat{\Sigma}_1 \sqrt{\frac{2}{m}} \mathbf{W}(0) \cdots \sqrt{\frac{2}{m}} \mathbf{W}(0) \widehat{\Sigma}_1 \right) \widehat{\Sigma} \cdots \widehat{\Sigma} \\ & \quad \times \left(\widehat{\Sigma}_1 \sqrt{\frac{2}{m}} \mathbf{W}(0) \cdots \Sigma_0 \sqrt{\frac{2}{m}} \mathbf{W}(0) \right). \end{aligned}$$

Then, with probability at least $1 - \delta$, there holds:

- By Lemma 9, $\left\| \Sigma_0 \sqrt{\frac{2}{m}} \mathbf{W}(0) \cdots \widehat{\Sigma}_1 \right\|_2 \lesssim 1$.
- By Lemma 11, $\left\| \widehat{\Sigma}_1 \sqrt{\frac{2}{m}} \mathbf{W}(0) \cdots \sqrt{\frac{2}{m}} \mathbf{W}(0) \widehat{\Sigma}_1 \right\|_2 \lesssim 1/L$.
- By (28), $\left\| \widehat{\Sigma}_1 \sqrt{\frac{2}{m}} \mathbf{W}(0) \cdots \Sigma_0 \sqrt{\frac{2}{m}} \mathbf{W}(0) \right\|_2 \lesssim \sqrt{\log m}$.
- When there is no $\widehat{\Sigma}$, by Lemma 10, $\left\| \Sigma_0 \sqrt{\frac{2}{m}} \mathbf{W}(0) \cdots \Sigma_0 \sqrt{\frac{2}{m}} \mathbf{W}(0) \right\|_2 \lesssim L \sqrt{\log m}$.

Combined with these results, counting the number of $\widehat{\Sigma}$, we obtain

$$\begin{aligned} & \left\| \sqrt{\frac{2}{m}}(\Sigma_0^b(\mathbf{x}_i) + \widehat{\Sigma}^b(\mathbf{x}_i))\mathbf{W}^b(0) \cdots \sqrt{\frac{2}{m}}(\Sigma_0^{a+1}(\mathbf{x}_i) + \widehat{\Sigma}^{a+1}(\mathbf{x}_i))\mathbf{W}^{a+1}(0) \right\|_2 \\ & \lesssim L \sqrt{\log m} + \sum_{j=1}^{b-a} \binom{b-a}{j} \left(\frac{1}{L} \right)^{j-1} 1^j \sqrt{\log m} \\ & \leq L \sqrt{\log m} \left(1 + \sum_{j=1}^L \left(\frac{eL}{j} \right)^j \left(\frac{1}{L} \right)^j \right) \lesssim L \sqrt{\log m}, \end{aligned}$$

where in the second inequality we have used $\binom{b-a}{j} \leq (e(b-a)/j)^j \leq (eL/j)^j$, the last inequality is due to $\sum_{j=1}^L (eL/j)^j$ converges and it is bounded by a constant. Now we have proved (30).

Denote $\Sigma' = \Sigma_0 + \widehat{\Sigma}$, through similar expansion, $\Sigma' \sqrt{\frac{2}{m}}(\mathbf{W}(0) + \widehat{\mathbf{W}}) \cdots (\Sigma') \sqrt{\frac{2}{m}}(\mathbf{W}(0) + \widehat{\mathbf{W}})$

is the sum of following terms

$$\begin{aligned} & \left(\Sigma' \sqrt{\frac{2}{m}} \mathbf{W}(0) \cdots \Sigma' \right) \sqrt{\frac{2}{m}} \widehat{\mathbf{W}} \left(\Sigma' \sqrt{\frac{2}{m}} \mathbf{W}(0) \cdots \Sigma' \right) \sqrt{\frac{2}{m}} \widehat{\mathbf{W}} \cdots \sqrt{\frac{2}{m}} \widehat{\mathbf{W}} \\ & \quad \times \left(\Sigma' \sqrt{\frac{2}{m}} \mathbf{W}(0) \cdots \Sigma' \sqrt{\frac{2}{m}} \mathbf{W}(0) \right). \end{aligned}$$

Since $\|\Sigma'\|_2 \lesssim 1$, using Eq. (30), we have

$$\begin{aligned} \left\| \Sigma' \sqrt{\frac{2}{m}} \mathbf{W}(0) \cdots \Sigma' \right\|_2 & \lesssim L \sqrt{\log m}, \\ \left\| \Sigma' \sqrt{\frac{2}{m}} \mathbf{W}(0) \cdots \Sigma' \sqrt{\frac{2}{m}} \mathbf{W}(0) \right\|_2 & \lesssim L \sqrt{\log m}. \end{aligned}$$

Note that $\max_{l \in [L]} \|\widehat{\mathbf{W}}^l\|_2 \leq R \lesssim \sqrt{m}/(L^2 \sqrt{\log m})$, then by counting the number of $\widehat{\mathbf{W}}$, we have

$$\begin{aligned} \left\| \widehat{\mathbf{G}}_b^a(\mathbf{x}_i) \right\|_2 & \lesssim \sqrt{\frac{1}{m}} \left(L \sqrt{\log m} + \sum_{j=1}^{b-a} \binom{b-a}{j} \left(\frac{1}{L^2} \sqrt{\frac{1}{\log m}} \right)^j (L \sqrt{\log m})^{j+1} \right) \\ & = L \sqrt{\frac{\log m}{m}} \left(1 + \sum_{j=1}^{b-a} \binom{b-a}{j} \left(\frac{1}{L} \right)^j \right) \lesssim L \sqrt{\frac{\log m}{m}}. \end{aligned}$$

The proof is completed. □

Denote $\widetilde{\Sigma}(\mathbf{x})$, $\widetilde{h}^l(\mathbf{x})$, $\widetilde{\mathbf{G}}_b^a(\mathbf{x})$ as (10), (12), (13) when $\mathbf{W} = \widetilde{\mathbf{W}}$.

Lemma 13 (Claim 11.2 and Proposition 11.3 in Allen-Zhu et al. [2019]). *For any $\mathbf{W}, \widetilde{\mathbf{W}} \in \mathcal{B}_R(\mathbf{W}(0))$. There exists a series of diagonal matrices $\{(\Sigma'')^l \in \mathbb{R}^{m \times m}\}_{l \in [L]}$ with entries in $[-1, 1]$ such that for any $l \in [L]$, there holds*

$$\begin{aligned} (a) \quad & h^l(\mathbf{x}) - \widetilde{h}^l(\mathbf{x}) = \sum_{k=1}^l \left[\prod_{j=k+1}^l \sqrt{\frac{2}{m}} (\widetilde{\Sigma}^j(\mathbf{x}) + (\Sigma'')^j) \widetilde{\mathbf{W}}^j \right] \sqrt{\frac{2}{m}} (\widetilde{\Sigma}^k(\mathbf{x}) + (\Sigma'')^k) (\mathbf{W}^k - \widetilde{\mathbf{W}}^k) h^{k-1}(\mathbf{x}). \\ (b) \quad & \|(\Sigma'')^l\|_0 \leq \|\Sigma^l(\mathbf{x}) - \widetilde{\Sigma}^l(\mathbf{x})\|_0. \end{aligned}$$

Now we introduce the following Bernstein inequality under bounded distributions.

Lemma 14 (Theorem 2.8.4 in Vershynin [2018]). *Let $X_1, \dots, X_N$ be independent, mean-zero random variables, such that $|X_i| \leq K$ for all i . Then for every $t \geq 0$, we have*

$$\mathbb{P} \left(\left| \sum_{i=1}^N X_i \right| \geq t \right) \leq 2 \exp \left(- \frac{t^2/2}{\lambda^2 + Kt/3} \right),$$

where $\lambda^2 = \sum_{i=1}^N \mathbb{E} X_i^2$ is the sum of the variance.

Lemma 15. *Suppose $m \gtrsim L^{10} \log(nL/\delta) (\log m)^4 R^2$. Then with probability at least $1 - \delta$, for any $\mathbf{W} \in \mathcal{B}_R(\mathbf{W}(0))$, $i \in [n]$ and $l \in [L]$, there holds*

$$\|h^l(\mathbf{x}_i) - h_0^l(\mathbf{x}_i)\|_2 \lesssim L^2 \sqrt{\frac{\log m}{m}} R \text{ and } \|\Sigma^l(\mathbf{x}_i) - \Sigma_0^l(\mathbf{x}_i)\|_0 \lesssim L^{4/3} (\log m)^{1/3} (mR)^{2/3}. \quad (31)$$

Proof. We prove these two inequalities by induction. Note that (31) holds for $l = 0$. Now we suppose (31) holds for $l - 1$. Let $\kappa > 0$ be a constant. For $i \in [n]$ and $l \in [L]$, we define $A^l(\mathbf{x}_i) = \{r \in$

$[m] : \mathbb{I}\{\langle \mathbf{w}_r^l, h^{l-1}(\mathbf{x}_i) \rangle \geq 0\} \neq \mathbb{I}\{\langle \mathbf{w}_r^l(0), h_0^{l-1}(\mathbf{x}_i) \rangle \geq 0\}$, then $\|\Sigma^l(x_i) - \Sigma_0^l(x_i)\|_0 = |A^l(\mathbf{x}_i)|$. Furthermore, we decompose $A^l(\mathbf{x}_i)$ into two parts based on the behavior of $\mathbf{w}_r^l(0)$:

$$A_1^l(\mathbf{x}_i) = \{r \in A^l(\mathbf{x}_i) : |\langle \mathbf{w}_r^l(0), h_0^{l-1}(\mathbf{x}_i) \rangle| \leq \kappa\} \quad \text{and} \quad A_2^l(\mathbf{x}_i) = \{r \in A^l(\mathbf{x}_i) : |\langle \mathbf{w}_r^l(0), h_0^{l-1}(\mathbf{x}_i) \rangle| > \kappa\}.$$

We will control $|A_1^l(\mathbf{x}_i)|$ and $|A_2^l(\mathbf{x}_i)|$ respectively.

For $r \in [m]$, we define $F_{r,i}^l = \mathbb{I}\{|\langle \mathbf{w}_r^l(0), h_0^{l-1}(\mathbf{x}_i) \rangle| \leq \kappa\}$. From Lemma 6 we know that $2/3 \leq \|h_0^{l-1}(\mathbf{x}_i)\|_2^2 \leq 4/3$. Condition on $h_0^{l-1}(\mathbf{x}_i)$, $\langle \mathbf{w}_r^l(0), h_0^{l-1}(\mathbf{x}_i) \rangle \sim \mathcal{N}(0, \|h_0^{l-1}(\mathbf{x}_i)\|_2^2)$, then we have

$$\begin{aligned} \text{var}(F_{r,i}^l) &\leq \mathbb{E}(F_{r,i}^l)^2 = \mathbb{E}(F_{r,i}^l) = \mathbb{P}(-\kappa \leq \langle \mathbf{W}_r^l(0), h_0^{l-1}(\mathbf{x}_i) \rangle \leq \kappa) \\ &\leq \frac{3}{2\sqrt{\pi}} \int_{-\kappa}^{\kappa} e^{-3x^2/8} dx \leq C\kappa. \end{aligned}$$

Then by Lemma 14, choose $K = 1, t = mC\kappa, \lambda^2 \leq mC\kappa$, it then follows that

$$\mathbb{P}\left(\left|\sum_{r=1}^m F_{r,i}^l - m\mathbb{E}(F_{r,i}^l)\right| \leq mC\kappa|h_0^{l-1}(\mathbf{x}_i)|\right) \geq 1 - 2\exp\left(-\frac{(mC\kappa)^2/2}{mC\kappa + mC\kappa/3}\right).$$

Hence, taking union bounds over l, i , with probability at least $1 - CnL\exp(-m\kappa)$, there holds for all i, l ,

$$|A_1^l(\mathbf{x}_i)| \leq \sum_{r=1}^m F_{r,i}^l \lesssim m\kappa. \quad (32)$$

For $r \in A_2^l(\mathbf{x}_i)$, since $\mathbb{I}\{\langle \mathbf{w}_r^l, h^{l-1}(\mathbf{x}_i) \rangle \geq 0\} \neq \mathbb{I}\{\langle \mathbf{w}_r^l(0), h_0^{l-1}(\mathbf{x}_i) \rangle \geq 0\}$, we have

$$(\langle \mathbf{w}_r^l, h^{l-1}(\mathbf{x}_i) \rangle - \langle \mathbf{w}_r^l(0), h_0^{l-1}(\mathbf{x}_i) \rangle)^2 \geq |\langle \mathbf{w}_r^l(0), h_0^{l-1}(\mathbf{x}_i) \rangle|^2 > \kappa^2.$$

We deduce that

$$\begin{aligned} \|\mathbf{W}^l h^{l-1}(\mathbf{x}_i) - \mathbf{W}^l(0) h_0^{l-1}(\mathbf{x}_i)\|_2^2 &\geq \sum_{r \in A_2^l(\mathbf{x}_i)} (\langle \mathbf{w}_r^l, h^{l-1}(\mathbf{x}_i) \rangle - \langle \mathbf{w}_r^l(0), h_0^{l-1}(\mathbf{x}_i) \rangle)^2 \\ &> \sum_{r \in A_2^l(\mathbf{x}_i)} \kappa^2 = \kappa^2 |A_2^l(\mathbf{x}_i)|. \end{aligned} \quad (33)$$

By assumption $\|h^{l-1}(\mathbf{x}_i) - h_0^{l-1}(\mathbf{x}_i)\|_2 \lesssim L^2 R \sqrt{\log m/m}$ and Lemma 6, we get

$$\begin{aligned} &\|\mathbf{W}^l h^{l-1}(\mathbf{x}_i) - \mathbf{W}^l(0) h_0^{l-1}(\mathbf{x}_i)\|_2^2 \\ &\leq (\|\mathbf{W}^l - \mathbf{W}^l(0)\|_2 \|h^{l-1}(\mathbf{x}_i) - h_0^{l-1}(\mathbf{x}_i) + h_0^{l-1}(\mathbf{x}_i)\|_2 + \|\mathbf{W}^l(0)\|_2 \|h^{l-1}(\mathbf{x}_i) - h_0^{l-1}(\mathbf{x}_i)\|_2)^2 \\ &\lesssim (R(\|h^{l-1}(\mathbf{x}_i) - h_0^{l-1}(\mathbf{x}_i)\|_2 + 1) + \sqrt{m} \|h^{l-1}(\mathbf{x}_i) - h_0^{l-1}(\mathbf{x}_i)\|_2)^2 \lesssim L^4 R^2 \log m. \end{aligned}$$

Combined with (33), we have

$$|A_2^l(\mathbf{x}_i)| \lesssim \frac{L^4 R^2 \log m}{\kappa^2}. \quad (34)$$

From (32) and (34) we know that

$$\begin{aligned} \|\Sigma^l(\mathbf{x}_i) - \Sigma_0^l(\mathbf{x}_i)\|_0 &= |A^l(\mathbf{x}_i)| = |A_1^l(\mathbf{x}_i)| + |A_2^l(\mathbf{x}_i)| \\ &\lesssim m\kappa + \frac{L^4 R^2 \log m}{(\kappa)^2} \lesssim L^{4/3} (\log m)^{1/3} (mR)^{2/3}, \end{aligned}$$

where in the last inequality we choose $\kappa = L^{4/3} (\log m)^{1/3} R^{2/3} m^{-1/3}$. Hence, due to the overparameterization of m , we have with probability at least $1 - \delta$, for $i \in [n], l \in [L]$,

$$\|\Sigma^l(\mathbf{x}_i) - \Sigma_0^l(\mathbf{x}_i)\|_0 \lesssim L^{4/3} (\log m)^{1/3} (mR)^{2/3} \lesssim \frac{m}{L^2 \log m}.$$

Applying Lemma 13, we have

$$h^l(\mathbf{x}_i) - h_0^l(\mathbf{x}_i) = \sum_{k=1}^l \hat{\mathbf{G}}_{l,0}^k(\mathbf{x}_i)(\mathbf{W}^k - \mathbf{W}^k(0))h_0^{k-1}(\mathbf{x}_i), \quad (35)$$

where $\hat{\mathbf{G}}_{l,0}^k(\mathbf{x}_i)$ is defined as

$$\hat{\mathbf{G}}_{l,0}^k(\mathbf{x}_i) = \left[\prod_{j=k+1}^l \sqrt{\frac{2}{m}} (\Sigma^j(\mathbf{x}_i) + (\Sigma'')^j) \mathbf{W}^j \right] \sqrt{\frac{2}{m}} (\Sigma^k(\mathbf{x}_i) + (\Sigma'')^k). \quad (36)$$

It then follows that

$$\begin{aligned} & \|\Sigma^j(\mathbf{x}_i) + (\Sigma'')^j - \Sigma_0^j(\mathbf{x}_i)\|_0 \\ & \leq \|\Sigma^j(\mathbf{x}_i) - \Sigma_0^j(\mathbf{x}_i)\|_0 + \|(\Sigma'')^j\|_0 \leq 2\|\Sigma^j(\mathbf{x}_i) - \Sigma_0^j(\mathbf{x}_i)\|_0 \lesssim \frac{m}{L^2 \log m}. \end{aligned}$$

Our overparameterization requirement implies that $R \lesssim \sqrt{m}/(L^2 \sqrt{\log m})$. Hence, by Lemma 12, we have

$$\|\hat{\mathbf{G}}_{l,0}^k(\mathbf{x}_i)\|_2 \lesssim L \sqrt{\frac{\log m}{m}}. \quad (37)$$

Therefore,

$$\begin{aligned} & \|h^l(\mathbf{x}_i) - h_0^l(\mathbf{x}_i)\|_2 \\ & = \left\| \sum_{k=1}^l \hat{\mathbf{G}}_{l,0}^k(\mathbf{x}_i)(\mathbf{W}^k - \mathbf{W}^k(0))h_0^{k-1}(\mathbf{x}_i) \right\|_2 \\ & \lesssim \sum_{k=1}^l L \sqrt{\frac{\log m}{m}} R \|h_0^{k-1}(\mathbf{x}_i)\|_2 \lesssim L^2 \sqrt{\frac{\log m}{m}} R, \end{aligned}$$

where the last inequality results from Lemma 6. As a result, (31) holds for l . We have completed the proof of the lemma. $\square$

The above lemma and Lemma 6 imply that with probability at least $1 - \delta$, for all $l \in [L], i \in [n]$,

$$\|h^l(\mathbf{x}_i)\|_2 \lesssim L^2 \sqrt{\frac{\log m}{m}} R + 1 \lesssim 1. \quad (38)$$

Remark 10. Although our approach shares similarities with Lemma B.3 in Zou et al. [2018], our analysis significantly relaxes the required conditions. Specifically, we only require $R/\sqrt{m} = \tilde{O}(L^{-5})$, whereas their result demands the stricter scaling $R/\sqrt{m} = \tilde{O}(L^{-11})$.

Furthermore, compared to Lemma 8.2 in Allen-Zhu et al. [2019], they derive the bound $\|h^l(\mathbf{x}_i) - h_0^l(\mathbf{x}_i)\|_2 \lesssim RL^{5/2} \sqrt{\log m}/\sqrt{m}$, which is worse than our result by a factor of $\sqrt{L}$.

The following lemma shows the uniform concentration property of deep ReLU property, which is crucial in the generalization analysis.

Lemma 16. *Let $R \geq 1$ be a constant. Assume $m \gtrsim L^{11} d(\log m)^5 \log(L/\delta) R^2$. Then with probability at least $1 - \delta$, for $\mathbf{W} \in \mathcal{B}_R(\mathbf{W}(0))$, $l \in [L]$, we have*

$$\sup_{\mathbf{x} \in \mathcal{X}} \|h^l(\mathbf{x}) - h_0^l(\mathbf{x})\|_2 \lesssim L^2 \sqrt{\frac{\log m}{m}} R. \quad (39)$$

Proof. We consider the $1/(C^L \sqrt{m})$ -cover of S^{d-1} and denote it by $D = \{\mathbf{x}^1, \dots, \mathbf{x}^{|D|}\}$. By Lemma 4.2.13 in Vershynin [2018],

$$|D| \leq (1 + 2C^L \sqrt{m})^d.$$

Note that Lemma 15 holds for n points $\mathbf{x}_1, \dots, \mathbf{x}_n$. We consider $|D|$ points in D , and obtain that if $m \gtrsim L^{11} d(\log m)^5 \log(L/\delta) R^2$, then $m \gtrsim L^{10} \log(|D|L/\delta)(\log m)^4 R^2$. Hence,

$$\|h^l(\mathbf{x}^j) - h_0^l(\mathbf{x}^j)\|_2 \lesssim L^2 \sqrt{\frac{\log m}{m}} R, \quad 1 \leq j \leq |D|. \quad (40)$$

For any $\mathbf{x} \in \mathcal{X}$, there exists $\mathbf{x}^j \in D$ with $\|\mathbf{x} - \mathbf{x}^j\|_2 \leq 1/(C^L \sqrt{m})$. It then follows that

$$\begin{aligned} & \|h^l(\mathbf{x}) - h^l(\mathbf{x}^j)\|_2^2 \\ &= \frac{2}{m} \sum_{r=1}^m (\sigma(\langle \mathbf{w}_r^l, h^{l-1}(\mathbf{x}) \rangle) - \sigma(\langle \mathbf{w}_r^l, h^{l-1}(\mathbf{x}^j) \rangle))^2 \\ &\leq \frac{2}{m} \sum_{r=1}^m (\langle \mathbf{w}_r^l, h^{l-1}(\mathbf{x}) \rangle - \langle \mathbf{w}_r^l, h^{l-1}(\mathbf{x}^j) \rangle)^2 \\ &= \frac{2}{m} \|\mathbf{W}^l(h^{l-1}(\mathbf{x}) - h^{l-1}(\mathbf{x}^j))\|_2^2 \leq C \|h^{l-1}(\mathbf{x}) - h^{l-1}(\mathbf{x}^j)\|_2^2 \\ &\leq C^L \|\mathbf{x} - \mathbf{x}^j\|_2^2 \leq \frac{1}{m}, \end{aligned}$$

where the first inequality is due to $\sigma(\cdot)$ is 1-Lipschitz. In the second inequality we have used $\|\mathbf{W}^l\|_2 \leq \|\mathbf{W}^l(0)\|_2 + R \lesssim \sqrt{m}$ due to Lemma 2. Similarly, we derive that

$$\|h_0^l(\mathbf{x}) - h_0^l(\mathbf{x}^j)\|_2^2 \leq \frac{1}{m}.$$

Therefore, combined with (40), we have

$$\begin{aligned} & \|h^l(\mathbf{x}) - h_0^l(\mathbf{x})\|_2 \\ &\leq \|h^l(\mathbf{x}) - h^l(\mathbf{x}^j)\|_2 + \|h^l(\mathbf{x}^j) - h_0^l(\mathbf{x}^j)\|_2 + \|h_0^l(\mathbf{x}) - h_0^l(\mathbf{x}^j)\|_2 \\ &\lesssim \frac{1}{\sqrt{m}} + L^2 \sqrt{\frac{\log m}{m}} R + \frac{1}{\sqrt{m}} \lesssim L^2 \sqrt{\frac{\log m}{m}} R, \end{aligned}$$

where the last inequality results from $R \geq 1$.

The proof is completed. $\square$

Remark 9. Compared to prior work, while Allen-Zhu et al. [2019], Zou et al. [2018] only establishes bounds for the training data, we prove the uniform convergence over the entire input space. Previous work on uniform concentration demonstrated that $\sup_{\mathbf{x} \in \mathcal{X}} \|h^l(\mathbf{x}) - h_0^l(\mathbf{x})\|_2 \lesssim C^L R / \sqrt{m}$ [Xu and Zhu, 2024]. We present a significant improvement, reducing the dependence on L from exponential to polynomial.

Lemma 17. Suppose $m \gtrsim L^{10} \log(nL/\delta)(\log m)^4 R^2$. Then with probability at least $1 - \delta$ for all $\mathbf{W} \in \mathcal{B}_R(\mathbf{W}(0))$, $l \in [L]$, $i \in [n]$

$$\|\mathbf{a}^\top (\mathbf{G}_L^l(\mathbf{x}_i) - \mathbf{G}_{L,0}^l(\mathbf{x}_i))\|_2 \lesssim \frac{L^{5/3} (\log m)^{2/3} R^{1/3}}{m^{1/6}}$$

Proof. For the case $l = L$,

$$\begin{aligned} & \|\mathbf{a}^\top (\mathbf{G}_L^L(\mathbf{x}_i) - \mathbf{G}_{L,0}^L(\mathbf{x}_i))\|_2 = \sqrt{\frac{2}{m}} \|\mathbf{a}^\top (\Sigma^L(\mathbf{x}_i) - \Sigma_0^L(\mathbf{x}_i))\|_2 \\ &= \sqrt{\frac{2}{m}} \sqrt{\sum_{r=1}^m a_r^2 (\mathbb{I}\{\langle \mathbf{w}_r^L, h^{L-1}(\mathbf{x}_i) \rangle \geq 0\} - \mathbb{I}\{\langle \mathbf{w}_r^L(0), h_0^{L-1}(\mathbf{x}_i) \rangle \geq 0\})^2} \\ &= \sqrt{\frac{2}{m}} \sqrt{\sum_{r \in A^L(\mathbf{x}_i)} |a_r|} = \sqrt{\frac{2|A^L(\mathbf{x}_i)|}{m}} \lesssim \frac{L^{2/3} (\log m)^{1/6} R^{1/3}}{m^{1/6}}, \end{aligned} \quad (41)$$

where the last inequality is due to Lemma 15.

Now we suppose $l < L$, then

$$\begin{aligned}
& \mathbf{a}^\top (\mathbf{G}_L^l(\mathbf{x}_i) - \mathbf{G}_{L,0}^l(\mathbf{x}_i)) \\
&= \mathbf{a}^\top \sqrt{\frac{2}{m}} (\Sigma^L(\mathbf{x}_i) \mathbf{W}^L \mathbf{G}_{L-1}^l(\mathbf{x}_i) - \Sigma_0^L(\mathbf{x}_i) \mathbf{W}^L(0) \mathbf{G}_{L-1,0}^l(\mathbf{x}_i)) \\
&= \sqrt{\frac{2}{m}} \mathbf{a}^\top (\Sigma^L(\mathbf{x}_i) - \Sigma_0^L(\mathbf{x}_i)) \mathbf{W}^L \mathbf{G}_{L-1}^l(\mathbf{x}_i) + \sqrt{\frac{2}{m}} \mathbf{a}^\top \Sigma_0^L(\mathbf{x}_i) (\mathbf{W}^L - \mathbf{W}^L(0)) \mathbf{G}_{L-1}^l(\mathbf{x}_i) \\
&+ \sqrt{\frac{2}{m}} \mathbf{a}^\top \Sigma_0^L(\mathbf{x}_i) \mathbf{W}^L(0) (\mathbf{G}_{L-1}^l(\mathbf{x}_i) - \mathbf{G}_{L-1,0}^l(\mathbf{x}_i)) \\
&= \sqrt{\frac{2}{m}} (\mathbf{a}^\top (\Sigma^L(\mathbf{x}_i) - \Sigma_0^L(\mathbf{x}_i)) \mathbf{W}^L \mathbf{G}_{L-1}^l(\mathbf{x}_i) + \mathbf{a}^\top \Sigma_0^L(\mathbf{x}_i) (\mathbf{W}^L - \mathbf{W}^L(0)) \mathbf{G}_{L-1}^l(\mathbf{x}_i)),
\end{aligned}$$

where the last equality is according to Lemma 7. Applying Lemma 12 and Lemma 15, there holds

$$\|\mathbf{G}_{L-1}^l(\mathbf{x}_i)\|_2 \lesssim L \sqrt{\frac{\log m}{m}}.$$

This implies that

$$\begin{aligned}
& \|\mathbf{a}^\top (\mathbf{G}_L^l(\mathbf{x}_i) - \mathbf{G}_{L,0}^l(\mathbf{x}_i))\|_2 \\
&\lesssim \sqrt{\frac{2}{m}} (\|\mathbf{a}^\top (\Sigma^L(\mathbf{x}_i) - \Sigma_0^L(\mathbf{x}_i)) \mathbf{W}^L \mathbf{G}_{L-1}^l(\mathbf{x}_i)\|_2 + \|\mathbf{a}^\top \Sigma_0^L(\mathbf{x}_i) (\mathbf{W}^L - \mathbf{W}^L(0)) \mathbf{G}_{L-1}^l(\mathbf{x}_i)\|_2) \\
&\leq \sqrt{\frac{2}{m}} \|\mathbf{a}^\top (\Sigma^L(\mathbf{x}_i) - \Sigma_0^L(\mathbf{x}_i))\|_2 \|\mathbf{W}^L\|_2 \|\mathbf{G}_{L-1}^l(\mathbf{x}_i)\|_2 \\
&+ \sqrt{\frac{2}{m}} \|\mathbf{a}\|_2 \|\Sigma_0^L(\mathbf{x}_i)\|_2 \|\mathbf{W}^L - \mathbf{W}^L(0)\|_2 \|\mathbf{G}_{L-1}^l(\mathbf{x}_i)\|_2 \\
&\lesssim \frac{L^{2/3} (\log m)^{1/6} R^{1/3}}{m^{1/6}} \sqrt{m} L \sqrt{\frac{\log m}{m}} + L \sqrt{\frac{\log m}{m}} R \\
&\lesssim \frac{L^{5/3} (\log m)^{2/3} R^{1/3}}{m^{1/6}},
\end{aligned}$$

where we have used (41) in the third inequality. The proof is completed. $\square$

Lemma 18. Assume $m \gtrsim L^{10} \log(nL/\delta) (\log m)^4 R^2$. Then with probability at least $1 - \delta$, for any $\mathbf{W} \in \mathcal{B}_R(\mathbf{W}(0))$, $i \in [n]$ and $l \in [L]$, there holds

$$\left\| \frac{\partial f_{\mathbf{W}}(\mathbf{x}_i)}{\partial \mathbf{W}^l} - \frac{\partial f_{\mathbf{W}(0)}(\mathbf{x}_i)}{\partial \mathbf{W}^l(0)} \right\|_F \lesssim \frac{L^{5/3} (\log m)^{2/3} R^{1/3}}{m^{1/6}}. \quad (42)$$

Proof. Since $\|xy^\top\|_F = \|x\|_2 \|y\|_2$ for two vectors x, y , we have

$$\begin{aligned}
& \left\| \frac{\partial f_{\mathbf{W}}(\mathbf{x}_i)}{\partial \mathbf{W}^l} - \frac{\partial f_{\mathbf{W}(0)}(\mathbf{x}_i)}{\partial \mathbf{W}^l(0)} \right\|_F \\
&= \|h^{l-1}(\mathbf{x}_i) \mathbf{a}^\top \mathbf{G}_L^l(\mathbf{x}_i) - h_0^{l-1}(\mathbf{x}_i) \mathbf{a}^\top \mathbf{G}_{L,0}^l(\mathbf{x}_i)\|_F \\
&\leq \|h^{l-1}(\mathbf{x}_i) \mathbf{a}^\top (\mathbf{G}_L^l(\mathbf{x}_i) - \mathbf{G}_{L,0}^l(\mathbf{x}_i))\|_F + \|(h^{l-1}(\mathbf{x}_i) - h_0^{l-1}(\mathbf{x}_i)) \mathbf{a}^\top \mathbf{G}_{L,0}^l(\mathbf{x}_i)\|_F \\
&= \|h^{l-1}(\mathbf{x}_i)\|_2 \|\mathbf{a}^\top (\mathbf{G}_L^l(\mathbf{x}_i) - \mathbf{G}_{L,0}^l(\mathbf{x}_i))\|_2 + \|h^{l-1}(\mathbf{x}_i) - h_0^{l-1}(\mathbf{x}_i)\|_2 \|\mathbf{a}^\top \mathbf{G}_{L,0}^l(\mathbf{x}_i)\|_2.
\end{aligned}$$

Using (38) and Lemma 17, we have

$$\|h^{l-1}(\mathbf{x}_i)\|_2 \|\mathbf{a}^\top (\mathbf{G}_L^l(\mathbf{x}_i) - \mathbf{G}_{L,0}^l(\mathbf{x}_i))\|_2 \lesssim \frac{L^{5/3} (\log m)^{2/3} R^{1/3}}{m^{1/6}}.$$

Applying Lemma 15 and (24), we obtain

$$\|h^{l-1}(\mathbf{x}_i) - h_0^{l-1}(\mathbf{x}_i)\|_2 \|\mathbf{a}^\top \mathbf{G}_{L,0}^l(\mathbf{x}_i)\|_2 \lesssim L^2 \sqrt{\frac{\log m}{m}} R \sqrt{m} L \sqrt{\frac{\log m}{m}} = \frac{L^3 R \log m}{\sqrt{m}}.$$

It then follows that

$$\left\| \frac{\partial f_{\mathbf{W}}(\mathbf{x}_i)}{\partial \mathbf{W}^l} - \frac{\partial f_{\mathbf{W}(0)}(\mathbf{x}_i)}{\partial \mathbf{W}^l(0)} \right\|_F \lesssim \frac{L^{5/3} (\log m)^{2/3} R^{1/3}}{m^{1/6}} + \frac{L^3 R \log m}{\sqrt{m}} \lesssim \frac{L^{5/3} (\log m)^{2/3} R^{1/3}}{m^{1/6}}.$$

The proof is completed. $\square$

B Proofs for Optimization

Lemma 19. Suppose $m \gtrsim L^{10} \log(nL/\delta)(\log m)^4 R^2$, then with probability at least $1 - \delta$, for $i \in [n]$, $\widetilde{\mathbf{W}}, \mathbf{W} \in \mathcal{B}_R(\mathbf{W}(0))$, we have

$$\left| f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i) - f_{\mathbf{W}}(\mathbf{x}_i) - \left\langle \frac{\partial f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i)}{\partial \widetilde{\mathbf{W}}}, \widetilde{\mathbf{W}} - \mathbf{W} \right\rangle \right| \lesssim \frac{L^{8/3} R^{4/3} (\log m)^{2/3}}{m^{1/6}}.$$

The left term plays a key role in the optimization analysis. Note that it is slightly different from the following term:

$$f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i) - f_{\mathbf{W}}(\mathbf{x}_i) - \left\langle \frac{\partial f_{\mathbf{W}}(\mathbf{x}_i)}{\partial \mathbf{W}}, \widetilde{\mathbf{W}} - \mathbf{W} \right\rangle,$$

which has been widely studied in the literature (Allen-Zhu et al. [2019], Zou et al. [2018]). We could get similar results by combining it with Lemma 18. Moreover, previous works derived the bound $\tilde{O}\left(\frac{L^3 R^{4/3}}{m^{1/6}}\right)$ [Cao and Gu, 2019].

Proof. Note that

$$\begin{aligned} \left\langle \frac{\partial f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i)}{\partial \widetilde{\mathbf{W}}}, \widetilde{\mathbf{W}} - \mathbf{W} \right\rangle &= \sum_{l=1}^L \left\langle \frac{\partial f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i)}{\partial \widetilde{\mathbf{W}}^l}, \widetilde{\mathbf{W}}^l - \mathbf{W}^l \right\rangle \\ &= \sum_{l=1}^L \left\langle (\tilde{\mathbf{G}}_L^l(\mathbf{x}_i))^\top \mathbf{a}(\tilde{h}^{l-1}(\mathbf{x}_i))^\top, \widetilde{\mathbf{W}}^l - \mathbf{W}^l \right\rangle \\ &= \sum_{l=1}^L \mathbf{a}^\top \tilde{\mathbf{G}}_L^l(\mathbf{x}_i) (\widetilde{\mathbf{W}}^l - \mathbf{W}^l) \tilde{h}^{l-1}(\mathbf{x}_i). \end{aligned}$$

Since $f_{\mathbf{W}}(\mathbf{x}_i) = \mathbf{a}^\top h^L(\mathbf{x}_i)$, applying Lemma 13, we obtain

$$\begin{aligned} f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i) - f_{\mathbf{W}}(\mathbf{x}_i) &= \mathbf{a}^\top (\tilde{h}^L(\mathbf{x}_i) - h^L(\mathbf{x}_i)) \\ &= \sum_{l=1}^L \left[\prod_{j=l+1}^L \sqrt{\frac{2}{m}} (\Sigma^j(\mathbf{x}_i) + (\Sigma'')^j) \mathbf{W}^j \right] \sqrt{\frac{2}{m}} (\Sigma^l(\mathbf{x}_i) + (\Sigma'')^l) (\widetilde{\mathbf{W}}^l - \mathbf{W}^l) \tilde{h}^{l-1}(\mathbf{x}_i) \end{aligned}$$

with $\|(\Sigma'')^l\|_0 \leq \|\Sigma^l(\mathbf{x}_i) - \tilde{\Sigma}^l(\mathbf{x}_i)\|_0$. Then

$$\begin{aligned} \|\Sigma^l(\mathbf{x}_i) + (\Sigma'')^l - \Sigma_0^l(\mathbf{x}_i)\|_0 &\leq \|\tilde{\Sigma}^l(\mathbf{x}_i) - \Sigma_0^l(\mathbf{x}_i)\|_0 + 2\|\Sigma^l(\mathbf{x}_i) - \Sigma_0^l(\mathbf{x}_i)\|_0 \\ &\lesssim L^{4/3} (\log m)^{1/3} (mR)^{\frac{2}{3}}, \end{aligned}$$

the last inequality is due to Lemma 15. We further let

$$\hat{\mathbf{G}}_L^l(\mathbf{x}_i) = \left[\prod_{j=l+1}^L \sqrt{\frac{2}{m}} (\Sigma^j(\mathbf{x}_i) + (\Sigma'')^j) \mathbf{W}^j \right] \sqrt{\frac{2}{m}} (\Sigma^l(\mathbf{x}_i) + (\Sigma'')^l).$$

Following the proof of Lemma 17, we have

$$\|\mathbf{a}^\top (\hat{\mathbf{G}}_L^l(\mathbf{x}_i) - \mathbf{G}_{L,0}^l(\mathbf{x}_i))\|_2 \leq \frac{L^{5/3} (\log m)^{2/3} R^{1/3}}{m^{1/6}},$$

which implies that

$$\begin{aligned} \|\mathbf{a}^\top (\hat{\mathbf{G}}_L^l(\mathbf{x}_i) - \tilde{\mathbf{G}}_L^l(\mathbf{x}_i))\|_2 &\leq \|\mathbf{a}^\top (\hat{\mathbf{G}}_L^l(\mathbf{x}_i) - \mathbf{G}_{L,0}^l(\mathbf{x}_i))\|_2 + \|\mathbf{a}^\top (\mathbf{G}_{L,0}^l(\mathbf{x}_i) - \tilde{\mathbf{G}}_L^l(\mathbf{x}_i))\|_2 \\ &\lesssim \frac{L^{5/3} (\log m)^{2/3} R^{1/3}}{m^{1/6}}. \end{aligned}$$

Hence,

$$\begin{aligned}
& \left| f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i) - f_{\mathbf{W}}(\mathbf{x}_i) - \left\langle \frac{\partial f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i)}{\partial \widetilde{\mathbf{W}}}, \widetilde{\mathbf{W}} - \mathbf{W} \right\rangle \right| \\
&= \left| \sum_{l=1}^L \mathbf{a}^\top (\widehat{\mathbf{G}}_L^l(\mathbf{x}_i) - \widetilde{\mathbf{G}}_L^l(\mathbf{x}_i)) (\widetilde{\mathbf{W}}^l - \mathbf{W}^l) \tilde{h}^{l-1}(\mathbf{x}_i) \right| \\
&\leq \sum_{l=1}^L \|\mathbf{a}^\top (\widehat{\mathbf{G}}_L^l(\mathbf{x}_i) - \widetilde{\mathbf{G}}_L^l(\mathbf{x}_i))^\top\|_2 \|\widetilde{\mathbf{W}}^l - \mathbf{W}^l\|_2 \|\tilde{h}^{l-1}(\mathbf{x}_i)\|_2 \\
&\lesssim L \frac{L^{5/3}(\log m)^{2/3} R^{1/3}}{m^{1/6}} R = \frac{L^{8/3} R^{4/3} (\log m)^{2/3}}{m^{1/6}},
\end{aligned}$$

where in the last inequality we have used (38). The proof is completed. $\square$

Lemma 20. Suppose $m \gtrsim L^{10} \log(nL/\delta)(\log m)^4 R^2$, then with probability at least $1 - \delta$, we have for $\widetilde{\mathbf{W}}, \mathbf{W} \in \mathcal{B}_R(\mathbf{W}(0))$,

$$\begin{aligned}
\left\langle \widetilde{\mathbf{W}} - \mathbf{W}, \frac{\partial \mathcal{L}_S(\widetilde{\mathbf{W}})}{\partial \widetilde{\mathbf{W}}} \right\rangle &\geq \mathcal{L}_S(\widetilde{\mathbf{W}}) - \mathcal{L}_S(\mathbf{W}) + \frac{2}{n} \sum_{i=1}^n (l'(y_i f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i)) - l'(y_i f_{\mathbf{W}}(\mathbf{x}_i)))^2 \\
&\quad - \frac{CL^{8/3}(\log m)^{2/3} R^{4/3}}{m^{1/6}} \mathcal{L}_S(\widetilde{\mathbf{W}}).
\end{aligned}$$

Proof. Since ℓ is $1/4$ -smooth, it enjoys the co-coercivity, i.e., $\ell(a) \geq \ell(b) + (a-b)\ell'(b) + 2(\ell'(a) - \ell'(b))^2$, which implies that

$$\begin{aligned}
& y_i \ell'(y_i f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i)) (f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i) - f_{\mathbf{W}}(\mathbf{x}_i)) \\
&\geq \ell(y_i f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i)) - \ell(y_i f_{\mathbf{W}}(\mathbf{x}_i)) + 2(\ell'(y_i f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i)) - \ell'(y_i f_{\mathbf{W}}(\mathbf{x}_i)))^2.
\end{aligned}$$

We combine the above inequality with Lemma 19 and obtain

$$\begin{aligned}
& \left\langle \widetilde{\mathbf{W}} - \mathbf{W}, \frac{\partial \mathcal{L}_S(\widetilde{\mathbf{W}})}{\partial \widetilde{\mathbf{W}}} \right\rangle \\
&= \frac{1}{n} \sum_{i=1}^n \left\langle \widetilde{\mathbf{W}} - \mathbf{W}, \frac{\partial f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i)}{\partial \widetilde{\mathbf{W}}} \right\rangle y_i \ell'(y_i f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i)) \\
&= \frac{1}{n} \sum_{i=1}^n y_i \ell'(y_i f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i)) \left(f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i) - f_{\mathbf{W}}(\mathbf{x}_i) - \left(f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i) - f_{\mathbf{W}}(\mathbf{x}_i) - \left\langle \frac{\partial f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i)}{\partial \widetilde{\mathbf{W}}}, \widetilde{\mathbf{W}} - \mathbf{W} \right\rangle \right) \right) \\
&\geq \frac{1}{n} \sum_{i=1}^n (\ell(y_i f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i)) - \ell(y_i f_{\mathbf{W}}(\mathbf{x}_i)) + 2(\ell'(y_i f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i)) - \ell'(y_i f_{\mathbf{W}}(\mathbf{x}_i)))^2) \\
&\quad - \frac{1}{n} \sum_{i=1}^n \left| f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i) - f_{\mathbf{W}}(\mathbf{x}_i) - \left\langle \frac{\partial f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i)}{\partial \widetilde{\mathbf{W}}}, \widetilde{\mathbf{W}} - \mathbf{W} \right\rangle \right| \ell(y_i f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i)) \\
&\geq \mathcal{L}_S(\widetilde{\mathbf{W}}) - \mathcal{L}_S(\mathbf{W}) + \frac{2}{n} \sum_{i=1}^n (l'(y_i f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i)) - l'(y_i f_{\mathbf{W}}(\mathbf{x}_i)))^2 \\
&\quad - CL^{8/3}(\log m)^{2/3} R^{4/3} m^{-1/6} \mathcal{L}_S(\widetilde{\mathbf{W}}),
\end{aligned}$$

where in the first inequality we have used $|y_i \ell'(y_i f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i))| \leq \ell(y_i f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i))$. $\square$

Lemma 21. Suppose $m \gtrsim L^{10} \log(nL/\delta)(\log m)^4 R^2$. Then with probability at least $1 - \delta$, for $\eta \leq 4/(5L)$ and $\widetilde{\mathbf{W}}, \mathbf{W} \in \mathcal{B}_R(\mathbf{W}(0))$,

$$\begin{aligned}
\left\| \mathbf{W} - \eta \frac{\partial \mathcal{L}_S(\mathbf{W})}{\partial \mathbf{W}} - \widetilde{\mathbf{W}} \right\|_F^2 &\leq \|\mathbf{W} - \widetilde{\mathbf{W}}\|_F^2 - 2\eta(\mathcal{L}_S(\mathbf{W}) - \mathcal{L}_S(\widetilde{\mathbf{W}})) \\
&\quad + 2\eta CL^{8/3}(\log m)^{2/3} R^{4/3} m^{-1/6} \mathcal{L}_S(\mathbf{W}) + 20\eta^2 L \tilde{F}_S^2(\widetilde{\mathbf{W}}).
\end{aligned}$$

Proof. By Lemma 8 and Lemma 18 we know that $\left\| \frac{\partial f_{\mathbf{W}}(\mathbf{x}_i)}{\partial \mathbf{W}^l} \right\|_F \leq 2$, hence $\left\| \frac{\partial f_{\mathbf{W}}(\mathbf{x}_i)}{\partial \mathbf{W}} \right\|_F \leq 2\sqrt{L}$, which implies that

$$\begin{aligned} \left\| \frac{\partial \mathcal{L}_S(\mathbf{W})}{\partial \mathbf{W}} \right\|_F^2 &= \left\| \frac{1}{n} \sum_{i=1}^n y_i \ell'(y_i f_{\mathbf{W}}(\mathbf{x}_i)) \frac{\partial f_{\mathbf{W}}(\mathbf{x}_i)}{\partial \mathbf{W}} \right\|_F^2 \\ &\leq \left(\frac{1}{n} \sum_{i=1}^n |\ell'(y_i f_{\mathbf{W}}(\mathbf{x}_i))| \left\| \frac{\partial f_{\mathbf{W}}(\mathbf{x}_i)}{\partial \mathbf{W}} \right\|_F \right)^2 \\ &\leq 4L \left(\frac{1}{n} \sum_{i=1}^n |\ell'(y_i f_{\mathbf{W}}(\mathbf{x}_i))| \right)^2 \\ &\leq 5L \left(\frac{1}{n} \sum_{i=1}^n (|\ell'(y_i f_{\mathbf{W}}(\mathbf{x}_i))| - |\ell'(y_i f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i))|) \right)^2 + 20L \left(\frac{1}{n} \sum_{i=1}^n |\ell'(y_i f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i))| \right)^2, \end{aligned}$$

where we have used the standard inequality $(a+b)^2 \leq 5a^2 + 5b^2/4$. Then by applying Lemma 20 we have

$$\begin{aligned} &\left\| \mathbf{W} - \eta \frac{\partial \mathcal{L}_S(\mathbf{W})}{\partial \mathbf{W}} - \widetilde{\mathbf{W}} \right\|_F^2 \\ &= \|\mathbf{W} - \widetilde{\mathbf{W}}\|_F^2 + \eta^2 \left\| \frac{\partial \mathcal{L}_S(\mathbf{W})}{\partial \mathbf{W}} \right\|_F^2 - 2\eta \left\langle \mathbf{W} - \widetilde{\mathbf{W}}, \frac{\partial \mathcal{L}_S(\mathbf{W})}{\partial \mathbf{W}} \right\rangle \\ &\leq \|\mathbf{W} - \widetilde{\mathbf{W}}\|_F^2 + 5\eta^2 L \left(\frac{1}{n} \sum_{i=1}^n (|\ell'(y_i f_{\mathbf{W}}(\mathbf{x}_i))| - |\ell'(y_i f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i))|) \right)^2 \\ &\quad + 20\eta^2 L \left(\frac{1}{n} \sum_{i=1}^n |\ell'(y_i f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i))| \right)^2 - 2\eta (\mathcal{L}_S(\mathbf{W}) - \mathcal{L}_S(\widetilde{\mathbf{W}})) - \frac{4\eta}{n} \sum_{i=1}^n (l'(y_i f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i)) - l'(y_i f_{\mathbf{W}}(\mathbf{x}_i)))^2 \\ &\quad + \eta CL^{8/3} (\log m)^{2/3} R^{4/3} m^{-1/6} \mathcal{L}_S(\mathbf{W}). \end{aligned}$$

Since $\eta \leq 4/(5L)$ and

$$\left(\frac{1}{n} \sum_{i=1}^n (|\ell'(y_i f_{\mathbf{W}}(\mathbf{x}_i))| - |\ell'(y_i f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i))|) \right)^2 \leq \frac{1}{n} \sum_{i=1}^n (l'(y_i f_{\widetilde{\mathbf{W}}}(\mathbf{x}_i)) - l'(y_i f_{\mathbf{W}}(\mathbf{x}_i)))^2,$$

the proof is completed. $\square$

Proof of Theorem 1 We prove it through induction. It holds for $t = 0$. Suppose it holds for $0, \dots, t-1$. Then for $l \in [L]$,

$$\begin{aligned} \|\mathbf{W}^l(t) - \mathbf{W}^l(0)\|_2 &\leq \|\mathbf{W}^l(t) - \mathbf{W}^l(0)\|_F \\ &\leq \sqrt{\|\mathbf{W}(t) - \overline{\mathbf{W}}\|_F^2} + \sqrt{\|\mathbf{W}(0) - \overline{\mathbf{W}}\|_F^2} \leq 2\sqrt{F_S(\overline{\mathbf{W}})}. \end{aligned}$$

Thus, from our overparameterization of m and plugging $R = 2\sqrt{F_S(\overline{\mathbf{W}})}$ into Lemma 21, we have

$$\begin{aligned} \|\mathbf{W}(t) - \overline{\mathbf{W}}\|_F^2 &\leq \|\mathbf{W}(t-1) - \overline{\mathbf{W}}\|_F^2 - 2\eta (\mathcal{L}_S(\mathbf{W}(t-1)) - \mathcal{L}_S(\overline{\mathbf{W}})) \\ &\quad + 2\eta CL^{8/3} (\log m)^{2/3} F_S^{2/3}(\overline{\mathbf{W}}) m^{-1/6} \mathcal{L}_S(\mathbf{W}(t-1)) + 20\eta^2 L \tilde{F}_S^2(\overline{\mathbf{W}}). \end{aligned}$$

Telescoping and note that $\tilde{F}_S(\overline{\mathbf{W}}) \leq \mathcal{L}_S(\overline{\mathbf{W}})$, we obtain

$$\begin{aligned} \|\mathbf{W}(t) - \overline{\mathbf{W}}\|_F^2 &+ 2\eta \sum_{k=0}^{t-1} (\mathcal{L}_S(\mathbf{W}(k)) - \mathcal{L}_S(\overline{\mathbf{W}})) \leq \|\mathbf{W}(0) - \overline{\mathbf{W}}\|_F^2 \\ &+ 2\eta CL^{8/3} (\log m)^{2/3} F_S^{2/3}(\overline{\mathbf{W}}) m^{-1/6} \sum_{k=0}^{t-1} \mathcal{L}_S(\mathbf{W}(k)) + 20\eta^2 L T \tilde{F}_S(\overline{\mathbf{W}}) \mathcal{L}_S(\overline{\mathbf{W}}), \end{aligned}$$

which implies

$$\begin{aligned} \|\mathbf{W}(t) - \bar{\mathbf{W}}\|_F^2 + 2\eta \sum_{k=0}^{t-1} (\mathcal{L}_S(\mathbf{W}(k))(1 - CL^{8/3}(\log m)^{2/3} F_S^{2/3}(\bar{\mathbf{W}})m^{-1/6}) \\ \leq \|\mathbf{W}(0) - \bar{\mathbf{W}}\|_F^2 + (2 + 20\eta L \tilde{F}_S(\bar{\mathbf{W}}))\eta T \mathcal{L}_S(\bar{\mathbf{W}}). \end{aligned}$$

Hence, when $m \gtrsim F_S^4(\bar{\mathbf{W}})(\log m)^4 L^{16}$, $\eta \leq 1/(20L\tilde{F}_S(\bar{\mathbf{W}}))$, there holds

$$\|\mathbf{W}(t) - \bar{\mathbf{W}}\|_F^2 + \eta \sum_{k=0}^{t-1} (\mathcal{L}_S(\mathbf{W}(k))) \leq F_S(\bar{\mathbf{W}}).$$

The proof is completed.

C Proofs for Generalization

From (35), we know that there exists $\hat{\mathbf{G}}_{L,0}^l(\mathbf{x}_i)$, $1 \leq l \leq L, i \in [n]$, such that

$$h^L(\mathbf{x}_i) - h_0^L(\mathbf{x}_i) = \sum_{l=1}^L \hat{\mathbf{G}}_{L,0}^l(\mathbf{x}_i)(\mathbf{W}^l - \mathbf{W}^l(0))h_0^{l-1}(\mathbf{x}_i).$$

Let E_1 be the event (w.r.t. $\mathbf{W}(0)$) that $\|\hat{\mathbf{G}}_{L,0}^l(\mathbf{x}_i)\|_2 \leq CL\sqrt{\log m/m}$ for all $\mathbf{W} \in \mathcal{W}_1$. Let E_2 be the event such that $\|h_0^{l-1}(\mathbf{x}_i)\|_2^2 \leq 4/3$, $1 \leq l \leq L, i \in [n]$. Then we have the following bound on Rademacher complexity:

Lemma 22. *Let $\mathcal{F}$ and $\mathcal{W}_1$ be defined in (7) and (8), respectively. If the events E_1, E_2 hold, then*

$$\mathfrak{R}_{S_1,n}(\mathcal{F}) \leq CL^2 \sqrt{\frac{F(\bar{\mathbf{W}}) \log m}{n}}, \quad (43)$$

where

$$\mathfrak{R}_{S_1,n}(\mathcal{F}) = \sup_{\tilde{S} \subset S_1: |\tilde{S}|=n} \mathfrak{R}_{\tilde{S}}(\mathcal{F}).$$

Proof. Let $\tilde{S} = \{\tilde{\mathbf{x}}_1, \dots, \tilde{\mathbf{x}}_n\}$. Then we have

$$\begin{aligned} \mathfrak{R}_{\tilde{S}}(\mathcal{F}) &= \mathbb{E}_\epsilon \left[\sup_{\mathbf{W} \in \mathcal{W}_1} \frac{1}{n} \sum_{i=1}^n \epsilon_i f_{\mathbf{W}}(\tilde{\mathbf{x}}_i) \right] \\ &\leq \mathbb{E}_\epsilon \left[\sup_{\mathbf{W} \in \mathcal{W}_1} \frac{1}{n} \sum_{i=1}^n \epsilon_i (f_{\mathbf{W}}(\tilde{\mathbf{x}}_i) - f_{\mathbf{W}(0)}(\tilde{\mathbf{x}}_i)) \right] + \mathbb{E}_\epsilon \left[\sup_{\mathbf{W} \in \mathcal{W}_1} \frac{1}{n} \sum_{i=1}^n \epsilon_i f_{\mathbf{W}(0)}(\tilde{\mathbf{x}}_i) \right] \\ &= \mathbb{E}_\epsilon \left[\sup_{\mathbf{W} \in \mathcal{W}_1} \frac{1}{n} \sum_{i=1}^n \epsilon_i (f_{\mathbf{W}}(\tilde{\mathbf{x}}_i) - f_{\mathbf{W}(0)}(\tilde{\mathbf{x}}_i)) \right] = \mathbb{E}_\epsilon \left[\sup_{\mathbf{W} \in \mathcal{W}_1} \frac{1}{n} \sum_{i=1}^n \epsilon_i \mathbf{a}^\top (h^L(\tilde{\mathbf{x}}_i) - h_0^L(\tilde{\mathbf{x}}_i)) \right] \\ &= \frac{1}{n} \mathbb{E}_\epsilon \left[\sup_{\mathbf{W} \in \mathcal{W}_1} \mathbf{a}^\top \sum_{i=1}^n \epsilon_i (h^L(\tilde{\mathbf{x}}_i) - h_0^L(\tilde{\mathbf{x}}_i)) \right] \\ &\leq \frac{\|\mathbf{a}\|_2}{n} \mathbb{E}_\epsilon \sup_{\mathbf{W} \in \mathcal{W}_1} \left\| \sum_{i=1}^n \epsilon_i (h^L(\tilde{\mathbf{x}}_i) - h_0^L(\tilde{\mathbf{x}}_i)) \right\|_2, \end{aligned} \quad (44)$$

where the last inequality is due to Cauchy-Schwartz inequality. Since the event E_1 holds, we have for all $\mathbf{W} \in \mathcal{W}_1$, $1 \leq k \leq l \leq L, i \in [n]$,

$$\|\hat{\mathbf{G}}_{L,0}^l(\tilde{\mathbf{x}}_i)(\mathbf{W}^l - \mathbf{W}^l(0))\|_F \leq \|\hat{\mathbf{G}}_{L,0}^l(\tilde{\mathbf{x}}_i)\|_2 \|\mathbf{W}^l - \mathbf{W}^l(0)\|_F \leq CL \sqrt{\frac{F(\bar{\mathbf{W}}) \log m}{m}},$$

where we have used $\|AB\|_F \leq \|A\|_2 \|B\|_F$ for two matrices A, B . Therefore,

$$\begin{aligned}
& \mathbb{E}_\epsilon \sup_{\mathbf{W} \in \mathcal{W}_1} \left\| \sum_{i=1}^n \epsilon_i (h^L(\tilde{\mathbf{x}}_i) - h_0^L(\tilde{\mathbf{x}}_i)) \right\|_2 \\
&= \mathbb{E}_\epsilon \sup_{\mathbf{W} \in \mathcal{W}_1} \left\| \sum_{i=1}^n \epsilon_i \sum_{l=1}^L \hat{\mathbf{G}}_{L,0}^l(\tilde{\mathbf{x}}_i) (\mathbf{W}^l - \mathbf{W}^l(0)) h_0^{l-1}(\tilde{\mathbf{x}}_i) \right\|_2 \\
&\leq \sum_{l=1}^L \mathbb{E}_\epsilon \sup_{\|\hat{\mathbf{G}}_{L,0}^l(\tilde{\mathbf{x}}_i) (\mathbf{W}^l - \mathbf{W}^l(0))\|_F \leq CL\sqrt{F(\bar{\mathbf{W}}) \log m/m}} \left\| \sum_{i=1}^n \epsilon_i \hat{\mathbf{G}}_{L,0}^l(\tilde{\mathbf{x}}_i) (\mathbf{W}^l - \mathbf{W}^l(0)) h_0^{l-1}(\tilde{\mathbf{x}}_i) \right\|_2.
\end{aligned}$$

Letting $g_1^l, \dots, g_m^l$ be the rows of matrix $\hat{\mathbf{G}}_{L,0}^l(\tilde{\mathbf{x}}_i) (\mathbf{W}^l - \mathbf{W}^l(0))$, we obtain

$$\left\| \sum_{i=1}^n \epsilon_i \hat{\mathbf{G}}_{L,0}^l(\tilde{\mathbf{x}}_i) (\mathbf{W}^l - \mathbf{W}^l(0)) h_0^{l-1}(\tilde{\mathbf{x}}_i) \right\|_2^2 = \sum_{r=1}^m \|g_r^l\|_2^2 \left(\sum_{i=1}^n \epsilon_i \frac{(g_r^l)^\top}{\|g_r^l\|_2} h_0^{l-1}(\tilde{\mathbf{x}}_i) \right)^2.$$

The supremum of this over all $g_1^l, \dots, g_m^l$ must be obtained when $\|g_r^l\|_2 = CL\sqrt{F(\bar{\mathbf{W}}) \log m/m}$ for some r . Hence

$$\begin{aligned}
& \mathbb{E}_\epsilon \sup_{\|\hat{\mathbf{G}}_{L,0}^l(\tilde{\mathbf{x}}_i) (\mathbf{W}^l - \mathbf{W}^l(0))\|_F \leq CL\sqrt{F(\bar{\mathbf{W}}) \log m/m}} \left\| \sum_{i=1}^n \epsilon_i \hat{\mathbf{G}}_{L,0}^l(\tilde{\mathbf{x}}_i) (\mathbf{W}^l - \mathbf{W}^l(0)) h_0^{l-1}(\tilde{\mathbf{x}}_i) \right\|_2 \\
&= CL\sqrt{\frac{F(\bar{\mathbf{W}}) \log m}{m}} \mathbb{E}_\epsilon \left\| \sum_{i=1}^n \epsilon_i h_0^{l-1}(\tilde{\mathbf{x}}_i) \right\|_2.
\end{aligned}$$

This implies that

$$\begin{aligned}
& \mathbb{E}_\epsilon \sup_{\mathbf{W} \in \mathcal{W}_1} \left\| \sum_{i=1}^n \epsilon_i (h^L(\tilde{\mathbf{x}}_i) - h_0^L(\tilde{\mathbf{x}}_i)) \right\|_2 \\
&\leq CL\sqrt{\frac{F(\bar{\mathbf{W}}) \log m}{m}} \sum_{l=1}^L \mathbb{E}_\epsilon \left\| \sum_{i=1}^n \epsilon_i h_0^{l-1}(\tilde{\mathbf{x}}_i) \right\|_2 \\
&\leq CL\sqrt{\frac{F(\bar{\mathbf{W}}) \log m}{m}} \sum_{l=1}^L \sqrt{\mathbb{E}_\epsilon \left\| \sum_{i=1}^n \epsilon_i h_0^{l-1}(\tilde{\mathbf{x}}_i) \right\|_2^2} \\
&= CL\sqrt{\frac{F(\bar{\mathbf{W}}) \log m}{m}} \sum_{l=1}^L \sqrt{\sum_{i=1}^n \|h_0^{l-1}(\tilde{\mathbf{x}}_i)\|_2^2} \leq CL^2 \sqrt{\frac{F(\bar{\mathbf{W}}) n \log m}{m}}.
\end{aligned}$$

The last inequality is due to the event E_2 . Combined with (44), we have

$$\mathfrak{R}_{\tilde{\mathcal{S}}}(\mathcal{F}) \leq \frac{\|\mathbf{a}\|_2}{n} \mathbb{E}_\epsilon \sup_{\mathbf{W} \in \mathcal{W}_1} \left\| \sum_{i=1}^n \epsilon_i (h^L(\tilde{\mathbf{x}}_i) - h_0^L(\tilde{\mathbf{x}}_i)) \right\|_2 \leq CL^2 \sqrt{\frac{F(\bar{\mathbf{W}}) \log m}{n}}.$$

The proof is completed. $\square$

Proof of Theorem 2 Using Lemma 16, we have

$$\begin{aligned}
& (f_{\mathbf{W}}(\mathbf{x}) - f_{\bar{\mathbf{W}}}(\mathbf{x}))^2 = (\mathbf{a}^\top (h^L(\mathbf{x}) - \bar{h}^L(\mathbf{x})))^2 \leq m \|h^L(\mathbf{x}) - \bar{h}^L(\mathbf{x})\|_2^2 \\
&\leq 2m (\|h^L(\mathbf{x}) - h_0^L(\mathbf{x})\|_2^2 + \|h_0^L(\mathbf{x}) - \bar{h}^L(\mathbf{x})\|_2^2) \\
&\leq CL^4 \log m F(\bar{\mathbf{W}}),
\end{aligned}$$

where the last inequality is due to $\|\mathbf{W}^l - \mathbf{W}^l(0)\|_2 \leq \|\mathbf{W} - \mathbf{W}(0)\|_F \leq \sqrt{F(\overline{\mathbf{W}})}$ and $\|\overline{\mathbf{W}}^l - \mathbf{W}^l(0)\|_2 \leq \|\mathbf{W}^l - \mathbf{W}(0)\|_2 + \|\mathbf{W}^l - \overline{\mathbf{W}}\|_2 \leq C\sqrt{F(\overline{\mathbf{W}})}$ for all $l \in [L]$. Since logistic loss ℓ is $1/4$ -smooth, the following property holds,

$$|\ell'(x)| \leq \sqrt{\ell(x)/2}, \quad x \in \mathbb{R}.$$

It then follows that for any $\mathbf{W} \in \mathcal{W}_1, z \in \mathcal{Z}$,

$$\begin{aligned} \ell(yf_{\mathbf{W}}(\mathbf{x})) &\leq \ell(yf_{\overline{\mathbf{W}}}(\mathbf{x})) + y(f_{\mathbf{W}}(\mathbf{x}) - f_{\overline{\mathbf{W}}}(\mathbf{x}))\ell'(yf_{\overline{\mathbf{W}}}(\mathbf{x})) + \frac{(f_{\mathbf{W}}(\mathbf{x}) - f_{\overline{\mathbf{W}}}(\mathbf{x}))^2}{8} \\ &\leq \ell(yf_{\overline{\mathbf{W}}}(\mathbf{x})) + 2|\ell'(yf_{\overline{\mathbf{W}}}(\mathbf{x}))|^2 + \frac{(f_{\mathbf{W}}(\mathbf{x}) - f_{\overline{\mathbf{W}}}(\mathbf{x}))^2}{4} \\ &\leq 2\ell(yf_{\overline{\mathbf{W}}}(\mathbf{x})) + CL^4 \log m F(\overline{\mathbf{W}}). \end{aligned}$$

Then we can choose $G' = 2G + CL^4 \log m F(\overline{\mathbf{W}})$. According to Lemma 14, we have with probability at least $1 - \delta$,

$$|\mathcal{L}_S(\overline{\mathbf{W}}) - \mathcal{L}(\overline{\mathbf{W}})| \leq \left(\frac{2G\mathcal{L}(\overline{\mathbf{W}}) \log(2/\delta)}{n} \right)^{1/2} + \frac{2G \log(2/\delta)}{3n}. \quad (45)$$

It then follows that

$$\mathcal{L}_S(\overline{\mathbf{W}}) \leq 2\mathcal{L}(\overline{\mathbf{W}}) + \frac{7G \log(2/\delta)}{6n},$$

which implies that $F_S(\overline{\mathbf{W}}) \leq F(\overline{\mathbf{W}})$. Combined with Theorem 1, with probability at least $1 - \delta$, we have $\mathbf{W}(t) \in \mathcal{W}_1$. Furthermore, events E_1, E_2 hold due to Lemma 6 and Lemma 15. Hence, by Lemma 1 and Lemma 22, there holds

$$\begin{aligned} \mathcal{L}(\mathbf{W}(t)) - 2\mathcal{L}_S(\mathbf{W}(t)) &\lesssim (\log n)^3 \mathfrak{R}_{S,n}^2(\mathcal{F}) + \frac{G' \log(2/\delta)}{n} \\ &= \tilde{O} \left(\frac{L^4 F(\overline{\mathbf{W}}) + G \log(2/\delta)}{n} \right). \end{aligned}$$

As a result,

$$\begin{aligned} \eta \sum_{t=0}^{T-1} \mathcal{L}(\mathbf{W}(t)) &= \eta \sum_{t=0}^{T-1} (L(\mathbf{W}(t)) - 2\mathcal{L}_S(\mathbf{W}(t)) + 2\eta \sum_{t=0}^{T-1} \mathcal{L}_S(\mathbf{W}(t))) \\ &= \tilde{O} \left(\frac{(\eta T L^4 + n) F(\overline{\mathbf{W}}) + \eta T G \log(2/\delta)}{n} \right), \end{aligned}$$

from which we derive

$$\frac{1}{T} \sum_{t=0}^{T-1} \mathcal{L}(\mathbf{W}(t)) = \tilde{O} \left(\frac{L^4 F(\overline{\mathbf{W}}) + G \log(2/\delta)}{n} \right),$$

where we have used $\eta T \asymp n$. The proof is completed.

D Proofs on NTK separability

We denote $\bar{h}$ be (12) when $\mathbf{W} = \overline{\mathbf{W}}$. Let $\overline{\mathbf{W}} = \mathbf{W}(0) + \lambda \mathbf{W}_*$. Choose $\lambda = 2 \log T / \gamma$. Applying Lemma 19 and letting $R = \lambda$, we know that if $m \gtrsim L^{16} d (\log m)^5 \log(nL/\delta) (\log T)^2 / \gamma^8$, then with probability at least $1 - \delta$, for all $i \in [n]$, there holds

$$\begin{aligned} &\left| \left\langle \lambda \mathbf{W}_*, \frac{\partial f_{\mathbf{W}(0)}(\mathbf{x}_i)}{\partial \mathbf{W}(0)} \right\rangle - f_{\overline{\mathbf{W}}}(\mathbf{x}_i) \right| \\ &= \left| f_{\mathbf{W}(0)}(\mathbf{x}_i) - f_{\overline{\mathbf{W}}}(\mathbf{x}_i) - \left\langle \mathbf{W}(0) - \overline{\mathbf{W}}, \frac{\partial f_{\mathbf{W}(0)}(\mathbf{x}_i)}{\partial \mathbf{W}(0)} \right\rangle \right| \\ &\leq CL^{8/3} \lambda^{4/3} m^{-1/6} (\log m)^{2/3} \leq \frac{\lambda \gamma}{2}, \end{aligned}$$

where we have used $f_{\mathbf{W}(0)}(\mathbf{x}) = 0$ for any $\mathbf{x} \in \mathcal{X}$ due to Lemma 7. Therefore, by Assumption 3, we have

$$\begin{aligned} y_i f_{\bar{\mathbf{W}}}(\mathbf{x}_i) &= y_i \left\langle \lambda \mathbf{W}_*, \frac{\partial f_{\mathbf{W}(0)}(\mathbf{x}_i)}{\partial \mathbf{W}(0)} \right\rangle - y_i \left(\left\langle \lambda \mathbf{W}_*, \frac{\partial f_{\mathbf{W}(0)}(\mathbf{x}_i)}{\partial \mathbf{W}(0)} \right\rangle - f_{\bar{\mathbf{W}}}(\mathbf{x}_i) \right) \\ &\geq \lambda y_i \left\langle \mathbf{W}_*, \frac{\partial f_{\mathbf{W}(0)}(\mathbf{x}_i)}{\partial \mathbf{W}(0)} \right\rangle - \left| \left\langle \lambda \mathbf{W}_*, \frac{\partial f_{\mathbf{W}(0)}(\mathbf{x}_i)}{\partial \mathbf{W}(0)} \right\rangle - f_{\bar{\mathbf{W}}}(\mathbf{x}_i) \right| \\ &\geq \lambda \gamma - \frac{\lambda \gamma}{2} = \frac{\lambda \gamma}{2} = \log T. \end{aligned}$$

As a result,

$$\mathcal{L}_S(\bar{\mathbf{W}}) = \frac{1}{n} \sum_{i=1}^n \ell(-y_i f_{\bar{\mathbf{W}}}(\mathbf{x}_i)) \leq \log(1 + \exp(-\log T)) \leq \frac{1}{T}.$$

For any $\mathbf{x} \in \mathcal{X}$, Lemma 16 implies that

$$\begin{aligned} y f_{\bar{\mathbf{W}}}(\mathbf{x}) &\geq -|f_{\bar{\mathbf{W}}}(\mathbf{x})| \geq -|f_{\mathbf{W}(0)}(\mathbf{x})| - |f_{\bar{\mathbf{W}}}(\mathbf{x}) - f_{\mathbf{W}(0)}(\mathbf{x})| \\ &\geq -\mathbf{a}^\top \|\bar{h}^L(\mathbf{x}) - h_0^L(\mathbf{x})\|_2 \geq -CL^2 \sqrt{\log m} \lambda \end{aligned}$$

Hence,

$$G = \sup_z \ell(y f_{\bar{\mathbf{W}}}(\mathbf{x})) \lesssim \log(1 + \exp(L^2 \sqrt{\log m} \log T / \gamma)) \lesssim \frac{L^2 \sqrt{\log m} \log T}{\gamma}.$$

Note that if $x^2 \leq \alpha x + \beta$, then $x^2 \leq \alpha^2 + 2\beta$. Combined with (45), it then follows that

$$\mathcal{L}(\bar{\mathbf{W}}) \leq 2\mathcal{L}_S(\bar{\mathbf{W}}) + \frac{4G \log(2/\delta)}{3n} + \frac{2G \log(2/\delta)}{n} \lesssim \frac{1}{T} + \frac{L^2 \sqrt{\log m} \log T \log(2/\delta)}{\gamma n}.$$

Thus, we have

$$F(\bar{\mathbf{W}}) \lesssim 3\eta T \left(\frac{1}{T} + \frac{L^2 \sqrt{\log m} \log T \log(2/\delta)}{\gamma n} \right) + \lambda^2 = \tilde{O} \left(\frac{(\log T)^2 (1 + \gamma L^2)}{\gamma^2} \right). \quad (46)$$

Applying Theorem 2 and note that $\tilde{F}_S(\bar{\mathbf{W}}) \leq \mathcal{L}_S(\bar{\mathbf{W}}) \leq \frac{1}{T}$, there holds

$$\frac{1}{T} \sum_{t=0}^{T-1} \mathcal{L}(\mathbf{W}(t)) = \tilde{O} \left(\frac{L^4 (\log T)^2 (1 + \gamma L^2)}{n \gamma^2} \right).$$

The proof is completed.