

the 1990s, the number of people with a diagnosis of schizophrenia has increased in the United Kingdom (Meltzer 1999). The prevalence of schizophrenia in the United Kingdom is estimated to be 1.2% (Meltzer 1999). The prevalence of schizophrenia in the United States is estimated to be 1.1% (Meltzer 1999).

There is a growing awareness of the need to improve the lives of people with schizophrenia. The World Health Organization (WHO) has developed a set of guidelines for the management of schizophrenia (WHO 1993). The guidelines recommend that people with schizophrenia should be treated with a combination of medication and psychosocial interventions. The guidelines also recommend that people with schizophrenia should be treated in a community setting, rather than in a hospital.

The guidelines also recommend that people with schizophrenia should be treated with a combination of medication and psychosocial interventions. The guidelines also recommend that people with schizophrenia should be treated in a community setting, rather than in a hospital. The guidelines also recommend that people with schizophrenia should be treated with a combination of medication and psychosocial interventions.

The guidelines also recommend that people with schizophrenia should be treated in a community setting, rather than in a hospital. The guidelines also recommend that people with schizophrenia should be treated with a combination of medication and psychosocial interventions. The guidelines also recommend that people with schizophrenia should be treated in a community setting, rather than in a hospital.

The guidelines also recommend that people with schizophrenia should be treated with a combination of medication and psychosocial interventions. The guidelines also recommend that people with schizophrenia should be treated in a community setting, rather than in a hospital. The guidelines also recommend that people with schizophrenia should be treated with a combination of medication and psychosocial interventions.

The guidelines also recommend that people with schizophrenia should be treated in a community setting, rather than in a hospital. The guidelines also recommend that people with schizophrenia should be treated with a combination of medication and psychosocial interventions. The guidelines also recommend that people with schizophrenia should be treated in a community setting, rather than in a hospital.

The guidelines also recommend that people with schizophrenia should be treated with a combination of medication and psychosocial interventions. The guidelines also recommend that people with schizophrenia should be treated in a community setting, rather than in a hospital. The guidelines also recommend that people with schizophrenia should be treated with a combination of medication and psychosocial interventions.

The guidelines also recommend that people with schizophrenia should be treated in a community setting, rather than in a hospital. The guidelines also recommend that people with schizophrenia should be treated with a combination of medication and psychosocial interventions. The guidelines also recommend that people with schizophrenia should be treated in a community setting, rather than in a hospital.

The guidelines also recommend that people with schizophrenia should be treated with a combination of medication and psychosocial interventions. The guidelines also recommend that people with schizophrenia should be treated in a community setting, rather than in a hospital. The guidelines also recommend that people with schizophrenia should be treated with a combination of medication and psychosocial interventions.