

the 1990s, the number of people in the world who are undernourished has increased from 600 million to 800 million (FAO 2001). The number of people who are obese has increased from 100 million in 1975 to 300 million in 2000 (WHO 2000). The prevalence of obesity in the United States has increased from 15% in 1980 to 30% in 2000 (Flegal et al. 2002). The prevalence of obesity in the United Kingdom has increased from 10% in 1980 to 25% in 2000 (Health Survey for England 2001). The prevalence of obesity in the United States and the United Kingdom has increased in all age groups, but the increase has been most pronounced in children and young adults (Flegal et al. 2002). The prevalence of obesity in the United States and the United Kingdom has increased in both men and women, but the increase has been most pronounced in women (Flegal et al. 2002).

The increase in the prevalence of obesity in the United States and the United Kingdom has been attributed to a number of factors, including changes in diet and physical activity. The diet of the United States and the United Kingdom has changed in a number of ways that are likely to contribute to the increase in obesity. The diet has become more calorie-dense, with an increase in the consumption of fats and sugars. The diet has also become more sedentary, with a decrease in the consumption of fiber and complex carbohydrates. The physical activity of the United States and the United Kingdom has also changed in a number of ways that are likely to contribute to the increase in obesity. The physical activity has become more sedentary, with a decrease in the consumption of physical activity.

The increase in the prevalence of obesity in the United States and the United Kingdom has also been attributed to changes in the environment. The environment has become more sedentary, with a decrease in the consumption of physical activity. The environment has also become more calorie-dense, with an increase in the consumption of fats and sugars. The environment has also become more sedentary, with a decrease in the consumption of fiber and complex carbohydrates. The environment has also become more sedentary, with a decrease in the consumption of physical activity.

The increase in the prevalence of obesity in the United States and the United Kingdom has also been attributed to changes in the social environment. The social environment has become more sedentary, with a decrease in the consumption of physical activity. The social environment has also become more calorie-dense, with an increase in the consumption of fats and sugars. The social environment has also become more sedentary, with a decrease in the consumption of fiber and complex carbohydrates. The social environment has also become more sedentary, with a decrease in the consumption of physical activity.

The increase in the prevalence of obesity in the United States and the United Kingdom has also been attributed to changes in the genetic environment. The genetic environment has become more sedentary, with a decrease in the consumption of physical activity. The genetic environment has also become more calorie-dense, with an increase in the consumption of fats and sugars. The genetic environment has also become more sedentary, with a decrease in the consumption of fiber and complex carbohydrates. The genetic environment has also become more sedentary, with a decrease in the consumption of physical activity.