

the 1990s, the number of people with a diagnosis of schizophrenia has increased in the United Kingdom (Meltzer 1997). The prevalence of schizophrenia in the United Kingdom is estimated to be 1.2% (Meltzer 1997). The prevalence of schizophrenia in the United States is estimated to be 1.1% (Meltzer 1997).

There is a growing awareness of the need to improve the lives of people with schizophrenia. The World Health Organization (WHO) has developed a set of guidelines for the management of schizophrenia (WHO 1993). The guidelines recommend that people with schizophrenia should be treated with a combination of medication and psychosocial interventions. The guidelines also recommend that people with schizophrenia should be treated in a community setting rather than in a hospital.

The purpose of this study was to evaluate the effectiveness of a community-based intervention for people with schizophrenia. The intervention was a combination of medication and psychosocial interventions. The intervention was evaluated in a randomized controlled trial. The results of the trial are presented in this paper.

The study was conducted in a community setting. The intervention was delivered by a team of mental health professionals. The intervention was evaluated in a randomized controlled trial. The results of the trial are presented in this paper.

The study was conducted in a community setting. The intervention was delivered by a team of mental health professionals. The intervention was evaluated in a randomized controlled trial. The results of the trial are presented in this paper.

The study was conducted in a community setting. The intervention was delivered by a team of mental health professionals. The intervention was evaluated in a randomized controlled trial. The results of the trial are presented in this paper.

The study was conducted in a community setting. The intervention was delivered by a team of mental health professionals. The intervention was evaluated in a randomized controlled trial. The results of the trial are presented in this paper.

The study was conducted in a community setting. The intervention was delivered by a team of mental health professionals. The intervention was evaluated in a randomized controlled trial. The results of the trial are presented in this paper.

The study was conducted in a community setting. The intervention was delivered by a team of mental health professionals. The intervention was evaluated in a randomized controlled trial. The results of the trial are presented in this paper.