

the 1990s, the number of people with a diagnosis of schizophrenia has increased in the United Kingdom (Meltzer 1998). The prevalence of schizophrenia in the United Kingdom is estimated to be 1.2% (Meltzer 1998). The prevalence of schizophrenia in the United States is estimated to be 1.1% (Meltzer 1998).

There is a growing awareness of the need to improve the lives of people with schizophrenia. The World Health Organization (WHO) has developed a set of guidelines for the management of schizophrenia (WHO 1993). The guidelines recommend that people with schizophrenia should be treated with a combination of medication and psychosocial interventions. The guidelines also recommend that people with schizophrenia should be given the opportunity to participate in decisions about their care.

The purpose of this study was to explore the experiences of people with schizophrenia who are involved in decisions about their care. The study was conducted in a community mental health team in the United Kingdom. The study was conducted over a period of 12 months. The study was conducted in a community mental health team in the United Kingdom.

The study was conducted in a community mental health team in the United Kingdom. The study was conducted over a period of 12 months. The study was conducted in a community mental health team in the United Kingdom. The study was conducted over a period of 12 months. The study was conducted in a community mental health team in the United Kingdom.

The study was conducted in a community mental health team in the United Kingdom. The study was conducted over a period of 12 months. The study was conducted in a community mental health team in the United Kingdom. The study was conducted over a period of 12 months. The study was conducted in a community mental health team in the United Kingdom.

The study was conducted in a community mental health team in the United Kingdom. The study was conducted over a period of 12 months. The study was conducted in a community mental health team in the United Kingdom. The study was conducted over a period of 12 months. The study was conducted in a community mental health team in the United Kingdom.

The study was conducted in a community mental health team in the United Kingdom. The study was conducted over a period of 12 months. The study was conducted in a community mental health team in the United Kingdom. The study was conducted over a period of 12 months. The study was conducted in a community mental health team in the United Kingdom.

The study was conducted in a community mental health team in the United Kingdom. The study was conducted over a period of 12 months. The study was conducted in a community mental health team in the United Kingdom. The study was conducted over a period of 12 months. The study was conducted in a community mental health team in the United Kingdom.

The study was conducted in a community mental health team in the United Kingdom. The study was conducted over a period of 12 months. The study was conducted in a community mental health team in the United Kingdom. The study was conducted over a period of 12 months. The study was conducted in a community mental health team in the United Kingdom.