

Problem Description


Theorem (Erdős' Pentagon Conjecture): The number of pentagons (5-cycles) in a simple triangle free graph G on n vertices is $\leq \left(\frac{n}{5}\right)^5$.

We are interest in adding the requirement that G is regular, so we ask: Given a simple, triangle free, regular graph G what is the maximum number of pentagons in the graph.

Conjecture (Regular Erdős' Pentagon): The total number of pentagons in a regular triangle free graph is

$$\lesssim \frac{n}{5} \left(\frac{\Delta(G)}{2} \right)^4 = \frac{n}{5} \frac{\Delta(G)^4}{16}.$$

Tightness: For any $k \in \mathbb{N}$ even we can construct a blowup of the 5-cycle with $k/2$ vertices:



Each "supernode" consists of an independent set of $k/2$ vertices which are connected densely to the vertices in the adjacent supernode.

Any vertex in this graph G has degree exactly k and G is triangle free, so G is a regular, triangle free graph on $n = \frac{5k}{2}$ vertices.

Consider any 5-cycle in this graph. If the 5-cycle contains only vertices from ≤ 4 then consider the subgraph induced by those supernodes. This subgraph is clearly bipartite so cannot contain an odd length cycle, which is a contradiction. Therefore this 5-cycle must contain exactly one vertex from each of the 5 supernodes. This choice of nodes uniquely identifies this cycle. Conversely, any choice of a vertex from each supernode forms a valid 5-cycle. Hence the set of 5-cycles is in 1-1 correspondence with all possible choices of 1 vertex from each supernode. Hence there are exactly $\left(\frac{k}{2}\right)^5$ 5-cycles.

The graph G has $\Delta(G) = k$ and $n = 5k/2$. We re-write

$$\left(\frac{k}{2}\right)^5 = \frac{k}{2} \left(\frac{k}{2}\right)^4 = \frac{n}{5} \left(\frac{\Delta(G)}{2}\right)^4$$

which shows G meets the bound of the conjecture.

We can construct an infinite increasing (in both n and $\Delta(G)$) sequence of such graphs. Hence if the conjecture is true it is tight.

Flag Argument

Locality

The factor of $\frac{n}{5}$ means we cannot apply local flags directly. Instead of bounding the total number of pentagons in the graph we instead try to bound the number of pentagons containing some fixed vertex v . Call this P_v .

If we show that $P_v \lesssim \frac{\Delta(G)^4}{k}$ then as each pentagon contains 5 such vertices $\sum_{v \in V(G)} P_v$ will count each pentagon 5-times, giving us a bound of $\lesssim \frac{n}{5} \frac{\Delta(G)^4}{k}$ on the total number of pentagons. If our tight example above is correct then we expect the extremal graphs to be vertex-transitive, meaning this approach should be effective.

Note


If a graph is triangle free, then C_5 is a subgraph iff it is an induced subgraph, hence counting induced pentagons suffices.

Transformation


Let G be a simple, regular, triangle free graph and let $v \in V(G)$ be an arbitrary vertex. As G is triangle free $N(v)$ is an independent set. G is regular so we also have $|N(v)| = \Delta(G)$.

Consider a pentagon containing v : $\{v, u_1, u_2, u_3, u_4\}$. Two of u_i must be $\in N(v)$ and two must be $\notin N(v) \cup \{v\}$

If we construct a 3-coloured graph H which is a copy of G where v is coloured green, $N(v)$ is coloured black and the rest are coloured red then the number of pentagons in G containing v is the same as the number of pentagons in H containing 1 green node, 2 black nodes and 2 red nodes.



In fact, it suffices to count how many paths of length 4 there are in H starting and ending with black nodes and using only red nodes as middle nodes as the green node is densely connected to the black nodes.



If we remove v (the solitary green node) from the graph then this count is unchanged. This also reduces the degree of some small number of vertices by 1 but this is asymptotically irrelevant.

We have then taken counting the number of pentagons containing v in G and turned it into an equivalent problem of counting paths of length 4 starting and ending with black nodes in a black/red coloured graph H . This graph H has $\Delta(G)$ black nodes and is $\Delta(G)$ regular. The set of black nodes is independent.

Let \mathcal{G} be the family of all red-black vertex coloured, regular graphs which are triangle free, have $\Delta(G)$ black nodes and the set of black nodes are independent. If we can bound the number of paths of length 4 starting and ending at black nodes in this family \mathcal{G} then we have a bound on our number of pentagons containing G .

Local Flags

Given the above family of graphs \mathcal{G} , what are the local flags?

Lemma: F is a local \emptyset -flag iff each connected component of $F \in \mathcal{G}$ contains a black node.

🕒 Todo

Proof. (Only the \Leftarrow direction actually matters for the application).

In particular this tells us what our local types σ are.