

# **2× Freezing Medium (R0 + 10% FCS + 20% DMSO)**

**Abdullah Lab, IMMEI, University Hospital Bonn**

**Buffer ID:** BUF-FREEZE-MIX-2x-001

**Version:** v1.0

**Author:** Dillon Corvino

**Date:** 2025-11-21

## **Purpose**

This buffer is a **2× concentrated freezing medium** used for the cryopreservation of PBMCs and other lymphocytes. When mixed 1:1 with a cell suspension in R10 or R0, it yields a standard **final freezing medium containing 10% DMSO and 5% FCS** (or 10% FCS if starting from R10), suitable for controlled-rate freezing.

## **Working buffer composition**

Component	Final concentration (2×)	Purpose
RPMI 1640 (R0)	—	Base medium
Fetal calf serum	10% (v/v)	Protects cells during freezing
DMSO (cell-grade)	20% (v/v)	Cryoprotectant

When mixed 1:1 with cells in R10 or R0, the **final freezing medium** contains:

- **10% DMSO**
- **5–10% FCS** (depending on suspension medium)
- RPMI base medium

## Preparation

### Stock reagents

Reagent	Stock concentration	Notes
RPMI 1640	—	Sterile-filtered
DMSO, cell-grade	100%	Use fresh or freshly opened
Fetal calf serum	—	Heat-inactivated recommended

### Preparation of 10 mL 2× freezing medium

1. In a sterile 15 mL tube, add:
  - **7 mL R0 medium** (RPMI without serum)
  - **1 mL heat-inactivated FCS** (10% v/v)
  - **2 mL DMSO** (20% v/v)
2. Mix gently by inversion; **do not vortex**.
3. Keep on ice until use.

#### Warning

DMSO is toxic at room temperature. Keep the 2× freezing medium on **ice**, and **mix with cells immediately** after preparation.

## Usage

1. Adjust PBMCs or lymphocytes to **20 × 10<sup>6</sup> cells/mL** in R10 or R0.
2. Add an **equal volume** of this 2× freezing medium.
3. Mix gently by pipetting.
4. Dispense **1 mL per cryovial** (final 10% DMSO).
5. Freeze using a **controlled-rate freezing container** (e.g. Mr Frosty) at  $-70/-80\text{ }^{\circ}\text{C}$ .
6. Transfer to liquid nitrogen after 12 hours.

## Storage and stability

- Use **fresh on the day of preparation**.

- Keep on **ice** during use.
- **Do not store** after the experiment; discard any remaining buffer.
- DMSO-containing mixtures degrade rapidly and should never be refrozen.

## Reagent details

Reagent	Supplier	Cat. #	Notes
RPMI 1640	Various	—	Base medium
Fetal calf serum	Various	—	Heat-inactivated
DMSO, cell-grade	Sigma/etc.	—	Highly hygroscopic; handle aseptically

## Safety

- Handle DMSO with gloves; it enhances dermal absorption of other chemicals.
- Prepare freezing medium in a biosafety cabinet using aseptic technique.
- Dispose of DMSO-containing waste according to institutional chemical waste protocols.
- Avoid eye/skin contact and inhalation.

## Version history

Version	Date	Author	Changes
v1.0	2025-11-21	Dillon Corvino	Initial Quarto buffer document for 2× freezing medium.