

**Table 3.** Bivariable Associations of Consistent Blood and Urogenital STI Screening with Various Exposures Among PrEP-Eligible ARTnet Participants who Have Ever Used PrEP

	Blood STI Screening (N=730)						Urogenital STI Screening (N=742)					
	Consistent <sup>a</sup>		Inconsistent		OR	95% CI	Consistent		Inconsistent		OR	95% CI
	N	%	N	%			N	%	N	%		
<b>Total Sample</b>	635	87.0	95	13.0			575	77.5	167	22.5		
<b>Age</b>												
15-24	75	83.3	15	16.7	1.00	-	73	80.2	18	19.8	1.00	-
25-34	215	87.8	30	12.2	1.43	0.72, 2.77	201	80.1	50	19.9	0.99	0.53, 1.78
35-44	123	90.4	13	9.6	1.89	0.85, 4.25	104	75.4	34	24.6	0.75	0.39, 1.42
45-54	140	85.9	23	14.1	1.22	0.59, 2.45	127	77.9	36	22.1	0.87	0.45, 1.62
55-65	82	85.4	14	14.6	1.17	0.53, 2.61	70	70.7	29	29.3	0.60	0.30, 1.16
<b>Race/Ethnicity</b>												
Black (Non-Hispanic)	30	93.8	2	6.3	1.00	-	28	87.5	4	12.5	1.00	-
Hispanic	84	91.3	8	8.7	0.70	0.10, 2.99	78	84.8	14	15.2	0.80	0.21, 2.44
Other (Non-Hispanic)	53	91.4	5	8.6	0.71	0.10, 3.50	45	77.6	13	22.4	0.49	0.13, 1.56
White (Non-Hispanic)	468	85.4	80	14.6	0.39	0.06, 1.33	424	75.7	136	24.3	0.45	0.13, 1.16
<b>Health Insurance</b>												
None	28	84.8	5	15.2	1.00	-	23	69.7	10	30.3	1.00	-
Public	98	87.5	14	12.5	1.25	0.38, 3.59	91	79.8	23	20.2	1.72	0.70, 4.06
Private	507	87.1	75	12.9	1.21	0.40, 2.97	459	77.5	133	22.5	1.50	0.67, 3.15
<b>Census Division</b>												
New England	36	92.3	3	7.7	1.00	-	33	80.5	8	19.5	1.00	-
Middle Atlantic	89	85.6	15	14.4	0.49	0.11, 1.61	86	79.6	22	20.4	0.95	0.37, 2.27
East North Central	89	84.0	17	16.0	0.44	0.10, 1.40	83	77.6	24	22.4	0.84	0.33, 1.99
West North Central	29	74.4	10	25.6	0.24	0.05, 0.87	24	60.0	16	40.0	0.36	0.13, 0.96
South Atlantic	119	86.9	18	13.1	0.55	0.12, 1.75	93	67.9	44	32.1	0.51	0.21, 1.15
East South Central	18	85.7	3	14.3	0.50	0.08, 2.94	13	61.9	8	38.1	0.39	0.12, 1.28
West South Central	62	89.9	7	10.1	0.74	0.15, 2.84	57	82.6	12	17.4	1.15	0.41, 3.08
Mountain	34	79.1	9	20.9	0.31	0.07, 1.16	32	74.4	11	25.6	0.71	0.24, 1.97
Pacific	159	92.4	13	7.6	1.02	0.23, 3.36	154	87.5	22	12.5	1.70	0.66, 4.02
<b>Deep South<sup>b</sup></b>												
No	556	87.3	81	12.7	1.00	-	513	79.2	135	20.8	1.00	-
Yes	79	84.9	14	15.1	0.82	0.46, 1.58	62	66.0	32	34.0	0.51	0.32, 0.82
<b>Exposure at Site<sup>c</sup></b>												
No							15	68.2	7	31.8	1.00	-
Yes							560	77.8	160	22.2	1.63	0.61, 3.94

Abbreviations: STI, sexually transmitted infection; PrEP, preexposure prophylaxis; OR, odds ratio; CI, confidence interval

<sup>a</sup> Consistent screening defined as always or sometimes (versus rarely or never) being screened at PrEP care visits compared

<sup>b</sup> Residence in Alabama, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina

<sup>c</sup> Defined as any insertive anal intercourse or receiving oral sex for urogenital STI screening