

Table 4. Bivariable Associations of Consistent Rectal and Pharyngeal STI Screening with Various Exposures Among PrEP-Eligible ARTnet Participants who Have Ever Used PrEP

	Rectal STI Screening (N=746)						Pharyngeal STI Screening (N=744)					
	Consistent ^a		Inconsistent		OR	95% CI	Consistent		Inconsistent		OR	95% CI
	N	%	N	%			N	%	N	%		
Total Sample	425	57.0	321	43.0			477	64.1	267	35.9		
Age												
15-24	61	67.0	30	33.0	1.00	-	63	70.0	27	30.0	1.00	-
25-34	146	57.9	106	42.1	0.68	0.41, 1.11	163	64.9	88	35.1	0.79	0.47, 1.32
35-44	74	53.6	64	46.4	0.57	0.33, 0.98	89	64.5	49	35.5	0.78	0.44, 1.37
45-54	91	55.2	74	44.8	0.60	0.35, 1.03	102	61.8	63	38.2	0.69	0.40, 1.19
55-65	53	53.0	47	47.0	0.55	0.31, 0.99	60	60.0	40	40.0	0.64	0.35, 1.17
Race/Ethnicity												
Black (Non-Hispanic)	16	50.0	16	50.0	1.00	-	19	59.4	13	40.6	1.00	-
Hispanic	57	62.0	35	38.0	1.63	0.72, 3.69	61	66.3	31	33.7	1.35	0.58, 3.07
Other (Non-Hispanic)	39	67.2	19	32.8	2.05	0.85, 5.02	43	74.1	15	25.9	1.96	0.78, 4.95
White (Non-Hispanic)	313	55.5	251	44.5	1.25	0.61, 2.56	354	63.0	208	37.0	1.16	0.55, 2.39
Health Insurance^a												
None	19	57.6	14	42.4	1.00	-	22	66.7	11	33.3	1.00	-
Public	58	50.0	58	50.0	0.74	0.33, 1.60	74	63.8	42	36.2	0.88	0.38, 1.96
Private	347	58.4	247	41.6	1.04	0.50, 2.09	380	64.2	212	35.8	0.90	0.41, 1.85
Census Division												
New England	29	70.7	12	29.3	1.00	-	30	73.2	11	26.8	1.00	-
Middle Atlantic	65	60.2	43	39.8	0.63	0.28, 1.33	74	69.2	33	30.8	0.82	0.36, 1.80
East North Central	57	52.8	51	47.2	0.46	0.21, 0.98	59	54.6	49	45.4	0.44	0.19, 0.95
West North Central	21	52.5	19	47.5	0.46	0.18, 1.13	24	61.5	15	38.5	0.59	0.22, 1.50
South Atlantic	65	46.1	76	53.9	0.35	0.16, 0.73	79	56.0	62	44.0	0.47	0.21, 0.98
East South Central	7	33.3	14	66.7	0.21	0.06, 0.62	9	42.9	12	57.1	0.28	0.09, 0.82
West South Central	40	58.0	29	42.0	0.57	0.24, 1.28	47	68.1	22	31.9	0.78	0.32, 1.82
Mountain	19	45.2	23	54.8	0.34	0.13, 0.83	20	47.6	22	52.4	0.33	0.13, 0.82
Pacific	122	69.3	54	30.7	0.93	0.43, 1.93	135	76.7	41	23.3	1.21	0.54, 2.56
Deep South^b												
No	384	59.1	266	40.9	1.00	-	424	65.4	224	34.6	1.00	-
Yes	41	42.7	55	57.3	0.52	0.33, 0.79	53	55.2	43	44.8	0.65	0.42, 1.01
Exposure at Site^c												
No	61	44.2	77	55.8	1.00	-	15	62.5	9	37.5	1.00	-
Yes	364	59.9	244	40.1	1.88	1.30, 2.74	462	64.2	258	35.8	1.07	0.45, 2.45

Abbreviations: STI, sexually transmitted infection; PrEP, preexposure prophylaxis; OR, odds ratio; CI, confidence interval

^a Consistent screening defined as always or sometimes (versus rarely or never) being screened at PrEP care visits compared

^b Residence in Alabama, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina

^c Defined as any receptive anal intercourse for rectal STI screening and any performance of oral sex on a partner for pharyngeal STI screening within the past year