

Table 5. Multivariable Associations of Consistent^a STI Screening with Residence in the Southeast U.S.^b Among PrEP-Eligible ARTnet Participants who Have Ever Used PrEP

	Adjusted PR ^c	95% CI
Syphilis Screening^d		
Residence Outside Southeast	1.00	-
Residence in Southeast	1.02	0.95, 1.10
Urogenital STI Screening		
Residence Outside Southeast	1.00	-
Residence in Southeast	0.86	0.76, 0.98
Rectal STI Screening		
Residence Outside Southeast	1.00	-
Residence in Southeast	0.76	0.62, 0.93
Pharyngeal STI Screening		
Residence Outside Southeast	1.00	-
Residence in Southeast	0.87	0.74, 1.03

Abbreviations: STI, sexually transmitted infection; PrEP, preexposure prophylaxis; PR, prevalence ratio; CI, confidence interval

^a Consistent screening defined as always or sometimes (versus rarely or never) being screened at PrEP care visits compared

^b Residence in Alabama, Arkansas, Florida, Georgia, Kentucky, Louisiana, Mississippi, North Carolina, South Carolina, Tennessee, Virginia, and West Virginia

^c Adjusted for age, race/ethnicity, annual household income, and sexual exposure at anatomical site

^d Not adjusted for sexual exposure at anatomical site