

Table 5. Multivariable associations of STI screening with living in Deep South states^a among HIV-negative ARTnet participants who have ever used PrEP

	Adjusted OR ^a	95% CI
HIV Screening (Always vs. Not Always)		
Not living in Deep South	<i>Ref</i>	
Living in Deep South	0.59	(0.32, 1.18)
Blood STI Screening (Always/Sometimes vs. Rarely/Never)		
Not living in Deep South	<i>Ref</i>	
Living in Deep South	0.84	(0.47, 1.62)
Urogenital STI Screening (Always/Sometimes vs. Rarely/Never)		
Not living in Deep South	<i>Ref</i>	
Living in Deep South	0.51	(0.32, 0.82)
Rectal STI Screening (Always/Sometimes vs. Rarely/Never)		
Not living in Deep South	<i>Ref</i>	
Living in Deep South	0.51	(0.33, 0.79)
Pharyngeal STI Screening (Always/Sometimes vs. Rarely/Never)		
Not living in Deep South	<i>Ref</i>	
Living in Deep South	0.66	(0.43, 1.02)

Abbreviations: STI, sexually transmitted infection; PrEP, pre-exposure prophylaxis; OR, odds ratio; CI, confidence interval

^aResidence in North Carolina, South Carolina, Georgia, Alabama, Mississippi, Louisiana, Florida

^bAdjusted for age, race/ethnicity, and exposure at site for urogenital, rectal, and pharyngeal STI screening