

Table 2. STI screening among HIV-negative ARTnet participants who have ever used PrEP

	Total		Current PrEP Use		Non-Current PrEP Use	
	N	%	N	%	N	%
Total Sample	809	100.0	631	100.0	178	100.0
Frequency of PrEP-related Check-ups^a						
Every month	23	2.9	13	2.1	10	5.7
Every 3 months	629	78.3	514	82.0	115	65.3
Every 6 months	83	10.3	75	12.0	8	4.5
Every 9 months	3	0.4	1	0.2	2	1.1
Every 12 months	8	1.0	8	1.3	0	0.0
Returned only once	21	2.6	3	0.5	18	10.2
Returned at some other interval of time	11	1.4	6	1.0	5	2.8
Did not return regularly	25	3.1	7	1.1	18	10.2
Frequency of Throat Swab at PrEP Visit for STI Screening						
Always	316	39.1	267	42.3	49	27.5
Sometimes	161	19.9	134	21.2	27	15.2
Rarely	56	6.9	44	7.0	12	6.7
Never	211	26.1	160	25.4	51	28.7
Missing ^a	65	8.0	26	4.1	39	21.9
Frequency of Rectal Swab at PrEP Visit for STI Screening						
Always	270	33.4	226	35.8	44	24.7
Sometimes	155	19.2	128	20.3	27	15.2
Rarely	58	7.2	50	7.9	8	4.5
Never	263	32.5	203	32.2	60	33.7
Missing ^a	63	7.8	24	3.8	39	21.9
Frequency of Blood Sample at PrEP Visit for STI Screening						
Always	492	60.8	415	65.8	77	43.3
Sometimes	143	17.7	114	18.1	29	16.3
Rarely	32	4.0	21	3.3	11	6.2
Never	63	7.8	44	7.0	19	10.7
Missing ^a	79	9.8	37	5.9	42	23.6
Frequency of Urine Sample or Swab at PrEP Visit for STI Screening						
Always	425	52.5	356	56.4	69	38.8
Sometimes	150	18.5	128	20.3	22	12.4
Rarely	42	5.2	33	5.2	9	5.1
Never	125	15.5	86	13.6	39	21.9
Missing ^a	67	8.3	28	4.4	39	21.9
Frequency of HIV screening during PrEP visits						
Always	685	84.7	566	89.7	119	66.9
Sometimes	50	6.2	40	6.3	10	5.6
Rarely	7	0.9	3	0.5	4	2.2
Never	10	1.2	3	0.5	7	3.9
Missing ^a	57	7.0	19	3.0	38	21.3

Abbreviations: STI, sexually transmitted infection; PrEP, pre-exposure prophylaxis

^aMissing data excluded due to questions left unanswered