

Lecture 14: February 10

Lecturer: Jonathan Taylor

Scribes: Kenneth Tay

14.1 Gibbs Sampling for Probit Model

We work through the Gibbs sampler for the probit model to show how data augmentation can make Gibbs sampling easier.

We first specify the model:

- Prior $g(\beta) \propto \exp(-\beta^T \Sigma^{-1} \beta / 2)$ (i.e. normal),
- Likelihood $P(Y_i = 1 | X_i, \beta) = \Phi(X_i^T \beta)$. We can also think of Y_i in terms of a latent variable ε_i :

$$Y_i = \begin{cases} 1 & \text{if } \varepsilon_i \leq X_i^T \beta, \\ 0 & \text{otherwise.} \end{cases}$$

Here, $\varepsilon_i \stackrel{iid}{\sim} \mathcal{N}(0, 1)$.

With these two pieces, we can compute the posterior density for β :

$$\begin{aligned} h(\beta | Y) &= h(\beta | \varepsilon \leq X\beta) \\ &\propto \left[\int_{\{\varepsilon \leq X\beta\}} e^{-\|\varepsilon\|^2/2} d\varepsilon \right] \cdot g(\beta), \\ h(\varepsilon, \beta | \varepsilon \leq X\beta) &\propto e^{-\|\varepsilon\|^2/2} \cdot e^{-\beta^T \Sigma^{-1} \beta / 2} \mathbf{1}_{\{\varepsilon \leq X\beta\}}. \end{aligned}$$

Note that if we ignore the indicator variable, the remainder corresponds to the distribution $\mathbb{N} \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} I & 0 \\ 0 & \Sigma \end{pmatrix} \right)$.

Goal: Sample $(\varepsilon, \beta) | \varepsilon \leq X\beta$.

Recall the Gibbs sampler from the previous lecture. WLOG, assume that $\Sigma = I$. (If not, we can write $\beta = \Sigma^{-1/2} \gamma$ with $\gamma \sim \mathcal{N}(0, I)$.)

Consider what happens to each step of choosing a new ε_i or β_i .

- When moving a particular ε_i , fix ε_{-i} and β . Then the density is $\propto e^{-\varepsilon_i^2/2} \mathbf{1}_{\{\varepsilon_i \leq X_i^T \beta\}}$. We can draw ε_i^{new} from this density, i.e. $\sim \mathcal{N}(0, 1) | (-\infty, X_i^T \beta)$.
Note that all the ε_i^{new} 's can be drawn simultaneously (no need to be sequential as in the usual Gibbs sampler)!
- When moving a particular β_i , fix ε and β_{-i} . The constraints from the indicator variable become

$$\begin{aligned}
\varepsilon_j &\leq X_j^T \beta && \text{for } 1 \leq j \leq n, \\
\varepsilon_j &\leq \sum_{l \neq i} X_{jl} \beta_l + X_{ji} \beta_i && \text{for } 1 \leq j \leq n, \\
X_{ji} \beta_i &\geq \varepsilon_j - \sum_{l \neq i} X_{jl} \beta_l. && \text{for } 1 \leq j \leq n,
\end{aligned}$$

These constraints bound β_i on either side, depending on the sign of X_{ji} :

$$\begin{aligned}
\max_{j: X_{ji} > 0} \frac{1}{X_{ji}} \left[\varepsilon_j - \sum_{l \neq i} X_{jl} \beta_l \right] &\leq \beta_i \leq \min_{j: X_{ji} < 0} \frac{1}{X_{ji}} \left[\varepsilon_j - \sum_{l \neq i} X_{jl} \beta_l \right], \\
L(\beta_{-i}, \varepsilon) &\leq \beta_i \leq U(\beta_{-i}, \varepsilon).
\end{aligned}$$

We draw β_i^{new} from $\mathcal{N}(0, 1) \mid [L, U]$.

14.2 (Agresti 8) Multinomial Regression

Recall the multinomial distribution: If $Y \sim \text{Multinom}(N, \pi)$, where $\pi \in \mathbb{R}_0^k$, $\sum \pi_i = 1$, then it has mass function

$$f(y_1, \dots, y_k \mid \pi) = \binom{N}{y_1, \dots, y_k} \prod_{i=1}^k \pi_i^{y_i},$$

supported on $y_1, \dots, y_k \in \mathbb{Z}_0$, $\sum y_i = N$. Note that there are actually only $k-1$ free parameters in this model. If we take the k^{th} category as the baseline category, we can write the mass function as

$$f(y_1, \dots, y_{k-1} \mid \pi) = \binom{N}{y_1, \dots, y_{k-1}} \prod_{i=1}^{k-1} \pi_i^{y_i},$$

where $y_k = N - \sum_{i=1}^{k-1} y_i$, $\pi_k = 1 - \sum_{i=1}^{k-1} \pi_i$. This mass function is supported on $y_1, \dots, y_{k-1} \in \mathbb{Z}_0$, $\sum_{i=1}^{k-1} y_i \leq N$.

We can compute the log likelihood for this model:

$$\begin{aligned}
\log L(\pi_1, \dots, \pi_{k-1} \mid Y) &= C + \sum_{i=1}^{k-1} y_i \log \pi_i + \left(N - \sum_{i=1}^{k-1} y_i \right) \log \left(1 - \sum_{i=1}^{k-1} \pi_i \right) \\
&= C + \sum_{i=1}^{k-1} y_i \log \left(\frac{\pi_i}{1 - \sum_{i=1}^{k-1} \pi_i} \right) + N \log \left(1 - \sum_{j=1}^{k-1} \pi_j \right),
\end{aligned}$$

where C is some constant that does not depend on π . The identity above shows that this model is an exponential family with sufficient statistic (y_1, \dots, y_{k-1}) and natural parameters $\eta_j = \log \left(\frac{\pi_j}{1 - \sum_{l=1}^{k-1} \pi_l} \right)$.

We can invert the relationship between η and π to obtain $\pi_j = \frac{e^{\eta_j}}{1 + \sum_{l=1}^{k-1} e^{\eta_l}}$. This allows us to rewrite the log likelihood in terms of the natural parameters:

$$\log L(\eta_1, \dots, \eta_{k-1} \mid Y) = \sum_{i=1}^n \eta_i y_i - N \log \left(1 + \sum_{i=1}^{k-1} e^{\eta_i} \right).$$

14.2.1 Baseline Multinomial Logit

In this model, we have $Y_i \stackrel{iid}{\sim} \text{Multinom}(N_i, \pi_\beta(X_i))$, where for $1 \leq j \leq k-1$,

$$[\pi_\beta(X_i)]_j = \frac{\exp(X_i^T \beta_j)}{1 + \sum_{l=1}^{k-1} \exp(X_i^T \beta_l)}, \quad \text{or equivalently,} \quad \eta_i = X_i^T \beta.$$

Note here that β is a $p \times (k-1)$ matrix. (In logistic regression, β was just a $p \times 1$ matrix.) In the above expression for $[\pi_\beta(X_i)]_j$, β_j refers to the j^{th} column of β .

We can compute the log likelihood and its gradient for this model:

$$\begin{aligned} \log L(\beta \mid Y) &= \sum_{i=1}^n \left[\sum_{l=1}^{k-1} Y_{il} (X_i^T \beta_l) - N_i \log \left(1 + \sum_{l=1}^{k-1} e^{X_i^T \beta_l} \right) \right], \\ \nabla \log L(\beta \mid Y) &= X^T [Y - \mathbb{E}_\beta(Y)] \in \mathbb{R}^{p \times (k-1)}. \end{aligned}$$

Note that the gradient of the log likelihood has the exact same form that we had for logistic and loglinear regression.

The Hessian $\nabla^2 \log L(\beta \mid Y)$ is some kind of tensor. It can also be thought of as $\text{Cov}(\nabla \log L(\beta \mid Y))$. It is given by

$$[\nabla^2 \log L(\beta \mid Y)]_{ijkl} = \sum_{c=1}^n X_{ci} X_{ck} \text{Cov}_\beta(Y_c)_{jl}.$$