CC - Test

Jean Pierre Decrorps 31 août 2017

R Markdown

data(Tiramisu)
DF <- Tiramisu
#kable(str(DF))</pre>

CC ill - tira

```
ret <- CC(DF, "ill", "tira", exact = FALSE)</pre>
ret
## $df1
                     Cases Controls Total
##
## Exposed
                                 27
                                       121
## Unexposed
                         7
                                 158
                                       165
## Total
                       101
                                185
                                       286
## Proportion exposed 0.93
                                0.15 0.42
##
## $df2
##
                  Point estimate 95%CI-ll 95%CI-ul
## Odds ratio
                           78.58
                                    31.45
                                            217.15
## Attr. frac. ex.
                            0.99
                                    0.97
                                             1.00
## Attr. frac. pop
                             0.92
                                       NA
                                                 NA
## chi2(1)
                           164.83
                                       NA
                                                 NA
## Pr>chi2
                            0.000
                                       NA
                                                 NA
kable(ret$df1, align=ret$df1.align)
```

| | Cases | Controls | Total |
|--------------------|-------|----------|-------|
| Exposed | 94 | 27 | 121 |
| Unexposed | 7 | 158 | 165 |
| Total | 101 | 185 | 286 |
| Proportion exposed | 0.93 | 0.15 | 0.42 |

kable(ret\$df2, digits=ret\$df2.digits)

| | Point estimate | 95%CI-ll | 95%CI-ul |
|-----------------|----------------|----------|----------|
| Odds ratio | 78.58 | 31.45 | 217.15 |
| Attr. frac. ex. | 0.99 | 0.97 | 1.00 |
| Attr. frac. pop | 0.92 | NA | NA |
| chi2(1) | 164.83 | NA | NA |
| Pr>chi2 | 0.000 | NA | NA |

CC ill - beer

```
result <- CC(DF, ill, beer, exact = TRUE)
kable(result$df1, align=result$df1.align)</pre>
```

| | Cases | Controls | Total |
|--------------------|-------|----------|-------|
| Exposed | 30 | 76 | 106 |
| Unexposed | 69 | 96 | 165 |
| Total | 99 | 172 | 271 |
| Proportion exposed | 0.30 | 0.44 | 0.39 |

kable(result\$df2, digits=result\$df2.digits)

Warning in rep(digits, length.out = m): 'x' is NULL so the result will be ## NULL

| | Point estimate | 95%CI-ll | 95%CI-ul |
|-----------------|----------------|----------|----------|
| Odds ratio | 0.55 | 0.31 | 0.95 |
| Prev. frac. ex. | 0.45 | 0.05 | 0.69 |
| Prev. frac. pop | 0.20 | NA | NA |
| chi2(1) | 5.09 | NA | NA |
| Pr>chi2 | 0.024 | NA | NA |
| Fisher p.value | 0.028 | NA | NA |

#result\$st