Thomas Clarke Quantum Computing Technical Foundations September 15, 2022

### **Assignment 1: Maths for Quantum Computing**

Assignment Due: Monday 12th September

Solutions can be handwritten on a separate sheet of paper, typed or done on a tablet. You may print this, write the solutions on it, and then scan and upload it.

Send the completed assignment to tclarke@asesoftware.com If you have any questions or difficulties, please do reach out to the same email.

Challenge Questions are Optional

#### 1. Complex Numbers

#### Question 1. Complex number algebra

Simplify the following into the form a + bi

- 1) (6+4i)+(3+5i)
- 2) (-6+4i)+(-3+5i)
- 3) i(2+3i)
- 4) (6+4i)(6-4i)

## Question 2. Complex conjugate

Find the complex conjugate for your answers to the previous question

Hint: the complex conjugate of z = a + ib is  $z^* = a - ib$ 

### Question 3. Euler's identity

Complex numbers can be represented using Euler's Formula as

$$a + bi = e^{i\varphi}$$


Where a, b are real numbers that satisfy  $a^2 + b^2 = 1$ , and  $\varphi$  is an angle in radians  $0 < \varphi \le 2\pi$ . The angle  $\varphi$  is related to a & b through the following relations:

$$a = \cos(\varphi), b = \sin(\varphi)$$

For example, if  $\varphi = \pi/3$ 

$$e^{i\pi/3} = \cos(\pi/3) + i\sin(\pi/3)$$
  
=  $\frac{1}{2}(1 + i\sqrt{3})$ 

This can be represented on the Argand diagram as



Represent the following in the form a + ib

- 1)  $e^{i0}$
- $2) e^{i\pi}$
- 3)  $e^{i\frac{\pi}{2}}$
- 4)  $e^{i\frac{\pi}{4}}$

Sketch all of these on the Argand plane to check your answers

Question 4. Challenge Question 1: Third root of unity

Multiply the following complex numbers

$$e^{i2\pi\frac{1}{3}} \times e^{i2\pi\frac{2}{3}} \times e^{i2\pi\frac{3}{3}}$$

This is extremely easy if you use indices rules.

#### 4.1. Linear Algebra

### Question 5. Vector addition

Compute the following:

$$1) \begin{bmatrix} 1 \\ 2 \end{bmatrix} + \begin{bmatrix} 3 \\ 4i \end{bmatrix}$$

$$2) \ \frac{1}{5} \begin{bmatrix} 3 \\ 7 \end{bmatrix} - \begin{bmatrix} 1 \\ 7 \end{bmatrix}$$

3) 
$$[9 \ 8] - [7 \ 6]$$

### Question 6. Conjugate Transpose

Write down the conjugate transpose for your answers to the previous question

Reminder: for any column vector  $|a\rangle = \begin{bmatrix} a_0 \\ a_1 \end{bmatrix}$ , the conjugate transpose is  $\langle a| = \begin{bmatrix} a_0 * & a_1 * \end{bmatrix}$  Where \* indicates the complex conjugate. The same process goes the other way from row vector back to column.

### Question 7. Inner products

Compute the following inner product  $\langle a|b\rangle$ 

Where 
$$|a\rangle = \begin{bmatrix} 1 \\ -i \end{bmatrix}$$
,  $|b\rangle = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ 

And verify  $\langle b|b\rangle = 1^{-1}$ 

## Question 8. Matrix-vector products

Compute the following matrix-vector products

1) 
$$\frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1\\ 1 & -1 \end{bmatrix} \begin{bmatrix} 1\\ 0 \end{bmatrix}$$
2) 
$$\begin{bmatrix} 0 & 1\\ 1 & 0 \end{bmatrix} \begin{bmatrix} 0\\ 1 \end{bmatrix}$$
3) 
$$\begin{bmatrix} 1 & 0\\ 0 & 1 \end{bmatrix} \begin{bmatrix} \alpha\\ \beta \end{bmatrix}$$

$$2) \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$3) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix}$$

Where  $\alpha, \beta$  can be any complex numbers

# Question 9. Challenge Question 2: Entanglement

Compute the matrix-vector product of  $CNOT_{0,1} |0+\rangle_{1,0}^2 = |\Phi^+\rangle_{1,0}$  (i.e. compute the vector  $|\Phi^+\rangle$  Where

$$CNOT_{0,1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}; |0+\rangle_{1,0} \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}$$

Further challenge:

Express your answer in terms of the vectors

 $<sup>^{1}\</sup>text{Originally this read }\langle a|a\rangle,$  but was changed because  $|a\rangle$  is not normalised

<sup>&</sup>lt;sup>2</sup>Here qubit 0 is the control qubit, qubit 1 is the target qubit. It is better explained in chapter 6.

$$|00\rangle = \begin{bmatrix} 1\\0\\0\\0 \end{bmatrix} |10\rangle = \begin{bmatrix} 0\\1\\0\\0 \end{bmatrix} |01\rangle = \begin{bmatrix} 0\\0\\1\\0 \end{bmatrix} |11\rangle = \begin{bmatrix} 0\\0\\0\\1 \end{bmatrix}$$

This matrix-vector product refers to the process of generating entanglement between two qubits. It will be covered in Week 4: Multiple Qubits

THOMAS CLARKE