



0 0 ECON0021 Microeconometrics Index Splited W56789

Table of Content

Table of Contents

1. Table of Content
2. Week 5: MLE & QMLE
 1. 5 MLE & QMLE
 1. Linear Probability Model
 2. MLE
 1. General (conditional) MLE Steps - My Personal Workflow
 2. General MLE Steps:
 3. Consistency
 4. Score
 5. Fisher Information and AVAR
 6. Hessian and Information Matrix Equality
 3. For Scalars
 4. Quasi MLE
 5. Conditional MLE
 3. Week 6: Binary Choice Model
 1. 6 Binary Choice Model - A
 1. Binary Choice Targets of Interest
 2. Linear Probability Model
 3. Reduced Form Approach

4. Structural Threshold Crossing Models
 1. Threshold Crossing Model
 2. Specifications
 3. Identification
 4. Maximum Likelihood
 5. Marginal Effects
4. Week 7: Random Utility Model
1. 7 Random Utility Model. Welfare Analysis. Binary Choice and Endogeneity - A
 1. Random Utility Model
 1. Latent Utility Formulation
 2. Intra/Inter-Personal Random Utility
 2. Welfare Analysis and Compensating Variation
 1. Framework
 2. Estimation of Compensation Variation
 3. Estimation Procedures in Action
 3. Endogeneity in Binary Regressions
 1. Endogeneity
 2. Bivariate Probit Model
 5. Week 8: Multinomial Choice (Vanilla Mn Logit & Too Flexible Mn Probit)
 1. 8 Multinomial Choice - A
 1. Notations
 2. Multinomial Logit Model (Vanilla, Not Realistic)
 1. Framework
 2. Estimation: MLE
 3. Marginal Effects and Elasticities
 4. Drawback: Substitution Patterns (IIA) and Own-Elasticities
 3. Flexible Substitution Pattern
 1. Multinomial Probit (Too Flexible)
 6. Week 9: More on Multinomial Choice (Nested Logit & Random Coefficients Logit)
 1. 9 More On Multinomial Choice - A
 1. Nested Logit (Tractable way to include correlations in ϵ I)
 1. Setup
 2. Joint CDF
 3. Likelihood and Estimation
 2. Random Coefficients Logit Model (Tractable way to include correlations in ϵ II)
 1. Setup
 2. Likelihood and Estimation
 3. Estimation with Aggregate Data

1. Setup
 2. Estimation

 7. Others
 8. Revisions
-

Week 5: MLE & QMLE

5 MLE & QMLE

Linear Probability Model

- Linear Probability Model:

$$Pr(Y = 1|X) = \mathbb{E}[Y|X] = X'\beta \iff Y = X'\beta + u, \mathbb{E}[u|X] = 0$$

- Problems with Linear Probability Model

- Fitted Pr outside [1, 0]: $\mathbb{E}[Y_i|X_i] = X'_i\beta \notin [0, 1]$
- Constant marginal effects (unrealistic)
- Heteroskedasticity

MLE

- Maximum Likelihood
 - Parametric: the population distribution belongs to a family of distributions fully characterised by a finite number of unknown parameters $\theta \in \mathbb{R}^p$
 - i.e. if we know a finite number of parameters specify the distribution, we know the full distribution
 - We want to estimate θ_0 , which is the true vector of parameters in the DGP
 - Given a sample, MLE Estimator is defined by the parameter value that maximises the likelihood (the probability of observing the sample)

General (conditional) MLE Steps - My Personal Workflow

1. Specify a parametric distribution $f_z(z; \theta)$
2. Specify a functional form for conditional likelihood $f_{Y|X}(y|x; \theta)$
3. Calculate individual log likelihood (pdf/pmf)

$$\log f_{Y|X}(y_i|x_i; \theta)$$

4. Log likelihood function for iid data:

$$\log \mathcal{L}_n(y_i|x_i; \theta) = \sum_{i=1}^n \log f_{Y|X}(y_i|x_i; \theta)$$

5. Calculate the **score** from *individual* likelihood

$$s(y_i, x_i; \theta) = \frac{\partial \log f_{Y|X}(y_i | x_i, \theta)}{\partial \theta}$$

caveat: differentiate to the *exact* parameters of the distribution (e.g. σ^2 instead of σ for a Normal distribution)

6. Obtain the **MLE estimator** $\hat{\theta}$ from FOC: set expectation of score to be 0

$$\begin{aligned} \frac{\partial}{\partial \theta} \mathbb{E}[\log f_{Y|X}(y_i | x_i, \theta_0)] &= \mathbb{E}[s(y_i | x_i; \theta_0)] = 0 \\ \implies 0 &= \frac{\partial}{\partial \theta} \left\{ \frac{1}{n} \sum_{i=1}^n \log f_{Y|X}(y_i | x_i; \hat{\theta}_{MLE}) \right\} = \frac{1}{n} \sum_{i=1}^n s(y_i | x_i; \hat{\theta}_{MLE}) = 0 \end{aligned}$$

(alternatively: use FOC: $\frac{\partial \log \mathcal{L}_n(Y_i | x_i; \theta)}{\partial \theta} = 0$)

7. Calculate **Hessian** from score

$$H(y_i, x_i; \theta) = \frac{\partial s(y_i, x_i; \theta)}{\partial \theta'}$$

8. Calculate **Fisher Information Matrix** from Hessian using the Information Matrix Equality (not valid in QMLE)

$$\mathcal{I}(\theta) = -\mathbb{E}_\theta[H(y_i, x_i; \theta)]$$

9. Calculate **asymptotic variance** from Hessian (use true parameter θ_0 here!)

$$Avar\left(\sqrt{n}\left(\hat{\theta}_{MLE} - \theta_0\right)\right) = \mathcal{I}(\theta_0)^{-1}; \widehat{Avar}\left(\hat{\theta}_{MLE} - \theta_0\right) = \frac{\mathcal{I}(\hat{\theta}_0)^{-1}}{n}$$

10. Obtain asymptotic distribution

$$\sqrt{n}(\hat{\theta}_{MLE} - \theta) \xrightarrow{d} \mathcal{N}(0, \mathcal{I}(\theta_0)^{-1})$$

General MLE Steps:

1. Specify a parametric distribution $f_z(z; \theta)$
2. Calculate the joint likelihood function (sample likelihood)

$$\mathcal{L}_n(z; \theta) = \prod_{i=1}^n f_z(z_i, \theta)$$

another way is to calculate the log of individual likelihood, and then sum up

3. Take log (monotonic transformation)

$$\log \mathcal{L}_n(z; \theta) = \sum_{i=1}^n \log f_z(z_i, \theta)$$

4. Maximise with respect to θ (often take FOCs)

$$\hat{\theta}_{MLE} = \arg \max_{\theta \in \Theta} \log \mathcal{L}_n(z, \theta)$$

caveat: differentiate to the *exact* parameters of the distribution (e.g. σ^2 instead of σ for a Normal distribution)

5. Calculate the Fisher Information Matrix \rightarrow Asymptotic distribution of MLE using following step below

Consistency

- Consistency

- Identification (Requirement): different parameters generate different distributions:

$$\theta \neq \theta' \implies f_z(z, \theta) \neq f_z(z, \theta')$$

- If identification hold, $\theta_0 = \arg \max_{\theta \in \Theta} \mathbb{E}[\log f_Z(z_i, \theta)]$ uniquely maximises $\mathbb{E}[\log f_z(z_i; \theta)]$ (a mild additional requirement: $\mathbb{E}[\log f_z(z_i; \theta)] < \infty$) and its sample analogue:

$$\hat{\theta}_{MLE} = \arg \max_{\theta \in \Theta} \frac{1}{n} \sum_{i=1}^n \log f_z(z_i, \theta) \xrightarrow{p} \arg \max_{\theta \in \Theta} \mathbb{E}[\log f_z(z_i, \theta)]$$

i.e. $\hat{\theta}_{MLE} \xrightarrow{p} \theta_0$

Score

- Score

- Score

$$s(z_i, \theta) = \frac{\partial \log f_z(z_i, \theta)}{\partial \theta} = \begin{bmatrix} \frac{\partial \log f_z(z_i, \theta)}{\partial \theta_1} \\ \vdots \\ \frac{\partial \log f_z(z_i, \theta)}{\partial \theta_p} \end{bmatrix}_{p \times 1}$$

- FOC implies that (for population)

$$0 = \frac{\partial}{\partial \theta} \mathbb{E}[\log f_z(z_i, \theta_0)] = \mathbb{E}[s(z_i, \theta_0)]$$

for sample:

$$0 = \frac{\partial}{\partial \theta} \left\{ \frac{1}{n} \sum_{i=1}^n \log f_z(z_i, \hat{\theta}_{MLE}) \right\} = \frac{1}{n} \sum_{i=1}^n s(z_i, \hat{\theta}_{MLE})$$

- If the samples perfectly fit the estimated distribution, then each score = 0

Fisher Information and AVAR

- Fisher Information and Asymptotic Variance

- Fisher Information Matrix is the expectation of the outer product of the score vector:

$$\mathcal{I}_{p \times p}(\theta) = \mathbb{E}_\theta[s(z_i, \theta)s(z_i, \theta)'] = \text{Var}_\theta[s(z_i, \theta)] = \int s(z_i, \theta)s(z_i, \theta)'f_z(z_i, \theta)dz_i$$

- Asymptotic variance of MLE is the inverse of the information matrix evaluated at the *true* θ_0 :

$$Avar\left(\sqrt{n}\left(\hat{\theta}_{MLE} - \theta_0\right)\right) \xrightarrow{p} \mathcal{I}(\theta_0)^{-1}$$

and the Asymptotic Distribution of MLE is

$$\sqrt{n}(\hat{\theta}_{MLE} - \theta_0) \xrightarrow{d} \mathcal{N}(0, \mathcal{I}(\theta_0)^{-1})$$

Hessian and Information Matrix Equality

- Hessian and Information Matrix Equality

- Hessian

$$H(z_i, \theta) = \frac{\partial^2 \log f_z(z_i, \theta)}{\partial \theta \partial \theta'} = \begin{bmatrix} \frac{\partial^2 \log f_z(z_i, \theta)}{\partial \theta_1^2} & \dots & \frac{\partial^2 \log f_z(z_i, \theta)}{\partial \theta_1 \partial \theta_p} \\ \dots & \dots & \dots \\ \frac{\partial^2 \log f_z(z_i, \theta)}{\partial \theta_p \partial \theta_1} & \dots & \frac{\partial^2 \log f_z(z_i, \theta)}{\partial \theta_p^2} \end{bmatrix}$$

- and note that Hessian is the derivative of Score:

$$H(z_i, \theta) = \frac{\partial}{\partial \theta'} s(z_i, \theta)$$

- Information Matrix Equality: Fisher Information Matrix = -Expectation of Hessian

$$\mathcal{I}(\theta) = -\mathbb{E}_\theta[H(z_i, \theta)] = -\int H(z_i, \theta)f_z(z_i, \theta)dz_i$$

- Inference

- $\sqrt{n}(\hat{\theta}_{MLE} - \theta_0) \xrightarrow{d} \mathcal{N}(0, \mathcal{I}(\theta_0)^{-1})$

- We can estimate $\mathcal{I}(\theta_0)$ by:

1. $\mathcal{I}(\hat{\theta}) = -\frac{1}{n} \sum_{i=1}^n H(z_i, \hat{\theta}_{MLE})$ (Information Matrix Equality)

2. or $\mathcal{I}(\hat{\theta}) = \frac{1}{n} \sum_{i=1}^n s(z_i, \hat{\theta}_{MLE})s(z_i, \hat{\theta}_{MLE})'$

- and calculate $\widehat{Avar}(\hat{\theta}_{MLE}) = \hat{I}(\theta_0)^{-1}$

For Scalars

- Score: $s(z_i, \theta) = \frac{\partial \log f_z(z_i, \theta)}{\partial \theta}$
- Hessian: $H(z_i, \theta) = \frac{\partial^2 \log f_z(z_i, \theta)}{\partial \theta^2} = \frac{\partial s(z_i, \theta)}{\partial \theta}$

- Fisher Information Matrix: $\mathcal{I}(\theta) = \underbrace{\text{Var}_{\theta}[s(z_i, \theta)]}_{=E_{\theta}[(s(z_i, \theta))^2]} = \underbrace{-\mathbb{E}_{\theta}[H(z_i, \theta)]}_{\text{Information Matrix Equality}}$
- Asymptotic Variance: $Avar\left(\sqrt{n}\left(\hat{\theta}_{MLE} - \theta_0\right)\right) = \mathcal{I}(\theta_0)^{-1}$

Quasi MLE

- Quasi Maximum Likelihood Estimator (QMLE)
- $\sqrt{n}\left(\hat{\theta}_{MLE} - \theta_0\right) \xrightarrow{d} \mathcal{N}(0, \Sigma)$

where

$$\Sigma = \{\mathbb{E}[H(z_i, \theta_0)]\}^{-1} \text{Var}[s(z_i, \theta_0)] \{\mathbb{E}[H(z_i, \theta_0)]\}^{-1}$$

is the Misspecification Robust SE with its sample analogue:

$$\hat{\Sigma} = \left[\frac{1}{n} \sum_{i=1}^n H(z_i, \hat{\theta}) \right]^{-1} \left[\frac{1}{n} \sum_{i=1}^n s(z_i, \hat{\theta}) s(z_i, \hat{\theta})' \right] \left[\frac{1}{n} \sum_{i=1}^n H(z_i, \hat{\theta}) \right]^{-1}$$

- The *information matrix equality does not hold for QMLE*, so it can be used as a test for misspecification: **information matrix test**:

$$\text{Correct Specification} \implies -\frac{1}{n} \sum_{i=1}^N H(w_i, \hat{\theta}) = \frac{1}{n} \sum_{i=1}^N s(w_i, \hat{\theta}) s'(w_i, \hat{\theta})'$$

Conditional MLE

- **Conditional MLE**: often $z_i = (y_i, x_i')'$ and we only specify $f_{Y|X}(y_i|x_i; \theta)$ (allows for different distributions for different x_i)
- In reality, we make the following assumptions:
 1. Distribution Assumption: specify a distribution $f_z(z_i, \theta)$
 2. Functional Form Assumption: specify $f_{Y|X}(f_i|x_i, \theta)$
- Relationship between con/uncon likelihood/pdf:

$$\underbrace{L_N(x_i, y_i, \theta)}_{\text{Uncon. Likelihood}} = \prod_{i=1}^N \underbrace{f_{XY}(y_i, x_i, \theta)}_{\text{Uncon. PDF}} = \prod_{i=1}^N \underbrace{f_{Y|X}(y_i|x_i, \theta)}_{\text{Cond. PDF}} \times \underbrace{f_X(x_i)}_{\text{Marginal Dist. of } x}$$

- Conditional MLE maximises **conditional log-likelihood** (because $\log f_Z(z_i; \theta) = \log f_{Y|X}(y_i|x_i; \theta) + \log f_X(x_i)$ and $\log f_X(x_i)$ does not depend on θ):

$$\hat{\theta}_{MLE} = \arg \max_{\theta \in \Theta} \sum_{i=1}^n \log f_{Y|X}(f_i|x_i, \theta)$$

- Chain Rule for Vectors

$$\frac{\partial}{\partial \beta} F(x^\top \beta) = x F'(x^\top \beta); \frac{\partial}{\partial \beta^T} F(x^\top \beta) = x^\top F'(x^\top \beta)$$

Week 6: Binary Choice Model

6 Binary Choice Model - A

Binary Choice Targets of Interest

- Binary Choice
- We want to learn:
 - $Pr(Y = 1|X = x) = \mathbb{E}[Y|X = x]$: the conditional probability given an individual's characteristics
 - $\frac{\partial Pr(Y=1|X=x)}{\partial x_l}$: the partial marginal effect of changes in x_l

Linear Probability Model

- Linear Probability Model:

$$y_i = x_i' \beta + \epsilon_i, \mathbb{E}[\epsilon_i|x_i] = 0 \quad \text{or} \quad \mathbb{E}[y_i|x_i] = x_i' \beta$$

- Note that because y_i is binomial (conditional on x_i), so:

$$\begin{cases} x_i' \beta = Pr(y_i = 1|x_i) \\ Var[\epsilon_i|x_i] = Pr(y_i = 1|x_i)(1 - Pr(y_i = 1|x_i)) \end{cases}$$

- Problems: Heteroskedasticity, Out-of-bound Predictions, Constant Marginal Effects

Reduced Form Approach

- Reduced Form Approach
- Use some increasing functions $\mathcal{F} : \mathbb{R} \rightarrow [0, 1]$ to model:

$$Pr(y_i = 1|x_i) = \mathcal{F}(x_i' \beta)$$

- A natural choice is the CDF: $F(z) = Pr(\epsilon \leq z)$ of a random variable ϵ , and the **Conditional Likelihood** takes the form of:

$$Pr(y_i|x_i, \beta) = F(x_i' \beta)^{y_i} [1 - F(x_i' \beta)]^{1-y_i}$$

- ↓ adds economics to such approach

Structural Threshold Crossing Models

Threshold Crossing Model

- **Threshold Crossing Model:** Suppose

$$y_i^* = x_i' \beta - \epsilon_i, \quad y_i = \begin{cases} 1 & \text{if } y_i^* \geq 0 \\ 0 & \text{if } y_i^* < 0 \end{cases}$$

then

$$Pr(y_i = 1|x_i) = F(x_i' \beta)$$

where F is the CDF for ϵ

- Conditional likelihood takes the form of:

$$Pr(y_i|x_i, \beta) = F(x_i' \beta)^{y_i} [1 - F(x_i' \beta)]^{1-y_i}$$

- This can be motivated by the [Random Utility Model](#) (an example of threshold crossing model):

$$Pr(y_i = 1|x_i) = Pr(\underbrace{U_{i1} - U_{i0}}_{y^*} \geq 0|x_i)$$

where U_{i1} and U_{i0} are utilities for $y_i = 1$ and $y_i = 0$

Specifications

- **Specifications**
- **Probit Model:** $\epsilon \sim \mathcal{N}(0, 1)$ with CDF $F(z) = \Phi(z)$:

$$Pr(y_i = 1|x_i) = \Phi(x_i' \beta) \text{ where } \Phi(z) = \int_{-\infty}^z \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{z^2}{2}\right) dz$$

- **Logit Model:** $\epsilon \sim$ Logistic distribution with CDF $F(z) = \Lambda(z)$:

$$Pr(y_i = 1|x_i) = \Lambda(x_i' \beta) = \frac{\exp(x_i' \beta)}{1 + \exp(x_i' \beta)} \text{ where } \Lambda(z) = \frac{\exp(z)}{1 + \exp(z)}$$

- Logistic distribution is similar to Normal distribution except with $Var = \frac{\pi^2}{3}$

Identification

- **Identification:** we must impose location and scale normalisations to fit the logit/probit
- Our actual model could be:

$$y_i^* = \beta_0 + x_i' \beta - \epsilon_i, \quad y_i = \begin{cases} 1 & \text{if } y_i^* \geq 0 \\ 0 & \text{if } y_i^* < 0 \end{cases}$$

where $E[\epsilon_i] = \mu$, $Var[\epsilon_i] = \sigma^2$ and its distribution is specified

- **Location normalisation:** adding the same constant to β_0 and to ϵ_i will not change y_i , so there is no generality to assume $\mu = 0$
- **Scale normalisation:** multiplying (β_0, β') and ϵ_i by the same positive constant will not change y_i , so there is no loss of generality to assume $\sigma = 1$

- Therefore, β_0, β and ϵ_i are not uniquely identified, and it's meaningless to look/compare at their corresponding estimates directly. (Note that the sign does have an immediate interpretation)

Maximum Likelihood

- Maximum Likelihood
- Likelihood:

$$L_n(\beta) = \prod_{i=1}^n F(x'_i \beta)^{y_i} [1 - F(x'_i \beta)]^{1-y_i}$$

- Log-likelihood:

$$\log L_n(\beta) = \sum_{i=1}^n y_i \log F(x'_i \beta) + (1 - y_i) \log [1 - F(x'_i \beta)]$$

- FOC:

$$0 = \frac{\partial \log L_n(\hat{\beta})}{\partial \beta} = \sum_{i=1}^n \frac{F'(x'_i \hat{\beta})}{F(x'_i \beta)(1 - F(x'_i \beta))} (y_i - F(x'_i \beta)) x_i$$

- Logit FOC:

$$0 = \sum_{i=1}^n (y_i - \Lambda(x'_i \hat{\beta})) x_i$$

Marginal Effects

- Marginal Effects
 - Conditional Probability

$$Pr(y_i = 1|x_i) = \begin{cases} x'_i \beta & \text{, for LPM} \\ \underbrace{F(x'_i \beta_l)}_{cdf} = \begin{cases} \Phi(x'_i \beta) & \text{, for Probit} \\ \Lambda(x'_i \beta) = \frac{\exp(x'_i \beta)}{1 + \exp(x'_i \beta)} & \text{, for Logit} \end{cases} & \end{cases}$$

- Marginal Effects for continuous x_l :

$$\frac{\partial Pr(y_i = 1|x_i)}{\partial x_{il}} = \begin{cases} \beta_l & \text{, for LPM} \\ \underbrace{f(x'_i \beta) \beta_l}_{pdf} = \begin{cases} \phi(x'_i \beta) \beta_l & \text{, for Probit} \\ \frac{\exp(x'_i \beta)}{[1 + \exp(x'_i \beta)]^2} \beta_l & \text{, for Logit} \end{cases} & \end{cases}$$

- Marginal Effect for Dummy x_l :

$$Pr(y_i = 1|x_{il} = 1, x_{i,others}) - Pr(y_i = 1|x_{il} = 0, x_{i,others})$$

- Average Marginal Effects

- Since marginal effects are heterogenous, and magnitude of β_l has no immediate interpretation (because of Identification Issue), we often report average marginal effects:

$$\mathbb{E} \left[\frac{\partial \Pr(y_i = 1 | x_i)}{\partial x_{il}} \right] = \mathbb{E}[f(x'_i \beta) \beta_l]$$

with its sample analogue:

$$\frac{1}{n} \sum_{i=1}^n f(x_i \hat{\beta}) \hat{\beta}_l$$

Week 7: Random Utility Model

7 Random Utility Model. Welfare Analysis. Binary Choice and Endogeneity - A

Random Utility Model

Latent Utility Formulation

- **Random Utility Hypothesis:** An individual has random utility that is drawn from a certain distribution each time a decision is made, and the individual makes a choice of $Y = 0$ or $Y = 1$ depending on which option provides a higher utility to her/him
- **Latent Utility Formulation:**

$$\begin{cases} U_1 = u_1(X) + \epsilon_1 \\ U_2 = u_0(X) + \epsilon_0 \end{cases}$$

where $u_1(X) = X' \gamma_1$, $u_0(X) = X' \gamma_0$ are non-random/deterministic latent utility dependent on observable characteristics X , and ϵ_1, ϵ_0 are random components of utility

- Then the choice probability of an individual with characteristic $X = x$ is:

$$\begin{aligned} \Pr(Y = 1 | X = x) &= \Pr(U_1 \geq U_0 | X = x) \\ &= \Pr(\underbrace{x'(\gamma_1 - \gamma_0)}_{\beta} \geq \underbrace{\epsilon_0 - \epsilon_1}_{\Delta\epsilon} | X = x) \\ &= \Pr(x' \beta \geq \Delta\epsilon | X = x) \end{aligned}$$

- assume $\Delta\epsilon \perp X$, this equals to $CDF_{\Delta\epsilon}(x' \beta)$
- with further distribution assumptions of $\Delta\epsilon$:

$$\Pr(Y = 1 | X = x) = \begin{cases} \Phi(x' \beta) & , \text{if } \epsilon_0, \epsilon_1 \sim \mathcal{N}(0, \frac{1}{2}) \rightarrow \Delta\epsilon \sim \mathcal{N}(0, 1) \\ \Lambda(x' \beta) & , \text{if } \epsilon_0, \epsilon_1 \sim \text{Type I extreme value} \rightarrow \epsilon \sim \text{Logit} \end{cases}$$

- **Exogeneity:** $\epsilon_1, \epsilon_2 \perp X$ (Statistical independence)

- **Identification:** parameters appear in both equations will be jointly identified (their differences are identified); *only parameters appear in only one of the equations will be uniquely identified*

Intra/Inter-Personal Random Utility

- Intra- or Inter-personal random utility
- **Intra-individual random utility:** Even under a given choice situation and his/her characteristics, his/her choice of Y is not deterministic due to the random utility (random state of mind)
- **Inter-individual random utility:** Each individual has a nonrandom choice at each choice situation, while individuals with the same observed characteristics can make different choices due to heterogeneity in tastes.
- With cross-sectional data, those two are indistinguishable.

Welfare Analysis and Compensating Variation

Framework

- Welfare Analysis
- Consumer i choose between options $j \in \{0, 1\}$:

$$U_{ij} = u_j(I_i - p_{ij}, x_{ij}) + \epsilon_{ij}$$

where I_i is income, p_{ij} is the price of option j , x_{ij} are consumer and products' characteristics

- Before the intervention, the utility of consumer i is:

$$\max_{j \in \{0,1\}} \{u_j(I_i - p_{ij}, x_{ij}) + \epsilon_{ij}\}$$

- Intervention: new prices $p_{ij} \rightarrow \tilde{p}_{ij}$ and new characteristics $x_{ij} \rightarrow \tilde{x}_{ij}$
- Then **Compensation Variation** $CV_i(I_i, p_i, x_i, \epsilon_i)$ is the change in income needed to hold utility at its initial level:

$$\max_{j \in \{0,1\}} \{u_j(I_i - p_{ij}, x_{ij}) + \epsilon_{ij}\} = \max_{j \in \{0,1\}} \{u_j(I_i + CV_i - \tilde{p}_{ij}, \tilde{x}_{ij}) + \epsilon_{ij}\}$$

- Compensation variations are paid *ex-ante* (before choices are made)!

Estimation of Compensation Variation

- Deal with heterogeneity in ϵ_i : For every individual i, we compute:

$$\begin{aligned} \mathbb{E}[CV_i|I_i, p_i, x_i] &= \mathbb{E}[CV(I_i, p_i, x_i, \epsilon_i)|I_i, p_i, x_i] \\ &= \int CV(I_i, p_i, x_i, \epsilon_i) f_\epsilon(\epsilon) d\epsilon \end{aligned}$$

- This is often approximated by numerical integration ([Monte-Carlo average](#)):

$$\frac{1}{S} \sum_{s=1}^S CV(I_i, p_i, x_i, \epsilon_i^s)$$

where $\epsilon_e^s = (\epsilon_0^s, \epsilon_1^s)'$ are i.i.d. draws from the underlying distribution (e.g. probit $\mathcal{N}(0, \frac{1}{2})$ or Type-1 extreme-value dist.) for a large number S

- With *known utility functions*, we *divide consumers according to their choice before/after intervention, and calculate the CV one by one*

- Deal with heterogeneity in individual characteristics I_i, p_i, x_i : LIE

$$\mathbb{E}[CV_i] = \mathbb{E}[\mathbb{E}[CV_i | I_i, p_i, x_i]]$$

- Thus, we can estimate $\mathbb{E}[CV_i]$ by:

$$\frac{1}{n} \sum_{i=1}^n \left\{ \underbrace{\frac{1}{S} \sum_{s=1}^S CV(I_i, p_i, x_i, \epsilon_i^s)}_{\hat{\mathbb{E}}[CV_i | I_i, p_i, x_i]} \right\}$$

Estimation Procedures in Action

- Estimate all parameters by MLE
- Divide individuals into: always takers of each option and switchers
- Calculate the threshold of $\Delta\epsilon_i$ for each kind of individuals
- Calculate the *CV* for each kind of individuals
 - For switchers, we need them to have the same utility with new options as before with old options
- Calculate with estimated model parameters:

$$\hat{\mathbb{E}}[CV_i | x_i] = \int CV(x_i) f_\epsilon(\epsilon) d\epsilon$$

- Calculate:

$$\mathbb{E}[CV_i] = \frac{1}{n} \sum_{i=1}^n \hat{\mathbb{E}}[CV_i | x_i]$$

Endogeneity in Binary Regressions

Endogeneity

- *Binary regression with Endogeneity*: individual observable characteristics X and the unobservable U are correlated:

$$Y = \mathbb{1}[\beta X + U \geq 0], X \not\perp U$$

- Example: C is endogenous (selection on unobservable that affects Y)

$$Y = \mathbb{1}[\alpha C + \beta' W + U \geq 0], C \not\perp U$$

Bivariate Probit Model

- Our model is:

$$\begin{cases} Y = \mathbb{1}[\alpha C + \beta' W + U \geq 0] \\ C = \mathbb{1}[\gamma' Z + V \geq 0] \end{cases}$$

where $\dim(Z) > \dim(W)$: Z should contain at least 1 **Instrumental Variable** that is not included in W : it provides exogenous variations in regressor C without causally affecting the outcome of interest Y (The instrument has to satisfy:

1. **Exclusion**: $IV_i \perp (U_i, V_i)$
2. **Relevance**: IV_i affects C_i

- Assume:

$$\begin{pmatrix} U \\ V \end{pmatrix} \sim \mathcal{N} \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 & \rho \\ \rho & 1 \end{pmatrix} \right)$$

where $\rho \neq 0$ (otherwise C will be exogenous)

- **Likelihood of Bivariate Probit**: $p(Y, C|W, Z; \theta)$ here $\theta = (\alpha, \beta, \gamma, \rho)'$
 - Conditional on W, Z , the realisation of (Y, C) is determined by realisation of (U, V)

(Y, C)	Threshold	Likelihood
$(0,0)$	$U \leq -\beta' W, V \leq -\gamma' Z$	$p_{00}(W, Z; \theta) = \int_{-\infty}^{-\beta' W} \int_{-\infty}^{-\gamma' Z} f_{U,V}(u, v; \rho) du dv$
$(1,0)$	$U \geq -\beta' W, V \leq -\gamma' Z$	$p_{10}(W, Z; \theta) = \int_{-\beta' W}^{+\infty} \int_{-\infty}^{-\gamma' Z} f_{U,V}(u, v; \rho) du dv$
$(0,1)$	$U \leq -\alpha - \beta' W, V \geq -\gamma' Z$	$p_{01}(W, Z; \theta) = \int_{-\infty}^{-\alpha - \beta' W} \int_{-\gamma' Z}^{+\infty} f_{U,V}(u, v; \rho) du dv$
$(1,1)$	$U \geq -\alpha - \beta' W, V \geq -\gamma' Z$	$p_{11}(W, Z; \theta) = \int_{-\alpha - \beta' W}^{+\infty} \int_{-\gamma' Z}^{+\infty} f_{U,V}(u, v; \rho) du dv$

- Joint Likelihood Function:

$$L_n(\theta) = \prod_{i=1}^n \{p_{11}(w_i, z_i; \theta)^{y_i c_i}\} \{p_{10}(w_i, z_i; \theta)^{y_i(1-c_i)}\} \{p_{01}(w_i, z_i; \theta)^{(1-y_i)c_i}\} \{p_{00}(w_i, z_i; \theta)^{(1-y_i)(1-c_i)}\}$$

and estimate $\theta = (\alpha, \beta, \gamma, \rho)'$ by MLE

- We can then estimate the Average Partial Effect:

$$\frac{1}{n} \sum_{i=1}^n \left(\hat{Pr}(Y = 1|C = 1, w_i) - \hat{Pr}(Y = 1|C = 0, w_i) \right)$$

Week 8: Multinomial Choice (Vanilla Mn Logit & Too Flexible Mn Probit)

8 Multinomial Choice - A

Notations

- W_i are individual-specific characteristics
- W_{ij} are individual- and choice-specific characteristics
- $X_i = (W_i, W_{i1}, \dots, W_{iJ})'$

Multinomial Logit Model (Vanilla, Not Realistic)

Framework

- Multinomial Logit Model
- Latent Utility Formulation: The utility that individual i would get from choice j :

$$U_{ij} = V_j(X_i) + \epsilon_{ij} = V_{ij} + \epsilon_{ij}$$

- Assume: $\epsilon_{ij} \sim (\text{i.i.d. in } i \text{ and } j)$ Type-I extreme value distribution:

$$\begin{aligned} Pr(Y_i = j | X_i = x) &= Pr(V_{ij} - V_{ik} \geq \epsilon_{ik} - \epsilon_{ij} \forall k | X_i = x) \\ &= \frac{\exp(V_{ij})}{\sum_{k=1}^J \exp(V_{ik})} \\ &= \frac{\exp(V_{ij} - V_{iJ})}{1 + \sum_{k=1}^{J-1} \exp(V_{ik} - V_{iJ})} \end{aligned}$$

- 如果所有选项的 V 有common term, 可以在第一步 \exp 之后就约掉

- Parameterise $V_{ij} = \alpha_j + W_i' \beta_j + W_{ij}' \gamma$, then:

$$V_{ij} - V_{iJ} = (\alpha_j - \alpha_J) + W_i'(\beta_j - \beta_J) + (W_{ij} - W_{iJ})' \gamma$$

- Naturally, we normalise $\alpha_J = 0, \beta_J = 0$ and, for the outside option: $V_{iJ} = 0$, so:

$$V_{ij} - V_{iJ} = \alpha_j + W_i' \beta_j + W_{ij} \gamma$$

- Note that we cannot incorporate products' characteristics that does not vary over individuals because they will be absorbed by α_j

Estimation: MLE

- Estimation: MLE

- Likelihood function:

$$\begin{aligned} L_n(\theta) &= \prod_{i=1}^n Pr(Y_i = y_i | X_i = x_i; \theta) \\ &= \prod_{i=1}^n \frac{\exp(V_{y_i}(x_i; \theta))}{\sum_{k=1}^J \exp(V_k(x_i; \theta))} \end{aligned}$$

where $\theta = (\alpha_1, \dots, \alpha_{J-1}, \beta_1, \dots, \beta_{J-1}, \gamma')'$

Marginal Effects and Elasticities

- Denote $P_j(x) = Pr(Y_i = j | X_i = x)$
- Marginal effects:

$$\frac{\partial P_j(x)}{\partial w_{kl}} = \begin{cases} \gamma_l P_j(x)(1 - P_j(x)) & , j = k \\ -\gamma_l P_j(x)P_k(x) & , \text{otherwise} \end{cases}$$

- Price elasticities ($w_{kl} = p_k, \gamma_l = -\alpha$):

$$\frac{\partial P_j(x)}{\partial p_k} \frac{p_k}{P_j(x)} = \begin{cases} -\alpha p_j(1 - P_j(x)) & , j = k \\ \alpha p_k P_k(x) & , \text{otherwise} \end{cases}$$

this shows the IIA property

Drawback: Substitution Patterns (IIA) and Own-Elasticities

- Problem 1: Independence from Irrelevant Alternatives (IIA): for any two choices j and k , the ratio of the choice probabilities:

$$\frac{Pr(Y_i = j|x)}{Pr(Y_i = k|x)} = \exp(V_{ij} - V_{ik})$$

does not depend on the characteristics of other alternatives

- Problem 2: Own-elasticities: most markets have many options, so $(1 - P_j(x))$ is close to a constant, so the *own-elasticity will be proportional to price*:

$$\frac{\partial P_j(x)}{\partial p_j} \frac{p_j}{P_j(x)} = -\alpha p_j \underbrace{(1 - P_j(x))}_{\text{close to constant}}$$

i.e. low price \implies low elasticity (not realistic)

Flexible Substitution Pattern

- Introduce correlations among the random components in the utility ϵ_{ij}
- Methods: Multinomial Probit, Nested Logit, Random Coefficient Logit

Multinomial Probit (Too Flexible)

- Very flexible

- Setup:

$$U_{ij} = V_{ij} + \epsilon_{ij}, \quad \epsilon_i = \begin{pmatrix} \epsilon_{i1} \\ \dots \\ \epsilon_{iJ} \end{pmatrix} \sim N(0, \Sigma)$$

where $\Sigma = \begin{pmatrix} 1 & \dots & \dots & \dots \\ \dots & 1 & \dots & \dots \\ \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & 1 \end{pmatrix}_{J \times J}$ (there will be $J \times (J - 1)$ cov.s need to be estimated)

- Individual likelihood

$$\begin{aligned} Pr(Y_j = j | X_i = x) &= Pr(U_{ij} \geq U_{ik} \forall k | X_i = x) \\ &= Pr(\epsilon_{ij} + V_{ij} \geq \epsilon_{ik} + V_{ik} \forall k | X_i = x) \end{aligned}$$

is not available in closed form, and can be only be calculated by Monte-Carlo simulation, which is computationally infeasible due to the high dimensions of Σ

Week 9: More on Multinomial Choice (Nested Logit & Random Coefficients Logit)

9 More On Multinomial Choice - A

Nested Logit (Tractable way to include correlations in ϵ)

- Nested Logit

Setup

- Allows ϵ_{ij} to be correlated within nests but not across nests
- (*drop index i for simplicity*)
- Still assume ϵ_j to follow Type-I Extreme Value distribution with CDF: $F(\epsilon) = \exp(-\exp(-\epsilon))$, but they can be *correlated within nests*

Joint CDF

- Joint CDF for Nested Logit with J nests and K_j options within each nest:

$$\begin{aligned} F(\epsilon_{11}, \dots, \epsilon_{JK_J}) &= \prod_{j=1}^J \exp \left(- \left[\sum_{k=1}^{K_j} \exp \left(-\frac{\epsilon_{jk}}{\tau_j} \right) \right]^{\tau_j} \right) \\ &= \exp \left(- \sum_{j=1}^J \left[\sum_{k=1}^{K_j} \exp \left(-\frac{\epsilon_{jk}}{\tau_j} \right) \right]^{\tau_j} \right) \end{aligned}$$

- τ_j allows ϵ_{jk} to be correlated within nest j with $\rho_j = 1 - \tau_j^2$
 - $\tau_j \uparrow$ correlation \downarrow

- $\tau_j = 1 \implies \epsilon_{jk}$ are independent (like in the multinomial logit) / or there is just 1 option in nest j
- $\tau_j = 0 \implies \epsilon_{jk}$ are perfectly correlated
- We assume/pre-specify the nests (correlation structure) but need to estimate τ 's

Likelihood and Estimation

- Individual Likelihood: For $U_{jk} = V_{jk} + \epsilon_{jk}$, we have:

$$P_{jk} = P_{k|j} \times P_j$$

where

$$\begin{cases} P_{k|j} = \frac{\exp\left(\frac{V_{jk}}{\tau_j}\right)}{\sum_{m=1}^{K_j} \exp\left(\frac{V_{jm}}{\tau_j}\right)} \\ P_j = \frac{\left(\sum_{m=1}^{K_j} \exp\left(\frac{V_{jm}}{\tau_j}\right)\right)^{\tau_j}}{\sum_{l=1}^J \left(\sum_{m=1}^{K_l} \exp\left(\frac{V_{lm}}{\tau_l}\right)\right)^{\tau_l}} \end{cases}$$

- θ (including τ_1, \dots, τ_J) is estimated by MLE
- *IIA no longer holds*
- Analytically and computationally tractable
- This is not a sequential choice!

Random Coefficients Logit Model (Tractable way to include correlations in ϵ II)

- Random Coefficients Logit Model

Setup

- Allows for preferences γ_i to be heterogeneous across consumers:

$$U_{ij} = W_i' \beta_j + W_{ij}' \gamma_i + \epsilon_{ij}$$

and we need to parameterise the distribution of γ_i

- a common choice is $\gamma_i \sim N(\gamma, \Sigma)$ (so λ includes γ, Σ ; θ includes λ, β) with Σ restricted to be diagonal

Likelihood and Estimation

- Conditional on γ_i , this is just the standard multinomial logit:

$$Pr(Y_i = j | x_i, \gamma_i; \beta) = \frac{\exp(W_i' \beta_j + W_{ij}' \gamma_i)}{\sum_{k=1}^J \exp(W_i' \beta_k + W_{ik}' \gamma_i)}$$

- To get $Pr(Y_i = j|x_i; \beta, \lambda)$, we integrate over γ_i :

$$Pr(Y_i = j|x_i; \beta, \lambda) = \int \underbrace{\frac{\exp(W_i' \beta_j + W_{ij}' \gamma_i)}{\sum_{k=1}^J \exp(W_i' \beta_k + W_{ik}' \gamma_i)}}_{Pr(Y_i=j|x_i, \gamma_i; \beta)} f(\gamma; \lambda) d\gamma$$

and is usually calculated by Monte-Carlo integration:

$$\frac{1}{S} \sum_{s=1}^S Pr(Y_i = j|X_i, \gamma_s; \beta)$$

for some large S , where γ_s are iid draws from $F(\lambda)$

- β, λ can be estimated by MLE with likelihood function:

$$L_n(\beta, \lambda) = \prod_{i=1}^n Pr(Y_i = y_i|x_i; \beta, \lambda)$$

- *IIA no longer holds*

Estimation with Aggregate Data

- Estimation with Aggregate Data

Setup

- $U_{ij} = \underbrace{X_j' \gamma + \xi_j}_{\delta_j} + \epsilon_{ij}$
 - X_j includes the price and observed characteristics of product j
 - ξ_j captures unobserved attributes of the product, unquantifiable factors, systematic shocks, etc.
 - ϵ_{ij} are iid Type I Extreme Values
- Apart from J goods, there is an outside good/option with $\delta_0 = 0$

Estimation

- The market shares are given by:

$$s_j = Pr(Y_i = j) = \frac{\exp(\delta_j)}{1 + \sum_{k=1}^J \exp(\delta_k)}$$

- Then:

$$\log(s_j) - \log(s_l) = \log \left(\frac{\frac{\exp(\delta_j)}{1 + \sum_{k=1}^J \exp(\delta_k)}}{\frac{\exp(\delta_l)}{1 + \sum_{k=1}^J \exp(\delta_k)}} \right) = \log \left(\frac{\exp(\delta_j)}{\exp(\delta_l)} \right) = \delta_j - \delta_l$$

- Since (assume with no loss in generality) $\delta_0 = 0$, we can *back out δ_j from the market shares*:

$$\delta_j = \log(s_j) - \log(s_0)$$

and focus on:

$$\delta_j = X'_j \gamma + \xi_j$$

- To *tackle with endogeneity of price* (p_j is likely to be correlated with ξ_j), we need Instrument Variables

- This can be generalised to a Random Coefficient version, but the steps are always: 1. back out δ_j from market shares; 2. Use IV
-

Others

- ECON0021 Week 7 Question
- Estimand is a population concept; Estimator is its finite sample analogue
- Inverse of a 2×2 matrix:

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

- Statistical independence:

$$\begin{aligned} A \perp B &\Leftrightarrow F_{A|B}(a|b) = F_A(a) \\ &\Leftrightarrow Pr(A < a|b) = Pr(A < a) \\ &\Leftrightarrow P(A = a, B = b) = P(A = a)P(B = b) \\ &\Leftrightarrow F_{AB}(a, b) = F_a(a)F_B(b) \\ &\Leftrightarrow f_{AB}(a, b) = f_A(a)f_B(b) \end{aligned}$$

- Conditional Expectation:

$$E[A|B] = \int_{-\infty}^{\infty} af_{A|B}(a|b) da$$

and $f_{A|B}(a|b) = \frac{f_{A,B}(a,b)}{f_B(b)}$

- Central Limit Theorem:** Let $X_i, i = 1, \dots, n$ are random samples independently drawn from a distribution with mean μ and variance σ^2 . Let $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$ be the sample average, then as $n \rightarrow \infty$:

$$\sqrt{n}(\bar{X}_n - \mu) \sim^d \mathcal{N}(0, \sigma^2)$$

- Confidence Interval (95%):

$$\left[-1.96 \frac{\sigma}{\sqrt{n}} + \bar{X}_n < \mu < 1.96 \frac{\sigma}{\sqrt{n}} + \bar{X}_n \right]$$

- OLS Estimand in Matrices:

$$\beta = E[X'X]^{-1}E[X'Y]$$

- OLS Estimator in Matrices:

$$\hat{\beta} = \left(\sum_{i=1}^n X_i X'_i \right)^{-1} \left(\sum_{i=1}^n X_i Y_i \right)$$

- pdf of $\mathcal{N}(0, \sigma^2)$:

$$f_X(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{x^2}{2\sigma^2}\right)$$

- Chain Rule for Vectors

$$\frac{\partial}{\partial \beta} F(x^\top \beta) = x F'(x^\top \beta); \frac{\partial}{\partial \beta^T} F(x^\top \beta) = x^\top F'(x^\top \beta)$$

- Jensen Inequality

- For $f(x) = x^2$, we have a nice proof:

$$Var(X) = E[X^2] - (E[X])^2 > 0 \implies E[X^2] > (E[X])^2$$

Revisions

- pdf of continuous distributed variable:

$$f_x(x) = Pr(x \in (x - \epsilon, x + \epsilon))$$

- conditional to unconditional pdf

$$Pr(X|Y) = \frac{Pr(X, Y)}{Pr(Y)} \rightsquigarrow (\text{apply to cdf and pdf})$$

- expectation as integrals

$$\mathbb{E}[X] = \begin{cases} \sum_k k \cdot Pr(X = k) & \text{for discrete X} \\ \rightarrow \int k \cdot f_X(x = k) dx & \text{as } k \rightarrow \infty \text{ for continuous X} \end{cases}$$

- conditional expectations

$$\mathbb{E}[X|Y] = \int k f_X(x = k) dk$$

- LIE in integrals

$$\mathbb{E}[X] = \mathbb{E}[\mathbb{E}[X|Y]] = \int \int x f_{X|Y}(k|y) dx f_Y dy$$

- *make brief explanations of the steps*

- make a cheatsheet of the full MLE workflow of possible distributions
- MLE: sample likelihood -> take log -> FOC with respect to parameters -> score -> Hessian -> asymptotic Var
- Asymptotic var:

$$AsympVar(X) = \sqrt{n} \cdot Var(x)$$