



## 0 0 ECON0022 Metrics for Macro and Finance Index

### Table of Contents

### Table of Contents

1. Table of Contents
2. Week 1: Basic Time Series and AR Models
  1. 1 1 Basic TS Notations & Autocorrelations
    1. Notations
    2. Autocorrelations
  2. 1 2 OLS Estimators in TS Models
    1. OLS Estimator in AR(1)
      1. Intro and Estimation
      2. Forecasting in AR(1)
    2. OLS Estimator in AR(p)
      1. Intro and Estimation
      2. Forecasting in AR(p)
  3. Week 2: Large-Sample & Inference in Stationary AR Models
    1. 2 1 Asymptotic Theories for Stationary & G.Mixing TS
      1. TS Concepts
        1. Stationarity
        2. Mixing / Weakly Dependent
      2. LLN and CLT for Time Series, HAC SE
    2. 2 2 Asymptotic Theory for OLS Estimator in Stationary AR Models

1. Properties of OLS Estimator in Stationary AR(1) Models
  1. Assumptions
  2. Asymptotic Analysis
2. Testing AR Assumptions
3. Properties of OLS Estimator in Stationary AR(p) Models
  1. Assumptions
  2. Asymptotic Analysis
4. Week 3: Stationarity of AR Process and ADL Models
  1. 3 1 Stationarity of MA & AR Models
    1. Stationarity of Moving Average (MA) Process
      1. MA(1)
      2. MA( $\infty$ )
      3. ARMA Models
    2. Stationarity of AR Process
      1. Stationarity of AR(1)
      2. Stationarity of AR(p)
      3. Primitive Conditions for OLS to be Normal in LN
  2. 3 2 Stationarity of AR Models and ADL Model
    1. Stationarity of AR Models - Detailed Derivation
    2. Autoregressive Distributed Lag (ADL) Models
5. Week 4: Forecasting and Model Selection
  1. 4 1 Forecast Uncertainty and Model Selection
    1. Forecast Uncertainty and Forecast Intervals
      1. Forecast Uncertainty - Concept
      2. Forecast Uncertainty - Estimation
      3. Forecast Intervals
    2. Multi-Step Forecasts
    3. AIC/BIC: Lag Length Selection using Information Criteria
      1. Bayes Information Criterion (BIC)
      2. Akaike Information Criterion (AIC)
  2. 4 2 [Forecasting and Model Selection II] - A
6. Week 5: Trends
  1. 5 1 [Trends I] - A
    1. Trends
    2. Random Walk / Unit Root I(1)
      1. UR/RW in AR(a) and Error Correction
      2. UR/RW in AR(2) Error Correction
      3. UR/RW in AR(p) Error Correction
    3. Problems of Trends

4. Dickey-Fuller (DF) Test for Stochastic Trends
  1. Dickey-Fuller Test
  2. 5 2 [Trends II] - A
    1. Integrated Stochastic Trends (Multiple Unit Roots)
7. Week 6: Asset Return Predictability
  1. 6 1 [Asset Return Predictability I] - A
    1. Basics
    2. Asset Return Predictability & Market Efficiency Hypothesis
    3. Trading Rules
  2. 6 2 [Asset Return Predictability II] - A
    1. Testing Weak Form  $H_{RW}$  against AR(p)
    2. Model-Free Test of Weak Form  $H_{RW}$ 
      1. Ljung-Box Q Test
      2. Variance-Ratio (VR) Test
    3. Testing Semi-Strong  $H_{RW}$  by Regressions
      1. Test based on DL Model
      2. Issues
8. Week 7: Structural Breaks (Another Type of Non-Stationarity)
  1. 7 1 [Structural Breaks I] - A
    1. Structural Breaks
    2. Testing for Structural Break when Break Date is Known
    3. Testing for Structural Breaks when Break Date is Unknown
      1. Quandt Likelihood (QLR) Test for One Break
      2. Testing Structural Breaks at the End of the TS
  2. 7 2 [Structural Breaks II] - A
    1. Testing for Structural Breaks when Break Date is Unknown: QLR Test Part II
      1. Quandt Likelihood (QLR) Test for Multiple Breaks
    2. Rolling-window estimates (brief)
9. Week 8: Estimation of Dynamic Causal Effects
  1. 8 1 [Dynamic Causal Effects I] - A
    1. Dynamic Causal Effects
    2. Distributed Lag Model
      1. DL Model (A Dynamic Model)
      2. DL Assumptions
      3. Properties of OLS under DL Assumptions
    3. Different Kinds of Exogeneity (PPT II)
    4. Standard Errors in ADL Models
    5. Estimators of Cumulative Multipliers and Their SE
10. Week 9: CAPM and APT

1. 9 1 [CAPM] - A
    1. Setup, Notations, and Assumptions
    2. CAPM
    3. Estimation and Testing of CAPM
      1. CAPM with Constant Betas
      4. Reasons for Rejection
  2. 9 2 CCAPM and APT - A
    1. Conditional CAPM (Varying Beta)
      1. Setup
      2. Estimation Method 1: Instrumenting the Betas (Not IV)
      3. Estimation Method 2: Model Time-Variation as Structural Breaks
      4. Estimation Method 3: Rolling-Window Estimator
    2. Arbitrage Pricing Theory (APT)
      1. Setup
      2. Choice of factors:
      3. Testing
      4. Critical Issues
  11. Week 10: Volatility Models
    1. 10 1 [Volatility Models] - A
      1. Heteroskedasticity & Notations
      2. Multiplicative Volatility Models (including all we discussed after)
      3. ARCH Model
        1. ARCH(1)
        2. ARCH(q)
      4. GARCH Model
      5. Forecasting Volatility and Evaluation
        1. One-Step Forecast
        2. Multi-Step Forecast
        3. Forecast Errors
        4. Forecast Evaluation
    2. 10 2 [Volatility Models II] - A
      1. OLS as Gaussian MLE in AR
      2. Gaussian MLE of ARCH(1): Estimation
      3. Inference in ARCH Models Estimated by MLE
        1. Asymptotic Distribution
        2. Hypothesis Testing
  12. Others
-

# Week 1: Basic Time Series and AR Models

## 1 1 Basic TS Notations & Autocorrelations

### Notations

- Basic TS Notations
- Lags, First Difference, Percentage Change

$$\% \Delta Y_t \approx 100 \Delta \log(Y_t) = 100(\log Y_t - \log Y_{t-1})$$

### Autocorrelations

- Population Autocorrelations

$$\begin{aligned}\rho_j &= \frac{Cov(Y_t, Y_{t-j})}{\sqrt{Var(Y_t) \cdot Var(Y_{t-j})}} \\ &= \frac{Cov(Y_t, Y_{t-j})}{Var(Y_t)}\end{aligned}$$

- Sample estimations of Autocorrelations

$$\hat{\rho}_j = \frac{\widehat{Cov}(Y_t, Y_{t-j})}{\widehat{Var}(Y_t)} = \frac{\frac{1}{T} \sum_{t=j+1}^T (Y_t - \bar{Y})(Y_{t-j} - \bar{Y})}{\frac{1}{T} \sum_{t=1}^T (Y_t - \bar{Y})^2}$$

where  $\bar{Y} = \frac{1}{T} \sum_{t=1}^T Y_t$

- This always offers a *consistent* estimator of autocorrelations as long as we have stationarity and mixing, but without enough lags as the actual DGP, we need HAC SE
- (PS1) **Partial Autocorrelations:**  $\phi_p$  in the regression model:

$$Y_t = \mu + \sum_{i=1}^p \phi_i Y_{t-i} + \epsilon_t$$

It measures the correlation between  $Y_t$  and  $Y_{t-p}$  after controlling for  $Y_t$ 's correlation with  $Y_{t-1} \dots Y_{t-p+1}$

## 1 2 OLS Estimators in TS Models

- Casual Inference / Forecasting: OLS gives you the best linear prediction ( $\min MSFE$ ) regardless of actual DGP

### OLS Estimator in AR(1)

#### *Intro and Estimation*

- Intro:

$$Y_t = \mu + \phi Y_{t-1} + \epsilon_t, \quad t = 2, 3, \dots, T$$

- Solving backward:

$$Y_t = \mu \sum_{i=0}^{t-2} \phi^i + \phi^{t-1} Y_1 + \sum_{i=0}^{t-2} \phi^i \epsilon_{t-i}$$

- OLS Estimator in AR(1)

$$\hat{\phi}_{AR1} = \frac{\sum_{t=2}^T (Y_t - \bar{Y})(Y_{t-1} - \bar{Y}_{-1})}{\sum_{t=2}^T (Y_{t-1} - \bar{Y}_{-1})^2}$$

where  $\bar{Y} = \frac{1}{T-1} \sum_{t=1}^T Y_t$  and  $\bar{Y}_{-1} = \frac{1}{T-1} \sum_{t=2}^T Y_{t-1}$

- $\hat{\phi}_{AR1} \approx \hat{\rho}_1$  (we distinguish  $\bar{Y}, \bar{Y}_{-1}$  here, but not in  $\hat{\rho}_1$ )
- Note that:

$$\hat{\mu} = \bar{Y} - \hat{\phi} \bar{Y}_{-1} \approx (1 - \hat{\phi}) \bar{Y} \iff \bar{Y} \approx \frac{\hat{\mu}}{1 - \hat{\phi}}$$

- (PS2) A general property for AR(1):

$$\rho_Y(k) = \phi^k$$

## Forecasting in AR(1)

- 1-period ahead forecast using population coefficients:  $Y_{T+1|T} = \mu + \phi Y_T$
- Feasible 1-period ahead forecast using estimated coefficients:  $\hat{Y}_{T+1|T} = \hat{\mu} + \hat{\phi} Y_T$
- 1-period ahead forecast error:  $Y_{T+1} - \hat{Y}_{T+1|T}$ 
  - Forecast errors are out-of-sample (unobserved); residuals are in-sample (observed)

## OLS Estimator in AR(p)

### Intro and Estimation

- Intro:

$$Y_t = \mu + \sum_{i=1}^p \phi_i Y_{t-i} + \epsilon_t \quad t = p+1, \dots, T$$

- OLS Estimator in AR(p)

$$\hat{\phi}_i = \frac{\sum_{t=p+1}^T \hat{r}_{i,t} Y_t}{\sum_{t=p+1}^T \hat{r}_{i,t}^2}$$

where  $\hat{r}_{i,t} = Y_{i,t} - \sum_{j \neq i} \hat{\alpha}_j Y_{j,t}$  are first-stage residuals (partialling out other lags)

## Forecasting in AR( $p$ )

- Feasible 1-period ahead forecast:

$$\hat{Y}_{T+1|T} = \hat{\mu} + \sum_{i=1}^p \hat{\phi}_i Y_{T+1-i}$$


---

## Week 2: Large-Sample & Inference in Stationary AR Models

### 2.1 Asymptotic Theories for Stationary & G.Mixing TS

#### TS Concepts

##### Stationarity

- Definition: A time series  $\{Y_t : t = 1, 2, \dots\}$  is **stationary** if its probability distribution does not change over time:

$$(Y_s, Y_{s-1}, \dots, Y_{s-p}) =^d (Y_t, Y_{t-1}, \dots, Y_{t-p}) \quad \forall s, t = 1, 2, \dots$$

- Implications:

- $\forall s, t$ ,  $Y_s$  and  $Y_t$  have the same joint distribution:

$$\mathbb{E}[Y_s] = \mathbb{E}[Y_t], \text{Var}[Y_s] = \text{Var}[Y_t] \quad \forall s, t$$

- $(Y_s, Y_{s-k})$  and  $(Y_t, Y_{t-k})$  have the same joint distribution:

$$\text{Cov}(Y_s, Y_{s-k}) = \text{Cov}(Y_t, Y_{t-k}) \quad \forall s, t, k$$

- $\bar{Y} = \frac{\sum_{t=1}^T Y_t}{T}$  is an unbiased estimator of  $\mathbb{E}[Y_t]$ , but NOT necessarily consistent (data can be highly autocorrelated  $\rightsquigarrow \text{Var}(\hat{Y}) \rightarrow \infty$ )

- Theorem:

- If a time series  $\{Y_t\}$  is stationary:
  - Any transformation  $f(Y_t)$  will also be stationary
  - For any  $p \geq 1$ ,  $Z_t = (Y_t, Y_{t-1}, \dots, Y_{t-p})$  will also be stationary

- Stationarity allows for (high) autocorrelations, iid is a special case of stationarity
- A pair of time series  $\{Z_t\} = \{(X_t, Y_t)\}$  is stationary if  $\forall p \geq 0$ , the joint distribution of  $(Z_t, Z_{t-1}, \dots, Z_{t-p})$  does not depend on  $t$

##### Mixing / Weakly Dependent

- **Mixing** restricts the dependence in data, requiring  $\{Y_t\}$  to have "short memory"

- Implication:

- Long run forecasts  $\rightarrow^p$  unconditional mean: for a function of  $y$  ( $\phi(Y)$ ):

$$\phi(Y_{T+k|T}) = \mathbb{E}[\phi(Y_{T+k})|Y_T, Y_{T-1}, \dots] \rightarrow^p \mathbb{E}[\phi_{T+k}] \text{ as } k \rightarrow \infty$$

- Long-run auto-covariance = 0:  $Cov(Y_{T+k}, Y_T) \rightarrow 0$  as  $k \rightarrow \infty$

## LLN and CLT for Time Series, HAC SE

- Law of Large Numbers (LLN) for Stationary and Mixing TS: suppose that  $\{Y_t\}$  is stationary and geometrically mixing with a valid 2nd moment  $\mathbb{E}[|Y_t|] < \infty$ ,

$$\text{as } T \rightarrow \infty, \bar{Y} = \frac{1}{T} \sum_{t=1}^T Y_t \xrightarrow{p} \mu_Y = \mathbb{E}[Y_t]$$

- Central Limit Theorem (CLT) for Stationary and Mixing TS: suppose that  $\{Y_t\}$  is stationary and geometrically mixing with a valid 4th moment  $\mathbb{E}[|Y_t|^4] < \infty$ ,

$$\text{as } T \rightarrow \infty, \sqrt{T}(\bar{Y} - \mu_Y) \xrightarrow{d} \mathcal{N}(0, \bar{\sigma}_Y^2)$$

where  $\bar{\sigma}_Y^2$  is the Long-run Variance:

$$\begin{aligned} \bar{\sigma}_Y^2 &= Var(Y_t) + 2 \sum_{k=1}^{T-1} \frac{T-k}{T} Cov(Y_t, Y_{t-k}) \\ &= Var(Y_t) \left[ 1 + 2 \sum_{k=1}^{T-1} \frac{T-k}{T} \rho_Y(k) \right] \end{aligned}$$

- Long-run Var can be estimated by HAC SE: Newey-West Estimator

$$\hat{\sigma}_{Y,T}^2 = \hat{\sigma}_Y^2 \left[ 1 + 2 \sum_{k=1}^{m-1} \frac{T-k}{T} \hat{\rho}_Y(k) \right]$$

where  $1 \leq m \leq T$  is a truncation parameter (S&W recommends  $0.75T^{\frac{1}{3}}$  and Wooldridge recommends  $0.025T^{\frac{1}{4}}$ )

- A caveat of using CLT/LLN for time-series data: we always use  $\frac{1}{\sqrt{T}}$  or  $\frac{1}{T}$  because we can freely discard segments of sample in time-series asymptotic analysis:

$$\frac{1}{T} \sum_{t=k}^T Y_t = \frac{1}{T} \sum_{t=1}^T Y_t - \underbrace{\frac{1}{T} \sum_{t=1}^k Y_t}_{=0 \text{ as } T \rightarrow \infty} = \frac{1}{T} \sum_{t=1}^T Y_t$$

## 2.2 Asymptotic Theory for OLS Estimator in Stationary AR Models

- Note that these theories also apply to TS regressions with regressors other than lags of  $Y_t$

## Properties of OLS Estimator in Stationary AR(1) Models

### Assumptions

- Because TS data are not iid in general, OLS only has well-defined distribution in large samples

- AR(1).1: Errors are Unpredictable:

$$Y_t = \mu + \phi Y_{t-1} + \epsilon_t$$

with:

- Strong AR(1).1:  $\mathbb{E}[\epsilon_t | Y_{t-1}, Y_{t-2}, \dots] = 0$  if this holds, standard SE can be used (because all autocorrelations of  $\epsilon_t$  are 0), and AR(1) is the optimal model for forecasting
  - Equivalent to  $\mathbb{E}[\epsilon_t | \epsilon_{t-1}, \epsilon_{t-2}, \dots] = 0$  or  $\mathbb{E}[\epsilon_t | Y_{t-1}, \epsilon_{t-1}, Y_{t-2}, \epsilon_{t-2}, \dots] = 0$  or  $\rho_\epsilon(k) = 0 \forall k > 0$
  - Weaker than saying errors are iid (e.g. allowing for heteroskedasticity)
- Weak AR(1).1:  $\mathbb{E}[\epsilon_t | Y_{t-1}] = 0$  if only this holds (e.g. selected less lags than DGP), then HAC SE will be needed, and the model is not optimal for forecasting
- AR(1).2:  $\{Y_t\}$  is Stationary and Geometrically Mixing with a Valid 4th Moment  $\mathbb{E}[||Y_t||^4]$ 
  - This provides access to LLN & CLT
- AR(1).3: Variation in Regressor  $Var(Y_t) > 0$

### Asymptotic Analysis

- Asymptotic theory of OLS in AR(1)
  - Under AR(1).1-3 (*strong AR(1).1*)  $\implies$  AR(1) estimator will be consistent and follow an asymptotic normal distribution, *standard SE* can be used:

$$\sqrt{T}(\hat{\phi} - \phi) \xrightarrow{d} \mathcal{N}\left(0, \frac{\mathbb{E}[(Y_{t-1} - \mu_Y)^2 \epsilon_t^2]}{Var(Y_t)^2}\right)$$

and  $Avar(\hat{\phi}) = \frac{1}{T} \frac{\mathbb{E}[(Y_{t-1} - \mu_Y)^2 \epsilon_t^2]}{Var(Y_t)^2}$

- *Weaker AR(1).1*  $\implies$  AR(1) estimator will still be consistent and follow an asymptotic normal distribution, but *HAC SE* needed: let  $v_t = (Y_{t-1} - \mu_Y)\epsilon_t$ :

$$Avar(\hat{\phi}) = \frac{1}{T} \frac{Var(v_t) \left\{ 1 + 2 \sum_{k=1}^{T-1} \frac{T-k}{T} \rho_{v_t, v_{t-k}} \right\}}{Var(Y_t)^2}$$

- 如果regressor是X的话，那么分母换成 $Var(X_t)^2$

- Proof of theorem:

$$\begin{aligned}
\hat{\phi} &= \frac{\sum_{t=2}^T (Y_{t-1} - \bar{Y}_{-1}) Y_t}{\sum_{t=2}^T (Y_{t-1} - \bar{Y}_{-1})^2} \\
&= \mu \underbrace{\frac{\sum_{t=2}^T (Y_{t-1} - \bar{Y}_{-1})}{\sum_{t=2}^T (Y_{t-1} - \bar{Y}_{-1})^2}}_{=0} + \phi \underbrace{\frac{\sum_{t=2}^T (Y_{t-1} - \bar{Y}_{-1}) Y_{t-1}}{\sum_{t=2}^T (Y_{t-1} - \bar{Y}_{-1})^2}}_{=1} + \frac{\sum_{t=2}^T (Y_{t-1} - \bar{Y}_{-1}) \epsilon_{t-1}}{\sum_{t=2}^T (Y_{t-1} - \bar{Y}_{-1})^2} \\
&= \phi + \frac{\sum_{t=2}^T (Y_{t-1} - \bar{Y}_{-1}) \epsilon_{t-1}}{\sum_{t=2}^T (Y_{t-1} - \bar{Y}_{-1})^2} \\
\implies \sqrt{T}(\hat{\phi} - \phi) &= \frac{\frac{1}{\sqrt{T}} \sum_{t=2}^T (Y_{t-1} - \bar{Y}_{-1}) \epsilon_{t-1}}{\frac{1}{T} \sum_{t=2}^T (Y_{t-1} - \bar{Y}_{-1})^2}
\end{aligned}$$

- With Strong AR(1).1:

$$\mathbb{E}[(Y_{t-1} - \mu_Y) \epsilon_t] = ^{LIE} \mathbb{E}\left[(Y_{t-1} - \mu_Y) \underbrace{\mathbb{E}[\epsilon_t | Y_{t-1}, Y_{t-2}, \dots]}_{=0}\right] = 0$$

- and there will be no covariance terms, so normal SE estimator can be used:

$$\begin{aligned}
\forall k \geq 1 : \mathbb{E}\left[[(Y_{t-1} - \mu_Y) \epsilon_t] \cdot [(Y_{t-k-1} - \mu_Y) \epsilon_{t-k}]\right] \\
&= \mathbb{E}\left[[(Y_{t-1} - \mu_Y) \underbrace{\mathbb{E}[\epsilon_t | Y_{t-1}, Y_{t-2}, \dots]}_{=0}] \cdot [(Y_{t-k-1} - \mu_Y) \epsilon_{t-k}]\right] \\
&= 0
\end{aligned}$$

- Thus:

$$\begin{cases} \frac{1}{\sqrt{T}} \sum_{t=2}^T (Y_{t-1} - \bar{Y}_{-1}) \epsilon_{t-1} & \xrightarrow{d} \mathcal{N}\left(0, \mathbb{E}[(Y_{t-1} - \mu_Y)^2 \epsilon_t^2]\right) \text{ by CLT} \\ \frac{1}{T} \sum_{t=2}^T (Y_{t-1} - \bar{Y}_{-1})^2 & \xrightarrow{p} \text{Var}(Y_t) \text{ by LLN} \end{cases}$$

- By Slutsky's Theorem:

$$\sqrt{T}(\hat{\phi} - \phi) \xrightarrow{d} \mathcal{N}\left(0, \frac{\mathbb{E}[(Y_{t-1} - \mu_Y)^2 \epsilon_t^2]}{\text{Var}(Y_t)^2}\right)$$

- Importance of AR.1: Strong AR(1).1  $\implies$  no covariance term (above)  $\implies$  normal SE estimator can be used
  - If we only have weak AR(1).1, then replace  $\mathbb{E}[(Y_{t-1} - \mu_Y)^2 \epsilon_t^2]$  with HAC SE
- See [OLS in DL](#) for asymptotic analysis of OLS estimators in DL models

## Testing AR Assumptions

- [Testing Strong AR.1 \(Errors are Unpredictable / Zero Conditional Mean\)](#): strong AR.1 implies all autocorrelations in  $\epsilon_t$  are 0, so we can:
  - Plot residuals and look for outliers
  - Check/test autocorrelation in residuals

- Examine their marginal distributions (ideally close to Normal)
- Testing AR.2 (Stationarity): All roots of the characteristic polynomial lie outside the unit circle (see Stationarity of AR Process)

## Properties of OLS Estimator in Stationary AR(p) Models

### Assumptions

- AR(p).1: Errors are Unpredictable  $Y_t = \mu + \sum_{i=1}^p \phi_i Y_{t-i} + \epsilon_t$  with
  - *Strong* AR(p).1:  $\mathbb{E}[\epsilon_t | Y_{t-1}, Y_{t-2}, \dots] = 0$  if this holds, standard se can be used (because all autocorrelations of  $\epsilon_t$  are 0)
  - *Weak* AR(p).1:  $\mathbb{E}[\epsilon_t | Y_{t-1}, \dots, Y_{t-p}] = 0$  if only this holds (e.g. selected less lags than DGP), then HAC SE will be needed
- AR(p).2:  $\{Y_t\}$  is Stationary and Geometrically Mixing with a Valid 4th Moment  $E[|Y_t|^4]$ 
  - This provides access to LLN & CLT
  - Requires all roots lie outside the unit circle and errors are iid PrimitiveConditions
  - No structural breaks
- AR(p).3: Variation in Regressor  $Var(Y_t) > 0$  and there is No Perfect Multicollinearity between  $Y_{t-1}, \dots, Y_{t-p}$

### Asymptotic Analysis

- Asymptotic theory of OLS in AR(p)
  - Under AR(p).1-3 with:
  - Strong AR(p).1  $\implies$  Standard SE:

$$\sqrt{T}(\hat{\phi}_k - \phi_k) \xrightarrow{d} \mathcal{N}\left(0, \frac{\mathbb{E}[r_{k,t}^2 \epsilon_t^2]}{Var(r_{k,t})^2}\right), k = 1, \dots, p$$

and  $Avar(\hat{\phi}_k) = \frac{1}{T} \frac{\mathbb{E}[r_{k,t}^2 \epsilon_t^2]}{Var(r_{k,t})^2}$

- Weaker AR(p).1  $\implies$  HAC SE needed: let  $v_t = r_{k,t} \epsilon_t$ ,

$$Avar(\hat{\phi}) = \frac{1}{T} \frac{Var(v_t) \left\{ 1 + 2 \sum_{k=1}^{T-1} \frac{T-k}{T} \rho_{v_t, v_{t-k}} \right\}}{Var(r_{k,t})^2}$$

- Is Assumption AR.1 (Strong) plausible in AR(p) model?
  - Wold's Decomposition: any stationary TS can be well-approximated (i.e. Strong AR(p).1 satisfied / no autocorrelation in  $\epsilon_t$ ) by an AR(p) if p is large enough

## Week 3: Stationarity of AR Process and ADL Models

### 3 1 Stationarity of MA & AR Models

#### Stationarity of Moving Average (MA) Process

##### MA(1)

- Setup:

$$Y_t = \mu + \epsilon_t + \alpha\epsilon_{t-1}$$

where  $\epsilon_t$  are i.i.d. distributed with  $\mathbb{E}[\epsilon_t] = 0, \text{Var}(\epsilon_t) = \sigma_\epsilon^2$

- Stationarity of MA(1) process:  $(\epsilon_t, \epsilon_{t-1})$  is stationary by construction,  $Y_t$  is a function of  $(\epsilon_t, \epsilon_{t-1})$ , so it's also stationary
- Moments of MA(1) process:

$$\begin{cases} \mathbb{E}[Y_t] &= \mu \\ \text{Var}(Y_t) &= (1 + \alpha^2)\sigma_\epsilon^2 \\ \text{Cov}(Y_t, Y_{t-k}) &= \begin{cases} \alpha\sigma_\epsilon^2 &, k = 1 \\ 0 &, k > 1 \end{cases} \end{cases}$$

##### MA( $\infty$ )

- Setup:

$$Y_t = \mu + \sum_{i=0}^{\infty} \alpha_i \epsilon_{t-i}$$

where  $\sum_{i=1}^{\infty} \alpha_i^2 < \infty$  and  $\{\epsilon_t\}$  are i.i.d. distributed with  $\mathbb{E}[\epsilon_t] = 0, \text{Var}(\epsilon_t) = \sigma_\epsilon^2$

- Stationarity of MA process of infinite order:  $(\epsilon_t, \epsilon_{t-1}, \dots, \epsilon_{t-\infty})$  is stationary by construction,  $Y_t$  is a function of  $(\epsilon_t, \epsilon_{t-1})$ , so it's also stationary
- $\sum_{i=1}^{\infty} \alpha_i^2 < \infty$  ensures MA( $\infty$ ) has a well-defined 2nd moment:

$$\mathbb{E} \left[ \left( \sum_{i=0}^{\infty} \alpha_i \epsilon_{t-i} \right)^2 \right] = \sigma_\epsilon^2 \sum_{i=1}^{\infty} \alpha_i^2 < \infty$$

#### ARMA Models

- ARMA( $p, q$ ):

$$Y_t = \mu + \sum_{i=1}^p \phi_i Y_{t-i} + \epsilon_t + \sum_{i=1}^q \alpha_i \epsilon_{t-i}$$

#### Stationarity of AR Process

##### Stationarity of AR(1)

- Condition:  $|\phi| < 1$  and  $\{\epsilon_t\}$  are i.i.d.

- Under these conditions,  $AR(1) \rightarrow MA(\infty)$  by iterating backwards, hence stationary:

$$\begin{aligned}
Y_t &= \mu + \phi Y_{t-1} + \epsilon_t \\
&= \phi^k Y_{t-k} + \mu \sum_{i=0}^{k-1} \phi^i + \sum_{i=0}^{k-1} \phi^i \epsilon_{t-i} \\
(as k \rightarrow \infty) &= \cancel{\phi^k Y_{t-k}}^0 + \mu \sum_{i=0}^{k-1} \phi^i \overset{\frac{\mu}{1-\phi}}{\cancel{\phi^i}} + \sum_{i=0}^{k-1} \phi^i \epsilon_{t-i} \\
&= 0 + \frac{\mu}{1-\phi} + \underbrace{\sum_{i=0}^{\infty} \phi^i \epsilon_{t-i}}_{MA(\infty)}
\end{aligned}$$

### Stationarity of AR(p)

- Condition:
  - All roots of the Characteristic Polynomial  $\phi(z) = 1 - \phi_1 z - \phi_2 z^2 - \dots - \phi_p z^p$  lie outside the unit circle:  $|z^*| > 1$
  - $\{\epsilon_t\}$  are i.i.d. ( $\Rightarrow$  standard SE)
- Under these conditions,  $AR(p) \rightarrow MA(\infty)$  by iterating backwards, hence stationary:

$$\begin{aligned}
Y_t &= \mu + \sum_{i=1}^p \phi_i Y_{t-i} + \epsilon_t \\
&= \frac{\mu}{1 - \phi_1 - \dots - \phi_p} + \sum_{i=0}^{\infty} \alpha_i \epsilon_{t-1}
\end{aligned}$$

where  $\alpha_1, \alpha_2, \dots$  are such that  $\sum_{i=0}^{\infty} \alpha_i z^i = \frac{1}{\phi(z)}$  and  $\phi(z) = 1 - \phi_1 z - \phi_2 z^2 - \dots - \phi_p z^p$

### Primitive Conditions for OLS to be Normal in LN

- Primitive conditions
  - Roots of  $\phi(z) = 1 - \phi_1 z - \phi_2 z^2 - \dots - \phi_p z^p$  are outside of the unit circle ( $|z^*| > 1$ )
    - Note that "eigenvalues" are reciprocal values of roots, so they need to be within the unit circle
    - Formal test: Dickey-Fuller Test
  - If further: errors  $\epsilon_t$  are i.i.d., then we can use standard SE. Otherwise, HAC SE
    - Check by residual analysis, see Testing AR Assumptions

## 3 2 Stationarity of AR Models and ADL Model

### Stationarity of AR Models - Detailed Derivation

- Stationarity of AR(p) process
- The role of the characteristic polynomial

### Autoregressive Distributed Lag (ADL) Models

- Setup:

$$Y_t = \mu + \sum_{i=1}^p \phi_i Y_{t-i} + \sum_{j=1}^q \psi_j X_{t-j} + \epsilon_t$$

- Stationarity of ADL model:

- All  $\{X_t\}$  needs to be stationary (fit AR models to check, same techniques as below)
  - The AR component needs to be stationary (check roots of characteristic polynomial, test by Dickey-Fuller Test)
  - Estimation and inference for ADL model: if stationary and mixing, OLS follows an asymptotic Normal distribution
  - Granger Causality
    - A F-test:  $H_0$ : coefficients of all value of one of the variable  $X_{a,t-1}, X_{a,t-2}, \dots, X_{a,t-q_a}$  are 0
    - It is a test for marginal predictability (not really causality)
  - Forecasting
- 

## Week 4: Forecasting and Model Selection

### 4.1 Forecast Uncertainty and Model Selection

#### Forecast Uncertainty and Forecast Intervals

##### Forecast Uncertainty - Concept

- Optimality in terms of MSFE:

$$\mathbb{E}[Y|X] = \arg \min_{m(X)} \underbrace{\mathbb{E}[(Y - m(X))^2]}_{MSFE}$$

and OLS consistently estimates the best linear predictor (if  $m(X)$  is linear):

$$\min_{\beta_0, \beta_1} \mathbb{E}[(Y - \beta_0 - \beta_1 X)^2]$$

- MSFE of OLS forecast under \*correct\* specification:
  - A combination of:
    - forecast error distribution ( $\epsilon_{T+1} = Y_{T+1} - Y_{T+1|T}$ )
    - uncertainty from using estimated parameter ( $(Y_{T+1|T} - \hat{Y}_{T+1|T})$ )
  - Total Forecast Error:

$$\begin{aligned} e_{T+1} &= Y_{T+1} - \hat{Y}_{T+1|T} \\ &= \underbrace{Y_{T+1} - Y_{T+1|T}}_{\epsilon_{T+1}} + \underbrace{Y_{T+1|T} - \hat{Y}_{T+1|T}}_{\epsilon_{T+1}} \\ &= \epsilon_{T+1} + (Y_{T+1|T} - \hat{Y}_{T+1|T}) \end{aligned}$$

- Mean Square Forecast Error (MSFE):

$$\begin{aligned}
 MSFE : \mathbb{E}[e_{T+1}^2] &= \mathbb{E}[(Y_{T+1} - \hat{Y}_{T+1|T})^2] \\
 &= \sigma_\epsilon^2 + \underbrace{\mathbb{E}[(Y_{T+1|T} - \hat{Y}_{T+1|T})^2]}_{\text{Error from Parameter Uncertainty}} \\
 &\quad (T \rightarrow \infty) = \sigma_\epsilon^2
 \end{aligned}$$

- RMSFE:  $RMSFE = \sqrt{MSFE}$

## Forecast Uncertainty - Estimation

- Three ways to estimate RMSFE

1. Large sample approximation ( $\widehat{RMSFE} \approx \underbrace{\hat{\sigma}_\epsilon}_{SER}$ )

2. Use actual forecast history (not practical)

3. Pseudo out-of-sample forecasting (POOs)

- Re-estimate the model every period for  $t = t_1, \dots, T-1$
- Compute the forecast for  $t+1$  using model estimated using data up to time  $t$ :  $\hat{Y}_{t+1|t}$
- Compute the poos forecast error:  $Y_{t+1} - \hat{Y}_{t+1|t}$
- POOS MSFE:

$$\widehat{MSFE} = \frac{1}{T-t_1} \sum_{t=t_1}^{T-1} (Y_{t+1} - \hat{Y}_{t+1|t})^2$$

- Idea: re-estimate the model every period from  $t_1$  to  $T-1$ , compute forecast error  $Y_{t+1} - \hat{Y}_{t+1|t}$ , and calculate  $\widehat{MSFE}$

## Forecast Intervals

- Using the RMSFE to construct forecast intervals: If  $\epsilon_{T+1}$  is normally distributed:

$$\begin{cases} 95\% FI : \hat{Y}_{T+1|T} \pm 1.96 \times \widehat{RMSFE} \\ 67\% FI : \hat{Y}_{T+1|T} \pm \widehat{RMSFE} \end{cases}$$

- Forecast intervals are not confidence intervals because  $Y_{T+1}$  is random, and FI is valid only if  $\epsilon_{T+1}$  is normal

## Multi-Step Forecasts

- Iterated multi-period forecast:

$$\begin{aligned}
 \hat{Y}_{T+1|T} &= \hat{\mu} + \hat{\phi} Y_T \\
 \rightsquigarrow \hat{Y}_{T+2|T} &= \hat{\mu} + \hat{\phi} \hat{Y}_{T+1|T} \\
 &\quad \dots \\
 \rightsquigarrow \hat{Y}_{T+h|T} &= \hat{\mu} + \hat{\phi} \hat{Y}_{T+h-1|T}
 \end{aligned}$$

- Direct multi-period forecast: exclude lags of order below  $h$  in the model:

$$\begin{aligned} h = 1 : \hat{Y}_t &= \hat{\mu}_1 + \hat{\phi}_1 Y_{t-1} \implies \hat{Y}_{T+1|T} = \hat{\mu}_1 + \hat{\phi}_1 Y_T \\ h = 2 : \hat{Y}_t &= \hat{\mu}_2 + \hat{\phi}_2 Y_{t-2} \implies \hat{Y}_{T+2|T} = \hat{\mu}_2 + \hat{\phi}_2 Y_T \\ &\dots \end{aligned}$$

- Which to use?

- Iterated method is generally preferred, especially when the original AR(p) model is correctly specified
- In ADL models, direct forecasting may be more suitable because we do not need to build forecast model for each  $X$

## AIC/BIC: Lag Length Selection using Information Criteria

- Typically, we don't shrink model by consecutive F-tests
  - see why not using F-test for details
- Lag Length Selection Using Information Criteria
  - Idea: balance the bias-variance tradeoff: bias/model fit (too few lags) and estimation variance/uncertainty (too many lags)

$$MSFE = \mathbb{E} \left[ \underbrace{(Y_{T+1} - Y_{T+1|T}^*)^2}_{\epsilon_T^2} \right] + \mathbb{E} \left[ (Y_{T+1}^* - \hat{Y}_{T+1|T}^*)^2 \right]$$

- First term in AIC/BIC (related to  $SSR$  or  $MSR$ ) is designed to penalise "bias" / forecast error, which is  $\mathbb{E} \left[ \underbrace{(Y_{T+1} - Y_{T+1|T}^*)^2}_{\epsilon_T^2} \right]$  in MSFE
- Second term (related to number of lags  $p$ ) is designed to penalise estimation uncertainty and overfitting, which is  $\mathbb{E} \left[ (Y_{T+1}^* - \hat{Y}_{T+1|T}^*)^2 \right]$  in MSFE

## Bayes Information Criterion (BIC)

- Bayes Information Criterion (BIC)

$$\begin{aligned} \hat{p}_{BIC} &= \arg \min_p BIC(p) \\ &= \arg \min_p \left\{ \underbrace{\log \left( \frac{SSR(p)}{T} \right)}_{\downarrow \text{in } p} + \underbrace{(p+1) \frac{\log(T)}{T}}_{\uparrow \text{in } p} \right\} \\ &= \arg \min_p \left\{ \underbrace{\log (MSR)}_{\downarrow \text{in } p} + \underbrace{(p+1) \frac{\log(T)}{T}}_{\uparrow \text{in } p} \right\} \end{aligned}$$

- If  $\{Y_t\}$  is indeed generated by an AR(p), then  $\hat{p}_{BIC} \xrightarrow{p} p$

## Akaike Information Criterion (AIC)

- Akaike Information Criterion (AIC)

$$\begin{aligned}\hat{p}_{AIC} &= \arg \min_p AIC(p) \\ &= \arg \min_p \left\{ \underbrace{\log \left( \frac{SSR(p)}{T} \right)}_{\downarrow \text{in } p} + \underbrace{(p+1) \frac{2}{T}}_{\uparrow \text{in } p} \right\} \\ &= \arg \min_p \left\{ \underbrace{\log (MSR)}_{\downarrow \text{in } p} + \underbrace{(p+1) \frac{2}{T}}_{\uparrow \text{in } p} \right\}\end{aligned}$$

- $\hat{p}_{AIC} \geq \hat{p}_{BIC}$
- If the DGP is indeed an AR(p), AIC is not consistent (it *overestimates* lags)
- Drawback of AIC/BIC and the alternative -- POOs (see [Alternative lag length selection](#))
- Generalization of BIC to ADL Models

$$BIC(p) = \log \left( \frac{SSR(p, q)}{T} \right) + (p+q+1) \frac{\log(T)}{T}$$

- SATAT command `varsoc` restricts  $p = q$
- so alternatively, you can: choose lags of Y by BIC, and decide whether X should be included by the Granger causality test
- Choosing the right lag length: Important for inference
  - For *forecasting*, incorrect lag length  $\rightsquigarrow$  HAC SE needed; not optimal forecast; but still consistent as long as the errors are mean-independent
  - For *causal inference / estimating the true parameter of DGP*, incorrect lag length  $\rightsquigarrow$  Bias and inconsistency
    - see [Week 8: Estimating Dynamic Causal Effects](#)

## 4 2 [Forecasting and Model Selection II] - A

- Population forecasts: Choosing loss function: we can choose other loss functions  $\mathcal{L}(Y_{T+1} - Y_{T+1|T})$  other than MSFE
- Optimal forecast:

$$MSFE = \mathbb{E} \left[ \underbrace{(Y_{T+1} - Y_{T+1|T}^*)^2}_{\epsilon_T^2} \right] + \mathbb{E} \left[ (Y_{T+1}^* - \hat{Y}_{T+1|T}^*)^2 \right]$$

- Bias-variance trade-off: shrinking the complexity of possible forecasts (by choosing finite lags / parametric form) will increase the first term ("bias"  $\uparrow$ ) and decrease the second term ("variance/estimation uncertainty"  $\downarrow$ )
- Dimension reduction

- Finite # lags
- Parametric form
- Bayes' and Akaike's Information Criteria
- Why not use F and t test to choose correct model?
  - It relies on repeated hypothesis testing with small effective sample size -- *size control is very difficult* (for  $k$  potential models, we need  $k!$  tests)
  - *Overall confidence level can be very low* (every time we have 5% chance of rejecting a correct null, so probability of type-1 errors will accumulate)  $\rightsquigarrow$  tends to select too many lags
- Alternative model selection rule: Poos:
  - *AIC and BIC both rely on large sample approximation of the exact MSFE*, which could be imprecise in small/moderate samples. Thus, we may consider POOS for small/moderate samples
  - Select a  $t_1$
  - For each model, re-estimate the model every period from  $t_1$  to  $T - 1$ , compute forecast error  $Y_{t+1} - \hat{Y}_{t+1|t}$ , and calculate POOS MSFE:

$$\widehat{\text{MSFE}} = \frac{1}{T - t_1} \sum_{t=t_1}^{T-1} (Y_{t+1} - \hat{Y}_{t+1|t})^2$$

- Compare different models in terms of their  $\widehat{\text{MSFE}}$
- Forecast interval: If  $e_{T,h} \sim \mathcal{N}(0, \sigma_e^2)$ , then  $\hat{e}_{T,h} \sim \mathcal{N}(0, \widehat{\text{MSFE}})$  (approximately) so:

$$Y_{T+1} - \hat{Y}_{T+1|T} \sim \mathcal{N}(0, \widehat{\text{MSFE}})$$

and

$$\begin{cases} 95\% \text{ FI: } \hat{Y}_{T+1|T} \pm 1.96 \widehat{\text{MSFE}} \\ 67\% \text{ FI: } \hat{Y}_{T+1|T} \pm \widehat{\text{MSFE}} \end{cases}$$

- Note that FI is only valid when the distribution of error term is Normal!
  - Iterative Forecasting in STATA
- 

## Week 5: Trends

### 5.1 [Trends I] - A

- Two important types of non-stationarity
  - Trends / Unit roots (this week)
  - Structural breaks ([Week 7: Structural Breaks](#))

### *Trends*

- Trends and Cycles: Most economic TS can be decomposed into:

$$Y_t = \underbrace{\text{Trend}_t}_{\text{non-stationary}} + \underbrace{\text{Cycle}_t}_{\text{stationary}}$$

- Deterministic and stochastic trends
  - Deterministic trend: non-random function of time (time drift)
  - Stochastic trend: random and varies over time (unit roots)

## Random Walk / Unit Root I(1)

### UR/RW in AR(a) and Error Correction

- Setup:

$$Y_t = Y_{t-1} + \epsilon_t, \quad \epsilon_t \text{ are serially uncorrelated}$$

- Two key features of a random walk

- Best prediction is current value:

$$\mathbb{E}[Y_{T+h|T}] = Y_T$$

- Unstable/exploding 2nd moment: suppose  $Y_0 = 0$ ,

$$\text{Var}(Y_t) = t\sigma_\epsilon^2$$

(variance grows linearly with time)

- A random walk with drift:

$$Y_t = \mu + Y_{t-1} + \epsilon_t, \quad \epsilon_t \text{ are serially uncorrelated}$$

- $\mu \neq 0$  induces a linear trend/drift in  $Y$ :  $Y_{T+h|T} = \mu h + Y_T$

- Stochastic trends and unit autoregressive roots:

- Random walk is an AR(1) with  $\phi = 1 \iff$  characteristic polynomial  $\phi(z) = 1 - z \iff \phi(1) = 0$ , so First Differencing it makes it stationary ([Error-correction Representation of AR\(1\)](#)):

$$\Delta Y_t = \mu + \underbrace{\delta}_{=0 \text{ if RW}} Y_{t-1} + \epsilon_t$$

### UR/RW in AR(2) Error Correction

- Unit roots in an AR(2):

$$Y_t = \mu + \phi_1 Y_{t-1} + \phi_2 Y_{t-2} + \epsilon_t$$

with its characteristic polynomial  $\phi(z) = 1 - \phi_1 z - \phi_2 z^2$  has 1 root at  $z = 1$ :

$$\phi(1) = 1 - \phi_1 - \phi_2 = 0 \iff \phi_1 + \phi_2 = 1$$

- Error-correction representation:

- Derivation (最好上来先左右同时减 $Y_{t-1}$ , 然后再从后往前迭) :

$$\begin{aligned}
 Y_t &= \mu + \phi_1 Y_{t-1} + \phi_2 Y_{t-2} + \epsilon_t \\
 &= \mu + \phi_1 Y_{t-1} + \cancel{\phi_2 Y_{t-2}} - \cancel{\phi_2 Y_{t-2}} + \phi_2 Y_{t-2} + \epsilon_t \\
 &= \mu + (\phi_1 + \phi_2) Y_{t-1} + \phi_2 \underbrace{(Y_{t-1} - Y_{t-2})}_{\Delta Y_{t-1}} + \epsilon_t \\
 &= \mu + (\phi_1 + \phi_2) Y_{t-1} + \phi_2 \Delta Y_{t-1} + \epsilon_t \\
 \underbrace{Y_t - Y_{t-1}}_{\Delta Y_t} &= \mu + \underbrace{(\phi_1 + \phi_2 - 1)}_{\delta} Y_{t-1} + \underbrace{\phi_2}_{\gamma_1} \Delta Y_{t-1} + \epsilon_t \\
 \Delta Y_t &= \mu + \delta Y_{t-1} + \gamma_1 \Delta Y_{t-1} + \epsilon_t
 \end{aligned}$$

- Result - Error-correction form of AR(2):

$$\Delta Y_t = \mu + \delta Y_{t-1} + \gamma_1 \Delta Y_{t-1} + \epsilon_t$$

where  $\delta = \phi_1 + \phi_2 - 1, \gamma_1 = -\phi_2$

- Unit root:  $\delta = 0$

- If indeed  $\delta = 0$ , then:

-  $\Delta Y_t$  is stationary  $\iff |\gamma_1| < 1$

- If  $|\gamma_1| = 1$ , then  $\Delta Y_t$  is still non-stationary (the process is integrated of order 2)

### UR/RW in AR(p) Error Correction

- Unit roots and error correction form in AR(p) models

- Setup:

$$Y_t = \mu + \sum_{i=1}^p \phi_i Y_{t-i} + \epsilon_t$$

with  $\sum_{i=1}^p \phi_i = 1$

- Error-correction form of AR(p):

$$\Delta Y_t = \mu + \delta Y_{t-1} + \underbrace{\gamma_1 \Delta Y_{t-1} + \dots + \gamma_{p-1} \Delta Y_{t-p+1}}_{\sum_{i=1}^{p-1} \gamma_i \Delta Y_{t-i}} + \epsilon_t$$

where

$$\begin{cases} 
 \delta = (\sum_{i=1}^p \phi_i) - 1 \quad (= 0 \text{ if unit root}) \\ 
 \gamma_1 = -\sum_{i=2}^p \phi_i \\ 
 \gamma_2 = -\sum_{i=3}^p \phi_i \\ 
 \dots \\ 
 \gamma_{p-1} = -\phi_p 
 \end{cases}$$

### Problems of Trends

- What problems are caused by trends?

- **Biasedness:** OLS of AR coefficient will be strongly biased towards 0 ↵ poor forecasts
- **Not well-behaving:** t-stat does not follow standard Normal distribution, even in large samples ↵ standard inference tools cannot be used
- If both Y and X have stochastic trends, we may find a "spurious relationship"

## Dickey-Fuller (DF) Test for Stochastic Trends

- How do you detect stochastic trends?
  - Eyeball: plot the data and see whether it is highly persistent
  - Check autocorrelations: slow decay in autocorrelations?

### Dickey-Fuller Test

- Fit the error-correction form
  - If the original model is AR(1), then:

$$\hat{\delta} = \frac{\sum_{t=1}^T (\Delta Y_t - \bar{\Delta Y})(Y_{t-1} - \bar{Y})}{\sum_{t=1}^T (Y_{t-1} - \bar{Y})^2}$$

- Test the null (RW/UR):

$$\begin{cases} H_0 : \delta = 0 & \text{Has RW/UR (Non-Stationary)} \\ H_1 : \delta < 0 & \text{No RW/UR (Stationary)} \end{cases}$$

- This is a **one-side test!**  $\{Y_t\}$  is stationary only when all roots are outside the unit circle)
- Compare the  $t_{\hat{\delta}} = \frac{\hat{\delta}}{se(\hat{\delta})}$  with the **critical values of the Dickey-Fuller distribution** (because regressor  $Y_{t-1}$  is non-stationary under  $H_0$ ):

**TABLE 14.5 Large-Sample Critical Values of the Augmented Dickey-Fuller Statistic**

Deterministic Regressors	10%	5%	1%
Intercept only	-2.57	-2.86	-3.43
Intercept and time trend	-3.12	-3.41	-3.96

- When should you include a time trend in the DF test?:
- Error-correction forms of AR(p) with/out time trend

$$\begin{cases} \text{Intercept only : } & \Delta Y_t = \mu + \delta Y_{t-1} + \sum_{i=1}^{p-1} \gamma_i \Delta Y_{t-i} + \epsilon_t \\ \text{Intercept & Time Trend : } & \Delta Y_t = \mu + \alpha t + \delta Y_{t-1} + \sum_{i=1}^{p-1} \{\gamma_i \Delta Y_{t-i}\} + \epsilon_t \end{cases}$$

- **Intercept only:**  $H_0$ : UR/RW;  $H_1$ :  $\{Y_t\}$  is stationary around a constant
- **Intercept & Time Trend:**  $H_0$ : UR/RW;  $H_1$ :  $\{Y_t\}$  is stationary around a linear time trend (overall still non-stationary), i.e. the TS has long-term growth
- How to address/mitigate problems raised by unit roots:

- A UR/RW in  $\{Y_t\} \implies$  First-differencing (i.e.  $\Delta Y_t = Y_t - Y_{t-1}$ ) will be stationary

## 5.2 [Trends II] - A

- Random walk with drift
  - Random walk with drift are explosive
  - RW with drift:

$$Y_t = \mu + Y_{t-1} + \epsilon_t$$

- Solving backwards:

$$Y_t = Y_0 + \mu t + \sum_{i=1}^t \epsilon_i \implies \begin{cases} \mathbb{E}[Y_t|Y_0] &= Y_0 + \mu t \rightarrow \infty \text{ as } t \rightarrow \infty \\ \text{Var}(Y_t|Y_0) &= t\sigma_\epsilon^2 \rightarrow \infty \text{ as } t \rightarrow \infty \end{cases}$$

- Thus, the MA weights are not square summable  $\implies MA(\infty)$  representation will not be available  $\implies$  non-stationary
- Testing Random Walk hypothesis
  - Analysis of t statistic
  - Dickey—Fuller (DF) distribution of t-stat

## Integrated Stochastic Trends (Multiple Unit Roots)

- General Stochastic Trends
- Integrated processes - multiple unit roots:
  - **I(1)** process (RW/UR): AR process with 1 unit root in its characteristic polynomial;  $\Delta Y_t$  is stationary:

$$Y_t = \mu + Y_{t-1} + \epsilon_t \iff \Delta Y_t = \mu + \epsilon_t$$

- **I(2)** process: AR process with 2 unit roots in its characteristic polynomial;  $\Delta Y_t$  still contains a RW/UR;  $\Delta^2 Y_t$  is stationary:

$$\begin{aligned} Y_t &= \mu + 2Y_{t-1} - Y_{t-2} + \epsilon_t \iff \Delta Y_t = \mu + \Delta Y_{t-1} + \epsilon_t \\ &\iff \Delta^2 Y_t = \mu + \epsilon_t \end{aligned}$$

- **I(d)** process: AR process with  $d$  unit roots in its characteristic polynomial; we have to difference  $d$  times to remove the unit roots:
  - $\Delta^d Y_t$  is stationary
  - $\Delta^i Y_t$  is non-stationary  $\forall i < d$
- Testing for I(0) versus I(1) versus I(2)
  - Sequential testing procedure:
    1. Test (I(1)) whether  $\delta = 0$  (UR) against  $\delta < 0$  (stationary) in FD:

$$\Delta Y_t = \mu + \delta Y_{t-1} + \sum_{i=1}^{p-1} \gamma_i \Delta Y_{t-i} + \epsilon_t$$

- 2. If accept, test ( $I(2)$ ) whether  $\delta = 0$  (UR) against  $\delta < 0$  (stationary) in SD:

$$\Delta^2 Y_t = \mu + \delta \Delta Y_{t-1} + \sum_{i=1}^{p-2} \gamma_i \Delta^2 Y_{t-i} + \epsilon_t$$

- 3. ...

- Summary: detecting and addressing unit roots/stochastic trends
- 

## Week 6: Asset Return Predictability

### 6.1 [Asset Return Predictability I] - A

#### Basics

- Asset Prices and Returns
  - Continuously compounded return or log-return

$$r_t = \log(1 + R_t) = p_t - p_{t-1}$$

where  $p_t = \log(P_t)$

- and  $r_t \approx R_t$  for small price changes
- Log-compounding formula

$$r_t(h) = \log(1 + R_t(h)) = \sum_{i=0}^{h-1} r_{t-i}$$

- Excess return:

$$\bar{R}_t = R_t - R_{0,t}, \bar{r}_t = r_t - r_{0,t}$$

- where  $R_{0,t}, r_{0,t}$  are risk-free returns
- Risk premium = Expected excess return:

$$\mathbb{E}[\bar{R}_{t+1} | \mathcal{I}_t] = \mathbb{E}[R_{t+1} | \mathcal{I}_t] - R_{f,t+1}$$

#### Asset Return Predictability & Market Efficiency Hypothesis

- Asset Return Predictability
- Martingale Hypothesis: current price contains all relevant information (any assets have *0 expected payoff* / no RP and risk-free rate is 0)

$$\mathbb{E}[r_{t+1} | \mathcal{I}_t] = 0 \iff \mathbb{E}[p_{t+1} | \mathcal{I}_t] = p_t$$

- Stochastic processes with this property are known as **Martingales**

- Random Walk Hypothesis: *allows risk premium* captured by  $\mu$ :

$$H_{RW} : \mathbb{E}[r_{t+1} | \mathcal{I}_t] = \mu \iff \mathbb{E}[p_{t+1} | \mathcal{I}_t] = p_t + \mu$$

or

$$p_t = \mu + p_{t-1} + \epsilon_t, \mathbb{E}[\epsilon_t | \mathcal{I}_{t-1}] = 0$$

- $\mathcal{I}_t$  is the information set used to forecast future price
- $\epsilon_t$  is the forecast error, capturing surprise movements and new information
- $\mu$  is expected log-return (RP)
- Riskier assets have higher variance ( $\sigma_\epsilon^2$ ) and higher risk premium  $\mu$
- Efficient Market Hypothesis:  $H_{RW}$  above with different information sets:
  - Weak-form efficiency:  $\mathcal{I}_t$  includes only prices
  - Semi-strong efficiency:  $\mathcal{I}_t$  includes all publicly available information
  - Strong efficiency:  $\mathcal{I}_t$  includes all information (even private information)
  - EMH *Allows a predictable component (risk premium) in excess return if investors are risk-averse*
    - EMH only requires that a trading strategy cannot earn abnormal returns relative to a benchmark model for the market

## Trading Rules

- "Technical" Trading Rules
  - Moving average (MA) rules
  - Do They "Work"?
  - Data Snooping - a multiple testing problem
    - e.g. testing at 5% level, then 5% of useless trading rules will be considered significantly effective
    - Reality-check test quantifies the effect of data snooping by evaluating the performance of the best trading rules in the context of the full "universe" of rules
      - Sullivan et al, 1999 found the best-performing rules still out-perform the benchmark

## 6 2 [Asset Return Predictability II] - A

- Some Stylized Facts of Returns - the S&P 100
- Random Walk Hypothesis
- Testing the RW Hypothesis - choice of information set and alternative

## Testing Weak Form $H_{RW}$ against $AR(p)$

- Random Walk Hypothesis

- "Weak test" of the RW Hypothesis with parametric alternative (AR(p))

- Test RW/UR:

$$H_{RW} : \mathbb{E}[p_t | p_{t-1}, p_{t-2}, \dots] = \mu + p_{t-1}$$

- Assume  $p_t$  follows an AR(p) process:

$$p_t = \mu + \sum_{i=1}^p \phi_i p_{t-i} + \epsilon_t$$

- Use the Error Correction Form of AR(p)

$$r_t = \mu + \delta p_{t-1} + \sum_{i=1}^{p-1} \gamma_i r_{t-i} + \epsilon_t$$

1. Use the DF test to test Unit Root (with drift) in  $\{p_t\}$ :  $H_0 : \delta = 0; H_1 : \delta < 0$
2. (If accept) Set  $\delta = 0$  and test  $H_{RW} : \gamma_1 = \gamma_2 = \dots = \gamma_{p-1} = 0$ 
  - If both are accepted, then we accept  $H_{RW}$

- Is AR(p) the right alternative?

- Model misspecification (non-linear predictability)
- Choose p

### Model-Free Test of Weak Form $H_{RW}$

#### Ljung-Box Q Test

- The Q (or Ljung-Box) Test based on *autocorrelations*:

$$H_{RW}/H_0 : \rho_r(h) = \text{corr}(r_t, r_{t+h}) = 0 \quad \forall h \in \mathbb{N}$$

- Q-statistic

$$Q_m = T(T+2) \sum_{h=1}^m \frac{1}{T-h} \hat{\rho}_r^2(h) \sim \chi_m^2 \text{ under } H_{0/RW} \text{ (stationary and mixing returns)}$$

- Size control (choosing  $m$ ): we need to choose  $m$  to achieve a balance between determining power (large  $m \rightsquigarrow$  more likely to find autocorrelation) and size of the test (small  $m \rightsquigarrow$  smaller estimation uncertainty)

#### Variance-Ratio (VR) Test

- The Variance-Ratio (VR) Test by Lo and MacKinlay (1988) based on *autocovariances*:
- Denote the variance of h-step return as

$$V(h) = \text{Var}(r_t(h)) = \text{Var}(r_t + r_{t-1} + \dots + r_{t-h+1})$$

- $H_{RW}$  implies that all autocovariance are zero, so:

$$V(h) = \sum_{i=0}^{h-1} Var(r_{t-i}) = h \times V(1) \iff VR(h) = \frac{V(h)}{hV(1)} = 1$$

- Sample Analogue:

$$\widehat{VR}(h) = \frac{\hat{V}(h)}{h\hat{V}(1)} \text{ where } \hat{V}(h) = \underbrace{\frac{T}{(T-h)(T-h+1)}}_{\approx \frac{1}{T}} \sum_{t=h+1}^T \underbrace{\left[ r_t(h) - \bar{r}(h) \right]^2}_{v_t(h)}$$

under  $H_{RW}$ :

$$t_{VR}(h) = \frac{\widehat{VR}(h) - 1}{\hat{\sigma}_{VR}(h)} \xrightarrow{d} \mathcal{N}(0, 1)$$

where  $\hat{\sigma}_{VR}^2(h) = \frac{1}{T} \frac{\hat{\Omega}_{VR}(h)}{h^2 \hat{V}^2(1)}$  and  $\hat{\Omega}_{VR}(h)$  is the HAC estimator of the long-run variance of  $VR(h)$

- Size control (choosing  $h$ ):

- we need to choose  $h$  to achieve a balance between determining power (large  $h \rightsquigarrow$  more likely to reject correctly) and size of the test (small  $h \rightsquigarrow$  smaller estimation uncertainty)
- In principle we should check all possible  $h$  by constructing a F-test
- In finite samples, the distribution of VR-stat can be far from normal

## Testing Semi-Strong $H_{RW}$ by Regressions

### Test based on DL Model

- Regression-based Tests: adding more variables to  $\mathcal{I}_t$
- simple distributed lag (DL) model:

$$r_{t+h}(h) = p_{t+h} - p_t = \beta_{h,0} + \beta_{h,1} X_t + \epsilon_{h,t}$$

where  $h \geq 1$  is the forecast horizon,  $X_t$  are potential predictors, and  $\epsilon_{h,t}$  is forecast error

- When  $h$  is large, this is known as a long-horizon regression
- In-Sample Evidence of Predictability
- Out-of-sample Evidence of Predictability

### Issues

- Some Critical Issues
  - Persistent Regressors
    - If  $\{X_t\}$  is highly persistent, then OLS estimator will be a mixture of Augmented Dickey-Fuller (ADF) distribution and  $\mathcal{N}(0, 1)$ :

$$t = \frac{\beta_{h,1}}{\hat{\sigma}_{\beta_{h,1}}} \sim [\lambda ADF + (1 - \lambda)\mathcal{N}(0, 1)]$$

- Long-Horizon Regressions

1. Size control: As  $h$  becomes larger, there will be fewer observations left for estimation (effective sample size become smaller)
  2.  $r_{t+h}(h)$  (or  $\epsilon_t(h)$ ) becomes more persistent as  $h$  gets larger, because it is a rolling summation of  $r_t$   $\rightsquigarrow$  OLS will have a Dickey-Fuller type distribution and we need a HAC-type estimator for the SE; the finite-sample distribution will be far from normal
  3. Lack of strict exogeneity of predictors -- a shock to returns will affect future values of predictors
- DL with a single predictor is too weak. We should use an ADL with more predictors
- 

## Week 7: Structural Breaks (Another Type of Non-Stationarity)

### 7.1 [Structural Breaks I] - A

#### Structural Breaks

- Structural break: another type of non-stationarity
  - our model

$$\begin{cases} Y_t = \mu + \phi Y_{t-1} + \psi X_{t-1} + \epsilon_t & , t = 1, \dots, \tau \\ Y_t = \mu' + \phi' Y_{t-1} + \psi' X_{t-1} + \epsilon_t & , t = \tau, \dots, T \end{cases}$$

- bias-variance trade-off

#### Testing for Structural Break when Break Date is Known

- Test for structural break when break date is known
- New variables
  - Dummy:  $D_t(\tau) = \mathbb{1}\{t > \tau\} = \begin{cases} 0, & t \leq \tau \\ 1, & t > \tau \end{cases}$
  - Deltas:  $\delta_0 = \mu' - \mu, \delta_1 = \phi' - \phi, \delta_2 = \psi' - \psi$
- Rewrite our model

$$Y_t = \mu + \phi Y_{t-1} + \psi X_{t-1} + \delta_0 D_t(\tau) + \delta_1 Y_{t-1} D_t(\tau) + \delta_2 X_{t-1} D_t(\tau) + \epsilon_t$$

- Chow test of structural break
  - $H_0 : \delta_0 = \delta_1 = \delta_2 = 0; H_1 : \text{at least one } \delta \neq 0$
  - Use Chow test: (heteroskedastic-robust)  $F\text{-stat} \sim F_{q,\infty}$  *when data is stationary in each of the segments* ( $|\phi| < 1, |\phi'| < 1, \{X_t\}$  is stationary + mixing) and  $q$  is the number of restriction

## Testing for Structural Breaks when Break Date is Unknown

### Quandt Likelihood (QLR) Test for One Break

- Quandt Likelihood Ratio (QLR) Statistic: test for

$$\begin{cases} H_0 & : \text{No break in } \{\tau_0, \dots, \tau_1\} \\ H_1 & : \text{One break in } \{\tau_0, \dots, \tau_1\} \end{cases}$$

- convention is to use the mid 70% data
- QLR statistic:

$$QLR = \max\{F(\tau_0), F(\tau_0 + 1), \dots, F(\tau_1)\}$$

where  $F(\cdot)$  is the Chow F-stat testing  $H_0$  : no break at  $\tau$

- Critical values of QLR statistic

- Critical values of QLR-stat are *larger* than that of individual F-stats ( $F_{q,\infty}$ ) because it depends on multiple F-distributions:

$$QLR \sim \max_{s_0 \leq s \leq s_1} \left\{ \frac{1}{q} \sum_{i=1}^q \frac{B_i^2(s)}{s(1-s)} \right\}$$

if  $CLR > cv_\alpha$ , we reject  $H_0$  and our estimated break date is

$$\hat{\tau} = \arg \max_{\tau_0 \leq \tau \leq \tau_1} F(\tau)$$

- QLR Stat for 15% trimming (keep the middle 70%)

**TABLE 14.6 Critical Values of the QLR Statistic with 15% Trimming**

Number of Restrictions (q)	10%	5%	1%
1	7.12	8.68	12.16
2	5.00	5.86	7.78
3	4.09	4.71	6.02
4	3.59	4.09	5.12
5	3.26	3.66	4.53
6	3.02	3.37	4.12
7	2.84	3.15	3.82
8	2.69	2.98	3.57
9	2.58	2.84	3.38
10	2.48	2.71	3.23

- Limits of QLR:
  - unable to detect breaks in the last 15% of the sample
  - *allows for only one break*; hard to handle multiple breaks (Discussed in part II below)

## Testing Structural Breaks at the End of the TS

- pseudo out-of-sample (poos)
  - Want to check whether there is a break from  $\tau_1$  to  $T$ :
  - 1. Found a break at  $\tau$  using QLR test
  - 2. Discard all data from  $t = 1 \dots \tau$
  - 3. Choose some "burn-in" samples from  $P$  to  $\tau_1$  where  $\tau < P < \tau_1$
  - 4. For  $t = P, \dots, T$ :
    - 1. Estimate the model using data from  $\tau$  to  $t$
    - 2. Calculate the in-sample  $\widehat{SER}$
    - 3. Compute  $\hat{Y}_{t+1|t}$  and  $\widehat{MSFE} = \frac{1}{T-P} \sum_{t=P}^T (Y_{t+1} - \hat{Y}_{t+1|t})^2$
  - 5. Compare the  $\widehat{MSFE}$  with the in-sample  $\widehat{SER}$  from the "burn in" regression: if they are similar, then it is likely that there's no break
- Rolling-window estimates
  - Pros: 1. can detect multiple breaks, 2. this is more flexible (no parametric assumption on coefficients); Cons: not a formal statistical test
  - 1. Choose a fixed time window  $H \geq 1$
  - 2. Estimate the model while rolling the window with  $H$  samples included:
    - For  $t = H + 1, \dots, T$ : estimate the model using data  $Y_t, \dots, Y_{t+H}; X_t, \dots, X_{t+H}$
    - Store the estimates  $\hat{\mu}_t, \hat{\phi}_t, \hat{\psi}_t$
  - 3. Plot the time series of estimates  $\hat{\mu}_t, \hat{\phi}_t, \hat{\psi}_t$  together with their CIs and compare them with overall estimates

## 7.2 [Structural Breaks II] - A

### Testing for Structural Breaks when Break Date is Unknown: QLR Test Part II

- Chow and QLR test of structural break

#### Quandt Likelihood (QLR) Test for Multiple Breaks

- Multiple breaks
- Method 1: Sequential One-Break QLR Test:
  - Test for single break using QLR Test for One Break and obtain  $\hat{\tau}$
  - Split the sample at  $\hat{\tau}$  and conduct QLR Test for One Break for each subsample:

$$QLR_1 = \max_{\tau_0 \leq \tau \leq \hat{\tau}} F(\tau), QLR_2 = \max_{\hat{\tau}+1 \leq \tau \leq \tau_1} F(\tau)$$

- If  $QLR_k > cv_\alpha$ , we conclude that SB also occurs in the  $k^{th}$  subsample
- Repeat
  - Pros: computationally easy; Cons: lose size control - the common problem with multiple hypothesis testing

- Method 2: Single Joint QLR Test

1. If we suppose there are  $k$  breaks, then we use  $k$  dummies to reconstruct the equation.  
e.g. when we suppose SB for intercepts:

$$Y_t = \mu + \delta_0 D_t(\tau_1) + \delta_1 D_t(\tau_2) + \cdots + \delta_k D_t(\tau_k)$$

2. Compute QLR:

$$QLR = \max_{\tau_1, \dots, \tau_k} \{F(\tau_1, \dots, \tau_k)\}$$

- Pros: less statistical problem; Cons: computationally intensive due to the large number of possible combinations of  $\tau$

## *Rolling-window estimates (brief)*

---

## Week 8: Estimation of Dynamic Causal Effects

### 8.1 [Dynamic Causal Effects I] - A

#### Dynamic Causal Effects

- Dynamic causal effects is the effect of a change in  $X$  on  $Y$  over time
- Ideal setting: Randomized experiment

#### Distributed Lag Model

##### DL Model (A Dynamic Model)

- DL estimators are OLS estimator in distributed lag model:

$$Y_t = \mu + \psi_0 X_t + \cdots + \psi_q X_{t-q} + \epsilon_t$$

where:

-  $\psi_0$  is the Immediate Impact Effect of change in  $X_t$ : effect of change in  $X_t$  on  $Y_t$ , holding  $(X_{t-1}, \dots, X_{t-q})$  constant (under weak exogeneity)

-  $\psi_1$  is the 1-period Dynamic Multiplier: effect of change in  $X_{t-1}$  on  $Y_t$ , holding  $(X_t, X_{t-2}, \dots, X_{t-q})$  constant (under weak exogeneity)

- ...

- h-period Dynamic Multiplier:

$$\psi_h = \frac{\partial \mathbb{E}[Y_t | X_t, X_{t-1}, \dots]}{\partial X_{t-h}} = \frac{\partial \mathbb{E}[Y_{t+h} | X_{t+h}, X_{t+h-1}, \dots]}{\partial X_t}$$

- k-period Cumulative Dynamic Multipliers measure the accumulated effects =  $\sum_{i=1}^k \psi_i$
- (PPT II) Estimators of cumulative multipliers and their se's: rewrite the model so that coefficients are the cumulative multipliers

- Example, a DL(2) Model:

$$\begin{aligned} Y_t &= \mu + \psi_0 X_t + \psi_1 X_{t-1} + \psi_2 X_{t-2} + \epsilon_t \\ &= \mu + \psi_0 (X_t - X_{t-1}) + \underbrace{(\psi_0 + \psi_1)(X_{t-1} - X_{t-2})}_{\delta_1} + \underbrace{(\psi_0 + \psi_1 + \psi_2)X_{t-2}}_{\delta_2} + \epsilon_t \end{aligned}$$

### *DL Assumptions*

- Distributed Lag Model Assumptions

1.  $X$  is **exogenous**:  $\mathbb{E}[\epsilon_t | X_t, X_{t-1}, X_{t-2} \dots] = 0$ 
  - This allows autocorrelations in  $\epsilon$ , but if so, HAC SE will be needed (see below)
2.  $\{Y_t, X_t\}$  is stationary and geometrically mixing
3. Eighth moments exist:  $\mathbb{E}[Y^8] < \infty$  and  $\mathbb{E}[X^8] < \infty$
4. No perfect multicollinearity

### *Properties of OLS under DL Assumptions*

- W2: Properties of OLS under DL assumptions
- OLS estimators of  $\mu, \psi_0, \dots, \psi_q$  are
  - *consistent (but biased)*
  - have *asymptotic Normal distribution* (so corresponding t/F statistics can be used), but variance needs to be adjusted for autocorrelations and heteroskedasticity ↓
- DL OLS with autocorrelated errors
  - *Math derivation* of Distribution with Weak Exogeneity:
  - Model:

$$Y_t = \mu + \psi_0 X_t + \epsilon_t, \begin{cases} \mathbb{E}[\epsilon_t | X_t, X_{t-1}, X_{t-2} \dots] = 0 \\ \mathbb{E}[\epsilon_t | \epsilon_{t-1}, \epsilon_{t-2} \dots] \neq 0 \end{cases}$$

- Estimator:

$$\hat{\phi}_0 = \frac{\sum_{t=1}^T (X_t - \bar{X}) Y_t}{\sum_{t=1}^T (X_t - \bar{X})^2} = \phi_0 + \frac{\sum_{t=1}^T (X_t - \bar{X}) \epsilon_t}{(X_t - \bar{X})^2}$$

- Use LLN for stationary and mixing TS:

$$\frac{1}{T} \sum_{t=1}^T (X_t - \bar{X})^2 \xrightarrow{p} Var(X_t) = \sigma_X^2$$

- Denote  $v_t = (X_t - \mu_X) \epsilon_t$

- Use CLT for stationary and mixing TS and  $\mathbb{E}[v_t] = 0$ :

$$\frac{1}{T} \sum_{t=1}^T (X_t - \bar{X})\epsilon_t \approx \frac{1}{T} \sum_{t=1}^T v_t \xrightarrow{d} \mathcal{N} \left( 0, \frac{\bar{\sigma}_v^2}{T} \right)$$

where  $\bar{\sigma}_v^2 = \sigma_v^2 \left\{ 1 + 2 \sum_{k=1}^{T-1} \frac{T-k}{T} \rho_{(v_t, v_{t-k})} \right\}$  (see below for estimator)

- **Result:** Distribution with Standard Exogeneity: in the simple model  $Y_t = \mu + \psi_0 X_t + \epsilon_t$  with  $\mathbb{E}[\epsilon_t | X_t, X_{t-1}, X_{t-2}, \dots] = 0$ :

$$\hat{\psi}_0 = \psi_0 + \frac{\sum_{t=1}^T (X_t - \bar{X})\epsilon_t}{\sum_{t=1}^T (X_t - \bar{X})^2} \sim^a \mathcal{N} \left( \psi_0, \frac{1}{T} \frac{\bar{\sigma}_v^2}{\sigma_X^4} \right)$$

where  $v_t = (X_t - \mu_X)\epsilon_t$  and  $\bar{\sigma}_v^2$  can be estimated by the [Newey-West Estimator](#):

$$\hat{\sigma}_v = \hat{\sigma}_v^2 + 2 \sum_{k=1}^m \frac{T-k}{T} \widehat{Cov}(v_t, v_{t-k})$$

and [autocovariance functions](#) can be estimated by:

$$\widehat{Cov}(v_t, v_{t-k}) = \frac{1}{T} \sum_{t=k}^T \hat{v}_t \hat{v}_{t-k}$$

- With [Strong Exogeneity Restriction \(Dynamically Complete\)](#), we can use standard SE: in the simple model  $Y_t = \mu + \psi_0 X_t + \epsilon_t$  with  $\mathbb{E}[\epsilon_t | X_t, \epsilon_{t-1}, X_{t-1}, \epsilon_{t-2}, X_{t-2}, \dots] = 0$ :

$$\hat{\psi}_0 = \psi_0 + \frac{\sum_{t=1}^T (X_t - \bar{X})\epsilon_t}{\sum_{t=1}^T (X_t - \bar{X})^2} \sim^a \mathcal{N} \left( \psi_0, \frac{1}{T} \frac{\mathbb{E}[(X_t - \mu_X)^2 \epsilon_t^2]}{\sigma_X^4} \right)$$

## Different Kinds of Exogeneity (PPT II)

- Exogeneity in time series regression
- Standard Exogeneity restriction: error is unpredictable with past and present values of the predictor:

$$\mathbb{E}[\epsilon_t | X_t, X_{t-1}, X_{t-2} \dots] = 0$$

- Interpretation of dynamic multiplier:

$$\psi_h = \frac{\partial \mathbb{E}[Y_t | X_t, X_{t-1}, \dots]}{\partial X_{t-h}} = \frac{\partial \mathbb{E}[Y_{t+h} | X_{t+h}, X_{t+h-1}, \dots]}{\partial X_t}$$

- Weaker Exogeneity restriction:

$$\mathbb{E}[\epsilon_t | X_t, X_{t-1}, \dots, X_{t-q}] = 0$$

- With this, we can only interpret  $\psi_h$  as a *finite-horizon* effect:

$$\psi_h = \frac{\partial \mathbb{E}[Y_t | X_t, X_{t-1}, \dots, X_{t-q}]}{\partial X_{t-h}}$$

- Stronger Exogeneity restriction:

$$\mathbb{E}[\epsilon_t | X_t, \epsilon_{t-1}, X_{t-1}, \epsilon_{t-2}, X_{t-2}, \dots] = 0$$

- With this, the model is *dynamically complete*: first q lags of X explains all the dynamics of Y (and HAC SE is not needed)
- Strict Exogeneity restriction: error is unpredictable with past, present, and future values of the predictor:

$$\mathbb{E}[\epsilon_t | \dots, X_{t+2}, X_{t+1}, X_t, X_{t-1}, X_{t-2}, \dots] = 0$$

- Estimation of Dynamic Causal Effects with Strictly Exogenous Regressors
  - With strict exogeneity, there are more efficient ways to estimate dynamic causal effects than DL estimators
    - GLS estimation
    - ADL estimation
- When is Exogeneity Plausible?
  - Think about whether  $Cov(\epsilon_{t+h}, X_t) = 0 \forall h \geq 0$
  - Reasons that exogeneity fails:
    - Simultaneity:  $\begin{cases} Y_t = \mu + \psi_0 X_t + \epsilon_t \\ X_t = \mu_X + \psi_X Y_t + \epsilon_{X,t} \end{cases}$
    - Feedback:  $Y_{t-q} \rightarrow X_t$

## Standard Errors in ADL Models

- In ADL/AR models, with enough lags of  $Y_t$ , the error term cannot be predicted using past  $Y_t$ 's, which is equivalent to that  $\epsilon_t$  cannot be predicted by past  $\epsilon_t$ 's (i.e.  $\epsilon_t$  is not serially correlated)
- But if there're not enough lags,  $\epsilon_t$  will still be serially correlated, so always check autocorrelations for residuals

## Estimators of Cumulative Multipliers and Their SE

- Rewrite the regression with coefficients equal to cumulative multipliers
- Example: DL(2):

$$\begin{aligned} Y_t &= \mu + \psi_0 X_t + \psi_2 X_{t-1} + \psi_2 X_{t-2} + \epsilon_t \\ &= \mu + \psi_0(X_t - X_{t-1}) + \underbrace{(\psi_0 + \psi_1)(X_{t-1} - X_{t-2})}_{\delta_1} + \underbrace{(\psi_0 + \psi_1 + \psi_2)X_{t-2}}_{\delta_2} + \epsilon_t \end{aligned}$$

## Week 9: CAPM and APT

### 9.1 [CAPM] - A

#### Setup, Notations, and Assumptions

- Mean-Variance Portfolio Selection: use variance to measure risk
- The Security Market
  - Risk-free asset:  $P_{t+1} = (1 + R_{0,t+1})P_{0,t}$
  - Risky asset:  $P_{t+1} = (1 + R_{t+1})P_t$
  - Set  $P_{0,t} = 1$
- Investor's Portfolio
  - Units of risk-free / risky asset:  $w_{0,t}, w_t$
  - Acquisition cost  $V_t = w_{0,t} + w_t P_t$
  - Value of portfolio 1 period later  $V_{t+1} = w_{0,t}(1 + R_{0,t+1}) + w_t P_{t+1}$
  - Conditional expected value  $\mathbb{E}_t[V_{t+1}] = w_{0,t}(1 + R_{0,t+1}) + w_t \mathbb{E}_t[P_{t+1}]$
  - Conditional variance  $\sigma_t^2(V_{t+1}) = w_t^2 \sigma_t^2(P_{t+1})$
- Investor's Preferences - assumption
  - Expected utility is quadratic:

$$\begin{aligned} U_t &= E_t[V_{t+1}] - \frac{\lambda}{2} \sigma_t^2(V_{t+1}) \\ &= w_{0,t}(1 + R_{0,t+1}) + w_t E_t[P_{t+1}] - \frac{\lambda}{2} w_t^2 \sigma_t^2(P_{t+1}) \end{aligned}$$

- Investor's Decision Problem

- Initial wealth at t:  $W_t$
  - Maximisation:

$$\max_{w_{0,t}, w_t} \left\{ w_{0,t}(1 + R_{0,t+1}) + w_t E_t[P_{t+1}] - \frac{\lambda}{2} w_t^2 \sigma_t^2(P_{t+1}) \right\} \text{ s.t. } w_{0,t} + w_t P_t = W_t$$

- Solution:  $W_t^* = \frac{1}{AP_t} \frac{E_t[\bar{R}_{t+1}]}{\sigma_t^2(\bar{R}_{t+1})}$ ,  $w_{0,t}^* = W_t - w_t^* P_t$  where  $\bar{R}_{t+1} = R_{t+1} - R_{0,t+1}$  is the excess return of the asset

#### CAPM

- The Capital Asset Pricing Model (CAPM)
  - Investor  $i \in \{1, \dots, M\}$  has risk aversion  $A_i$ , initial wealth  $W_{i,t}$ , and chooses his/her portfolio according to the above mean-var analysis
  - The Market Portfolio:

$$\text{Total Unit Held} = \begin{cases} \bar{w}_t = \sum_{i=1}^M w_{i,t}^* = \frac{1}{\bar{A}P_t} \frac{E_t[\bar{R}_{t+1}]}{\sigma_t^2(\bar{R}_{t+1})} \\ \bar{w}_{0,t} = \sum_{i=1}^M W_{i,t} - \bar{w}_t P_t = \bar{W}_t - \bar{w}_t P_t \end{cases}$$

$$\text{where } \bar{A} = \left[ \sum_{i=1}^M \frac{1}{A_i} \right]^{-1}$$

- General CAPM:

$$\mathbb{E}_t[\bar{R}_{i,t+1}] = \beta_{i,t} \mathbb{E}_t[\bar{R}_{m,t+1}]$$

where  $\begin{cases} \beta_{i,t} = \frac{\text{Cov}_t[\bar{R}_{i,t+1}, \bar{R}_{m,t+1}]}{\text{Var}_t[\bar{R}_{m,t+1}]}, \forall i \in \{1, \dots, n\} \\ \bar{R}_{m,t+1} = \sum_{i=0}^N \pi_i \bar{R}_{i,t+1} \text{ with } \pi_i = \text{share of total wealth in asset } i \end{cases}$

- "Bar" means excess return here
- $\bar{R}_{m,t+1}$  and  $\beta_{i,t}$ 's fully explain individual assets' expected excess returns
- Expected returns are linear to betas
- Market RP / expected excess return  $\mathbb{E}_t[\bar{R}_{m,t+1}] > 0$ : in equilibrium, asset prices must be such that all assets are held, and risk-averse investors will only hold risky assets if they have positive excess returns

## Estimation and Testing of CAPM

- Collecting data

### CAPM with Constant Betas

- CAPM with Constant Betas

- Rewrite the CAPM as a regression model:

$$\bar{R}_{i,t} = \beta_{i,t} \bar{R}_{m,t} + \epsilon_{i,t}, \quad \mathbb{E}[\epsilon_{i,t} | \mathcal{I}_{t-1}] = 0$$

- Assume *time-unvarying*  $\beta_{i,t} = \beta_i$ , this can be estimated using CAPM Time Series Regression:

$$\bar{R}_{i,t} = \alpha_i + \beta_i \bar{R}_{m,t} + \epsilon_{i,t}, \quad i = 1, \dots, n$$

which is a system of distributed lag models

- CAPM suggests:  $H_0 : \alpha_i = 0 \forall i \in \{1, \dots, n\}$ , which is a cross-equation restriction. We need to account for covariance across assets, so we stack all CAPM equations into a vector DL model:

$$\bar{R}_t = \alpha + \beta' \bar{R}_{m,t} + \epsilon_t$$

where  $\bar{R}_t = (\bar{R}_{1,t}, \dots, \bar{R}_{n,t})'$ ,  $\alpha = (\alpha_1, \dots, \alpha_n)'$ ,  $\beta = (\beta_1, \dots, \beta_n)'$ ,  $\epsilon_t = (\epsilon_{1,t}, \dots, \epsilon_{n,t})'$

- This can be estimated by OLS:

$$\hat{\beta} = \frac{\sum_{t=1}^T (\bar{R}_{m,t} - \hat{\mu}_{\bar{R}_m})(\bar{R}_t - \hat{\mu}_{\bar{R}})}{\sum_{t=1}^T (\bar{R}_{m,t} - \hat{\mu}_{\bar{R}_m})^2}, \hat{\alpha} = \hat{\mu}_{\bar{R}} - \hat{\beta} \hat{\mu}_{\bar{R}_m}$$

where  $\hat{\mu}_{\hat{R}_m}, \hat{\mu}_{\hat{R}}$  are sample means, and the var-cov matrix is:

$$\hat{\Omega}_\epsilon = \frac{1}{T} \sum_{t=1}^T \hat{\epsilon}_t \hat{\epsilon}'_t, \quad \hat{\epsilon} = \hat{R}_t - \hat{\alpha} - \hat{\beta} \bar{R}_{m,t}$$

- We can test CAPM:  $H_0 : \alpha_i = 0, i = 1, \dots, n$  using an F-stat with HAC SE

## *Reasons for Rejection*

- Possible Explanations for Rejection
  - Unrealistic assumptions (model is too simple)
    - Mean-var preference does not hold
    - Asymmetric information
    - Behaviour
    - Market incompleteness
  - Time-varying betas
  - Data collection: Roll's critique (stock index is not complete representation of the market portfolio), data snooping, etc.

## *9.2 CCAPM and APT - A*

### *Conditional CAPM (Varying Beta)*

- Estimation and Testing of Conditional CAPM

#### *Setup*

- Allows for  $\beta_{i,t}$  to vary over time:

$$\mathbb{E}_t[\bar{R}_{i,t+1}] = \beta_{i,t+1} \mathbb{E}_t[\bar{R}_{m,t+1}]$$

or

$$\bar{R}_{i,t} = \beta_{i,t} \bar{R}_{m,t} + \epsilon_{i,t}, \quad \mathbb{E}[\epsilon_{i,t} | \mathcal{I}_{t-1}] = 0$$

#### *Estimation Method 1: Instrumenting the Betas (Not IV)*

- **Instrumenting the Betas:** specify the underlying structures of  $\beta$
- Suppose variations in  $\beta_{i,t}$  are caused by a single underlying factor  $Z_t$ :

$$\beta_{i,t} = b_{0,i} + b_{1,i} Z_t$$

- If we can identify and observe  $Z_t$ , we can run the following regression:

$$\bar{R}_{i,t} = \alpha_i + b_{0,i} \bar{R}_{m,t} + b_{1,i} (Z_t \bar{R}_{m,t}) + \epsilon_{i,t}$$

and test  $H_0 : \alpha_i = 0$  for  $i = 1, \dots, n$

- Shortcomings:
  - Require correct choice of  $Z_t$ 's, which is hard
  - Many standard choices of  $Z_t$ 's are only available at low frequencies, so we have small effective sample size

### *Estimation Method 2: Model Time-Variation as Structural Breaks*

- Model time-variation as structural change
- We can model CCAPM as a CAPM regression with structural breaks (treat  $\alpha_{i,t}, \beta_{i,t}$  as potentially changing coefficients)
  - and we could use [QLR Test](#) to identify structural changes in  $\beta_{i,t}$
- Advantage: more robust because it's model-free (no misspecification issues about  $Z_t$ )
- Disadvantage: less efficient and provides no explanation for time-variation

### *Estimation Method 3: Rolling-Window Estimator*

- Rolling-window estimator
- Setup:

$$\bar{R}_{i,t} = \beta_{i,t} \bar{R}_{m,t} + \epsilon_{i,t}$$

and

$$\begin{cases} \hat{\beta}_{i,t} = \frac{\sum_{s=1}^T w_{s,t} (\bar{R}_{m,s} - \hat{\mu}_{\bar{R}_{m,t}}) \bar{R}_{i,s}}{\sum_{s=1}^T w_{s,t} (\bar{R}_{m,s} - \hat{\mu}_{\bar{R}_{m,t}})^2} \\ \hat{\alpha}_{i,t} = \hat{\mu}_{\bar{R}_{i,t}} - \hat{\beta}_{i,t} \hat{\mu}_{\bar{R}_{m,t}} \\ \hat{\mu}_{\bar{R}_{i,t}} = \sum_{s=1}^T w_{s,t} \bar{R}_{i,s}, \hat{\mu}_{\bar{R}_{m,t}} = \sum_{s=1}^T w_{s,t} \bar{R}_{m,s} \end{cases}$$

where weights satisfy  $\sum_s w_{s,t} = 1$ . For example:

$$w_{s,t} = \begin{cases} \frac{1}{M} & , t - M \leq s \leq t \\ 0 & , \text{otherwise} \end{cases}$$

- The choice of  $M$  is critical, and we can use POOs method.

## *Arbitrage Pricing Theory (APT)*

### *Setup*

- Arbitrage Pricing Theory (APT)
- CAPM assumes the only source of risk is from the covariance with the overall market portfolio; APT is a [multi-factor model](#) allowing for multiple sources of risk
- Does not rely on specific assumptions of utilities, but simply on no-arbitrage
- A financial market:

$$\bar{R}_{i,t} = \alpha_i + \beta_{i,1} F_{1,t} + \cdots + \beta_{i,K} F_{K,t} + \epsilon_{i,t} \quad (\text{APT})$$

where

- $F_{1,t}, \dots, F_{K,t}$  are a set of common factors
- Each asset has a set of factor loadings  $\beta_i = (\beta_{1,i} \dots \beta_{K,i})'$  quantifying the impact of each factor on the excess return of asset  $i$
- $\epsilon_{i,t}$  are idiosyncratic noise/risk satisfying  $\mathbb{E}[\epsilon_{i,t}|F_{1,t}, \dots, F_{K,t}] = 0$
- or:

$$\mathbb{E}[\bar{R}_{i,t}] = \beta_{i,1}\lambda_1 + \dots + \beta_{i,K}\lambda_K$$

where:

- $\lambda_k = \mathbb{E}[F_{k,t}]$  is the risk premium associated with the  $k$ th factor (price of risk)
- $\beta_{i,k}$  is the exposure of asset  $i$  to risk factor  $k$  (quantity of risk)

### *Choice of factors:*

- **Choice of Factors**
- Macro factors: variables that links "real economy" (production) to nominal one (financial markets); e.g. growth rate of industrial production (business cycle), spread between corporate and government bond yields (changes in aggregate level of risk)
- Returns of traded portfolios: Factors chosen as returns of a set of portfolios; e.g. Fama-French factor model
- Observable characteristics (aka "fundamentals"): Factors are unobserved while beta's are modelled as observable functions of firm-specific characteristics; e.g. size of firm, past performance

### *Testing*

- Any testing of multi-factor models are conditional on having chosen the correct factors
- Under no-arbitrage:

$$\mathbb{E}[\bar{R}_{i,t}] = \beta_{i,1}\lambda_1 + \dots + \beta_{i,K}\lambda_K$$

where

- $\lambda_k = \mathbb{E}[F_{k,t}]$  is the risk premium associated with the  $k$ th factor (price of risk)
- $\beta_{i,k}$  is the exposure of asset  $i$  to risk factor  $k$  (quantity of risk)
- Therefore, we can test APT by running regression (APT) and test  $H_0 : \alpha_i = 0 \forall i$
- **Testing Observed factor models:**
  - Suppose we have data on  $n$  different assets/portfolios
  - Collect their individual excess returns in  $\bar{R}_t = (R_{1,t}, \dots, R_{n,t})'$
  - Rewrite joint APT model as a vector regression:

$$\bar{R}_t = \alpha + \beta' F_t + \epsilon_t, \quad \mathbb{E}[\epsilon_t|F_t] = 0$$

- Estimate the  $\alpha$ 's and  $\beta$ 's jointly
- Test  $H_0 : \alpha_1, \dots, \alpha_n = 0$

- Testing Characteristic-based factor models:

- Setup:

$$\bar{R}_{i,t} = \alpha_i + \beta_{i,1}F_{1,t} + \cdots + \beta_{i,K}F_{K,t} + \epsilon_{i,t}$$

but we don't know the true factors. Instead, we assume:

$$\beta_{i,j} = b_{i,j}X_i$$

where  $X_i$  are observed characteristics of firm  $i$  and  $b_{i,j}$  are loadings of these characteristics

- Rewrite the model (treat  $F_t^* = b'_i F$  as unknown and use  $X_i$  as regressors):

$$\bar{R}_{i,t} = \alpha_i + (b_{i,1}F_{1,t} + \cdots + b_{i,K}F_{K,t})X_i + \epsilon_{i,t} = \alpha_i + F_t^*X_i + \epsilon_{i,t}$$

- Estimate the model and test  $\alpha_i = 0 \forall i$
- *If rejected, reasons:*
  - Consumers are irrational / EMH violated
  - Mis-specification: wrong factors have been included / important ones have been excluded

## Critical Issues

- Some Critical Issues
  - In addition to CAPM, APT is more subject to **data-snooping**: If a factor is included using standard statistical tools (e.g. t-statistic), then with 5% probability a given factor will be included even though it is insignificant
- 

## Week 10: Volatility Models

### 10.1 [Volatility Models] - A

#### Heteroskedasticity & Notations

- Heteroskedasticity
- Conditional variance:

$$\sigma_t^2 = \sigma^2(\mathcal{I}_{t-1}) = \text{Var}(\epsilon_t | \mathcal{I}_{t-1}) = \text{Var}(Y_t | \mathcal{I}_{t-1})$$

- We use  $\epsilon_t$  instead of  $Y_t$  as inputs (conditional on  $\mathcal{I}_{t-1}$ , they are the same: knowing  $\epsilon_t \iff Y_t$ )

#### Multiplicative Volatility Models (including all we discussed after)

- Multiplicative volatility model

- rescaled error term:

$$z_t = \frac{\epsilon_t}{\sigma_t} \iff \epsilon_t = \sigma_t z_t$$

and  $z_t$  captures *surprise movements* in  $\epsilon_t$  after controlling for the level of volatility  $\sigma_t$

- By construction:  $\mathbb{E}[z_t | \mathcal{I}_{t-1}] = 0, \text{Var}(z_t | \mathcal{I}_{t-1}) = 1$

## ARCH Model

- The ARCH Model: autoregressive conditional heteroskedastic model

- *Assumptions*:

- ARCH.1: Volatility  $\sigma_t^2$  depends on a finite number of lags  $p$ :

$$\sigma_t^2 = \sigma^2(\epsilon_{t-1}, \epsilon_{t-2}, \dots, \epsilon_{t-p})$$

- ARCH.2:  $\sigma^2(\epsilon_{t-1}, \epsilon_{t-2}, \dots, \epsilon_{t-p})$  has a simple linear form. In ARCH(1), this is just

$$\sigma^2(\epsilon_{t-1}, \epsilon_{t-2}, \dots, \epsilon_{t-p}) = w + \alpha \epsilon_{t-1}^2$$

## ARCH(1)

- Setup of ARCH(1):

$$\epsilon_t = \sigma_t z_t, \quad \sigma_t^2 = w + \alpha \epsilon_{t-1}^2 \quad (\text{ARCH}(1))$$

with  $w > 0, \alpha \geq 0$

- Conditional variance in ARCH(1):

$$\text{Var}(Y_t | \mathcal{I}_{t-1}) = \text{Var}(\epsilon_t | \mathcal{I}_{t-1}) = \mathbb{E}[\sigma_t^2 z_t^2 | \mathcal{I}_{t-1}] = \sigma_t^2 = w + \alpha \epsilon_{t-1}^2$$

- Time series properties:

- Rewrite  $\sigma_t^2$  as an *AR(1)* process:

$$\sigma_t^2 = w + \alpha \sigma_{t-1}^2 + u_t, \quad u_t = \alpha \sigma_{t-1}^2 (z_{t-1}^2 - 1)$$

where  $\mathbb{E}[u_t | \sigma_{t-1}^2] = 0$  and  $\mathbb{E}[u_t u_{t+h}] = 0$

- $\rightsquigarrow$  we can use all properties from AR(1), such as:

$$\rho_{\sigma_t^2}(k) = \alpha^k$$

- Long-run/Unconditional variance: taking expectation on both sides, we get:

$$\sigma_\epsilon^2 = \frac{w}{1 - \alpha}$$

- If  $\alpha < 1$ , then  $\sigma_\epsilon^2$  exists and  $\epsilon_t = \sigma_t z_t$  is stationary
- This is not the unconditional variance of  $Y_t$ , which needs to be analysed with the model of  $Y_t$  itself

## ARCH( $q$ )

- Setup:

$$\epsilon_t = \sigma_t z_t, \quad \sigma_t^2 = w + \sum_{i=1}^q \alpha_i \epsilon_{t-1}^2 \quad (\text{ARCH(q)})$$

- Long-run/Unconditional variance:

$$\sigma_Y^2 = \sigma_\epsilon^2 = \frac{w}{1 - \sum_{i=1}^q \alpha_i}$$

- If  $\sum_{i=1}^q \alpha_i < 1$ , ARCH( $q$ ) process is stationary with a well-defined 2nd moment

## GARCH Model

- GARCH(1,1):

$$\epsilon_t = \sigma_t z_t, \quad \sigma_t^2 = w + \alpha \epsilon_{t-1}^2 + \beta \sigma_{t-1}^2 \quad (\text{GARCH}(1,1))$$

- GARCH(1,1) = restricted ARCH( $\infty$ ): by backward recursion (注意只 iterate  $\sigma$ , 不要 iterate  $\epsilon$ ) :

$$\begin{aligned} \sigma_t^2 &= w + \alpha \epsilon_{t-1}^2 + \beta(w + \alpha \epsilon_{t-2}^2 + \beta \sigma_{t-2}^2) \\ &\dots \\ &= w \underbrace{\sum_{i=0}^{\infty} \beta^i}_{\frac{w}{1-\beta}} + \alpha \sum_{i=1}^{\infty} \beta^{i-1} \epsilon_{t-i}^2 + \underbrace{\beta^\infty \sigma_{t-\infty}^2}_0 \\ &= \frac{w}{1-\beta} + \alpha \sum_{i=1}^{\infty} \beta^{i-1} \epsilon_{t-i}^2 \end{aligned}$$

- It allows all lags of  $\epsilon$  to affect current volatility
- Long-run/Unconditional Variance:

$$\mathbb{E}[\sigma_t^2] = \frac{w}{1 - \alpha - \beta}$$

- $\implies$  Requirement for a valid (non-exploding & positive) long-run variance:

$$w > 0, \alpha \geq 0, \beta \geq 0, \alpha + \beta < 1$$

## Forecasting Volatility and Evaluation

### One-Step Forecast

- ARCH/GARCH deliver exact one-step ahead forecasts ( $\sigma_{T+1|T}^2$ ). Ignoring parameter uncertainty, there is no forecast error ( $\sigma_{T+1|T}^2 = \sigma_{T+1}^2$ )
- Plugging in estimated parameters, we get the feasible forecast. Example: for ARCH(1):

$$\hat{\sigma}_{T+1}^2 = \hat{w} + \hat{\alpha} \epsilon_T^2$$

## Multi-Step Forecast

- The optimal square-loss forecast:

$$\sigma_{T+h|T}^2 = \mathbb{E} [\sigma_{T+h}^2 | \mathcal{I}_T]$$

- Example: ARCH(1):

$$\sigma_{T+h|T}^2 = w + \alpha \mathbb{E} [\epsilon_{T+h-1}^2 | \mathcal{I}_T] = w + \alpha \sigma_{T+h-1|T}^2 = \dots$$

- Iterating backward, this is equal to

$$w(1 + \alpha + \dots + \alpha^{h-1}) + \alpha^h \epsilon_T^2$$

- Plugging in estimated parameters, we get the **feasible h-step forecast**:

$$\hat{\sigma}_{T+h|T}^2 = \hat{w}(1 + \hat{\alpha} + \dots + \hat{\alpha}^{h-1}) + \hat{\alpha}^h \epsilon_T^2$$

## Forecast Errors

- Arises only in multi-step forecast  $h > 1$
- Volatility forecast error (*no parameter uncertainty*):

$$e_{T+h} = \sigma_{T+h}^2 - \sigma_{T+h|T}^2$$

- Volatility forecast error (*with parameter uncertainty*):

$$\hat{e}_{T+h} = \sigma_{T+h}^2 - \hat{\sigma}_{T+h|T}^2 = e_{T+h} + (\sigma_{T+h|T}^2 - \hat{\sigma}_{T+h|T}^2)$$

- Parameter uncertainty vanishes as  $T \rightarrow \infty$ , so we typically ignore it:

$$\lim_{T \rightarrow \infty} \hat{e}_{T+h} = e_{T+h}$$

- Example: 2-step ahead forecast error of ARCH(1):

$$e_{T+2} = \sigma_{T+2}^2 - \sigma_{T+2|T}^2 = (w + \alpha \epsilon_{T+1}^2) - (w + \alpha \sigma_{T+1}^2) = \alpha \sigma_{T+1}^2 (z_{T+1}^2 - 1)$$

- This implies our forecast is unbiased:

$$\mathbb{E}[e_{T+2} | \mathcal{I}_T] = 0, \quad \underbrace{\mathbb{E}[e_{T+2}^2 | \mathcal{I}_T]}_{MSFE} = \alpha^2 \kappa_4 \sigma_{T+1}^4$$

$$\text{where } \kappa_4 = \mathbb{E} [(z_{T+1}^2 - 1)^2]$$

- MSFE:

$$MSFE = \alpha^2 \kappa_4 \mathbb{E}[\sigma_{T+1}^4]$$

## Forecast Evaluation

- We don't observe the actual volatility  $\sigma_t^2$ . Instead, we use a noisy measure of volatility  $\epsilon_t^2$  to proxy it:

$$\epsilon_t^2 = \sigma_t^2 + \underbrace{(\epsilon_t^2 - \sigma_t^2)}_{u_t}$$

where  $u_t = \sigma_t^2(z_t^2 - 1)$  is the measurement error:

$$\mathbb{E}[u_t | \mathcal{I}_{t-1}] = 0, \quad \mathbb{E}[u_t^2 | \mathcal{I}_{t-1}] = \sigma_t^4 \kappa_4$$

where  $\kappa_4 = \mathbb{E}[(z_{t+1}^2 - 1)^2]$

- Thus,  $\epsilon_t^2$  is a noisy but unbiased proxy for  $\sigma_t^2$  with error variance  $\sigma_t^2 \kappa_4$
- Forecast evaluation with proxy: For a given forecast  $\hat{\sigma}_{t+1|t}^2$ , the mean square forecast error can be calculated by POOs:

$$\widehat{MSFE} = \frac{1}{T - t_1 - 1} \sum_{t=t_1}^{T-1} (\epsilon_{t+1}^2 - \hat{\sigma}_{t+1|t}^2)^2$$

## 10 2 [Volatility Models II] - A

### OLS as Gaussian MLE in AR

- Gaussian MLE in AR

### Gaussian MLE of ARCH(1): Estimation

- Gaussian MLE is more efficient than OLS in estimating ARCH
- Example: ARCH(1)
- Let:

$$\begin{cases} Y_t = \mu + \epsilon_t \\ \epsilon_t = \sigma_t z_t, \quad \sigma_t^2 = w + \alpha \epsilon_{t-1}^2 \end{cases}$$

and suppose  $z_t \sim^{iid} \mathcal{N}(0, 1) \implies Y_t | Y_{t-1} \sim \mathcal{N}(\mu, \sigma_t^2)$

- The conditional density (likelihood) of individual observation is:

$$f_\theta(Y_t | Y_{t-1}) = \frac{1}{\sqrt{2\pi\sigma_t^2}} \exp \left[ -\frac{(Y_t - \mu)^2}{2\sigma_t^2} \right]$$

- Then the log-likelihood of a single observation is:

$$\log f_\theta(Y_t | Y_{t-1}) = -\frac{1}{2} \log(2\pi) - \frac{1}{2} \log(\sigma_t^2) - \frac{(Y_t - \mu)^2}{2\sigma_t^2}$$

where  $\sigma_t^2 = w + \alpha(Y_{t-1} - \mu)^2$

- Gaussian MLE of  $\theta = (\mu, w, \alpha)$  maximised the joint log-likelihood:

$$\begin{aligned}\hat{\theta}_{MLE} &= \arg \max_{\theta} \sum_{t=1}^T \log f_{\theta}(Y_t | Y_{t-1}) \\ &= \arg \max_{\theta} \sum_{t=1}^T \left\{ -\frac{1}{2} \log(2\pi) - \frac{1}{2} \log(\sigma_t^2) - \frac{(Y_t - \mu)^2}{2\sigma_t^2} \right\}\end{aligned}$$

## Inference in ARCH Models Estimated by MLE

### Asymptotic Distribution

- Inference in ARCH Model
- If data is stationary and mixing, the MLE satisfies:

$$\hat{\theta}_{MLE} \sim^a N \left( \theta, \underbrace{\frac{1}{T} H^{-1} \Omega H^{-1} U}_{\text{Robust SE}} \right)$$

and if indeed  $z_t \sim N(0, 1)$  then  $\Omega = H$ :

$$\hat{\theta}_{MLE} \sim^a N \left( \theta, \frac{1}{T} H^{-1} \right)$$

### Hypothesis Testing

- Hypothesis Testing
- To test individual hypothesis, we use t-statistics:

$$t = \frac{\hat{\theta}_{MLE,i} - \theta_i}{\hat{\sigma}_{ii}}$$

where  $\hat{\sigma}_{ii}$  is the  $(i, i)$ th element of  $\frac{1}{T} \hat{H}^{-1} \hat{\Omega} \hat{H}^{-1}$

- Joint hypothesis test can be done with Likelihood-Ratio (LR) statistic:

$$LR = 2T \left\{ \underbrace{\log f_{\hat{\theta}_{MLE}}(Y_2, \dots, Y_T | Y_1)}_{\text{Unrestricted}} - \underbrace{\log f_{\tilde{\theta}_{MLE}}(Y_2, \dots, Y_T | Y_1)}_{\text{Restricted}} \right\}$$

and under the null of restricted model:

$$LR \xrightarrow{d} \chi_m^2$$

and  $m$  is the number of restrictions

## Others

- ECON0022 Review Questions
- ECON0022 Week 7 Questions

- F-Statistics:

$$F = \frac{SSR_R - SSR_U}{SSR_U} \times \frac{n - k}{q}$$

- Log-Normal Expectation:

$$\ln X \sim \mathcal{N}(\mu, \sigma^2) \implies \mathbb{E}[X] = \exp\left(\mu + \frac{\sigma^2}{2}\right)$$