

## DM-Quiz-2020-Q2

52.94% (9/17)

✗ 1. Select the intercept-only models, if any:

- ☒ A  $y = b_0 + b_1 \cdot x$
- ☐ B  $y = b_0 + b_1 \cdot x_1 + b_2 \cdot x_2$
- ☐ C  $\ln(y) = b_0$
- ☐ D  $y = e^{(b_1 \cdot x)}$
- ☐ E I do not know

✓ 2. The interpretation of adjusted  $R^2$  for multiple linear regression is the same as the interpretation of  $R^2$  for simple linear regression.

- ☐ A True
- ☒ B False
- ☐ C I do not know

✓ 3. Which one is observable ("visible")?

- ☒ A  $e$  (residuals)
- ☐ B  $\varepsilon$  (regression error)
- ☐ C neither
- ☐ D I do not know

✗ 4. The estimation of  $\beta$  is distributed as:

- ☐ A  $b \sim N(0, \sigma^2)$
- ☐ B  $b \sim N(\beta, \sigma^2)$
- ☐ C  $b \sim N(\beta, \sigma^2(X^T X)^{-1})$
- ☒ D It does not have distribution
- ☐ E I do not know

✗ 5. How many parameters are estimated by OLS in the case of simple linear regression?

- ☐ A 1
- ☒ B 2
- ☐ C 3
- ☐ D I do not know

✓ 6. Estimation of  $e$  (residuals) is distributed as:

- ☐ A  $e \sim N(0, \sigma^2)$
- ☒ B  $e \sim N(0, \sigma^2 M)$  ( $M$  is a matrix)
- ☐ C It does not have distribution
- ☐ D I do not know

✗ 7.  $b$  parameters can be computed using only

- ☒ A OLS
- ☐ B ML
- ☐ C Neither
- ☐ D I do not know

✗ 8. In the case of multiple linear regression

- ☐ A  $\text{adj } R^2 < 1$  (always)
- ☒ B  $\text{adj } R^2 \leq 1$  (always)
- ☐ C I do not know

✓ 9.  $b =$

- ☒ A  $(X'X)^{-1}X'Y$
- ☐ B  $(X'X)^{-1}Y'X$
- ☐ C  $(Y'X)^{-1}Y'X$
- ☐ D I do not know

✓ 10. The total sum of squares equals

- ☒ A  $\text{Sum}((y - \text{mean}(y))^2)$
- ☐ B  $\text{Sum}((\hat{y} - \text{mean}(y))^2)$
- ☐ C  $\text{Sum}((y - \hat{y})^2)$
- ☐ D I do not know

✓ 11. The regression sum of squares equals

- ☐ A  $\text{Sum}((y - \text{mean}(y))^2)$
- ☒ B  $\text{Sum}((\hat{y} - \text{mean}(y))^2)$
- ☐ C  $\text{Sum}((y - \hat{y})^2)$
- ☐ D I do not know

✓ 12. If RSS is the regression sum of squares and ESS is the error sum of squares then

- ☒ A  $R^2 = 1 - \text{ESS}/\text{TSS}$
- ☐ B  $R^2 = \text{ESS}/\text{TSS}$
- ☐ C  $R^2 = \text{ESS}/\text{RSS}$
- ☐ D I do not know

✗ 13. Multicollinearity occurs when

- ☐ A  $\text{rank}(X) < m$  ( $m$  is the number of explanatory variables)
- ☒ B  $\text{var}(\epsilon) = \sigma^2 I$
- ☐ C  $E(\epsilon) = 0$
- ☐ D  $\text{cov}(\epsilon_i, \epsilon_j) = \text{const}$
- ☐ E I do not know

✓ 14. In simple linear regression model response variable ( $y$ ) can be

- ☐ A binary
- ☐ B categorical
- ☒ C numeric
- ☐ D ordinal
- ☐ E I do not know

✗ 15. In a simple linear regression model, explanatory variables can be

- ☐ A binary
- ☐ B categorical
- ☒ C numeric
- ☐ D ordinal
- ☐ E I do not know
- ☐ F all answers are correct
- ☐ G Neither

✗ 16. If  $A$  is a matrix,  $X$  is the vector of random variables, then  $\text{var}(AX) =$

- ☐ A  $A'\text{var}(X)A$
- ☒ B  $A^2\text{var}(X)$
- ☐ C  $\text{var}(x)$
- ☐ D Can not be calculated
- ☐ E I do not know

✓ 17. Which of the answers can be used to conclude about the significance of variables (if any)?

- ☒ A t values
- ☐ B Estimated coefficients (only)
- ☐ C SE of estimated coefficients (only)
- ☐ D I do not know