



# Cation-Chloride Cotransporters and Neuronal Function

Peter Blaesse,<sup>1</sup> Matti S. Airaksinen,<sup>2</sup> Claudio Rivera,<sup>3</sup> and Kai Kaila<sup>1,2,\*</sup>

<sup>1</sup>Department of Biological and Environmental Sciences, University of Helsinki, Viikinkaari 1, FIN-00014, Helsinki, Finland

<sup>2</sup>Neuroscience Center, University of Helsinki, Viikinkaari 4, FIN-00014, Helsinki, Finland

<sup>3</sup>Institute of Biotechnology, University of Helsinki, Viikinkaari 9, FIN-00014, Helsinki, Finland

\*Correspondence: kai.kaila@helsinki.fi

DOI 10.1016/j.neuron.2009.03.003

Recent years have witnessed a steep increase in studies on the diverse roles of neuronal cation-chloride cotransporters (CCCs). The versatility of CCC gene transcription, posttranslational modification, and trafficking are on par with what is known about ion channels. The cell-specific and subcellular expression patterns of different CCC isoforms have a key role in modifying a neuron's electrophysiological phenotype during development, synaptic plasticity, and disease. While having a major role in controlling responses mediated by GABA<sub>A</sub> and glycine receptors, CCCs also show close interactions with glutamatergic signaling. A crosstalk among CCCs and trophic factors is important in short-term and long-term modification of neuronal properties. CCCs appear to be multifunctional proteins that are also involved in shaping neuronal structure at various stages of development, from stem cells to synaptogenesis. The rapidly expanding work on CCCs promotes our understanding of fundamental mechanisms that control brain development and functions under normal and pathophysiological conditions.

#### Introduction

Ion trafficking across biological membranes is mediated by channels and transporters, and these two types of molecules form the basis of electrical signaling in the nervous system. However, the volume of research on voltage- and ligand-gated channels is orders of magnitude higher than what has been devoted to the study of neuronal ion transporters. This bias has no rational foundation: it does not reflect a difference in the impact or versatility of their functions. What is often ignored is that ion transporters are as important as channels in the generation of electrical signals.

During the recent years, there has been a sharp increase in the interest of neurobiologists in plasmalemmal ion-transport mechanisms, especially in cation chloride cotransporters (CCCs). Initially, CCCs were studied in nonneuronal cells in order to elucidate their role in volume regulation (Payne et al., 2003). More recently, it has turned out that these transporters are crucial in the control of the electrochemical Cl<sup>-</sup> gradient that is required for "classical" hyperpolarizing postsynaptic inhibition mediated by GABA<sub>A</sub> receptors (GABA<sub>A</sub>Rs) and glycine receptors (Figure 1). The rapidly expanding research within this field has shown that CCCs play a fundamental role in neuronal proliferation, differentiation, disease, damage, and recovery, as well as in a variety of phenomena traditionally associated with plasmalemmal receptor- and channel-mediated effects, such as those involved in synaptic plasticity. Different CCCs show distinct cellular and subcellular expression patterns, and their functions are modulated via short-term and long-term changes at the level of transcription, translation, posttranslational modifications, and trafficking. Furthermore, CCCs are involved in intricate signaling cascades with growth factors and hormones. In view of their functional versatility, CCCs are likely to gain importance in drug research.

The milieu intérieur of the central nervous system (CNS) is tightly regulated by the blood-brain barrier and other epithelial structures, such as the choroid plexus, which involves a wide spectrum of ion transporters, including CCCs (Pedersen et al., 2006; Praetorius, 2007). These topics will not be discussed in the present review, and, with our main focus on neuronal CCCs, only some basic aspects of the ion-regulatory functions of glial cells will be touched upon. Outside the nervous system, CCCs are expressed in all organ systems, hence playing a key role in, for instance, salt transport and the control of blood pressure (Pedersen et al., 2006; Russell, 2000; Flatman, 2008). It is important to recognize the wide spectrum of CCC functions in experiments on transporters.


#### **Structure and Molecular Diversity**

The CCC family in mammals consists of nine members encoded by the genes *Slc12a1-9*. The CCC proteins are glycoproteins with apparent molecular weights of 120–200 kDa. Seven out of the nine CCCs described so far are plasmalemmal ion transporters (Gamba, 2005; Mercado et al., 2004; Payne et al., 2003), and in terms of function they fall into three categories: two members are Na-K-2Cl cotransporters (NKCCs; isoforms NKCC1 and NKCC2), one is an Na-Cl cotransporter (NCC), and four are K-Cl cotransporters (KCCs; isoforms KCC1-4). The physiological roles of the remaining two CCCs (CIP1 or *Slc12a8*, and CCC9 or *Slc12a9*) are yet unknown.

The predicted secondary structure of CCCs, confirmed only for NKCC1 so far (Gerelsaikhan and Turner, 2000), consists of 12 membrane spanning segments that are flanked by intracellular termini that constitute about half of the protein. Heteroand homo-oligomers of the different CCCs have been described for nearly all CCCs (Blaesse et al., 2006; Pedersen et al., 2008;

### Review








Figure 1. Ion-Transport Mechanisms Underlying GABAA and Glycine Receptor-Mediated Responses in Immature and Mature Neurons (Left) NKCC1 mediates CI uptake in immature hippocampal and cortical neurons, while an as yet unidentified CI transporter denoted by X operates in some neurons either in parallel with NKCC1 or in its absence. (Right) In mature CNS neurons, KCC2 is the principal K-CI cotransporter, and it is often coexpressed with KCC3. CI<sup>-</sup> transport by all CCCs is fuelled by the Na<sup>+</sup> and K<sup>+</sup> gradients generated by the Na-K ATPase. Because the intracellular concentration of CI<sup>-</sup> is high in immature neurons (indicated by the shade of blue), EGABA and EGIV are not directly affected by the intracellular HCO3<sup>-</sup> concentration, while in mature neurons, the channel-mediated net efflux of HCO<sub>2</sub> can have a major effect on the reversal potentials. For further details, see text.

Simard et al., 2007; and references cited therein). However, the available data are not conclusive regarding the mechanisms whereby oligomerization affects CCC function.

With the exception of NKCC2 and NCC, which are predominantly found in kidney, all CCCs are expressed in neurons or glial cells-or both-at least at some stage of CNS development. KCC1 is considered a "household" isoform in various cell types, including glial cells, but present data suggest little or no expression in central neurons (Kanaka et al., 2001; Li et al., 2002; Payne et al., 1996; Rivera et al., 1999; Rust et al., 2007). In contrast, KCC2 is an exceptional CCC, as it is exclusively expressed in CNS neurons (Karadsheh and Delpire, 2001; Payne et al., 1996; Uvarov et al., 2005; Williams et al., 1999). There has been surprisingly little work on KCC3 despite its presence in various brain regions (Becker et al., 2003; Boettger et al., 2003; Le Rouzic et al., 2006; Pearson et al., 2001). Interestingly, KCC3 protein levels show a developmental increase (Boettger et al., 2003; Pearson et al., 2001) that is similar to the extensively studied upregulation of KCC2, a topic to be discussed in detail below. Even less is known about the possible role of KCC4 in the nervous system, an isoform that is expressed at a high level in the embryonic brain (Li et al., 2002). Another CCC that has attracted considerable interest among neurobiologists is NKCC1 (Gamba, 2005; Payne et al., 2003).

#### **Evolution of CCCs**

Transporter proteins, such as CCCs, may have arisen by duplication of genes encoding channel-type protein subunits with a smaller number of transmembrane segments (Saier, 2003). The existence of putative CCC homologs in C. elegans and in a cyanobacterium points to an early evolutionary origin of this protein family (Xu et al., 1994). Consistent with this, homologs of CCCs are present in diverse organisms. Many plant species have a CCC gene that seems to encode for a functional burnetanide-sensitive Na-K-2Cl cotransporter (Colmenero-Flores et al., 2007). Given that Drosophila apparently has only one KCC gene, and that its expression is neuron specific (Hekmat-Scafe et al., 2006), it is tempting to speculate that the first KCC evolved

in animals with a nervous system to serve inhibitory neurotransmission.

#### Alternative Splicing

Alternative splicing greatly increases the diversity of the proteome. Alternative transcripts have often distinct expression patterns, and they can be translated into proteins with different structures, phosphorylation sites, or subcellular localizations. With regard to CCCs expressed in the CNS, splice variants are known for KCC1, KCC2, KCC3, and NKCC1 (Gamba, 2005; Mercado et al., 2004; Uvarov et al., 2007). KCC3a is expressed in the brain, whereas KCC3b is restricted to organs outside the CNS. In the adult rat brain, alternative splicing of KCC3a has been reported to create a neuron-specific KCC3c form, whereas the native KCC3a is restricted to glia (Le Rouzic et al., 2006). Two functional isoforms with distinct expression patterns were described for NKCC1 (Randall et al., 1997; Vibat et al., 2001). NKCC1a and NKCC1b are both expressed ubiquitously, but the relative expression level for NKCC1b is higher in the adult brain. The difference between NKCC1a and NKCC1b is an NKCC1a-specific 16 amino acid peptide encoded by exon-21 that contains a putative PKA phosphorylation site. In three out of four NKCC1 mutant mice described so far, both isoforms are disrupted (Table 1). However, the mouse characterized by Dixon et al. (1999) is an NKCC1a-specific knockout.

The mammalian KCC2 (Slc12a5) gene generates two isoforms, KCC2a and KCC2b, with different N termini (Uvarov et al., 2007). The mRNA levels of the two isoforms are similar in the neonatal mouse. While the overall expression of KCC2a remains relatively constant during postnatal development, the expression of KCC2b is strongly upregulated, especially in the cortex (Uvarov et al., 2007). This indicates that KCC2b is responsible for the extensively studied "developmental shift" from depolarizing to hyperpolarizing GABAergic responses, a conclusion supported by the finding that GABAergic responses remain depolarizing in cortical cultures from knockout mice (Zhu et al., 2005), which are now known to lack KCC2b only (Uvarov et al., 2007).



Table 1. Major Phenotypes of Mice with Disruptions of Genes Encoding CCCs			
	Affected Isoforms	Mice Described by	Major Characteristics of Phenotype
KCC1	KCC1 knockout	Rust et al., 2007	no obvious phenotype
KCC2	KCC2a and KCC2b knockout	Hübner et al., 2001b	die after birth due to respiratory failure
	KCC2b knockout	Woo et al., 2002	survival up to P17; generalized seizures; low body weight
	hypomorphic mice; ~17% of normal KCC2 (KCC2a and KCC2b)	Tornberg et al., 2005	increased anxiety-like behavior; increased seizure susceptibility; impaired spatial learning and memory; reduced sensitivity to thermal and mechanical stimuli
KCC3	KCC3a-c knockout	Howard et al., 2002; Boettger et al., 2003	peripheral neuropathy; deafness (inner ear defects); neurodegeneration; reduced seizure threshold; hypertension (neurogenic mechanism; Rust et al., 2006)
KCC4	KCC4 knockout	Boettger et al., 2002	deafness (inner ear defects); renal tubular acidosis
NKCC1	NKCC1a and NKCC1b knockout	Delpire et al., 1999; Flagella et al., 1999; Pace et al., 2000	deafness and imbalance (inner ear defects); impaired blood pressure; growth retardation; defective spermatogenesis; alterations in locomotion; increased threshold to thermal stimulation
	NKCC1a knockout	Dixon et al., 1999	deafness and imbalance (inner ear defects)

Analyzing the phenotypes of knockout mice has shed some light on the physiological functions of the two KCC2 isoforms, which have distinct functions in the brain (Uvarov et al., 2007). Mice that completely lack KCC2 expression die immediately after birth due to severe motor defects, including respiratory failure (Hübner et al., 2001b). In contrast, KCC2b-specific knockout mice are viable, but show abnormal posture and gait and frequent seizures within a couple of days after birth and die at around 2 postnatal weeks, perhaps because of the recurrent seizures (Table 1; Woo et al., 2002). KCC2a is obviously important for some basic functions of lower brain structures, and it is sufficient to promote survival for up to 3 postnatal weeks, but the absence of KCC2b leads to seizures during this period.

Most published data on the spatial and temporal expression patterns of KCC2 reflect the expression of both KCC2a and KCC2b, because the mRNA probes and antibodies used in these studies detect both isoforms. Hence, in most of what is discussed below, "KCC2" will refer to both splice variants.

#### **Expression Patterns of CCCs**

A wealth of data indicate that one of the major factors underlying phenotypic variation among distinct types of neurons is based on the cell-specific as well as subcellular patterns of CCC expression.

#### **Neuron-Specific Expression of KCC2**

The expression of only a few genes is as tightly restricted to central neurons as *Slc12a5* (encoding KCC2). KCC2 mRNA is abundant in most neurons of the mature CNS but shows negligible expression in nonneuronal cell types, neuronal progenitors, or peripheral neurons. The CNS neuron-specific expression of KCC2 seems, at least in mammals, to be ensured by multiple, redundant mechanisms, including binding sites for neuron-enriched transcription factors such as Egr4 and neuron-restrictive silencing elements (Uvarov et al., 2005, 2006) (Figure 2).

KCC2 is located in the plasma membrane of somata and dendrites in various brain regions, e.g., cerebellar granule cells (Williams et al., 1999), thalamic relay cells (Bartho et al., 2004), auditory brainstem neurons (Blaesse et al., 2006), and cortical

neurons (Gulyas et al., 2001; Szabadics et al., 2006). Freund and coworkers demonstrated high levels of KCC2 in the dendritic spines of rat cortical pyramidal neurons and interneurons (Gulyas et al., 2001), a puzzling observation in light of the preferential localization of glutamatergic synapses to spines and GABAergic synapses to somata and dendritic shafts. However, this paradox is at least partly resolved by a recent study that demonstrated a morphogenic role of KCC2 in spine formation independent of its role in ion transport (Li et al., 2007; see below).

In immature neurons in the auditory brainstem, KCC2 is initially expressed at a high level in a functionally inactive form, and the neurons generate depolarizing GABA<sub>A</sub> and glycine responses due to the presence of an unidentified Cl<sup>-</sup> uptake transporter (denoted as "X" in Figure 1; Balakrishnan et al., 2003; Blaesse et al., 2006). Functionally inactive KCC2 has also been observed in primary cortical cultures during the first few days in vitro (Khirug et al., 2005). These results imply that the available data supporting a key role of NKCC1 in Cl<sup>-</sup> uptake in immature neocortical and hippocampal neurons cannot be directly extrapolated to other types of neurons; and further, the mere presence of KCC2 protein in a neuron does not necessarily imply that it is functionally active.

#### **Expression Patterns of NKCC1**

Because of inconsistent and contradictory results, general statements about developmental changes or about the cellular and subcellular distribution of NKCC1 in CNS neurons are hardly possible. Even the widespread idea that neuronal NKCC1 expression decreases during postnatal rodent development (Hübner et al., 2001a; Yamada et al., 2004) is challenged by data that show a developmental increase in the NKCC1 mRNA and protein levels (Clayton et al., 1998; Mikawa et al., 2002; Wang et al., 2002; Yan et al., 2001a). A paper by Delpire and coworkers (Plotkin et al., 1997), often cited as evidence for a postnatal downregulation of NKCC1 in the rat forebrain, actually demonstrates roughly similar levels at P0 and P21. As described below, NKCC1 is functional in a wide variety of both juvenile and mature central neurons.






Figure 2. Regulation of CCC Functionality

CCC functions are regulated by transcriptional control, alternative splicing, subcellular targeting, and posttranslational modifications that lead to cell-specific as well as discrete subcellular expression patterns of the distinct CCC proteins. (Upper left) An example of transcriptional control for KCC2. Neuron-enriched transcription factors (e.g., Egr4) and NRSEs are responsible for the neuron-specific expression of KCC2 and regulate the expression level (Uvarov et al., 2005, 2006). (Upper right panel) Alternative splicing of KCC2 and NKCC1 where the various splice variants have distinct binding sites for kinases such as SPAK/OSR and PKA (Randall et al., 1997; Uvarov et al., 2007). (Lower left) An example of subcellular targeting of NKCC1 into the axon (axon initial segment; Khirug et al., 2008) and of KCC2 into the dendritic shafts and spines. The mechanisms whereby various CCCs are targeted into axons versus dendrites and somata are not known. While protein 4.1N is involved in the anchoring of spine KCC2 (Li et al., 2007), it is not clear whether this is true for KCC2 located in dendritic shafts and neuronal somata (indicated by a question mark). (Lower right) Expression levels of plasmalemmal functional CCCs are influenced, among other things, by the rate of insertion from secretory (exocytic) vesicles and removal by endocytic vesicles. The illustration shows how kinases can modify CCC function by influencing the relative rates of endo- and exocytosis. In this specific example, the KCC2 endocytosis is suppressed by a PKC-mediated phosphorylation of KCC2 (Lee et al., 2007). The scheme also highlights the lack of evidence (indicated by a question mark) for direct modulation of the intrinsic rate of ion transport by neuronal CCCs.

An essential role of NKCC1 in cell proliferation has been proposed in many studies (for review, see Russell, 2000). The high expression of NKCC1 in embryonic ventricular zones (Hübner et al., 2001a; Li et al., 2002; but see Clayton et al., 1998) points to a role in neuronal proliferation. NKCC1 is also expressed in the inner ear, where a lack of this protein leads to a defect in the production of endolymph and hence to deafness that is not neuronal in origin (Delpire et al., 1999; Dixon et al., 1999). Similar effects have been observed in KCC3 and KCC4 knockout mice (Boettger et al., 2002, 2003). NKCC1 is also expressed in dorsal root ganglion (DRG) cells (Rocha-Gonzalez et al., 2008) and in olfactory receptor neurons (Kaneko et al., 2004; Reisert et al., 2005).

Much of the published data on NKCC1 protein expression in CNS neurons has been obtained using a monoclonal NKCC antibody, T4 (Lytle et al., 1995). While T4 is a useful tool in immunoblots (Lytle et al., 1995; Zhang et al., 2006), interpretation of these data are complicated by NKCC1 expression in both neurons and glia (Kanaka et al., 2001; Hübner et al., 2001a). Further, perhaps because of differences in the methods of epitope retrieval, there are conflicting data on T4's capability to reliably detect NKCC1 immunohistochemically (Chen et al., 2005; Dixon et al., 1999; Li et al., 2008; Zhang et al., 2006, 2007). An antibody generated by Forbush and coworkers detects the phosphorylated NKCC1, thereby providing data on both the presence and functional state of NKCC1 (Flemmer et al., 2002). In the absence of reliable immunohistochemical techniques, other methods, including specific pharmacological block with bumetanide and analysis of NKCC1 knockout mice, have been used to examine the expression patterns of functional NKCC1 (Khirug et al., 2008).

#### Coexpression of KCC2 and NKCC1

There are no data that would suggest K-Cl cotransport in axons and no evidence for the presence of axonal KCC2 protein (e.g.,





Hübner et al., 2001b; Szabadics et al., 2006). The terminals of retinal ON bipolar neurons have been reported to contain KCC2 (Vardi et al., 2000), but it is perhaps not appropriate to call these neurites "axons," as they do not generate action potentials. In contrast to this, functional evidence indicates that NKCC1 is expressed in axons (e.g., Jang et al., 2001; Khirug et al., 2008; see below). The mechanisms that underlie the differential subcellular targeting of distinct CCCs to the axonal and somatodendritic compartments are not known, but recent evidence from work on NKCC1 in epithelial cells suggests that alternative splicing can play a major role (Carmosino et al., 2008). The coexpression of NKCC1 and KCC2 in specific neurons that is evident on the basis of structural and functional data (e.g., Duebel et al., 2006; Gavrikov et al., 2003; Khirug et al., 2008; Martina et al., 2001; Marty and Llano, 2005; Romo-Parra et al., 2008; Szabadics et al., 2006; Trevino and Gutierrez, 2005; Vardi et al., 2000) indicates a tight control of [Cl<sup>-</sup>]<sub>i</sub> that can result in the generation of intraneuronal CI<sup>-</sup> gradients (Figure 2; and below).

#### KCC3 and KCC4 Expression Patterns

There is not much data on the neuronal expression patterns of KCC3 and KCC4, but what is available clearly indicates that further work on these two isoforms is highly motivated. In immunoblots, KCC3 was detected in all adult CNS regions, and KCC3 immunoreactivity was mainly located in myelinated tracts of the spinal cord and brain, although cells in the choroid plexus, pyramidal neurons in the hippocampus and cortex, and cerebellar Purkinje cells were also labeled (Pearson et al., 2001). In contrast, two other studies failed to observe significant KCC3 expression in the CNS white matter in adult mice (Boettger et al., 2003; Howard et al., 2002). KCC3 seems, however, to be expressed by the Schwann cells in peripheral nerves of juvenile mice (Byun and Delpire, 2007). KCC4 mRNA is highly expressed in the ventricular zone during embryonic development (Li et al., 2002), and in adult brain tissue it has been detected in the suprachiasmatic nucleus (Le Rouzic et al., 2006).

#### **Basic Properties and Modes of Operation of CCCs**

With regard to their source of energy, ion transporters fall into two categories: primary active transporters are directly fueled by ATP, while secondary active transporters take the energy for transport of the driven ion from the electrochemical gradient of another ion species. The ubiquitous primary active transporter, the Na-K ATPase, generates plasmalemmal K<sup>+</sup> and Na<sup>+</sup> gradients that provide the main source of energy for most secondary active transporters, including all CCCs (Figure 1). Extrusion of Cl<sup>-</sup> via KCC-mediated K-Cl cotransport is driven by the K<sup>+</sup> gradient, while uptake of Cl<sup>-</sup> in many (but not all; e.g., Balakrishnan et al., 2003; Zhang et al., 2007) neurons is driven by NKCC1 that exploits the plasmalemmal Na+ gradient as its energy source for Na-K-2Cl cotransport. The KCC2 cotransporter operates at near-equilibrium (Payne et al., 2003), which permits maximization of the efficiency of harvesting energy from the K<sup>+</sup> gradient to fuel the extrusion of Cl<sup>-</sup>. The transport stoichiometry of CCCs renders them electrically silent, thus they do not directly influence the neuronal membrane potential. Yet it should be noted that electroneutrality as such does not provide any information on the possible presence or absence

of effects of membrane voltage on the transport process (see Kaila, 1994).

The electroneutrality of CCCs is an obvious obstacle in electrophysiological work on CCCs, precluding direct monitoring of their function. Therefore, electrophysiological experiments on CCCs exploit anion-permeable channels (mainly GABA<sub>A</sub> and glycine receptors) as "read-out devices," either by monitoring the reversal potential of the channel-mediated currents or using more sophisticated Cl<sup>-</sup> loading techniques that will be described below. Additional techniques for analyzing ion transport by CCCs are based on assays of <sup>86</sup>Rb fluxes in cultured cells (Payne, 1997) and on the use of Cl<sup>-</sup>-sensitive indicators (Berglund et al., 2006; Verkman, 1990).

NKCC1 can be selectively blocked by a diuretic compound, bumetanide (1-10 μM), which has an up to 500-fold higher affinity for NKCC1 ( $K_i \sim 0.1 \mu M$ ) than for KCC2 (Payne et al., 2003). Furosemide, another diuretic, blocks both NKCC1 and KCCs with an equal potency, and it is often used at 100  $\mu M$  or higher to inhibit KCC2 in experiments on neuronal cultures and slices. 100 µM does not provide a full block, and an obvious confounding factor is the simultaneous effect on NKCC1. In addition, furosemide has inhibitory effects on N-methyl D-aspartate (NMDA) and GABAA receptors (Staley, 2002). Dihydroindenyloxy-alkanoic acid (DIOA) is yet another nonspecific drug that has been extensively used to block K-Cl cotransport (Payne et al., 2003). The lack of drugs with a specific action on KCC2 (and KCCs in general) has been a major problem in experimental work on neuronal CCCs. However, in a recent high-throughput screening study based on 234,000 compounds, a number of molecules that inhibit KCC2 with little effect on NKCC1 were identified (Delpire et al., 2009). Notably, an effective block of KCCs is achieved (sometimes on purpose but often unintentionally) in electrophysiological experiments based on whole-cell or gramicidin perforated-patch clamp recordings by replacing intracellular K+ with Cs+, a standard procedure to obtain a better space clamp. Cs<sup>+</sup> blocks KCCs because the cation selectivity sequence of K-Cl cotransport is  $K^+ \approx Rb^+ \approx NH_4^+ >> Cs^+$ (Payne et al., 2003; Williams and Payne, 2004).

#### **Neuronal Volume Regulation**

In the brain, cell swelling and the concomitant decrease in the interstitial space can lead to pathophysiological activity as well as to ischemia due to the compression of blood vessels. While cellular swelling induced by hypotonia (as often used in experiments on nonneuronal cells) causes a net influx of water and thereby leads to a fall in [Cl7], neuronal swelling is usually brought about by intense synaptic activity and a concomitant massive channel-mediated net influx of cations and Cl- that drives an osmotically obliged net influx of water. Hence, the mechanisms of swelling and KCC-mediated recovery are radically distinct under these two conditions. In case of neuronal swelling, recovery can be based on two distinct mechanisms: a genuine volume sensor that is able to kinetically activate K-Cl cotransport, and/or an enhanced driving force for K-Cl cotransport due to elevated intraneuronal Cl<sup>-</sup>. Interestingly, neurons are thought to be devoid of water channels (aquaporins; Amiry-Moghaddam and Ottersen, 2003), which lowers their sensitivity to osmotic challenges. Consistent with a role for KCC3 in neuronal volume regulation, recovery of cell volume after hypotonic



swelling is compromised in cortical neurons from KCC3 knockout mice (Boettger et al., 2003). In contrast, KCC2's role in neuronal volume regulation is probably secondary to the steep dependence of its net Cl<sup>-</sup> extrusion on the momentary intracellular chloride load. Nevertheless, this does not exclude an additional regulatory role for KCC2 in response to an osmotic challenge (Bergeron et al., 2006).

#### Setting E<sub>CI</sub> and E<sub>GABA-A</sub>

One of the main tasks of CCCs is to maintain a constant [CI]; in the face of channel-mediated fluxes that tend to dissipate the transmembrane electrochemical Cl<sup>-</sup> gradient, i.e., the driving force of  $GABA_AR$ -mediated  $CI^-$  currents.  $[CI^-]_i$  is also affected by chloride-dependent secondary active ion transporters involved in neuronal HCO<sub>3</sub><sup>-</sup>/pH regulation (Chesler, 2003; Hentschke et al., 2006; Romero et al., 2004). The sodium-independent CI/HCO<sub>3</sub> exchanger AE3 may act as a significant CI<sup>-</sup> uptake mechanism in neurons (Gonzalez-Islas et al., 2008), and in some neurons might represent transporter X in Figure 1. In addition to its role in neuronal Cl<sup>-</sup> homeostasis, HCO<sub>3</sub><sup>-</sup> acts as a significant carrier of depolarizing current across GABAARs (Kaila and Voipio, 1987). This is because neuronal pH regulation leads to a much higher intracellular level of HCO<sub>3</sub><sup>-</sup> than would be expected on the basis of a passive distribution. Hence, the reversal potential of GABAA receptor-mediated responses  $(E_{GABA-A})$  is not identical to the equilibrium potential of  $Cl^{-}(E_{Cl})$ , but shows a deviation to more positive values (Kaila and Voipio, 1987). The bicarbonate-mediated current component can be larger than the one mediated by Cl<sup>-</sup> in adult mammalian neurons (Gulledge and Stuart, 2003; Kaila et al., 1993). Notably, the reversal potential of the potassium-selective channels activated by GABA<sub>B</sub> receptors ( $E_{GABA-B}$ ) would be identical to  $E_{GABA-A}$  if (1) K-Cl cotransport were at equilibrium (see below) and (2) GABA<sub>A</sub> receptors were not permeable to bicarbonate. This is clearly not the case, and  $E_{GABA-B}$  attains values that are significantly more negative than E<sub>GABA-A</sub> (Farrant and Kaila, 2007; Kaila, 1994).

#### **Functional Regulation of CCCs** Thermodynamics versus Kinetics

Thermodynamics can be used when examining the energetic conditions that are a prerequisite for a change in a system from state A to state B, but the physical mechanisms and the time course of this change play no role in this context. The energetic conditions required for transmembrane ion movements to occur can be derived from thermodynamics, which allows, for example, calculating Nernstian ionic equilibrium potentials (in the case of channel-mediated movements) as well as the equilibrium conditions for a given ion transporter. It should be emphasized that a thermodynamic analysis does not tell anything about transport kinetics (the temporal, dynamic aspects of ion movements). Hence, even a steep, energetically favorable ion gradient does not "activate" an ion transporter. In a manner analogous to a closed ion channel, an ion transporter may be fully inactive despite the presence of a strong energetic driving force of its substrate ions. The kinetics of transport is determined by the number of functional plasmalemmal transporters as well as their intrinsic transport rates.

#### **Equilibrium Conditions for KCCs and NKCCs**

The reversal potential of a current mediated by an ion channel defines the electrochemical equilibrium for the current: despite a free pathway provided by open channels for electrodiffusion, no net movement of ions takes place. In an analogous manner, an ion transporter is at thermodynamic equilibrium when no net flux of ions is mediated by the transporter even if it is fully activated. Under equlibrium conditions, ion fluxes do take place, but the net influx and efflux of any given ion species are exactly equal.

For KCCs, the equilibrium is defined by  $[K^+]_0[CI^-]_0 =$ [K<sup>+</sup>]<sub>i</sub>[Cl<sup>-</sup>]<sub>i</sub>. This simple equation has important consequences. First, it implies that  $E_{K} = E_{CI}$ , which means that  $E_{GABA-A}$  and  $E_{GABA-B}$  would be equal under KCC equilibrium if  $GABA_ARs$ and the channels activated by GABABRs were ideally selective for Cl<sup>-</sup> and K<sup>+</sup>, respectively. Second, it shows that when KCCs are active,  $[Cl]_i$  is highly sensitive to changes in  $[K^+]_0$ , and a modest increase in [K<sup>+</sup>]<sub>o</sub> will lead to net uptake of Cl<sup>-</sup> by KCC2. Finally, it also implies that the driving force for extrusion of CI by KCCs is enhanced during an elevation of [CI], and therefore KCCs may also contribute to extracellular K+ transients (Voipio and Kaila, 2000). While it is worth re-emphasizing that thermodynamic considerations as such are not sufficient to make predictions about changes in the magnitude of ion transport in response to an increase in the deviation from equilibrium (i.e., the driving force acting on the transporter), we used the somewhat fuzzy term "thermodynamic regulation" (Payne et al., 2003) to define a situation where net fluxes mediated by a close-to-equilibrium transporter such as KCC2 are directly affected by the driving forces of the substrate ions. However, even KCC2 appears not to be constitutively active under all conditions, and an increasing amount of evidence indicates that kinetic regulation plays a crucial role.

The equilibrium of NKCCs is defined by  $[Na^+]_o[K^+]_o[Cl^-]_o^2 =$  $[Na^+]_i[K^+]_i[Cl^-]_i^2$ . The second power of  $[Cl^-]_i$  looks intriguing, and at first sight one might think that slight changes in [Cl-]i might exert an effect on NKCC1 via "thermodynamic regulation." However, under physiological conditions, NKCC1 is not close to equilibrium (Russell, 2000), but rather it is controlled by kinetic regulatory mechanisms that include a [Cl<sup>-</sup>]<sub>i</sub>-sensitive step (see below). For instance, in mammalian DRG neurons, kinetic constraints inactivate CI<sup>-</sup> uptake by NKCC1 even in the presence of a substantial thermodynamic driving force (Rocha-Gonzalez et al., 2008).

#### The Plasmalemmal Na-K ATPase

It is immediately clear from the close functional relationship between the plasmalemmal Na-K ATPase and CCCs (see Figure 1) that any factor that has an effect on the Na-K ATPase will have an influence on the operation of CCCs. This means that endogenous ligands (e.g., endogenous ouabains and agrin) that regulate neuronal Na-K ATPase isoforms will also affect CCC functions (Hilgenberg et al., 2006). Moreover, manipulations that lead to altered KCC or NKCC function may in fact reflect a change in Na-K ATPase function. For instance, when examining changes in CCC function following neuronal stimulation, the tight coupling between Na-K ATPases, CCCs, and  $E_{GABA-A}$  should be kept in mind. Interestingly, structural interactions have been demonstrated between the Na-K ATPase and KCC2 via coimmunoprecipitation of the α2 subunit and KCC2 in synaptic membrane fractions (Ikeda et al., 2004). Furthermore, the brain-type creatine kinase, an enzyme involved in the



production of ATP, has been shown to activate KCC2 and KCC3 via a protein-protein interaction (Inoue et al., 2006; Salin-Cantegrel et al., 2008).

#### **Kinetic Regulation of CCCs**

As pointed out before, the ion-translocating capacity of CCCs cannot be directly monitored using electrophysiological techniques, and there are no data available on the transport capacity of any CCC at the molecular level. Furthermore, except for a few cases, published data on the kinetic activation or inactivation of a given neuronal CCC do not provide information on whether this reflects a change in the number of membrane-located transporters, a change in their intrinsic rates of ion transport, or both (Figure 2). In the following, "kinetics" and "kinetic regulation" are terms that lump together the membrane trafficking of the CCCs as well as the (putative) changes in their unitary rates of ion transport.

Phosphorylation mechanisms play a pivotal role in the kinetic regulation of CCCs. In pioneering studies, it was shown that phosphorylation activates NKCC1 (Lytle and Forbush, 1992). A fall in [CI], below a homeostatic "set point" promoted direct phosphorylation and activation of NKCC1, leading to an active uptake of CI<sup>-</sup> (Russell, 2000). In contrast, the conventional view is that phosphorylation inactivates KCCs (Payne et al., 2003). However, there is evidence that tyrosine phosphorylation leads to activation of KCC2 (Vale et al., 2005; Wake et al., 2007; but see Stein et al., 2004) and that KCC2 is also directly phosphorylated and thereby activated by the protein kinase C (PKC; Lee et al., 2007; Banke and Gegelashvili, 2008). The activation by PKC is probably due to increased surface stability of the phosphorylated protein (Lee et al., 2007). In addition to PKC, several other kinases can regulate CCCs (Delpire and Gagnon, 2008; Kahle et al., 2008b). The serine-threonine kinases Wnks and SPAK/OSR1 are of particular interest in that they seem to be coregulators of NKCCs and KCCs. Studies with NKCCs in nonneuronal cells suggest that chloride-sensitive Wnks phosphorylate and activate the SPAK/OSR1, which in turn phosphorylates and activates NKCCs. While KCC2a and KCC3a contain a SPAK-binding motif, KCC2b and KCC3b do not, and the biological significance of this difference remains unknown. Most of the work above has been done in nonneuronal cells, and for neurobiologists, a crucial question is which of these mechanisms are relevant for CNS function.

## CCCs and the Polarity of $GABA_A$ Receptor-Mediated Responses

#### Chloride Extrusion and Hyperpolarizing GABA Responses

The pioneering work on the role of K-Cl cotransport in conventional hyperpolarizing GABAergic transmission was done in crayfish preparations (Aickin et al., 1982) and mammalian cortical neurons (Misgeld et al., 1986; Thompson and Gähwiler, 1989). In neurons,  $E_K$  is more negative than the resting  $V_m$ , which allows K-Cl cotransport to support conventional hyperpolarizing IPSPs (recall that when KCC is close to equilibrium,  $E_{Cl}$  is close to  $E_K$ ). However, the HCO $_3^-$  permeability of GABA $_A$ Rs implies that  $E_{GABA-A} > E_{Cl}$ , and hence active Cl $^-$  extrusion is a *necessary* but not a *sufficient* condition for hyperpolarizing GABA $_A$  responses (Kaila et al., 1993; Kaila, 1994; Rivera et al., 1999).

In most central neurons studied so far, the main chloride extruder is KCC2, but in some neurons KCC3 appears to have a significant role in setting  $E_{\rm GABA-A}$  (Boettger et al., 2003).

#### **Chloride Uptake and Depolarizing GABA Responses**

NKCC1 appears to be the main Cl<sup>-</sup> uptake mechanism responsible for GABA-mediated depolarizing currents mediated by postsynaptic and extrasynaptic receptors in developing hippocampal and neocortical neurons (Achilles et al., 2007; Sipilä et al., 2006; Yamada et al., 2004). A prominent exception from the conventional hyperpolarizing or slightly depolarizing responses evoked by GABA in mature cortical neurons was reported by Szabadics et al. (2006; but see Glickfeld et al., 2009). They showed that the GABA-mediated postsynaptic potentials evoked by chandelier cells (a class of interneurons that exclusively target the axon initial segment of principal neurons) had a depolarizing driving force of 15 mV. The underlying Cl<sup>-</sup> uptake mechanism was identified as NKCC1 in subsequent work, which showed that the GABA-mediated depolarization in the axon initial segment was abolished by bumetanide and in NKCC1 knockout mice (Khirug et al., 2008). The role of NKCC1 in neuronal Cl<sup>-</sup> uptake is not restricted to cortical structures. For instance, in olfactory receptor neurons, a drastic reduction in the driving force of the depolarizing odor-activated Cl current occurs in the presence of bumetanide and in NKCC1 knockout mice (Reisert et al., 2005; see also Kaneko et al., 2004). In contrast, the bumetanide-insensitive transporter(s) accounting for depolarizing GABA and glycine actions in the immature auditory brainstem and retinal ganglion and amacrine cells have not been identified (Balakrishnan et al., 2003; Zhang et al., 2007).

Depolarizing actions by GABA and glycine receptors have also been observed in presynaptic terminals. A classical example is the primary afferent depolarization (PAD) in DRG neurons (Rudomin and Schmidt, 1999; Russell, 2000), which seems to be largely but not fully attributable to Cl uptake mediated by NKCC1 (Rocha-Gonzalez et al., 2008). However, all of the available data are based on measurements carried out in the cell body, which may not be representative of other cellular compartments, and it will be critical to obtain an estimate of [Cl<sup>-</sup>]<sub>i</sub> in the central terminals of primary afferent fibers. Within the CNS, Cl uptake has been shown to produce depolarizing presynaptic GABA or glycine responses with a consequent increase in glutamate release in the ventromedial hypothalamus (Jang et al., 2001), the calyx of Held (Price and Trussell, 2006), terminals on isolated hippocampal CA3 neurons (Jang et al., 2006), and parallel fibers in the cerebellum (Stell et al., 2007). In these studies, presynaptic NKCC1-mediated Cl<sup>-</sup> accumulation was evident in the hypothalamus, but not in the calyx of

#### Intraneuronal CF Gradients

NKCC1 and KCCs are coexpressed in certain types of mature neurons, and spatially distinct expression patterns can result in steady-state chloride gradients and compartmentalization of  $E_{\rm GABA-A}$  within an individual neuron (e.g., Duebel et al., 2006; Gavrikov et al., 2003; Khirug et al., 2008; Price and Trussell, 2006; Szabadics et al., 2006). The steady-state  $E_{\rm GABA-A}$  gradients can be surprisingly large; in cortical principal neurons, the axosomato-dendritic CI<sup>-</sup> gradient attains a value up to 15–20 mV (Khirug et al., 2008; Szabadics et al., 2006). A similar gradient



is seen between the presynaptic terminals and the soma of globular bushy cells in the auditory brainstem (Price and Trussell, 2006). Given that distinct GABAergic interneurons in various brain structures target anatomically distinct sites in the postsynaptic neurons (Freund and Buzsaki, 1996), it is not correct to assign a singular  $E_{\text{GABA-A}}$  value to a given neuron—a more appropriate approach is to specify the  $E_{\text{GABA-A}}$  level of a given GABAergic input in the postsynaptic neuron.

#### Assessing the Efficacy of Neuronal Cl<sup>-</sup> Extrusion

The gramicidin-perforated patch clamp technique is often thought to be an ideal electrophysiological technique to study the presence and efficacy of neuronal Cl<sup>-</sup> extrusion. Because the gramicidin-induced pores are not permeable to CI-, there is no flux of Cl<sup>-</sup> between the recording pipette and the intracellular compartment (Kyrozis and Reichling, 1995). A novel approach where the cell membrane is kept intact relies on measuring  $E_{GABA-A}$  on the basis of the reversal potentials of single  $GABA_A$ and NMDA receptor channels (Tyzio et al., 2006, 2008). However, even if an ideal technique were available, measuring  $E_{GABA-A}$  in a resting neuron can at best verify the presence of Cl extrusion, but this provides no information on the efficacy of Cl<sup>-</sup> transport. Even a very inefficient Cl<sup>-</sup> extrusion mechanism will be able to maintain a hyperpolarizing E<sub>GABA-A</sub> if the net Cl<sup>-</sup> influx into a neuron (i.e., the cellular Cl<sup>-</sup> load) is small. Momentary changes in a target neuron's ionic load are among the most important context-dependent variables that influence GABAergic signaling (Buzsaki et al., 2007; Farrant and Kaila, 2007). Hence, in order to gain information that is physiologically more relevant than an estimate of the steady-state  $E_{GABA-A}$  in a resting neuron, it is important to design experiments to enable an assessment of the efficacy of Cl extrusion. Quantitative assessments of the efficacy of Cl extrusion are based on a procedure where a defined Cl<sup>-</sup> load is imposed on a cell, and the cell's capability to maintain the [Cl<sup>-</sup>]<sub>i</sub> provides a valid estimate of the efficacy of extrusion (e.g., Jarolimek et al., 1999; Khirug et al., 2005; Rivera et al., 2002). Ion-loading techniques have been routinely used in the study of neuronal pH regulation (Chesler, 2003), but the resting steady-state value of intraneuronal pH does not provide an idea of the efficacy of acid extrusion. By analogy, E<sub>GABA-A</sub> is not a reliable indicator of a neuron's capacity to regulate its [Cl], as was nicely illustrated in a recent study on chronically injured cortex, which showed that a marked decrease in KCC2 function was not associated with a detectable change in  $E_{GABA-A}$  (Jin et al., 2005).

### The "Developmental Shift" in GABA<sub>A</sub> Receptor-Mediated Signaling

It is frequently stated in both reviews and primary research papers that "during the neonatal period," KCC2 is upregulated, leading to a negative shift in  $E_{\rm GABA-A}$  from depolarizing to hyperpolarizing. This is a misleading generalization from work on rodent hippocampal and cortical pyramidal neurons, and there are pronounced *species-specific differences* in the temporal expression patterns of KCC2 within a given neuronal population. At the time of birth, "altricial" species such as rats show delayed KCC2 expression in the hippocampus and neocortex, while in the guinea pig (a species with precocious neonates) KCC2 expression has already reached a high level (Rivera et al.,

1999). This is also true for the perinatal human hippocampus and neocortex (Vanhatalo et al., 2005; but see Dzhala et al., 2005), and a very early onset of KCC2 expression is seen in the cortical subplate (Bayatti et al., 2008). On the other hand, within a single species, high levels of KCC2 expression are first seen in the spinal cord and subcortical brain regions, and thereafter in the cortex (Li et al., 2002; Rivera et al., 1999; Vinay and Jean-Xavier, 2008; see also Blaesse et al., 2006). The novel data on KCC2 isoforms (Uvarov et al., 2007) indicate that KCC2b is responsible for the developmental increase in neuronal CI<sup>-</sup> extrusion, at least in the cortex and in the auditory brainstem (Balakrishnan et al., 2003; Zhu et al., 2005). Interestingly, the shift from GABA<sub>A</sub> receptor depolarization to hyperpolarization of gonadotropin-releasing hormone neurons is delayed until the time of puberty (Clarkson and Herbison, 2006).

A relatively high  $[C\Gamma]_i$  is the "default" state of a wide variety of fully differentiated nonneuronal mammalian cells (Pedersen et al., 2006; Russell, 2000). Neurobiologists tend to regard immature neurons with their high  $[C\Gamma]_i$  and rather depolarized  $E_{\text{GABA-A}}$  as something exceptional while, in fact, it is the mature central neurons with their low  $[C\Gamma]_i$  that are the "aberrant" ones.  $[C\Gamma]_i$  in immature CNS neurons is typically in the range of 25–40 mM (e.g., Achilles et al., 2007; Balakrishnan et al., 2003; Kakazu et al., 1999; Kilb et al., 2002; Rohrbough and Spitzer, 1996; Sipilä et al., 2006; Yamada et al., 2004) and values typically of  $\sim$ 5 mM are seen in neuronal somata or dendrites after maturation (e.g., Khirug et al., 2008; Tyzio et al., 2008).

#### Inhibitory and Excitatory Actions of GABA

Unfortunately, the current literature on the developmental shift of E<sub>GABA-A</sub> and its consequences is burdened by a major misconception: it is often assumed that depolarizing GABA actions imply excitation and that a necessary and sufficient condition for a genuinely inhibitory GABA action is an  $E_{\text{GABA-A}}$  that is more negative than resting V<sub>m</sub>. This idea is a misleading simplification, which emphasizes the role of voltage inhibition (the hyperpolarizing response associated to an inhibitory postsynaptic potential) and ignores the crucial role played by shunting inhibition (based on the conductance of postsynaptic GABAARs). Shunting inhibition acts by decreasing the temporal and spatial summation of excitatory inputs. Hence, several points should be kept in mind when considering the relationships between the steady-state values of  $E_{GABA-A}$  and the efficacy of inhibition. The opening of GABAARs will have a shunting inhibitory action regardless of the value of  $E_{GABA-A}$ . Moderately depolarizing GABA actions can be functionally inhibitory and sometimes even more effective than hyperpolarizing responses because of the intrinsic outward rectification of GABAA currents and because depolarization leads to the inactivation of Na<sup>+</sup> channels and activation of K<sup>+</sup> channels. Finally, GABAergic transmission is not necessarily excitatory even if the postsynaptic response would be more positive than spiking threshold. This is because the spike voltage threshold is not a fixed parameter, but depends on the rate of change of the membrane potential, and also on the background conductance (e.g., Katz, 1966).

In conclusion, the steady-state levels of  $E_{\rm GABA-A}$  do not give a reliable estimate of the physiologically relevant efficacy of postsynaptic inhibition. This does not, of course, imply that  ${\rm Cl}^-$  transport is irrelevant here. It is important to recognize that



neurons in vivo undergo continuous fluctuations in their membrane potential, and fast changes take place in the driving force of the GABAAR current as well as in the momentary Cl load a neuron experiences. Hence, functional inhibition is largely determined by the efficacy of Cl extrusion, which has to be high enough to maintain a level of  $E_{\rm GABA-A}$  sufficiently negative (but not necessarily hyperpolarizing) to prevent the cell from firing.

In view of the large number of variables described above, it is not surprising that GABA can have "dual" actions in developing neurons (Khalilov et al., 1999; Lamsa et al., 2000). The duration of the depolarizing postsynaptic potential outlasts the local conductance increase, i.e., the "shunt." Hence, depolarizing GABAergic responses can facilitate the triggering of action potentials by glutamatergic neurons (Gao et al., 1998). This mode of action is consistent with a facilitatory or "permissive" role for depolarizing GABAergic transmission in the generation of early network events (see Sipilä et al., 2005; Zheng et al., 2006; and below). In an elegant study on neonatal rodent motoneurons, Jean-Xavier et al. (2007) first showed that depolarizing IPSPs (dIPSPs) were able to facilitate spike triggering by subthreshold excitatory events in their late phase. Notably, the window of facilitation became wider as  $E_{CI}$  was more depolarized, and started earlier when the site of dIPSP generation was moved away from the excitatory input. Therefore, in addition to  $E_{\text{GABA-A}}/E_{\text{Gly}}$ , the neuronal and synaptic architecture as well as the temporal relationships among GABA/glycinergic and glutamatergic postsynaptic events are critical factors that control the qualitative postsynaptic effects of depolarizing GABA/ glycinergic transmission. In summary, the widely spread idea that a dichotomous "depolarizing-to-hyperpolarizing switch" would control the inhibitory or excitatory nature of GABA/ glycinergic transmission in developing circuits is a profound oversimplification.

#### The Developmental Shift in EGABA-A

A large shift of  $E_{\rm GABA-A}$  from depolarizing to more negative and often hyperpolarizing values appears to be a ubiquitous feature of central neurons. A pioneering electrophysiological study on this topic was carried out by Obata et al. (1978) in chick spinal motor neurons, and the first observations related to the molecular mechanisms of the shift were made by Rivera et al. (1999), who showed that upregulation of KCC2 renders GABA hyperpolarizing in rat CA1 hippocampal neurons. The key observation was an increase in KCC2 expression during the first 2 postnatal weeks, a time window that was known to be associated with a negative shift in  $E_{\rm GABA-A}$ . Because there are no selective drugs to block CCCs in an isoform-specific manner, gene knockdown experiments were done to confirm the role of KCC2.

In various regions of the brain, the appearance of hyperpolarizing GABA<sub>A</sub>R signaling occurs earlier in the female and has been attributed to higher KCC2 and/or lower NKCC1 activity (Galanopoulou, 2008; Perrot-Sinal et al., 2007). The sex-specific GABAergic responses appear to promote sexual differentiation of the brain (Auger et al., 2001; Galanopoulou, 2008). In a recent study on rat pup hippocampi, Tyzio et al. (2006) reported that maternal oxytocin induces a perinatal hyperpolarizing shift in  $E_{\rm GABA-A}$  to astonishingly negative values (–100 mV and beyond). The authors suggested that oxytocin blocks NKCC1, thereby protecting the newborn brain from anoxic-ischemic damage,

a condition that is a major cause of neurological dysfunctions in humans. However, the implications of these findings for the human condition call for further studies.

## Neuronal Maturation and Emerging Networks Depolarizing GABA and the Generation of Endogenous Activity in Developing Networks

There is a wide consensus that endogenous activity generated prior to the maturation of sensory inputs plays a crucial role in the development of neuronal networks (Ben-Ari et al., 2007; Katz and Shatz, 1996; O'Donovan, 1999; Spitzer, 2006). According to a rather widespread and simplistic view, this kind of early activity is directly driven by depolarizing GABAergic transmission, and the "depolarization-hyperpolarization switch" puts an end to it. Notably, Minlebaev et al. (2007) have shown that in the neonatal neocortex in vivo, blocking GABAergic transmission has little effect on the basic patterns of the generation of spontaneous events known as "spindle bursts" (Khazipov et al., 2004). In contrast, the dominating pattern of neocortical spontaneous activity in vitro, which is reminiscent of hippocampal sharp potentials, is sensitive to GABAA receptor antagonists (Rheims et al., 2008). This type of activity, originally termed "giant depolarizing potentials" (GDPs), has been studied in detail in the developing hippocampus, and there are two hypotheses regarding the roles of depolarizing GABA in their generation. In the scenario proposed by Ben-Ari and coworkers (Ben-Ari et al., 2007), GABAergic interneurons are mainly responsible for pacing GDPs in a phasic manner (Ben-Ari et al., 1989). While all hippocampal regions are able to produce GDP-like activity, the CA3 region shows the highest propensity for GDP generation. Concerning the role of depolarizing GABA, a more recent view is that the GDPs are primarily paced by the CA3 glutamatergic pyramidal neurons, and depolarizing GABAergic signaling (both synaptic and tonic) has a facilitatory role in their generation (Sipilä et al., 2005). In other words, depolarizing GABAergic transmission is temporally nonpatterned and has a permissive or facilitatory role, while the intrinsic bursting properties of the CA3 glutamatergic neurons are responsible for the temporal patterning of GDPs. An analogous mechanism has recently been proposed for starburst cells in the developing retina (Zheng et al., 2006). Notably, GDPs are blocked by burnetanide both in vitro and in vivo, pointing to a crucial role for NKCC1 (Dzhala et al., 2005; Sipilä et al., 2006).

### Crosstalk among Neurotrophic Mechanisms, CCCs, and GABAergic Signaling

The concerted actions of CCCs and GABAergic signaling in brain development comprise an extremely wide spectrum of phenomena, and these will of course manifest themselves in neuronal network function and behavior. In fact, GABA's role in the formation of neuronal connectivity starts already during the early stages of neurogenesis (Owens and Kriegstein, 2002). Moreover, GABAergic transmission plays a key role during "critical" periods, when the nervous system is particularly prone to both normal and aberrant types of input (Hensch, 2005; Kanold and Shatz, 2006).

GABA<sub>A</sub>Rs are functional in neuronal precursors and immature neurons, long before the formation of GABAergic synapses (Demarque et al., 2002; LoTurco et al., 1995; Owens et al., 1999).



During neuronal development, depolarizing extrasynaptic and synaptic GABAergic signaling promotes action potential activity, opening of voltage-gated Ca2+ channels, and activation of NMDA receptors (Ben-Ari et al., 2007; Fukuda et al., 1998; Hennou et al., 2002; Kandler and Gillespie, 2005; Serafini et al., 1995). These responses lead to transient elevations of intracellular Ca2+ levels and activation of downstream intracellular signaling cascades, which are central in mediating the trophic effects of GABA during development (Ben-Ari et al., 2007; Owens and Kriegstein, 2002). Trophic effects of GABA have been observed in vitro at various levels of neuronal and network development, including DNA synthesis, migration, morphological maturation of individual neurons, and synaptogenesis (Akerman and Cline, 2006; Liu et al., 2005, 2006; LoTurco et al., 1995; Marty et al., 1996, 2000; Owens and Kriegstein, 2002; Represa and Ben Ari, 2005; Reynolds et al., 2008; Simat et al., 2007; Wang and Kriegstein, 2008; see also Andang et al., 2008). Interestingly, the developmental patterns of GABAergic signaling are repeated during adult neurogenesis in the dentate gyrus in vivo (Esposito et al., 2005; Ge et al., 2006; Laplagne et al., 2007; Tozuka et al., 2005; see also Toni et al., 2008). BDNF has been ascribed a key role in a number of the studies cited above. It is a well-known fact that the release of trophic factors is enhanced during synchronous neuronal activity (Lessmann et al., 2003), and one of the functions of the early spontaneous network events described above may be to promote the release of these substances.

Despite the above findings related to depolarizing GABA actions, their significance in normal neuronal development in vivo is not clear (Wang and Kriegstein, 2009). Somewhat surprisingly, synaptogenesis and early brain development are hardly affected in knockout mice where GABA synthesis, vesicular transport, or vesicular release are eliminated (Jin et al., 1999; Varoqueaux et al., 2002; Verhage et al., 2000; Wojcik et al., 2006). On the other hand, using in utero electroporation to overexpress KCC2 in proliferating neurons, Cancedda et al. (2007) found that while neuronal migration was not affected (but see Heck et al., 2007), morphological maturation was markedly impaired in immature neurons that were devoid of depolarizing GABAergic responses.

An intriguing hypothesis suggests that presynaptically released GABA acts as a self-limiting trophic factor in a negative feedback loop where depolarizing postsynaptic GABAAR responses are required for the developmental upregulation of KCC2 (Ganguly et al., 2001; see also Leitch et al., 2005). However, subsequent studies showed that developmental KCC2 upregulation can take place in hippocampal and midbrain cultures in the complete absence of GABAAR signaling (Ludwig et al., 2003; Titz et al., 2003). Furthermore, during embryonic development in VIAAT knockout mice, KCC2 mRNA and protein levels are unaffected, despite the absence of GABAergic synaptic transmission (Wojcik et al., 2006).

#### Role of CCCs in the Formation of Synapses

Depolarizing actions of GABA play at least a permissive role in excitatory and inhibitory synaptogenesis (Akerman and Cline, 2007). In the Xenopus tadpole visual system in vivo, early overexpression of KCC2 led to a reduction in neuronal [Cl7], levels, interfering with the maturation of the glutamatergic system and enhancing the development of GABAergic synapses (Akerman and Cline, 2006). Overexpression of KCC2 in early hippocampal cultures selectively increased the number of GABAergic synapses and the frequency of GABAergic miniature postsynaptic currents (Chudotvorova et al., 2005). Consistent with the above work, Riekki et al. (2008) found that miniature IPSC frequency is lower in mice with reduced KCC2 levels (Tornberg et al., 2005).

However, the various roles of CCCs in neuronal development and connectivity may cover a much wider scope than what is caused by their actions on GABAergic and glycinergic responses. Recently, Li et al. (2007) obtained evidence that KCC2 may play a morphogenic role in the formation of excitatory synapses during cortical development. Cultured KCC2-/neurons were found to have long, morphologically aberrant "filopodia-like" spines, but this phenotype could be rescued by transfecting the neurons with an N-terminal deletion construct that is not capable of mediating K-Cl cotransport. This suggested that the role of KCC2 in spine maturation is not based on its transport activity, but might be mediated by interactions with cytoskeletal proteins. KCC2 immunoprecipitation assays from adult mouse brain homogenate suggest an interaction between the KCC2 C-terminal domain and the 4.1N protein, which is known to mediate structural interactions between the cytoskeleton, transmembrane proteins, and adhesion molecules (Denker and Barber, 2002). While it remains to be verified whether KCC2 has a role in spine formation in vivo, neurons from mice with reduced KCC2 protein levels show an increase in the length of dendritic protrusions (Tornberg et al., 2005). Thus, KCC2 may function as a synchronizing factor in the development of inhibitory and excitatory neurotransmission. In this scenario, the progressive developmental increase in the number of functional glutamatergic synapses that target spines will be paralleled by an increase in the KCC2-dependent neuronal CIT extrusion capacity, and consequently, by an increase in the efficacy of GABAergic inhibition (Li et al., 2007).

#### **Neuronal Plasticity**

Postsynaptic GABA<sub>A</sub> signaling has the unique property of "ionic plasticity," which is based on short- and long-term changes in the transmembrane anion gradients. A robust manifestation of short-term GABAergic plasticity is apparent in CA1 pyramidal neurons following high-frequency stimulation of Shaffer collaterals, a classical experimental paradigm to evoke long-term potentiation. Under these conditions, the membrane potential of CA1 pyramidal neurons shows a biphasic response, with a prompt hyperpolarization followed by a large and prolonged depolarization. The depolarizing phase of this GABAergic response is attributable to two mechanisms: an initial and fast positive shift in  $E_{\text{GABA-A}}$ , caused by a bicarbonate-dependent uptake of Cl<sup>-</sup> (see Kaila and Voipio, 1987) that partly overlaps with and is followed by an increase in [K<sup>+</sup>]<sub>o</sub>, which has a direct depolarizing action on the neuronal membrane potential (Fiumelli and Woodin, 2007; Kaila et al., 1997; Voipio and Kaila, 2000). Hence, pronounced short-term changes that affect the efficacy and sometimes even the qualitative (inhibitory versus excitatory) action of GABAergic transmission can take place in an activitydependent manner without the involvement of changes in the kinetic modulation, trafficking and/or expression of CCCs.



There is evidence that KCC2 has an extremely high rate of turnover (recycling) at the plasmalemma (Lee et al., 2007), which suggests that the transporter is subject to continuous kinetic modulation. Recent work has shown that coincident firing of pre- and postsynaptic neurons or prolonged postsynaptic spiking alone can result in positive shifts in  $E_{\mathsf{GABA-A}}$  (Woodin et al., 2003; Fiumelli et al., 2005). These shifts are dependent on intracellular Ca2+, and a PKC-dependent mechanism was suggested to mediate the functional downregulation of KCC2 (Fiumelli et al., 2005). This conclusion contradicts Lee et al. (2007), who reported that PKC inhibits the endocytosis of KCC2, resulting in functional upregulation of KCC2, and it has been recently suggested (Brumback and Staley, 2008) that changes in Na-K ATPase activity are responsible for the activity-induced changes in E<sub>GABA-A</sub> that were observed by Fiumelli and coworkers.

In hippocampal slices exposed to a Mg<sup>2+</sup>-free solution, the ensuing interictal activity was shown to downregulate KCC2 mRNA and protein in a BDNF-dependent manner, and it was concluded that this pathophysiological plasticity is attributable to transcriptional changes in KCC2 expression levels (Rivera et al., 2004). However, the authors could not exclude the possibility that the decreased efficacy of Cl<sup>-</sup> extrusion was caused by changes in posttranscriptional mechanisms. Indeed, there is evidence that BDNF can act on KCC2 function within a time window that is too brief to be mediated by transcriptional effects (Wardle and Poo, 2003). Hence, the overall picture looks rather similar to what is known about the plasticity of glutamatergic transmission: fast changes are mediated by posttranscriptional mechanisms, but at the same time changes in gene expression take place which appear to "consolidate" the initial effects.

#### Disease Mechanisms Mutations of CCCs

While a large number of genetic defects in ion channels (channelopathies) are known to produce severe dysfunctions of neuronal signaling in diseases such as epilepsy, migraine, and pain (Mulley et al., 2003; Waxman, 2007), little data are available for ion transporters. However, a striking example of a genetic "transporteropathy" is provided by hereditary motor and sensory neuropathy with agenesis of the corpus callosum (HMSN/ACC; Salin-Cantegrel et al., 2008), an autosomal recessive disease characterized by progressive sensory neuropathy, mental retardation, and dysmorphic features such as a high arched palate and agenesis of the corpus callosum. All of the four identified mutations that underlie this disease involve C-terminal deletion of KCC3, which renders the transporter nonfunctional (Howard et al., 2002). The mutations disrupt activation of KCC3 by the brain-type creatine kinase, which suggests the first proteinprotein interaction involved in HMSN/ACC pathophysiology (Salin-Cantegrel et al., 2008). KCC3 knockout mice have a phenotype that is largely consistent with the human disease (Boettger et al., 2003; Delpire and Mount, 2002; Howard et al., 2002).

A substantial fraction of idiopathic epilepsies are thought to have a polygenic basis (Mulley et al., 2005). Here, it is of interest that *kazachoc* mutations of *Drosophila kcc* that reduce the level of kcc protein are susceptible to epileptic-like seizures (Hekmat-

Scafe et al., 2006). However, no mutations or polymorphisms in human CCC genes related to seizure susceptibility have been published so far.

#### Neuronal Trauma

The brain is highly susceptible to a wide spectrum of acute and chronic pathophysiological conditions, including anoxicischemic states, mechanical damage, and epileptic activity. A depolarizing shift in  $E_{GABA-A}$  in response to neuronal trauma has been observed following anoxia/ischemia, neurite transection, and osmotic imbalance (Katchman et al., 1994; Pond et al., 2006; van den Pol et al., 1996). The trauma-induced effects on E<sub>GABA-A</sub> are largely attributable to a downregulation of KCC2 with a consequent increase in [Cl<sup>-</sup>]<sub>i</sub> (Bonislawski et al., 2007; Fukuda et al., 1998; Nabekura et al., 2002; Papp et al., 2008), and some data point to a concurrent functional upregulation of NKCC1 (Pond et al., 2006; Yan et al., 2001b). There is no doubt that fast changes in CCC gene expression can take place in response to neuronal trauma (De Koninck, 2007; Rivera et al., 2002; Shulga et al., 2008), but the precise roles of transcriptional, posttranscriptional, and posttranslational mechanisms in the pathophysiological effects mediated by CCCs on [Cl7], and  $E_{\text{GABA-A}}$  remain to be identified (see Figure 2).

#### **Epilepsy**

While the large majority ( $\sim$ 70%) of epilepsies is idiopathic, the remaining cases are largely attributable to cerebrovascular diseases, traumatic brain injury, and tumors. A downregulation of KCC2 associated with epileptiform activity has been observed in a number of in vivo and in vitro studies in animal models (Jin et al., 2005; Pathak et al., 2007; Rivera et al., 2002, 2004), and this mechanism may act as one of the links between neuronal injury and epilepsy (Pathak et al., 2007). In a pivotal study, Miles and coworkers showed that hippocampal slices obtained from temporal lobe epilepsy patients generated interictal activity that was attributable to depolarizing GABAA-mediated transmission in a subpopulation of principal neurons (Cohen et al., 2002; see also Kohling et al., 1998). A follow-up study showed that the depolarization was related to a downregulation of KCC2 and that, interestingly, the interictal activity was blocked by burnetanide (Huberfeld et al., 2007; see also Munoz et al., 2007). The molecular cascades that lead to KCC2 downregulation in various pathophysiological states include those mediated by BDNF-TrkB signaling (Rivera et al., 2004).

GABA<sub>A</sub>-mediated transmission is not suppressed in epileptic tissue (Avoli et al., 2005; Kohling et al., 1998), but the Cl<sup>-</sup> extrusion capacity seems to play a key role in setting the susceptibility of neurons to epileptiform activity. In transgenic mice, the propensity to epilepsy in vivo and hyperexcitability in vitro are enhanced with an inverse relationship to the level of KCC2 expression (Riekki et al., 2008; Tornberg et al., 2005; Woo et al., 2002; Zhu et al., 2005). GABAergic inhibition can also revert into excitation without any change in the functional expression of KCCs, as mentioned above (Kaila et al., 1997). Indeed, epileptic seizures are associated with large extracellular K<sup>+</sup> transients, which may have a substantial GABAergic component (Avoli et al., 2005; Kaila et al., 1997). A pathophysiological increase in [K<sup>+</sup>]<sub>o</sub> can lead to a vicious circle that involves a further depolarization of both the membrane potential and  $E_{\mathsf{GABA-A}}$  (Kakazu et al., 2000; Payne et al., 2003; Thompson and Gähwiler,



1989) and a cell swelling that enhances proepileptic ephaptic signaling (Haglund and Hochman, 2005).

#### **CCC Blockers as Putative Anticonvulsant Drugs**

A number of studies have examined the possible utility of diuretic drugs, especially furosemide and bumetanide, in the treatment of epileptic disorders. While some observations indicate an anticonvulsant action (e.g., Haglund and Hochman, 2005; Hochman et al., 1995), the interpretation is complicated by the wide diversity of the drugs' cellular targets within and outside the brain and, in the case of furosemide, because of its lack of specificity at the molecular level (Staley, 2002). The search for novel drugs is particularly important in the case of neonatal seizures, where standard anticonvulsants have poor efficacy (Rennie and Boylan, 2007). At the moment, bumetanide has attracted much attention as a putative drug to suppress seizures in human neonates (Dzhala et al., 2005, 2008; Kahle et al., 2008a), an idea that is largely based on the assumption that seizure susceptibility is linked to depolarizing actions of GABA in the neonate brain (e.g., Ben Ari and Holmes, 2006). However, the expression patterns of KCC2 and the parallel developmental characteristics of the human EEG do not suggest that GABA would be overtly depolarizing in fullterm newborn cortex (Vanhatalo et al., 2005), but the situation is probably different for preterm neonates (Dzhala et al., 2005; Vanhatalo et al., 2005). In addition, work in vitro has resulted in conflicting data regarding the anticonvulsant properties of bumetanide (Kilb et al., 2007). Clearly, much more research is needed to test the hypothesis of a connection from nonpathophysiological depolarizing actions of GABA to seizure generation in neonates and its possible clinical implications.

#### Neuropathic Pain

Neuropathic pain is a common and highly debilitating condition resulting from damage or dysfunction of peripheral nerves. This type of pain is often refractory to standard analgesic agents, and opioid treatment can result in unacceptable side effects. Symptomatic management of neuropathic pain by suppressing neuronal activity with antiepileptic drugs can also have undesirable side effects (Blackburn-Munro and Erichsen, 2005; Jensen, 2002; Rogawski and Loscher, 2004). Hence, neuropathic pain represents a major unmet medical need. GABA and glycinergic inhibition play a pivotal role via modulatation of the afferent input into nociceptive neurons in the superficial dorsal horn and control of their output to higher areas of the central nervous system (Price et al., 2005). Loss of inhibitory neurotransmission may underlie several forms of chronic pain. De Koninck and coworkers have proposed that suppression of inhibition by a collapse of the chloride gradient in dorsal horn lamina I neurons is a cellular substrate of neuropathic pain (Coull et al., 2003; De Koninck, 2007). They reported that peripheral nerve injury decreased KCC2 expression and produced a depolarizing shift in  $E_{\mathsf{GABA-A}}$  in a subpopulation of lamina I neurons in rats. This loss of inhibition appeared to be mediated via BDNF released from activated spinal microglia (Coull et al., 2005).

#### Pathogenic versus Protective Mechanisms

At first sight, downregulation of neuronal CI extrusion and a consequent decrease in the efficacy of inhibition appears to be maladaptive for the survival of diseased neuronal tissue. Here, one should note that the responses of neurons to trauma can be causally related to the pathophysiological sequelae per se, or to intrinsic protective mechanisms that promote neuronal survival and re-establishment of functional connectivity (Payne et al., 2003). It has been proposed that the posttraumatic downregulation of neuronal CI<sup>-</sup> extrusion and the consequent depolarizing shift in  $E_{\mathsf{GABA-A}}$  are aspects of dedifferentiation of neurons and neuronal networks that are permissive for the "recapitulation" of developmental programs for the rewiring of surviving neurons (Payne et al., 2003; Pieraut et al., 2007; Shulga et al., 2008). Furthermore, in view of the tight link between the Na-K ATPase and CCC functions, it is interesting that seizure activity and neuronal damage have been shown to lead to a downregulation of the sodium pump (e.g., Pylova et al., 1989; Ross and Soltesz, 2000), which may well be a neuroprotective response to minimize local oxygen and glucose demand during traumarelated hyperexcitability and plasticity.

#### **Conclusions and Topics for Future Work**

Neurons typically have exceedingly complex morphological properties, and the data reviewed above suggest that many have steady-state intracellular CIT gradients that are set by cell-type-specific, subcellular expression patterns of functional CCCs. An immediate question that arises here is the role of such gradients in neuronal signaling during ongoing activity (e.g., oscillations) in the brain. Spatial "sign changes" (inhibitory versus excitatory) in GABAergic signaling have been identified under steady-state conditions, for instance in mature neurons in the retina (Gavrikov et al., 2003). Hence, it is possible that the spectrum of context-dependent actions of GABA include analogous sign changes that take place within a timescale of perhaps tens of milliseconds, thereby adding a further degree of freedom to the computational properties of neurons. Testing this hypothesis will require fast and reliable monitoring of neuronal membrane potential in parallel with measurements of [Cl<sup>-</sup>]<sub>i</sub>, a task that can be achieved only after further developments of Cl<sup>-</sup> imaging techniques.

It is likely that in vivo at least some functional CCCs are homooligomers (Blaesse et al., 2006; Casula et al., 2001), and work in vitro shows that monomers of distinct CCC isoforms can combine to hetero-oligomers (Simard et al., 2007). But does hetero-oligomerization also take place in vivo? And if it does, what are the subunit compositions of native CCC hetero-oligomers? Information of this kind is physiologically important, and it will have immediate consequences on work aimed at designing drugs (such as antiepileptic or analgesic compounds) that target CCCs in the living organism. Obtaining insight into CCC molecular structure should help in solving these problems. Future studies on polymorphisms will add to the spectrum of both functional and dysfunctional CCCs, a topic that has much potential in the search for novel disease genes.

Current data also demonstrate that the CCCs are not only working as Cl<sup>-</sup> transporters. This is in line with a number of previous observations showing that plasmalemmal channels and ion transporters are often multifunctional and that they can exert actions at the level of cellular structure (Denker and Barber, 2002). The role that KCC2 has on spine formation may well be a facet in a much broader picture where CCCs act as morphogenic factors. Little is known about the anchoring of CCCs,



and future work is likely to uncover novel protein-protein interactions that influence the functional and structural roles of CCCs. In addition to the search for ion-regulatory metabolons incorporating CCCs, an intriguing question is whether "plasticity" of anchoring proteins is a determinant of subcellular CCC expression and functionality.

In addition to the electrophysiologically obvious link between CCCs and anion-permeable channels gated by GABA and glycine, CCCs are involved in intricate, bidirectional interactions with trophic factors as well as intracellular signaling cascades. However, we are still at a loss regarding the precise mechanisms whereby, and to what extent, trophic factors control CCC expression and functionality. Clearly, this kind of knowledge is needed to unravel the roles of CCCs in brain development, plasticity, and disease.

While research on the diverse functions on CCCs in the CNS is still in its infancy, the overall picture is exciting and holds promise for major breakthroughs in our understanding of the most fundamental mechanisms underlying the workings of the nervous system. The wide and rapidly increasing research investments employing a wide spectrum of powerful molecular, electrophysiological, and imaging techniques is likely to lead to a fast narrowing of the gap between the level of our knowledge of ion transporters and ion channels.

#### **ACKNOWLEDGMENTS**

The original research work by the authors was supported by grants from the Academy of Finland, from the Sigrid Jusélius Foundation, and from the European Integrated Project "EPICURE" EFP6-037315. M.S.A., C.R., and K.K. are members of the Finnish CoE of Molecular and Integrative Neuroscience, and K.K. is a member of the Nordic CoE "WIRED." P.B. is a postdoctoral fellow of the German Academic Exchange Service (DAAD). We thank Drs. Eric Delpire, Christian A. Hübner, John A. Payne, and Theodore J. Price for constructive comments on the manuscript.

#### **REFERENCES**

Achilles, K., Okabe, A., Ikeda, M., Shimizu-Okabe, C., Yamada, J., Fukuda, A., Luhmann, H.J., and Kilb, W. (2007). Kinetic properties of Cl uptake mediated by Na+-dependent K+-2Cl cotransport in immature rat neocortical neurons. J. Neurosci. 27, 8616-8627.

Aickin, C.C., Deisz, R.A., and Lux, H.D. (1982). Ammonium action on postsynaptic inhibition in crayfish neurones: implications for the mechanism of chloride extrusion. J. Physiol. 329, 319-339.

Akerman, C.J., and Cline, H.T. (2006). Depolarizing GABAergic conductances regulate the balance of excitation to inhibition in the developing retinotectal circuit in vivo. J. Neurosci. 26, 5117-5130.

Akerman, C.J., and Cline, H.T. (2007). Refining the roles of GABAergic signaling during neural circuit formation. Trends Neurosci. 30, 382-389.

Amiry-Moghaddam, M., and Ottersen, O.P. (2003). The molecular basis of water transport in the brain. Nat. Rev. Neurosci. 4, 991-1001.

Andang, M., Hjerling-Leffler, J., Moliner, A., Lundgren, T.K., Castelo-Branco, G., Nanou, E., Pozas, E., Bryja, V., Halliez, S., Nishimaru, H., et al. (2008). Histone H2AX-dependent GABA(A) receptor regulation of stem cell proliferation. Nature 451, 460-464.

Auger, A.P., Perrot-Sinal, T.S., and McCarthy, M.M. (2001). Excitatory versus inhibitory GABA as a divergence point in steroid-mediated sexual differentiation of the brain. Proc. Natl. Acad. Sci. USA 98, 8059-8064.

Avoli, M., Louvel, J., Pumain, R., and Kohling, R. (2005). Cellular and molecular mechanisms of epilepsy in the human brain. Prog. Neurobiol. 77, 166-200.

Balakrishnan, V., Becker, M., Löhrke, S., Nothwang, H.G., Güresir, E., and Friauf, E. (2003). Expression and function of chloride transporters during development of inhibitory neurotransmission in the auditory brainstem. J. Neurosci. 23, 4134-4145.

Banke, T.G., and Gegelashvili, G. (2008). Tonic activation of group I mGluRs modulates inhibitory synaptic strength by regulating KCC2 activity. J. Physiol.

Bartho, P., Payne, J.A., Freund, T.F., and Acsady, L. (2004). Differential distribution of the KCI cotransporter KCC2 in thalamic relay and reticular nuclei. Eur. J. Neurosci. 20, 965-975.

Bayatti, N., Moss, J.A., Sun, L., Ambrose, P., Ward, J.F.H., Lindsay, S., and Clowry, G.J. (2008). A molecular neuroanatomical study of the developing human neocortex from 8 to 17 postconceptional weeks revealing the early differentiation of the subplate and subventricular zone. Cereb. Cortex 18, 1536-1548.

Becker, M., Nothwang, H.G., and Friauf, E. (2003). Differential expression pattern of chloride transporters NCC, NKCC2, KCC1, KCC3, KCC4, and AE3 in the developing rat auditory brainstem. Cell Tissue Res. 312, 155-165.

Ben Ari, Y., and Holmes, G.L. (2006). Effects of seizures on developmental processes in the immature brain. Lancet Neurol. 5, 1055-1063.

Ben-Ari, Y., Cherubini, E., Corradetti, R., and Gaiarsa, J.L. (1989). Giant synaptic potentials in immature rat CA3 hippocampal neurons. J. Physiol.

Ben-Ari, Y., Gaiarsa, J.L., Tyzio, R., and Khazipov, R. (2007). GABA: a pioneer transmitter that excites immature neurons and generates primitive oscillations. Physiol. Rev. 87, 1215-1284.

Bergeron, M.J., Gagnon, E., Caron, L., and Isenring, P. (2006). Identification of key functional domains in the C terminus of the K+-Cl- cotransporters. J. Biol. Chem. 281, 15959-15969.

Berglund, K., Schleich, W., Krieger, P., Loo, L., Wang, D., Cant, N., Feng, G., Augustine, G., and Kuner, T. (2006). Imaging synaptic inhibition in transgenic mice expressing the chloride indicator, Clomeleon. Brain Cell Biol. 35, 207-

Blackburn-Munro, G., and Erichsen, H.K. (2005). Antiepileptics and the treatment of neuropathic pain: Evidence from animal models. Curr. Pharm. Des. 11,

Blaesse, P., Guillemin, I., Schindler, J., Schweizer, M., Delpire, E., Khiroug, L., Friauf, E., and Nothwang, H.G. (2006). Oligomerization of KCC2 correlates with development of inhibitory neurotransmission. J. Neurosci. 26, 10407-10419.

Boettger, T., Hübner, C.A., Maier, H., Rust, M.B., Beck, F.X., and Jentsch, T.J. (2002). Deafness and renal tubular acidosis in mice lacking the K-Cl co-transporter Kcc4. Nature 416, 874-878.

Boettger, T., Rust, M.B., Maier, H., Seidenbecher, T., Schweizer, M., Keating, D.J., Faulhaber, J., Ehmke, H., Pfeffer, C., Scheel, O., et al. (2003). Loss of K-Cl co-transporter KCC3 causes deafness, neurodegeneration and reduced seizure threshold. EMBO J. 22, 5422-5434.

Bonislawski, D.P., Schwarzbach, E.P., and Cohen, A.S. (2007). Brain injury impairs dentate gyrus inhibitory efficacy. Neurobiol. Dis. 25, 163-169.

Brumback, A.C., and Staley, K.J. (2008). Thermodynamic regulation of NKCC1-mediated CI- cotransport underlies plasticity of GABAA signaling in neonatal neurons. J. Neurosci. 28, 1301-1312.

Buzsaki, G., Kaila, K., and Raichle, M. (2007). Inhibition and brain work. Neuron

Byun, N., and Delpire, E. (2007). Axonal and periaxonal swelling precede peripheral neurodegeneration in KCC3 knockout mice. Neurobiol. Dis. 28,

Cancedda, L., Fiumelli, H., Chen, K., and Poo, M.M. (2007). Excitatory GABA action is essential for morphological maturation of cortical neurons in vivo. J. Neurosci. 27, 5224-5235.

Carmosino, M., Gimenez, I., Caplan, M., and Forbush, B. (2008). Exon loss accounts for differential sorting of Na-K-Cl cotransporters in polarized epithelial cells. Mol. Biol. Cell 19, 4341-4351.

### Neuron Review



Casula, S., Shmukler, B.E., Wilhelm, S., Stuart-Tilley, A.K., Su, W., Chernova, M.N., Brugnara, C., and Alper, S.L. (2001). A dominant negative mutant of the KCC1 K-Cl cotransporter: both N- and C-terminal cytoplasmic domains are required for K-Cl cotransport activity. J. Biol. Chem. 276, 41870-41878.

Chen, H., Luo, J., Kintner, D.B., Shull, G.E., and Sun, D. (2005). Na+-dependent chloride transporter (NKCC1)-null mice exhibit less gray and white matter damage after focal cerebral ischemia. J. Cereb. Blood Flow Metab. 25, 54-66.

Chesler, M. (2003). Regulation and modulation of pH in the brain. Physiol. Rev. 83, 1183-1221.

Chudotvorova, I., Ivanov, A., Rama, S., Hübner, C.A., Pellegrino, C., Ben Ari, Y., and Medina, I. (2005). Early expression of KCC2 in rat hippocampal cultures augments expression of functional GABA synapses. J. Physiol. 566, 671–679.

Clarkson, J., and Herbison, A.E. (2006). Development of GABA and glutamate signaling at the GnRH neuron in relation to puberty. Mol. Cell. Endocrinol. 254,

Clayton, G.H., Owens, G.C., Wolff, J.S., and Smith, R.L. (1998). Ontogeny of cation-Cl- cotransporter expression in rat neocortex. Brain Res. Dev. Brain Res. 109, 281-292.

Cohen, I., Navarro, V., Clemenceau, S., Baulac, M., and Miles, R. (2002). On the origin of interictal activity in human temporal lobe epilepsy in vitro. Science 298, 1418-1421.

Colmenero-Flores, J.M., Martinez, G., Gamba, G., Vazquez, N., Iglesias, D.J., Brumos, J., and Talon, M. (2007). Identification and functional characterization of cation-chloride cotransporters in plants. Plant J. 50, 278-292.

Coull, J.A., Boudreau, D., Bachand, K., Prescott, S.A., Nault, F., Sik, A., De Koninck, P., and De Koninck, Y. (2003). Trans-synaptic shift in anion gradient in spinal lamina I neurons as a mechanism of neuropathic pain. Nature 424,

Coull, J.A.M., Beggs, S., Boudreau, D., Boivin, D., Tsuda, M., Inoue, K., Gravel, C., Salter, M.W., and De Koninck, Y. (2005). BDNF from microglia causes the shift in neuronal anion gradient underlying neuropathic pain. Nature 438, 1017-

De Koninck, Y. (2007). Altered chloride homeostasis in neurological disorders: a new target. Curr. Opin. Pharmacol. 7, 93-99.

Delpire, E., and Mount, D.B. (2002). Human and murine phenotypes associated with defects in cation-chloride cotransport. Annu. Rev. Physiol. 64, 803-843.

Delpire, E., and Gagnon, K.B.E. (2008). SPAK and OSR1: STE20 kinases involved in the regulation of ion homoeostasis and volume control in mammalian cells. Biochem. J. 409, 321-331.

Delpire, E., Lu, J.M., England, R., Dull, C., and Thorne, T. (1999). Deafness and imbalance associated with inactivation of the secretory Na-K-2Cl co-transporter. Nat. Genet. 22, 192-195.

Delpire, E., Days, E., Lewis, M., Mi, D., Kim, K., Lindsley, C.W., and Weaver, C.D. (2009). Small molecule screen identifies inhibitors for the neuronal K-Cl cotransporter KCC2. Proc. Natl. Acad. Sci. USA, in press 10.1073/pnas. 0812756106.

Demarque, M., Represa, A., Becq, H., Khalilov, I., Ben Ari, Y., and Aniksztejn, L. (2002). Paracrine intercellular communication by a Ca2+- and SNARE-independent release of GABA and glutamate prior to synapse formation. Neuron 36, 1051-1061.

Denker, S.P., and Barber, D.L. (2002). Ion transport proteins anchor and regulate the cytoskeleton. Curr. Opin. Cell Biol. 14, 214-220.

Dixon, M.J., Gazzard, J., Chaudhry, S.S., Sampson, N., Schulte, B.A., and Steel, K.P. (1999). Mutation of the Na-K-Cl co-transporter gene Slc12a2 results in deafness in mice. Hum. Mol. Genet. 8, 1579-1584.

Duebel, J., Haverkamp, S., Schleich, W., Feng, G., Augustine, G.J., Kuner, T., and Euler, T. (2006). Two-photon imaging reveals somatodendritic chloride gradient in retinal ON-type bipolar cells expressing the biosensor clomeleon. Neuron 49, 81-94.

Dzhala, V.I., Talos, D.M., Sdrulla, D.A., Brumback, A.C., Mathews, G.C., Benke, T.A., Delpire, E., Jensen, F.E., and Staley, K.J. (2005). NKCC1 transporter facilitates seizures in the developing brain. Nat. Med. 11, 1205-1213.

Dzhala, V.I., Brumback, A.C., and Staley, K.J. (2008). Bumetanide enhances phenobarbital efficacy in a neonatal seizure model. Ann. Neurol. 63, 222-235.

Esposito, M.S., Piatti, V.C., Laplagne, D.A., Morgenstern, N.A., Ferrari, C.C., Pitossi, F.J., and Schinder, A.F. (2005). Neuronal differentiation in the adult hippocampus recapitulates embryonic development. J. Neurosci. 25, 10074-10086.

Farrant, M., and Kaila, K. (2007). The cellular, molecular and ionic basis of GABAA receptor signalling. Prog. Brain Res. 160, 59-87.

Fiumelli, H., and Woodin, M.A. (2007). Role of activity-dependent regulation of neuronal chloride homeostasis in development. Curr. Opin. Neurol. 17, 81-86.

Fiumelli, H., Cancedda, L., and Poo, M.M. (2005). Modulation of GABAerqic transmission by activity via postsynaptic Ca2+-dependent regulation of KCC2 function. Neuron 48, 773-786.

Flagella, M., Clarke, L.L., Miller, M.L., Erway, L.C., Giannella, R.A., Andringa, A., Gawenis, L.R., Kramer, J., Duffy, J.J., Doetschman, T., et al. (1999). Mice lacking the basolateral Na-K-2Cl cotransporter have impaired epithelial chloride secretion and are profoundly deaf. J. Biol. Chem. 274, 26946-26955.

Flatman, P.W. (2008). Cotransporters, WNKs and hypertension: an update. Curr. Opin. Nephrol. Hypertens. 17, 186-192.

Flemmer, A.W., Gimenez, I., Dowd, B.F.X., Darman, R.B., and Forbush, B. (2002). Activation of the Na-K-Cl cotransporter NKCC1 detected with a phospho-specific antibody. J. Biol. Chem. 277, 37551-37558.

Freund, T.F., and Buzsaki, G. (1996). Interneurons of the hippocampus. Hippocampus 6, 347-470.

Fukuda, A., Muramatsu, K., Okabe, A., Shimano, Y., Hida, H., Fujimoto, I., and Nishino, H. (1998). Changes in intracellular Ca2+ induced by GABAA receptor activation and reduction in CI- gradient in neonatal rat neocortex. J. Neurophysiol. 79. 439-446.

Galanopoulou, A.S. (2008). Dissociated gender-specific effects of recurrent seizures on GABA signaling in CA1 pyramidal neurons: role of GABAA receptors. J. Neurosci. 28. 1557-1567.

Gamba, G. (2005). Molecular physiology and pathophysiology of electroneutral cation-chloride cotransporters. Physiol. Rev. 85, 423-493.

Ganguly, K., Schinder, A.F., Wong, S.T., and Poo, M.M. (2001). GABA itself promotes the developmental switch of neuronal GABAergic responses from excitation to inhibition. Cell 105, 521-532.

Gao, X.B., Chen, G., and van den Pol, A.N. (1998). GABA-dependent firing of glutamate-evoked action potentials at AMPA/kainate receptors in developing hypothalamic neurons. J. Neurophysiol. 79, 716-726.

Gavrikov, K.E., Dmitriev, A.V., Keyser, K.T., and Mangel, S.C. (2003). Cationchloride cotransporters mediate neural computation in the retina. Proc. Natl. Acad. Sci. USA 100, 16047-16052.

Ge, S., Goh, E.L.K., Sailor, K.A., Kitabatake, Y., Ming, G.L., and Song, H. (2006). GABA regulates synaptic integration of newly generated neurons in the adult brain. Nature 439, 589-593.

Gerelsaikhan, T., and Turner, R.J. (2000). Transmembrane topology of the secretory Na+-K+-2Cl- cotransporter NKCC1 studied by in vitro translation. J. Biol. Chem. 275, 40471-40477.

Glickfeld, L.L., Roberts, J.D., Somogyi, P., and Scanziani, M. (2009). Interneurons hyperpolarize pyramidal cells along their entire somatodendritic axis. Nat. Neurosci. 12, 21-23.

Gonzalez-Islas, C.E., Chub, N.L., and Wenner, P. (2008). NKCC1 and AE3 appear to accumulate chloride in embryonic motoneurons. J. Neurophysiol. 101. 507-518.

Gulledge, A.T., and Stuart, G.J. (2003). Excitatory actions of GABA in the cortex. Neuron 37, 299-309.





Gulyas, A.I., Sik, A., Payne, J.A., Kaila, K., and Freund, T.F. (2001). The KCI cotransporter, KCC2, is highly expressed in the vicinity of excitatory synapses in the rat hippocampus. Eur. J. Neurosci. 13, 2205-2217.

Haglund, M.M., and Hochman, D.W. (2005). Furosemide and mannitol suppression of epileptic activity in the human brain. J. Neurophysiol. 94,

Heck, N., Kilb, W., Reiprich, P., Kubota, H., Furukawa, T., Fukuda, A., and Luhmann, H.J. (2007). GABA-A receptors regulate neocortical neuronal migration in vitro and in vivo. Cereb. Cortex 17, 138-148.

Hekmat-Scafe, D.S., Lundy, M.Y., Ranga, R., and Tanouye, M.A. (2006). Mutations in the K+/Cl- cotransporter gene kazachoc (kcc) increase seizure susceptibility in Drosophila. J. Neurosci. 26, 8943-8954.

Hennou, S., Khalilov, I., Diabira, D., Ben Ari, Y., and Gozlan, H. (2002). Early sequential formation of functional GABA(A) and glutamatergic synapses on CA1 interneurons of the rat foetal hippocampus. Eur. J. Neurosci. 16, 197-208.

Hensch, T.K. (2005). Critical period plasticity in local cortical circuits. Nat. Rev. Neurosci. 6, 877-888.

Hentschke, M., Wiemann, M., Hentschke, S., Kurth, I., Hermans-Borgmeyer, I., Seidenbecher, T., Jentsch, T.J., Gal, A., and Hübner, C.A. (2006). Mice with a targeted disruption of the CI-/HCO3- exchanger AE3 display a reduced seizure threshold. Mol. Cell. Biol. 26, 182-191.

Hilgenberg, L.G.W., Su, H., Gu, H., O'Dowd, D.K., and Smith, M.A. (2006). [alpha]3Na+/K+-ATPase is a neuronal receptor for agrin. Cell 125, 359-369.

Hochman, D.W., Baraban, S.C., Owens, J.W., and Schwartzkroin, P.A. (1995). Dissociation of synchronization and excitability in furosemide blockade of epileptiform activity. Science 270, 99-102.

Howard, H.C., Mount, D.B., Rochefort, D., Byun, N., Dupre, N., Lu, J., Fan, X., Song, L., Riviere, J.B., Prevost, C., et al. (2002). The K-Cl cotransporter KCC3 is mutant in a severe peripheral neuropathy associated with agenesis of the corpus callosum. Nat. Genet. 32, 384-392.

Huberfeld, G., Wittner, L., Clemenceau, S., Baulac, M., Kaila, K., Miles, R., and Rivera, C. (2007). Perturbed chloride homeostasis and GABAergic signaling in human temporal lobe epilepsy. J. Neurosci. 27, 9866-9873.

Hübner, C.A., Lorke, D.E., and Hermans-Borgmeyer, I. (2001a). Expression of the Na-K-2Cl-cotransporter NKCC1 during mouse development. Mech. Dev. 102, 267-269.

Hübner, C.A., Stein, V., Hermans-Borgmeyer, I., Meyer, T., Ballanyi, K., and Jentsch, T.J. (2001b). Disruption of KCC2 reveals an essential role of K-Cl cotransport already in early synaptic inhibition. Neuron 30, 515-524.

Ikeda, K., Onimaru, H., Yamada, J., Inoue, K., Ueno, S., Onaka, T., Toyoda, H., Arata, A., Ishikawa, T., Taketo, M.M., et al. (2004). Malfunction of respiratoryrelated neuronal activity in Na+, K+-ATPase alpha 2 subunit-deficient mice is attributable to abnormal CI- homeostasis in brainstem neurons. J. Neurosci. 24, 10693-10701.

Inoue, K., Yamada, J., Ueno, S., and Fukuda, A. (2006). Brain-type creatine kinase activates neuron-specific K+-Cl- co-transporter KCC2. J. Neurochem.

Jang, I.S., Jeong, H.J., and Akaike, N. (2001). Contribution of the Na-K-Cl cotransporter on GABAA receptor-mediated presynaptic depolarization in excitatory nerve terminals. J. Neurosci. 21, 5962-5972.

Jang, I.S., Nakamura, M., Ito, Y., and Akaike, N. (2006). Presynaptic GABAA receptors facilitate spontaneous glutamate release from presynaptic terminals on mechanically dissociated rat CA3 pyramidal neurons. Neuroscience 138, 25-35.

Jarolimek, W., Lewen, A., and Misgeld, U. (1999). A furosemide-sensitive K+-Cl- cotransporter counteracts intracellular Cl- accumulation and depletion in cultured rat midbrain neurons. J. Neurosci. 19, 4695-4704.

Jean-Xavier, C., Mentis, G.Z., O'Donovan, M.J., Cattaert, D., and Vinay, L. (2007). Dual personality of GABA/glycine-mediated depolarizations in immature spinal cord. Proc. Natl. Acad. Sci. USA 104, 11477-11482.

Jensen, T.S. (2002). Anticonvulsants in neuropathic pain: rationale and clinical evidence. Eur. J. Pain 6, 61-68.

Jin, Y., Jorgensen, E., Hartwieg, E., and Horvitz, H.R. (1999). The Caenorhabditis elegans gene unc-25 encodes glutamic acid decarboxylase and is required for synaptic transmission but not synaptic development. J. Neurosci. 19.539-548.

Jin, X., Huguenard, J.R., and Prince, D.A. (2005). Impaired CI- extrusion in layer V pyramidal neurons of chronically injured epileptogenic neocortex. J. Neurophysiol. 93, 2117-2126.

Kahle, K.T., Staley, K.J., Nahed, B.V., Gamba, G., Hebert, S.C., Lifton, R.P., and Mount, D.B. (2008a). Roles of the cation-chloride cotransporters in neurological disease. Nat. Clin. Pract. Neurol. 4, 490-503.

Kahle, K.T., Ring, A.M., and Lifton, R.P. (2008b). Molecular physiology of the WNK kinases. Annu. Rev. Physiol. 70, 329-355.

Kaila, K. (1994). Ionic basis of GABA(A) receptor channel function in the nervous system. Prog. Neurobiol. 42, 489-537.

Kaila, K., and Voipio, J. (1987). Postsynaptic fall in intracellular pH induced by GABA-activated bicarbonate conductance, Nature 330, 163-165.

Kaila, K., Voipio, J., Paalasmaa, P., Pasternack, M., and Deisz, R.A. (1993). The role of bicarbonate in GABAA receptor-mediated IPSPs of rat neocortical neurones. J. Physiol. 464, 273-289.

Kaila, K., Lamsa, K., Smirnov, S., Taira, T., and Voipio, J. (1997). Long-lasting GABA-mediated depolarization evoked by high-frequency stimulation in pyramidal neurons of rat hippocampal slice is attributable to a network-driven, bicarbonate-dependent K+ transient. J. Neurosci. 17, 7662-7672.

Kakazu, Y., Akaike, N., Komiyama, S., and Nabekura, J. (1999). Regulation of intracellular chloride by cotransporters in developing lateral superior olive neurons. J. Neurosci. 19, 2843-2851.

Kakazu, Y., Uchida, S., Nakagawa, T., Akaike, N., and Nabekura, J. (2000). Reversibility and cation selectivity of the K+-Cl- cotransport in rat central neurons. J. Neurophysiol. 84, 281-288.

Kanaka, C., Ohno, K., Okabe, A., Kuriyama, K., Itoh, T., Fukuda, A., and Sato, K. (2001). The differential expression patterns of messenger RNAs encoding K-Cl  $\,$ cotransporters (KCC1,2) and Na-K-2Cl cotransporter (NKCC1) in the rat nervous system. Neuroscience 104, 933-946.

Kandler, K., and Gillespie, D.C. (2005). Developmental refinement of inhibitory sound-localization circuits. Trends Neurosci. 28, 290-296.

Kaneko, H., Putzier, I., Frings, S., Kaupp, U.B., and Gensch, T. (2004). Chloride accumulation in mammalian olfactory sensory neurons. J. Neurosci. 24, 7931-7938.

Kanold, P.O., and Shatz, C.J. (2006). Subplate neurons regulate maturation of cortical inhibition and outcome of ocular dominance plasticity. Neuron 51, 627-638.

Karadsheh, M.F., and Delpire, E. (2001). Neuronal restrictive silencing element is found in the KCC2 gene: molecular basis for KCC2-specific expression in neurons. J. Neurophysiol. 85, 995-997.

Katchman, A.N., Vicini, S., and Hershkowitz, N. (1994). Mechanism of early anoxia-induced suppression of the GABAA- mediated inhibitory postsynaptic current. J. Neurophysiol. 71, 1128-1138.

Katz, B. (1966). Nerve, Muscle and Synapse (New York: McGraw-Hill).

Katz, L.C., and Shatz, C.J. (1996). Synaptic activity and the construction of cortical circuits. Science 274, 1133-1138.

Khalilov, I., Dzhala, V., Ben Ari, Y., and Khazipov, R. (1999). Dual role of GABA in the neonatal rat hippocampus. Dev. Neurosci. 21, 310-319.

Khazipov, R., Sirota, A., Leinekugel, X., Holmes, G.L., Ben Ari, Y., and Buzsaki, G. (2004). Early motor activity drives spindle bursts in the developing somatosensory cortex. Nature 432, 758-761.

Khirug, S., Huttu, K., Ludwig, A., Smirnov, S., Voipio, J., Rivera, C., Kaila, K., and Khiroug, L. (2005). Distinct properties of functional KCC2 expression in immature mouse hippocampal neurons in culture and in acute slices. Eur. J. Neurosci. 21, 899-904.

Khirug, S., Yamada, J., Afzalov, R., Voipio, J., Khiroug, L., and Kaila, K. (2008). GABAergic depolarization of the axon initial segment in cortical principal



- neurons is caused by the Na-K-2Cl cotransporter NKCC1. J. Neurosci. 28,
- Kilb, W., Ikeda, M., Uchida, K., Okabe, A., Fukuda, A., and Luhmann, H.J. (2002). Depolarizing glycine responses in Cajal-Retzius cells of neonatal rat cerebral cortex. Neuroscience 112, 299-307.
- Kilb, W., Sinning, A., and Luhmann, H.J. (2007). Model-specific effects of burnetanide on epileptiform activity in the in-vitro intact hippocampus of the newborn mouse. Neuropharmacology 53, 524-533.
- Kohling, R., Lucke, A., Straub, H., Speckmann, E.J., Tuxhorn, I., Wolf, P., Pannek, H., and Oppel, F. (1998). Spontaneous sharp waves in human neocortical slices excised from epileptic patients. Brain 121, 1073-1087.
- Kyrozis, A., and Reichling, D.B. (1995). Perforated-patch recording with gramicidin avoids artifactual changes in intracellular chloride concentration. J. Neurosci. Methods 57, 27-35.
- Lamsa, K., Palva, J.M., Ruusuvuori, E., Kaila, K., and Taira, T. (2000). Synaptic GABA(A) activation inhibits AMPA-kainate receptor-mediated bursting in the newborn (P0-P2) rat hippocampus. J. Neurophysiol. 83, 359-366.
- Laplagne, D.A., Kamienkowski, J.E., Esposito, M.S., Piatti, V.C., Zhao, C., Gage, F.H., and Schinder, A.F. (2007). Similar GABAergic inputs in dentate granule cells born during embryonic and adult neurogenesis. Eur. J. Neurosci. 25, 2973-2981.
- Le Rouzic, P., Ivanov, T.R., Stanley, P.J., Baudoin, F.M.H., Chan, F., Pinteaux, E., Brown, P.D., and Luckman, S.M. (2006). KCC3 and KCC4 expression in rat adult forebrain. Brain Res. 1110, 39-45.
- Lee, H.H.C., Walker, J.A., Williams, J.R., Goodier, R.J., Payne, J.A., and Moss, S.J. (2007). Direct protein kinase C-dependent phosphorylation regulates the cell surface stability and activity of the potassium chloride cotransporter KCC2. J. Biol. Chem. 282, 29777-29784.
- Leitch, E., Coaker, J., Young, C., Mehta, V., and Sernagor, E. (2005). GABA type-A activity controls its own developmental polarity switch in the maturing retina. J. Neurosci. 25, 4801-4805.
- Lessmann, V., Gottmann, K., and Malcangio, M. (2003). Neurotrophin secretion: current facts and future prospects. Prog. Neurobiol. 69, 341-374.
- Li, H., Tornberg, J., Kaila, K., Airaksinen, M.S., and Rivera, C. (2002). Patterns of cation-chloride cotransporter expression during embryonic rodent CNS development, Eur. J. Neurosci, 16, 2358-2370.
- Li, H., Khirug, S., Cai, C., Ludwig, A., Blaesse, P., Kolikova, J., Afzalov, R., Coleman, S.K., Lauri, S., Airaksinen, M.S., et al. (2007). KCC2 interacts with the dendritic cytoskeleton to promote spine development. Neuron 56, 1019-1033.
- Li, B.Q., McKernan, K., and Shen, W. (2008). Spatial and temporal distribution patterns of Na-K-2C1 cotransporter in adult and developing mouse retinas. Vis. Neurosci. 25, 109-123.
- Liu, X.X., Wang, Q., Haydar, T.F., and Bordey, A. (2005). Nonsynaptic GABA signaling in postnatal subventricular zone controls proliferation of GFAPexpressing progenitors. Nat. Neurosci. 8, 1179-1187.
- Liu, Z., Neff, R.A., and Berg, D.K. (2006). Sequential interplay of nicotinic and GABAergic signaling guides neuronal development. Science 314, 1610–1613.
- LoTurco, J.J., Owens, D.F., Heath, M.J.S., Davis, M.B.E., and Kriegstein, A.R. (1995). GABA and glutamate depolarize cortical progenitor cells and inhibit DNA synthesis, Neuron 15, 1287-1298.
- Ludwig, A., Li, H., Saarma, M., Kaila, K., and Rivera, C. (2003). Developmental up-regulation of KCC2 in the absence of GABAergic and glutamatergic transmission. Eur. J. Neurosci. 18, 3199-3206.
- Lytle, C., and Forbush, B. (1992). The Na-K-Cl cotransport protein of shark rectal gland. II. Regulation by direct phosphorylation. J. Biol. Chem. 267, 25438-25443.
- Lytle, C., Xu, J.C., Biemesderfer, D., and Forbush, B. (1995). Distribution and diversity of Na-K-Cl cotransport proteins: a study with monoclonal antibodies. Am. J. Physiol. 269, C1496-C1505.
- Martina, M., Royer, S., and Pare, D. (2001). Cell-type-specific GABA responses and chloride homeostasis in the cortex and amygdala. J. Neurophysiol. 86, 2887-2895.

- Marty, A., and Llano, I. (2005). Excitatory effects of GABA in established brain networks. Trends Neurosci. 28, 284-289.
- Marty, S., Berninger, B., Carroll, P., and Thoenen, H. (1996). GABAergic stimulation regulates the phenotype of hippocampal interneurons through the regulation of brain-derived neurotrophic factor. Neuron 16, 565-570.
- Marty, S., Wehrle, R., and Sotelo, C. (2000). Neuronal activity and brainderived neurotrophic factor regulate the density of inhibitory synapses in organotypic slice cultures of postnatal hippocampus. J. Neurosci. 20, 8087-8095.
- Mercado, A., Mount, D.B., and Gamba, G. (2004). Electroneutral cation-chloride cotransporters in the central nervous system. Neurochem. Res. 29, 17-25.
- Mikawa, S., Wang, C., Shu, F., Wang, T., Fukuda, A., and Sato, K. (2002). Developmental changes in KCC1, KCC2 and NKCC1 mRNAs in the rat cerebellum. Brain Res. Dev. Brain Res. 136, 93-100.
- Minlebaev, M., Ben Ari, Y., and Khazipov, R. (2007). Network mechanisms of spindle-burst oscillations in the neonatal rat barrel cortex in vivo. J. Neurophysiol. 97, 692-700.
- Misgeld, U., Deisz, R.A., Dodt, H.U., and Lux, H.D. (1986). The role of chloride transport in postsynaptic inhibition of hippocampal neurons. Science 232, 1413-1415.
- Mulley, J.C., Scheffer, I.E., Petrou, S., and Berkovic, S.F. (2003). Channelopathies as a genetic cause of epilepsy. Curr. Opin. Neurol. 16, 171-176.
- Mulley, J.C., Scheffer, I.E., Harkin, L.A., Berkovic, S.F., and Dibbens, L.M. (2005). Susceptibility genes for complex epilepsy. Hum. Mol. Genet. 14, R243-R249.
- Munoz, A., Mendez, P., DeFelipe, J., and Alvarez-Leefmans, F.J. (2007). Cation-chloride cotransporters and GABA-ergic innervation in the human epileptic hippocampus. Epilepsia 48, 663-673.
- Nabekura, J., Ueno, T., Okabe, A., Furuta, A., Iwaki, T., Shimizu-Okabe, C., Fukuda, A., and Akaike, N. (2002). Reduction of KCC2 expression and GABA(A) receptor-mediated excitation after in vivo axonal injury. J. Neurosci. 22, 4412-
- O'Donovan, M.J. (1999). The origin of spontaneous activity in developing networks of the vertebrate nervous system. Curr. Opin. Neurol. 9, 94-104.
- Obata, K., Oide, M., and Tanaka, H. (1978). Excitatory and inhibitory actions of GABA and glycine on embryonic chick spinal neurons in culture. Brain Res. 144. 179-184.
- Owens, D.F., and Kriegstein, A.R. (2002). Is there more to GABA than synaptic inhibiton? Nat. Rev. Neurosci. 3, 715-727.
- Owens, D.F., Liu, X.L., and Kriegstein, A.R. (1999). Changing properties of GABA(A) receptor-mediated signaling during early neocortical development. J. Neurophysiol. 82, 570-583.
- Pace, A.J., Lee, E., Athirakul, K., Coffman, T.M., O'Brien, D.A., and Koller, B.H. (2000). Failure of spermatogenesis in mouse lines deficient in the Na+-K+-2CI(-) cotransporter. J. Clin. Invest. 105, 441-450.
- Papp, E., Rivera, C., Kaila, K., and Freund, T.F. (2008). Relationship between neuronal vulnerability and potassium-chloride cotransporter 2 immunoreactivity in hippocampus following transient forebrain ischemia. Neuroscience 154, 677-689.
- Pathak, H.R., Weissinger, F., Terunuma, M., Carlson, G.C., Hsu, F.C., Moss, S.J., and Coulter, D.A. (2007). Disrupted dentate granule cell chloride regulation enhances synaptic excitability during development of temporal lobe epilepsy. J. Neurosci. 27, 14012-14022.
- Payne, J.A. (1997). Functional characterization of the neuronal-specific K-Cl cotransporter - implications for [K+](o) regulation. Am. J. Physiol. 42, C1516-C1525
- Payne, J.A., Stevenson, T.J., and Donaldson, L.F. (1996). Molecular characterization of a putative K-Cl cotransporter in rat brain. A neuronal-specific isoform. J. Biol. Chem. 271, 16245-16252.
- Payne, J.A., Rivera, C., Voipio, J., and Kaila, K. (2003). Cation-chloride co-transporters in neuronal communication, development and trauma. Trends Neurosci. 26, 199-206.



Pearson, M.M., Lu, J., Mount, D.B., and Delpire, E. (2001). Localization of the K-Cl-cotransporter, KCC3, in the central and peripheral nervous systems: expression in the choroid plexus, large neurons and white matter tracts. Neuroscience *103*, 481–491.

Pedersen, S.F., O'Donnell, M.E., Anderson, S.E., and Cala, P.M. (2006). Physiology and pathophysiology of Na+/H+ exchange and Na+-K+-2Cl- cotransport in the heart, brain, and blood. Am. J. Physiol. 291, R1–R25.

Pedersen, M., Carmosino, M., and Forbush, B. (2008). Intramolecular and intermolecular fluorescence resonance energy transfer in fluorescent protein-tagged Na-K-Cl cotransporter (NKCC1): sensitivity to regulatory conformational change and cell volume. J. Biol. Chem. 283, 2663–2674.

Perrot-Sinal, T.S., Sinal, C.J., Reader, J.C., Speert, D.B., and McCarthy, M.M. (2007). Sex differences in the chloride cotransporters, NKCC1 and KCC2, in the developing hypothalamus. J. Neuroendocrinol. *19*, 302–308.

Pieraut, S., Laurent-Matha, V., Sar, C., Hubert, T., Mechaly, I., Hilaire, C., Mersel, M., Delpire, E., Valmier, J., and Scamps, F. (2007). NKCC1 phosphorylation stimulates neurite growth of injured adult sensory neurons. J. Neurosci. 27. 6751–6759.

Plotkin, M.D., Snyder, E.Y., Hebert, S.C., and Delpire, E. (1997). Expression of the Na-K-2Cl cotransporter is developmentally regulated in postnatal rat brains: a possible mechanism underlying GABA's excitatory role in immature brain. J. Neurobiol. 33, 781–795.

Pond, B.B., Berglund, K., Kuner, T., Feng, G.P., Augustine, G.J., and Schwartz-Bloom, R.D. (2006). The chloride transporter Na+-K+-Cl- cotransporter isoform-1 contributes to intracellular chloride increases after in vitro ischemia. J. Neurosci. *26*, 1396–1406.

Praetorius, J. (2007). Water and solute secretion by the choroid plexus. Pflugers Arch. 454, 1–18.

Price, G.D., and Trussell, L.O. (2006). Estimate of the chloride concentration in a central glutamatergic terminal: a gramicidin perforated-patch study on the Calyx of Held. J. Neurosci. 26, 11432–11436.

Price, T.J., Cervero, F., and De Koninck, Y. (2005). Role of cation-chloride-cotransporters (CCC) in pain and hyperalgesia. Curr. Top. Med. Chem. 5, 547–555

Pylova, S.I., Majkowska, J., Hilgier, W., Kapuscinski, A., and Albrecht, J. (1989). Rapid decrease of high-affinity ouabain binding-sites in hippocampal CA1 region following short-term global cerebral-ischemia in rat. Brain Res. 490, 170–173.

Randall, J., Thorne, T., and Delpire, E. (1997). Partial cloning and characterization of Slc12a2: the gene encoding the secretory Na+-K+-2Cl- cotransporter. Am. J. Physiol. 273, C1267–C1277.

Reisert, J., Lai, J., Yau, K.W., and Bradley, J. (2005). Mechanism of the excitatory CI- response in mouse olfactory receptor neurons. Neuron 45, 553–561.

Rennie, J., and Boylan, G. (2007). Treatment of neonatal seizures. Arch. Dis. Child. 92, F148–F150.

Represa, A., and Ben Ari, Y. (2005). Trophic actions of GABA on neuronal development. Trends Neurosci. 28, 278–283.

Reynolds, A., Brustein, E., Liao, M., Mercado, A., Babilonia, E., Mount, D.B., and Drapeau, P. (2008). Neurogenic role of the depolarizing chloride gradient revealed by global overexpression of KCC2 from the onset of development. J. Neurosci. 28, 1588–1597.

Rheims, S., Minlebaev, M., Ivanov, A., Represa, A., Khazipov, R., Holmes, G.L., Ben Ari, Y., and Zilberter, Y. (2008). Excitatory GABA in rodent developing neocortex in vitro. J. Neurophysiol. *100*, 609–619.

Riekki, R., Pavlov, I., Tornberg, J., Lauri, S.E., Airaksinen, M.S., and Taira, T. (2008). Altered Synaptic Dynamics and Hippocampal Excitability but Normal Long-Term Plasticity in Mice Lacking Hyperpolarizing GABAA Receptor-Mediated Inhibition in CA1 Pyramidal Neurons. J. Neurophysiol. 99, 3075–3089.

Rivera, C., Voipio, J., Payne, J.A., Ruusuvuori, E., Lahtinen, H., Lamsa, K., Pirvola, U., Saarma, M., and Kaila, K. (1999). The K+/Cl- co-transporter KCC2 renders GABA hyperpolarizing during neuronal maturation. Nature 397, 251–255.

Rivera, C., Li, H., Thomas-Crusells, J., Lahtinen, H., Viitanen, T., Nanobashvili, A., Kokaia, Z., Airaksinen, M.S., Voipio, J., Kaila, K., and Saarma, M. (2002). BDNF-induced TrkB activation down-regulates the K+-Cl- cotransporter KCC2 and impairs neuronal Cl- extrusion. J. Cell Biol. *159*, 747–752.

Rivera, C., Voipio, J., Thomas-Crusells, J., Li, H., Emri, Z., Sipilä, S., Payne, J.A., Minichiello, L., Saarma, M., and Kaila, K. (2004). Mechanism of activity-dependent downregulation of the neuron-specific K-Cl cotransporter KCC2. J. Neurosci. 24. 4683–4691.

Rocha-Gonzalez, H.I., Mao, S., and Alvarez-Leefmans, F.J. (2008). Na+, K+, 2Cl(-) cotransport and intracellular chloride regulation in rat primary sensory neurons: Thermodynamic and kinetic aspects. J. Neurophysiol. *100*, 169–184.

Rogawski, M.A., and Loscher, W. (2004). The neurobiology of antiepileptic drugs for the treatment of nonepileptic conditions. Nat. Med. *10*, 685–692.

Rohrbough, J., and Spitzer, N.C. (1996). Regulation of intracellular CI- levels by Na(+)-dependent CI- cotransport distinguishes depolarizing from hyperpolarizing GABAA receptor-mediated responses in spinal neurons. J. Neurosci. 16, 82–91.

Romero, M.F., Fulton, C.M., and Boron, W.F. (2004). The SLC4 family of HCO3- transporters. Pflugers Arch. 447, 495–509.

Romo-Parra, H., Trevino, M., Heinemann, U., and Gutierrez, R. (2008). GABA actions in hippocampal area CA3 during postnatal development: Differential shift from depolarizing to hyperpolarizing in somatic and dendritic compartments. J. Neurophysiol. 99, 1523–1534.

Ross, S.T., and Soltesz, I. (2000). Selective depolarization of interneurons in the early posttraumatic dentate gyrus: involvement of the Na(+)/K(+)-ATPase. J. Neurophysiol. 83, 2916–2930.

Rudomin, P., and Schmidt, R.F. (1999). Presynaptic inhibition in the vertebrate spinal cord revisited. Exp. Brain Res. 129, 1–37.

Russell, J.M. (2000). Sodium-potassium-chloride cotransport. Physiol. Rev. 80, 211–276.

Rust, M.B., Faulhaber, J., Budack, M.K., Pfeffer, C., Maritzen, T., Didie, M., Beck, F.X., Boettger, T., Schubert, R., Ehmke, H., et al. (2006). Neurogenic mechanisms contribute to hypertension in mice with disruption of the K-Cl cotransporter KCC3. Circ. Res. 98, 549–556.

Rust, M.B., Alper, S.L., Rudhard, Y., Shmukler, B.E., Vicente, R., Brugnara, C., Trudel, M., Jentsch, T.J., and Hübner, C.A. (2007). Disruption of erythroid K-Cl cotransporters alters erythrocyte volume and partially rescues erythrocyte dehydration in SAD mice. J. Clin. Invest. 117, 1708–1717.

Saier, M.H. (2003). Tracing pathways of transport protein evolution. Mol. Microbiol. 48, 1145–1156.

Salin-Cantegrel, A., Shekarabi, M., Holbert, S., Dion, P., Rochefort, D., Laganiere, J., Dacal, S., Hince, P., Karemera, L., Gaspar, C., et al. (2008). HMSN/ACC truncation mutations disrupt brain-type creatine kinase-dependant activation of K+/Cl- cotransporter 3. Hum. Mol. Genet. *17*, 2703–2711.

Serafini, R., Valeyev, A.Y., Barker, J.L., and Poulter, M.O. (1995). Depolarizing GABA-activated CI- channels in embryonic rat spinal and olfactory bulb cells. J. Physiol. 488, 371–386.

Shulga, A., Thomas-Crusells, J., Sigl, T., Blaesse, A., Mestres, P., Meyer, M., Yan, Q., Kaila, K., Saarma, M., Rivera, C., and Giehl, K.M. (2008). Posttraumatic GABAA-mediated [Ca2+]i increase is essential for the induction of brain-derived neurotrophic factor-dependent survival of mature central neurons. J. Neurosci. 28, 6996–7005.

Simard, C.F., Bergeron, M.J., Frenette-Cotton, R., Carpentier, G.A., Pelchat, M.E., Caron, L., and Isenring, P. (2007). Homooligomeric and heterooligomeric associations between K+-Cl- Cotransporter isoforms and between K+-Cl- and Na+-K+-Cl- cotransporters. J. Biol. Chem. 282, 18083–18093.

Simat, M., Ambrosetti, L., Lardi-Studler, B., and Fritschy, J.M. (2007). GABAergic synaptogenesis marks the onset of differentiation of basket and stellate cells in mouse cerebellum. Eur. J. Neurosci. 26, 2239–2256.

Sipilä, S.T., Huttu, K., Soltesz, I., Voipio, J., and Kaila, K. (2005). Depolarizing GABA acts on intrinsically bursting pyramidal neurons to drive giant depolarizing potentials in the immature hippocampus. J. Neurosci. *25*, 5280–5289.



- Sipilä, S.T., Schuchmann, S., Voipio, J., Yamada, J., and Kaila, K. (2006). The cation-chloride cotransporter NKCC1 promotes sharp waves in the neonatal rat hippocampus. J. Physiol. 573, 765-773.
- Spitzer, N.C. (2006). Electrical activity in early neuronal development. Nature 444, 707-712.
- Staley, K.J. (2002). Diuretics as antiepileptic drugs: should we go with the flow? Epilepsy Curr. 2, 35-38.
- Stein, V., Hermans-Borgmeyer, I., Jentsch, T.J., and Hübner, C.A. (2004). Expression of the KC1 cotransporter KCC2 parallels neuronal maturation and the emergence of low intracellular chloride. J. Comp. Neurol. 468, 57-64.
- Stell, B.M., Rostaing, P., Triller, A., and Marty, A. (2007). Activation of presynaptic GABAA receptors induces glutamate release from parallel fiber synapses. J. Neurosci. 27, 9022-9031.
- Szabadics, J., Varga, C., Molnar, G., Olah, S., Barzo, P., and Tamas, G. (2006). Excitatory effect of GABAergic axo-axonic cells in cortical microcircuits. Science 311, 233-235.
- Thompson, S.M., and Gähwiler, B.H. (1989). Activity-dependent disinhibition. II. Effects of extracellular potassium, furosemide, and membrane potential on ECI- in hippocampal CA3 neurons. J. Neurophysiol. 61, 512-523.
- Titz, S., Hans, M., Kelsch, W., Lewen, A., Swandulla, D., and Misgeld, U. (2003). Hyperpolarizing inhibition develops without trophic support by GABA in cultured rat midbrain neurons. J. Physiol. 550, 719-730.
- Toni, N., Laplagne, D.A., Zhao, C., Lombardi, G., Ribak, C.E., Gage, F.H., and Schinder, A.F. (2008). Neurons born in the adult dentate gyrus form functional synapses with target cells. Nat. Neurosci. 11, 901-907.
- Tornberg, J., Voikar, V., Savilahti, H., Rauvala, H., and Airaksinen, M.S. (2005). Behavioural phenotypes of hypomorphic KCC2-deficient mice. Eur. J. Neurosci. 21, 1327-1337.
- Tozuka, Y., Fukuda, S., Namba, T., Seki, T., and Hisatsune, T. (2005). GABAergic excitation promotes neuronal differentiation in adult hippocampal progenitor cells. Neuron 47, 803-815.
- Trevino, M., and Gutierrez, R. (2005). The GABAergic projection of the dentate gyrus to hippocampal area CA3 of the rat: pre- and postsynaptic actions after seizures. J. Physiol. 567, 939-949.
- Tyzio, R., Cossart, R., Khalilov, I., Minlebaev, M., Hübner, C.A., Represa, A., Ben Ari, Y., and Khazipov, R. (2006). Maternal oxytocin triggers a transient inhibitory switch in GABA signaling in the fetal brain during delivery. Science 314, 1788-1792.
- Tyzio, R., Minlebaev, M., Rheims, S., Ivanov, A., Jorquera, I., Holmes, G.L., Zilberter, Y., Ben Ari, Y., and Khazipov, R. (2008). Postnatal changes in somatic gamma-aminobutyric acid signalling in the rat hippocampus. Eur. J. Neurosci. 27. 2515-2528.
- Uvarov, P., Pruunsild, P., Timmusk, T., and Airaksinen, M.S. (2005). Neuronal K+/Cl- co-transporter (KCC2) transgenes lacking neurone restrictive silencer element recapitulate CNS neurone-specific expression and developmental up-regulation of endogenous KCC2 gene. J. Neurochem. 95, 1144-1155.
- Uvarov, P., Ludwig, A., Markkanen, M., Rivera, C., and Airaksinen, M.S. (2006). Upregulation of the neuron-specific K+/Cl- cotransporter expression by transcription factor early growth response 4. J. Neurosci. 26, 13463-13473.
- Uvarov, P., Ludwig, A., Markkanen, M., Pruunsild, P., Kaila, K., Delpire, E., Timmusk, T., Rivera, C., and Airaksinen, M.S. (2007). A novel N-terminal isoform of the neuron-specific K-Cl cotransporter KCC2. J. Biol. Chem. 282, 30570-30576.
- Vale, C., Caminos, E., Martinez-Galan, J.R., and Juiz, J.M. (2005). Expression and developmental regulation of the K+-Cl- cotransporter KCC2 in the cochlear nucleus. Hear. Res. 206, 107-115.
- van den Pol, A.N., Obrietan, K., and Chen, G. (1996). Excitatory actions of GABA after neuronal trauma. J. Neurosci. 16, 4283-4292.
- Vanhatalo, S., Palva, J.M., Andersson, S., Rivera, C., Voipio, J., and Kaila, K. (2005). Slow endogenous activity transients and developmental expression of K+-Cl- cotransporter 2 in the immature human cortex. Eur. J. Neurosci. 22, 2799-2804.

- Vardi, N., Zhang, L.L., Payne, J.A., and Sterling, P. (2000). Evidence that different cation chloride cotransporters in retinal neurons allow opposite responses to GABA. J. Neurosci. 20, 7657-7663.
- Varoqueaux, F., Sigler, A., Rhee, J.S., Brose, N., Enk, C., Reim, K., and Rosenmund, C. (2002). Total arrest of spontaneous and evoked synaptic transmission but normal synaptogenesis in the absence of Munc13-mediated vesicle priming. Proc. Natl. Acad. Sci. USA 99, 9037-9042.
- Verhage, M., Maia, A.S., Plomp, J.J., Brussaard, A.B., Heeroma, J.H., Vermeer, H., Toonen, R.F., Hammer, R.E., van den Berg, T.K., Missler, M., et al. (2000). Synaptic assembly of the brain in the absence of neurotransmitter secretion. Science 287, 864-869.
- Verkman, A.S. (1990). Development and biological applications of chloridesensitive fluorescent indicators. Am. J. Physiol. 259, C375-C388.
- Vibat, C.R.T., Holland, M.J., Kang, J.J., Putney, L.K., and O'Donnell, M.E. (2001). Quantitation of Na+-K+-2Cl(-) cotransport splice variants in human tissues using kinetic polymerase chain reaction. Anal. Biochem. 298, 218-230.
- Vinay, L., and Jean-Xavier, C. (2008). Plasticity of spinal cord locomotor networks and contribution of cation-chloride cotransporters. Brain Res. Brain Res. Rev. 57, 103-110.
- Voipio, J., and Kaila, K. (2000). GABAergic excitation and K(+)-mediated volume transmission in the hippocampus. Prog. Brain Res. 125, 329-338.
- Wake, H., Watanabe, M., Moorhouse, A.J., Kanematsu, T., Horibe, S., Matsukawa, N., Asai, K., Ojika, K., Hirata, M., and Nabekura, J. (2007). Early changes in KCC2 phosphorylation in response to neuronal stress result in functional downregulation. J. Neurosci. 27, 1642-1650.
- Wang, D.D., and Kriegstein, A.R. (2008). GABA regulates excitatory synapse formation in the neocortex via NMDA receptor activation. J. Neurosci. 28, 5547-5558.
- Wang, D.D., and Kriegstein, A.R. (2009). Defining the Role of GABA in Cortical Development. J. Physiol., in press. Published online January 19, 2009. jphysiol.2008.167635v1.
- Wang, C., Shimizu-Okabe, C., Watanabe, K., Okabe, A., Matsuzaki, H., Ogawa, T., Mori, N., Fukuda, A., and Sato, K. (2002). Developmental changes in KCC1, KCC2, and NKCC1 mRNA expressions in the rat brain. Brain Res. Dev. Brain Res. 139, 59-66.
- Wardle, R.A., and Poo, M.M. (2003). Brain-derived neurotrophic factor modulation of GABAergic synapses by postsynaptic regulation of chloride transport. J. Neurosci. 23, 8722-8732.
- Waxman, S.G. (2007). Channel, neuronal and clinical function in sodium channelopathies: from genotype to phenotype. Nat. Neurosci. 10, 405-409.
- Williams, J.R., and Payne, J.A. (2004). Cation transport by the neuronal K+-Clcotransporter KCC2: thermodynamics and kinetics of alternate transport modes. Am. J. Physiol. 287, C919-C931.
- Williams, J.R., Sharp, J.W., Kumari, V.G., Wilson, M., and Payne, J.A. (1999). The neuron-specific K-Cl cotransporter, KCC2. Antibody development and initial characterization of the protein. J. Biol. Chem. 274, 12656-12664.
- Wojcik, S.M., Katsurabayashi, S., Guillemin, I., Friauf, E., Rosenmund, C., Brose, N., and Rhee, J.S. (2006). A shared vesicular carrier allows synaptic corelease of GABA and glycine. Neuron 50, 575-587.
- Woo, N.S., Lu, J.M., England, R., McClellan, R., Dufour, S., Mount, D.B., Deutch, A.Y., Lovinger, D.M., and Delpire, E. (2002). Hyperexcitability and epilepsy associated with disruption of the mouse neuronal-specific K-Cl cotransporter gene. Hippocampus 12, 258-268.
- Woodin, M.A., Ganguly, K., and Poo, M.M. (2003). Coincident pre- and postsynaptic activity modifies GABAergic synapses by postsynaptic changes in Cl- transporter activity. Neuron 39, 807-820.
- Xu, J.C., Lytle, C., Zhu, T.T., Payne, J.A., Benz, E.J., and Forbush, B. (1994). Molecular cloning and functional expression of the bumetanide-sensitive Na-K-Cl cotransporter. Proc. Natl. Acad. Sci. USA 91, 2201-2205.
- Yamada, J., Okabe, A., Toyoda, H., Kilb, W., Luhmann, H.J., and Fukuda, A. (2004). CI- uptake promoting depolarizing GABA actions in immature rat neocortical neurones is mediated by NKCC1. J. Physiol. 557, 829-841.





Yan, Y., Dempsey, R.J., and Sun, D. (2001a). Expression of Na+-K+-Clcotransporter in rat brain during development and its localization in mature astrocytes. Brain Res. 911, 43-55.

Yan, Y.P., Dempsey, R.J., and Sun, D. (2001b). Na+-K+-Cl- cotransporter in rat focal cerebral ischemia. J. Cereb. Blood Flow Metab. 21, 711-721.

Zhang, L.L., Fina, M.E., and Vardi, N. (2006). Regulation of KCC2 and NKCC during development: Membrane insertion and differences between cell types. J. Comp. Neurol. 499, 132-143.

Zhang, L.L., Delpire, E., and Vardi, N. (2007). NKCC1 does not accumulate chloride in developing retinal neurons. J. Neurophysiol. 98, 266-277.

Zheng, J.J., Lee, S., and Zhou, Z.J. (2006). A transient network of intrinsically bursting starburst cells underlies the generation of retinal waves. Nat. Neurosci. 9, 363-371.

Zhu, L., Lovinger, D., and Delpire, E. (2005). Cortical neurons lacking KCC2 expression show impaired regulation of intracellular chloride. J. Neurophysiol. 93, 1557-1568.