

MEDICAL PROGRESS

BRAIN PEPTIDES (Second of Two Parts)

DOROTHY T. KRIEGER, M.D., AND JOSEPH B. MARTIN, M.D., PH.D.

PEPTIDE FUNCTIONS IN THE CENTRAL NERVOUS SYSTEM

Brain Peptides as Neuronal Markers

Most degenerative neurologic disorders are characterized by premature cell death in discrete neuronal populations. In some cases, the cellular components affected can be characterized partly by their functional roles in neuronal systems. Further delineation of the cellular basis of such neurologic disorders may contribute important new information about their pathogeneses. Immunocytochemical characterization of the cell types that are commonly affected in degenerative diseases, together with measurement of regional peptide concentrations, may provide clues to the neuronal specificity (or lack of it) involved in a given disorder. It is conceivable that one or more of the metabolic processes subserving synthesis, processing, or degradation of specific peptides is the locus of a pathologic process resulting in or contributing to cell death. An example is the recent report by Brandt et al.⁹² postulating a hyper-endorphin syndrome in a child with fatal necrotizing encephalomyelopathy (Leigh's disease). Neuronal dysfunction may also result in defects in peptide release leading to loss of transsynaptic or trophic effects that are normally exerted on other neurons.

The idea that a specific neurotransmitter deficiency is important in the pathophysiology and clinical manifestations of a neuronal-system degeneration has been developed most completely in Parkinson's disease and Huntington's chorea. Both disorders have specific clinical and neuropathologic features. Their pathogeneses are different, since Parkinson's disease is usually sporadic, whereas Huntington's disease is an autosomal-dominant disorder with a high degree of penetrance.

The documented loss of dopaminergic cells in the substantia nigra in Parkinson's disease appears to account for most of the clinical manifestations, including slowness of movement, rigidity, and tremor, and also for its successful, though temporary, alleviation with L-dopa. However, the recognition of the loss of nigral

dopaminergic cells has not provided any clues to the fundamental nature of the degenerative process. Hypothalamic dopaminergic neurons are largely preserved in this disorder, as is their neuroendocrine function, so that a generalized dopaminergic cellular disorder does not appear to occur.⁹³ Undefined additional factors must contribute to the selectivity of the degenerative process. Several peptides, including somatostatin, substance P, cholecystokinin (CCK), and the enkephalins, are present in high concentration in the basal ganglia (Table 3, Part 1).^{12,13} Immunocytochemical studies and measurement of peptide changes in Parkinson's disease may contribute to a further definition of the disorder and possibly to a clarification of its pathogenesis.

Huntington's disease affects other neuronal populations; cellular loss is most extensive in the putamen and caudate nucleus and less widespread in the lower laminae of the cerebral cortex. Other areas of the basal ganglia are less severely affected. There is no explanation for this pattern of cell death. Neurons in these regions originate from different areas of the brain during development, differ markedly in cytologic features and connectivity, have dissimilar functions, and contain several neurotransmitters. The cellular loss in the putamen and caudate is accompanied by a loss of the neurotransmitter gamma-aminobutyric acid (GABA) and of glutamic acid decarboxylase, its synthesizing enzyme.⁹⁴ However, it is clear that the deficiency of this neurotransmitter is not itself the cause of cell death, because GABA-containing neurons elsewhere in the brain survive.

The neuronal loss in Huntington's disease is also accompanied by changes in the regional content of acetylcholine and certain peptides. The concentration of substance P is decreased in the globus pallidus and the putamen.⁹⁴ Levels of CCK⁹⁵ and met-enkephalin⁹⁶ are reported to be decreased in the basal ganglia in Huntington's disease, whereas levels of vasoactive in-

Abbreviations Used:

ACTH	Adrenocorticotrophic hormone
β -LPH	β -Lipotropin
CCK	Cholecystokinin
CNS	Central nervous system
CSF	Cerebrospinal fluid
GABA	Gamma-aminobutyric acid
MSH	Melanocyte-stimulating hormone (α , β , and γ types)
PAG	Periaqueductal gray
TRH	Thyrotropin-releasing hormone
VIP	Vasoactive intestinal peptide

From the Department of Medicine, Mount Sinai Medical Center, New York, and the Department of Neurology, Massachusetts General Hospital and Harvard Medical School, Boston. Address reprint requests to Dr. Martin at the Massachusetts General Hospital, Fruit St., Boston, MA 02114.

Supported by grants (AM 26252 and NS 16367 [Huntington's Disease Center Without Walls], NB 02893, and AM 20337) from the U.S. Public Health Service and by the Lita Annenberg Hazin Charitable Trust Fund and the Dewey D. Stone and S. Robert Stone, Jr., Funds for Research in Parkinson's Disease.

testinal peptide (VIP) are normal.⁹⁷ We have recently found that concentrations of somatostatin are considerably increased in the caudate, putamen, and globus pallidus.⁹⁸ Spindel et al. have also found an increase in thyrotropin-releasing hormone (TRH) in tissue at autopsy of patients with Huntington's disease.⁹⁹ These findings point to a degree of neuronal selectivity in the degenerative process.

Alzheimer's disease, the most common cause of dementia in the United States, is characterized by loss of neurons in specific areas of the cerebral cortex, including the association areas of the orbital frontal, parietal, and temporal lobes. Changes are particularly extensive in the hippocampus. The disorder is usually sporadic but occasionally familial, and it occurs with increasing incidence in the seventh decade and beyond. The nature of the degenerative process is entirely unknown. The earliest cellular lesion, the neurofibrillary tangle, consists of an ultrastructurally identifiable alteration in the neurofilaments and neurotubules. Several studies have documented a decrease in acetylcholine content and choline acetyltransferase activity in the cortical regions affected. A recent report provided evidence for a selective decrease in somatostatin concentration in the hippocampus and other cortical regions in this disease.¹⁰⁰

It will also be of interest to investigate a group of sensory disorders in which disorders of peptidergic function seem probable. One of these, congenital insensitivity or indifference to pain, is a condition in which the patient seems unresponsive to painful stimuli. Such patients may also have anhidrosis, poor temperature regulation, and orthostatic hypotension — features that point to dysfunction of the autonomic nervous system. The cellular defect in this disorder includes loss of small neurons in the dorsal-root ganglions, degeneration in the tract of Lissauer, and diminishment of the spinal tract of the trigeminal nerve. Interestingly enough, the neuropathologic changes recognized give no explanation for the commonly associated mental subnormality. Another disorder of potential importance for students of peptide function is familial dysautonomia (Riley-Day syndrome). This disorder, which is inherited as an autosomal-recessive trait, is recognizable by failure to thrive early in infancy, episodes of autonomic dysfunction, hyporeflexia, and impaired sensation of pain and temperature. Peripheral-nerve biopsy reveals a loss of small myelinated and unmyelinated fibers. The disorder is thought to be the result of failure in neural-crest cellular migration or differentiation, giving rise to a combination of cell loss in the dorsal-root ganglions and autonomic disturbance. Associated peptide dysfunction remains to be determined.

Perhaps the most intriguing development concerning peptide functions in relation to premature neuronal-cell death will be the evolution of molecular-biologic probes that measure cellular secretory products for tagging of cell dysfunctions. Purification of messenger RNA with preserved translation capabil-

ities has been achieved in brain tissue post mortem.¹⁰¹ Attempts to determine abnormalities of cell products in single neurons obtained from brain tissue affected by Alzheimer's disease and Huntington's disease may soon be feasible.

Trophic Functions of Peptides in Neural Tissue

The possibility that peptides have trophic effects in the central nervous system (CNS) has been considered.¹⁰² In general terms, substances of small molecular weight, such as acetylcholine, GABA, glycine, and in most cases serotonin and norepinephrine, exert specific effects through receptors that are identifiable on either presynaptic or postsynaptic membranes. Many of these effects, but not all, involve alterations in ionic channels that secondarily influence transmembrane potentials and cellular electrical excitability. Some, such as dopamine, also exert effects by activation of membrane-bound adenylate cyclases. The effects of small peptides on neurons have been much more difficult to elucidate. They almost certainly include effects that are mediated by specific receptors (e.g., opioid peptides)⁵⁰ or by adenylate cyclase (e.g., VIP).¹⁰³ Other biochemical effects at the cellular level are presumed to occur. Some of these reactions may mediate chemotrophic functions that are important for intercellular connectivity during development and potentially for nervous-system regeneration. The extent to which peptidergic substances of low molecular weight may contribute to developmental structure-function relations and to plasticity within the brain are important areas for future investigation.

Peptides in Cerebrospinal Fluid

Various peptides have been detected by radioimmunoassay in cerebrospinal fluid (CSF)¹⁰⁴ (Table 4). The origin and fate of peptides found in the CSF are entirely unknown. The CSF is generally considered to be an important route for the removal of substances from the brain extracellular space, which is continuous with CSF, but it is also possible that such substances may be transported from the CSF to the brain extracellular space. Secretion of a peptide into the CSF could represent one mechanism for dissemination of the peptide to the brain parenchyma, although such a mechanism has not yet been demonstrated.

It is of interest that peptides' behavioral effects, which are often specific for an individual peptide, occur after administration through either the cerebroventricular or cisternal route. Although the concentrations given generally exceed those normally

Table 4. Peptides Detected in Human Cerebrospinal Fluid.

Thyrotropin-releasing hormone	Enkephalins
Luteinizing hormone-releasing hormone	Neurohypophyseal hormones
Somatostatin	Gastrin
Prolactin	Cholecystokinin
Growth hormone	Vasoactive intestinal peptide
ACTH	Angiotensin II
β -Endorphin	Substance P

found in CSF under physiologic conditions, the specificity of such effects is striking, and it implies that the peptide is capable of reaching specific target receptor sites and evoking selective responses. Immunoreactivity of peptides in CSF appears to be stable even when the fluid is kept at room temperature for several hours, suggesting that degradation is minimal. However, full characterization of the molecules that show peptide immunoreactivity in the CSF has not been accomplished, and further investigations are necessary.

Some fundamental observations must be made before a full understanding of peptides in CSF is achieved. These investigations would include sequential CSF sampling from lumbar cisternal and ventricular sites in human beings and primates in order to establish the dynamic characteristics of peptide secretion into CSF, further studies to document transport (active or passive) from blood to CSF and from CSF to blood, and anatomic studies of peptide penetration into the brain extracellular space after CSF instillation. The methods for these and other studies are now available and should provide important information.

Because CSF is accessible to study, it will be important to pursue analysis of peptide content in a variety of neurologic and psychiatric disorders that are unexplained at present. A few reports have documented changes in CSF concentrations of peptides either in neurologic disorders or after experimental manipulation. Levels of somatostatin are increased in lumbar CSF in various destructive and degenerative disorders of the CNS.¹⁰⁵ CSF levels of substance P are reported to be low in certain peripheral neuropathies and in the Shy-Drager syndrome¹⁰⁶ and to be elevated in spinal arachnoiditis.¹⁰⁷ Concentrations of immunoreactive β -endorphin and ACTH in CSF are reported to be normal in schizophrenia and acromegaly; a marked reduction in levels of both occurs in Cushing's disease.¹⁰⁸ Additional CSF studies in human beings are now feasible and may be pursued in association with efforts to stimulate or inhibit release of peptides selectively through peripheral-nerve stimulation or administration of neuropharmacologically active drugs. It is possible that assessment of peptide concentrations in chronic pain, narcolepsy, dystonia musculorum deformans, schizophrenia, and manic-depressive illness may contribute useful clinical information. A limiting factor, as in measurements of CSF concentrations of the metabolic products of conventional neurotransmitters, is the fact that lumbar CSF samples may principally reflect the release of spinal-cord rather than brain peptides.

Pain

The relay of painful sensations from peripheral tissues to the CNS occurs through unmyelinated or thinly myelinated nerve fibers that terminate in the outer laminae of the dorsal horn of the spinal cord. Fibers from the face end in the medulla oblongata in the spi-

nal nucleus of the fifth cranial nerve. The pain fibers for the body arise from small bipolar neurons of the dorsal-root ganglions; these neurons send processes outward that end in peripheral tissues, primarily as free nerve endings, and centrally through the dorsal root to end in the substantia gelatinosa. The neurotransmitters that relay nociception have been extensively investigated, and recent experiments have implicated several peptides, notably substance P. In addition to the initial processing of nociceptive inputs that occurs at the spinal-cord level, there are also descending inputs from brain-stem centers to the spinal cord that exert effects over local circuits.

Immunocytochemical studies indicate that substance P, somatostatin, CCK, angiotensin II, and VIP are present in spinal-ganglion neurons.¹¹ It has been estimated that 20 per cent of all ganglion cells in the cat and rat contain substance P. A smaller proportion contain somatostatin. Both cell types have extensive terminations in the substantia gelatinosa. Dense-core granules located in primary afferent terminals in the monkey show a positive reaction product for substance P.¹⁰⁹ These terminations appear to occur largely by axosomatic and axodendritic contacts on spinothalamic relay cells of the dorsal horn. Some of these cells can be identified as enkephalin-containing cells.

The evidence that substance P is a primary sensory neurotransmitter is convincing. The substance is released from the spinal cord after depolarization by potassium or electrical stimulation.¹¹⁰ Peripheral-nerve stimulation of nociceptive fibers elicits release of the peptide into the CSF.¹¹⁰ Microiontophoretic application of substance P results in prolonged excitation of dorsal-horn neurons, which can also be shown to be activated by noxious peripheral stimuli.¹¹¹

Analgesia can be induced by agents that cause depletion of substance P. Capsaicin, a substance that is obtained from red pepper and has been shown to be structurally related to homovanillic acid, causes release and nerve-terminal depletion of substance P from the spinal cord, accompanied by elevated peripheral thresholds to painful stimuli.¹¹² Long-term administration of capsaicin leads to degeneration of small-diameter sensory neurons that contain substance P, and the concentration of the peptide in the spinal cord is reduced by 50 per cent. These observations provide evidence for a function of the peptide in pain transmission at the spinal-cord level. An interaction of substance P with enkephalins at the spinal-cord level is suggested by evidence that enkephalins inhibit release of substance P from sensory neurons in culture.¹¹³

Peptides also function in pain pathways at other sites in the CNS. In the early 1970s it was shown that electrical stimulation of discrete areas of the brain stem, especially the ventrolateral periaqueductal gray (PAG) matter of the midbrain, produces analgesia in animals¹¹⁴ and human beings.¹¹⁵ The presence of opioid receptors has been documented in similar brain regions. Considerable overlap is found between the

sites of stimulation-produced analgesia and of analgesia induced by intracerebral opiate injection; the PAG is the most effective site for both. These data suggest that endogenous opioids may be active in this system. In addition to high receptor concentration in the PAG, immunohistochemical studies show a rich field of enkephalin terminations. Further evidence that opioid peptides are involved in the pain relay at this level is the finding that analgesia produced by PAG stimulation in human beings is partially blocked by naloxone¹¹⁵ and is associated with a rise in β -endorphin immunoreactivity in the CSF.¹¹⁶ Intraventricular administration of β -endorphin is effective in causing analgesia in laboratory animals, whereas the intravenous route is not, presumably because of failure to cross the blood-brain barrier.⁸³ Intrathecal injection of β -endorphin in human beings is effective in causing profound analgesia.¹¹⁷

The question of whether acupuncture is effective in altering nociceptive inputs and whether it acts by means of endogenous opioids is still unresolved. Electroacupuncture is reported to stimulate release of met-enkephalin into the CSF in human beings.¹¹⁸ As recently reviewed by Wall and Woolf,¹¹⁹ there are few controlled clinical trials of the efficacy of acupuncture in the treatment of chronic pain, and the results are inconclusive. There is no doubt, however, that acupuncture can induce powerful, though transient, analgesic effects in selected persons. The effectiveness of transcutaneous stimulation in alleviation of certain kinds of pain in the segmental distribution of the stimulation provides support for a role of sensory convergence and interdependence at the spinal-cord level. The role of peptides in mediation of these responses requires further study. Naloxone, an opiate antagonist, completely blocks the analgesic effects of exogenously administered narcotics. However, naloxone has little effect on pain thresholds in human beings, although it is postulated to be effective when given in high doses to block placebo-induced analgesia in some subjects.¹²⁰ In some but not all studies, naloxone has been reported to block acupuncture-induced analgesia in animals and human beings.¹¹⁹

Peptides and Cerebral Circulation

Autoregulation of the cerebral circulation is only partly understood. Blood flow to the brain parenchyma is regulated within a narrow range despite large fluctuations in systemic blood pressure. Neuronal activity is known to increase local blood flow. Peptides with known vasoactive properties are found within the CNS in axonal processes that terminate on or near blood vessels.

VIP, a 28-amino acid peptide, is found in highest concentrations in the cerebral cortex rather than the hypothalamus, unlike most other peptides. A well-developed plexus of VIP-immunoreactive nerve fibers is present around the anterior cerebral, proximal middle cerebral, and posterior communicating arteries.¹²¹ Terminals of these fibers penetrate the adven-

titia-media border and terminate in a morphologic arrangement that closely resembles the pattern of noradrenergic and cholinergic fibers.

In vitro studies support a function of VIP in vasomotor regulation. Larsson et al.¹²¹ found that VIP applied to a segment of cat middle-cerebral artery in an organ bath had no effect on resting tone; however, in the presence of tonic contraction induced by 5-hydroxytryptamine, VIP produced a dose-dependent dilatation. This observation suggests that the effects of VIP on cerebral blood-vessel tone are modulatory rather than direct.

There is no definitive evidence to define the physiologic role of peptides in the regulation of cerebral blood flow or oxygen metabolism. A recent report indicates that intracarotid injection of VIP increases cerebral blood flow in anesthetized animals.¹²² The fact that peptides with known vasoactive properties, such as VIP, substance P, neurotensin, bradykinin, oxytocin, and angiotensin, are found within the CNS in nerve terminals, some of which end on or near blood vessels, suggests that these peptides may contribute to regulation of cerebral vascular tone.

Opioid Peptides and Epilepsy

Parenteral administration of high doses of morphine or withdrawal of morphine in habituated persons is associated with changes in seizure threshold or the appearance of epileptiform discharges on the electroencephalogram.¹²³⁻¹²⁵ Opioid peptides applied iontophoretically usually inhibit neuronal firing,¹²⁶⁻¹²⁹ although an excitatory action is seen in the hippocampus.¹²⁹ Cortical epileptic seizures can be induced in rats by the cerebroventricular administration of met-enkephalin or leu-enkephalin.^{84,130} Chronic motor seizures in rats resulted in elevated levels of met-enkephalin in the hippocampus.¹³¹ Interestingly enough, enkephalin induces seizures in doses that are devoid of analgesic activity, and the seizures are attenuated or prevented by naloxone.⁸⁴ Conversely, morphine produces analgesia at low doses, and much higher doses are required to produce seizures.¹²⁵ A recent report indicates that anticonvulsants that are most effective in petit mal epilepsy, such as ethosuximide, trimethadione, and valproate sodium, are effective in blocking leu-enkephalin seizures, whereas major anticonvulsants such as phenytoin and phenobarbital, are ineffective.¹³²

The epileptogenic and analgesic properties of the enkephalins appear to be both dose-specific and site-specific. The same dose of enkephalin that causes analgesia without seizures when injected near the PAG causes seizures without analgesia when administered near the dorsomedial nucleus of the thalamus.^{84,132} Moreover, β -endorphin induces epileptiform activity in the limbic cortex when administered into the CSF of rats.⁸³ This activity is induced by doses that are devoid of analgesic activity, and the epileptiform activity is not associated with discernible behavioral effects.

In the baboon, *Papio papio*, which is photosensitive for seizures, the opioid peptides are without epileptogenic effect.¹³³ Cerebroventricular injection of met-enkephalin and leu-enkephalin, β -endorphin, FK 33.824 (a long-acting met-enkephalin analogue), and morphine all failed to produce epileptic effects. Although the susceptibility of met-enkephalin and leu-enkephalin to rapid biologic degradation could explain their lack of activity in the baboon, if diffusion through periventricular tissue were required to reach epileptogenic sites, this argument does not apply to the other compounds, which are stable.

In the rat, the epileptogenic effects of opiates and opioid peptides may depend on a selective inhibition of inhibitory neurons in the hippocampus.¹³⁴ Since the hippocampus does not participate in the genesis of photically induced seizures in the baboon,¹³⁵ these data are consistent with the failure of morphinomimetic peptides to facilitate such seizures. The absence of any signs of hippocampal seizures after administration of morphine or opioid peptides suggests that the baboon may be less sensitive than the rat to the indirect excitant effect of enkephalins on the hippocampus. The importance of these substances in human epilepsy remains to be determined.

Effects of Peptides on Behavior

Evidence has accumulated over the past two decades to indicate that impairment of conditioned avoidance behavior, which is present in hypophysectomized animals, is corrected by systemic or cerebroventricular administration of ACTH, α -melanotropin (α -MSH), and vasopressin. This effect can also be induced by fragments or analogues of these hormones that are devoid of any endocrine effects.¹³⁶ ACTH fragments 4-7 and 4-10, which are sequences common to ACTH, α -MSH, and the " β -MSH-like" part of β -lipotropin (β -LPH) and of γ -MSH, are effective in this regard. Similar behavioral effects of these peptides can also be observed in intact animals. The data suggest that these peptides can enhance the rate of acquisition of avoidance behavior — an effect interpreted as memory formation — and can inhibit the extinction of such avoidance behavior on stimulus withdrawal — an effect interpreted as persistence of a learned response. Further studies suggest that ACTH fragments are involved in short-term memory processes, whereas vasopressin is involved in long-term memory. It has subsequently been shown that vasopressin and oxytocin have opposite effects on avoidance behavior. Some of the effects of vasopressin in correcting deficient passive-avoidance behavior in Brattleboro rats with hereditary vasopressin deficiency may be due to enhanced central availability of oxytocin, since oxytocin levels in the CSF of these rats are elevated.¹³⁷

More recently, endorphins have also been reported to affect avoidance behavior.¹³⁸ Subcutaneously administered α -endorphin is more effective than γ -endorphin in facilitating acquisition of avoidance be-

havior and delaying extinction, whereas γ -endorphin facilitates extinction, and des-tyrosine γ -endorphin is even more active.⁷⁶ In view of the inability of the latter derivative to bind to opiate receptors, such observations suggest a dissociation of behavioral effects from opiate binding.

The effects of vasopressin on learning may be mediated by modulation of neurotransmission in distinct catecholamine systems distributed through the dorsal noradrenergic bundle.¹³⁸ Evidence from electrophysiologic studies suggests that the mechanism responsible for the effects of ACTH on behavior involves facilitation of a selective arousal state of limbic-midbrain structures, resulting in increased attention and perception and enhancement of stimulus-specific behavioral responses.¹³⁹ The biochemical basis for these effects, whether through alterations in cellular second-messenger systems such as cyclic AMP, protein synthesis, membrane phosphorylation, or neurotransmitter turnover, remains to be established.

When behavioral effects of exogenously administered peptides were initially reported, it was concluded that they might be derived from endogenous pituitary hormones secreted into plasma, with degradation fragments entering the brain. However, to date there has been no evidence of generation of such fragments in plasma; it is also doubtful, in view of the short half-life of such fragments in the blood after exogenous administration, that such a mechanism could be important. The demonstration of the synthesis of the ACTH precursor molecule in the brain suggests the possibility that these fragments are derived locally; anatomic pathways from the site of synthesis in the hypothalamic arcuate nucleus to the limbic system have been reported.¹⁴⁰ The postulated existence of vascular pathways providing for retrograde flow from the pituitary to the brain has also been suggested as the route whereby such peptides may reach the CNS.^{3,4} The latter explanation implies that conditions that affect pituitary secretion, such as stress, are involved in learning. The more attractive postulation of a CNS source for these peptides provides a means for integration of diverse types of neural mechanisms.

Initially, the physiologic relevance of these animal studies was questioned, since the major conditioned avoidance response studied was that motivated by fear. Subsequent studies have indicated the effectiveness of these peptides in operant-conditioning situations in which animals were motivated by hunger or sex.¹⁴¹ To date, few studies have been performed in a double-blind manner; there is still limited information on dose-response characteristics, the relation of the dose administered to physiologic concentrations of peptides present in the animal, sex differences in response, and circadian variability of response. Questions also arise as to whether the active ACTH 4-7 fragment (Met-Glu-His-Phe) is derived from the ACTH/MSH molecule or whether this sequence occurs in another uncharacterized peptide. Since the *dextro*-7 isomer of ACTH 4-7 has effects similar to

those of the *levo*- isomer in avoidance acquisition but opposite effects on extinction, this observation raises questions about the nature of the receptor involved. There are also reports indicating an affinity of ACTH and its fragments for opiate receptors.¹⁴² It is apparent that although a role for these peptides in cognitive function appears to have been shown, the specificity of these responses and their biochemical and physiologic basis require extended investigation.

The appearance of these studies has occasioned questions on their applicability to human learning and memory. There are reports that systemic administration of ACTH 4-10 improved the attention of mentally retarded subjects¹⁴³ and increased visual discrimination in normal men, whereas it augmented verbal skills in women.¹⁴⁴ Negative findings have also been reported,¹⁴⁵ and the number of clinical observations thus far is insufficient to yield any definite conclusions. Most of the studies to date have been performed in young, healthy volunteers; studies in persons in whom cognitive functions are disturbed, such as the elderly, may provide more consequential information.

Studies with vasopressin or its long-acting analogue 1-desamino-8-D-arginine vasopressin have reported improvements of ability to learn new information,¹⁴⁶ and beneficial effects in patients with long-term amnesia resulting from car accidents.¹⁴⁷ Beneficial effects of vasopressin have also been noted in several tests of attention and memory in elderly persons.¹⁴⁸

Feeding Behavior

The relation of food intake to caloric needs represents one of the body's major homeostatic mechanisms. Previous studies have described satiety centers located in the ventromedial nucleus, which receives stimulatory serotonergic inputs from the raphe nuclei and inhibitory inputs from the ventral adrenergic bundles, and feeding centers in the lateral hypothalamus, which receives a prominent dopaminergic input.¹⁴⁹ There is controversy over whether dopamine has a stimulatory or inhibitory role.¹⁵⁰ There appears to be a reciprocal interaction between the ventromedial hypothalamus and the lateral hypothalamus, characterized by mutual inhibition — that is, activation of one area results in inhibition of the other, whereas suppression of one results in activation of the other.

Recent studies in animals have implicated several peptides in the regulation of feeding behavior.¹⁵¹ CCK,¹⁵² TRH,¹⁵³ and insulin¹⁵⁴ are reported to be satiety factors, decreasing food intake, whereas β -endorphin has been implicated in states of increased food ingestion.¹⁵⁵ The site of action of these peptides in affecting feeding behavior has been presumed to be within the CNS, with controversy over whether this effect is mediated through the ventromedial hypothalamus¹⁵⁶; interaction with the noradrenergic system has been proposed.¹⁵⁷ Suppression of feeding by CCK

has also been reported to be mediated through a par-enteral abdominal site.¹⁵⁸

Peptidergic effects on glucoregulation have also been described. Systemic administration of neurotensin and bombesin and intraventricular administration of CCK have been shown to produce hyperglycemia.^{159,160} Studies in human subjects, however, using physiologically relevant doses of neurotensin, which were fivefold lower than those employed in animal studies, have failed to reveal a hyperglycemic effect.¹⁶¹

It has been suggested that the low levels of CCK reported to be present in the brains of genetically obese (ob/ob) mice may be causally related to the hyperphagia present in these animals.¹⁶² Other studies,¹⁶³⁻¹⁶⁵ however, have failed to show a notable alteration in brain CCK in obese animals. It has also been suggested that the elevated pituitary β -endorphin concentrations present in genetically obese animals¹⁵⁵ are causally related to such obesity. This hypothesis has been questioned,¹⁶⁶ since such elevations follow but do not precede the development of obesity. In addition, there are reports of reversal of hyperphagia by naloxone in several experimental models of obesity, not all of which were characterized by increased pituitary β -endorphin concentrations.¹⁶⁷ It would be expected, however, that central rather than pituitary β -endorphin concentrations would provide a more meaningful insight into the role of β -endorphin in obese states, and there are recent reports that brain β -endorphin concentrations are increased in genetically obese rats¹⁶⁸ and that starvation decreases such concentrations.¹⁶⁹ In view of the known interaction of β -endorphin with hypothalamic serotonin and dopamine turnover, it is indeed possible that interactions between the peptidergic and monoaminergic systems are involved in feeding regulation. There is an intriguing report that the body weight of the genetically obese rat is greatly reduced after long-term ingestion of L-dopa.¹⁷⁰

Temperature Regulation

Temperature regulation represents a complex interplay between peripheral and central receptors.^{149,171} There have been numerous studies of the effects of changes of temperature on neurotransmitter release (centrally and peripherally), as well as on hormone release and, conversely, on the effect of neurotransmitters (usually administered intracisternally) on temperature regulation.¹⁴⁹ With regard to the central mechanisms involved in temperature regulation, those studies have emphasized the role of noradrenergic and serotonergic involvement, although controversy exists over the specific effects of these substances on body temperature among various species. The effects of central cooling on peripheral release of norepinephrine and thyroid hormone also appear to be well substantiated.

Recent studies have implicated several brain peptides in the regulation of body temperature. These ef-

fects have yet to be fully integrated into any of the above schemes. Intracisternal injection of TRH causes hyperthermia, whereas small amounts (10^{-10} to 10^{-15} M) of β -endorphin, neurotensin, and bombesin result in hypothermia.¹⁷²⁻¹⁷⁴ There is some evidence of an interaction of neurotensin with dopaminergic systems and with the hypothalamic-pituitary-thyroid axis.¹⁷⁵ Bombesin has different effects in different species, and it is still uncertain whether the hypothermic effect occurs at ambient temperatures¹⁷² or only in animals exposed to cold.¹⁷⁶ The effects of bombesin are reversed by administration of either TRH, prostaglandins E_1 and E_2 , or naloxone.¹⁷⁴ TRH also reverses the hypothermic effect of endorphin, but not the antinociceptive effect of morphine.¹⁷⁷ The effect of TRH is observed in hypophysectomized animals, implying a central rather than a peripheral effect. A recent report has described a function of vasopressin in suppression of the febrile response in newborn sheep.¹⁷⁸

A possible common mechanism responsible for all the above effects is suggested by findings that a decrease in hypothalamic calcium concentration or an increase in sodium concentration elevates the core temperature,¹⁷⁹ whereas an increase in hypothalamic calcium concentration leads to a drop in core temperature.¹⁸⁰ Prostaglandins, which are reported to be thermogenic, facilitate mitochondrial uptake of calcium, thereby decreasing local concentrations.

We are indebted to Kathy Sullivan and Martha Conant for assistance in the preparation of this manuscript.

REFERENCES

92. Brandt NJ, Terenius L, Jacobsen BB, et al. Hyper-endorphin syndrome in a child with necrotizing encephalomyelopathy. *N Engl J Med.* 1980; 303:914-6.
93. Thorner MO, Login IS. Prolactin secretion as an index of brain dopaminergic function. In: Martin JB, Reichlin S, Bick KL, eds. *Neurosecretion and brain peptides: implications for brain functions and neurological disease.* New York: Raven Press, 1981:503-20.
94. Bird ED. Chemical pathology of Huntington's disease. *Annu Rev Pharmacol Toxicol.* 1980; 20:533-51.
95. Rehfeld J. Cholecystokinin. *Trends Neurosci.* 1980; 3:65-7.
96. Emson PC, Arregui A, Clement-Jones V, Sandberg BEB, Rossor M. Regional distribution of methionine-enkephalin and substance P-like immunoreactivity in normal human brain and in Huntington's disease. *Brain Res.* 1980; 199:147-60.
97. Emson PC, Fahrenkrug J, Spokes EGS. Vasoactive intestinal polypeptide (VIP): distribution in normal human brain and in Huntington's disease. *Brain Res.* 1979; 173:174-8.
98. Cooper PE, Aronin N, Bird ED, Leeman SE, Martin JB. Increased somatostatin in basal ganglia of Huntington's disease. *Neurology (Minneapolis).* (in press).
99. Spindel ER, Wurtman RJ, Bird ED. Increased TRH content of the basal ganglia in Huntington's disease. *N Engl J Med.* 1980; 303:1235-6.
100. Davies P, Katzman R, Terry RD. Reduced somatostatin-like immunoreactivity in cerebral cortex from cases of Alzheimer disease and Alzheimer senile dementia. *Nature.* 1980; 288:279-80.
101. Gilbert JM, Brown BA, Stocchi P, Bird ED, Marotta EA. The preparation of biologically active messenger RNA from human post-mortem brain tissue. *J Neurochem.* (in press).
102. Martin JB, Landis DMD. Potential implications of brain peptides in neurological disease. In: Martin JB, Reichlin S, Bick KL, eds. *Neurosecretion and brain peptides: implications for brain functions and neurological disease.* New York: Raven Press, 1981:673-90.
103. Quik M, Iversen LL, Bloom SR. Effect of vasoactive intestinal peptide (VIP) and other peptides on cyclic cAMP accumulation in rat brain. *Biochem Pharmacol.* 1978; 27:2209-13.
104. Jackson IMD. Significance and function of neuropeptides in cerebrospinal fluid. In: Wood JH, ed. *Neurobiology of cerebrospinal fluid.* New York: Plenum Press, 1980:625-50.
105. Patel YC, Rao K, Reichlin S. Somatostatin in human cerebrospinal fluid. *N Engl J Med.* 1977; 296:529-33.
106. Nutt JG, Mroz EA, Leeman SE, Williams AC, Engel WK, Chase TN. Substance P in human cerebrospinal fluid: reduction in peripheral neuropathy and autonomic dysfunction. *Neurology (Minneapolis).* 1980; 30:1280-5.
107. Jackson IMD. Neural peptides in the cerebrospinal fluid. In: Martin JB, Reichlin S, Bick KL, eds. *Neurosecretion and brain peptides: implications for brain functions and neurological disease.* New York: Raven Press, 1981:337-55.
108. Nakao K, Oki S, Tanaka I, et al. Immunoreactive β -endorphin and adrenocorticotropin in human cerebrospinal fluid. *J Clin Invest.* 1980; 66:1383-90.
109. Leeman SE, DiFiglia M, Aronin N. Neurotensin (NT) and substance P (SP) localization in monkey and rat spinal cord: light and electron microscopy. *Soc Neurosci Abstract.* 1980; 10:354.
110. Jessell TM. The role of substance P in sensory transmission and pain perception. In: Martin JB, Reichlin S, Bick KL, eds. *Neurosecretion and brain peptides: implications for brain functions and neurological disease.* New York: Raven Press, 1981:189-98.
111. Nicoll RA, Schenker C, Leeman SE. Substance P as a transmitter candidate. *Annu Rev Neurosci.* 1980; 3:227-68.
112. Yaksh TL, Farb DH, Leeman SE, Jessell TM. Intrathecal capsaicin depletes substance P in the rat spinal cord and produces prolonged thermal analgesia. *Science.* 1979; 206:481-3.
113. Mudge AW, Leeman SE, Fischbach GD. Enkephalin inhibits release of substance P from sensory neurons in culture and decreases action potential duration. *Proc Natl Acad Sci USA.* 1979; 76:526-30.
114. Mayer DJ, Wolfle TL, Akil H, Carder B, Liebeskind JC. Analgesia from electrical stimulation in the brainstem of the rat. *Science.* 1971; 174:1351-4.
115. Hosobuchi Y, Adams JE, Linchitz R. Pain relief by electrical stimulation of the central gray matter in humans and its reversal by naloxone. *Science.* 1977; 197:183-6.
116. Akil H, Richardson DE, Barchas JD, Li CH. Appearance of β -endorphin-like immunoreactivity in human ventricular cerebrospinal fluid upon analgesic electrical stimulation. *Proc Natl Acad Sci USA.* 1978; 75:5170-2.
117. Oyama T, Jin T, Yamaya R, Ling N, Guillemin R. Profound analgesic effects of β -endorphin in man. *Lancet.* 1980; 1:122-4.
118. Clement-Jones V, McLoughlin L, Lowry PJ, Besser GM, Rees LH, Wen HL. Acupuncture in heroin addicts: changes in met-enkephalin and β -endorphin in blood and cerebrospinal fluid. *Lancet.* 1979; 2:380-3.
119. Wall PD, Woolf CJ. What we don't know about pain. *Nature.* 1980; 287:185-6.
120. Basbaum AI, Fields HL. Endogenous pain control mechanisms: review and hypothesis. *Ann Neurol.* 1978; 4:451-62.
121. Larsson L-I, Edvinsson L, Fahrenkrug J, et al. Immunohistochemical localization of a vasodilatory polypeptide (VIP) in cerebrovascular nerves. *Brain Res.* 1976; 113:400-4.
122. Heistad DD, Marcus ML, Said SI, Gross PM. Effect of acetylcholine and vasoactive intestinal peptide on cerebral blood flow. *Am J Physiol.* 1980; 239:H73-80.
123. Corrado AP, Longo VG. An electrophysiological analysis of the convulsant action of morphine, codeine and thebaine. *Arch Int Pharmacodyn Ther.* 1961; 132:255-69.
124. Le Gal La Salle G, Calvino B, Ben-Ari Y. Morphine enhances amygdaloid seizures and increases inter-ictal spike frequency in kindled rats. *Neurosci Lett.* 1977; 6:255-60.
125. Urca G, Frenk H, Liebeskind JC, Taylor AN. Morphine and enkephalin: analgesic and epileptic properties. *Science.* 1977; 197:83-6.
126. Denavit-Saubié M, Champagnat J, Zieglängsberger W. Effects of opiates and methionine-enkephalin on pontine and bulbar respiratory neurones of the cat. *Brain Res.* 1978; 155:55-67.
127. Fredrickson RCA, Norris FH. Enkephalin-induced depression of single neurons in brain areas with opiate receptors — antagonism by naloxone. *Science.* 1976; 194:440-2.
128. Zieglängsberger W, Fry JP, Herz A, Moroder L, Wunsch E. Enkephalin-induced inhibition of cortical neurons and the lack of this effect in morphine tolerance/dependent rats. *Brain Res.* 1976; 115:160-4.
129. Nicoll RA, Siggins GR, Ling N, Bloom FE, Guillemin R. Neuronal actions of endorphins and enkephalins among brain regions: a comparative microiontophoretic study. *Proc Natl Acad Sci USA.* 1977; 74:2584-8.

130. Frenk H, Urca G, Liebeskind JC. Enkephalin-induced cortical seizures in the rat. *Soc Neurosci Abstract*. 1977; 3:291.
131. Hong JS, Wood PL, Gillin JC, Yang HYT, Costa E. Changes of hippocampal Met-enkephalin content after recurrent motor seizures. *Nature*. 1980; 285:231-2.
132. Snead OC III, Bearden LJ. Anticonvulsants specific for petit mal antagonize epileptogenic effect of leucine enkephalin. *Science*. 1980; 210:1031-3.
133. Meldrum BS, Menini C, Stutzmann JM, Naquet R. Effects of opiate-like peptides, morphine, and naloxone in the photosensitive baboon, *Papio papio*. *Brain Res*. 1979; 170:333-48.
134. Zieglängsberger W, French ED, Siggins GR, Bloom FE. Opioid peptides may excite hippocampal pyramidal neurons by inhibiting adjacent inhibitory interneurons. *Science*. 1979; 205:415-7.
135. Fischer-Williams M, Poncet M, Riche D, Naquet R. Light-induced epilepsy in the baboon, *Papio papio*: cortical and depth recordings. *Electroencephalogr Clin Neurophysiol*. 1968; 25:557-69.
136. de Kloet R, de Wied D. The brain as target tissue for hormones of pituitary origin: behavioral and biochemical studies. In: Martini L, Ganong WF, eds. *Frontiers in neuroendocrinology*. Vol. 6. New York: Raven Press, 1980:157-201.
137. Dogterom J, van Wimersma Greidanus JB, Swaab DF. Evidence for the release of vasopressin and oxytocin into cerebrospinal fluid: measurements in plasma and CSF of intact and hypophysectomized rats. *Neuroendocrinology*. 1977; 24:108-18.
138. Kovács GL, Bohus B, Versteeg DHG. The effects of vasopressin on memory processes: the role of noradrenergic neurotransmission. *Neuroscience*. 1979; 4:1529-37.
139. Bohus B. Effects of ACTH-like neuropeptides on animal behavior and man. *Pharmacology*. 1979; 18:113-22.
140. Watson SJ, Richard CW III, Barchas JD. Adrenocorticotropin in rat brain: immunocytochemical localization in cells and axons. *Science*. 1978; 200:1180-2.
141. Beckwith BE, Sandman CA. Behavioral influences of the neuropeptides ACTH and MSH: a methodological review. *Neurosci Biobehav Rev*. 1978; 2:311-38.
142. Terenius L. Effect of peptides and amino acids on dihydromorphine binding to the opiate receptor. *J Pharm Pharmacol*. 1975; 27:450-2.
143. Sandman CA, George J, Walker BB, Nolan JD, Kastin AJ. Neuropeptide MSH/ACTH 4-10 enhances attention in the mentally retarded. *Pharmacol Biochem Behav*. 1976; 5: Suppl 1:23-8.
144. Sandman CA, Kastin AJ. A behavioral strategy for the CNS actions of the neuropeptides. In: *Current studies of hypothalamic function*. Vol. 2. Metabolism and behaviour. Basel: S Karger, 1978:163-74.
145. Bohus B. Effects of ACTH-like neuropeptides on animal behavior and man. *Pharmacology*. 1979; 18:113-22.
146. Weingartner H, Gold P, Ballenger JC, et al. Effects of vasopressin on human memory functions. *Science*. 1981; 211:601-3.
147. Oliveros JC, Jandali MK, Timsit-Berthier M, et al. Vasopressin in amnesia. *Lancet*. 1978; 1:42-3.
148. Legros JJ, Gilot P, Seron X, et al. Influence of vasopressin on learning and memory. *Lancet*. 1978; 1:41-2.
149. Martin JB, Reichlin S, Brown GM. *Clinical neuroendocrinology*. Philadelphia: FA Davis, 1977.
150. Leibowitz SF, Rossakis C. Mapping study of brain dopamine- and epinephrine-sensitive sites which cause feeding suppression in the rat. *Brain Res*. 1979; 172:101-13.
151. Morley JE. The neuroendocrine control of the appetite: the role of the endogenous opiates, cholecystokinin, TRH, gamma-amino-butyric acid and the diazepam receptor. *Life Sci*. 1980; 27:355-68.
152. Gibbs J, Young RC, Smith GP. Cholecystokinin elicits satiety in rats with open gastric fistulas. *Nature*. 1973; 245:323-5.
153. Vijayan E, McCann SM. Suppression of feeding and drinking activity in rats following intraventricular injection of thyrotropin releasing hormone (TRH). *Endocrinology*. 1979; 100:1727-30.
154. Woods SC, Lotter EC, McKay LD, Porte D Jr. Chronic intracerebroventricular infusion of insulin reduces food intake and body weight of baboons. *Nature*. 1979; 282:503-5.
155. Margules DL, Moisset B, Lewis MJ, Shibuya H, Pert CB. β -Endorphin is associated with overeating in genetically obese mice (*ob/ob*) and rats (*fa/fa*). *Science*. 1978; 202:988-91.
156. Kulkosky PJ, Breckenridge C, Krinsky R, Woods SC. Satiety elicited by the C-terminal octapeptide of cholecystokinin-pancreozymin in normal and VMH-lesioned rats. *Behav Biol*. 1976; 18:227-34.
157. McCaleb ML, Myers RD. Cholecystokinin acts on the hypothalamic "noradrenergic system" involved in feeding. *Peptides*. 1980; 1:47-50.
158. Smith GP, Gibbs J. Brain-gut peptides and the control of food intake. In: Martin JB, Reichlin S, Bick KL, eds. *Neurosecretion and brain peptides: implications for brain functions and neurological disease*. New York: Raven Press, 1981:389-96.
159. Morley JE, Levine AS. Intraventricular cholecystokinin-octapeptide (CCK-8) produces hyperglycemia and hypothermia. *Clin Res*. 1980; 28:721A. abstract.
160. Carraway RE, Demers LM, Leeman SE. Hyperglycemia effect of neurotensin, a hypothalamic peptide. *Endocrinology*. 1976; 99:1452-62.
161. Blackburn AM, Fletcher DR, Adrian TE, Bloom SR. Neurotensin infusion in man. Pharmacokinetics and effect on gastrointestinal and pituitary hormones. *J Clin Endocrinol Metab*. 1980; 51:1257-61.
162. Straus E, Yalow RS. Cholecystokinin in the brains of obese and non-obese mice. *Science*. 1979; 203:68-9.
163. Schneider BS, Monahan JW, Hirsch J. Brain cholecystokinin and nutritional status in rats and mice. *J Clin Invest*. 1979; 64:1348-56.
164. Oku J, Glick Z, Shimomura Y, Inoue S, Bray GA, Walsh J. Cholecystokinin and obesity. *Clin Res*. 1980; 28:281A. abstract.
165. Ho P, Hansky J. Cholecystokinin (CCK)-like peptide in gut and brain of normal and genetically obese mice. *Gastroenterology*. 1979; 76:1155. abstract.
166. Rossier J, Rogers J, Shibasaki T, Guillemin R, Bloom FE. Opioid peptides and α -melanocyte-stimulating hormone in genetically obese (*ob/ob*) mice during development. *Proc Natl Acad Sci USA*. 1979; 76:2077-80.
167. Gunion MW, Peters RH. Pituitary beta-endorphin levels and naloxone effects on food intake in several obesity syndromes. *Soc Neurosci*. 1980; 6:527. abstract.
168. Gibson MJ, Krieger DT. Altered CNS-pituitary function in Zucker (*fa/fa*) fatty rats: absent circadian periodicity of activity, feeding, and plasma corticosterone (B) concentrations and elevated brain and pituitary neurointermediate (NI) lobe beta-endorphin (β -EP) concentrations. *Soc Neurosci*. 1980; 6:118. abstract.
169. Gamber SR, Garthwaite TL, Pontzer CH, Hagen TC. Fasting associated with decrease in hypothalamic β -endorphin. *Science*. 1980; 210:1271-2.
170. Hemmes RB, Pack HM, Hirsch J. Chronic ingestion of L-dopa dramatically reduces body weight of the genetically obese Zucker rat. *Fed Proc*. 1979; 38:277. abstract.
171. Benzinger TH. Heat regulation: homeostasis of central temperature in man. *Physiol Rev*. 1969; 49:671-759.
172. Nemeroff CB, Osbahr AJ III, Manberg PJ, Ervin GN, Prange AJ Jr. Alterations in nociception and body temperature after intracisternal administration of neurotensin, β -endorphin, other endogenous peptides, and morphine. *Proc Natl Acad Sci USA*. 1979; 76:5368-71.
173. Brown MR, Rivier JE, Kobayashi R, Vale WW. Neurotensin-like and bombesin-like peptides: CNS distribution and actions. In: Bloom SR, ed. *Gut hormones*. Edinburgh: Churchill-Livingstone, 1978:550-8.
174. Brown MR, Tache Y, Rivier J, Pittman G. Peptides and regulation of body temperature. In: Martin JB, Reichlin S, Bick KL, eds. *Neurosecretion and brain peptides: implications for brain functions and neurological disease*. New York: Raven Press, 1981:397-408.
175. Nemeroff CB, Bissette G, Manberg PJ, Osbahr AJ III, Breese GR, Prange AJ Jr. Neurotensin-induced hypothermia: evidence for an interaction with dopaminergic and the hypothalamic-pituitary-thyroid axis. *Brain Res*. 1980; 195:69-84.
176. Brown M, Rivier J, Vale W. Bombesin: potent effects on thermoregulation in the rat. *Science*. 1977; 196:998-1000.
177. Holaday JW, Tseng L-F, Loh HH, Li CH. Thyrotropin releasing hormone antagonizes β endorphin hypothermia and catalepsy. *Life Sci*. 1977; 22:1537-43.
178. Cooper KE, Kasting NW, Lederis K, Veale WL. Evidence supporting a role for endogenous vasopressin in natural suppression of fever in the sheep. *J Physiol (Lond)*. 1979; 295:33-45.
179. Myers RD, Tytell M. Fever: reciprocal shift in brain sodium to calcium ratio as the set-point temperature rises. *Science*. 1972; 178:765-7.
180. Hanagan JL, Williams BA. Brain calcium: role in temperature regulation. *Science*. 1973; 181:663-4.