

**Topic: Equation of a hyperbolic conic section****Question: Which function defines the same hyperbola?**

$$x = \frac{4}{\cos t} + 1$$

$$y = 2 \sin t \left( 1 + \tan t \tan \frac{t}{2} \right) - 1$$

**Answer choices:**

A  $\frac{(x-4)^2}{1^2} - \frac{(y+1)^2}{2^2} = 1$

B  $\frac{(x-1)^2}{2^2} - \frac{(y+1)^2}{4^2} = 1$

C  $\frac{(x-1)^2}{4^2} - \frac{(y+1)^2}{2^2} = 1$

D  $\frac{(x+1)^2}{4^2} - \frac{(y-1)^2}{2^2} = 1$



**Solution: C**

Rewrite both equations.

$$x = \frac{4}{\cos t} + 1$$

$$x - 1 = \frac{4}{\cos t}$$

$$\frac{(x - 1)^2}{4^2} = \frac{1}{\cos^2 t} = 1 + \tan^2 t$$

and

$$y = 2 \sin t \left( 1 + \tan t \tan \frac{t}{2} \right) - 1$$

$$y + 1 = 2 \sin t \left( 1 + \tan t \tan \frac{t}{2} \right)$$

$$\frac{y + 1}{2} = \sin t \left( 1 + \tan t \tan \frac{t}{2} \right)$$

$$\frac{y + 1}{2} = \tan t$$

$$\frac{(y + 1)^2}{2^2} = \tan^2 t$$

Subtract this second equation from the first.

$$\frac{(x - 1)^2}{4^2} - \frac{(y + 1)^2}{2^2} = 1 + \tan^2 t - \tan^2 t$$



$$\frac{(x-1)^2}{4^2} - \frac{(y+1)^2}{2^2} = 1$$



**Topic: Equation of a hyperbolic conic section****Question: Which parametric functions represent the hyperbola?**

$$3x^2 - y^2 + 6x + 4y - 10 = 0$$

**Answer choices:**

- A  $x = \sec t - 1$  and  $y = 3 \tan t - 2$
- B  $x = -1 + \sqrt{3} \sec t$  and  $y = 2 \tan t - 3$
- C  $x = \csc t - 1$  and  $y = 2 + 3 \tan t$
- D  $x = -1 + \sqrt{3} \sec t$  and  $y = 2 + 3 \tan t$



**Solution: D**

Complete the square with respect to both variables.

$$3x^2 - y^2 + 6x + 4y - 10 = 0$$

$$3(x^2 + 2x + 1) - (y^2 - 4y + 4) = 9$$

$$3(x + 1)^2 - (y - 2)^2 = 9$$

Divide out the coefficients.

$$\frac{(x + 1)^2}{3} - \frac{(y - 2)^2}{9} = \frac{9}{9}$$

$$\frac{(x + 1)^2}{3} - \frac{(y - 2)^2}{9} = 1$$

For a hyperbola in the form

$$\frac{(x - h)^2}{a^2} - \frac{(y - k)^2}{b^2} = 1$$

we can parametrize the curve using the equations  $x = h + a \sec t$  and  $y = k + b \tan t$ . In our case, we get

$$x = -1 + \sqrt{3} \sec t$$

and

$$y = 2 + 3 \tan t$$



**Topic:** Equation of a hyperbolic conic section**Question:** Which set of parametric equations represent the hyperbola?

$$4x^2 - y^2 - 8mx + 2ny + 4m^2 - n^2 = 16$$

**Answer choices:**

A  $x = \frac{2}{\cos t} - m$  and  $y = 4 \tan t - n$

B  $x = \frac{2}{\cos t} + m$  and  $y = 4 \tan t + n$

C  $x = \frac{4}{\cos t} + m$  and  $y = 4 \tan t + n$

D  $x = \frac{16}{\cos t} + m$  and  $y = 4 \tan t + n$



**Solution: B**

Choose the functions from answer choice B.

$$x = \frac{2}{\cos t} + m$$

$$y = 4 \tan t + n$$

Rewrite both equations.

$$x = \frac{2}{\cos t} + m$$

$$x - m = \frac{2}{\cos t}$$

$$\frac{x - m}{2} = \frac{1}{\cos t}$$

$$\frac{(x - m)^2}{2^2} = \frac{1}{\cos^2 t}$$

$$\frac{(x - m)^2}{2^2} = 1 + \tan^2 t$$

and

$$y = 4 \tan t + n$$

$$y - n = 4 \tan t$$

$$\frac{y - n}{4} = \tan t$$



$$\frac{(y - n)^2}{4^2} = \tan^2 t$$

Subtract this second equation from the first.

$$\frac{(x - m)^2}{4} - \frac{(y^2 - n)^2}{16} = 1 + \tan^2 t - \tan^2 t$$

$$\frac{(x - m)^2}{4} - \frac{(y^2 - n)^2}{16} = 1$$

Expand the equation above, and simplify.

$$\frac{x^2 - 2mx + m^2}{4} - \frac{y^2 - 2ny + n^2}{16} = 1$$

$$4(x^2 - 2mx + m^2) - y^2 - 2ny + n^2 = 16$$

$$4x^2 - 8mx + 4m^2 - y^2 - 2ny + n^2 = 16$$

$$4x^2 - y^2 - 8mx + 2ny + 4m^2 - n^2 = 16$$

