



Antiepileptic therapy with bromides— historical and actual importance

B.J. Steinhoff

To cite this article: B.J. Steinhoff (1992) Antiepileptic therapy with bromides— historical and actual importance, *Journal of the History of the Neurosciences*, 1:2, 119-123, DOI: [10.1080/09647049209525523](https://doi.org/10.1080/09647049209525523)

To link to this article: <http://dx.doi.org/10.1080/09647049209525523>



Published online: 23 Sep 2009.



Submit your article to this journal [↗](#)



Article views: 20



View related articles [↗](#)



Citing articles: 4 View citing articles [↗](#)

Antiepileptic therapy with bromides— historical and actual importance

B.J. Steinhoff

*Neurologische Klinik, Ludwig-Maximilians-Universität München, Klinikum
Großhadern, Germany*

THE ERA OF DOMINANCE

The introduction of potassium bromide as an antiepileptic drug (AED) by Sir Charles Locock in 1857 (see also Friedlander, 1986) was a milestone in the history of epileptology. Up to that time anticonvulsive treatments included such drugs as turpentine, silver salts, indigo, belladonna and chloroform (Donath, 1914) but without avail. Other remedies tried included *Adonis vernalis* (Bechterew, 1894), hyoscyanus (Daly, 1915), thyroid extract (Dercum, 1916), pancreatic extract (Riggs, 1916), sodium bilborate (Kimber, 1923), calcium lactate (Kaufmann, 1916), other calcium containing compounds (Ivanov, 1937), benzedrine, amytal and ephedrine (Minski & Gillen, 1937), again in vain (Steinhoff, 1988). Such dangerous and ineffective invasive methods as cranial trepanation were occasionally performed (Lennox, 1940), expressing the general helplessness of antiepileptic therapy.

Sir Charles Locock, chairman of the Royal Medical and Chirurgical Society in London and obstetrician to the Royal Family (Lennox, 1960), reported that the application of 30 grains of potassium bromide daily had completely suppressed epileptic seizures in 14 of 15 female patients (Locock, 1857). It was tried, because epileptic attacks were thought to be related to masturbation and hysteria, and the sedative and antiaphrodisiac properties of bromides were well known (Dreifuss, 1989).

Only one year later (1858) Radcliffe, Seé and Wilks definitively used bromides as AEDs (Steinhoff, 1988) and Wilks especially was responsible for its rapidly increasing use in the following years (Temkin, 1971).

The experience of some success at last in the treatment of epileptic disorders after such a long time of frustration and helplessness soon led to such enthusiastic comments as that of Radcliffe, only 4 years after Sir Charles Locock's first report: "The name of Sir Charles Locock ought to be remembered with gratitude by every epileptic" (Radcliffe, 1861).

Soon, not only in epileptology bromides became widely used. In the middle 1870s, the annual bromide consumption at the National Hospital London was 2.5 tons (Lennox, 1960). Not only potassium bromide, but other salts such as sodium, calcium, strontium, nickel, rubidium, rubidium ammonium, arsenic, zinc, gold sodium, caesium, or cadmium bromides were tried in antiepileptic treatment (Steinhoff, 1988). Although some authors tried to prove the superiority of some over others, there was no significant difference between the various compounds, due to the fact that it is the bromine component that is the effective antiepileptic factor (Lennox, 1960).

For 55 years, there had been no real alternative antiepileptic treatment. Chloralhydrate (Ulrich, 1914) and the ketogenic diet (Schoenborn, 1930) were the only antiepileptic therapeutic possibilities which are still of limited importance but used in the period between the appearance of bromides and phenobarbitone (PB).

Bromide therapy was optimized. Toulouse and Richet, for example, developed the concept of bromide application together with limited salt intake, so that the average dosage of bromides could be reduced without lowering its efficacy (Richet, 1903; Richet & Toulouse, 1899; Toulouse, 1900), because high chloride intake shortens the half life of bromides (Dreifuss, 1989). This was an important step since undesired side-effects had become obvious.

Critical comments regarding the side-effects of bromides became more frequent especially the chronic, sometimes even lethal, bromide intoxication. In 1905, for instance, Peterson mentioned dementia, intoxication, depression, mania and paranoia as typical undesirable effect of bromides. In spite of their remarkable antiepileptic efficacy he expressed the opinion that the epileptic patient would probably have been better off, if bromides had not been discovered.

THE APPEARANCE OF NEW ALTERNATIVE ANTIEPILEPTIC DRUGS

In 1912 Hauptmann introduced PB as a new AED with very good efficacy. Although bromides lost their importance, many authors as late as the 1930s, such as Rodhe, Winterseel, Sutherland, Szecsödy and Reitmann recommended them (Steinhoff, 1988).

With the appearance of phenytoin (PHT) in 1938 (Meritt & Putnam, 1938), antiepileptic therapy was enriched by a drug which was non-hypnotic. By 1940 Lennox suggested PHT as the only first-line AED. Table 1 shows the introduction of the main AEDs.

The introduction of carbamazepine (CBZ) and valproic acid (VPA) added further new drugs to worldwide accepted first-line treatments.

Bromides virtually vanished, mainly because of their side-effects and assumed lower efficacy compared with modern alternatives (Steinhoff, 1988). Twenty years ago, the Swiss epileptologist Schweingruber (1971) stated that bromides had no more than historical importance, and this was

Table 1. First announcement of antiepileptic efficacy of important antiepileptic drugs.

Bromides (BR)	1857 (Locock)
Phenobarbitone (PB)	1912 (Hauptmann)
Phenytoin (PHT)	1938 (Merrit & Putnam)
Primidone (PRM)	1952 (Richards & Everett)*
Ethosuximide (ESM)	1958 (Zimmermann & Burgemeister)*
Carbamazepine (CBZ)	1963 (Lorge)*
Valproic acid (VPA)	1964 (Meunier & Carraz)*

*See Kugler & Spatz (1984).

the common opinion of that time. But, this judgement was based either on case reports or non-controlled investigations and in some specialized epilepsy centres dealing with intractable epilepsies bromides were still used as add-on drugs in selected cases, mainly with generalized tonic-clonic seizures (GTCS) (Steinhoff, 1988).

THE COMEBACK

In spite of the use of modern first-line AEDs approximately 20–30% of all patients with epilepsies will not be completely seizure-free (Schmidt, 1988).

The observation of good bromide efficacy in selected cases encouraged the undertaking of a retrospective, controlled investigation (Steinhoff & Kruse, 1988). Antiepileptic treatment of 60 patients with GTCS, who had been treated with bromides alone (5 cases) or as an add-on drug, was analyzed. The results were compared with a very similar group of patients who had been on PHT and/or PB as basic medication. Surprisingly no significant difference was found regarding the antiepileptic efficacy and the frequency of side-effects between the bromide and the control groups. Of the 60 patients, 58% became free of GTCS or had a reduction of the GTCS-frequency of more than 50% without loss of efficacy for at least 3 years. All side-effects were reversible with dosage reduction or complete bromide withdrawal in a single case. Common side-effects were loss of appetite and weight, apathy, tiredness, psychomotor retardation and asynergic coordination. The success rate was 10% higher than in the control group. It was therefore claimed that there is an indication for bromides in every case of epilepsy with GTCS resistant to therapy with first line AEDs.

These observations are supported by various investigations of other epileptologists all published during recent years and reporting between 27% (Scheunemann, 1983) and 55% (Woody, 1990) improvement in the treatment of epilepsies with intractable seizures, so that several authors have very recently recommended bromides as an antiepileptic drug alter-

native (Boenigk *et al.*, 1985; Dreifuss & Bertram, 1986; Ernst *et al.*, 1988; Scheunemann, 1983; Steinhoff & Kruse, 1988; Woody, 1990).

The rather low rate of side-effects might be a result of the conditions of modern antiepileptic therapy with regular clinical and laboratory examinations of the patients, which allow the detection of undesired drug effects early enough to prevent serious complications.

CONCLUSION

More than 130 years after its introduction, bromide treatment in some patients with epilepsy of GTCS type has been re-established as an additional therapy in selected cases.

REFERENCES

- Bechterew W (1894): Über die Bedeutung des gleichzeitigen Gebrauchs der Bromide und des Adonis vernalis bei Epilepsie. *Neurologisches Centralblatt Leipzig* 13, 838–843.
- Boenigk HE, Lorenz J-H, Jürgens U (1985): Bromide – heute als antiepileptische Substanzen noch nützlich? *Nervenarzt* 56, 579–582.
- Davy AH (1915): Bromides in epilepsy. *Br Med J* 2, 692.
- Dercum FX (1916): Epilepsy with special reference to treatment. *JAMA* 67, 247–253.
- Donath J (1914): Sedobrol in der Behandlung der Epilepsie. *Wiener Klinische Wochenschrift* 27, 182–185.
- Dreifuss FE (1989): Bromides. In *Antiepileptic drugs*, ed R Levy, R Mattson, B Meldrum, JK Penry, FE Dreifuss, 3rd edn, New York: Raven Press.
- Dreifuss FE, Bertram EH (1986): Bromide therapy for intractable seizures. *Epilepsia* 27, 593 (Abstract).
- Ernst J-P, Dooze H, Baier WK (1988): Bromides were effective in intractable epilepsy with generalized tonic-clonic seizures and onset in early childhood. *Brain Dev* 10, 385–388.
- Friedlander WJ (1986): Who was the father of bromide treatment of epilepsy? *Arch Neurol* 43, 505–507.
- Hauptmann A (1912): Luminal bei Epilepsie. *Münch Med Wschr* 59, 1907.
- Ivanov V (1937): Erfahrungen mit Calcibronat bei Epilepsie. *Ärztliche Korrespondenz: Sonderdruck* 1–5.
- Kauffmann L (1916): Discussion of paper by Dercum, FX: Epilepsy with special reference to treatment. *JAMA* 67, 252.
- Kimber WJT (1923): Observations on the treatment of epilepsy with borax and bromide. *J Ment Sci* 114, 205–207.
- Kugler J, Spatz R (1984): *Spektrum Antiepileptika*. Zug: Aesopus Verlag Zug AG.
- Lennox WG (1940): Drug therapy (bromide, phenobarbital and sodium diphenyl hydantoin): the pharmacopeia and the physician. *JAMA* 114, 1347–1354.
- Lennox WG (1960): *Epilepsy and related disorders*. Boston: Little, Brown and Company.
- Locock C (1857): Discussion of paper by Sieveking, EH: Analysis of fifty-two cases of epilepsy observed by the author. *Lancet* 1, 528.

- Merritt HH, Putnam TJ (1938): Sodium diphenyl hydantoinate in the treatment of convulsive disorders. *JAMA* **111**, 1068.
- Minski L, Gillen JB (1937): Blood bromide investigations in psychic epileptics. *Br Med J* **2**, 850–851.
- Peterson F (1905): The bromides in epilepsy. *Am Med* **9**, 109.
- Radcliffe CB (1861): Epileptic and other convulsive affections of the nervous system—their pathology and treatment. 3rd edn, London: John Churchill.
- Richet C (1903): L'Hypochloruration dans le traitement de l'épilepsie par le bromure de potassium. *Comptes rendus des Séances de la Société de Biologie et des ses Filiales* **55**, 374.
- Richet C, Toulouse E (1899): Effets d'une alimentation pauvre en chlorures sur le traitement de l'épilepsie par le bromure de sodium. *Comptes rendus d'Académie des Sciences* **850**.
- Riggs CE (1916): Discussion of paper by Dercum, FX: Epilepsy with special reference to treatment. *JAMA* **67**, 251.
- Scheunemann W (1983): Brom – eine therapeutische Chance bei frühkindlicher Grand-mal-Epilepsie. In *Epilepsie* **83**, eds O Hallen, JG Meyer-Wahl, J Braun, Reinbek bei Hamburg: Einhorn Presse Verlag.
- Schmidt D (1988): Epilepsien. In *Therapie und Verlauf neurologischer Erkrankungen*, eds Th Brandt, J Dichgans, HC Diener, Stuttgart: Verlag W. Kohlhammer.
- Schoenborn S (1930): Luminal – oder Brombehandlung bei Epilepsie. *Fortschr Ther* **6**, 178–180.
- Schweingruber R (1971): 1st Brom als Antiepileptikum noch zeitgemäß? In *Deutsche Sektion der Internationalen Liga gegen Epilepsie: Rundbrief Nr. 40 der Deutschen Sektion der Internationalen*.
- Steinhoff B (1988): Zur antiepileptischen Therapie mit Bromiden: Darstellung in der Literatur – Röntgenfluoreszenzanalytische Bestimmung von Bromiden in Körperflüssigkeiten – Klinisch Ergebnisse der Behandlung von 60 Grand-mal-Epilepsien mit Bromiden. Medical thesis, University of Heidelberg.
- Steinhoff B, Kruse R (1988): Zur antiepileptischen Therapie mit Bromiden – eine klinische Studie. In *Epilepsie* **83**, ed P Wolf, Reinbeck bei Hamburg: Einhorn Presse Verlag.
- Temkin O (1971): *The falling sickness*, 2nd edn, Baltimore: The Johns Hopkins Press.
- Toulouse E (1900): *Traitement de l'épilepsie par les bromures et l'hypochloruration*. *Rev Psychiat* **3**, 1–14.
- Ulrich A (1914): Beitrag zur Technik der wirksamen Brombehandlung der Epilepsie. *Correspondenzblatt der Schweizer Ärzte* **44**, 641–652.
- Woody RC (1990): Bromide therapy for pediatric seizure disorder intractable to other antiepileptic drugs. *J Child Neurol* **5/1**, 65–67.