

Alternating Sign Matrices and Plane Partitions

Manjil P. Saikia

Cardiff University

11 October, 2022

ASMs

ASMs

An alternating sign matrix (ASM) of size n is an $n \times n$ matrix with entries in the set $\{0, 1, -1\}$

ASMs

An alternating sign matrix (ASM) of size n is an $n \times n$ matrix with entries in the set $\{0, 1, -1\}$ such that

- ▶ all row and column sums are equal to 1,

ASMs

An alternating sign matrix (ASM) of size n is an $n \times n$ matrix with entries in the set $\{0, 1, -1\}$ such that

- ▶ all row and column sums are equal to 1,
- ▶ and the non-zero entries alternate in each row and column.

ASMs

An alternating sign matrix (ASM) of size n is an $n \times n$ matrix with entries in the set $\{0, 1, -1\}$ such that

- ▶ all row and column sums are equal to 1,
- ▶ and the non-zero entries alternate in each row and column.

Clearly all permutation matrices are also ASMs.

ASMs

An alternating sign matrix (ASM) of size n is an $n \times n$ matrix with entries in the set $\{0, 1, -1\}$ such that

- ▶ all row and column sums are equal to 1,
- ▶ and the non-zero entries alternate in each row and column.

Clearly all permutation matrices are also ASMs.

For instance, there are 7 ASMs of order 3, these are the six permutation matrices

ASMs

An alternating sign matrix (ASM) of size n is an $n \times n$ matrix with entries in the set $\{0, 1, -1\}$ such that

- ▶ all row and column sums are equal to 1,
- ▶ and the non-zero entries alternate in each row and column.

Clearly all permutation matrices are also ASMs.

For instance, there are 7 ASMs of order 3, these are the six permutation matrices and the matrix

$$\begin{pmatrix} 0 & 1 & 0 \\ 1 & -1 & 1 \\ 0 & 1 & 0 \end{pmatrix}.$$

ASMs Enumeration

ASMs Enumeration

It was conjectured by Mills, Robbins and Rumsey that the number of ASMs of order n is

$$\prod_{j=0}^{n-1} \frac{(3j+1)!}{(n+j)!}.$$

ASMs Enumeration

It was conjectured by Mills, Robbins and Rumsey that the number of ASMs of order n is

$$\prod_{j=0}^{n-1} \frac{(3j+1)!}{(n+j)!}.$$

This conjecture was proved by Doron Zeilberger and independently by Greg Kuperberg.

ASMs Enumeration

It was conjectured by Mills, Robbins and Rumsey that the number of ASMs of order n is

$$\prod_{j=0}^{n-1} \frac{(3j+1)!}{(n+j)!}.$$

This conjecture was proved by Doron Zeilberger and independently by Greg Kuperberg.

Zeilberger's proved a stronger statement using constant term identities, while Kuperberg exploited a connection with models from statistical mechanics.

How were they first defined?

How were they first defined?

Given a matrix A , we let A_j^i denote the matrix that remains when the i th row and j th column of A are deleted.

How were they first defined?

Given a matrix A , we let A_j^i denote the matrix that remains when the i th row and j th column of A are deleted. If we remove more than one row or column, then the indices corresponding to those are added to the super- and sub- scripts.

How were they first defined?

Given a matrix A , we let A_j^i denote the matrix that remains when the i th row and j th column of A are deleted. If we remove more than one row or column, then the indices corresponding to those are added to the super- and sub- scripts.

Theorem (Desnanot-Jacobi adjoint matrix theorem)

If A is an $n \times n$ matrix, then

$$\det(A) \det(A_{1,n}^{1,n}) = \det(A_1^1) \det(A_n^n) - \det(A_n^1) \det(A_1^n)$$

How were they first defined?

Given a matrix A , we let A_j^i denote the matrix that remains when the i th row and j th column of A are deleted. If we remove more than one row or column, then the indices corresponding to those are added to the super- and sub- scripts.

Theorem (Desnanot-Jacobi adjoint matrix theorem)

If A is an $n \times n$ matrix, then

$$\det(A) \det(A_{1,n}^{1,n}) = \det(A_1^1) \det(A_n^n) - \det(A_n^1) \det(A_1^n)$$

or

How were they first defined?

Given a matrix A , we let A_j^i denote the matrix that remains when the i th row and j th column of A are deleted. If we remove more than one row or column, then the indices corresponding to those are added to the super- and sub- scripts.

Theorem (Desnanot-Jacobi adjoint matrix theorem)

If A is an $n \times n$ matrix, then

$$\det(A) \det(A_{1,n}^{1,n}) = \det(A_1^1) \det(A_n^n) - \det(A_n^1) \det(A_1^n)$$

or

$$\det(A) = \frac{1}{\det(A_{1,n}^{1,n})} \times \det \begin{pmatrix} \det(A_1^1) & \det(A_n^1) \\ \det(A_1^n) & \det(A_n^n) \end{pmatrix}.$$

How were they first defined?

Given a matrix A , we let A_j^i denote the matrix that remains when the i th row and j th column of A are deleted. If we remove more than one row or column, then the indices corresponding to those are added to the super- and sub- scripts.

Theorem (Desnanot-Jacobi adjoint matrix theorem)

If A is an $n \times n$ matrix, then

$$\det(A) \det(A_{1,n}^{1,n}) = \det(A_1^1) \det(A_n^n) - \det(A_n^1) \det(A_1^n)$$

or

$$\det(A) = \frac{1}{\det(A_{1,n}^{1,n})} \times \det \begin{pmatrix} \det(A_1^1) & \det(A_n^1) \\ \det(A_1^n) & \det(A_n^n) \end{pmatrix}.$$

This gives us a way of evaluating determinants, in terms of smaller determinants.

More determinants

More determinants

Reverend Charles L. Dodgson

More determinants

Reverend Charles L. Dodgson, better known by his pen name of Lewis Carroll used Desnanot-Jacobi theorem to give an algorithm for evaluating determinants in terms of 2×2 determinants.

More determinants

Reverend Charles L. Dodgson, better known by his pen name of Lewis Carroll used Desnanot-Jacobi theorem to give an algorithm for evaluating determinants in terms of 2×2 determinants.
For instance, we get

More determinants

Reverend Charles L. Dodgson, better known by his pen name of Lewis Carroll used Desnanot-Jacobi theorem to give an algorithm for evaluating determinants in terms of 2×2 determinants.

For instance, we get

$$\det \begin{pmatrix} a_{1,1} & a_{1,2} & a_{1,3} \\ a_{2,1} & a_{2,2} & a_{2,3} \\ a_{3,1} & a_{3,2} & a_{3,3} \end{pmatrix} = \frac{1}{a_{2,2}} \times \det \begin{pmatrix} \det \begin{pmatrix} a_{2,2} & a_{2,3} \\ a_{3,2} & a_{3,3} \end{pmatrix} & \det \begin{pmatrix} a_{2,1} & a_{2,2} \\ a_{3,1} & a_{3,2} \end{pmatrix} \\ \det \begin{pmatrix} a_{1,2} & a_{1,3} \\ a_{2,2} & a_{2,3} \end{pmatrix} & \det \begin{pmatrix} a_{1,1} & a_{1,2} \\ a_{2,1} & a_{2,2} \end{pmatrix} \end{pmatrix}.$$

Even more determinants

Even more determinants

$$\det \begin{pmatrix} a_{1,1} & a_{1,2} & a_{1,3} & a_{1,4} \\ a_{2,1} & a_{2,2} & a_{2,3} & a_{2,4} \\ a_{3,1} & a_{3,2} & a_{3,3} & a_{3,4} \\ a_{4,1} & a_{4,2} & a_{4,3} & a_{4,4} \end{pmatrix} = \frac{1}{\det \begin{pmatrix} a_{2,2} & a_{2,3} \\ a_{3,2} & a_{3,3} \end{pmatrix}}$$

$$\times \det \begin{pmatrix} \det \begin{pmatrix} a_{2,2} & a_{2,3} & a_{2,4} \\ a_{3,2} & a_{3,3} & a_{3,4} \\ a_{4,2} & a_{4,3} & a_{4,4} \end{pmatrix} & \det \begin{pmatrix} a_{2,1} & a_{2,2} & a_{2,3} \\ a_{3,1} & a_{3,2} & a_{3,3} \\ a_{4,1} & a_{4,2} & a_{4,3} \end{pmatrix} \\ \det \begin{pmatrix} a_{1,2} & a_{1,3} & a_{1,4} \\ a_{2,2} & a_{2,3} & a_{2,4} \\ a_{3,2} & a_{3,3} & a_{3,4} \end{pmatrix} & \det \begin{pmatrix} a_{1,1} & a_{1,2} & a_{1,3} \\ a_{2,1} & a_{2,2} & a_{2,3} \\ a_{3,1} & a_{3,2} & a_{3,3} \end{pmatrix} \end{pmatrix}.$$

Generalizing the determinant

Generalizing the determinant

In the 1980s, Robbins and Rumsey looked at a generalization of the 2×2 determinant

Generalizing the determinant

In the 1980s, Robbins and Rumsey looked at a generalization of the 2×2 determinant, which they called the λ -determinant.

Generalizing the determinant

In the 1980s, Robbins and Rumsey looked at a generalization of the 2×2 determinant, which they called the λ -determinant.

They defined

$$\det_{\lambda} \begin{pmatrix} a_{1,1} & a_{1,2} \\ a_{2,1} & a_{2,2} \end{pmatrix} = a_{1,1}a_{2,2} + \lambda a_{2,1}a_{1,2}.$$

Generalizing the determinant

In the 1980s, Robbins and Rumsey looked at a generalization of the 2×2 determinant, which they called the λ -determinant.

They defined

$$\det_{\lambda} \begin{pmatrix} a_{1,1} & a_{1,2} \\ a_{2,1} & a_{2,2} \end{pmatrix} = a_{1,1}a_{2,2} + \lambda a_{2,1}a_{1,2}.$$

Using the previous observations, they generalized it to an $n \times n$ determinant.

ASMs and Determinants

ASMs and Determinants

Their main result in this direction was

ASMs and Determinants

Their main result in this direction was

Theorem (Robbins-Rumsey)

Let A be an $n \times n$ matrix with entries $a_{i,j}$, \mathcal{A}_n be the set of all ASMs, $\mathcal{I}(B)$ be the inversion number of B and $\mathcal{N}(B)$ be the number of -1 's in B . Then

$$\det_{\lambda}(A) = \sum_{B \in \mathcal{A}_n} \lambda^{\mathcal{I}(B)} (1 + \lambda^{-1})^{\mathcal{N}(B)} \prod_{i,j=1}^n a_{i,j}^{B_{i,j}}.$$

ASMs and Determinants

Their main result in this direction was

Theorem (Robbins-Rumsey)

Let A be an $n \times n$ matrix with entries $a_{i,j}$, \mathcal{A}_n be the set of all ASMs, $\mathcal{I}(B)$ be the inversion number of B and $\mathcal{N}(B)$ be the number of -1 's in B . Then

$$\det_{\lambda}(A) = \sum_{B \in \mathcal{A}_n} \lambda^{\mathcal{I}(B)} (1 + \lambda^{-1})^{\mathcal{N}(B)} \prod_{i,j=1}^n a_{i,j}^{B_{i,j}}.$$

This was the first appearance of an ASM in the literature.

What's the inversion number?

What's the inversion number?

An easy way to calculate the inversion number is to take products of all pairs of entries for which one of them lies to the right and above the other, and then adding them all up.

What's the inversion number?

An easy way to calculate the inversion number is to take products of all pairs of entries for which one of them lies to the right and above the other, and then adding them all up.

$$\begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 1 & -1 & 0 & 0 & 1 \\ 0 & 1 & -1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{pmatrix}$$

What's the inversion number?

An easy way to calculate the inversion number is to take products of all pairs of entries for which one of them lies to the right and above the other, and then adding them all up.

$$\begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 1 & -1 & 0 & 0 & 1 \\ 0 & 1 & -1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{pmatrix}$$

There are seven pairs here whose product is $+1$ and two pairs whose product is -1 .

What's the inversion number?

An easy way to calculate the inversion number is to take products of all pairs of entries for which one of them lies to the right and above the other, and then adding them all up.

$$\begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 1 & -1 & 0 & 0 & 1 \\ 0 & 1 & -1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{pmatrix}$$

There are seven pairs here whose product is $+1$ and two pairs whose product is -1 . So the inversion number is 5 .

How does one get a formula from this?

How does one get a formula from this?

First some observations

How does one get a formula from this?

First some observations

$$\begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & -1 & 0 & 1 \\ 0 & 1 & 0 & 0 \end{pmatrix}$$

There can be only one **1** in the top row (or, first column).

How does one get a formula from this?

First some observations

$$\begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & -1 & 0 & 1 \\ 0 & 1 & 0 & 0 \end{pmatrix}$$

There can be only one **1** in the top row (or, first column). Let $A_{n,k}$ be the number of $n \times n$ ASMs with a **1** at the top of top k th column.

How does one get a formula from this?

First some observations

$$\begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & -1 & 0 & 1 \\ 0 & 1 & 0 & 0 \end{pmatrix}$$

There can be only one **1** in the top row (or, first column). Let $A_{n,k}$ be the number of $n \times n$ ASMs with a **1** at the top of top k th column. Some thought will give us, $A_{n,k} = A_{n,n+1-k}$

How does one get a formula from this?

First some observations

$$\begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & -1 & 0 & 1 \\ 0 & 1 & 0 & 0 \end{pmatrix}$$

There can be only one **1** in the top row (or, first column). Let $A_{n,k}$ be the number of $n \times n$ ASMs with a **1** at the top of top k th column. Some thought will give us, $A_{n,k} = A_{n,n+1-k}$ (symmetry).

How does one get a formula from this?

First some observations

$$\begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & -1 & 0 & 1 \\ 0 & 1 & 0 & 0 \end{pmatrix}$$

There can be only one **1** in the top row (or, first column). Let $A_{n,k}$ be the number of $n \times n$ ASMs with a **1** at the top of top k th column. Some thought will give us, $A_{n,k} = A_{n,n+1-k}$ (symmetry).

Further, if A_n is the number of $n \times n$ ASMs, then $A_{n,1} = A_{n,n} = A_{n-1}$.

Guessing the formula

Guessing the formula

This allows one to check small values to get a formula.

Guessing the formula

This allows one to check small values to get a formula. Mills, Robbins and Rumsey did exactly that.

Guessing the formula

This allows one to check small values to get a formula. Mills, Robbins and Rumsey did exactly that.

They first conjectured

Guessing the formula

This allows one to check small values to get a formula. Mills, Robbins and Rumsey did exactly that.

They first conjectured

$$\frac{A_{n,k}}{A_{n,k+1}} = \frac{k(2n - k - 1)}{(n - k)(n + k - 1)}.$$

Guessing the formula

This allows one to check small values to get a formula. Mills, Robbins and Rumsey did exactly that.

They first conjectured

$$\frac{A_{n,k}}{A_{n,k+1}} = \frac{k(2n - k - 1)}{(n - k)(n + k - 1)}.$$

This means that the $A_{n,k}$'s are uniquely determined by the $A_{n,k-1}$'s when $k > 1$ and by $A_{n,1} = \sum_{k=1}^{n-1} A_{n-1,k}$.

Towards a final formula

Towards a final formula

The above conjecture can be reformulated as

Towards a final formula

The above conjecture can be reformulated as

$$A_{n,k} = \binom{n+k-2}{k-1} \frac{(2n-k-1)!}{(n-k)!} \prod_{j=0}^{n-2} \frac{(3j+1)!}{(n+j)!}.$$

Towards a final formula

The above conjecture can be reformulated as

$$A_{n,k} = \binom{n+k-2}{k-1} \frac{(2n-k-1)!}{(n-k)!} \prod_{j=0}^{n-2} \frac{(3j+1)!}{(n+j)!}.$$

From here, knowing that $A_n = A_{n+1,1}$ allows one to conjecture the ASM enumeration formula.

The Many Faces of ASMs

The Many Faces of ASMs

Theorem (Andrews, Zeilberger, Ayer-Behrend-Fischer)

The following combinatorial objects are counted by the formula

$$\prod_{j=0}^{n-1} \frac{(3j+1)!}{(n+j)!}$$

The Many Faces of ASMs

Theorem (Andrews, Zeilberger, Ayer-Behrend-Fischer)

The following combinatorial objects are counted by the formula

$$\prod_{j=0}^{n-1} \frac{(3j+1)!}{(n+j)!}$$

► *Alternating Sign Matrices,*

The Many Faces of ASMs

Theorem (Andrews, Zeilberger, Ayer-Behrend-Fischer)

The following combinatorial objects are counted by the formula

$$\prod_{j=0}^{n-1} \frac{(3j+1)!}{(n+j)!}$$

- ▶ *Alternating Sign Matrices,*
- ▶ *Descending Plane Partitions,*

The Many Faces of ASMs

Theorem (Andrews, Zeilberger, Ayer-Behrend-Fischer)

The following combinatorial objects are counted by the formula

$$\prod_{j=0}^{n-1} \frac{(3j+1)!}{(n+j)!}$$

- ▶ *Alternating Sign Matrices,*
- ▶ *Descending Plane Partitions,*
- ▶ *Totally Symmetric Self-Complementary Plane Partitions,*

The Many Faces of ASMs

Theorem (Andrews, Zeilberger, Ayer-Behrend-Fischer)

The following combinatorial objects are counted by the formula

$$\prod_{j=0}^{n-1} \frac{(3j+1)!}{(n+j)!}$$

- ▶ *Alternating Sign Matrices,*
- ▶ *Descending Plane Partitions,*
- ▶ *Totally Symmetric Self-Complementary Plane Partitions,*
- ▶ *Alternating Sign Triangles,*

The Many Faces of ASMs

Theorem (Andrews, Zeilberger, Ayer-Behrend-Fischer)

The following combinatorial objects are counted by the formula

$$\prod_{j=0}^{n-1} \frac{(3j+1)!}{(n+j)!}$$

- ▶ *Alternating Sign Matrices,*
- ▶ *Descending Plane Partitions,*
- ▶ *Totally Symmetric Self-Complementary Plane Partitions,*
- ▶ *Alternating Sign Triangles,*
- ▶ *...and a couple more.*

Plane Partitions

Plane Partitions

A *plane partition* in an $a \times b \times c$ box is a subset

$$PP \subset \{1, 2, \dots, a\} \times \{1, 2, \dots, b\} \times \{1, 2, \dots, c\}$$


with $(i', j', k') \in PP$ if $(i, j, k) \in PP$ and $(i', j', k') \leq (i, j, k)$.

Plane Partitions

A *plane partition* in an $a \times b \times c$ box is a subset

$$PP \subset \{1, 2, \dots, a\} \times \{1, 2, \dots, b\} \times \{1, 2, \dots, c\}$$

with $(i', j', k') \in PP$ if $(i, j, k) \in PP$ and $(i', j', k') \leq (i, j, k)$.



Totally Symmetric Self-Complementary Plane Partitions

Totally Symmetric Self-Complementary Plane Partitions

If a plane partition has all the symmetries (that is, $(i, j, k) \in PP$ if and only if all six permutations of (i, j, k) are also in PP)

Totally Symmetric Self-Complementary Plane Partitions


If a plane partition has all the symmetries (that is, $(i, j, k) \in PP$ if and only if all six permutations of (i, j, k) are also in PP) and is its own complement (that is, if $(i, j, k) \in PP$ then $(2n + 1 - i, 2n + 1 - j, 2n + 1 - k) \notin PP$)

Totally Symmetric Self-Complementary Plane Partitions

If a plane partition has all the symmetries (that is, $(i, j, k) \in PP$ if and only if all six permutations of (i, j, k) are also in PP) and is its own complement (that is, if $(i, j, k) \in PP$ then $(2n + 1 - i, 2n + 1 - j, 2n + 1 - k) \notin PP$), then it is called *totally symmetric self-complementary plane partitions* (TSSCPP).


Totally Symmetric Self-Complementary Plane Partitions

If a plane partition has all the symmetries (that is, $(i, j, k) \in PP$ if and only if all six permutations of (i, j, k) are also in PP) and is its own complement (that is, if $(i, j, k) \in PP$ then $(2n+1-i, 2n+1-j, 2n+1-k) \notin PP$), then it is called *totally symmetric self-complementary plane partitions* (TSSCPP).



Totally Symmetric Self-Complementary Plane Partitions

If a plane partition has all the symmetries (that is, $(i, j, k) \in PP$ if and only if all six permutations of (i, j, k) are also in PP) and is its own complement (that is, if $(i, j, k) \in PP$ then $(2n+1-i, 2n+1-j, 2n+1-k) \notin PP$), then it is called *totally symmetric self-complementary plane partitions* (TSSCPP).



The class of TSSCPPs inside a $2n \times 2n \times 2n$ box are equinumerous with $n \times n$ ASMs.

Does the story end here?

Does the story end here?

In the late 1980's Richard Stanley suggested the study of various symmetry classes of ASMs

Does the story end here?

In the late 1980's Richard Stanley suggested the study of various symmetry classes of ASMs; this let Robbins to conjecture formulas for many of these classes.

Does the story end here?

In the late 1980's Richard Stanley suggested the study of various symmetry classes of ASMs; this let Robbins to conjecture formulas for many of these classes.

It turned out to be as difficult as enumerating ASMs,

Does the story end here?

In the late 1980's Richard Stanley suggested the study of various symmetry classes of ASMs; this let Robbins to conjecture formulas for many of these classes.

It turned out to be as difficult as enumerating ASMs, and this study was only recently completed in 2017.

Symmetry Classes

Symmetry Classes

- ▶ Vertically Symmetric ASMs: $a_{i,j} = a_{i,n+1-j}$, n odd (Kuperberg 2002)

Symmetry Classes

- ▶ Vertically Symmetric ASMs: $a_{i,j} = a_{i,n+1-j}$, n odd (Kuperberg 2002)
- ▶ Half-turn Symmetric ASMs: $a_{i,j} = a_{n+1-i,n+1-j}$, n odd (Razumov-Stroganov 2005), n even (Kuperberg 2002)

Symmetry Classes

- ▶ Vertically Symmetric ASMs: $a_{i,j} = a_{i,n+1-j}$, n odd (Kuperberg 2002)
- ▶ Half-turn Symmetric ASMs: $a_{i,j} = a_{n+1-i,n+1-j}$, n odd (Razumov-Stroganov 2005), n even (Kuperberg 2002)
- ▶ Diagonally Symmetric ASMs: $a_{i,j} = a_{j,i}$, no 'nice' formula (Behrend, Fischer & Koutschan, 2022+)

Symmetry Classes

- ▶ Vertically Symmetric ASMs: $a_{i,j} = a_{i,n+1-j}$, n odd (Kuperberg 2002)
- ▶ Half-turn Symmetric ASMs: $a_{i,j} = a_{n+1-i,n+1-j}$, n odd (Razumov-Stroganov 2005), n even (Kuperberg 2002)
- ▶ Diagonally Symmetric ASMs: $a_{i,j} = a_{j,i}$, no 'nice' formula (Behrend, Fischer & Koutschan, 2022+)
- ▶ Quarter-turn Symmetric ASMs: $a_{i,j} = a_{j,n+1-i}$, n odd (Razumov-Stroganov 2005), n even (Kuperberg 2002)

Symmetry Classes

- ▶ Vertically Symmetric ASMs: $a_{i,j} = a_{i,n+1-j}$, n odd (Kuperberg 2002)
- ▶ Half-turn Symmetric ASMs: $a_{i,j} = a_{n+1-i,n+1-j}$, n odd (Razumov-Stroganov 2005), n even (Kuperberg 2002)
- ▶ Diagonally Symmetric ASMs: $a_{i,j} = a_{j,i}$, no 'nice' formula (Behrend, Fischer & Koutschan, 2022+)
- ▶ Quarter-turn Symmetric ASMs: $a_{i,j} = a_{j,n+1-i}$, n odd (Razumov-Stroganov 2005), n even (Kuperberg 2002)
- ▶ Horizontally and vertically Symmetric ASMs:
 $a_{i,j} = a_{i,n+1-j} = a_{n+1-i,j}$, n odd (Okada 2004)

Symmetry Classes

- ▶ Vertically Symmetric ASMs: $a_{i,j} = a_{i,n+1-j}$, n odd (Kuperberg 2002)
- ▶ Half-turn Symmetric ASMs: $a_{i,j} = a_{n+1-i,n+1-j}$, n odd (Razumov-Stroganov 2005), n even (Kuperberg 2002)
- ▶ Diagonally Symmetric ASMs: $a_{i,j} = a_{j,i}$, no 'nice' formula (Behrend, Fischer & Koutschan, 2022+)
- ▶ Quarter-turn Symmetric ASMs: $a_{i,j} = a_{j,n+1-i}$, n odd (Razumov-Stroganov 2005), n even (Kuperberg 2002)
- ▶ Horizontally and vertically Symmetric ASMs:
 $a_{i,j} = a_{i,n+1-j} = a_{n+1-i,j}$, n odd (Okada 2004)
- ▶ Diagonally and Antidiagonally Symmetric ASMs:
 $a_{i,j} = a_{j,i} = a_{n+1-j,n+1-i}$, n odd (Behrend, Fischer & Konvalinka 2017)

Symmetry Classes

- ▶ Vertically Symmetric ASMs: $a_{i,j} = a_{i,n+1-j}$, n odd (Kuperberg 2002)
- ▶ Half-turn Symmetric ASMs: $a_{i,j} = a_{n+1-i,n+1-j}$, n odd (Razumov-Stroganov 2005), n even (Kuperberg 2002)
- ▶ Diagonally Symmetric ASMs: $a_{i,j} = a_{j,i}$, no 'nice' formula (Behrend, Fischer & Koutschan, 2022+)
- ▶ Quarter-turn Symmetric ASMs: $a_{i,j} = a_{j,n+1-i}$, n odd (Razumov-Stroganov 2005), n even (Kuperberg 2002)
- ▶ Horizontally and vertically Symmetric ASMs:
 $a_{i,j} = a_{i,n+1-j} = a_{n+1-i,j}$, n odd (Okada 2004)
- ▶ Diagonally and Antidiagonally Symmetric ASMs:
 $a_{i,j} = a_{j,i} = a_{n+1-j,n+1-i}$, n odd (Behrend, Fischer & Konvalinka 2017)
- ▶ All symmetries: $a_{i,j} = a_{j,i} = a_{i,n+1-j}$, no 'nice' formula.



Refined Enumeration of ASMs

Refined Enumeration of ASMs

Some observations are in order.

Refined Enumeration of ASMs

Some observations are in order.

$$\begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 & 0 & 1 & 0 \\ 1 & -1 & 0 & 1 & 0 & -1 & 1 \\ 0 & 0 & 1 & -1 & 1 & 0 & 0 \\ 0 & 1 & -1 & 1 & -1 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}$$

Refined Enumeration of ASMs

Some observations are in order.

$$\begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 & 0 & 1 & 0 \\ 1 & -1 & 0 & 1 & 0 & -1 & 1 \\ 0 & 0 & 1 & -1 & 1 & 0 & 0 \\ 0 & 1 & -1 & 1 & -1 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}$$

- There is only one 1 in any boundary row/column of an ASM.

Refined Enumeration of ASMs

Some observations are in order.

$$\begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 & 0 & 1 & 0 \\ 1 & -1 & 0 & 1 & 0 & -1 & 1 \\ 0 & 0 & 1 & -1 & 1 & 0 & 0 \\ 0 & 1 & -1 & 1 & -1 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}$$

- ▶ There is only one 1 in any boundary row/column of an ASM.
- ▶ This suggests the question: how many ASMs with the position of the 1 fixed at a certain row/column exist?

Refined Enumeration of ASMs

Some observations are in order.

$$\begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 & 0 & 1 & 0 \\ 1 & -1 & 0 & 1 & 0 & -1 & 1 \\ 0 & 0 & 1 & -1 & 1 & 0 & 0 \\ 0 & 1 & -1 & 1 & -1 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}$$

- ▶ There is only one 1 in any boundary row/column of an ASM.
- ▶ This suggests the question: how many ASMs with the position of the 1 fixed at a certain row/column exist?
- ▶ These are called refined enumeration of ASMs.

Refined Enumeration of ASMs

Refined Enumeration of ASMs

The study began with conjectures by Robbins about the number of ASMs of order n with the position of the 1 in the first row at the k th column is given by

Refined Enumeration of ASMs

The study began with conjectures by Robbins about the number of ASMs of order n with the position of the 1 in the first row at the k th column is given by

$$A_{n,k} = \binom{n+k-2}{k-1} \frac{(2n-k-1)!}{(n-k)!} \prod_{j=0}^{n-2} \frac{3j+1)!}{(n+j)!}.$$

Refined Enumeration of ASMs

The study began with conjectures by Robbins about the number of ASMs of order n with the position of the 1 in the first row at the k th column is given by

$$A_{n,k} = \binom{n+k-2}{k-1} \frac{(2n-k-1)!}{(n-k)!} \prod_{j=0}^{n-2} \frac{3j+1)!}{(n+j)!}.$$

This was proved by Zeilberger (1996).

Refined Enumeration of ASMs

The study began with conjectures by Robbins about the number of ASMs of order n with the position of the 1 in the first row at the k th column is given by

$$A_{n,k} = \binom{n+k-2}{k-1} \frac{(2n-k-1)!}{(n-k)!} \prod_{j=0}^{n-2} \frac{3j+1!}{(n+j)!}.$$

This was proved by Zeilberger (1996).

Several people have worked on the refined enumeration of ASMs as well as their symmetry classes

Refined Enumeration of ASMs

The study began with conjectures by Robbins about the number of ASMs of order n with the position of the 1 in the first row at the k th column is given by

$$A_{n,k} = \binom{n+k-2}{k-1} \frac{(2n-k-1)!}{(n-k)!} \prod_{j=0}^{n-2} \frac{3j+1!}{(n+j)!}.$$

This was proved by Zeilberger (1996).

Several people have worked on the refined enumeration of ASMs as well as their symmetry classes: Behrend, Fischer, Romik, Razumov-Stroganov, Ayyer-Romik, Romik-Karlinisky, etc.

Some (ex-) conjectures

Some (ex-) conjectures

Several conjectures on refined enumeration of ASMs existed.

Some (ex-) conjectures

Several conjectures on refined enumeration of ASMs existed.

- ▶ Fischer (2009) conjectured a formula for the number of VSASMs with the position of the 1's in the second row fixed.

Some (ex-) conjectures

Several conjectures on refined enumeration of ASMs existed.

- ▶ Fischer (2009) conjectured a formula for the number of VSASMs with the position of the 1's in the second row fixed.
- ▶ Robbins (late 1980s) conjectured formulas for refined enumeration of QTSASMs.

Some (ex-) conjectures

Several conjectures on refined enumeration of ASMs existed.

- ▶ Fischer (2009) conjectured a formula for the number of VSASMs with the position of the 1's in the second row fixed.
- ▶ Robbins (late 1980s) conjectured formulas for refined enumeration of QTSASMs.
- ▶ Duchon (2008) conjectured a formula for the refined enumeration of quasi-QTSASMs.

Some (ex-) conjectures

Several conjectures on refined enumeration of ASMs existed.

- ▶ Fischer (2009) conjectured a formula for the number of VSASMs with the position of the 1's in the second row fixed.
- ▶ Robbins (late 1980s) conjectured formulas for refined enumeration of QTSASMs.
- ▶ Duchon (2008) conjectured a formula for the refined enumeration of quasi-QTSASMs.
- ▶ We proved these conjectures – and more – in joint work with Ilse Fischer (2020).

VSASMs

VSASMs

In the case of VSASMs, we can make the following observations:

VSASMs

In the case of VSASMs, we can make the following observations:

- ▶ They exist for odd order.

VSASMs

In the case of VSASMs, we can make the following observations:

- ▶ They exist for odd order.
- ▶ The middle column is always $(1, -1, 1, \dots, -1, 1)^T$.

VSASMs

In the case of VSASMs, we can make the following observations:

- ▶ They exist for odd order.
- ▶ The middle column is always $(1, -1, 1, \dots, -1, 1)^T$.
- ▶ The second row has exactly two 1's.

$$\begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 & 0 & 1 & 0 \\ 1 & -1 & 0 & 1 & 0 & -1 & 1 \\ 0 & 0 & 1 & -1 & 1 & 0 & 0 \\ 0 & 1 & -1 & 1 & -1 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}$$

VSASMs

In the case of VSASMs, we can make the following observations:

- ▶ They exist for odd order.
- ▶ The middle column is always $(1, -1, 1, \dots, -1, 1)^T$.
- ▶ The second row has exactly two 1's.

$$\begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 & 0 & 1 & 0 \\ 1 & -1 & 0 & 1 & 0 & -1 & 1 \\ 0 & 0 & 1 & -1 & 1 & 0 & 0 \\ 0 & 1 & -1 & 1 & -1 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}$$

So we can ask for refined enumeration w.r.t. the position of the 1's in the second row.

Fischer's Conjecture

Fischer's Conjecture

Razumov and Stroganov (2004) has a formula counting the number of VSASMs with a fixed one in the first column.

$$A_{VC}(2n+1, i) = \frac{1}{2^{n-1}} \prod_{k=1}^{n-1} \frac{(6k-2)!(2k-1)!}{(4k-1)!(4k-2)!} \\ \times \sum_{k=1}^{i-1} (-1)^{i+k-1} \frac{(2n+k-2)!(4n-k-1)!}{(4n-2)!(k-1)!(2n-k)!}$$

Fischer's Conjecture

Razumov and Stroganov (2004) has a formula counting the number of VSASMs with a fixed one in the first column.

$$A_{VC}(2n+1, i) = \frac{1}{2^{n-1}} \prod_{k=1}^{n-1} \frac{(6k-2)!(2k-1)!}{(4k-1)!(4k-2)!} \\ \times \sum_{k=1}^{i-1} (-1)^{i+k-1} \frac{(2n+k-2)!(4n-k-1)!}{(4n-2)!(k-1)!(2n-k)!}$$

Ilse Fischer (2007) had conjectured that the number of $(2n+1) \times (2n+1)$ VSASMs, where the first one in the second row is in the i th column is equal to

Fischer's Conjecture

Razumov and Stroganov (2004) has a formula counting the number of VSASMs with a fixed one in the first column.

$$A_{VC}(2n+1, i) = \frac{1}{2^{n-1}} \prod_{k=1}^{n-1} \frac{(6k-2)!(2k-1)!}{(4k-1)!(4k-2)!} \\ \times \sum_{k=1}^{i-1} (-1)^{i+k-1} \frac{(2n+k-2)!(4n-k-1)!}{(4n-2)!(k-1)!(2n-k)!}$$

Ilse Fischer (2007) had conjectured that the number of $(2n+1) \times (2n+1)$ VSASMs, where the first one in the second row is in the i th column is equal to

$$\frac{(2n+i-2)!(4n-i-1)!}{2^{n-1}(4n-2)!(i-1)!(2n-i)!} \left(\prod_{j=1}^{n-1} \frac{(6j-2)!(2j-1)!}{(4j-1)!(4j-2)!} \right).$$

Fischer's Conjecture

Fischer's Conjecture

Theorem (Fischer-S., 2020)

The number of $(2n+1) \times (2n+1)$ VSASM with a 1 in the i -th position in it's second row is given by

$$\frac{(2n+i-2)!(4n-i-1)!}{2^{n-1}(4n-2)!(i-1)!(2n-i)!} \left(\prod_{j=1}^{n-1} \frac{(6j-2)!(2j-1)!}{(4j-1)!(4j-2)!} \right) \\ = A_{VC}(2n+1, i) + A_{VC}(2n+1, i+1).$$

Fischer's Conjecture

Theorem (Fischer-S., 2020)

The number of $(2n+1) \times (2n+1)$ VSASM with a 1 in the i -th position in it's second row is given by

$$\frac{(2n+i-2)!(4n-i-1)!}{2^{n-1}(4n-2)!(i-1)!(2n-i)!} \left(\prod_{j=1}^{n-1} \frac{(6j-2)!(2j-1)!}{(4j-1)!(4j-2)!} \right) \\ = A_{VC}(2n+1, i) + A_{VC}(2n+1, i+1).$$

A bijective proof of the last relation would be of interest.

Other Symmetry Classes?

Other Symmetry Classes?

- ▶ We have formulas for many of the symmetry classes as well as other type of ASMs.

Other Symmetry Classes?

- ▶ We have formulas for many of the symmetry classes as well as other type of ASMs.
- ▶ In some cases the results are in terms of generating functions.

Other Symmetry Classes?

- ▶ We have formulas for many of the symmetry classes as well as other type of ASMs.
- ▶ In some cases the results are in terms of generating functions.

For an ASM or quasi-ASM A in a class $XASM$, let

$\Theta_{XASM}(A)$ = number of -1 's in the fundamental region of A ,

$\psi_T^i(A)$ = (position of the 1 in the top row of A) $- i$,

$\overline{\psi}_T^i(A)$ = (position of the first 1 in the second-top row of A) $- i$,

$\psi_L^i(A)$ = (position of the 1 in the leftmost column of A) $- i$,

Generating Functions

Generating Functions

The definitions of certain ASM generating functions are then

$$Z_{2n+1}^{\text{VSASM}}(x, z_1, z_2) = \sum_{A \in \text{VSASM}(2n+1)} x^{\Theta_{\text{VSASM}}(A)} z_1^{\Psi_L^2(A)} z_2^{\bar{\Psi}_T^1(A)},$$

$$Z_{2n+1}^{\text{HVSASM}}(x, z) = \sum_{A \in \text{HVSASM}(2n+1)} x^{\Theta_{\text{HVSASM}}(A)} z^{\bar{\Psi}_T^2(A)},$$

$$Z_{2n+1}^{\text{HVPASM}}(x, z_1, z_2) = \sum_{A \in \text{HVPASM}(2n+1)} x^{\Theta_{\text{HVPASM}}(A)} z_1^{\Psi_T^1(A)} z_2^{\Psi_L^1(A)}.$$

Generating Functions

The definitions of certain ASM generating functions are then

$$Z_{2n+1}^{\text{VSASM}}(x, z_1, z_2) = \sum_{A \in \text{VSASM}(2n+1)} x^{\Theta_{\text{VSASM}}(A)} z_1^{\Psi_L^2(A)} z_2^{\bar{\Psi}_T^1(A)},$$

$$Z_{2n+1}^{\text{HVSASM}}(x, z) = \sum_{A \in \text{HVSASM}(2n+1)} x^{\Theta_{\text{HVSASM}}(A)} z^{\bar{\Psi}_T^2(A)},$$

$$Z_{2n+1}^{\text{HVPASM}}(x, z_1, z_2) = \sum_{A \in \text{HVPASM}(2n+1)} x^{\Theta_{\text{HVPASM}}(A)} z_1^{\Psi_T^1(A)} z_2^{\Psi_L^1(A)}.$$

We have not yet defined some classes of ASMs.

Fischer's Conjecture

Fischer's Conjecture

We can now reformulate our result in a succinct manner.

Fischer's Conjecture

We can now reformulate our result in a succinct manner.

Theorem (Fischer-S., 2020)

We have for all $n \geq 1$,

$$(1+z) Z_{2n+1}^{\text{VSASM}}(1, z, 1) = Z_{2n+1}^{\text{VSASM}}(1, 1, z) + z^{2n-1} Z_{2n+1}^{\text{VSASM}}(1, 1, \bar{z}).$$

Fischer's Conjecture

We can now reformulate our result in a succinct manner.

Theorem (Fischer-S., 2020)

We have for all $n \geq 1$,

$$(1+z) Z_{2n+1}^{\text{VSASM}}(1, z, 1) = Z_{2n+1}^{\text{VSASM}}(1, 1, z) + z^{2n-1} Z_{2n+1}^{\text{VSASM}}(1, 1, \bar{z}).$$

Recently, we proved a generalization of this.

Fischer's Conjecture

We can now reformulate our result in a succinct manner.

Theorem (Fischer-S., 2020)

We have for all $n \geq 1$,

$$(1+z) Z_{2n+1}^{\text{VSASM}}(1, z, 1) = Z_{2n+1}^{\text{VSASM}}(1, 1, z) + z^{2n-1} Z_{2n+1}^{\text{VSASM}}(1, 1, \bar{z}).$$

Recently, we proved a generalization of this.

Theorem (Behrend-S., 2022)

We have for all $n \geq 1$,

$$(1+z) Z_{2n+1}^{\text{VSASM}}(x, z, 1) = Z_{2n+1}^{\text{VSASM}}(x, 1, z) + z^{2n-1} Z_{2n+1}^{\text{VSASM}}(x, 1, \bar{z}).$$

Vertically and Horizontally Symmetric ASMs

Vertically and Horizontally Symmetric ASMs

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & -1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & -1 & 0 & 0 & 1 & 0 \\ 1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 & 1 \\ 0 & 1 & 0 & 0 & -1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & -1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

VHSASMs

VHSASMs

Theorem (Fischer-S., 2020)

We have for all $n \geq 1$,

$$\begin{aligned} Z_{4n+1}^{\text{HVSASM}}(1, z) + z^{4n-3} Z_{4n+1}^{\text{HVSASM}}(1, \bar{z}) \\ = (1+z) \tilde{Z}_{2n}^{\text{CSTCPP}}(1, z) Z_{2n+1}^{\text{VSASM}}(1, z, 1), \end{aligned}$$

and

$$\begin{aligned} Z_{4n+3}^{\text{HVSASM}}(1, z) + z^{4n-1} Z_{4n+3}^{\text{HVSASM}}(1, \bar{z}) \\ = (1+z) \tilde{Z}_{2n+2}^{\text{CSTCPP}}(1, z) Z_{2n+1}^{\text{VSASM}}(1, z, 1). \end{aligned}$$

VHSASMs

Theorem (Fischer-S., 2020)

We have for all $n \geq 1$,

$$\begin{aligned} Z_{4n+1}^{\text{HVSASM}}(1, z) + z^{4n-3} Z_{4n+1}^{\text{HVSASM}}(1, \bar{z}) \\ = (1+z) \tilde{Z}_{2n}^{\text{CSTCPP}}(1, z) Z_{2n+1}^{\text{VSASM}}(1, z, 1), \end{aligned}$$

and

$$\begin{aligned} Z_{4n+3}^{\text{HVSASM}}(1, z) + z^{4n-1} Z_{4n+3}^{\text{HVSASM}}(1, \bar{z}) \\ = (1+z) \tilde{Z}_{2n+2}^{\text{CSTCPP}}(1, z) Z_{2n+1}^{\text{VSASM}}(1, z, 1). \end{aligned}$$

Again, we have not yet defined one generating function in the above. We have generalized the above results, but we need some more background on plane partitions.

Vertically and Horizontally Perverse ASM

Vertically and Horizontally Perverse ASM

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & -1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & 1 & 0 & -1 & 0 & 1 & -1 & 1 & 0 \\ 1 & -1 & 1 & -1 & 1 & \star & 1 & -1 & 1 & -1 & 1 \\ 0 & 1 & -1 & 1 & 0 & -1 & 0 & 1 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & -1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

VHPASMs

Theorem (Fischer-S., 2020)

We have for all $n \geq 1$,

$$\begin{aligned} Z_{2n-1}^{\text{HVPASM}}(1, 1, z) + z^{4n-1} Z_{2n-1}^{\text{HVPASM}}(1, 1, \bar{z}) \\ = (1+z)(1-z+z^2) \left(Z_{2n+1}^{\text{VSASM}}(1, z, 1) \right)^2, \end{aligned}$$

and

$$\begin{aligned} Z_{2n-1}^{\text{HVPASM}}(1, z, 1) + z^{4n-3} Z_{2n-1}^{\text{HVPASM}}(1, \bar{z}, 1) \\ = (1+z) \left(Z_{2n+1}^{\text{VSASM}}(1, z, 1) \right)^2. \end{aligned}$$

VHPASMs

Theorem (Fischer-S., 2020)

We have for all $n \geq 1$,

$$\begin{aligned} Z_{2n-1}^{\text{HVPASM}}(1, 1, z) + z^{4n-1} Z_{2n-1}^{\text{HVPASM}}(1, 1, \bar{z}) \\ = (1+z)(1-z+z^2) \left(Z_{2n+1}^{\text{VSASM}}(1, z, 1) \right)^2, \end{aligned}$$

and

$$\begin{aligned} Z_{2n-1}^{\text{HVPASM}}(1, z, 1) + z^{4n-3} Z_{2n-1}^{\text{HVPASM}}(1, \bar{z}, 1) \\ = (1+z) \left(Z_{2n+1}^{\text{VSASM}}(1, z, 1) \right)^2. \end{aligned}$$

We have generalized the above results.


Plane Partitions

Plane Partitions

A *plane partition* in an $a \times b \times c$ box is a subset

$$PP \subset \{1, 2, \dots, a\} \times \{1, 2, \dots, b\} \times \{1, 2, \dots, c\}$$

with $(i', j', k') \in PP$ if $(i, j, k) \in PP$ and $(i', j', k') \leq (i, j, k)$.



Cyclically Symmetric Plane Partitions

Cyclically Symmetric Plane Partitions

A plane partition in an $n \times n \times n$ box is called *cyclically symmetric* if $(j, k, i) \in PP$ when $(i, j, k) \in PP$.

Cyclically Symmetric Plane Partitions

A plane partition in an $n \times n \times n$ box is called *cyclically symmetric* if $(j, k, i) \in PP$ when $(i, j, k) \in PP$.

George Andrews in 1979 proved that the number of cyclically symmetric plane partitions in an $n \times n \times n$ box is


$$\prod_{j=0}^{n-1} \frac{(3j+2)(3j)!}{(n+j)!}.$$

Cyclically Symmetric Plane Partitions

A plane partition in an $n \times n \times n$ box is called *cyclically symmetric* if $(j, k, i) \in PP$ when $(i, j, k) \in PP$.

George Andrews in 1979 proved that the number of cyclically symmetric plane partitions in an $n \times n \times n$ box is

$$\prod_{j=0}^{n-1} \frac{(3j+2)(3j)!}{(n+j)!}.$$



Cyclically Symmetric Transpose Complementary Plane Partitions

Cyclically Symmetric Transpose Complementary Plane Partitions

The complement of a plane partition π in a $r \times s \times t$ box is defined by the plane partition

$$\pi^c := \{(r+1-i, s+1-j, t+1-k) : (i, j, k) \notin \pi\}.$$

Cyclically Symmetric Transpose Complementary Plane Partitions

The complement of a plane partition π in a $r \times s \times t$ box is defined by the plane partition

$$\pi^c := \{(r+1-i, s+1-j, t+1-k) : (i, j, k) \notin \pi\}.$$

A plane partition π in an $2n \times 2n \times 2n$ box is called *cyclically symmetric transpose-complementary* if $(j, k, i) \in \pi$ when $(i, j, k) \in \pi$

Cyclically Symmetric Transpose Complementary Plane Partitions

The complement of a plane partition π in a $r \times s \times t$ box is defined by the plane partition

$$\pi^c := \{(r+1-i, s+1-j, t+1-k) : (i, j, k) \notin \pi\}.$$

A plane partition π in an $2n \times 2n \times 2n$ box is called *cyclically symmetric transpose-complementary* if $(j, k, i) \in \pi$ when $(i, j, k) \in \pi$ & $\pi^c = \pi$.

Cyclically Symmetric Transpose Complementary Plane Partitions

The complement of a plane partition π in a $r \times s \times t$ box is defined by the plane partition

$$\pi^c := \{(r+1-i, s+1-j, t+1-k) : (i, j, k) \notin \pi\}.$$


A plane partition π in an $2n \times 2n \times 2n$ box is called *cyclically symmetric transpose-complementary* if $(j, k, i) \in \pi$ when $(i, j, k) \in \pi$ & $\pi^c = \pi$.

Mills, Robbins and Rumsey in 1987 proved that the number of cyclically symmetric transpose-complementary plane partitions in an $2n \times 2n \times 2n$ box is

$$\prod_{i=0}^{n-1} \frac{(2i+1)(6i+2)!}{(6i+1)(2n+2i)!}.$$


Cyclically Symmetric Transpose Complementary Plane Partitions

Cyclically Symmetric Transpose Complementary Plane Partitions





Cyclically Symmetric Transpose Complementary Rhombus Tilings of a Hexagon

Cyclically Symmetric Transpose Complementary Rhombus Tilings of a Hexagon





Plane Partitions to Non Intersecting Lattice Paths

Plane Partitions to Non Intersecting Lattice Paths



Plane Partitions to Non Intersecting Lattice Paths



The starting points are $A_i(i, 2i)$ and the ending points are $E_j(2j, j)$.

Generating Functions

Generating Functions

We define the following generating functions

Generating Functions

We define the following generating functions

$$Z_{2n}^{\text{CSTCPP}}(x, z) = \sum_P x^{\text{special parts in } P} z^{\#(n)}$$

$$\tilde{Z}_{2n}^{\text{CSTCPP}}(x, z) = \sum_P x^{\text{quasi special parts in } P} z^{n+1+\#(n)_1-\#(n-1)_1},$$

Generating Functions

We define the following generating functions

$$Z_{2n}^{\text{CSTCPP}}(x, z) = \sum_P x^{\text{special parts in } P} z^{\#(n)}$$

$$\tilde{Z}_{2n}^{\text{CSTCPP}}(x, z) = \sum_P x^{\text{quasi special parts in } P} z^{n+1+\#(n)_1-\#(n-1)_1},$$

where

- ▶ special parts = number of horizontal steps below the diagonal
 $x = y$,

Generating Functions

We define the following generating functions

$$Z_{2n}^{\text{CSTCPP}}(x, z) = \sum_P x^{\text{special parts in } P} z^{\#(n)}$$

$$\tilde{Z}_{2n}^{\text{CSTCPP}}(x, z) = \sum_P x^{\text{quasi special parts in } P} z^{n+1+\#(n)_1-\#(n-1)_1},$$

where

- ▶ special parts = number of horizontal steps below the diagonal $x = y$,
- ▶ quasi special parts = number of horizontal steps below the diagonal $x = y + 1$,

Generating Functions

We define the following generating functions

$$Z_{2n}^{\text{CSTCPP}}(x, z) = \sum_P x^{\text{special parts in } P} z^{\#(n)}$$

$$\tilde{Z}_{2n}^{\text{CSTCPP}}(x, z) = \sum_P x^{\text{quasi special parts in } P} z^{n+1+\#(n)_1-\#(n-1)_1},$$

where

- ▶ special parts = number of horizontal steps below the diagonal $x = y$,
- ▶ quasi special parts = number of horizontal steps below the diagonal $x = y + 1$, and
- ▶ $\#(i)_j$ = number of i 's in row j .

Related Results

Related Results

Theorem (Behrend-S., 2022)

We have, for all $n \geq 1$

$$\begin{aligned} & \tilde{Z}_{2n}^{\text{CSTCPP}}(x, z) \\ = & \det_{1 \leq i, j \leq n-1} \left(\begin{array}{l} \sum_{k=0}^{\min(2i, 2j)} \binom{i}{k-i} \binom{j}{k-j} x^{2i-k}, \\ j \leq n-2 \\ \sum_{k=0}^{2i} \sum_{m=n-1}^k \sum_{l=n-1}^m \binom{2n-4-m}{k-m} \binom{i}{k-i} x^{2i-k} z^{2l-m}, \\ j = n-1 \end{array} \right). \end{aligned}$$

Related Results contd.

Related Results contd.

Theorem (Behrend-S., 2022)

We have, for all $n \geq 1$

$$(1+z)\tilde{Z}_{2n}^{\text{CSTCPP}}(1,z) = Z_{2n}^{\text{CSTCPP}}(1,z) + z^{2n-1}Z_{2n}^{\text{CSTCPP}}(1,\bar{z}).$$

Related Results contd.

Theorem (Behrend-S., 2022)

We have, for all $n \geq 1$

$$(1+z) \tilde{Z}_{2n}^{\text{CSTCPP}}(1, z) = Z_{2n}^{\text{CSTCPP}}(1, z) + z^{2n-1} Z_{2n}^{\text{CSTCPP}}(1, \bar{z}).$$

- There is a subset of VSASMs which are equinumerous with CSTCPPs (Behrend-S., 2022).

Related Results contd.

Theorem (Behrend-S., 2022)

We have, for all $n \geq 1$

$$(1+z) \tilde{Z}_{2n}^{\text{CSTCPP}}(1, z) = Z_{2n}^{\text{CSTCPP}}(1, z) + z^{2n-1} Z_{2n}^{\text{CSTCPP}}(1, \bar{z}).$$

- ▶ There is a subset of VSASMs which are equinumerous with CSTCPPs (Behrend-S., 2022).
- ▶ We call them quasi-VSASMs. The above result can also be written in terms of quasi-VSASMs.

Related Results contd.

Theorem (Behrend-S., 2022)

We have, for all $n \geq 1$

$$(1+z) \tilde{Z}_{2n}^{\text{CSTCPP}}(1, z) = Z_{2n}^{\text{CSTCPP}}(1, z) + z^{2n-1} Z_{2n}^{\text{CSTCPP}}(1, \bar{z}).$$

- ▶ There is a subset of VSASMs which are equinumerous with CSTCPPs (Behrend-S., 2022).
- ▶ We call them quasi-VSASMs. The above result can also be written in terms of quasi-VSASMs.

Theorem (Behrend-S., 2022)

The number of quasi-VSASMs of order $2n$ is equal to

$$\prod_{i=0}^{n-1} \frac{(2i+1)(6i+2)!}{(6i+1)(2n+2i)!}.$$

Generalized Results

Generalized Results

Theorem (Behrend-S., 2022)

We have, for all $n \geq 1$

$$\begin{aligned} Z_{4n+1}^{\text{HVSASM}}(x, z) + z^{4n-3} Z_{4n+1}^{\text{HVSASM}}(x, \bar{z}) \\ = (1 + z) \tilde{Z}_{2n}^{\text{CSTCPP}}(x, z) Z_{2n+1}^{\text{VSASM}}(x, z, 1), \end{aligned}$$

Generalized Results

Theorem (Behrend-S., 2022)

We have, for all $n \geq 1$

$$\begin{aligned} Z_{4n+1}^{\text{HVSASM}}(x, z) + z^{4n-3} Z_{4n+1}^{\text{HVSASM}}(x, \bar{z}) \\ = (1+z) \tilde{Z}_{2n}^{\text{CSTCPP}}(x, z) Z_{2n+1}^{\text{VSASM}}(x, z, 1), \end{aligned}$$

$$\begin{aligned} Z_{4n+3}^{\text{HVSASM}}(x, z) + z^{4n-1} Z_{4n+3}^{\text{HVSASM}}(x, \bar{z}) \\ = (1+z) \tilde{Z}_{2n+2}^{\text{CSTCPP}}(x, z) Z_{2n+1}^{\text{VSASM}}(x, z, 1), \end{aligned}$$

Generalized Results

Theorem (Behrend-S., 2022)

We have, for all $n \geq 1$

$$\begin{aligned} Z_{4n+1}^{\text{HVSASM}}(x, z) + z^{4n-3} Z_{4n+1}^{\text{HVSASM}}(x, \bar{z}) \\ = (1+z) \tilde{Z}_{2n}^{\text{CSTCPP}}(x, z) Z_{2n+1}^{\text{VSASM}}(x, z, 1), \end{aligned}$$

$$\begin{aligned} Z_{4n+3}^{\text{HVSASM}}(x, z) + z^{4n-1} Z_{4n+3}^{\text{HVSASM}}(x, \bar{z}) \\ = (1+z) \tilde{Z}_{2n+2}^{\text{CSTCPP}}(x, z) Z_{2n+1}^{\text{VSASM}}(x, z, 1), \end{aligned}$$

$$\begin{aligned} Z_{2n-1}^{\text{HVPASM}}(x, 1, z) + z^{4n-1} Z_{2n-1}^{\text{HVPASM}}(x, 1, \bar{z}) \\ = (1+z) \tilde{Z}_{2n+1}^{\text{VSASM}}(x, z) Z_{2n+1}^{\text{VSASM}}(x, z, 1), \end{aligned}$$

Generalized Results

Theorem (Behrend-S., 2022)

We have, for all $n \geq 1$

$$\begin{aligned} Z_{4n+1}^{\text{HVSASM}}(x, z) + z^{4n-3} Z_{4n+1}^{\text{HVSASM}}(x, \bar{z}) \\ = (1+z) \tilde{Z}_{2n}^{\text{CSTCPP}}(x, z) Z_{2n+1}^{\text{VSASM}}(x, z, 1), \end{aligned}$$

$$\begin{aligned} Z_{4n+3}^{\text{HVSASM}}(x, z) + z^{4n-1} Z_{4n+3}^{\text{HVSASM}}(x, \bar{z}) \\ = (1+z) \tilde{Z}_{2n+2}^{\text{CSTCPP}}(x, z) Z_{2n+1}^{\text{VSASM}}(x, z, 1), \end{aligned}$$

$$\begin{aligned} Z_{2n-1}^{\text{HVPASM}}(x, 1, z) + z^{4n-1} Z_{2n-1}^{\text{HVPASM}}(x, 1, \bar{z}) \\ = (1+z) \tilde{Z}_{2n+1}^{\text{VSASM}}(x, z) Z_{2n+1}^{\text{VSASM}}(x, z, 1), \end{aligned}$$

and

$$\begin{aligned} Z_{2n-1}^{\text{HVPASM}}(x, z, 1) + z^{4n-3} Z_{2n-1}^{\text{HVPASM}}(x, \bar{z}, 1) \\ = (1+z) \hat{Z}_{2n+1}^{\text{VSASM}}(x, z) Z_{2n+1}^{\text{VSASM}}(x, z, 1). \end{aligned}$$

Thank you for your attention!

Bijection between ASMs and Six Vertex Model

Bijection between ASMs and Six Vertex Model

Kuperberg's proof of the ASM conjecture was by exploiting a bijection between the ASMs and a model in statistical physics, called the six-vertex model.

Bijection between ASMs and Six Vertex Model

Kuperberg's proof of the ASM conjecture was by exploiting a bijection between the ASMs and a model in statistical physics, called the six-vertex model.




Figure: Six Vertex Model with Domain Wall Boundary Condition.

Bijection between ASMs and Six Vertex Model

A state of a corresponding six-vertex model is an orientation on the edges of this graph, such that both the in-degree and the out-degree of each vertex with degree 4 is 2.




Figure: Six Vertex Model with Domain Wall Boundary Condition.

The Bijection

The Bijection

If we associate to each of the degree 4 vertex in a six-vertex state with a number, as given in the figure below

The Bijection

If we associate to each of the degree 4 vertex in a six-vertex state with a number, as given in the figure below




Figure: The corresponding states of the six-vertex model and the entries of an ASM.

The Bijection

If we associate to each of the degree 4 vertex in a six-vertex state with a number, as given in the figure below




Figure: The corresponding states of the six-vertex model and the entries of an ASM.

then we obtain a matrix with entries in the set $\{0, 1, -1\}$.

The Bijection

If we associate to each of the degree 4 vertex in a six-vertex state with a number, as given in the figure below




Figure: The corresponding states of the six-vertex model and the entries of an ASM.

then we obtain a matrix with entries in the set $\{0, 1, -1\}$.

Such a matrix will be an ASM, and we get a bijection between ASMs and states of the six-vertex model.

Example

Example





Figure: Six Vertex Model with Domain Wall Boundary Condition.

Example



$$\begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & -1 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

Figure: Six Vertex Model with Domain Wall Boundary Condition.

Weighted Enumeration

Weighted Enumeration

- ▶ We assign to each vertex v , a weight $w(v)$.

Weighted Enumeration

- ▶ We assign to each vertex v , a weight $w(v)$.
- ▶ Weight of a state C is $W(C) = \prod_{v \in C} w(v)$.

Weighted Enumeration

- ▶ We assign to each vertex v , a weight $w(v)$.
- ▶ Weight of a state C is $W(C) = \prod_{v \in C} w(v)$.
- ▶ Generating function or the partition function $Z_n = \sum_C W(C)$.

Weighted Enumeration

- ▶ We assign to each vertex v , a weight $w(v)$.
- ▶ Weight of a state C is $W(C) = \prod_{v \in C} w(v)$.
- ▶ Generating function or the partition function $Z_n = \sum_C W(C)$.
- ▶ Specializing the parameters in Z_n , we get enumeration results.

Our Weights

Our Weights

q is a parameter, which we will specialize later, $\bar{x} = \frac{1}{x} = x^{-1}$ and $\sigma(x) = x - \bar{x}$.

Our Weights

q is a parameter, which we will specialize later, $\bar{x} = \frac{1}{x} = x^{-1}$ and $\sigma(x) = x - \bar{x}$.




Figure: The weights of the vertices of an ASM with spectral parameter u .

Our Weights

q is a parameter, which we will specialize later, $\bar{x} = \frac{1}{x} = x^{-1}$ and $\sigma(x) = x - \bar{x}$.





Figure: The weights of the vertices of an ASM with spectral parameter u .


We normalize further by $\sigma(q^2)$ so that we have all entries 1 and -1 to have weight 1 .

How are weights assigned?

How are weights assigned?



How are weights assigned?



A vertex lying at the intersection of a vertical line with parameter y_j and a horizontal line with parameter x_i is assigned the label $\frac{x_i}{y_j}$.

Another type of ASM

Another type of ASM

In order to study VSASMs we need what are called U-turn domain boundary wall conditions.

Another type of ASM

In order to study VSASMs we need what are called U-turn domain boundary wall conditions.

The set of VASAMs is a subset of what are called U-turn ASMs or ASMs with U-turn boundary (UASMs).

Another type of ASM

In order to study VSASMs we need what are called U-turn domain boundary wall conditions.

The set of VASAMs is a subset of what are called U-turn ASMs or ASMs with U-turn boundary (UASMs).

We will explain this connection shortly.

UASMs

UASMs

An U-turn ASM is an $2n \times n$ array which satisfies the usual properties of ASMs if one looks at it vertically.

UASMs

An U-turn ASM is an $2n \times n$ array which satisfies the usual properties of ASMs if one looks at it vertically.

However, if one looks at it horizontally then the 1's and -1 's alternate if we start along an odd numbered row from left to right and then continue along the next even numbered row from right to left.

UASMs

An U-turn ASM is an $2n \times n$ array which satisfies the usual properties of ASMs if one looks at it vertically.

However, if one looks at it horizontally then the 1's and -1 's alternate if we start along an odd numbered row from left to right and then continue along the next even numbered row from right to left.




Figure: An U-turn ASM with the corresponding six-vertex state.

New weights


New weights

As can be seen from the figure, we add an additional parameter on the U-turns.


New weights

As can be seen from the figure, we add an additional parameter on the U-turns.

This gives rise to two new type of vertices whose corresponding weights are given below.



$$\sigma(bu)$$



$$\sigma(b\bar{u})$$

Figure: Weights of the new vertices.

Partition Function

Partition Function

Tsuchiya was the first to consider a U-turn domain wall boundary condition, and gave a partition function for them.

Partition Function

Tsuchiya was the first to consider a U-turn domain wall boundary condition, and gave a partition function for them.

$$Z_U(n; x, y) = \frac{\sigma(q^2)^n \prod_i (\sigma(b\bar{y}_i) \sigma(q^2 x_i^2)) \prod_{i,j} (\sigma'(x_i \bar{y}_j) \sigma'(x_i y_j))}{\prod_{i < j} (\sigma(\bar{x}_i x_j) \sigma(y_i \bar{y}_j)) \prod_{i \leq j} (\sigma(\bar{x}_i \bar{x}_j) \sigma(y_i y_j))} \times \det M_U(n; x, y), \quad (1)$$

where $\sigma'(x) = \sigma(qx)\sigma(q\bar{x})$ and M_U is an $n \times n$ matrix defined as

$$M_U(n; x, y)_{i,j} = \frac{1}{\sigma'(x_i \bar{y}_j)} - \frac{1}{\sigma'(x_i y_j)}.$$

Some observations

Some observations

- ▶ VSASMs occur only for odd order.

Some observations

- ▶ VSASMs occur only for odd order.

$$\begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 & 0 & 1 & 0 \\ 1 & -1 & 0 & 1 & 0 & -1 & 1 \\ 0 & 0 & 1 & -1 & 1 & 0 & 0 \\ 0 & 1 & -1 & 1 & -1 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}.$$

Some observations

- ▶ VSASMs occur only for odd order.
- ▶ We need only the first $n + 1$ columns of the VSASM to know the full matrix.

Some observations

- ▶ VSASMs occur only for odd order.
- ▶ We need only the first $n + 1$ columns of the VSASM to know the full matrix.
- ▶ The middle column is an alternating row with 1 and -1.

Some observations

- ▶ VSASMs occur only for odd order.
- ▶ We need only the first $n + 1$ columns of the VSASM to know the full matrix.
- ▶ The middle column is an alternating row with 1 and -1.

$$\begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 & 0 & 1 & 0 \\ 1 & -1 & 0 & 1 & 0 & -1 & 1 \\ 0 & 0 & 1 & -1 & 1 & 0 & 0 \\ 0 & 1 & -1 & 1 & -1 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}.$$

Some observations

- ▶ VSASMs occur only for odd order.
- ▶ We need only the first $n + 1$ columns of the VSASM to know the full matrix.
- ▶ The middle column is an alternating row with 1 and -1.

Some observations

- ▶ VSASMs occur only for odd order.
- ▶ We need only the first $n + 1$ columns of the VSASM to know the full matrix.
- ▶ The middle column is an alternating row with 1 and -1.
- ▶ So, n columns are sufficient to know the whole matrix.

Some observations

- ▶ VSASMs occur only for odd order.
- ▶ We need only the first $n + 1$ columns of the VSASM to know the full matrix.
- ▶ The middle column is an alternating row with 1 and -1.
- ▶ So, n columns are sufficient to know the whole matrix.
- ▶ Moreover, the first and last rows are always the same.

Transformation

Transformation

We can transform a VSASM into an USASM in two steps:

Transformation

We can transform a VSASM into an USASM in two steps:

- ▶ Delete the last row.

Transformation

We can transform a VSASM into an USASM in two steps:

- ▶ Delete the last row.
- ▶ Connect pairwise the alternating edges on the right most column of the $2n \times n$ matrix.

Transformation

We can transform a VSASM into an USASM in two steps:

- ▶ Delete the last row.
- ▶ Connect pairwise the alternating edges on the right most column of the $2n \times n$ matrix.




Figure: Transformation of a VSASM into a USASM.

Transformation

We can transform a VSASM into an USASM in two steps:

- ▶ Delete the last row.
- ▶ Connect pairwise the alternating edges on the right most column of the $2n \times n$ matrix.




Figure: Transformation of a VSASM into a USASM.

Notice that all U-turns are downward pointing.

Thank you for your attention!