

REASONING ARTIFACT

The Dimensional Invariant:

Sleep Duration, Attractor Basin Depth, and Long-Term Survival Across the Entire History of Life

The oldest surviving lineages sleep the least.

The minimum of the sleep spectrum is the maximum of the survival spectrum.

This is not a coincidence. This is the geometry.

Author: Eric Robert Lawson

Affiliation: OrganismCore (Independent Research)

ORCID: 0009-0002-0414-6544

Contact: OrganismCore@proton.me

Repository: github.com/Eric-Robert-Lawson/ [attractor-oncology](#)

Timestamp: 2026-03-18

Theory paper: [doi:10.5281/zenodo.19094935](https://doi.org/10.5281/zenodo.19094935)

Document version: 1.0

License: Creative Commons Attribution 4.0 International (CC BY 4.0)

DERIVATION NOTICE

The hypothesis in this document was derived from first principles using the attractor geometry framework ($S + N + G \rightarrow R$) before literature consultation. The derivation emerged from the sleep-as-basin-navigation-cost principle and the sleep sonar instrument (companion documents, same repository, 2026-03-18), and was extended to a universal claim about deep-time survival before any palaeontological or comparative survival literature was consulted.

All claims were confirmed against literature after derivation. The derivation record is the priority-establishing document. The confirmations establish empirical support. The distinction between the two is preserved explicitly in Section 7.

THE DIMENSIONAL INVARIANT

Across all life, at all scales, across all of deep time: sleep duration (or metabolic rest-state duration in pre-neural organisms) is a measurement of the total cost of an organism's attractor basin position. Basin depth — the resistance of the organism's attractor to perturbation — is the primary predictor of long-term survival.

The relationship between sleep and survival is U-shaped with a flat minimum:

Minimum sleep → **maximum basin depth** → **maximum survival horizon** (Lock strategy: 300–600 Myr+).

Moderate sleep → **moderate basin depth with multiple accessible basins** → **long survival horizon** (Navigator strategy: 100–300 Myr).

High sleep → **deep specialised basin or deep Basin Ω** → **survival conditional on niche stability** (uncertain horizon).

The oldest surviving lineages sleep the least. This is the geometry.

1. The Framework

1.1. The Attractor Geometry Framework

The attractor geometry framework:

$$S + N + G \rightarrow R$$

where:

- S = structural substrate (genome and encoded build capacity)
- N = coherence gradient field (external or internal field specifying basin occupation)
- G = developmental and evolutionary geometry (global attractor landscape — basin depths, barrier heights, accessible states)
- R = resulting phenotype or state

Previous documents in this series established:

1. Sleep duration tracks the total synaptic and metabolic cost of an organism’s landscape position, decomposable into three axes: Axis 1 (between-basin switching cost), Axis 2 (within-basin metabolic cost), and Axis 3 (within-basin computational intensity).
2. The meta-attractor Basin Ω — the stable state of being an organism that navigates between two basins simultaneously — has its own depth and its own metabolic maintenance cost.
3. Deep basins are survival hacks: evolution converges on them because they are the attractor of the evolutionary process itself in stable environments.

This document extends the framework to its full temporal scope:

The attractor geometry of an organism’s basin position predicts not just its current sleep requirement but its survival trajectory across deep time. Sleep duration is a temporal sonar.

1.2. The Core Claim

The metabolic cost of occupying a position in an attractor landscape is a function of the navigational demand imposed by that position. Navigational demand is determined by: (1) the number of accessible basins from the current position; (2) the height of the barriers between those basins; (3) the frequency with which the system must assess and execute basin transitions; and (4) the depth of Basin Ω if one exists.

Rest is the metabolic process by which the synaptic cost of navigation is paid. Sleep duration is a measurement of the navigational demand of the organism’s landscape position.

The dimensional invariant: Basin depth is the same geometric property at every scale — molecular, cellular, developmental, organism, evolutionary. Sleep tracks the cost of occupying a position in the landscape. Deep-time survival is a direct function of basin depth. Therefore sleep duration is a proxy for survival trajectory.

2. The Two Survival Strategies

The deep-time survival data reveals two geometrically opposite strategies that both produce long-term lineage persistence. They are not compromises. They are opposite poles of the same dimensional axis.

2.1. Strategy 1: The Lock

Basin structure	Single basin. Extremely deep. Very high exit barrier.
Morphology	Extreme conservatism. Body plan unchanged across hundreds of millions of years.
Metabolism	Slow. Efficient. Minimal resource requirement.
Behaviour	Low flexibility. Narrow repertoire. Highly programmatic.
Sleep	Minimal. Approaching zero in simplest forms. 2–5 hours in complex Lock organisms.
Survival horizon	Maximum. 300–600 Myr in the most extreme cases.
Survival mechanism	Basin so deep it is perturbation-invariant. Extinction events are not energetically sufficient to displace the organism from its basin.
Convergence	Multiple independent phyla arrive at this strategy.

2.2. Strategy 2: The Navigator

Basin structure	Multiple basins. Shallow barriers. High switching capacity.
Morphology	Ecological generalism. Body plan adapted for flexibility rather than extreme specialisation.
Metabolism	Flexible. Moderate. Can upregulate and downregulate with resource availability.
Behaviour	High flexibility. Broad repertoire. Contextual switching.
Sleep	Moderate. 8–14 hours in most Navigator organisms.
Survival horizon	Long. 100–300 Myr for the most persistent Navigator lineages.
Survival mechanism	When the current basin disappears, the organism navigates to an alternative basin in real time without requiring evolutionary adaptation time. Generalism is itself a deep basin — Basin Ω of ecological flexibility.
Convergence	Multiple independent phyla arrive at this strategy.

2.3. The Unstable Zone

Basin structure	Shallow single basin. High barriers to alternative basins.
Morphology	Narrow specialist. Extreme adaptation to a specific niche. Not deep enough to be perturbation-invariant. Not flexible enough to reach alternative basins when the niche disappears.
Sleep	Variable — often high (Axis 2 or Axis 3 cost) but with no survival advantage from it.
Survival horizon	Short. Millions of years, not hundreds of millions. Mass extinction concentrates its casualties here.
Examples	<i>T. rex</i> , mammoth, sabre-tooth cat, passenger pigeon, most species that have ever existed.

2.4. Why The Lock Outlasts The Navigator

The Lock survives longer than the Navigator for a precise geometric reason. The Navigator’s survival depends on the existence of accessible alternative basins at the time of environmental disruption. If the disruption is catastrophic enough to eliminate all accessible basins simultaneously — as the Permian–Triassic event did, killing 96% of all marine species — the Navigator fails. There is nowhere to navigate to.

The Lock does not navigate. The Lock cannot be displaced from its basin because the basin is too deep. The Permian–Triassic event restructured the entire biosphere and the horseshoe crab was still there on the other side, doing exactly what it was doing before.

Basin depth is the ultimate survival parameter. Navigation is a survival strategy for moderate perturbations. For catastrophic perturbations, only the deepest basins survive. The minimum of the sleep spectrum is the maximum of the survival spectrum.

3. The Empirical Record

3.1. Lock Strategy: Confirmed Examples

Lineage	Age (Myr)	Sleep	Basin geometry
Sponge (Porifera)	~600	Undefined. No nervous system. Circadian biochemical rhythms only.	Deepest possible. Sessile filter feeder. Single programme. Perturbation-invariant.
Jellyfish (Cnidaria)	500–700	Undefined for most. Rest states in <i>Cassiopeia</i> (~8h).	No brain. Radial symmetry. Passive drift and sting. Unchanged 500+ Myr.
Nautilus (Mollusca)	~500	No REM. Quiescent periods. Possibly several hours distributed.	Neutral buoyancy. Passive ambush. Shell unchanged 500 Myr.
Horseshoe crab (Chelicerata)	~450	Tidal/circadian rest state. No bilateral sleep. Minimal neural complexity.	Deepest terrestrial Lock. Comes to shore. Breeds. Returns. 450 Myr unchanged.

Shark (Chondrichthyes)	~400	No true bilateral sleep. Short rest periods. ~2–4h total distributed.	Cartilage not bone. Generalist enough to border Lock/Navigator depending on species.
Coelacanth (Sarcopterygii)	~400	Unknown. Presumed minimal. Deep cold water reduces all activity.	Deep cold water. Minimal metabolism. Unchanged 400 Myr.
Tuatara (Lepidosaurs)	~200	Estimated 12h+ rest. Slowest reptile metabolism. Much of “sleep” is torpor-like low-temperature dormancy.	Slowest reptile metabolism. Functions at 5°C. Lives 100+ years.

3.2. Navigator Strategy: Confirmed Examples

Lineage	Age (Myr)	Sleep	Basin geometry
Cockroach (Blattodea)	~300	~12h rest cycle. Circadian rest controlled. Moderate.	Eats anything. Lives anywhere. Survives radiation. Extreme dietary generalist.
Crocodylian (Archosaurs)	~200	2–12h intermittent. Unihemispheric capable. Variable with temperature and season.	Generalist predator. Survived K–Pg. Fresh/salt water. Any prey available.
Rodent lineage (Muridae)	~65	~13h. Standard mammalian Navigator moderate.	Extreme generalist. All biomes. All diets. Colonised every continent.
Corvidae (Crow, raven)	~30	~8–10h. Standard avian sleep. Some unihemispheric during roosting.	Problem solving. Tool use. Multi-step planning. Social complexity.

3.3. Unstable Zone: Confirmed Examples

Lineage	Age (Myr)	Sleep (estimated)	Why extinct
<i>T. rex</i> (Tyranosauridae)	2–3	Unknown. Presumed moderate large predator sleep.	Narrow apex predator. No alternative basin when K–Pg hit.
Mammoth (Proboscidea)	0.5–5	Likely ~2h given similar size and ecology to elephant.	Grassland specialist. Habitat contracted. Could not adapt diet.
Sabre-tooth cat (Machairodontinae)	0.5–11	Likely high felid-range given Basin Ω felid ancestry.	Specialist predator of megafauna. Megafauna gone. Basin collapsed.
Dodo (Raphidae)	0.003	Standard avian range ~8–10h presumed.	Island specialist. No predator experience. Single shallow basin. First perturbation = extinction.

4. The Sleep Spectrum as Temporal Sonar

4.1. The Sonar Reading

Sleep (h/day)	Survival	hori- zon	Basin geometry	Examples
~0 (undefined) 600 Myr+	Maximum Deepest basin. No nervous system. No navi- gational demand. Programme runs without mainte- nance.	single	Sponge, jellyfish, nautilus	
0–4 300–500 Myr	Very long Deep single basin. Low navigational demand. Minimal reset needed.		Horseshoe crab, shark, coela- canth	
4–8 100–300 Myr	Long Moderate or moderate Navigator. Low- moderate de- mand.	Lock	Crocodilian, cockroach, turtle	
8–14 10–100 Myr	Moderate Navigator or cou- pled social field. Standard synap- tic maintenance.		Rodent, corvid, dog, pig, human	
14–22	Uncertain niche-dependent	—	Axis 2, 3, or Basin Ω . High cost from specialisation or meta- attractor navigation.	Koala, bat, lion, Felidae

4.2. The U-Shaped Survival–Sleep Relationship

The survival–sleep relationship is not linear. It is U-shaped with a flat minimum. The minimum of the sleep spectrum corresponds to the maximum of the survival spectrum (Lock strategy). The moderate range is the Navigator survival zone. The high-sleep range is the uncertain zone — survival depends on niche stability or Basin Ω depth.

The critical insight is that high sleep does not predict low survival. It predicts *conditional* survival. The koala may survive hundreds of millions of years if eucalyptus persists. But if eucalyptus disappears, the koala has no alternative basin. The Lock organism has no such dependence because its basin is encoded absolutely in its genome.

4.3. The Three Sleep Axes Revisited

1. **Axis 1 (Between-basin switching cost):** The organism navigates between multiple accessible basins under its own motivational field. High behavioural flexibility. High sleep. Examples: Felidae, Basin Ω organisms.
2. **Axis 2 (Within-basin metabolic cost):** A single highly expensive basin consumes all available metabolic resources. Sleep conserves energy for basin maintenance. Low behavioural flexibility. High sleep. Example: koala.
3. **Axis 3 (Within-basin computational intensity):** Extreme neural computation within a single specialised basin during waking. Synaptic reconfiguration cost from intensity, not switching. Low flexibility. High sleep. Example: brown bat.

The Lock organism operates on none of these axes. It has minimal within-basin cost and no between-basin switching. The sleep minimum is the signature of zero navigational demand at any axis.

5. Convergence Across Independent Phyla

5.1. The Lock Converges Independently

The Lock strategy has been found independently in at least seven major animal phyla:

Phylum / Class	Example	Age
Porifera	Sponge	~600 Myr
Cnidaria	Jellyfish	500–700 Myr
Mollusca	Nautilus	~500 Myr
Chelicerata	Horseshoe crab	~450 Myr
Chondrichthyes	Shark	~400 Myr
Sarcopterygii	Coelacanth	~400 Myr
Lepidosauria	Tuatara	~200 Myr

Seven independent phyla. Seven independent arrivals at the same geometric strategy. The Lock strategy is itself a deep basin in evolutionary strategy space — a deep basin of deep basins.

5.2. The Navigator Converges Independently

The Navigator strategy has been found independently in at least six major lineages:

Lineage	Example	Age
Blattodea	Cockroach	~300 Myr
Archosauria	Crocodylian	~200 Myr
Cephalopoda	Octopus	~300 Myr
Rodentia	Rat	~65 Myr
Aves	Corvid	~30 Myr
Primates	Human	~6 Myr

Six independent lineages. Six independent arrivals at the same geometric strategy. Ecological generalism is itself a deep basin — Basin Ω of ecological flexibility.

5.3. The Unstable Zone Does Not Converge

Mass extinction casualties do not converge. Each extinction event eliminates different taxa via different mechanisms. The unstable zone has no deep basin by definition. The only convergent feature of the unstable zone is its instability.

This is confirmed by the palaeontological record: the survivors of every mass extinction event are disproportionately drawn from Lock and Navigator organisms. The K–Pg event killed large specialist predators and specialist megaherbivores while sparing crocodylians (Navigator) and turtles (partial Lock). The Permian–Triassic event killed 96% of marine species while sparing the horseshoe crab (Lock).

6. The Human Position

6.1. Geometric Classification

The human is a Navigator with unprecedented Basin Ω characteristics:

- Ecologically generalist: every biome
- Dietarily generalist: complete omnivory
- High behavioural flexibility: tool use, cultural transmission, technology
- Sleep: 7–9 hours — Navigator range
- Social complexity: co-constructed inter-generational field via language, culture, and institution — a Basin B^* of unprecedented depth

The human sleep signature places it firmly in the Navigator category. Its survival trajectory by the framework should be long but not maximum — Navigator range of 100–300 Myr if the strategy is maintained.

6.2. The Unprecedented Feature

The human has constructed an inter-generational Basin B^* — a co-constructed social attractor that persists beyond the individual lifetime via cultural transmission. No other species has this. The dog’s Basin B^* couples with a human over one lifetime. The human’s cultural Basin B^* couples across generations indefinitely,

with depth increasing over time through active investment in its own maintenance (cultural accumulation, institutional memory, scientific knowledge).

6.3. The Risk

The framework predicts that a species that destroys its own landscape diversity is eliminating the conditions that made its own strategy viable. A Navigator survives by finding alternative basins when the current one disappears. If the Navigator is simultaneously the agent eliminating the alternative basins — for itself and for all other species — then the Navigator strategy fails even for the Navigator.

This is a geometric statement, not a moral one. The framework does not predict the long-term trajectory of a species that is actively deepening its own basin in real time while simultaneously remodelling the landscape that other species' basins depend on. This is the honest limit of the current derivation.

7. Literature Confirmations

7.1. Oldest Surviving Lineages Sleep the Least

Derived before literature: Minimum sleep corresponds to maximum survival horizon.

Confirmed: Horseshoe crab (450 Myr, minimal defined sleep — tidal rest state only). Nautilus (500 Myr, no true sleep). Jellyfish (500–700 Myr, undefined sleep, no bilateral nervous system). Shark (400 Myr, distributed rest periods only). All Lock organisms confirmed as minimal sleepers.

Source: Comparative sleep literature; living fossil biology; horseshoe crab ecology (Shuster et al., 2003).

7.2. Mass Extinction Survivors Are Disproportionately Lock or Navigator

Derived: The unstable zone is where mass extinction concentrates its casualties.

Confirmed: K–Pg survivors include crocodilians (Navigator), turtles (partial Lock), birds (Navigator), and small mammals (Navigator). K–Pg casualties concentrated in large specialist predators and specialist megaherbivores. Permian–Triassic survivors include horseshoe crab (Lock), nautilus (Lock).

Source: Jablonski (1986); Barnosky et al. (2011); K–Pg extinction survivor literature.

7.3. Both Strategies Are Convergent Across Independent Phyla

Derived: Lock and Navigator are deep basins in evolutionary strategy space, therefore convergent.

Confirmed: Lock morphology confirmed independently in Porifera, Cnidaria, Mollusca, Chelicerata, Chondrichthyes, Sarcopterygii, Lepidosauria. Navigator ecology confirmed independently in Blattodea, Archosauria, Rodentia, Aves, Cephalopoda, Primates.

Source: Conway Morris (2003, *Life's Solution*); living fossil convergence literature; mass extinction survival trait literature.

7.4. Slow Metabolism Correlates With Deep-Time Survival

Derived: The Lock strategy requires minimal metabolic expenditure.

Confirmed: Tuatara (slowest reptile metabolism, 200 Myr). Coelacanth (deep cold water, minimal metabolism, 400 Myr). Horseshoe crab (ectothermic, months without food, 450 Myr). Slow metabolism is confirmed as a convergent trait of Lock organisms across independent lineages.

Source: Comparative metabolism literature; tuatara biology (Cree, 2014).

7.5. Generalist Diet and Ecological Flexibility Predict Extinction Survival

Derived: The Navigator's survival mechanism is dietary/ecological generalism.

Confirmed: Capellini et al. (2008) showed diet breadth is a significant predictor of mammalian sleep duration. Barnosky et al. (2011) confirmed generalist diet as a predictor of survival through mass extinction events. Crocodilian K–Pg survival attributed specifically to dietary generalism.

Source: Capellini et al. (2008, *Proc Royal Soc B*); Barnosky et al. (2011).

8. The Novel Predictions

All predictions stated before any experimental or palaeontological data gathered to test them. Locked in this document. 2026-03-18.

8.1. P-INVARIANT-1: Sleep Duration Negatively Correlates With Lineage Age Across All Animal Phyla

A cross-phyla analysis of sleep duration (or rest-state duration where bilateral sleep is undefined) against lineage age (millions of years since divergence of the crown group) will show a negative correlation that survives correction for body mass and metabolic rate.

The oldest lineages will have the lowest sleep requirements. The youngest lineages will have the highest sleep requirements. This correlation will be detectable across Porifera, Cnidaria, Mollusca, Arthropoda, and Vertebrata simultaneously.

Discriminating value: This prediction is not generated by any existing sleep function theory (energy conservation, memory consolidation, immune function, or synaptic homeostasis alone). It requires the dimensional invariant framework to generate it.

8.2. P-INVARIANT-2: Mass Extinction Survivor Species Will Show Lower Mean Sleep Duration Than Mass Extinction Casualty Sister Taxa

For each major mass extinction event, comparing the sleep duration of survivor lineages to the estimated sleep duration of closely related casualty lineages will show that Lock survivors sleep significantly less than casualties, and Navigator survivors show significantly higher dietary breadth than casualties.

Testable using the fossil record for body plan conservatism proxy (Lock), dietary reconstruction from isotope and dental analysis (Navigator), and phylogenetic comparison.

8.3. P-INVARIANT-3: The Unstable Zone Boundary Is Predictable From Sleep Duration and Dietary Breadth Alone

A two-dimensional plot of sleep duration (y-axis) against dietary breadth (x-axis) will show a survival boundary:

- Low sleep + any dietary breadth = Lock strategy = high survival probability
- High sleep + high dietary breadth = Navigator strategy = high survival probability
- High sleep + low dietary breadth = Unstable zone = low survival probability
- Low sleep + low dietary breadth = Lock strategy = maximum survival probability

The boundary between survival and extinction risk is predictable from these two measurements alone.

8.4. P-INVARIANT-4: Extreme Specialists Show No Sleep–Flexibility Correlation

Extreme ecological specialists with high sleep requirements (Axis 2 or 3 cost) will show no correlation between sleep duration and cognitive flexibility or learning rate — unlike moderate- sleep Navigator organisms, where sleep duration tracks flexibility. The koala's high sleep pays for detoxification. The lion's high sleep pays for Basin Ω navigation. Same sleep duration. Different function. Distinguishable by flexibility measurement.

8.5. P-INVARIANT-5: Lock Organisms Are Undomesticable

Lock organisms — those in the deepest single basins with minimal sleep requirements — will show the greatest resistance to domestication of any animal group. No Lock organism has ever been domesticated. This is predicted by the dimensional invariant: the same basin depth that allows survival of mass extinctions renders the basin immovable by any domestication field that does not also destroy the organism.

8.6. P-INVARIANT-6: The Dimensional Invariant Is Detectable in Single-Celled Organisms via Metabolic Rest State Duration

Single-celled organisms show metabolic rest states whose duration tracks their niche breadth in the same direction as the invariant predicts for multicellular organisms. Specialist bacteria (obligate extremophiles) will show longer continuous activity periods with minimal rest relative to generalist bacteria (*E. coli*, *B. subtilis*).

This extends the dimensional invariant below the level of nervous system entirely, confirming it as a geometric property of basin topology, not of neural architecture.

9. The Dimensional Invariant: Complete Formal Statement

THE DIMENSIONAL INVARIANT: COMPLETE STATEMENT

Across all life, at all scales, across all of deep time:

Sleep duration (or metabolic rest-state duration in pre-neural organisms) is a measurement of the total cost of an organism's current landscape position, decomposable into:

Axis 1: Between-basin switching cost
 Axis 2: Within-basin metabolic cost
 Axis 3: Within-basin computational intensity

Basin depth — the resistance of the organism's attractor basin to perturbation — is the primary predictor of long-term survival.

The relationship between sleep and survival is U-shaped with a flat minimum:

Minimum sleep → maximum basin depth → maximum survival horizon (Lock strategy).

Moderate sleep → moderate basin depth + multiple accessible basins → long survival horizon (Navigator strategy).

High sleep → deep specialised basin (Axis 2/3) or deep Basin Ω → survival conditional on niche stability or meta- attractor depth.

The invariant holds because:

1. Basin depth is the same geometric property at every scale.
2. Sleep tracks the cost of occupying a landscape position.
3. Deep-time survival is a direct function of basin depth.
4. Therefore sleep duration is a proxy for survival trajectory.

The oldest surviving lineages sleep the least.

This is not a coincidence.

This is the geometry.

10. The Unified Derivation Chain

This document completes a derivation chain that began with a single cancer biology observation and has extended to a universal principle about all life across all of deep time.

Step	Claim	Record
1. Cancer	FOXA1/EZH2 ratio identifies whether a cancer cell is epigenetically arrested in the cancer attractor.	10.5281/zenodo.18883922
2. Domest- tication	The domestication syndrome is field-specified developmental arrest of NCC migration. The pig is the clean case.	10.5281/zenodo.19094935
3. Canine	Above a genomic fixation threshold, the landscape is remodelled. Basin A becomes inaccessible.	Repository 2026-03-18
4. Feline	The symmetric dual-basin landscape. The three-class domestication taxonomy. Basin Ω as meta-attractor.	Repository 2026-03-18
5. Sleep cost	Sleep duration tracks the total cost of an organism's landscape position. Three axes. Universal principle.	Repository 2026-03-18
6. Sleep sonar	The sleep spectrum maps onto the three-axis framework across all animals.	Repository 2026-03-18
7. This document	The dimensional invariant. Sleep predicts survival trajectory across deep time. The oldest surviving lineages sleep the least.	This document

THE UNIFIED STATEMENT

The attractor geometry framework is a single geometric principle operating identically at every scale at which life organises itself.

From a cancer cell deciding whether to differentiate or proliferate, to a domestic pig deciding whether to feral, to a cat choosing which basin to occupy this morning, to a horseshoe crab that has been in the same basin for 450 million years without deciding anything at all:

the geometry is the same.
The basin is the same concept.
The cost of navigation is the same cost.
The sleep is the same payment.
The survival is the same outcome.

Scale is the only variable.
The geometry is invariant.

Linked Records (Confirmed DOIs at Time of Deposit)

Record	DOI / Location
Theory paper	doi:10.5281/zenodo.19094935
P7 pre-registration	doi:10.5281/zenodo.19096128
P9 pre-registration	[DOI pending — insert after Zenodo confirmation]
FOXA1/EZH2 cancer	doi:10.5281/zenodo.18883922
Plant inversion	doi:10.5281/zenodo.19040399
LECA resurrection	doi:10.5281/zenodo.18986790
Canine / feline / sleep ar- tifacts	github.com/Eric-Robert-Lawson/ attractor-oncology (2026-03-18)

Document ID: DIMENSIONAL_INVARIANT_SLEEP_SURVIVAL_REASONING_ARTIFACT_v1.0

Date: 2026-03-18

Author: Eric Robert Lawson / OrganismCore

Status: REASONING ARTIFACT — DERIVATION RECORD. Locked at deposit. No modifications after Zenodo timestamp.

Primary theoretical contributions: (1) The dimensional invariant (oldest surviving lineages sleep the least; minimum sleep = maximum survival horizon; U-shaped survival-sleep relationship with flat minimum). (2) The two-strategy taxonomy (Lock and Navigator). (3) The unstable zone identification and boundary conditions. (4) Predictions P-INVARIANT-1 through P-INVARIANT-6.

Theory paper: [doi:10.5281/zenodo.19094935](https://doi.org/10.5281/zenodo.19094935)

Repository: github.com/Eric-Robert-Lawson/attractor-oncology

ORCID: [0009-0002-0414-6544](https://orcid.org/0009-0002-0414-6544)

Contact: OrganismCore@proton.me

License: CC BY 4.0

Version: 1.0 — LOCKED AT PRE-REGISTRATION.