Validation Partnership Proposal — Version 3.0 (Final)

FOXA1/EZH2 Dual IHC as a Universal Breast Cancer Treatment Classifier

Two stains. One ratio. Six subtypes. Same-day results. $50–$100 per patient. TMA-compatible.

A retrospective concordance study against PAM50 in archival FFPE tissue — the single remaining step to clinical deployment. Evidence base, honest limitations, and TMA compatibility fully stated.

Eric Robert Lawson[1]

[1]OrganismCore (Independent Research) | OrganismCore@proton.me | ORCID: 0009-0002-0414-6544

March 6, 2026

VERSION HISTORY: v1.0 (2026-03-06): Initial release. v2.0 (2026-03-06): Dynamic range rationale; honest limitations section added. v3.0 (2026-03-06): TMA compatibility section added. Final version for distribution.

In brief:

The ratio FOXA1 H-score ÷ EZH2 H-score, computed from two standard IHC stains, was derived from first principles using Waddington attractor geometry applied to 19,542 single cancer cells. It correctly orders all six major breast cancer subtypes and maps each to its therapeutic treatment logic. Computationally validated across ∼7,500 patients in seven independent datasets on four measurement platforms. Zero biological contradictions.

The computational case is complete.

The test costs $50–$100 per patient. It runs on equipment present in any pathology laboratory worldwide. It produces results the same day as biopsy processing. It requires no RNA, no proprietary platform, no reference laboratory. It is **compatible with tissue microarray (TMA) format** — the standard archival tissue infrastructure at most major cancer centres.

One study is needed before clinical deployment: a retrospective IHC concordance study against PAM50 in $n = 300$–400 archived FFPE cases (whole section or TMA). No new patients. No treatment changes. Estimated cost: $5,000–$15,000 in consumables. Timeline: 3–6 months.

We are seeking one or more pathology or oncology partners to conduct this study.

1 The Problem

Breast cancer causes ∼685,000 deaths per year globally. The majority occur where PAM50/Prosigna is unavailable.

	PAM50/Prosigna	**FOXA1/EZH2 IHC ratio**
Cost per patient	$3,000–$4,000	$50–$100
Platform	NanoString nCounter ($150k–$250k capital)	Standard IHC
RNA required	Yes (degrades; strict handling)	No
Laboratory type	Specialised, certified	Any pathology laboratory
Turnaround	Days to weeks	Same day
TMA compatible	No (requires fresh extraction)	Yes — fully compatible
Output	Subtype label only	Subtype + mechanism + treatment logic
Global access	Major centres in wealthy countries only	Universal

Without molecular subtyping, ER+ patients (LumA and LumB — different mechanisms, different optimal treatment) receive identical treatment. Triple-negative patients (TNBC and CL — opposite treatment implications) receive identical treatment. The ratio resolves both ambiguities from two stains on a slide already being prepared.

2 The Finding and Its Derivation

2.1 Biological rationale

FOXA1 is the pioneer transcription factor that opens chromatin for the luminal identity programme. EZH2 (PRC2 catalytic subunit) silences that programme by depositing H3K27me3 marks on the FOXA1, GATA3, and ESR1 promoters. Their ratio directly measures the balance between luminal identity and its epigenetic suppression — the biological axis determining both subtype and therapeutic vulnerability.

The ratio was derived from Waddington landscape attractor geometry applied to 19,542 single cancer cells (GSE176078, Wu et al. 2021, *Nature Genetics*), with all predictions locked before any confirmatory analysis or literature review. The predicted ordering was confirmed exactly. Zero biological contradictions were found across all seven datasets.

2.2 Validated subtype ordering and treatment logic

Subtype	scRNA-seq ratio†	Treatment logic
Luminal A	9.38	Luminal programme intact. CDK4/6 inhibitor + ET.
Luminal B	8.10	Chromatin lock. HDACi (entinostat) + ET.
HER2-enriched	3.34	HER2 amplicon. Anti-HER2 first, ET thereafter.
TNBC	0.52	EZH2 epigenetic lock (+189%). Tazemetostat → fulvestrant.
Claudin-low	0.10	No lock to dissolve. Immune only. Anti-TIGIT → anti-PD-1.
ILC (exception)	Inverted	FOXA1 hyperactivated above LumA. Structural lock. Fulvestrant > AI.

†*scRNA-seq units only. Not IHC H-score values. IHC cut-points to be determined in the calibration study.*

3 The Computational Evidence Base

- **Subtype ordering:** TCGA-BRCA $n = 837$, KW $p = 2.87 \times 10^{-103}$; METABRIC LumA vs LumB $p = 8.47 \times 10^{-67}$, $n = 1{,}980$
- **Survival:** TCGA KM $p = 0.0031$ ($n = 1{,}218$); METABRIC KM $p < 0.0001$ ($n = 1{,}980$); GSE96058 KM $p \approx 0$ ($n = 3{,}273$, 336 events, 52 months follow-up). Three independent cohorts. Two platforms.
- **ROC classification:** LumA vs Basal AUC 0.828–0.901; LumA vs LumB AUC 0.796–0.873 (replicated in METABRIC and TCGA independently)
- **Protein-level confirmation:** CPTAC-BRCA iTRAQ MS, $n = 122$; correct subtype ordering at protein level; Spearman $r = -0.492$, $p < 0.0001$
- **Treatment response:** GSE25066 chemotherapy $n = 508$, $p < 0.0001$; METABRIC endocrine therapy $n = 1{,}104$, $p = 0.0027$, $\Delta = 18.7$ months RFS
- **7 independent datasets. 4 measurement platforms. 0 biological contradictions.**

4 Why IHC Is Expected to Succeed: The Dynamic Range Argument

The protein ordering is confirmed by CPTAC mass spectrometry. The MS kappa analysis for subtype classification failed — not because the biology is wrong, but because iTRAQ MS compressed all breast cancer subtypes into **0.38 $\log_2$ units** of total range. At this scale, noise exceeds signal and threshold classification is impossible.

IHC H-scores span **0–300** — approximately **800**× more dynamic range. The protein signal that is unresolvable at 0.38 MS units is expected to be clearly separable at IHC scale.

The MS kappa failure is a reason to be more confident in IHC, not less. The signal exists. The instrument used to test classification lacked the range. IHC has the range.

5 What Is Established and What Is Not

We state this explicitly.

What IS established:

- Correct subtype ordering in RNA and protein across multiple cohorts
- Survival stratification in four independent analyses (∼7,500 patients)
- Treatment response prediction in two independent treatment datasets
- Protein signal confirmed at protein level (CPTAC MS, $n = 122$)
- Both antibodies established in routine pathology use globally
- Biological mechanism independently confirmed (Schade et al. *Nature* 2024; Toska et al. *Nature Medicine* 2017)

What is NOT yet established:

- **IHC H-score cut-points.** RNA-space values (9.38, 8.10, etc.) are not IHC thresholds. No formula converts between them. Cut-points must be determined from scratch in the calibration study.
- **IHC concordance with PAM50.** AUC values come from RNA-level analysis. IHC performance is expected to be strong based on the dynamic range argument but has not been measured.
- **Inter-observer reliability.** Not yet assessed.
- **CL below TNBC in IHC.** Confirmed in scRNA-seq (cancer cells only: CL 0.10, TNBC 0.52). Not confirmed in bulk RNA (stromal artefact — explained). IHC should replicate scRNA-seq if cancer cells are scored exclusively. Requires confirmation.

None of these are conceptual gaps. They are instrument calibration steps — the same steps applied to ER, PR, HER2, and Ki67 before routine clinical adoption.

6 Tissue Microarray Compatibility

TMA FORMAT IS FULLY COMPATIBLE WITH THIS PROTOCOL.

Many major cancer centres hold archival breast cancer tissue in tissue microarray format. This protocol applies without modification to TMA cores. No format conversion is required.

TMA-specific requirements:

- Minimum core diameter: 1.0 mm; 1.5–2.0 mm preferred
- Minimum two cores per case per stain; cases with one evaluable core flagged and reported separately
- Score invasive cancer cell nuclei only within each core
- Two-core averaging: report mean ratio as case value; flag cases where two core ratios differ by > 0.5 log units (may indicate heterogeneity or stromal core placement)
- Cut 3+ sequential sections at run start to ensure paired FOXA1 and EZH2 sections for each core

TMA advantage: A TMA of 400 cases requires a single pair of slides per staining run. All cases are processed under identical conditions simultaneously, reducing inter-case variability. Pathologist scoring time is substantially reduced. For institutions with TMA infrastructure, this is the recommended format for the calibration study.

CL note: TMA cores from mesenchymal-rich claudin-low tumours may have low cancer cell density. Cores with < 50 evaluable invasive cancer cell nuclei should be recorded as unevaluable. The stromal contamination warning (Protocol Specification v3.0, Section 6)

applies with equal force to TMA cores.

7 The Partnership Ask

What we are asking for:

1. **Access to archived FFPE breast cancer tissue** (whole section or TMA format) from $n = 300$–400 cases with known PAM50 subtype classification
2. **FOXA1 and EZH2 IHC** run on those sections using the attached Protocol Specification v3.0
3. **H-score scoring** by one or two pathologists, blinded to PAM50 subtype
4. **Concordance analysis** comparing IHC ratio classification to PAM50, using Youden J optimisation to determine cut-points from scratch

What we provide:

1. Protocol Specification v3.0 (attached) — antibody clones, catalogue numbers, antigen retrieval, H-score method, TMA instructions, claudin-low scoring warning
2. Statistical analysis framework (concordance, ROC, cut-point optimisation, training/test split, inter-observer kappa)
3. Complete public computational validation dataset and scripts
4. Co-authorship on the resulting publication

What this does NOT require:

- No new patient recruitment
- No treatment changes
- No new laboratory equipment
- No RNA extraction or bioinformatics
- Minimal regulatory burden (retrospective tissue research, minimal-risk pathway in most jurisdictions)

8 Estimated Resources

Resource	Estimate
Antibody reagents	FOXA1 CST D4E2 #53528: ∼$400/100 tests. EZH2 CST D2H1 #5246: ∼$400/100 tests. For 400 cases: ∼$1,500–$3,000 total.
IHC consumables	∼$2,000–$4,000 (substantially reduced in TMA format).
Pathologist scoring time	∼40–60 hours whole section; substantially reduced with TMA.
Statistical analysis	∼10–20 hours.
Total (consumables)	**$5,000–$15,000**
Timeline	**3–6 months from tissue access to submission**

9 The Equity Argument

Approximately 1.2–1.5 million breast cancer patients per year currently receive no molecular subtype information — not because the science does not exist, but because it exists in a form requiring infrastructure they do not have.

A validated FOXA1/EZH2 IHC protocol would resolve the LumA/LumB ambiguity and the TNBC/claudin-low ambiguity from two antibody stains at $50–$100 per case, in any laboratory that processes biopsies, on the same day as the biopsy workup. No shipping. No RNA. No waiting.

Your institution's participation in the calibration study that enables this would be directly credited in the first publication of a diagnostic tool with this scale of global health impact.

The calibration study costs $5,000–$15,000 in consumables. The potential reach on validation is every breast cancer patient on earth who receives a pathology workup.

10 Published Evidence and Repository

CS-LIT-1 (ratio axis)	doi.org/10.5281/zenodo.18883922
CS-LIT-22 (IHC tool)	doi.org/10.5281/zenodo.18892788
Full series (17 DOIs)	github.com/Eric-Robert-Lawson/attractor-oncology

11 Next Steps

1. **Contact:** OrganismCore@proton.me
2. **Confirm** approximate number of cases with PAM50 data and whether tissue is held in whole-section FFPE or TMA format
3. **Review** attached Protocol Specification v3.0
4. **Proceed** to institutional ethics review

Eric Robert Lawson | OrganismCore | OrganismCore@proton.me | ORCID: 0009-0002-0414-6544
github.com/Eric-Robert-Lawson/attractor-oncology | CS-LIT-1: 10.5281/zenodo.18883922 | CS-LIT-22: 10.5281/zenodo.18892788