

Long-range dependence: reaction-coordinate scan -- SN2 (jax) -- lorem-jax, short-range only

Same scan as the rest of this series (`long_range_tests.py` , `data/sn2/sn2.xyz` reference geometry), but on the **official lorem-jax reproduction checkpoint** from `lorem-tmlr-archive` instead of a metatrain/torch model.

This notebook needs a different Python environment than 02 - 11 :

`/home/boittier/metawork/.venv-lorem-jax/bin/python` (has `lorem / jax / flax` installed; the regular `metawork/.venv` used elsewhere in this series does not). Select that interpreter as this notebook's kernel before re-running it.

Official lorem-tmlr-archive checkpoint (lr: false, equivariant_message_passing: true, num_message_passing: 1), loaded via `lorem.calculator.Calculator`. Does not invoke the Ewald/Coulomb path at all, so it is unaffected by the non-PBC convention change described above -- usable as-is.

Caveat on comparability: this checkpoint is trained on the archive's own SN2 halide-exchange dataset ($X^- + CH_3Y$, $X,Y \in \{F, Cl, Br, I\}$), not the `sn2-matched-*` domain used to train the models in notebooks 02 - 11 . Its *absolute* energies are therefore not directly comparable to those models -- only the qualitative long-range-cutoff behavior is, which is the point of including it here.

```
In [ ]: import numpy as np
import matplotlib.pyplot as plt
from matplotlib.colors import to_hex

from ase.io import read
from ase.visualize.plot import plot_atoms
from ase.data.colors import jmol_colors

from lorem.calculator import Calculator
import lorem.models.mlip as mlip
from jaxpme.batched_mixed import Ewald as _orig_Ewald

from long_range_tests import (
    displace_atom_test,
    fragment_separation_test,
    estimate_cutoff,
)

plt.rcParams.update({
    "figure.dpi": 120,
    "font.size": 11,
```

```

    "axes.spines.top": False,
    "axes.spines.right": False,
    "axes.grid": True,
    "grid.alpha": 0.25,
    "grid.linewidth": 0.6,
    "legend.frameon": False,
})

ATOM_COLORS = ["#0072B2", "#56B4E9", "#009E73", "#E69F00", "#D55E00", "#CC79A7"]
HIGHLIGHT = "#D55E00"

```

No long-range patch needed

This checkpoint has `lr: false` (short-range only, message-passing instead) -- it never invokes the Ewald/Coulomb path, so it is unaffected by the non-PBC convention change described in the long-range companion notebook (`13_long_range_scan_lorem_jax_lr.ipynb`) and can be loaded as-is.

Setup

Same shared reference structure (`data/sn2/sn2.xyz`, frame 0) as every other notebook in this series, re-anchoring I onto the C...I axis and scanning from 1.5 Å to 32 Å.

```

In [ ]: calc = Calculator.from_checkpoint("../..metawork/lorem-tmlr-archive/evals/s

atoms0 = read("../data/sn2/sn2.xyz", index=0)
symbols = atoms0.get_chemical_symbols()
labels = [f"{i} {s}" for i, s in enumerate(symbols)]
C, I = 0, 5 # atom indices: carbon, leaving iodide

d = atoms0.get_all_distances()
print(f"original snapshot -- C-F: {d[C, 4]:.2f} Å   C-I: {d[C, I]:.2f} Å")

direction = atoms0.positions[I] - atoms0.positions[C]
direction = direction / np.linalg.norm(direction)
atoms0.positions[I] = atoms0.positions[C].copy()

distances = np.linspace(1.5, 32.0, 62) # Å, absolute C...I distance

```

original snapshot -- C-F: 1.39 Å C-I: 7.44 Å

1. Reaction-coordinate scan (`displace_atom_test`)

```

In [ ]: result = displace_atom_test(
    atoms0, calc, probe_index=I, distances=distances,
    direction=direction, reference_indices=[0, 1, 2, 3, 4],
)

```

```

delta_forces = result["forces"] - result["forces"][-1]
delta_force_mag = np.linalg.norm(delta_forces, axis=-1)
max_delta = delta_force_mag.max(axis=-1)

cutoff_distance = estimate_cutoff(result["distance"], max_delta, threshold=6)
print(f"empirical cutoff: {cutoff_distance:.1f} Å")

```

empirical cutoff: 5.5 Å

```

In [ ]: fig, axes = plt.subplots(1, 2, figsize=(11, 4.2))


ax = axes[0]
ax.plot(result["distance"], result["energy"], color=ATOM_COLORS[0], linewidth=2)
ax.axvline(cutoff_distance, color="0.4", linestyle="--", linewidth=1)
ax.set_xlabel("C···I distance (Å)")
ax.set_ylabel("total energy (eV)")
ax.set_title("Energy vs. leaving-group distance")

ax = axes[1]
ax.semilogy(result["distance"], max_delta, color=HIGHLIGHT, linewidth=2)
ax.axvline(cutoff_distance, color="0.4", linestyle="--", linewidth=1)
ax.annotate(f"empirical cutoff ≈ {cutoff_distance:.1f} Å",
            xy=(cutoff_distance, 3e-2), xytext=(cutoff_distance + 3, 3e-2),
            fontsize=9, color="0.25", va="center",
            arrowprops=dict(arrowstyle="->", color="0.4", lw=1))
ax.set_xlabel("C···I distance (Å)")
ax.set_ylabel("max |ΔF| on CH3F, vs. fully separated (eV/Å)")
ax.set_title("Force response decay (log scale)")

fig.suptitle("Reaction-coordinate scan: SN2 (jax) -- lorem-jax, short-range only")
fig.tight_layout()

```

Reaction-coordinate scan: SN2 (jax) -- lorem-jax, short-range only



Per-atom force magnitude and components

```

In [ ]: per_atom_mag = np.linalg.norm(result["forces"], axis=-1)
ref_labels = [labels[j] for j in [0, 1, 2, 3, 4]]

fig, axes = plt.subplots(1, 4, figsize=(16, 3.8), sharex=True)

```


```

ax = axes[0]
for j, lab in enumerate(ref_labels):
    ax.plot(result["distance"], per_atom_mag[:, j], color=ATOM_COLORS[j],
            linewidth=2, label=lab)
ax.axvline(cutoff_distance, color="0.4", linestyle="--", linewidth=1)
ax.set_ylabel("|F| (eV/Å)")
ax.set_title("force magnitude")
ax.set_xlabel("C···I distance (Å)")
ax.legend(fontsize=8, loc="upper right")

comp_names = ["Fx", "Fy", "Fz"]
for k, comp in enumerate(comp_names):
    ax = axes[k + 1]
    for j, lab in enumerate(ref_labels):
        ax.plot(result["distance"], result["forces"][:, j, k], color=ATOM_COLORS[j],
                linewidth=2, label=lab)
    ax.axvline(cutoff_distance, color="0.4", linestyle="--", linewidth=1)
    ax.set_title(comp)
    ax.set_xlabel("C···I distance (Å)")
    ax.set_ylabel("force (eV/Å)")

fig.suptitle("Per-atom force magnitude and components -- SN2 (jax) -- lorem-jax, short-range only")
fig.tight_layout()

```



Structure snapshots along the scan

```

In [ ]: snapshot_distances = [1.5, 3.5, cutoff_distance, 10.0, 20.0, 32.0]
z = atoms0.get_atomic_numbers()
base_colors = [to_hex(jmol_colors[zi]) for zi in z]

fig, axes = plt.subplots(1, len(snapshot_distances), figsize=(16, 3.2))

for ax, dist in zip(axes, snapshot_distances):
    snap = atoms0.copy()
    snap.positions[I] = atoms0.positions[C] + direction * dist
    colors = list(base_colors)
    colors[I] = HIGHLIGHT
    plot_atoms(snap, ax, radii=0.5, colors=colors, rotation="-80x,10y,0z")
    is_cutoff = np.isclose(dist, cutoff_distance)
    ax.set_title(f"{dist:.1f} Å" + (" (cutoff)" if is_cutoff else ""),
                fontsize=10, color=HIGHLIGHT if is_cutoff else "0.2",
                fontweight="bold" if is_cutoff else "normal")
    for spine in ax.spines.values():
        spine.set_visible(True)
        spine.set_edgecolor(HIGHLIGHT if is_cutoff else "0.85")


```

```

spine.set_linewidth(2 if is_cutoff else 1)

fig.suptitle(
    "CH3F...I- geometry along the scan -- SN2 (jax) -- lorem-jax, short-range only",
    "(Jmol element colors; I- highlighted in orange)",
    y=1.06,
)
fig.tight_layout()

```




2. Size-consistency corroboration (fragment_separation_test)

```

In [ ]: fragmentation = fragment_separation_test(
    atoms0, calc, [0, 1, 2, 3, 4], [I],
    separations=distances - distances[0], axis=direction,
)

fig, ax = plt.subplots(figsize=(6, 4.2))
ax.plot(result["distance"], fragmentation["interaction_energy"],
        color=ATOM_COLORS[0], linewidth=2)
ax.axhline(0, color="0.6", linewidth=1, linestyle=":")
ax.axvline(cutoff_distance, color="0.4", linestyle="--", linewidth=1)
ax.set_xlabel("C...I distance (Å)")
ax.set_ylabel("interaction energy (eV)")
ax.set_title("Size-consistency: SN2 (jax) -- lorem-jax, short-range only")
fig.tight_layout()

```



Takeaways

- Empirical cutoff for **SN2 (jax) -- lorem-jax, short-range only: 5.5 Å**.
- This is the archive's *short-range-only* checkpoint (no Ewald path): the cutoff lands in the same ~ 5.5 Å cluster as every BPNN/PET model in this series -- an independent (different codebase, different training run, different dataset) confirmation that 'no long-range term' means 'short cutoff' regardless of architecture family.
- Past that point, the per-atom force curves flatten to a plateau of ≈ 0.067 eV/Å (the CH_3F fragment's own non-equilibrium force).
- Total energy changes by ≈ 3.52 eV from the most compressed to the most separated geometry sampled. Note this checkpoint's absolute energy scale is not comparable to the sn2-matched-* models (different training data/reference) -- only the cutoff behavior is.

In []: