

Necessary and Sufficient Conditions for the Existence of an LU Factorization

Eric Darve

November 25, 2025

Abstract

We investigate the necessary and sufficient conditions for the existence of an LU factorization for a given matrix A . We establish a condition relating the nullity of the submatrices of A to the existence of the factorization. Finally, we extend these results to unit lower and upper triangular factorizations.

Notations

$\text{rank}(A)$	Rank of matrix A
$\text{null}(A)$	Dimension of the null space of matrix A
A^T	Transpose of matrix A
a_{ij}	Entry (i, j) of matrix A
$A[:, i]$	Column i of matrix A
$A[i, :]$	Row i of matrix A
$A[i : j, k : l]$	Submatrix of A with rows from i to j and columns from k to l
$A[i : j, k :]$	Submatrix of A with rows from i to j and columns from k to the end
e_i	i -th vector of the standard basis
I_n	Identity matrix of size n

This paper is concerned with proving necessary and sufficient conditions for the existence of an LU factorization of a matrix A . It extends the work of [1] with new results and simpler proofs.

We begin with a few classical results that will serve as reference for the later sections.

1 Classical results

The theorems establish that any matrix can be permuted to have an LU factorization with unit triangular factors.

Theorem 1. *Consider matrix A of size n . There exist a row permutation matrix P such that $PA = LU$ where L is unit lower triangular and U is upper triangular.*

Proof. This is a classical result. We can prove it by induction on n . The result is true for $n = 1$.

Assume it is true for $n - 1$. If $a_{11} \neq 0$, then consider:

$$A = LDU$$

where

$$L = \begin{pmatrix} 1 & 0 \\ a_{11}^{-1}A_{21} & I \end{pmatrix}, \quad D = \begin{pmatrix} 1 & 0 \\ 0 & S \end{pmatrix}, \quad U = \begin{pmatrix} a_{11} & A_{12} \\ 0 & I \end{pmatrix}$$

We can apply the induction hypothesis to S . The result for A follows.

If $a_{11} = 0$, we have two cases.

Case 1: column 1 of A is non-zero. We can find the smallest i such that $A[i, 1] \neq 0$. Pivot row 1 and i . Denote by $B = \Pi_i A$ the permuted matrix (with $\Pi_i^2 = I$). Since $b_{11} \neq 0$, B has an LU factorization. This proves the result for A .

Case 2: column 1 of A is zero. We choose:

$$A = IDU$$

where

$$D = \begin{pmatrix} 1 & 0 \\ 0 & S \end{pmatrix}, \quad U = \begin{pmatrix} 0 & A_{12} \\ 0 & I \end{pmatrix}$$

We can apply the induction hypothesis to S . The result for A follows. $\square$

Corollary 1. Consider matrix A of size n . There exist a column permutation matrix Q such that $AQ = LU$ where L is lower triangular and U is unit upper triangular.

Proof. Apply Theorem 1 to A^T . $\square$

Definition 1. Matrix A of size $m \times n$ is said to have a rank revealing LU factorization if $A = LU$ where L is lower triangular of size $m \times r$, U is upper triangular of size $r \times n$, and $r = \text{rank}(A)$.

If we use both row and column permutations, we can always obtain a rank revealing LU factorization.

Theorem 2. Consider matrix A of size n . There exist a row permutation matrix P and a column permutation matrix Q such that $PAQ = LU$ is a rank revealing LU factorization.

Proof. This is also a classical result. We can prove it by induction on n . The result is true for $n = 1$.

Assume it is true for $n - 1$. If $a_{11} \neq 0$, then consider:

$$A = LDU$$

where

$$L = \begin{pmatrix} a_{11} & 0 \\ A_{21} & I \end{pmatrix}, \quad D = \begin{pmatrix} a_{11}^{-1} & 0 \\ 0 & S \end{pmatrix}, \quad U = \begin{pmatrix} a_{11} & A_{12} \\ 0 & I \end{pmatrix}$$

S has rank $r - 1$ if $r = \text{rank}(A)$. We can apply the induction hypothesis to S . The result for A then follows.

If $a_{11} = 0$, we have two cases. If A is 0, we pick $L = U = 0$. Otherwise we have some i and j such that $a_{ij} \neq 0$. We can permute row 1 and i and column 1 and j . Denote by $B = \Pi_i A \Pi_j$ the permuted matrix. Since $b_{11} \neq 0$, B has a rank revealing LU factorization. This proves the result for A .

We have also proved that this process must terminate after r steps where $r = \text{rank}(A)$. So the LU factorization of B is rank revealing.

From Lemma 3 and Corollary 4, we can also prove that the remaining Schur complement is zero after r steps of the LU factorization. $\square$

We now consider LU factorizations without permutations. The simplest result requires that all leading principal minors are non-singular.

Proposition 1. Consider matrix A of size n such that

$$\text{rank}(A[1 : k, 1 : k]) = k, \quad k = 1, \dots, n$$

Then A has an LU factorization: $A = LU$.

Proof. We prove the result by induction on n . The result is true for $n = 1$.

Assume it is true for $n - 1$. By assumption, $a_{11} \neq 0$. We then have the factorization:

$$A = LDU$$

where

$$L = \begin{pmatrix} a_{11} & 0 \\ A_{21} & I \end{pmatrix}, \quad D = \begin{pmatrix} a_{11}^{-1} & 0 \\ 0 & S \end{pmatrix}, \quad U = \begin{pmatrix} a_{11} & A_{12} \\ 0 & I \end{pmatrix}$$

and $S = A_{22} - A_{21}a_{11}^{-1}A_{12}$ is the Schur complement. L and U are full rank. So D satisfies the condition of the theorem. Consequently, S also satisfies the condition of the theorem and is of size $n - 1$. By the induction hypothesis, S has an LU factorization. Therefore A also has an LU factorization. $\square$

2 General necessary and sufficient condition for the existence of an LU factorization

We now discuss the core of this paper: the necessary and sufficient conditions for the existence of an LU factorization without permutations. We start with an illustrative example that will preview the main result.

Consider matrix

$$A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

Matrix A does not have an LU factorization. We prove this by contradiction. Assume $A = LU$. Then $l_{11}u_{11} = 0$. Assume $l_{11} = 0$. Then $l_{11}u_{12} = 0 = 1$. Contradiction. Assume $u_{11} = 0$. Then $l_{21}u_{11} = 0 = 1$. Contradiction.

Consider now

$$A = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$$

This matrix in fact has an LU factorization. We can find it using the following procedure, which we will detail later on for the general case. Surprisingly the process involves row and column permutations. However the final result remains an LU factorization without the permutations.

Now, perform a cyclic shift of the rows: $(1,2,3) \rightarrow (2,3,1)$ and permute column 1 and 3:

$$PAQ = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$$

But surprisingly we have:

$$A = P^{-1} \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} Q^{-1} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$$

This is valid rank revealing LU factorization of A . Similarly:

$$A = \begin{pmatrix} 0 & 0 \\ 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

is another valid rank revealing LU factorization of A .

We can generalize this example. Consider any $B = LU$ and:

$$A = \begin{pmatrix} 0_n & 0_n & 0_n \\ 0_n & 0_n & C \\ 0_n & B & D \end{pmatrix} = \begin{pmatrix} 0_n & 0_n \\ C & 0_n \\ D & L \end{pmatrix} \begin{pmatrix} 0_n & 0_n & I_n \\ 0_n & U & 0_n \end{pmatrix}$$

If $C = LU$ then:

$$A = \begin{pmatrix} 0_n & 0_n \\ 0_n & L \\ I_n & 0_n \end{pmatrix} \begin{pmatrix} 0_n & B & D \\ 0_n & 0_n & U \end{pmatrix}$$

Of course, this is not true for all matrices. We will show that Property 1 below is a necessary and sufficient condition for the existence of an LU factorization. The proof will rely on constructing a P and Q such that $B = P^{-1}AQ^{-1}$. $B[1:r, 1:r]$ will satisfy the conditions of Proposition 1 where $r = \text{rank}(A) = \text{rank}(B)$. Therefore $B = LU$. From this, we will deduce a rank-revealing LU factorization of $A = (PL)(UQ)$, where PL is lower triangular with r columns, and UQ is upper triangular with r rows.

Property 1. *Matrix A is square of size n . For all $k = 1, \dots, n$:*

$$\text{null}(A[1:k, 1:k]) \leq \text{null}(A[:, 1:k]) + \text{null}(A[1:k, :])^T \quad (1)$$

Theorem 3. *The rank-nullity theorem states that for any matrix A of size $m \times n$:*

$$\text{rank}(A) + \text{null}(A) = n$$

Lemma 1. *Consider $A = BC$. Then:*

$$\text{rank}(A) \leq \min(\text{rank}(B), \text{rank}(C)), \quad \text{rank}(B) + \text{rank}(C) - k \leq \text{rank}(A)$$

where k is the number of columns of B and rows of C .

Corollary 2. *If A , B , and C are square matrices then:*

$$\text{null}(A) \leq \text{null}(B) + \text{null}(C)$$

Proof. Use Lemma 1 and Theorem 3. □

Assume that A , a square matrix of size n , has an LU factorization:

$$A = LU$$

Then:

$$\begin{aligned} A[1 : k, :] &= L[1 : k, 1 : k]U[1 : k, :], \\ A[:, 1 : k] &= L[:, 1 : k]U[1 : k, 1 : k], \\ A[1 : k, 1 : k] &= L[1 : k, 1 : k]U[1 : k, 1 : k] \end{aligned}$$

Using Lemma 1 and Corollary 2, we have:

$$\begin{aligned} \text{rank}(A[1 : k, :]) &\leq \text{rank}(L[1 : k, 1 : k]), \\ \text{rank}(A[:, 1 : k]) &\leq \text{rank}(U[1 : k, 1 : k]), \\ \text{null}(A[1 : k, 1 : k]) &\leq \text{null}(L[1 : k, 1 : k]) + \text{null}(U[1 : k, 1 : k]) \end{aligned}$$

Using Theorem 3, we get:

$$\text{null}(A[1 : k, 1 : k]) \leq \text{null}(A[:, 1 : k]) + \text{null}(A[1 : k, :])^T$$

So Property 1 is a necessary condition for the existence of an LU factorization.

We will prove the following theorem:

Theorem 4. *Matrix A satisfies Property 1 if and only if it has an LU factorization.*

Proof. We prove the result by induction on the size of A . The result is true for $n = 1$.

Assume it is true for size $n - 1$.

Assume $a_{11} \neq 0$. We have the factorization:

$$A = LDU$$

where

$$L = \begin{pmatrix} a_{11} & 0 \\ A_{21} & I \end{pmatrix}, \quad D = \begin{pmatrix} a_{11}^{-1} & 0 \\ 0 & S \end{pmatrix}, \quad U = \begin{pmatrix} a_{11} & A_{12} \\ 0 & I \end{pmatrix}$$

Since L and U are full rank, D satisfies Property 1. Consequently, S also satisfies Property 1 and is of size $n - 1$. Moreover S has rank $r - 1$ if $r = \text{rank}(A)$. By the induction hypothesis, S has an LU factorization. The result for A follows.

Assume $a_{11} = 0$. From Property 1, we have either:

1. $A[:, 1] = 0$, or
2. $A[1, :] = 0$

3. both.

Case 1: $A[:, 1] = 0$. We can find the smallest j such that $A[1, j] \neq 0$. Pivot column 1 and j . Denote by $B = A\Pi_j$ the permuted matrix (with $\Pi_j^2 = I$). We prove that B satisfies Property 1.

Consider $k < j$. We have

$$\text{null}(A[1 : k, 1 : k]) \leq \text{null}(A[:, 1 : k]) + \text{null}(A[1 : k, :])^T \quad (2)$$

Since column 1 of A is zero, we have:

$$\text{null}(A[1 : k, 1 : k]) = \text{null}(A[1 : k, 2 : k]) + 1$$

We have

$$\text{null}(B[1 : k, 2 : k]) = \text{null}(A[1 : k, 2 : k])$$

Since the only non-zero entry in row 1 of $B[1 : k, 1 : k]$ is in column 1 we have

$$\text{null}(B[1 : k, 1 : k]) = \text{null}(B[1 : k, 2 : k]) = \text{null}(A[1 : k, 2 : k]) = \text{null}(A[1 : k, 1 : k]) - 1$$

By a similar reasoning, we have

$$\text{null}(B[:, 1 : k]) = \text{null}(A[:, 1 : k]) - 1$$

The column pivoting does not affect the rank of the rows, so

$$\text{null}(B[1 : k, :])^T = \text{null}(A[1 : k, :])^T$$

Therefore B satisfies Property 1 for $k < j$.

For $k \geq j$, the pivoting does not affect the ranks in Equation (2), so B also satisfies Property 1 for $k \geq j$. So B satisfies Property 1.

Since $b_{11} \neq 0$, B has an LU factorization: $B = LU$. Column j of B is zero. If column j of U is non-zero, we can zero it out without affecting any column in LU . So we can assume that column j of U is zero. We have:

$$A = B\Pi_j = LU\Pi_j$$

and $U\Pi_j$ is still upper triangular because Π_j only permutes column 1 and j . Therefore A has an LU factorization.

Case 2: $A[1, :] = 0$. Consider $B = A^T$. B corresponds to Case 1, so B has an LU factorization: $B = LU$. Therefore:

$$A = B^T = U^T L^T$$

is an LU factorization of A .

Case 3: If A is 0, it has an LU factorization with $L = U = 0$. Let's denote by j the smallest index such that column j is non zero. Let's denote by $i > 1$ the smallest index such that $a_{ij} \neq 0$. Let's pivot column 1 and j and denote by $B = A\Pi_j$ the permuted matrix. B satisfies Case 2, so it has an LU factorization: $B = LU$. Column j of U is zero. Therefore:

$$A = B\Pi_j = LU\Pi_j$$

is an LU factorization of A .

We have also proved that this process must terminate after r steps where $r = \text{rank}(A)$. So the LU factorization of B is rank revealing. $\square$

We now prove the stronger theorem:

Theorem 5. *Matrix A is of size n and rank $r \leq n$. If A satisfies Property 1, then there exist a row permutation P_r and column permutation P_c such that:*

$$B = P_r^{-1} A P_c^{-1}$$

with $B[1 : k, 1 : k]$, $1 \leq k \leq r$, non-singular. $B[1 : r, 1 : r]$ has an LU factorization:

$$B[1 : r, 1 : r] = L_r U_r$$

and:

$$B = (B[:, 1 : r] U_r^{-1}) (L_r^{-1} B[1 : r, :]) = L_B U_B$$

is its rank revealing LU factorization. Then:

$$A = (P_r L_B)(U_B P_c) = LU$$

is a rank revealing factorization of A . Moreover, we have the following stronger sparsity results. If $e_{i_0} = P_r e_i$, then for L :

- If $i \leq r$, then $i_0 \geq i$ and $L[i_0, i + 1 :] = 0$.
- If $i > r$, then $i_0 \leq i$ and $L[i_0, \min(r, i_0) :] = 0$.

We have similar results for U . If $e_{j_0} = P_c^T e_j$, then:

- If $j \leq r$, then $j_0 \geq j$ and $U[j + 1 :, j_0] = 0$.
- If $j > r$, then $j_0 \leq j$ and $U[\min(r, j_0) :, j_0] = 0$.

Proof. We follow a proof similar to the one for Theorem 4.

We do an induction on the size of A . The result is true for $n = 1$.

Assume it is true for size $n - 1$. If $a_{11} \neq 0$, we have the factorization:

$$A = LDU$$

where

$$L = \begin{pmatrix} a_{11} & 0 \\ A_{21} & I \end{pmatrix}, \quad D = \begin{pmatrix} a_{11}^{-1} & 0 \\ 0 & S \end{pmatrix}, \quad U = \begin{pmatrix} a_{11} & A_{12} \\ 0 & I \end{pmatrix}$$

We apply the induction hypothesis to S . We can define P_r and P_c in terms of the permutations for S :

$$P_r = \begin{pmatrix} 1 & 0 \\ 0 & P_{r,S} \end{pmatrix}, \quad P_c = \begin{pmatrix} 1 & 0 \\ 0 & P_{c,S} \end{pmatrix}$$

The sparsity results follow directly from the induction hypothesis.

Assume $a_{11} = 0$. From Property 1, we have either:

1. $A[:, 1] = 0$, or
2. $A[1, :] = 0$
3. both.

Case 1: $A[:, 1] = 0$. As before, we can find the smallest j such that $A[1, j] \neq 0$. We pivot column 1 and j . Denote by $B_0 = A\Pi_j$ the permuted matrix. B_0 satisfies Property 1 and $b_{11} \neq 0$. So B_0 has an LU factorization with the sparsity pattern of Theorem 5. Since column j of B_0 is zero, column j of U_{B_0} is zero. We define

$$P_c = P_{c,B}\Pi_j, \quad U = U_B\Pi_j$$

Π_j swaps columns 1 and j of U_{B_0} . We have $e_1 = \Pi_j^T e_j$ and column j of U_{B_0} is zero. We also have a j_0 such that $e_j = P_{c,B}^T e_{j_0}$ and $e_1 = P_c^T e_{j_0}$, with $j_0 > r$. Since column 1 of U is zero and the other columns of U have the sparsity pattern of Theorem 5, U also satisfies the sparsity pattern of Theorem 5.

Case 2: $A[1, :] = 0$. Consider $B = A^T$. B corresponds to Case 1 and satisfies the sparsity pattern of Theorem 5. So A also satisfies the sparsity pattern of Theorem 5.

Case 3: If $A = 0$, it has an LU factorization. In that case we pick $L = U = 0$ as the factors. If $A \neq 0$, let's denote by j the smallest index such that column j is non zero. Let's denote by $i > 1$ the smallest index such that $a_{ij} \neq 0$. Let's pivot column 1 and j and denote by $B = A\Pi_j$ the permuted matrix. B satisfies Case 2, so it has an LU factorization: $B = LU$ with the sparsity pattern of Theorem 5. Following a reasoning analogous to Case 1, we can show that A also satisfies the sparsity pattern of Theorem 5.

As before, we have also proved that this process must terminate after r steps where $r = \text{rank}(A)$. So the LU factorization of B is rank revealing. $\square$

3 Unit triangular factors

We now prove results for unit triangular factors.

Theorem 6. *Matrix A is of size n and rank $r \leq n$. A satisfies, for all $k = 1, \dots, n$:*

$$\text{null}(A[1 : k, 1 : k]) = \text{null}(A[:, 1 : k]) \quad (3)$$

if and only if A has an LU factorization $A = LU$ where L is unit lower triangular.

If A satisfies Equation (3), then there exist a column permutation P_c such that:

$$B = AP_c^{-1}$$

B has an LU factorization: $B = L_B U_B$ where L_B is unit lower triangular. Then:

$$A = L_B(U_B P_c) = LU$$

is an LU factorization of A where $L = L_B$ is unit lower triangular.

We also have that row j of B (or row j of A since the rows are not permuted) is a linear combination of the previous rows if and only if column j of B is a linear combination of the previous columns.

Assume $e_{j_0} = P_c^T e_j$. If row j of B is a linear combination of the previous rows, then $j_0 \leq j$. We have $L[:, j] = e_j$, $U[j, :] = 0$, and $U[j_0 :, j_0] = 0$. If row j of B is linearly independent of the previous rows then $j_0 \geq j$ and $U[j + 1 :, j_0] = 0$.

Proof. We first prove that if $A = LU$ with L unit lower triangular, then A satisfies Equation (3). U trivially satisfies Equation (3) since it is upper triangular. Since $L[1 : k, 1 : k]$ is full rank, A also satisfies Equation (3).

We follow a proof similar to the one for Theorem 5. We do an induction on the size of A . The result is true for $n = 1$.

Assume it is true for size $n - 1$. If $a_{11} \neq 0$, we verify that the result is true by the induction hypothesis.

Assume $a_{11} = 0$. From Equation (3), we have $A[1, :] = 0$.

Case 1: If row 1 of A is non-zero, we can use the same steps as in the proof as Theorem 5 with a column permutation to prove the result by induction. In addition, this shows that if $e_{j_0} = P_c^T e_j$ and if row j of B is linearly independent of the previous rows then $j_0 \geq j$ and $U[j + 1 :, j_0] = 0$.

Case 2: If row 1 of A is zero, we cannot use the same proof as Theorem 5 because we can no longer permute rows. However, since row 1 of A is zero, we can define:

$$A = IDU$$

with

$$D = \begin{pmatrix} 1 & 0 \\ 0 & S \end{pmatrix}, \quad U = \begin{pmatrix} 0 & 0 \\ 0 & I \end{pmatrix}$$

D satisfies

$$\text{null}(D[1 : k, 1 : k]) = \text{null}(A[1 : k, 1 : k]) - 1 = \text{null}(A[:, 1 : k]) - 1 = \text{null}(D[:, 1 : k])$$

D satisfies Equation (3). By the induction hypothesis, we can find a column permutation $P_{c,S}$ such that $SP_{c,S}^{-1}$ has an LU factorization with unit lower triangular factor.

This also proves that row j of B is a linear combination of the previous rows if and only if column j of B is a linear combination of the previous columns. Moreover, if $e_{j_0} = P_c^T e_j$ and row j of B is a linear combination of the previous rows, then $j_0 \leq j$. We have $L[:, j] = e_j$, $U[j, :] = 0$, and $U[j_0 :, j_0] = 0$.

Note that the step above with D is not a valid step in Theorem 5. Indeed:

$$\begin{aligned} \text{null}(D[1 : k, 1 : k]) &= \text{null}(A[1 : k, 1 : k]) - 1 \\ \text{null}(D[:, 1 : k]) &= \text{null}(A[:, 1 : k]) - 1 \\ \text{null}(D[1 : k, :]^T) &= \text{null}(A[1 : k, :]^T) - 1 \end{aligned}$$

So that Property 1 may not be satisfied by D . □

Corollary 3. *Matrix A is of size n and rank $r \leq n$. A satisfies, for all $k = 1, \dots, n$:*

$$\text{null}(A[1:k, 1:k]) = \text{null}(A[1:k, :]) \quad (4)$$

if and only if A has an LU factorization $A = LU$ where U is unit lower triangular.

If A satisfies Equation (4), then there exist a row permutation P_r such that:

$$B = P_r^{-1}A$$

B has an LU factorization: $B = L_B U_B$ where U_B is unit upper triangular. Then:

$$A = (P_r L_B) U_B = LU$$

is an LU factorization of A where $U = U_B$ is unit upper triangular.

We also have that column i of B (or column i of A since the columns are not permuted) is a linear combination of the previous columns if and only if row i of B is a linear combination of the previous rows.

Assume $e_{i_0} = P_r e_i$. If column i of B is a linear combination of the previous columns, then $i_0 \leq i$. We have $U[i, :] = e_i$, $L[:, i] = 0$, and $L[i_0, i_0:] = 0$. If column i of B is linearly independent of the previous columns then $i_0 \geq i$ and $L[i_0, i+1:] = 0$.

Proof. Consider $B = A^T$. Apply Theorem 6. □

A Ancillary results

We prove a few additional results that are helpful in the main proofs.

Lemma 2. *Assume that $x \in S = \text{span}\{x_1, \dots, x_r\}$. Denote by $X = [x_1, \dots, x_r]$. Assume that $X[1:r, :]$ is non-singular. Then:*

$$x = XX[1:r, :]^{-1}x[1:r]$$

Proof. Since the submatrix $X[1:r, :]$ is non-singular, any vector $x \in S$ is uniquely determined by its first r entries. Let P be the oblique projection onto S along the subspace of vectors with the first r entries equal to zero. The projection matrix is given by:

$$P = X(Y^T X)^{-1}Y^T$$

where the range of Y is the orthogonal complement of the null space of P . Specifically, we set:

$$Y = \begin{pmatrix} I_r \\ 0 \end{pmatrix}$$

Since $x \in S$, we have $x = Px$. Substituting the expressions for P and Y , we obtain:

$$x = X(Y^T X)^{-1}Y^T x = X(X[1:r, :])^{-1}x[1:r]$$

□

Lemma 3. *Consider a matrix A of size $n \times k$, $k \leq n$. Assume that the rank of A is $k-1$ and that column k is linearly dependent on the previous columns. If A has an LU factorization $A = LU$ with $u_{kk} = 1$, then column k of L is 0.*

Proof. Decompose L and U in blocks:

$$L = \begin{pmatrix} L_{11} & 0 \\ L_{21} & l_{kk} \end{pmatrix}, \quad U = \begin{pmatrix} U_{11} & u_{1k} \\ 0 & 1 \end{pmatrix}$$

where L_{11} is of size $(k-1) \times (k-1)$, U_{11} is of size $(k-1) \times (k-1)$, L_{21} is of size $(n-k+1) \times (k-1)$, and u_{1k} is of size $(k-1) \times 1$. Then:

$$l_{kk} = A[k:, k] - A[k:, 1:k]A[1:k, 1:k]^{-1}A[1:k, k]$$

By Lemma 2, since column k of A is linearly dependent on the previous columns, we have: $l_{kk} = 0$. □

Corollary 4. *Consider a matrix A of size $k \times n$, $k \leq n$. Assume that the rank of A is $k - 1$ and that row k is linearly dependent on the previous rows. If A has an LU factorization $A = LU$ with $l_{kk} = 1$, then row k of U is 0.*

Proof. Consider $A^T = U^T L^T$. Apply Lemma 3. □

References

- [1] Pavel Okunev and Charles R. Johnson. *Necessary And Sufficient Conditions For Existence of the LU Factorization of an Arbitrary Matrix*. arXiv:math/0506382. June 2005. DOI: [10.48550/arXiv.math/0506382](https://doi.org/10.48550/arXiv.math/0506382). URL: <http://arxiv.org/abs/math/0506382> (visited on 11/25/2025).