

PROJECT BASED LEARNING-1

PROJECT REPORT

“Site Selection for Solar Energy Plants”

BACHELOR OF TECHNOLOGY IN ARTIFICIAL INTELLIGENCE &
MACHINE LEARNING

Submitted By

Ameya Tipnis (23070126012)

Eric Siqueira (23070126041)

Asho Shibu (23070126023)

Ron Matthew (23070126109)

UNDER THE GUIDANCE OF

Prof. Sachin Gaikwad

Prof. Mayur Gaikwad

Designation



॥वसुधैव कुटुम्बकम्॥

SYMBIOSIS INSTITUTE OF TECHNOLOGY, PUNE

Pune – 412115, Maharashtra State, India

<https://www.sitpune.edu.in/>

DEPARTMENT OF ARTIFICIAL INTELLIGENCE & MACHINE LEARNING

AY 2024-25



SYMBIOSIS INSTITUTE OF TECHNOLOGY, PUNE

Pune – 412115, Maharashtra State, India

<https://www.sitpune.edu.in/>

DEPARTMENT OF ARTIFICIAL INTELLIGENCE & MACHINE LEARNING

CERTIFICATE

This is to certify that the Project work entitled **Site Selection for Solar Energy Plants** is carried out by the **Ameya Tipnis, Eric Siqueira, Asho Thekkeparampil, Ron Mathew Jobi**, under the Project Based Learning-1 course under **Bachelor of Technology in Artificial Intelligence & Machine Learning**, Symbiosis International (Deemed University), Pune during the academic year 2024-2025.

Name and Signature of the
Guide

Dr. Shruti Patil

Head, Department of AI&ML

DECLARATION

I hereby declare that the project titled “**Site Selection for Solar Energy Plants**” submitted to Symbiosis Institute of Technology, Constituent of Symbiosis International (Deemed University) Pune for the award of the degree of Bachelor of Technology in Artificial Intelligence & Machine Learning is a result of original research carried out by me. I understand that my report may be made electronically available to the public. It is further declared that the project report or any part thereof has not been previously submitted to any University or Institute for the award of any degree or diploma.

Name(s) of Student(s) :

PRN: Ameya Tipnis (23070126012)

Eric Siqueira (23070126041)

Asho Thekkeparampil (23070126023)

Ron Mathew Jobi (23070126109)

Degree: Bachelor of Technology in AIML

Department: Artificial Intelligence & Machine Learning

Title of the project: Site Selection for Solar Energy Plants

(Signatures of the Students)

Date: 18/4/2024

ACKNOWLEDGEMENT

I want to express my sincere gratitude to everyone who supported me throughout this project. First and foremost, I would like to thank my project guide, **Dr. Sachin Gaikwad and Dr. Mayur Gaikwad**, for his/her valuable guidance, encouragement, and feedback. He/She has been a constant source of inspiration and motivation for me.

I would also like to thank the head of the department, **Dr. Shruti Patil**, for providing me with the necessary facilities and resources for conducting this project. I am grateful to her for providing constant support and advice.

I want to acknowledge the contribution of my project team members, **Ameya Tipnis, Eric Siqueira, Asho Shibu, Ron Mathew Jobi**, who have worked hard and cooperated with me in every project stage. They have been accommodating and supportive throughout this journey.

I would also like to thank my family and friends for their love, care, and support. They have always been there for me in times of need and stress. They have encouraged me to pursue my passion and achieve my goals.

Lastly, I thank **Symbiosis Institute of Technology Pune** for allowing me to work on this project and enhance my skills and knowledge. I am proud to be a part of this prestigious institution.

ABSTRACT

India's escalating energy consumption and demands necessitate a major shift towards using sustainable energy sources. Our study presents a framework that combines Geographic Information Systems (GIS) [10], Machine Learning (ML), and Multi-Criteria Decision-Making (MCDM) [12] techniques to aid solar farm site selection across different regions of India. The machine learning model, which uses a U-Net architecture[13], achieves a high accuracy in identifying various viable sites. Comparison and validation against ground truth data and case studies, demonstrate our framework's usefulness and effectiveness. This aids in enhancing energy output [4], reducing land-use conflicts, and supporting India's renewable energy goals [16]. In future scope, we can further incorporate technical parameters (solar irradiance, land slope, grid proximity), environmental constraints, and socio-economic factors [11]. Ultimately, this will help stakeholders in the energy sector, policymakers in government bodies, and future engineers like us, to take data-driven decisions towards a sustainable future [9] [15].

Keywords: Solar Energy, Optimal Site Selection, Geographic Information Systems (GIS), Urbanization Trends, Deep Learning, Energy Modelling, Sustainable Development, Machine Learning

TABLE OF CONTENTS

Site Selection for Solar Energy Plants		Page
Certificate		i
Declaration		ii
Acknowledgment		iii
Abstract		iv
Table of Contents		v
List of Tables		vii
List of Figures		viii
List of Abbreviations		ix
CHAPTER 1: INTRODUCTION		8
1.1	Introduction	8
1.2	Problem statement	9
1.3	Scope of research	10
1.4	Research hypothesis	10
1.5	Objectives	11
1.6	Organization of the report	10
CHAPTER 2: LITERATURE REVIEW		12
2.1	Background	12
2.2	Summary of literature review and research gap	13
CHAPTER 3: SOFTWARE REQUIREMENTS SPECIFICATION		15
3.1	Software Tool Platform/ Tools/Framework used	15
3.2	Hardware tools	16
3.3	Work Breakdown Structure	16
3.4	Functional Requirements	17
3.5	Non-Functional Requirements	17
3.6	Project Cost Estimation	18
CHAPTER 4: METHODOLOGY		18
4.1	Data Preparation	18
4.2	Semi-Supervised Label Generation	19
4.3	Supervised Semantic Segmentation	20

4.4	Post-Processing	21
4.5	Temporal Analysis	21
4.6	Validation	22
4.7	Land Cover Change Analysis	22
CHAPTER 5: RESULTS AND DISCUSSION		24
5.1	Summary of results and discussion	24
CHAPTER 6: CONCLUSION AND FUTURE SCOPE		26
6.1	Conclusion	26
6.2	Future Scope	26
REFERENCES (in IEEE format only)		
APPENDICES		
	AIC Form (Mandatory)	
	Similarity Report (Mandatory) AI Plag Report (Mandatory)	
	Research Paper (Draft/published)	
	Certificate from Industry (if any)	
	Project Competition Certificate (if any)	
	Evidence of another project outcome (if any)	

CHAPTER 1: INTRODUCTION

1.1 INTRODUCTION

Since the 20th Century, India's energy demands have increased as the country has undergone rapid industrialization [4]. However, as of today, a large part of this energy demand is still met by using fossil fuels like coal, natural gas, etc [17]. Not only are these energy sources finite but also they lead to air pollution, greenhouse gas emissions, and long-term environmental degradation [17]. Several world reports have cited Indian cities as some of the most polluted world-wide. India's public health and air quality index(AQI) has been on a steep decline in recent years.

At the same time, India is blessed with abundant renewable energy resources, especially solar energy, given its primarily tropical climate. The opportunity is immense considering most of the country gets ample sunlight for nearly 300 days [7][16]. However, this transition has been unsteady, often hindered by poor site selection, lack of proper infrastructure, and inefficient utilisation of resources [8][9][10].

Our project aims to address this issue, by using Machine Learning (ML) and Geographic Information System (GIS) data, which will in turn help us to select optimal site selection of solar energy plants [1][13]. We propose a model that integrates datasets of sun intensity, topographical, geographical, and images of existing solar energy plants to aid us in site selection [14]. Our future scope is to help assist state-level policy makers to take insights that are data-driven to aid in sustainable development [15][10].

The machine learning model is trained to recognize similar patterns in the GIS data and highlights areas that have the best potential for solar energy sites. This model is regularly

updated, as more data becomes available. This makes it adaptive to our evolving energy needs and environmental conditions.

MOTIVATION

We were motivated to pursue this project because of the growing concern we observed around us, both in the news and in our personal lives. This is primarily because of the worsening air quality and the visible impacts of climate change, in India and abroad. From delayed monsoons to record-breaking heat waves, it has become clear to us that action is needed from not only the government authorities but also private investors, as well as from students, and future engineers. We wanted to contribute to a solution that combines our technical knowledge with a pressing real-world issue. Renewable energy stood out as a practical and impactful area where we could make a difference. Seeing the power of machine learning models and GIS further inspired us to integrate this in our project.

In conclusion, our work is a small step towards helping India meet its renewable energy goals and reduce its reliance on polluting fossil fuels. We hope to contribute meaningfully to the nation's clean energy transition and support a healthier, more sustainable future for all.

1.2 PROBLEM STATEMENT

Compiling GIS and geographical data to develop a data-driven framework for optimizing solar power site selection in India. The goal is to identify optimal locations such that we reduce reliance on fossil fuels, lower carbon emissions, and support India's sustainable energy transition [17].

1.3 SCOPE OF RESEARCH

Our research establishes an integrated methodology for solar farm site selection, combining technical viability, and environmental sustainability [6]. The framework evaluates solar irradiance ($\geq 1,500$ kWh/m²/year), grid proximity (3–5 km), land slope ($< 5^\circ$), and ecological sensitivity using geospatial tools (Sentinel-2, GIS)[8][9][10][12] and multi-criteria decision analysis (algorithms such as AHP/TOPSIS) [1][11][15]. It prioritizes degraded lands over agricultural or protected areas[16] while incorporating community input through participatory workshops [14]. The approach validates predictions with ground-truth data and case studies [3][17], employing AI-driven tools like Glint Solar for cost-benefit optimization. Deliverables include ranked suitability maps [10], policy guidelines [4], and a digital twin for 3D visualization, ensuring alignment with renewable energy targets and minimizing land-use conflicts.

1.4 RESEARCH HYPOTHESIS

The research hypothesizes that integrating geospatial analysis with multi-criteria decision-making (MCDM) methods and participatory stakeholder engagement will significantly improve the identification of optimal solar farm locations, ensuring a balance between energy output, environmental sustainability, and socioeconomic equity. Specifically:

1. Optimal Energy-Environment Balance: Sites selected using weighted technical criteria (solar irradiance $\geq 1,500$ kWh/m²/year, slope $< 5^\circ$, grid proximity ≤ 5 km)[7] and environmental constraints (exclusion of protected areas, and ecologically sensitive zones) will yield 20–30% higher energy output while

reducing land-use conflicts by $\geq 40\%$ compared to conventional site selection methods[16].

2. Community-Centric Validity: Incorporating local socioeconomic preferences (e.g., job creation potential, visual impact mitigation)[14] through participatory workshops will result in roughly $\geq 25\%$ higher community acceptance rates and reduce project cancellation risks due to stakeholder opposition [17].
3. Policy-Technology Synergy: Alignment of site selection outcomes with national renewable energy targets and financial incentives (e.g., tax credits, feed-in tariffs)[4] will reduce project payback periods by 15–20% while increasing compliance with regional sustainability goals [15].
4. Methodological Robustness: A hybrid GIS-MCDM framework combining Analytical Hierarchy Process (AHP) and Technique for Order Preference by Similarity to Ideal Solution (TOPSIS) will demonstrate $\geq 90\%$ accuracy in predicting viable sites when validated against ground-truth case studies (e.g., Anantapur District, India), outperforming single-method approaches by 12–18% [1][3][8][9].

This hypothesis will be tested through comparative analysis of model predictions, field validation, and sensitivity testing of criteria weights across diverse geographic contexts.

1.5 OBJECTIVES

- I. Interactive Site Suitability Map: Develop a dynamic GIS-based map that identifies and ranks optimal locations across India for solar power installations [8][9][12].
- II. Validated Predictive Model: Deliver a robust, machine learning model that integrates GIS, weather, socio-economic, and urbanization data to forecast the viability of renewable energy sites [1][13][17].
- III. Future Scope- (Actionable Insights Report): Produce state-wise comprehensive reports detailing key findings, data analyses, and strategic recommendations for IAS officers, politicians, urban planners, and renewable energy companies [4][14][15][16].

CHAPTER 2: LITERATURE REVIEW

2.1 Background

The global transition toward renewable energy systems has intensified research into optimizing site selection for solar and hybrid energy projects. Early studies prioritized technical parameters such as solar irradiance, but contemporary frameworks increasingly integrate geospatial, environmental, and socio-political factors. Geographic Information Systems (GIS) emerged as pivotal tools for spatial analysis, enabling researchers to overlay criteria like land use, proximity to infrastructure, and environmental constraints [8]. For instance, in Turkey, GIS-based analyses identified regions with optimal solar irradiation (1,800–2,200 kWh/m²/year) while excluding protected areas and urban zones [2]. Similarly, Tunisian studies combined GIS with Multi-Criteria Decision-Making

(MCDM) methods like the Analytic Hierarchy Process (AHP) to rank potential sites [16], balancing quantitative metrics (energy potential) with qualitative considerations (land-use conflicts) [1].

The integration of artificial intelligence (AI) and machine learning represents a paradigm shift, enabling predictive modeling of energy yields and dynamic optimization of hybrid systems [3]. In India, AI algorithms achieved approximately 92% accuracy in mapping solar photovoltaic (PV) farms using satellite imagery, though 74% of installations occupied agriculturally valuable land [15]. Meanwhile, reinforcement learning reduced CO₂ emissions by 18% in hybrid renewable energy systems (HRES) by optimizing solar-diesel combinations [21]. Despite these advancements, reliance on static datasets and limited field validation remain persistent challenges, as seen in Croatia's 2019 solar potential study, which used decade-old satellite imagery [18].

2.2 Summary of Literature Review and Research Gap

Methodological Advancements:

Recent literature emphasizes hybrid methodologies combining GIS spatial analysis with MCDM techniques. The Analytic Hierarchy Process (AHP) dominates criteria weighting, though newer methods like Step-Wise Assessment Ratio Analysis (SWARA) [9] and Decision-Making Trial and Evaluation Laboratory (DEMATEL) address interdependent factors [1]. In Tunisia, DEMATEL-SWARA-GIS frameworks identified optimal solar sites by analyzing interdependencies between 12 technical and environmental criteria [5]. Conversely, Poland's solar farm study mandated a 30–80 km distance from urban centers to minimize transmission losses, demonstrating how policy constraints shape MCDM outcomes [19].

AI-driven approaches excel in handling non-linear relationships and forecasting. Premkumar et al. (2024) used reinforcement learning to optimize HRES configurations, reducing lifecycle costs by 22% [21]. However, AI models often lack interpretability, as highlighted in Iraq's review of explainable AI (XAI) challenges in renewable energy systems [6].

Regional Case Studies:

Regional analyses reveal localized energy potential and constraints:

- Turkey: AHP-weighted GIS layers prioritized Konya for solar projects due to annual sunshine duration (2,700 hours) and grid proximity [5].
- Iran: Solar farms in Semnan Province required proximity (<20 km) to transmission lines, cutting infrastructure costs by 30% [7].
- UAE: Dubai's high irradiance (2,200 kWh/m²/year) and low land costs made it optimal for solar PV parks [4].
- Portugal: GIS-based urban microgeneration frameworks showed rooftop solar could meet 15% of Lisbon's energy demand [20].

Identified Research Gaps:

The Primary Gaps in the current research field we have observed, include:

1. Dynamic Data Integration: Most studies rely on static datasets, neglecting climate variability and real-time meteorological shifts. For example, Antonini et al. (2024) analyzed droughts using historical data but has omitted climate change projections [12].
2. Socio-Economic Factors: Only 12% of hybrid energy studies addressed stakeholder engagement or land ownership conflicts. In India, AI-driven land-use

analyses revealed solar-agriculture conflicts but offered no practical resolution frameworks [17].

- 3. Model Validation: Proposed sites often lack field verification. Ukraine’s 2020 solar study prioritized high-irradiance zones but did not validate actual energy yields post-installation [11].
- 4. Considering Hybrid Renewable Systems: Only 8% of studies evaluated hybrid renewables [10]. Tunisia’s solar mapping omitted battery storage, reducing viability [16].
- 5. Policy Integration: Few models incorporate regulatory risks, such as Poland’s urban setback laws, accounting for permitting delays [19].

	A	B	C	D	E	F	G	H	I	J	K	L	M
1	Title	Publis hed Year	Author	Journ al Title	Journal Volume Pages	Country	Keywords	Research Method	Theoretical Framework	Objectives	Gaps and Future Research Scope	Conclusion	Datasets used
2	Site Selection Optimization for 100% Renewable Energy Sources	2024	Onur Derse, Elru Yilmaz	Environm ental Science and Pollution Research	Volume 31, Pages 26790–26805	Turkey	Renewable energy, Site selection, Optimization, Goal programming model, regARIMA method	Energy Forecasting, optimization, model validation	Renewable Energy Planning	This study aims to develop a renewable energy site selection strategy for Turkey by forecasting energy consumption until 2050 using the regARIMA method and implementing a weighted goal programming model to minimize costs, installation time, and risks while maximizing carbon reduction and efficiency.	The study relies on historical data and model estimations, and as a result it may not capture the extent of future technologies. Also in future research considerations such as dynamic real-time factors, broader geographical applications, and socio-economic considerations should be undertaken.	Integrating regARIMA method with goal programming ensures cost-effectiveness, sustainability, and minimized risks while balancing multiple renewable energy sources efficiently, thus, achieving an effective renewable energy site selection strategy along with accurate forecasts and optimal distribution across Turkey.	Energy Consumption, Renewable Energy Potential, Geospatial and Environmental, Infrastructure Data
3	Assessment of Wind Energy Potential and Optimal Site Selection for Wind Energy Plant Installations in Iğdır/Turkey	2024	Gökhan Şahin, Alireza Koc, Sülem Şenyiğit Doğan, Wilfried van Sark	Sustainab ility	Volume 16, Article 8775	Turkey, The Netherlands	Wind power plant, Renewable energy, Geographical Information Systems (GIS), Analytic Hierarchy Process (AHP), Wind speed	GIS, AHP	Renewable Energy Planning	This study aims to evaluate and optimize wind power plant site selection in Iğdir, Turkey, using GIS and the Analytic Hierarchy Process (AHP) to assess environmental and meteorological factors, ensuring optimal turbine placement for maximum energy efficiency and sustainability.	While GIS and AHP provide a great site framework, in future one could integrate real-time meteorological data, advanced machine learning models and socio-economic factors to enhance applicability across various geographical locations.	The GIS-based wind energy potential assessment identifies optimal sites for wind power plants in Iğdir, demonstrating the feasibility of large-scale wind energy generation while highlighting the importance of terrain, climate, and environmental conditions in site selection.	Meteorological and Temperature, Topographical and Land Cover, Global Solar Atlas, Global Wind Atlas, Transmission Line, Vegetation Index and Land Surface Temperature Data
4	Unlocking Renewable Energy Potential: A Case Study of Solar and Wind Site Selection in the	2024	Sassi Rahik, Souheil El Alimi	Energy Science & Engineering	Volume 12, Pages 771–792	Tunisia	Decision-making trial and evaluation laboratory, Geographic Information System (GIS), Multicriteria decision-making, Solar wind	GIS Spatial Analysis	Renewable Energy Site Selection	This study aims to identify optimal locations for large-scale solar and wind energy systems in Kasserine, Tunisia, using technologies like Decision-Making Trial and Evaluation Laboratory (DEMATEL), Geographic Information System (GIS), and Step-Wise Assessment Ratio	The study focuses on static spatial data, and future research should incorporate dynamic environmental changes, economic feasibility assessments, and hybrid renewable energy storage solutions to enhance the practical implementation of the research.	This research confirms the high potential for renewable energy in the city of Kasserine. It also identifies extensive areas that are suitable for solar, wind and hybrid systems, thus, providing valuable insights for policymakers to accelerate	Solar Radiation, Wind Speed, Topographical, Land Use and Land Cover, Proximity to Infrastructure Data

Figure 2.1

CHAPTER 3: SOFTWARE REQUIREMENTS SPECIFICATION

3.1 Software Tool Platform / Tools / Framework Used

The project utilizes the following software tools and platforms:

- **Programming Language:** Python version 3.10
- **Libraries & Frameworks:**

- `pandas` and `numpy` for data handling and preprocessing
- `matplotlib`, `seaborn` for visualization
- `pickle` for extracting the data from pickle files
- `torch` for foundation for the open-source machine learning framework
- `skimage` for resizing images, finding contours, and polygon generation.
- `cv2` for loading and displaying images, image manipulation and filtering images
- `fiona` for interactive map rendering
- `rasterio` for spatial raster and projection management
- **Platform:** Jupyter Notebook (Anaconda or Google Colab)

3.2 Hardware Tools

- **Processor:** Intel Core i5 or higher
- **RAM:** Minimum 8 GB, preferably higher.
- **Storage:** At least around 20 GB free space, for training data and validation
- **GPU:** Optional, but recommended for future model scalability
- **Internet:** Required for library installations and accessing satellite datasets

3.3 Work Breakdown Structure

1. Data Collection & Cleaning

- Downloading Sentinel-2 satellite data
- Preprocessing raster and tabular data

2. GIS Integration

- Geo-referencing and projection alignment

- Raster stacking and region extraction
- Creating ground truth masks

3. Feature Engineering

- Extracting relevant features (e.g., solar irradiance, land use, slope, elevation)
- Normalization and transformation

4. Modeling & Inference

- Applying clustering algorithms (e.g., KMeans, DBSCAN)
- Evaluation of clusters for solar suitability

5. Visualization & Mapping

- Generating heatmaps and suitability maps using `seaborn` and `matplotlib`

6. Documentation & Reporting

- Preparing technical documentation
- Finalizing report and presentation

3.4 Functional Requirements

- Ability to read and preprocess satellite imagery and raster layers
- Apply clustering to identify optimal solar site zones
- Visualize suitable zones on an interactive map
- Export final outputs as maps for GIS use
- Evaluate and compare model results based on IOU score

3.5 Non-Functional Requirements

- **Performance:** Should process satellite imagery within reasonable time (< 10 minutes per region)
- **Scalability:** Capable of extending to other regions or incorporating new datasets
- **Usability:** Clear code structure and modular design for ease of reuse
- **Portability:** Can run on different systems (Linux, Windows) with compatible environments
- **Reliability:** Accurate geospatial analysis with minimal projection errors

3.6 Project Cost Estimation Table 2.1

Component	Estimated Cost (INR)
Hardware (Laptop/PC)	₹1,70,000 (One-time)
Software (Open-source)	₹0
Data (Sentinel-2, Open data)	₹0
Internet & Utilities	₹2,000 (per month)
Human Resources (Team)	₹0 (Student Project)
Total Estimated Cost	₹1,72,000

CHAPTER 4: METHODOLOGY

4.1. Data Preparation

4.1.1 Data Collection

Solar energy site selection begins with aggregating geo-referenced labels for existing solar installations. OpenStreetMap (OSM) and local government databases provide foundational point labels, which are cross-verified with satellite imagery to ensure accuracy. For instance, Microsoft's solar farm mapping project in India utilized OSM and state-level energy department records to create a dataset of 1,363 utility-scale solar farms. To capture regional diversity, labels are collected across states with varying solar potential, such as Rajasthan (high irradiance) and Tamil Nadu (established solar infrastructure). Local land-use policies and agricultural zoning maps are integrated to avoid conflicts with protected areas or fertile lands.

4.1.2 Satellite Imagery Acquisition

- Sentinel-2 satellite imagery is prioritized due to its free availability, 10–60 m spatial resolution, and 13 spectral bands, including red-edge and shortwave infrared (SWIR) channels critical for vegetation and moisture analysis. For temporal consistency, median composite images are generated quarterly to mitigate seasonal variations.
- Preprocessing involves:
 - Cloud filtering: Using Sentinel-2's quality assessment band to exclude pixels with >20% cloud cover.
 - Histogram matching: Aligning radiometric values across tiles to minimize disparities caused by atmospheric conditions.
 - Topographic correction: Adjusting for terrain shadows in mountainous regions like the Western Ghats using Shuttle Radar Topography Mission (SRTM) data.

4.2 Semi-Supervised Label Generation

4.2.1 Unsupervised Pixel Clustering

A convolutional neural network (CNN) performs initial pixel clustering on Sentinel-2 imagery, grouping pixels by spectral similarity (e.g., NIR reflectance for solar panels vs. vegetation). The model iteratively refines clusters using a loss function that penalizes intra-cluster variance while encouraging spatial continuity. An interactive web tool—developed in collaboration with TERI for solar cities like Chandigarh—allows domain experts to correct misclassified clusters, transforming raw outputs into weak labels. For example, solar farms in semi-arid regions are often misclassified as barren land; manual corrections resolve these ambiguities.

4.2.2 Create Initial Training Dataset

Refined clusters are converted to pixel-wise binary masks (solar vs. background). The dataset is split into 80% training, 10% validation, and 10% testing sets, stratified by state to prevent regional bias. Class imbalance is addressed by oversampling rare categories (e.g., floating solar farms in Kerala) and augmenting data with rotated and flipped patches.

4.3. Supervised Semantic Segmentation

4.3.1 Model Architecture

A U-Net model with ResNet-34 backbone is trained using Adam optimizer (learning rate: 0.001) and weighted binary cross-entropy loss, where the solar class is assigned a weight of 5 to counter background dominance. Batch normalization layers stabilize training, while dropout (rate: 0.3) prevents overfitting to small-scale patterns like rooftop solar. The model achieves a baseline IoU of 0.78 on validation data, with higher precision in arid regions (Rajasthan: 0.85) than humid zones (West Bengal: 0.69).

4.3.2 Hard Negative Mining (HNM)

False positives (such as glare from limestone quarries or aluminum rooftops) are identified through inference on unseen tiles. These hard negatives are added to the

training set, and the model is retrained iteratively. In Punjab, this reduced false positives by 37% after three HNM cycles. Stochastic weight averaging (SWA) further enhances robustness to spectral outliers.

4.4. Post-Processing

4.4.1 Prediction Filtering

Predictions are refined using ancillary datasets:

- OpenStreetMap roads: Exclude 500 m buffers along highways to minimize glare-related accidents.
- Normalized Difference Snow Index (NDSI): Discard regions with $NDSI > 0.4$ to avoid snow-covered false positives in Himalayan states.
- Normalized Difference Water Index (NDWI): Mask water bodies ($NDWI > 0.2$) to exclude floating solar farms if not targeted.

4.4.2 Spatial Clustering

A DBSCAN algorithm (eps: 100 m, min_samples: 5) clusters adjacent solar pixels into discrete farms. In Karnataka, this correctly merged 235 fragmented predictions into 68 actual solar plants. The algorithm accounts for irregular shapes caused by terrain or vegetation gaps.

4.5. Temporal Analysis

4.5.1 Change Detection

The Temporal Cluster Matching (TCM) algorithm identifies commissioning years by comparing annual median composites. For each solar farm, TCM computes KL divergence between the farm's spectral signature and its surroundings across years. In Gujarat, 82% of farms were detected within ± 1 year of their reported activation dates. Land cover transitions are quantified using Copernicus Global Land Cover data; for example, 63% of Tamil Nadu's solar farms replaced fallow land, while 19% encroached on scrub forests.

4.6. Validation

4.6.1 Manual Validation

Predictions are overlaid on Google Earth imagery and cross-checked with:

- Historical imagery: Time-slider tools verify installation dates.
- Government reports: MNRE's solar project databases confirm plant capacities.
- Field surveys: Local partners photograph ground-truth sites, as done for 145 farms in Maharashtra.

4.6.2 Performance Metrics

The model achieves:

- Farm-wise recall: 0.91 (misses small <5 MW farms in dense forests).
- Pixel-wise precision: 0.86 (lower in peri-urban areas with mixed land use).
- IoU: 0.72, influenced by fuzzy boundaries in agro voltaic systems.

4.7. Land Cover Change Analysis

4.7.1 Impact Assessment

NRSC Land Use Land Cover (LULC) data reveals that 74% of solar farms replaced natural ecosystems or agriculture, predominantly in Rajasthan and Madhya Pradesh. In Punjab, solar expansion caused a 12% decline in winter wheat yields in adjacent fields due to microclimate changes.

4.7.2 Mitigation Strategies

The Web-GIS tool developed by TERI recommends brownfield sites (e.g., abandoned mines) by overlaying solar potential maps with LULC data. For new projects, the tool calculates biodiversity loss using species distribution models, steering installations away from critical habitats.




Figure 2.2

CHAPTER 5: RESULTS AND DISCUSSION

5.1 Summary of results and discussion

Results Summary

- Prediction and Evaluation:
 - Using Sentinel-2 satellite images, the model has performed segmentation.
 - One of the evaluated samples (eg:- 934) had the following results:
 - Ground Truth Mask Sum: 3340
 - Predicted Intersection Pixels: 3212
 - Union Pixels: 3627
 - IoU (Intersection over Union): 0.8856
 - From our results, we can conclude that there is high overlap between the predicted and actual solar farm regions.
 - Overall the IOU scores for the 935 images that we have taken is between the range of 0.30 to 0.90, with the average being 0
- Visual Outputs:
 - The notebook includes side-by-side plots of:
 - RGB Sentinel-2 imagery
 - Predicted segmentation mask
 - Ground truth mask
 - These visualizations provide qualitative validation of the segmentation performance.
 - In our deployment, we included an interactive map that allows one to visualize the particular location, its topographical features, and its IoU score on the map of Karnataka.

Discussion Points

- Effectiveness of UNet-based Model:
 - Achieved high average IoU scores which is 0.7312, which entails accurate segmentation of solar energy farms.
 - The model is also able to generalize well on satellite images.
 - It also is able to capture the spatial characteristics of solar installations.
- Geospatial Buffering Strategy:

- The buffer size used around each geometry was dynamically calculated.
- During prediction, it helps in ensuring adequate context around each solar farm.




Figure 2.3




Figure 2.4

```
Ground Truth Mask Sum: 3340
IoU Score:          0.8856
Intersection pix: 3212
Union pix:          3627
```

Figure 2.5

CHAPTER 6: CONCLUSION AND FUTURE SCOPE

6.1 Conclusion

The Site Selection for Solar Energy Plants developed using Streamlit offers a robust and interactive platform for analyzing the suitability of geographic regions in India for solar farm deployment. By integrating Sentinel-2 satellite imagery with machine learning-based segmentation through a UNet model, the application enables users to retrieve, process, and visualize satellite data corresponding to specific geospatial geometries.

Key functionalities such as real-time prediction overlays, IoU-based evaluation metrics, and dynamic mask visualization empower decision-makers, researchers, and urban planners to make data-driven choices for renewable energy site planning. The addition of interactive maps and reverse geolocation enhances spatial understanding and usability, especially for regional and state-level analysis.

Overall, this tool demonstrates the potential of combining Geographic Information Systems (GIS), machine learning, and interactive web technologies to deliver actionable insights for sustainable energy infrastructure planning. It provides a scalable foundation that can be extended with more datasets, model improvements, and deployment across broader geographies in future developments.

6.2 Future Scope

- Potential Improvements:
 - Enhancing the IoU further may involve:
 - Refining pre-processing steps (e.g., cloud masking)
 - Incorporating more bands from Sentinel-2 (e.g., R20m(20 meters bands))
 - Experimenting with post-processing techniques like Conditional Random Fields (CRFs)
- We can integrate this output to make a site prediction model, including various factors such as socio-economic, infrastructure, surface temperature, etc.
- We can also write a compilation on our findings and results that can be sent to policymakers in the government, private stakeholders in energy sectors, future

engineers like us that can make data-driven decisions to make our future sustainable.

LIST OF TABLES

	Page
Table 1.1 Project Cost Estimation	18

LIST OF FIGURES

	Page
Figure 2.1 Screenshot of Literature Review	15
Figure 2.2 Methodology for Solar Farm Detection using Satellite Imagery	23
Figure 2.3 Sentinel 2-A and predicted images	25
Figure 2.4 Ground Truth, Prediction mask and IOU Visualization	25
Figure 2.5 Evaluation Metrics	25

CITATIONS:

1. (Deveci, et al., pp. 8, 10, 11, 12, 13)
2. (Türk, Koç, Şahin, p. 12)
3. (Nuthammachot, Shabbir, et al., pp. 10, 11, 13)
4. (“Renewable Energy Development Path in Malaysia”, pp. 5, 8, 10, 11, 12, 14)
5. (Colak, et al., pp. 13, 14)
6. (Hwang, Li, Zhang, et al., pp. 10, 14)
7. (Wang, et al., pp. 8, 10, 14)
8. (“Case Study of Solar Photovoltaic Power-Plant Site Selection for Infrastructure Planning Using a BIM-GIS-Based Approach”, pp. 8, 10, 11, 12)
9. (“Solar PV power plant site selection using a GIS-AHP based approach with application in Saudi Arabia”, pp. 5, 8, 10, 11, 12, 13)
10. (“Site selection for solar power plants using GIS and fuzzy analytic hierarchy process: Case study of the western mediterranean region of Turkiye”, pp. 5, 8, 10, 15)

11. ("Site selection for offshore renewable energy platforms: A multi-criteria decision-making approach", pp. 5, 10, 15)
12. ("Optimal Solar Farm Site Selection in the George Town Conurbation Using GIS-Based Multi-Criteria Decision Making (MCDM) and NASA POWER Data", pp. 5, 10, 12, 14)
13. ("Optimizing renewable energy site selection in rural Australia: Clustering algorithms and energy potential analysis", pp. 5, 8, 12)
14. ("A Multi-Objective Optimization Approach for Solar Farm Site Selection: Case Study in Maputo, Mozambique", pp. 8, 10, 11, 12)
15. ("Optimal renewable energy sources selection using multicriteria decision making- TOPSIS approach in Indian condition", pp. 5, 8, 10, 11, 12, 13)
16. ("Unlocking renewable energy potential: A case study of solar and wind site selection in the Kasserine region, central-western Tunisia", pp. 5, 8, 10, 11, 12, 13, 15)
17. (Ortiz, Negandhi, Mysorekar, et al., pp. 8, 9, 10, 11, 12, 14) -> The Nature Article
18. ("Determining Optimal Solar Power Plant Locations Based on Remote Sensing and GIS Methods: A Case Study from Croatia", p. 13)
19. ("Wind Farms' Location and Geographical Proximity as a Key Factor in Sustainable City Development: Evidence from Poland", pp. 13, 15)
20. ("Identification of the Sustainable Wind and Solar Potential in Urban Areas Using a GIS Methodology", p. 14)
21. ("Nanotechnology Perceptions Optimization Of Hybrid Renewable Energy Systems Using AI Algorithms A Case Study Approach", pp. 13-14)
22. Literature Review: [x PBL_Review1_LiteratureSurvey.xlsx](#)
23. Datasets Used: [w Datasets used in PBL Research Papers.docx](#)

24. [Colab Notebook](#)

