# Machine Learning

Justin Grimmer

Associate Professor Department of Political Science University of Chicago

February 26th, 2018

- Suppose we have N units, i = 1, ..., N.

- Suppose we have N units, i = 1, ..., N.
- Each unit receives treatment  $T_i$  (potentially vector valued)

- Suppose we have N units, i = 1, ..., N.
- Each unit receives treatment  $T_i$  (potentially vector valued)
- Suppose we have (potentially vector valued) response  $\mathbf{Y}_i(\mathbf{T}_i)$

- Suppose we have N units, i = 1, ..., N.
- Each unit receives treatment  $T_i$  (potentially vector valued)
- Suppose we have (potentially vector valued) response  $\mathbf{Y}_i(\mathbf{T}_i)$ 
  - Unit i's response to treatment  $T_i$

- Suppose we have N units, i = 1, ..., N.
- Each unit receives treatment  $T_i$  (potentially vector valued)
- Suppose we have (potentially vector valued) response  $\boldsymbol{Y}_i(\boldsymbol{T}_i)$ 
  - Unit i's response to treatment  $T_i$
  - Function: maps from treatments to responses

- Suppose we have N units, i = 1, ..., N.
- Each unit receives treatment  $T_i$  (potentially vector valued)
- Suppose we have (potentially vector valued) response  $\mathbf{Y}_i(\mathbf{T}_i)$ 
  - Unit i's response to treatment  $T_i$
  - Function: maps from treatments to responses
- Fundamental problem of causal inference observe only one  $m{Y}_i(m{T}_i)$

- Suppose we have N units, i = 1, ..., N.
- Each unit receives treatment  $T_i$  (potentially vector valued)
- Suppose we have (potentially vector valued) response  $\mathbf{Y}_i(\mathbf{T}_i)$ 
  - Unit i's response to treatment  $T_i$
  - Function: maps from treatments to responses
- Fundamental problem of causal inference observe only one  $\boldsymbol{Y}_i(\boldsymbol{T}_i)$
- Quantity of interest:

- Suppose we have N units, i = 1, ..., N.
- Each unit receives treatment  $T_i$  (potentially vector valued)
- Suppose we have (potentially vector valued) response  $\boldsymbol{Y}_i(\boldsymbol{T}_i)$ 
  - Unit i's response to treatment  $T_i$
  - Function: maps from treatments to responses
- Fundamental problem of causal inference observe only one  $\boldsymbol{Y}_i(\boldsymbol{T}_i)$
- Quantity of interest:
  - Simplest (yet very powerful) case $\leadsto T_i \in \{0,1\}, \; Y_i(T_i) \in \{0,1\}$

- Suppose we have N units, i = 1, ..., N.
- Each unit receives treatment  $T_i$  (potentially vector valued)
- Suppose we have (potentially vector valued) response  $\boldsymbol{Y}_i(\boldsymbol{T}_i)$ 
  - Unit i's response to treatment  $T_i$
  - Function: maps from treatments to responses
- Fundamental problem of causal inference observe only one  $m{Y}_i(m{T}_i)$
- Quantity of interest:
  - Simplest (yet very powerful) case $\leadsto T_i \in \{0,1\}, \ Y_i(T_i) \in \{0,1\}$

$$ATE = E[Y(1) - Y(0)]$$

- Suppose we have N units, i = 1, ..., N.
- Each unit receives treatment  $T_i$  (potentially vector valued)
- Suppose we have (potentially vector valued) response  $\boldsymbol{Y}_i(\boldsymbol{T}_i)$ 
  - Unit i's response to treatment  $T_i$
  - Function: maps from treatments to responses
- Fundamental problem of causal inference observe only one  $m{Y}_i(m{T}_i)$
- Quantity of interest:
  - Simplest (yet very powerful) case $\leadsto T_i \in \{0,1\}, \; Y_i(T_i) \in \{0,1\}$

$$ATE = E[Y(1) - Y(0)]$$

- Estimate with:

- Suppose we have N units, i = 1, ..., N.
- Each unit receives treatment  $T_i$  (potentially vector valued)
- Suppose we have (potentially vector valued) response  $\boldsymbol{Y}_i(\boldsymbol{T}_i)$ 
  - Unit i's response to treatment  $T_i$
  - Function: maps from treatments to responses
- Fundamental problem of causal inference observe only one  $\boldsymbol{Y}_i(\boldsymbol{T}_i)$
- Quantity of interest:
  - Simplest (yet very powerful) case  $\leadsto$   $T_i \in \{0,1\}, \ Y_i(T_i) \in \{0,1\}$

$$ATE = E[Y(1) - Y(0)]$$

- Estimate with:

$$\widehat{ATE} = E[Y(1)|T=1] - E[Y(0)|T=0]$$

- Suppose we have N units, i = 1, ..., N.
- Each unit receives treatment  $T_i$  (potentially vector valued)
- Suppose we have (potentially vector valued) response  $\boldsymbol{Y}_i(\boldsymbol{T}_i)$ 
  - Unit i's response to treatment  $T_i$
  - Function: maps from treatments to responses
- Fundamental problem of causal inference observe only one  $m{Y}_i(m{T}_i)$
- Quantity of interest:
  - Simplest (yet very powerful) case $\leadsto T_i \in \{0,1\}, \ Y_i(T_i) \in \{0,1\}$

$$ATE = E[Y(1) - Y(0)]$$

- Estimate with:

$$\widehat{ATE} = \frac{\sum_{i=1}^{N} I(T_i = 1) Y_i}{\sum_{i=1}^{N} I(T_i = 1)} - \frac{\sum_{i=1}^{N} I(T_i = 0) Y_i}{\sum_{i=1}^{N} I(T_i = 0)}$$

- Suppose we have N units, i = 1, ..., N.
- Each unit receives treatment  $T_i$  (potentially vector valued)
- Suppose we have (potentially vector valued) response  $\boldsymbol{Y}_i(\boldsymbol{T}_i)$ 
  - Unit i's response to treatment  $T_i$
  - Function: maps from treatments to responses
- Fundamental problem of causal inference observe only one  $Y_i(T_i)$
- Quantity of interest:
  - Simplest (yet very powerful) case $\leadsto T_i \in \{0,1\}, \ Y_i(T_i) \in \{0,1\}$

$$ATE = E[Y(1) - Y(0)]$$

- Estimate with:

$$\widehat{\mathsf{ATE}} = \frac{\sum_{i=1}^{N} I(T_i = 1) Y_i}{\sum_{i=1}^{N} I(T_i = 1)} - \frac{\sum_{i=1}^{N} I(T_i = 0) Y_i}{\sum_{i=1}^{N} I(T_i = 0)}$$

Question: how do we accurately estimate quantities, like ATE?

## Our Plan for the Day

- Experimental design
- Conditional average treatment effects
- Methods for estimating heterogeneous treatment effects

Rep. Harold "Hal" Rogers (KY-05) announced today that Kentucky is slated to receive \$962,500 to protect critical infrastructure- power plants, chemical facilities, stadiums, and other high-risk assets, through the U.S. Department of Homeland Security's buffer zone protection program

A federal grant will help keep the Brainerd Lakes Airport operating in winter weather. Today, Congressman Jim Oberstar announced that the Federal Aviation Administration (FAA) will award \$528,873 to the Brainerd airport. The funding will be used to purchase new snow removal and deicing equipment.

Congresswoman Darlene Hooley (OR-5) and Congressmen Earl Blumenauer (OR-3), David Wu (OR-1) and Greg Walden (OR-2) joined together today in announcing \$375,000 in federal funding for the Oregon Partnership to combat methamphetamine abuse in Oregon.

What information in credit claiming messages affect evaluations?

Experiment: vary the recipient of money and the action reported in credit claiming statement (and many other features)

Experiment: vary the recipient of money and the action reported in credit claiming statement (and many other features)

Treatments: type

- 1) Planned Parenthood
- 2) Parks
- 3) Gun Range
- 4) Fire Department
- 5) Police
- 6) Roads

Experiment: vary the recipient of money and the action reported in credit claiming statement (and many other features)

Treatments: type, stage

- 1) Will request
- 2) Requested
- 3) Secured

Experiment: vary the recipient of money and the action reported in credit claiming statement (and many other features)

Treatments: type, stage, money

- 1) \$50 Thousand
- 2) \$20 Million

Experiment: vary the recipient of money and the action reported in credit claiming statement (and many other features)

- Treatments: type, stage, money, collaboration
  - 1) Alone
  - 2) w/ Senate Democrat
  - 3) w/ Senate Republican

Experiment: vary the recipient of money and the action reported in credit claiming statement (and many other features)

Treatments: type, stage, money, collaboration, partisanship

- 1) Democrat
- 2) Republican

Experiment: vary the recipient of money and the action reported in credit claiming statement (and many other features)

Treatments: type, stage, money, collaboration, partisanship

Control Condition:

Advertising press release

#### Example Treatment:

**Headline**: Representative [blackbox] secured \$50 Thousand to purchase safety equipment for local firefighters

**Body**: Representative [blackbox] (Democrat) and Senator [blackbox], a Democrat, secured \$50 Thousand to purchase safety equipment for local firefighters.

Rep. [blackbox] said "This money will help our brave firefighters stay safe as they protect our businesses and homes"

#### Example Treatment:

**Headline**: Representative [blackbox] will request \$20 million for medical equipment at the local Planned Parenthood.

**Body**: Representative [blackbox] (Democrat), will request \$20 million for medical equipment at the local Planned Parenthood.

Rep. [blackbox] said "This money would help provide state of the art care for women in our community."

214 other conditions

214 other conditions

Dependent variable: Approve of representative

214 other conditions

Dependent variable: Approve of representative

Goal → measure effect of credit claiming content on approval ratings

214 other conditions

Dependent variable: Approve of representative

Goal → measure effect of credit claiming content on approval ratings Mechanics → Mechanical Turk sample (Findings are replicated in representative samples, using real representatives/senators)

- Participant i (i = 1, ..., N), has treatment assignment  $T_i$ 

- Participant i (i = 1, ..., N), has treatment assignment  $T_i$
- If  $T_i = 0$  for control condition

- Participant i (i = 1, ..., N), has treatment assignment  $T_i$
- If  $T_i = 0$  for control condition
- $\mathbf{T}_i = (\mathsf{T}_{i,\mathsf{type}},\mathsf{T}_{i,\mathsf{stage}},\mathsf{T}_{i,\mathsf{money}},\mathsf{T}_{i,\mathsf{collab}},\mathsf{T}_{i,\mathsf{part.}})$

- Participant i (i = 1, ..., N), has treatment assignment  $T_i$
- If  $T_i = 0$  for control condition
- $\mathbf{T}_i = (\mathsf{T}_{i,\mathsf{type}},\mathsf{T}_{i,\mathsf{stage}},\mathsf{T}_{i,\mathsf{money}},\mathsf{T}_{i,\mathsf{collab.}},\mathsf{T}_{i,\mathsf{part.}})$
- $Y_i(\mathbf{T}_i)$ : participant i's Approval decision under treatment  $\mathbf{T}_i$

- Participant i (i = 1, ..., N), has treatment assignment  $T_i$
- If  $T_i = 0$  for control condition
- $\mathbf{T}_i = (\mathsf{T}_{i,\mathsf{type}},\mathsf{T}_{i,\mathsf{stage}},\mathsf{T}_{i,\mathsf{money}},\mathsf{T}_{i,\mathsf{collab.}},\mathsf{T}_{i,\mathsf{part.}})$
- $Y_i(\mathbf{T}_i)$ : participant i's Approval decision under treatment  $\mathbf{T}_i$
- Quantities of Interest

- Participant i (i = 1, ..., N), has treatment assignment  $T_i$
- If  $T_i = 0$  for control condition
- $\mathbf{T}_i = (\mathsf{T}_{i,\mathsf{type}}, \mathsf{T}_{i,\mathsf{stage}}, \mathsf{T}_{i,\mathsf{money}}, \mathsf{T}_{i,\mathsf{collab}}, \mathsf{T}_{i,\mathsf{part.}})$
- $Y_i(\mathbf{T}_i)$ : participant i's Approval decision under treatment  $\mathbf{T}_i$
- Quantities of Interest
- Effect of particular component of message:

- Participant i (i = 1, ..., N), has treatment assignment  $T_i$
- If  $T_i = 0$  for control condition
- $\mathbf{T}_i = (\mathsf{T}_{i,\mathsf{type}},\mathsf{T}_{i,\mathsf{stage}},\mathsf{T}_{i,\mathsf{money}},\mathsf{T}_{i,\mathsf{collab}},\mathsf{T}_{i,\mathsf{part.}})$
- $Y_i(\mathbf{T}_i)$ : participant i's Approval decision under treatment  $\mathbf{T}_i$
- Quantities of Interest
- Effect of particular component of message:
  - $T_{stage} = Secured$

- Participant i (i = 1, ..., N), has treatment assignment  $T_i$
- If  $T_i = 0$  for control condition
- $\mathbf{T}_i = (\mathsf{T}_{i,\mathsf{type}},\mathsf{T}_{i,\mathsf{stage}},\mathsf{T}_{i,\mathsf{money}},\mathsf{T}_{i,\mathsf{collab.}},\mathsf{T}_{i,\mathsf{part.}})$
- $Y_i(\mathbf{T}_i)$ : participant i's Approval decision under treatment  $\mathbf{T}_i$
- Quantities of Interest
- Effect of particular component of message:
  - $T_{stage} = Secured$
  - $T_{stage} = Requested$

- Participant i (i = 1, ..., N), has treatment assignment  $T_i$
- If  $T_i = 0$  for control condition
- $\mathbf{T}_i = (\mathsf{T}_{i,\mathsf{type}},\mathsf{T}_{i,\mathsf{stage}},\mathsf{T}_{i,\mathsf{money}},\mathsf{T}_{i,\mathsf{collab}},\mathsf{T}_{i,\mathsf{part.}})$
- $Y_i(\mathbf{T}_i)$ : participant i's Approval decision under treatment  $\mathbf{T}_i$
- Quantities of Interest
- Effect of particular component of message:
  - $T_{\text{stage}} = Secured$
  - $T_{stage} = Requested$
  - $T_{stage} = Will Request$

- Participant i (i = 1, ..., N), has treatment assignment  $T_i$
- If  $T_i = 0$  for control condition
- $\mathbf{T}_i = (\mathsf{T}_{i,\mathsf{type}}, \mathsf{T}_{i,\mathsf{stage}}, \mathsf{T}_{i,\mathsf{money}}, \mathsf{T}_{i,\mathsf{collab.}}, \mathsf{T}_{i,\mathsf{part.}})$
- $Y_i(\mathbf{T}_i)$ : participant i's Approval decision under treatment  $\mathbf{T}_i$
- Quantities of Interest
- Effect of particular component of message:
  - $T_{stage} = Secured$
  - $T_{stage} = Requested$
  - $T_{stage} = Will Request$
  - $T_i = k$

- Participant i (i = 1, ..., N), has treatment assignment  $T_i$
- If  $T_i = 0$  for control condition
- $\mathbf{T}_i = (\mathsf{T}_{i,\mathsf{type}},\mathsf{T}_{i,\mathsf{stage}},\mathsf{T}_{i,\mathsf{money}},\mathsf{T}_{i,\mathsf{collab}},\mathsf{T}_{i,\mathsf{part.}})$
- $Y_i(\mathbf{T}_i)$ : participant i's Approval decision under treatment  $\mathbf{T}_i$
- Quantities of Interest
- Marginal Average Treatment Effect (MATE $_{T_i=k}$ )

- Participant i (i = 1, ..., N), has treatment assignment  $T_i$
- If  $T_i = 0$  for control condition
- $\mathbf{T}_i = (\mathsf{T}_{i,\mathsf{type}},\mathsf{T}_{i,\mathsf{stage}},\mathsf{T}_{i,\mathsf{money}},\mathsf{T}_{i,\mathsf{collab.}},\mathsf{T}_{i,\mathsf{part.}})$
- $Y_i(\mathbf{T}_i)$ : participant i's Approval decision under treatment  $\mathbf{T}_i$
- Quantities of Interest
- Marginal Average Treatment Effect (MATE $_{\mathsf{T}_i=k}$ )

$$MATE_{T_{j}=k} = \int E[Y(T_{j}=k, \mathbf{T}_{-j}) - Y(0)]dF_{\mathbf{T}_{-j}|T_{j}=k}$$

- Participant i (i = 1, ..., N), has treatment assignment  $T_i$
- If  $T_i = 0$  for control condition
- $\mathbf{T}_i = (\mathsf{T}_{i,\mathsf{type}},\mathsf{T}_{i,\mathsf{stage}},\mathsf{T}_{i,\mathsf{money}},\mathsf{T}_{i,\mathsf{collab}},\mathsf{T}_{i,\mathsf{part.}})$
- $Y_i(\mathbf{T}_i)$ : participant i's Approval decision under treatment  $\mathbf{T}_i$
- Quantities of Interest
- Marginal Average Treatment Effect (MATE $_{\mathsf{T}_j=k}$ )

$$MATE_{T_{j}=k} = \int E[Y(T_{j}=k, \mathbf{T}_{-j}) - Y(0)] dF_{\mathbf{T}_{-j}|T_{j}=k}$$

$$MATE_{T_{j}=k} = E[Y(T_{j}=k) - Y(0)]$$

- Participant i (i = 1, ..., N), has treatment assignment  $T_i$
- If  $T_i = 0$  for control condition
- $\mathbf{T}_i = (\mathsf{T}_{i,\mathsf{type}},\mathsf{T}_{i,\mathsf{stage}},\mathsf{T}_{i,\mathsf{money}},\mathsf{T}_{i,\mathsf{collab}},\mathsf{T}_{i,\mathsf{part.}})$
- $Y_i(\mathbf{T}_i)$ : participant i's Approval decision under treatment  $\mathbf{T}_i$
- Quantities of Interest
- Marginal Average Treatment Effect (MATE $_{\mathsf{T}_i=k}$ )

$$MATE_{T_j=k} = E[Y(T_j = k)|T_j = k] - E[Y(0)|T = 0]$$

- Participant i (i = 1, ..., N), has treatment assignment  $T_i$
- If  $T_i = 0$  for control condition
- $\mathbf{T}_i = (\mathsf{T}_{i,\mathsf{type}}, \mathsf{T}_{i,\mathsf{stage}}, \mathsf{T}_{i,\mathsf{money}}, \mathsf{T}_{i,\mathsf{collab}}, \mathsf{T}_{i,\mathsf{part}})$
- $Y_i(\mathbf{T}_i)$ : participant i's Approval decision under treatment  $\mathbf{T}_i$
- Quantities of Interest
- Marginal Average Treatment Effect (MATE $_{T_i=k}$ )

$$\begin{aligned} \mathsf{MATE}_{\mathsf{T}_{j}=k} &= \mathsf{E}[Y(\mathsf{T}_{j}=k)|\mathsf{T}_{j}=k] - \mathsf{E}[Y(0)|\mathsf{T}=0] \\ \mathsf{MATE}_{\mathsf{T}_{j}=k} &= \frac{\sum_{i=1}^{N} Y_{i} I(\mathsf{T}_{ij}=k)}{\sum_{i=1}^{N} I(\mathsf{T}_{ij}=k)} - \frac{\sum_{i=1}^{N} Y_{i} I(\mathsf{T}_{i}=0)}{\sum_{i=1}^{N} I(\mathsf{T}_{i}=0)} \end{aligned}$$

- Response may be conditional on respondent characteristics **x** 

- Response may be conditional on respondent characteristics **x**
- For example  $\mathbf{x} = (Conservative, Republican)$

- Response may be conditional on respondent characteristics **x**
- For example  $\mathbf{x} = (Conservative, Republican)$
- Marginal Conditional Average Treatment Effect (MCATE $_{\mathsf{T}_i=k,\mathbf{x}}$ )

- Response may be conditional on respondent characteristics  ${\bf x}$
- For example  $\mathbf{x} = (Conservative, Republican)$
- Marginal Conditional Average Treatment Effect (MCATE $_{\mathsf{T}_j=k,\mathbf{x}}$ )

$$MCATE_{T_i=k,x} = E[Y(T_j=k) - Y(0)|x]$$

$$MCATE_{T_i=k,\mathbf{x}} = E[Y(T_j=k)|\mathbf{x}] - E[Y(0)|\mathbf{x}]$$

$$\begin{aligned} \mathsf{MCATE}_{\mathsf{T}_{j}=k,\mathbf{x}} &= & \mathsf{E}[Y(\mathsf{T}_{j}=k)|\mathbf{x}] - \mathsf{E}[Y(0)|\mathbf{x}] \\ \mathsf{MCATE}_{\mathsf{T}_{j}=k,\mathbf{x}} &= & \frac{\sum_{i=1}^{N} Y_{i} I(\mathsf{T}_{j}=k,\mathbf{x}_{i}=\mathbf{x})}{\sum_{i=1}^{N} I(\mathsf{T}_{i}=k,\mathbf{x}_{i}=\mathbf{x})} - \frac{\sum_{i=1}^{N} Y_{i} I(\mathsf{T}_{i}=0,\mathbf{x}_{i}=\mathbf{x})}{\sum_{i=1}^{N} I(\mathsf{T}_{i}=0,\mathbf{x}_{i}=\mathbf{x})} \end{aligned}$$

$$\begin{aligned} \mathsf{MCATE}_{\mathsf{T}_{j}=k,\mathbf{x}} &= & \mathsf{E}[Y(\mathsf{T}_{j}=k)|\mathbf{x}] - \mathsf{E}[Y(0)|\mathbf{x}] \\ \mathsf{MC}\widehat{\mathsf{ATE}}_{\mathsf{T}_{j}=k,\mathbf{x}} &= & & \frac{\sum_{i=1}^{N} Y_{i} I(\mathsf{T}_{j}=k,\mathbf{x}_{i}=\mathbf{x})}{\sum_{i=1}^{N} I(\mathsf{T}_{j}=k,\mathbf{x}_{i}=\mathbf{x})} - \frac{\sum_{i=1}^{N} Y_{i} I(\mathsf{T}_{i}=0,\mathbf{x}_{i}=\mathbf{x})}{\sum_{i=1}^{N} I(\mathsf{T}_{i}=0,\mathbf{x}_{i}=\mathbf{x})} \end{aligned}$$

 Curse of Dimensionality: highly variable estimates, (sometimes) empty strata

$$\begin{aligned} & \mathsf{MCATE}_{\mathsf{T}_{j}=k,\mathbf{x}} &= & \mathsf{E}[Y(\mathsf{T}_{j}=k)|\mathbf{x}] - \mathsf{E}[Y(0)|\mathbf{x}] \\ & \mathsf{MCATE}_{\mathsf{T}_{j}=k,\mathbf{x}} &= & \frac{\sum_{i=1}^{N} Y_{i} I(\mathsf{T}_{j}=k,\mathbf{x}_{i}=\mathbf{x})}{\sum_{i=1}^{N} I(\mathsf{T}_{j}=k,\mathbf{x}_{i}=\mathbf{x})} - \frac{\sum_{i=1}^{N} Y_{i} I(\mathsf{T}_{i}=0,\mathbf{x}_{i}=\mathbf{x})}{\sum_{i=1}^{N} I(\mathsf{T}_{i}=0,\mathbf{x}_{i}=\mathbf{x})} \end{aligned}$$

- Curse of Dimensionality: highly variable estimates, (sometimes) empty strata
- Separate systematic differences from noise → data and assumptions

$$\begin{aligned} \mathsf{MCATE}_{\mathsf{T}_j=k,\mathbf{x}} &=& \mathsf{E}[Y(\mathsf{T}_j=k)|\mathbf{x}] - \mathsf{E}[Y(0)|\mathbf{x}] \\ \mathsf{MC}\widehat{\mathsf{ATE}_{\mathsf{T}_j=k,\mathbf{x}}} &=& \frac{\sum_{i=1}^N Y_i I(\mathsf{T}_j=k,\mathbf{x}_i=\mathbf{x})}{\sum_{i=1}^N I(\mathsf{T}_j=k,\mathbf{x}_i=\mathbf{x})} - \frac{\sum_{i=1}^N Y_i I(\mathsf{T}_i=0,\mathbf{x}_i=\mathbf{x})}{\sum_{i=1}^N I(\mathsf{T}_i=0,\mathbf{x}_i=\mathbf{x})} \end{aligned}$$

- Curse of Dimensionality: highly variable estimates, (sometimes) empty strata
- Separate systematic differences from noise → data and assumptions Heterogeneous treatment effect methods

$$\begin{aligned} &\mathsf{MCATE}_{\mathsf{T}_j=k,\mathbf{x}} &=& \mathsf{E}[Y(\mathsf{T}_j=k)|\mathbf{x}] - \mathsf{E}[Y(0)|\mathbf{x}] \\ &\mathsf{MC}\widehat{\mathsf{ATE}_{\mathsf{T}_j=k,\mathbf{x}}} &=& \frac{\sum_{i=1}^N Y_i I(\mathsf{T}_j=k,\mathbf{x}_i=\mathbf{x})}{\sum_{i=1}^N I(\mathsf{T}_j=k,\mathbf{x}_i=\mathbf{x})} - \frac{\sum_{i=1}^N Y_i I(\mathsf{T}_i=0,\mathbf{x}_i=\mathbf{x})}{\sum_{i=1}^N I(\mathsf{T}_i=0,\mathbf{x}_i=\mathbf{x})} \end{aligned}$$

- Curse of Dimensionality: highly variable estimates, (sometimes) empty strata
- Separate systematic differences from noise → data and assumptions Heterogeneous treatment effect methods
  - LASSO, Find It (Imai and Ratkovic, 2013) → sparsity

$$\begin{aligned} & \mathsf{MCATE}_{\mathsf{T}_{j}=k,\mathbf{x}} &= & \mathsf{E}[Y(\mathsf{T}_{j}=k)|\mathbf{x}] - \mathsf{E}[Y(0)|\mathbf{x}] \\ & \mathsf{MCATE}_{\mathsf{T}_{j}=k,\mathbf{x}} &= & \frac{\sum_{i=1}^{N} Y_{i} I(\mathsf{T}_{j}=k,\mathbf{x}_{i}=\mathbf{x})}{\sum_{i=1}^{N} I(\mathsf{T}_{j}=k,\mathbf{x}_{i}=\mathbf{x})} - \frac{\sum_{i=1}^{N} Y_{i} I(\mathsf{T}_{i}=0,\mathbf{x}_{i}=\mathbf{x})}{\sum_{i=1}^{N} I(\mathsf{T}_{i}=0,\mathbf{x}_{i}=\mathbf{x})} \end{aligned}$$

- Curse of Dimensionality: highly variable estimates, (sometimes) empty strata
- Separate systematic differences from noise → data and assumptions Heterogeneous treatment effect methods
  - LASSO, Find It (Imai and Ratkovic, 2013) → sparsity
  - Ridge, KRLS (Hainmueller and Hazlett, 2013) → flexible surface, dense

$$\begin{aligned} &\mathsf{MCATE}_{\mathsf{T}_j=k,\mathbf{x}} &=& \mathsf{E}[Y(\mathsf{T}_j=k)|\mathbf{x}] - \mathsf{E}[Y(0)|\mathbf{x}] \\ &\mathsf{MC}\widehat{\mathsf{ATE}_{\mathsf{T}_j=k,\mathbf{x}}} &=& \frac{\sum_{i=1}^N Y_i I(\mathsf{T}_j=k,\mathbf{x}_i=\mathbf{x})}{\sum_{i=1}^N I(\mathsf{T}_j=k,\mathbf{x}_i=\mathbf{x})} - \frac{\sum_{i=1}^N Y_i I(\mathsf{T}_i=0,\mathbf{x}_i=\mathbf{x})}{\sum_{i=1}^N I(\mathsf{T}_i=0,\mathbf{x}_i=\mathbf{x})} \end{aligned}$$

- Curse of Dimensionality: highly variable estimates, (sometimes) empty strata
- Separate systematic differences from noise → data and assumptions Heterogeneous treatment effect methods
  - LASSO, Find It (Imai and Ratkovic, 2013) → sparsity
  - Ridge, KRLS (Hainmueller and Hazlett, 2013) → flexible surface, dense
  - Model m to estimate some function  $g_m(T_i = k, x)$

$$\begin{aligned} \mathsf{MCATE}_{\mathsf{T}_j = k, \mathbf{x}} &= \mathsf{E}[Y(\mathsf{T}_j = k) | \mathbf{x}] - \mathsf{E}[Y(0) | \mathbf{x}] \\ \mathsf{MC}\widehat{\mathsf{ATE}}_{\mathsf{T}_j = k, \mathbf{x}} &= \widehat{g}_m(\mathsf{T}_j = k, \mathbf{x}) - \widehat{g}_m(0, \mathbf{x}) \end{aligned}$$

- Curse of Dimensionality: highly variable estimates, (sometimes) empty strata
- Separate systematic differences from noise → data and assumptions Heterogeneous treatment effect methods
  - LASSO, Find It (Imai and Ratkovic, 2013) → sparsity
  - Ridge, KRLS (Hainmueller and Hazlett, 2013) → flexible surface, dense
  - Model m to estimate some function  $g_m(\mathsf{T}_j=k, \pmb{x})$
- Perform well:  $g_m(T_j = k, x)$  accurately estimates response surface  $(E[Y(T_j = k)|x])$

$$\begin{aligned} \mathsf{MCATE}_{\mathsf{T}_j = k, \mathbf{x}} &= \mathsf{E}[Y(\mathsf{T}_j = k) | \mathbf{x}] - \mathsf{E}[Y(0) | \mathbf{x}] \\ \mathsf{MC}\widehat{\mathsf{ATE}}_{\mathsf{T}_j = k, \mathbf{x}} &= \widehat{g}_m(\mathsf{T}_j = k, \mathbf{x}) - \widehat{g}_m(0, \mathbf{x}) \end{aligned}$$

- Curse of Dimensionality: highly variable estimates, (sometimes) empty strata
- Separate systematic differences from noise → data and assumptions Heterogeneous treatment effect methods
  - LASSO, Find It (Imai and Ratkovic, 2013) → sparsity
  - Ridge, KRLS (Hainmueller and Hazlett, 2013) → flexible surface, dense
  - Model m to estimate some function  $g_m(\mathsf{T}_j=k, \pmb{x})$
- Perform well:  $g_m(T_j = k, x)$  accurately estimates response surface  $(E[Y(T_j = k)|x])$
- Perform well: accurate out of sample prediction and classification (van der Laan et al 2007, Raftery et al 2005)

$$\begin{aligned} \mathsf{MCATE}_{\mathsf{T}_j = k, \mathbf{x}} &= \mathsf{E}[Y(\mathsf{T}_j = k) | \mathbf{x}] - \mathsf{E}[Y(0) | \mathbf{x}] \\ \mathsf{MC}\widehat{\mathsf{ATE}}_{\mathsf{T}_j = k, \mathbf{x}} &= \widehat{g}_m(\mathsf{T}_j = k, \mathbf{x}) - \widehat{g}_m(0, \mathbf{x}) \end{aligned}$$

- Curse of Dimensionality: highly variable estimates, (sometimes) empty strata
- Separate systematic differences from noise → data and assumptions Heterogeneous treatment effect methods
  - LASSO, Find It (Imai and Ratkovic, 2013) → sparsity
  - Ridge, KRLS (Hainmueller and Hazlett, 2013) → flexible surface, dense
  - Model m to estimate some function  $g_m(T_j = k, x)$
- Perform well:  $g_m(T_j = k, \mathbf{x})$  accurately estimates response surface  $(E[Y(T_j = k)|\mathbf{x}])$
- Perform well: accurate out of sample prediction and classification (van der Laan et al 2007, Raftery et al 2005)

Create ensemble: weighting methods by (unique) out of sample predictive performance

$$MC\widehat{ATE}_{\mathsf{T}_{j}=k,\mathbf{x}} = \sum_{m=1}^{M} \widehat{\pi}_{m}(\widehat{g}_{m}(\mathsf{T}_{j}=k,\mathbf{x}) - \widehat{g}_{m}(0,\mathbf{x}))$$

$$MC\widehat{\mathsf{ATE}}_{\mathsf{T}_j=k,\mathbf{x}} = \sum_{m=1}^{M} \widehat{\boldsymbol{\pi}}_m(\widehat{g}_m(\mathsf{T}_j=k,\mathbf{x}) - \widehat{g}_m(0,\mathbf{x}))$$

$$MC\widehat{\mathsf{ATE}}_{\mathsf{T}_j=k,\mathbf{x}} = \sum_{m=1}^M \widehat{\pi}_m(\widehat{g}_m(\mathsf{T}_j=k,\mathbf{x}) - \widehat{g}_m(0,\mathbf{x}))$$

- Suppose we have M (m = 1, ..., M) models.

$$MC\widehat{\mathsf{ATE}_{\mathsf{T}_j=k,\mathbf{x}}} = \sum_{m=1}^M \widehat{\pi}_m(\widehat{g}_m(\mathsf{T}_j=k,\mathbf{x}) - \widehat{g}_m(0,\mathbf{x}))$$

- Estimate weights  $(\widehat{\pi}_m)$ 

$$MC\widehat{ATE}_{\mathsf{T}_j=k,\mathbf{x}} = \sum_{m=1}^{M} \widehat{\pi}_m(\widehat{g}_m(\mathsf{T}_j=k,\mathbf{x}) - \widehat{g}_m(0,\mathbf{x}))$$

- Estimate weights  $(\widehat{\pi}_m)$ 
  - 10-fold cross validation: generate M out of sample predictions for each observation

$$MC\widehat{ATE}_{\mathsf{T}_j=k,\mathbf{x}} = \sum_{m=1}^{M} \widehat{\pi}_m(\widehat{g}_m(\mathsf{T}_j=k,\mathbf{x}) - \widehat{g}_m(0,\mathbf{x}))$$

- Estimate weights  $(\widehat{\pi}_m)$ 
  - 10-fold cross validation: generate M out of sample predictions for each observation

$$\widehat{\mathbf{Y}}_i = (\widehat{Y}_{i1}, \widehat{Y}_{i2}, \dots, \widehat{Y}_{iM})$$

- Suppose we have M (m = 1, ..., M) models.

$$MC\widehat{ATE}_{\mathsf{T}_j=k,\mathbf{x}} = \sum_{m=1}^{M} \widehat{\pi}_m(\widehat{g}_m(\mathsf{T}_j=k,\mathbf{x}) - \widehat{g}_m(0,\mathbf{x}))$$

- Estimate weights  $(\widehat{\pi}_m)$ 
  - 10-fold cross validation: generate M out of sample predictions for each observation

$$\widehat{\mathbf{Y}}_i = (\widehat{Y}_{i1}, \widehat{Y}_{i2}, \dots, \widehat{Y}_{iM})$$

- Estimate weights with constrained regression:

- Suppose we have M (m = 1, ..., M) models.

$$MC\widehat{ATE}_{\mathsf{T}_j=k,\mathbf{x}} = \sum_{m=1}^{M} \widehat{\pi}_m(\widehat{g}_m(\mathsf{T}_j=k,\mathbf{x}) - \widehat{g}_m(0,\mathbf{x}))$$

- Estimate weights  $(\widehat{\pi}_m)$ 
  - 10-fold cross validation: generate M out of sample predictions for each observation

$$\widehat{\mathbf{Y}}_i = (\widehat{Y}_{i1}, \widehat{Y}_{i2}, \dots, \widehat{Y}_{iM})$$

- Estimate weights with constrained regression:

$$Y_i = \sum_{m=1}^{M} \pi_m \hat{Y}_{im} + \epsilon_i$$

- Suppose we have M (m = 1, ..., M) models.

$$MC\widehat{ATE}_{\mathsf{T}_j=k,\mathbf{x}} = \sum_{m=1}^{M} \widehat{\pi}_m(\widehat{g}_m(\mathsf{T}_j=k,\mathbf{x}) - \widehat{g}_m(0,\mathbf{x}))$$

- Estimate weights  $(\widehat{\pi}_m)$ 
  - 10-fold cross validation: generate M out of sample predictions for each observation

$$\widehat{\mathbf{Y}}_i = (\widehat{Y}_{i1}, \widehat{Y}_{i2}, \dots, \widehat{Y}_{iM})$$

- Estimate weights with constrained regression:

$$Y_i = \sum_{m=1}^{M} \pi_m \hat{Y}_{im} + \epsilon_i$$

where we impose constraints:  $\pi_m \geq 0$  and  $\sum_{m=1}^{M} \pi_m = 1$ .

- Suppose we have M (m = 1, ..., M) models.

$$\widehat{\mathsf{MCATE}_{\mathsf{T}_j=k,\mathbf{x}}} = \sum_{m=1}^M \widehat{\pi}_m(\widehat{g}_m(\mathsf{T}_j=k,\mathbf{x}) - \widehat{g}_m(0,\mathbf{x}))$$

- Estimate weights  $(\widehat{\pi}_m)$ 
  - 10-fold cross validation: generate  ${\it M}$  out of sample predictions for each observation

$$\widehat{\mathbf{Y}}_i = (\widehat{Y}_{i1}, \widehat{Y}_{i2}, \dots, \widehat{Y}_{iM})$$

- Estimate weights with constrained regression:

$$Y_i = \sum_{m=1}^{M} \pi_m \hat{Y}_{im} + \epsilon_i$$

where we impose constraints:  $\pi_m \geq 0$  and  $\sum_{m=1}^{M} \pi_m = 1$ .

- Result  $\widehat{\pi}_m$  for each method

- Suppose we have M (m = 1, ..., M) models.

$$MC\widehat{ATE}_{\mathsf{T}_j=k,\mathbf{x}} = \sum_{m=1}^{M} \widehat{\pi}_m(\widehat{g}_m(\mathsf{T}_j=k,\mathbf{x}) - \widehat{g}_m(0,\mathbf{x}))$$

- Estimate weights  $(\widehat{\pi}_m)$ 
  - 10-fold cross validation: generate  ${\it M}$  out of sample predictions for each observation

$$\widehat{\mathbf{Y}}_i = (\widehat{Y}_{i1}, \widehat{Y}_{i2}, \dots, \widehat{Y}_{iM})$$

- (Alternatively) Estimate weights from mixture model (EBMA) (Raftery et al 2005; Montgomery, Hollenback, Ward 2012) → EM, Gibbs, Variational Approximation

- Suppose we have M (m = 1, ..., M) models.

$$MC\widehat{ATE}_{T_j=k,\mathbf{x}} = \sum_{m=1}^{M} \widehat{\pi}_m(\widehat{g}_m(T_j=k,\mathbf{x}) - \widehat{g}_m(0,\mathbf{x}))$$


- Estimate weights  $(\widehat{\pi}_m)$
- Estimate  $\widehat{g}_m(\mathsf{T}_j=k,\mathbf{x}) \leadsto \mathsf{Apply}$  all M models to entire data set

- Suppose we have M (m = 1, ..., M) models.


$$MCA\widehat{\mathsf{TE}_{\mathsf{T}_j=k,\mathsf{x}_{\mathsf{new}}}} = \sum_{m=1}^{M} \widehat{\pi}_m(\widehat{g}_m(\mathsf{T}_j=k,\mathsf{x}_{\mathsf{new}}) - \widehat{g}_m(0,\mathsf{x}_{\mathsf{new}}))$$

- Estimate weights  $(\widehat{\pi}_m)$
- Estimate  $\widehat{g}_m(\mathsf{T}_j=k,\mathbf{x}) \leadsto \mathsf{Apply}$  all M models to entire data set
- Generate effects of interest (perhaps weighting to other population)  $\mathbf{x}_{\text{new}}$


### Monte Carlo Evidence



### Monte Carlo Evidence



#### Monte Carlo Evidence



Ensembles outperform constituent methods  $\rightsquigarrow$  ensembles place weight on better performing method

Recall: experiment to assess effects of credit claiming on approval ↔ 1,074 participants (MTurk)

Recall: experiment to assess effects of credit claiming on approval  $\rightsquigarrow$  1,074 participants (MTurk)

Apply ensemble method (7 constituent methods, 10 fold cross validation), including treatments and Partisanship and Ideology.

Recall: experiment to assess effects of credit claiming on approval  $\rightsquigarrow$  1,074 participants (MTurk)

Apply ensemble method (7 constituent methods, 10 fold cross validation), including treatments and Partisanship and Ideology.

Positive weight on three methods:

Recall: experiment to assess effects of credit claiming on approval ↔ 1,074 participants (MTurk)

Apply ensemble method (7 constituent methods, 10 fold cross validation), including treatments and Partisanship and Ideology.

Positive weight on three methods:

1) LASSO (0.62)

Recall: experiment to assess effects of credit claiming on approval ↔ 1,074 participants (MTurk)

Apply ensemble method (7 constituent methods, 10 fold cross validation), including treatments and Partisanship and Ideology.

Positive weight on three methods:


- 1) LASSO (0.62)
- 2) KRLS (0.24)


Recall: experiment to assess effects of credit claiming on approval ↔ 1,074 participants (MTurk)


Apply ensemble method (7 constituent methods, 10 fold cross validation), including treatments and Partisanship and Ideology.

Positive weight on three methods:

- 1) LASSO (0.62)
- 2) KRLS (0.24)
- 3) Find it (0.14)










990


13 / 14







990







Treatment Effect



→ Constituents evaluate expenditures using qualitative information, rather than numerical facts

#### Issues with experimental design

- Treatments are conditional on what else is included: averaging other quantities  $\neq$  to excluding other quantities

- Treatments are conditional on what else is included: averaging other quantities  $\neq$  to excluding other quantities
- Potential for fishing is massive → pre analysis plan and split-sample design

- Treatments are conditional on what else is included: averaging other quantities  $\neq$  to excluding other quantities
- Potential for fishing is massive → pre analysis plan and split-sample design
- Assumption about the way information delivered:

- Treatments are conditional on what else is included: averaging other quantities  $\neq$  to excluding other quantities
- Potential for fishing is massive → pre analysis plan and split-sample design
- Assumption about the way information delivered:
  - 1) We know salient dimensions

- Treatments are conditional on what else is included: averaging other quantities  $\neq$  to excluding other quantities
- Potential for fishing is massive → pre analysis plan and split-sample design
- Assumption about the way information delivered:
  - 1) We know salient dimensions
  - 2) We're constructing effects that correspond with effects in reality

#### Issues with experimental design

- Treatments are conditional on what else is included: averaging other quantities  $\neq$  to excluding other quantities
- Potential for fishing is massive → pre analysis plan and split-sample design
- Assumption about the way information delivered:
  - 1) We know salient dimensions
  - 2) We're constructing effects that correspond with effects in reality

Wednesday's lecture will introduce discovery methods