

Vv255 Applied Calculus III

Recitation VI

LIU Xieyang

Teaching Assistant

University of Michigan - Shanghai Jiaotong University
Joint Institute

Summer Term 2015

Contents

Lecture 12: Differential

Lecture 13: Extreme values and saddle points

Lecture 14: Lagrange Multipliers

Linearization & Differential

Linearization:

The **linearization** of a function $f(x, y)$ at a point (x_0, y_0) , where f is differentiable,

$$L(x, y) = f(x_0, y_0) + f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0).$$

Differential: If we move from (x_0, y_0) to a point $(x_0 + dx, y_0 + dy)$ nearby, the resulting change

$$df = f_x(x_0, y_0)dx + f_y(x_0, y_0)dy$$

in the linearization of f is called the **total differential** of f .

Linearization & Differential (cont.)

- ▶ The plane $z = L(x, y)$ is tangent to the surface $z = f(x, y)$ at the point (x_0, y_0) .
- ▶ The linearization of a function of two variables is a **tangent-plane approximation** in the same way that the linearization of a function of a single variable is a **tangent line approximation**.

Contents

Lecture 12: Differential

Lecture 13: Extreme values and saddle points

Lecture 14: Lagrange Multipliers

Extreme Values for function of several variables

Let (a, b) be a point in the domain D of a function $f(x, y)$, then $f(a, b)$ is

- Relative **maximum** value of f if

$$f(a, b) \geq f(x, y) \quad \text{for some disk with center } (a, b).$$

- Relative **minimum** value of f if

$$f(a, b) \leq f(x, y) \quad \text{for some disk with center } (a, b).$$

- *Absolute* **maximum** value of f for an interval I if

$$f(a, b) \geq f(x, y) \quad \text{for all points } (x, y) \text{ in } R.$$

- *Absolute* **minimum** value of f for an interval I if

$$f(a, b) \leq f(x, y) \quad \text{for all points } (x, y) \text{ in } R.$$

Finding Local Extrema

Procedures for finding **local/relative** extrema:

Step 1: **Gradient Test: Finding Critical Points**

$$\text{critical points} \begin{cases} 1. & \nabla f \text{ does not exist.} \\ 2. & \nabla f = 0. \end{cases}$$




Figure: Saddle point: In general, we will say that a function has a saddle point P if there are two distinct vertical planes through P such that P in one of the planes is a local maximum and P in the other is a local minimum.

Finding Local Extrema (cont.)

Step 2: Second derivative Test: Hessian Matrix

$$\mathbf{H} = \begin{bmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{bmatrix} \quad \mathbf{H} = \begin{bmatrix} f_{xx} & f_{xy} & f_{xz} \\ f_{yx} & f_{yy} & f_{yz} \\ f_{zx} & f_{zy} & f_{zz} \end{bmatrix}$$

Suppose f is differentiable and $\nabla f = 0$ at a point P_0 , and if

all the eigenvalues of \mathbf{H} at P_0 are positive \implies local minimum

all the eigenvalues of \mathbf{H} at P_0 are negative \implies local maximum

\mathbf{H} at P_0 has both positive and negative eigenvalues \implies saddle point

One of the eigenvalues of \mathbf{H} is zero \implies inconclusive

Finding Global Extrema

Procedures for finding **global/Absolute** extrema:

1. Find the local extreme values of f in the domain D .
2. Find the local extreme values of f on boundary of the domain D .
 - ▶ Direct examination;
 - ▶ Lagrange multiplier (will be talked about later)
3. Compare values in step 1. and step 2., the largest of them is the global maximum, the smallest is the global minimum.

Contents

Lecture 12: Differential

Lecture 13: Extreme values and saddle points

Lecture 14: Lagrange Multipliers

Geometry Basis of Lagrange Multipliers




Figure: Maximum of f is 400; minimum of f is 200

To motivate the method of Lagrange multipliers, suppose that we are trying to maximize a function $f(x, y)$ subject to the constraint $g(x, y) = 0$. Geometrically, this means that we are looking for a point (x_0, y_0) on the graph of the constraint curve at which $f(x, y)$ is as large as possible.

Geometry Basis of Lagrange Multipliers (cont.)




Figure: Maximum of f is 400; minimum of f is 200

To help locate such a point, let us construct a contour plot of $f(x, y)$ in the same coordinate system as the graph of $g(x, y) = 0$. For example, the graph on the left shows some typical level curves of $f(x, y) = c$, which we have labeled $c = 100, 200, 300, 400, 500$ for purpose of illustration. In this figure, each point of intersection of $g(x, y) = 0$ with a level curve is a candidate for a solution, since these points lie on the constraint curve. Among the seven such intersections shown in the figure, the maximum value of $f(x, y)$ occurs at the intersection (x_0, y_0) where $f(x, y)$ has a value of 400.

Geometry Basis of Lagrange Multipliers (cont.)




Figure: Maximum of f is 400; minimum of f is 200

Note that at (x_0, y_0) the constraint curve and the level curve just **touch** and thus have a **common tangent line** at this point.

Since $\nabla f(x_0, y_0)$ is **normal** to the level curve $f(x, y) = 400$ at (x_0, y_0) , and since $\nabla g(x_0, y_0)$ is **normal** to the constraint curve $g(x, y) = 0$ at (x_0, y_0) , we conclude that the vectors $\nabla f(x_0, y_0)$ and $\nabla g(x_0, y_0)$ must be **parallel**. That is

$$\nabla f(x_0, y_0) = \lambda \nabla g(x_0, y_0) \quad \text{for some scalar } \lambda$$

Lagrange Multipliers

Ultimate Goal:

Minimize/Maximize f subject to the constraint(s) $g_1 = k_1, g_2 = k_2, \dots,$
 $g_n = k_n,$
 where f and g_1, \dots, g_n are functions of several variables.

Method:

Step 1: Solve the simultaneous equations:

$$\begin{aligned}\nabla f &= \lambda \nabla g_1 + \mu \nabla g_2 + \dots \\ g_1 &= k_1 \\ g_2 &= k_2 \\ &\dots\end{aligned}$$

Step 2:

Evaluate f at all the points (x, y, z, \dots) that result from step 1. The largest of these values is the maximum value of f ; the smallest is the minimum value of f .