

# Introduction to Dynamic Spatio-Temporal Models

Erick A. Chacon-Montalvan<sup>1,2</sup>, Erick A. Chacon-Montalvan<sup>2</sup>, Erick A. Chacon-Montalvan<sup>3</sup>, Erick A. Chacon-Montalvan<sup>3</sup>

<sup>1</sup>*CHICAS, Medical School, Lancaster University, Lancaster, United Kingdom*

<sup>2</sup>*CHICAS, Medical School, Lancaster University, Lancaster, United Kingdom*

May 15, 2019

## Abstract

In the geometric distribution,  $\theta$  represents the probability of succes in each trial. Furthermore, it should follow that  $y$  is countable and greater than one. In general, in the negative binomial distribution,  $y$  is the number of trials until get the  $k$  first success. For this reason, the negative binomial distribution has certain assumptions considering that each event come from a Bernoulli distribution.(i) Each  $x_i$  is independent of any. In the geometric distribution,  $\theta$  represents the probability of succes in each trial. Furthermore, it should follow that  $y$  is countable and greater than one. In general, in the negative binomial distribution,  $y$  is the number of trials until get the  $k$  first success. For this reason, the negative binomial distribution has certain assumptions considering that each event come from a Bernoulli distribution.(i) Each  $x_i$  is independent of any.

**Keywords:** Spatio-temporal, Climate variability, Malaria prevalence, geostatistics, Matern.

## 1. Introduction

In the geometric distribution,  $\theta$  represents the probability of succes in each trial. Furthermore, it should follow that  $y$  is countable and greater than one. In general, in the negative binomial distribution,  $y$  is the number of trials until get the  $k$  first success. In the geometric distribution,  $\theta$  represents the probability of succes in each trial. Furthermore, it should follow that  $y$  is countable and greater than one. In general, in the negative binomial distribution,  $y$  is the number of trials until get the  $k$  first success.

In the geometric distribution,  $\theta$  represents the probability of succes in each trial. Furthermore, it should follow that  $y$  is countable and greater than one. In general, in the negative binomial distribution,  $y$  is the number of trials until get the  $k$  first success. In the geometric distribution,  $\theta$  represents the probability of succes in each trial. Furthermore, it should follow that  $y$  is countable and greater than one. In general, in the negative binomial distribution,  $y$  is the number of trials until get the  $k$  first success. In the geometric distribution,  $\theta$  represents the probability of succes in each trial. Furthermore, it should follow that  $y$  is countable and greater than one. In general, in the negative binomial distribution,  $y$  is the

number of trials until get the  $k$  first success. In the geometric distribution,  $\theta$  represents the probability of succes in each trial. Furthermore, it should follow that  $y$  is countable and greater than one. In general, in the negative binomial distribution,  $y$  is the number of trials until get the  $k$  first success.

**Statement:**

$$p(y > r|\theta) = (1 - \theta)^r, \quad (1)$$

where  $r = 1, 2, \dots$

**Proof:**

$$p(y > r|\theta) = 1 - p(y \leq r|\theta)$$

$$p(y > r|\theta) = 1 - \sum_{k=1}^r (1 - \theta)^{k-1} \theta$$

$$p(y > r|\theta) = 1 - \theta \sum_{k=1}^r (1 - \theta)^{k-1}$$

$$p(y > r|\theta) = 1 - \theta \left[ \frac{1 - (1 - \theta)^r}{1 - (1 - \theta)} \right]$$

$$p(y > r|\theta) = 1 - \theta \left[ \frac{1 - (1 - \theta)^r}{\theta} \right]$$

For this reason, the negative binomial distribution has certain assumptions considering that each event come from a Bernoulli distribution. (i) Each  $x_i$  is independent of any other  $x_j$ , (ii). They are identically distributed  $x_i \sim \theta^{x_i} (1 - \theta)^{1-x_i}$ , where  $i = 1, 2, \dots$  (iii) The variance is greater than the mean. For this reason is used in overdispersed countable data. (iv) This distribution is unimodal.

For this reason, the negative binomial distribution has certain assumptions considering that each event come from a Bernoulli distribution. (i) Each  $x_i$  is independent of any other  $x_j$ , (ii). They are identically distributed  $x_i \sim \theta^{x_i} (1 - \theta)^{1-x_i}$ , where  $i = 1, 2, \dots$  (iii) The variance is greater than the mean. For this reason is used in overdispersed countable data. (iv) This distribution is unimodal.

For this reason, the negative binomial distribution has certain assumptions considering that each event come from a Bernoulli distribution. (i) Each  $x_i$  is independent of any other  $x_j$ , (ii). They are identically distributed  $x_i \sim \theta^{x_i} (1 - \theta)^{1-x_i}$ , where  $i = 1, 2, \dots$  (iii) The variance is greater than the mean. For this reason is used in overdispersed countable data. (iv) This distribution is unimodal.

For this reason, the negative binomial distribution has certain assumptions considering that each event come from a Bernoulli distribution. (i) Each  $x_i$  is independent of any other  $x_j$ , (ii). They are identically distributed  $x_i \sim \theta^{x_i} (1 - \theta)^{1-x_i}$ , where  $i = 1, 2, \dots$  (iii) The variance is greater than the mean. For this reason is used in overdispersed countable data. (iv) This distribution is unimodal.

In the geometric distribution,  $\theta$  represents the probability of succes in each trial. Fur-

thermore, it should follow that  $y$  is countable and greater than one. In general, in the negative binomial distribution,  $y$  is the number of trials until get the  $k$  first success. For this reason, the negative binomial distribution has certain assumptions considering that each event come from a Bernoulli distribution. (i) Each  $x_i$  is independent of any other  $x_j$ , (ii). They are identically distributed  $x_i \sim \theta^{x_i}(1 - \theta)^{1-x_i}$ , where  $i = 1, 2, \dots$  (iii) The variance is greater than the mean. For this reason is used in overdispersed countable data. (iv) This distribution is unimodal.

## 2. Section 2

In the geometric distribution,  $\theta$  represents the probability of succes in each trial. Furthermore, it should follow that  $y$  is countable and greater than one. In general, in the negative binomial distribution,  $y$  is the number of trials until get the  $k$  first success. For this reason, the negative binomial distribution has certain assumptions considering that each event come from a Bernoulli distribution. (i) Each  $x_i$  is independent of any other  $x_j$ , (ii) They are identically distributed  $x_i \sim \theta^{x_i}(1 - \theta)^{1-x_i}$ , where  $i = 1, 2, \dots$  (iii) The variance is greater than the mean. For this reason is used in overdispersed countable data. (iv) This distribution is unimodal.

## 3. Section 1

In the geometric distribution,  $\theta$  represents the probability of succes in each trial. Furthermore, it should follow that  $y$  is countable and greater than one. In general, in the negative binomial distribution,  $y$  is the number of trials until get the  $k$  first success. For this reason, the negative binomial distribution has certain assumptions considering that each event come from a Bernoulli distribution. (i) Each  $x_i$  is independent of any other  $x_j$ , (ii) They are identically distributed  $x_i \sim \theta^{x_i}(1 - \theta)^{1-x_i}$ , where  $i = 1, 2, \dots$  (iii) The variance is greater than the mean. For this reason is used in overdispersed countable data. (iv) This distribution is unimodal.

## 4. Section 1

In the geometric distribution,  $\theta$  represents the probability of succes in each trial. Furthermore, it should follow that  $y$  is countable and greater than one. In general, in the negative binomial distribution,  $y$  is the number of trials until get the  $k$  first success. For this reason, the negative binomial distribution has certain assumptions considering that each event come from a Bernoulli distribution. (i) Each  $x_i$  is independent of any other  $x_j$ , (ii) They are identically distributed  $x_i \sim \theta^{x_i}(1 - \theta)^{1-x_i}$ , where  $i = 1, 2, \dots$  (iii) The variance is greater than the mean. For this reason is used in overdispersed countable data. (iv) This distribution is unimodal.

## 5. Section 1

In the geometric distribution,  $\theta$  represents the probability of succes in each trial. Furthermore, it should follow that  $y$  is countable and greater than one. In general, in the negative binomial distribution,  $y$  is the number of trials until get the  $k$  first success. For this reason, the negative binomial distribution has certain assumptions considering that

each event come from a Bernoulli distribution. (i) Each  $x_i$  is independent of any other  $x_j$ , (ii) They are identically distributed  $x_i \sim \theta^{x_i}(1 - \theta)^{1-x_i}$ , where  $i = 1, 2, \dots$  (iii) The variance is greater than the mean. For this reason is used in overdispersed countable data. (iv) This distribution is unimodal.