

Research Notes:

Introduction to Dynamic Spatio-Temporal Models

ERICK A. CHACON-MONTALVAN¹, ERICK A. CHACON-MONTALVAN²

MARCH 11, 2017

Abstract:

In the geometric distribution, θ represents the probability of succes in each trial. Furthermore, it should follow that y is countable and greater than one. In general, in the negative binomial distribution, y is the number of trials until get the k first success. For this reason, the negative binomial distribution has certain assumptions considering that each event come from a Bernoulli distribution.(i) Each x_i is independent of any.

Keywords:

Spatiotemporal, Climate variability, Malaria prevalence.

¹CHICAS, Medical School, Lancaster University, Lancaster, United Kingdom

²CHICAS, Medical School, Lancaster University, Lancaster, United Kingdom

1. Section 1

In the geometric distribution, θ represents the probability of succes in each trial. Furthermore, it should follow that y is countable and greater than one. In general, in the negative binomial distribution, y is the number of trials until get the k first success.

Statement:

$p(y > r|\theta) = (1 - \theta)^r$, where $r = 1, 2, \dots$

Proof:

$$\begin{aligned}
 p(y > r|\theta) &= 1 - p(y \leq r|\theta) \\
 p(y > r|\theta) &= 1 - \sum_{k=1}^r (1 - \theta)^{k-1} \theta \\
 p(y > r|\theta) &= 1 - \theta \sum_{k=1}^r (1 - \theta)^{k-1} \\
 p(y > r|\theta) &= 1 - \theta \left[\frac{1 - (1 - \theta)^r}{1 - (1 - \theta)} \right] \\
 p(y > r|\theta) &= 1 - \theta \left[\frac{1 - (1 - \theta)^r}{\theta} \right] \\
 p(y > r|\theta) &= 1 - [1 - (1 - \theta)^r] = (1 - \theta)^r \quad \blacksquare
 \end{aligned}$$

For this reason, the negative binomial distribution has certain assumptions considering that each event come from a Bernoulli distribution.(i) Each x_i is independent of any other x_j , (ii). They are identically distributed $x_i \sim \theta^{x_i} (1 - \theta)^{1-x_i}$, where $i = 1, 2, \dots$ (iii) The

variance is greater than the mean. For this reason is used in overdispersed countable data. (iv) This distribution is unimodal.

For this reason, the negative binomial distribution has certain assumptions considering that each event come from a Bernoulli distribution.(i) Each x_i is independent of any other x_j , (ii). They are identically distributed $x_i \sim \theta^{x_i} (1 - \theta)^{1-x_i}$, where $i = 1, 2, \dots$ (iii) The variance is greater than the mean. For this reason is used in overdispersed countable data. (iv) This distribution is unimodal.

For this reason, the negative binomial distribution has certain assumptions considering that each event come from a Bernoulli distribution.(i) Each x_i is independent of any other x_j , (ii). They are identically distributed $x_i \sim \theta^{x_i} (1 - \theta)^{1-x_i}$, where $i = 1, 2, \dots$ (iii) The variance is greater than the mean. For this reason is used in overdispersed countable data. (iv) This distribution is unimodal.

For this reason, the negative binomial distribution has certain assumptions considering that each event come from a Bernoulli distribution.(i) Each x_i is independent of any other x_j , (ii). They are identically distributed $x_i \sim \theta^{x_i} (1 - \theta)^{1-x_i}$, where $i = 1, 2, \dots$ (iii) The variance is greater than the mean. For this reason is used in overdispersed countable data. (iv) This distribution is unimodal.

In the geometric distribution, θ represents the probability of succes in each trial. Furthermore, it should follow that y is countable and greater than one. In general, in the negative binomial distribution, y is the number of trials until get the k first success. For this reason, the negative binomial distribution has certain assumptions considering that each event come from a Bernoulli distribution.(i) Each x_i is independent of any other x_j , (ii). They are identically distributed $x_i \sim \theta^{x_i} (1 - \theta)^{1-x_i}$, where $i = 1, 2, \dots$ (iii) The

variance is greater than the mean. For this reason is used in overdispersed countable data. (iv) This distribution is unimodal.

2. Section 1

In the geometric distribution, θ represents the probability of succes in each trial. Furthermore, it should follow that y is countable and greater than one. In general, in the negative binomial distribution, y is the number of trials until get the k first success. For this reason, the negative binomial distribution has certain assumptions considering that each event come from a Bernoulli distribution.(i) Each x_i is independent of any other x_j , (ii) They are identically distributed $x_i \sim \theta^{x_i}(1 - \theta)^{1-x_i}$, where $i = 1, 2, \dots$ (iii) The variance is greater than the mean. For this reason is used in overdispersed countable data. (iv) This distribution is unimodal.

3. Section 1

In the geometric distribution, θ represents the probability of succes in each trial. Furthermore, it should follow that y is countable and greater than one. In general, in the negative binomial distribution, y is the number of trials until get the k first success. For this reason, the negative binomial distribution has certain assumptions considering that each event come from a Bernoulli distribution.(i) Each x_i is independent of any other x_j , (ii) They are identically distributed $x_i \sim \theta^{x_i}(1 - \theta)^{1-x_i}$, where $i = 1, 2, \dots$ (iii) The variance is greater than the mean. For this reason is used in overdispersed countable data. (iv) This distribution is unimodal.

4. Section 1

In the geometric distribution, θ represents the probability of succes in each trial. Furthermore, it should follow that y is countable and greater than one. In general, in the negative binomial distribution, y is the number of trials until get the k first success. For this reason, the negative binomial distribution has certain assumptions considering that each event come from a Bernoulli distribution.(i) Each x_i is independent of any other x_j , (ii) They are identically distributed $x_i \sim \theta^{x_i}(1 - \theta)^{1-x_i}$, where $i = 1, 2, \dots$ (iii) The variance is greater than the mean. For this reason is used in overdispersed countable data. (iv) This distribution is unimodal.

5. Section 1

In the geometric distribution, θ represents the probability of succes in each trial. Furthermore, it should follow that y is countable and greater than one. In general, in the negative binomial distribution, y is the number of trials until get the k first success. For this reason, the negative binomial distribution has certain assumptions considering that each event come from a Bernoulli distribution.(i) Each x_i is independent of any other x_j , (ii) They are identically distributed $x_i \sim \theta^{x_i}(1 - \theta)^{1-x_i}$, where $i = 1, 2, \dots$ (iii) The variance is greater than the mean. For this reason is used in overdispersed countable data. (iv) This distribution is unimodal.