HW1 Due: Feb 18th, 2022

### 1 GitHub setup

On GitHub

- Obtain student developer pack.
- Create a private repository math-6040-2022-spring (please substitute 6040 by 7260 if you are taking the graduate level). Add xji3 as your collaborators with write permission (instruction).

On your local machine:

- clone the repository: please refer to this webpage with instructions for your operating system.
- enter the folder: cd math-6040-2022-spring.
- after finishing the rest of the questions, save your file inside your git repository folder math-6040-2022-spring with name hw1.pdf. Please make it human-readable.
- now using git commands to stage this change: git add hw1.pdf
- commit: git commit -m "hw1 submission" (remember to replace the quotation mark)
- push to remote server: git push
- tag version hw1: git tag hw1 and push: git push -- tags.

Take a look at the tags on GitHub (instructions).

When submitting your hw, please email your instructor (xji4@tulane.edu) a link to your tag (instructions).

## 2 Show that for matrix $\mathbf{A} \in \mathbb{R}^{m \times n}$ :

$$rank(\mathbf{A}) = rank(\mathbf{A}^T) = rank(\mathbf{A}^T\mathbf{A}) = rank(\mathbf{A}\mathbf{A}^T)$$

#### 3 Show that for simple linear regression

$$y_i = \beta_0 + \beta_1 x_i + \epsilon_i,$$

the line  $(\hat{y}_i = \hat{\beta}_0 + \hat{\beta}_1 x_i)$  through the means of the variables (the point  $(\bar{x}, \bar{y})$ ) has  $\sum \epsilon_i = 0$ 

#### 4 JF excercise 5.1

Prove that the least-squares fit in simple regression analysis has the following properties:

- (a)  $\sum \hat{y}_i \epsilon_i = 0$ .
- (b)  $\sum (y_i \hat{y}_i)(\hat{y}_i \bar{y}) = \sum \epsilon_i(\hat{y}_i \bar{y}) = 0.$

# 5 JF excercise 5.6

Why is it the case that the multiple-correlation coefficient  $R^2$  can never get smaller when an explanatory variable is added to the regression equation? [Hint: Recall that the regression equation is fit by minimizing the residual sum of squares, which is equivalent as maximizing  $R^2$  (why?).]