

Clase-5.R

WINDOWS 10

2019-08-09

```
#Erick Raymundo Pérez Silva  
#Clase 5  
#09/08/19
```

```
# ANOVA -----
```

```
#ho no existe diferencia entre los tratamientos  
#ha al menos un grupo es diferente a los demas
```

```
arena <- c(6, 10, 8, 6, 14, 17, 9, 11, 7, 11)  
arcilla <- c(17, 15, 3, 11, 14, 12, 12, 8, 10, 13)  
limo <- c(13, 16, 9, 12, 15, 16, 17, 13, 18, 14)  
  
y.ton <- c(arena, arcilla, limo)  
suelo <- gl(3, 10, 30, labels =c("arena", "arcilla", "limo"))  
  
prod <- data.frame(suelo, y.ton)  
head(prod)
```

```
##      suelo y.ton  
## 1 arena      6  
## 2 arena     10  
## 3 arena      8  
## 4 arena      6  
## 5 arena     14  
## 6 arena     17
```

```
tapply(prod$y.ton, prod$suelo, mean)
```

```
##      arena arcilla      limo  
##      9.9      11.5      14.3
```

```
tapply(prod$y.ton, prod$suelo, var)
```

```
##      arena      arcilla      limo  
## 12.544444 15.388889  7.122222
```

```
shapiro.test(prod$y.ton)
```

```
##  
##  Shapiro-Wilk normality test  
##  
## data:  prod$y.ton  
## W = 0.97214, p-value = 0.5993
```


```
## Sirve para la homogeneidad de varianzas  
bartlett.test(prod$y.ton, prod$suelo)
```

```
##  
##  Bartlett test of homogeneity of variances
```

```
##
## data: prod$y.ton and prod$suelo
## Bartlett's K-squared = 1.2764, df = 2, p-value = 0.5283
fligner.test(prod$y.ton, prod$suelo)

##
## Fligner-Killeen test of homogeneity of variances
##
## data: prod$y.ton and prod$suelo
## Fligner-Killeen:med chi-squared = 0.36507, df = 2, p-value =
## 0.8332
## las varianzas son homogeneas

boxplot(prod$y.ton ~ prod$suelo, xlab = "Tipo de suelo",
        ylab = "Ton/ha", col = "blue")
```



```
##ANOVA
aov.suelo <- aov(prod$y.ton ~ prod$suelo)
aov.suelo

## Call:
## aov(formula = prod$y.ton ~ prod$suelo)
##
## Terms:
## prod$suelo Residuals
```


```
## Sum of Squares          99.2      315.5
## Deg. of Freedom         2         27
##
## Residual standard error: 3.41836
## Estimated effects may be unbalanced
```

```
summary(aov.suelo)
```

```
##              Df Sum Sq Mean Sq F value Pr(>F)
## prod$suelo    2   99.2   49.60   4.245  0.025 *
## Residuals    27  315.5   11.69
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
par(mfrow=c(2,2))
```

```
plot(aov(prod$y.ton ~ prod$suelo))
```



```
par(mfrow=c(1,1))
```

```
## si los datos estan cerca de la linea central en la grafica "nomal Q-Q"
## quiere decir que los datos vienen de una distribucion normal
```

```
##prueba de tukey sirve para saber cual de los tratamientos es diferente,
## 3 o mas medias
```


```
TukeyHSD(aov.suelo, conf.level = 0.95)
```

```
## Tukey multiple comparisons of means
```

```
##      95% family-wise confidence level
##
## Fit: aov(formula = prod$y.ton ~ prod$suelo)
##
## `$prod$suelo`
##           diff      lwr      upr      p adj
## arcilla-arena 1.6 -2.1903777 5.390378 0.5546301
## limo-arena    4.4  0.6096223 8.190378 0.0204414
## limo-arcilla  2.8 -0.9903777 6.590378 0.1785489
```

##lwr rango menor
##upr rango mayor

```
plot(TukeyHSD(aov.suelo))
```



```
summary.lm(aov.suelo)
```

```
##
## Call:
## aov(formula = prod$y.ton ~ prod$suelo)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
##    -8.5    -1.8     0.3     1.7     7.1
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
```

```
## (Intercept)          9.900      1.081    9.158 9.04e-10 ***
## prod$sueloarcilla    1.600      1.529    1.047 0.30456
## prod$suelolimo      4.400      1.529    2.878 0.00773 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 3.418 on 27 degrees of freedom
## Multiple R-squared:  0.2392, Adjusted R-squared:  0.1829
## F-statistic: 4.245 on 2 and 27 DF,  p-value: 0.02495
## se acepta la hipotesis alternativa ya que, al menos uno de los grupos
## es diferente a los demas
```