

INFERRING VEGETATION MEMORY FROM REMOTE SENSING DATA USING NOVEL CLIMATE RECONSTRUCTION PRODUCTS

M.Sc. Thesis Defense



UNIVERSITETET I BERGEN



Erik Kusch

erik@i-solution.de

Ecological and Environmental Change Research Group
University of Bergen

05/07/2019

- 1** Background
 - Motivation
 - Concept
- 2** Allocating and Preparing Data
 - Vegetation Data
 - Climate Data
 - Plant Functional Data
- 3** Delineating Vegetation Memory
- 4** Results
 - Coefficients of Vegetation Memory
 - Regional Aspects of Vegetation Memory
 - Functional Aspects to Vegetation Memory
- 5** Conclusion

Outline

1 Background

- Motivation
- Concept

2 Allocating and Preparing Data

- Vegetation Data
- Climate Data
- Plant Functional Data

3 Delineating Vegetation Memory

4 Results

- Coefficients of Vegetation Memory
- Regional Aspects of Vegetation Memory
- Functional Aspects to Vegetation Memory

5 Conclusion

What is this?

Vegetation Memory is the effect of **antecedent ecosystem/environmental anomalies** on **current vegetation performance**^[1].

Components of Memory:^[2]

- 1 *Intrinsic Memory* (e.g. antecedent vegetation characteristics)^[2,3]
- 2 *Extrinsic Memory* (antecedent climate characteristics)^[2-4]

Explaining Memory:

- 1 *Causal pathways* remain poorly understood^[5]
- 2 Expressions of *Plant Functional* as a possible solution

Can we distinguish between **intrinsic** and **extrinsic** memory effects?

What biological traits cause areas to exert **intrinsic** and **extrinsic** memory?

What is this?

Vegetation Memory is the effect of **antecedent ecosystem/environmental anomalies** on **current vegetation performance**^[1].

Components of Memory:^[2]

- 1 *Intrinsic Memory* (e.g. antecedent vegetation characteristics)^[2,3]
- 2 *Extrinsic Memory* (antecedent climate characteristics)^[2–4]

Explaining Memory:

- 1 *Causal pathways* remain poorly understood^[5]
- 2 Expressions of *Plant Functional* as a possible solution

Can we distinguish between **intrinsic** and **extrinsic** memory effects?

What biological traits cause areas to exert **intrinsic** and **extrinsic** memory?

What is this?

Vegetation Memory is the effect of **antecedent ecosystem/environmental anomalies** on **current vegetation performance**^[1].

Components of Memory:^[2]

- 1 *Intrinsic Memory* (e.g. antecedent vegetation characteristics)^[2,3]
- 2 *Extrinsic Memory* (antecedent climate characteristics)^[2–4]

Explaining Memory:

- 1 *Causal pathways* remain poorly understood^[5]
- 2 Expressions of *Plant Functional* as a possible solution

Can we **distinguish** between **intrinsic** and **extrinsic** memory effects?

What biological traits cause areas to exert **intrinsic** and **extrinsic** memory?

What is this?

Vegetation Memory is the effect of **antecedent ecosystem/environmental anomalies** on **current vegetation performance**^[1].

Components of Memory:^[2]

- 1 *Intrinsic Memory* (e.g. antecedent vegetation characteristics)^[2,3]
- 2 *Extrinsic Memory* (antecedent climate characteristics)^[2–4]

Explaining Memory:

- 1 *Causal pathways* remain poorly understood^[5]
- 2 Expressions of *Plant Functional* as a possible solution

Can we **distinguish** between **intrinsic** and **extrinsic** memory effects?

What biological traits cause areas to exert **intrinsic** and **extrinsic** memory?

What is this?

Vegetation Memory is the effect of **antecedent ecosystem/environmental anomalies** on **current vegetation performance**^[1].

Components of Memory:^[2]

- 1 *Intrinsic Memory* (e.g. antecedent vegetation characteristics)^[2,3]
- 2 *Extrinsic Memory* (antecedent climate characteristics)^[2–4]

Explaining Memory:


- 1 *Causal pathways* remain poorly understood^[5]
- 2 Expressions of *Plant Functional* as a possible solution

Can we **distinguish** between **intrinsic** and **extrinsic** memory effects?

What **biological traits** cause areas to exert **intrinsic** and **extrinsic memory**?

Why do this?


Vegetation Memory proposed to be a *proxy of engineering resilience in ecosystems* (*high memory ~ low resilience*)^[3].



How valid is this assumption?


Why do this?

Vegetation Memory proposed to be a *proxy of engineering resilience in ecosystems* (*high memory ~ low resilience*)^[3].




How valid is this assumption?

Intrinsic vs. Extrinsic Memory Components



→ Big emphasis on **dryland regions** due to demonstrated vegetation memory effects^[1,3,4,6], and the strong dependence of dryland vegetation on local water regimes^[5]

Intrinsic vs. Extrinsic Memory Components




→ Big emphasis on **dryland regions** due to demonstrated vegetation memory effects^[1,3,4,6], and the strong dependence of dryland vegetation on local water regimes^[5]

Study Region


The Iberian Region

Intrinsic Memory




De Keersmaecker et al. (2015). A model quantifying global vegetation resistance and resilience to short term climate anomalies and their relationship with vegetation cover. *Global Ecology and Biogeography*.


Intrinsic Memory



Seddon et al. (2016). Sensitivity of global terrestrial ecosystems to climate variability. *Nature*




Water Memory



Vicente-Serrano et al. (2013). Response of vegetation to drought time-scales across global land biomes. *Proceedings of the National Academy of Sciences*

Water Memory Length



Liu et al. (2018). Water memory effects and their impacts on global vegetation productivity and resilience. *Scientific Reports*

Additional Study Regions:

■ Caatinga, Brazil


■ Australia

■ Contiguous US

Study Region


The Iberian Region

Intrinsic Memory




De Keersmaecker et al. (2015). A model quantifying global vegetation resistance and resilience to short term climate anomalies and their relationship with vegetation cover. *Global Ecology and Biogeography*

Intrinsic Memory




Seddon et al. (2016). Sensitivity of global terrestrial ecosystems to climate variability. *Nature*

Computerized Bioclimatic Maps of the World: Ombrotypes of Europe and Mideast



Water Memory



Vicente-Serrano et al. (2013). Response of vegetation to drought time-scales across global land biomes. *Proceedings of the National Academy of Sciences*

Water Memory Length



Liu et al. (2018). Water memory effects and their impacts on global vegetation productivity and resilience. *Scientific Reports*

Additional Study Regions:

■ Caatinga, Brazil

■ Australia

■ Contiguous US

Outline

1 Background

- Motivation
- Concept

2 Allocating and Preparing Data

- Vegetation Data
- Climate Data
- Plant Functional Data

3 Delineating Vegetation Memory

4 Results

- Coefficients of Vegetation Memory
- Regional Aspects of Vegetation Memory
- Functional Aspects to Vegetation Memory

5 Conclusion

Normalised Difference Vegetation Index (NDVI)

We are using the **GIMMS NDVI3g** (v.1).

$$NDVI = \frac{\rho_{NIR} - \rho_{RED}}{\rho_{NIR} + \rho_{RED}} \quad (1)$$

■ Why:

- *Biological Relevance*: Proxy of biomass and vegetation cover
- *Comparability*: Has been used in other studies of vegetation memory

■ Core Characteristics:

- *Resolution*: $0.083^\circ \sim 9.27\text{km} \times 9.27\text{km}$
- *Availability*: 1982 - 2015

Normalised Difference Vegetation Index (NDVI)

We are using the **GIMMS NDVI3g** (v.1).

$$NDVI = \frac{\rho_{NIR} - \rho_{RED}}{\rho_{NIR} + \rho_{RED}} \quad (1)$$

■ Why:

- *Biological Relevance*: Proxy of biomass and vegetation cover
- *Comparability*: Has been used in other studies of vegetation memory

■ Core Characteristics:

- *Resolution*: $0.083^\circ \sim 9.27\text{km} \times 9.27\text{km}$
- *Availability*: 1982 - 2015

Normalised Difference Vegetation Index (NDVI)

We are using the **GIMMS NDVI3g** (v.1).

$$NDVI = \frac{\rho_{NIR} - \rho_{RED}}{\rho_{NIR} + \rho_{RED}} \quad (1)$$

■ Why:


- *Biological Relevance*: Proxy of biomass and vegetation cover
- *Comparability*: Has been used in other studies of vegetation memory

■ Core Characteristics:

- *Resolution*: $0.083^\circ \sim 9.27\text{km} \times 9.27\text{km}$
- *Availability*: 1982 - 2015

Monthly Composites

Mean NDVI 1982 – 2015



ERA5 & Variables I

We are using the **ERA5** Data Set^[7,8].

■ Why:

- Applicable globally
- Gap-less time series
- More sophisticated approach than previously utilised:
 - Worldclim - Superior Temporal Resolution (superior resolving of climate extremes)
 - CRU - Superior Spatial Resolution
- *Climatological Relevance*: Assessment of climate uncertainty possible

■ Core Characteristics:

- *Resolution*: $30\text{km} \times 30\text{km}$
- *Availability*: 1950 - Present

ERA5 & Variables I

We are using the **ERA5** Data Set^[7,8].

■ Why:

- Applicable globally
- Gap-less time series
- More sophisticated approach than previously utilised:
 - Worldclim - Superior Temporal Resolution (superior resolving of climate extremes)
 - CRU - Superior Spatial Resolution
- *Climatological Relevance*: Assessment of climate uncertainty possible

■ Core Characteristics:

- *Resolution*: $30\text{km} \times 30\text{km}$
- *Availability*: 1950 - Present

ERA5 & Variables I

We are using the **ERA5** Data Set^[7,8].

■ Why:

- Applicable globally
- Gap-less time series
- More sophisticated approach than previously utilised:
 - Worldclim - Superior Temporal Resolution (superior resolving of climate extremes)
 - CRU - Superior Spatial Resolution
- *Climatological Relevance*: Assessment of climate uncertainty possible

■ Core Characteristics:

- *Resolution*: $30\text{km} \times 30\text{km}$
- *Availability*: 1950 - Present

ERA5 & Variables II

We are using the following **climate variables**:

ERA5 & Variables II

We are using the following **climate variables**:

- **Soil Moisture** - Q_{soil}

- **Air Temperature** - T_{air}

ERA5 & Variables II

We are using the following **climate variables**:

■ **Soil Moisture** - Q_{soil}

- *Why:* Soil moisture effects on vegetation may indicate if/how the local ecosystem may deal with drought stress^[5,9,10].
- *How:* as different layers of depth
 - Q_{soil} - 0-7cm layer
 - Q_{soil2} - 7-28cm layer
 - Q_{soil3} - 28-100cm layer
 - Q_{soil4} - 100-255cm layer

■ **Air Temperature** - T_{air}

ERA5 & Variables II

We are using the following **climate variables**:


■ **Soil Moisture** - Q_{soil}

- *Why:* Soil moisture effects on vegetation may indicate if/how the local ecosystem may deal with drought stress^[5,9,10].
- *How:* as different layers of depth
 - Q_{soil} - 0-7cm layer
 - Q_{soil2} - 7-28cm layer
 - Q_{soil3} - 28-100cm layer
 - Q_{soil4} - 100-255cm layer

■ **Air Temperature** - T_{air}

- *Why:* Temperature drives plant physiology and drives levels of aridity^[3,9].
- *How:* As one single layer (2m above ground)

Kriging



Plant Functional Data

Plant Functional Traits (PFTs)

■ Why:

- *Biological Relevance*: Snapshots of functional reality
- *Comparability*: Have widely been used in functional ecology^[11,12]

■ Core Measures:

- *Vegetative Height*: Ability to pre-empt light
- *Leaf Nitrogen Content*: Measure of photosynthetic potential and nitrogen acquisition

Life History Traits (LHTs)

■ Why:

- *Biological Relevance*: Indices of plant behaviour through time
- *Comparability*: Capture much of natural life strategy variation^[13]

■ Core Measures:

- *Fast-Slow Continuum (FSC)*: Capture over 60% of the variation in plant life history strategies
 - FSC-1: Life History Speed
 - FSC-2: Reproductive Strategy/Output
- *Reactivity*: Instantaneous biological responses

Plant Functional Data

Plant Functional Traits (PFTs)

■ Why:

- *Biological Relevance*: Snapshots of functional reality
- *Comparability*: Have widely been used in functional ecology^[11,12]

■ Core Measures:

- *Vegetative Height*: Ability to pre-empt light
- *Leaf Nitrogen Content*: Measure of photosynthetic potential and nitrogen acquisition

Life History Traits (LHTs)

■ Why:

- *Biological Relevance*: Indices of plant behaviour through time
- *Comparability*: Capture much of natural life strategy variation^[13]

■ Core Measures:

- *Fast-Slow Continuum (FSC)*: Capture over 60% of the variation in plant life history strategies
 - FSC-1: Life History Speed
 - FSC-2: Reproductive Strategy/Output
- *Reactivity*: Instantaneous biological responses

Plant Functional Data

Plant Functional Traits (PFTs)

■ Why:

- *Biological Relevance*: Snapshots of functional reality
- *Comparability*: Have widely been used in functional ecology^[11,12]

■ Core Measures:

- *Vegetative Height*: Ability to pre-empt light
- *Leaf Nitrogen Content*: Measure of photosynthetic potential and nitrogen acquisition

Life History Traits (LHTs)

■ Why:

- *Biological Relevance*: Indices of plant behaviour through time
- *Comparability*: Capture much of natural life strategy variation^[13]

■ Core Measures:

- *Fast-Slow Continuum (FSC)*: Capture over 60% of the variation in plant life history strategies
 - FSC-1: Life History Speed
 - FSC-2: Reproductive Strategy/Output
- *Reactivity*: Instantaneous biological responses

Plant Functional Data

Plant Functional Traits (PFTs)

■ Why:

- *Biological Relevance*: Snapshots of functional reality
- *Comparability*: Have widely been used in functional ecology^[11,12]

■ Core Measures:

- *Vegetative Height*: Ability to pre-empt light
- *Leaf Nitrogen Content*: Measure of photosynthetic potential and nitrogen acquisition

Life History Traits (LHTs)

■ Why:

- *Biological Relevance*: Indices of plant behaviour through time
- *Comparability*: Capture much of natural life strategy variation^[13]

■ Core Measures:

- *Fast-Slow Continuum (FSC)*: Capture over 60% of the variation in plant life history strategies
 - FSC-1: Life History Speed
 - FSC-2: Reproductive Strategy/Output
- *Reactivity*: Instantaneous biological responses

Plant Functional Data

Plant Functional Traits (PFTs)

■ Why:

- *Biological Relevance*: Snapshots of functional reality
- *Comparability*: Have widely been used in functional ecology^[11,12]

■ Core Measures:

- *Vegetative Height*: Ability to pre-empt light
- *Leaf Nitrogen Content*: Measure of photosynthetic potential and nitrogen acquisition

Life History Traits (LHTs)

■ Why:

- *Biological Relevance*: Indices of plant behaviour through time
- *Comparability*: Capture much of natural life strategy variation^[13]

■ Core Measures:

- *Fast-Slow Continuum (FSC)*: Capture over 60% of the variation in plant life history strategies
 - FSC-1: Life History Speed
 - FSC-2: Reproductive Strategy/Output
- *Reactivity*: Instantaneous biological responses

Plant Functional Data

Plant Functional Traits (PFTs)

■ Why:

- *Biological Relevance*: Snapshots of functional reality
- *Comparability*: Have widely been used in functional ecology^[11,12]

■ Core Measures:

- *Vegetative Height*: Ability to pre-empt light
- *Leaf Nitrogen Content*: Measure of photosynthetic potential and nitrogen acquisition

Life History Traits (LHTs)

■ Why:

- *Biological Relevance*: Indices of plant behaviour through time
- *Comparability*: Capture much of natural life strategy variation^[13]

■ Core Measures:

- *Fast-Slow Continuum (FSC)*: Capture over 60% of the variation in plant life history strategies
 - FSC-1: Life History Speed
 - FSC-2: Reproductive Strategy/Output
- *Reactivity*: Instantaneous biological responses

Outline

1 Background

- Motivation
- Concept

2 Allocating and Preparing Data

- Vegetation Data
- Climate Data
- Plant Functional Data


3 Delineating Vegetation Memory

4 Results

- Coefficients of Vegetation Memory
- Regional Aspects of Vegetation Memory
- Functional Aspects to Vegetation Memory

5 Conclusion

Pixyel-Wise Model Building



PCA Regression

Model Formula: $NDVI_t = PC_1 + PC_2 + PC_3$

$NDVI_t$ NDVI anomaly at month t

PC_1 First principal component

PC_2 Second principal component

PC_3 Third principal component

	PC1	PC2	PC3	
$NDVI_{t-1}$	2.5	-1.14	1.95	
$Qsoil_{1,6}$	2.6	-0.73	-2.03	
$Tair$	-1.6	-3.00	-0.25	
Model Coefficients	1.77	-0.50	0.71	

Loadings

$$C_p = \sum_{i=1}^3 (L_{p;i} * RC_i)$$

C_p Coefficient of variable p

i Principal component counter

$L_{p;i}$ Loading of variable p on principal component i

RC_i Model coefficient for principal component i

	PC1	PC2	PC3	Coefficient
$NDVI_{t-1}$	4.41	0.57	1.39	6.4
$Qsoil_{1,6}$	4.57	0.37	-1.45	4.7
$Tair$	-2.79	1.50	-0.18	-1.5

Σ

Model Comparisons:

- Which variable exerts the greatest influence on vegetation anomalies?
- Which soil moisture layer is the most biologically influential?

PCA Regression

Model Formula: $NDVI_t = PC_1 + PC_2 + PC_3$

$NDVI_t$ NDVI anomaly at month t

PC_1 First principal component

PC_2 Second principal component

PC_3 Third principal component

	PC1	PC2	PC3	
$NDVI_{t-1}$	2.5	-1.14	1.95	
$Qsoil_{1:6}$	2.6	-0.73	-2.03	
$Tair$	-1.6	-3.00	-0.25	
Model Coefficients	1.77	-0.50	0.71	

Loadings

$$C_p = \sum_{i=1}^3 (L_{p;i} * RC_i)$$

C_p Coefficient of variable p

i Principal component counter

$L_{p;i}$ Loading of variable p on principal component i

RC_i Model coefficient for principal component i

	PC1	PC2	PC3	Coefficient
$NDVI_{t-1}$	4.41	0.57	1.39	6.4
$Qsoil_{1:6}$	4.57	0.37	-1.45	4.7
$Tair$	-2.79	1.50	-0.18	-1.5

Σ

Model Comparisons:

- Which variable exerts the greatest influence on vegetation anomalies?
- Which soil moisture layer is the most biologically influential?

Outline

1 Background

- Motivation
- Concept

2 Allocating and Preparing Data

- Vegetation Data
- Climate Data
- Plant Functional Data





3 Delineating Vegetation Memory

4 Results

- Coefficients of Vegetation Memory
- Regional Aspects of Vegetation Memory
- Functional Aspects to Vegetation Memory

5 Conclusion


Vegetation Memory Coefficients

SoilLag**NDVI t-1****Qsoil1****Tair**


→ Is *intrinsic* memory really *intrinsic*?

Vegetation Memory Coefficients


SoilLag




NDVI t-1



Qsoil1




Tair



→ Is **intrinsic** memory really *intrinsic*?


Distinguishing Intrinsic and Extrinsic Memory

Qsoil1 is the most informative of the soil moisture layers!




Identifying Underlying Extrinsic Patterns I

Uniform $NDVI[t - 1]$ effect across Australia **contrasts** with **other studies**.




Intrinsic Memory by Seddon et al.^[6]:




Identifying Underlying Extrinsic Patterns II

Uniform $NDVI[t - 1]$ effect across Australia **contrasts** with **other studies**.




Intrinsic Memory by Seddon et al.^[6]:




Identifying Underlying Extrinsic Patterns II

Uniform $NDVI[t - 1]$ effect across Australia **contrasts** with **other studies**.




Intrinsic Memory by Seddon et al.^[6]:




Previous $t - 1$ variation can
be understood through
extrinsic vegetation memory.

Vegetation Memory Adaptation



Relationship of $t - 1$ coefficient and extrinsic vegetation memory length
is not uniform within or between study regions.


Vegetation Memory Adaptation




Relationship of $t - 1$ coefficient and extrinsic vegetation memory length is not uniform within or between study regions.

Plant Function I

Linking **plant functional traits** and vegetation memory proved **non-conclusive** but life history traits showed **interesting patterns**:




Plant Function II



Some LHTs can be linked to some vegetation memory characteristics,

Plant Function II



Some LHTs can be linked to some vegetation memory characteristics,

Outline

1 Background

- Motivation
- Concept

2 Allocating and Preparing Data

- Vegetation Data
- Climate Data
- Plant Functional Data

3 Delineating Vegetation Memory

4 Results

- Coefficients of Vegetation Memory
- Regional Aspects of Vegetation Memory
- Functional Aspects to Vegetation Memory

5 Conclusion

Conclusion

Summary:

Extrinsic memory should not be neglected in favour of intrinsic memory.




Conclusion

Summary:

Extrinsic memory should not be neglected in favour of intrinsic memory.

- 1 Intrinsic vegetation memory as a proxy for engineering resilience may be an oversimplification.




Conclusion

Summary:

Extrinsic memory should not be neglected in favour of intrinsic memory.

- 1 Intrinsic vegetation memory as a proxy for engineering resilience may be an oversimplification.




Conclusion

Summary:

Extrinsic memory should not be neglected in favour of intrinsic memory.

- 1 Intrinsic vegetation memory as a proxy for engineering resilience may be an oversimplification.




Distinguishing intrinsic and extrinsic memory components remains challenging.

Conclusion

Summary:

- 1 Intrinsic vegetation memory as a proxy for engineering resilience may be an oversimplification.




Distinguishing intrinsic and extrinsic memory components remains challenging.

Conclusion

Summary:

- 1 Intrinsic vegetation memory as a proxy for engineering resilience may be an oversimplification.

Interactions of vegetation memory characteristics are region-dependant.




Conclusion

Summary:

- 1 Intrinsic vegetation memory as a proxy for engineering resilience may be an oversimplification.

Interactions of vegetation memory characteristics are region-dependant.

- 2 Vegetation memory processes differ greatly between regions.




Conclusion

Summary:

- 1 Intrinsic vegetation memory as a proxy for engineering resilience may be an oversimplification.
- 2 Vegetation memory processes differ greatly between regions.

Some measures of life history strategies are related to vegetation memory characteristics.




Conclusion

Summary:

- 1 Intrinsic vegetation memory as a proxy for engineering resilience may be an oversimplification.
- 2 Vegetation memory processes differ greatly between regions.

Some measures of life history strategies are related to vegetation memory characteristics.




Challenging to establish direct proxies of either intrinsic or extrinsic vegetation memory components.

Conclusion

Summary:

- 1 Intrinsic vegetation memory as a proxy for engineering resilience may be an oversimplification.
- 2 Vegetation memory processes differ greatly between regions.
- 3 The relationship of plant function and vegetation memory requires further research.



Challenging to establish direct proxies of either intrinsic or extrinsic vegetation memory components.

Conclusion

Summary:

- 1 Intrinsic vegetation memory as a proxy for engineering resilience may be an oversimplification.
- 2 Vegetation memory processes differ greatly between regions.
- 3 The relationship of plant function and vegetation memory requires further research.

Future Directions:

- 1 Identify additional relevant climate parameters (e.g. cloud cover and soil type)
- 2 Extend analyses beyond dryland regions
- 3 Analyse changes in vegetation memory over time
- 4 Use additional plant functional data (e.g. BIEN data base)
- 5 Implement advanced mapping methods for plant functional data

Conclusion

Summary:

- 1 Intrinsic vegetation memory as a proxy for engineering resilience may be an oversimplification.
- 2 Vegetation memory processes differ greatly between regions.
- 3 The relationship of plant function and vegetation memory requires further research.

Future Directions:

- 1 Identify additional relevant climate parameters (e.g. cloud cover and soil type)
- 2 Extend analyses beyond dryland regions
- 3 Analyse changes in vegetation memory over time
- 4 Use additional plant functional data (e.g. BIEN data base)
- 5 Implement advanced mapping methods for plant functional data

Conclusion

Summary:

- 1 Intrinsic vegetation memory as a proxy for engineering resilience may be an oversimplification.
- 2 Vegetation memory processes differ greatly between regions.
- 3 The relationship of plant function and vegetation memory requires further research.

Future Directions:

- 1 Identify additional relevant climate parameters (e.g. cloud cover and soil type)
- 2 Extend analyses beyond dryland regions
- 3 Analyse changes in vegetation memory over time
- 4 Use additional plant functional data (e.g. BIEN data base)
- 5 Implement advanced mapping methods for plant functional data

Conclusion

Summary:

- 1 Intrinsic vegetation memory as a proxy for engineering resilience may be an oversimplification.
- 2 Vegetation memory processes differ greatly between regions.
- 3 The relationship of plant function and vegetation memory requires further research.

Future Directions:

- 1 Identify additional relevant climate parameters (e.g. cloud cover and soil type)
- 2 Extend analyses beyond dryland regions
- 3 Analyse changes in vegetation memory over time
- 4 Use additional plant functional data (e.g. BIEN data base)
- 5 Implement advanced mapping methods for plant functional data

Conclusion

Summary:

- 1 Intrinsic vegetation memory as a proxy for engineering resilience may be an oversimplification.
- 2 Vegetation memory processes differ greatly between regions.
- 3 The relationship of plant function and vegetation memory requires further research.

Future Directions:

- 1 Identify additional relevant climate parameters (e.g. cloud cover and soil type)
- 2 Extend analyses beyond dryland regions
- 3 Analyse changes in vegetation memory over time
- 4 Use additional plant functional data (e.g. BIEN data base)
- 5 Implement advanced mapping methods for plant functional data

Conclusion

Summary:

- 1 Intrinsic vegetation memory as a proxy for engineering resilience may be an oversimplification.
- 2 Vegetation memory processes differ greatly between regions.
- 3 The relationship of plant function and vegetation memory requires further research.

Future Directions:

- 1 Identify additional relevant climate parameters (e.g. cloud cover and soil type)
- 2 Extend analyses beyond dryland regions
- 3 Analyse changes in vegetation memory over time
- 4 Use additional plant functional data (e.g. BIEN data base)
- 5 Implement advanced mapping methods for plant functional data

Sources

- [1] Liu, L., Zhang, Y., Wu, S., Li, S. & Qin, D. Water memory effects and their impacts on global vegetation productivity and resilience. *Scientific Reports* **8**, 1–9 (2018).
- [2] Ogle, K. *et al.* Quantifying ecological memory in plant and ecosystem processes. *Ecology Letters* **18**, 221–235 (2015).
- [3] De Keersmaecker, W. *et al.* A model quantifying global vegetation resistance and resilience to short-term climate anomalies and their relationship with vegetation cover. *Global Ecology and Biogeography* **24**, 539–548 (2015).
- [4] Vicente-Serrano, S. M. *et al.* Response of vegetation to drought time-scales across global land biomes. *Proceedings of the National Academy of Sciences* **110**, 52–57 (2013).
- [5] Smith, A. P. *et al.* Shifts in pore connectivity from precipitation versus groundwater rewetting increases soil carbon loss after drought. *Nature Communications* **8**, 1335 (2017).
- [6] Seddon, A. W. R., Macias-Fauria, M., Long, P. R., Benz, D. & Willis, K. J. Sensitivity of global terrestrial ecosystems to climate variability. *Nature* **531**, 229–232 (2016).
- [7] Albergel, C. *et al.* Skill and Global Trend Analysis of Soil Moisture from Reanalyses and Microwave Remote Sensing. *Journal of Hydrometeorology* **14**, 1259–1277 (2013).
- [8] Balsamo, G. *et al.* ERA-Interim/Land: a global land surface reanalysis data set. *Hydrology and Earth System Sciences* **19**, 389–407 (2015).
- [9] Rudgers, J. A. *et al.* Climate sensitivity functions and net primary production: A framework for incorporating climate mean and variability. *Ecology* **99**, 576–582 (2018). [0608246v3](#).
- [10] Papagiannopoulou, C. *et al.* Vegetation anomalies caused by antecedent precipitation in most of the world. *Environmental Research Letters* **12**, 074016 (2017).
- [11] Ordóñez, A. & Svenning, J.-C. Geographic patterns in functional diversity deficits are linked to glacial-interglacial climate stability and accessibility. *Global Ecology and Biogeography* **24**, 826–837 (2015).
- [12] Díaz, S. *et al.* The global spectrum of plant form and function. *Nature* **529**, 167–171 (2016).
- [13] Salguero-Gómez, R. *et al.* The COMPADRE Plant Matrix Database: an open online repository for plant demography. *Journal of Ecology* **103**, 202–218 (2015).