

# Supplementary information supporting the paper: Warnatzsch et al. (2020) Climate Change Impact on Aflatoxin Contamination Risk in Malawi's Maize Crops.

## Supplementary Tables

**Table 1:** Regional Climate Models (RCM) sources, the original calendar format, and the climatic variables used. All of the models other than CanRCM4\_r2 were accessed through The Earth System Grid Federation (ESGF) data index (ESGF, 2017). The CanRCM4\_r2 model was accessed through the Canadian Centre for Climate Modelling and Analysis website (CCCma, 2017).

RCM	Institution	Lateral Boundary Conditions	Climatic Variable	Original Calendar
<b>CCLM4-8-17_v1</b> (COSMO, 2017)	Climate Limited-area Modelling Community (CLMcom)	CNRM-CM5 r1i1p1	Tas, Pr	365-days
		HadGEM2-ES r1i1p1		360-days
		EC-EARTH r12i1p1		366-days
		MPI-ESM-LR r1i1p1		366-days
<b>HIRHAM5_v2</b> (Christensen et al, 2007)	Danmarks Meteorologiske Insitut (DMI)	EC-EARTH r3i1p1	Tas, Pr, Hurs	366-days
<b>RACMO22T_v1</b> (van Meijgaard et al, 2008)	Koninklijk Nederlands Meteorologisch Instituut (KNMI)	HadGEM2-ES r1i1p1	Tas, Pr, Hurs	360-days
		EC-EARTH r1i1p1		366-days
<b>RCA4_v1</b> (Samuelsson et al, 2015)	Sveriges Meteorologiska och Hydrologiska Institut (SMHI)	CanESM2 r1i1p1	Tas, Pr, Hurs	366-days
		CNRM-CM5 r1i1p1		366-days
		CSIRO-MK3-6-0 r1i1p1		365-days
		GFDL-ESM2M r1i1p1		365-days
		IPSL-CM5A-MR r1i1p1		365-days
		HadGEM2-ES r1i1p1		360-days
		EC-EARTH r12i1p1		366-days
		MIROC5 r1i1p1		365-days
		MPI-ESM-LR r1i1p1		366-days
		NORESML1-M r1i1p1		365-days
<b>REMO2009_v1</b> (Jacob et al, 2012)	Climate Service Centre Germany (CSC) and Max Planck Institut (MPI)	EC-EARTH r12i1p1	Tas, Pr	366-days
		MPI-ESM-LR r1i1p1		366-days
<b>CanRCM4_r2</b> (Scinocca et al, 2016)	Canadian Centre for Climate Modelling and Analysis (CCCma)	CanESM2 r1i1p1	Tas, Pr	365-days

**Table 2:** Observed data sources

<b>Dataset</b>	<b>Variable Used</b>	<b>Resolution</b>	<b>Time-Period Available</b>	<b>Data Source</b>
<b>Climate Research Unit (CRU) version 4.0</b> (Harris et al, 2014)	Tas, TasMin, TasMax and Pr	0.5° Monthly Land Only	1901-2015	Gridded Station Data
<b>University of Delaware (UDel) version 4.01</b> (Willmott & Matsuura, 2001)	Tas and Pr	0.5° Monthly Land Only	1901-2010	Gridded Station Data
<b>Global Precipitation Climatology Centre (GPCC) version 7</b> (Schneider et al, 2015)	Pr	1.0° Monthly	1901-2010	Satellite and Station Data

**Table 3:** List of data sources for the climate files used in AquaCrop. Note that all RCMs referred to in this table are listed in Table 1 and the observed data referred to in this table are from the sources listed in Table 2. Note also that a version of each of these climate files was created for the three regions of Malawi: Northern, Central and Southern. All climate data used in AquaCrop used a 365-day calendar averaged over the relevant time period.

File	Time Scale	RCP	Temperature	Reference Evapotranspiration	Precipitation Rate	CO <sub>2</sub> Concentration
1	1971-2000	N/A	Mean of observed monthly data for minimum and maximum temperature	Calculated using methodology described in Section 2	Observed monthly data for precipitation rates	AquaCrop Mauna Loa CO <sub>2</sub>
2	2020-2049	4.5	Projected ensemble mean daily minimum and maximum temperature		Projected ensemble mean precipitation rate	AquaCrop IPCC RCP 4.5
3		8.5				AquaCrop IPCC RCP 8.5
4	2040-2069	4.5				AquaCrop IPCC RCP 4.5
5		8.5				AquaCrop IPCC RCP 8.5

**Table 4:** List of data sources for the climate files used in AFLA-maize. Note that all hindcast and projected data referred to in this table are from the RCMs listed in Table 1 and the observed data referred to in this table are from the sources listed in Table 2. Note also that a version of each of these climate files was created for the three regions of Malawi: Northern, Central and Southern. All climate data used in AFLA-maize used a 366-day calendar with daily results for each date and year (i.e. 365/366 days (depending on leap year) x 30 years for each scenario).

File	Time Scale	RCP	Temperature	Relative Humidity	Precipitation Rate	Leaf Wetness
1	1971-2000	N/A	Mean of observed monthly data for minimum and maximum temperature	Hindcasted ensemble mean daily relative humidity	Observed monthly data for precipitation rates	Calculated using methodology described in main paper (Section 4.1)
2	2020-2049	4.5	Projected ensemble mean daily minimum and maximum temperature	Projected ensemble mean daily relative humidity	Projected ensemble mean precipitation rate	
3		8.5				
4	2040-2069	4.5				
5		8.5				

**Table 5:** Calibration of individual maize cultivars (Sutcliffe, 2014) compared with AquaCrop default values.

Parameter (in degree days)	AquaCrop Default	Fast-Development	Slow-Development
Emergence	80	60	98
Max Canopy	705	684	720
Senescence	1400	1008	1715
Maturity	1700	1332	2267
Length building up to Harvest Index (HI)	750	636	1140
Duration of flowering	180	132	220
Degree-Days for Flowering	880	636	1078
Time to maximum rooting depth	1409	1120	1722

**Table 6:** Common calibration of both maize varieties compared with AquaCrop default values.

Parameter		AquaCrop Default	Calibrated Maize	Reference
<b>Initial Canopy Cover</b>	Initial Plant Density (plants/ha)	75 000	47 000	(Wiyo et al, 1999)
<b>Canopy Development</b>	Maximum Canopy Cover (%)	96	75	(Fiwa, 2015)
<b>Root Deepening</b>	Maximum effective rooting depth (m)	2.3	0.6	
<b>Harvest Index</b>	Hio (%)	48	40	
<b>Air Temperature Stresses</b>	Base temperature (°C)	8	13	(Benson et al, 2016)
	Upper temperature (°C)	30	32	
<b>Soil Salinity Stress - Salt tolerance</b>	ECe lower threshold (dS/m)	2	4	adapted from (Benson et al, 2016)
	ECe upper threshold (dS/m)	10	9	
<b>Crop response to soil fertility stress</b>	Crop response to soil fertility stress	Not Considered	Considered	(Fiwa, 2015)
	Biomass Production (%)		69	
	Canopy Decline in Season		medium	
	Reduction of Canopy Expansion (%)		7	
	Average decline in Canopy Cover (%/day)		0.1	
	Reduction in water productivity (%)		47	

**Table 7:** Development Dates for the fast-development maize variety. The date of emergence and harvest dates are listed for each climatic scenario and time period, as well as the number of days after planting (DAP) those dates occur. The difference ( $\Delta$ ) between the development of the crop under historic climatic and future climatic conditions is also listed.

Sowing Date	Region	Climate Scenario	Emergence			Harvest		
			Date	DAP	$\Delta$	Date	DAP	$\Delta$
15 Nov.	Northern	1971-2000	21 Nov.	6	N/A	09 Apr.	145	N/A
		2020-2049 RCP 4.5	20 Nov.	5	-1	20 Mar.	126	-19
		2020-2049 RCP 8.5	20 Nov.	5	-1	19 Mar.	125	-20
		2040-2069 RCP 4.5	20 Nov.	5	-1	20 Mar.	126	-19
		2040-2069 RCP 8.5	20 Nov.	5	-1	19 Mar.	125	-20
	Central	1971-2000	20 Nov.	5	N/A	22 Mar.	127	N/A
		2020-2049 RCP 4.5	20 Nov.	5	0	07 Mar.	113	-14
		2020-2049 RCP 8.5	20 Nov.	5	0	05 Mar.	111	-16
		2040-2069 RCP 4.5	20 Nov.	5	0	07 Mar.	113	-14
		2040-2069 RCP 8.5	20 Nov.	5	0	05 Mar.	111	-16
	Southern	1971-2000	20 Nov.	5	N/A	04 Mar.	109	N/A
		2020-2049 RCP 4.5	20 Nov.	5	0	22 Feb.	99	-10
		2020-2049 RCP 8.5	20 Nov.	5	0	21 Feb.	98	-11
		2040-2069 RCP 4.5	20 Nov.	5	0	22 Feb.	99	-10
		2040-2069 RCP 8.5	20 Nov.	5	0	21 Feb.	98	-11
10 Dec.	Northern	1971-2000	17 Dec.	7	N/A	12 May	153	N/A
		2020-2049 RCP 4.5	16 Dec.	6	-1	19 Apr.	131	-22
		2020-2049 RCP 8.5	16 Dec.	6	-1	16 Apr.	128	-25
		2040-2069 RCP 4.5	16 Dec.	6	-1	19 Apr.	131	-22
		2040-2069 RCP 8.5	16 Dec.	6	-1	16 Apr.	128	-25
	Central	1971-2000	16 Dec.	6	N/A	22 Apr.	133	N/A
		2020-2049 RCP 4.5	15 Dec.	5	-1	05 Apr.	117	-16
		2020-2049 RCP 8.5	15 Dec.	5	-1	03 Apr.	115	-18
		2040-2069 RCP 4.5	15 Dec.	5	-1	05 Apr.	117	-16

		2040-2069 RCP 8.5	15 Dec.	5	-1	03 Apr.	115	-18
	Southern	1971-2000	15 Dec.	5	N/A	02 Apr.	113	N/A
		2020-2049 RCP 4.5	15 Dec.	5	0	22 Mar.	103	-10
		2020-2049 RCP 8.5	15 Dec.	5	0	20 Mar.	101	-12
		2040-2069 RCP 4.5	15 Dec.	5	0	22 Mar.	103	-10
		2040-2069 RCP 8.5	15 Dec.	5	0	20 Mar.	101	-12
<b>30 Dec.</b>	Northern	1971-2000	06 Jan.	7	N/A	14 Jun.	166	N/A
		2020-2049 RCP 4.5	05 Jan.	6	-1	13 May	135	-31
		2020-2049 RCP 8.5	05 Jan.	6	-1	10 May	132	-34
		2040-2069 RCP 4.5	05 Jan.	6	-1	13 May	135	-31
		2040-2069 RCP 8.5	05 Jan.	6	-1	10 May	132	-34
	Central	1971-2000	05 Jan.	6	N/A	20 May	141	N/A
		2020-2049 RCP 4.5	05 Jan.	6	0	29 Apr.	121	-20
		2020-2049 RCP 8.5	05 Jan.	6	0	27 Apr.	119	-22
		2040-2069 RCP 4.5	05 Jan.	6	0	29 Apr.	121	-20
		2040-2069 RCP 8.5	05 Jan.	6	0	27 Apr.	119	-22
	Southern	1971-2000	04 Jan.	5	N/A	28 Apr.	119	N/A
		2020-2049 RCP 4.5	04 Jan.	5	0	14 Apr.	106	-13
		2020-2049 RCP 8.5	04 Jan.	5	0	12 Apr.	104	-15
		2040-2069 RCP 4.5	04 Jan.	5	0	14 Apr.	106	-13
		2040-2069 RCP 8.5	04 Jan.	5	0	12 Apr.	104	-15

**Table 8:** Development Dates for the slow-development maize variety. The date of emergence and harvest dates are listed for each climatic scenario and time period, as well as the number of days after planting (DAP) those dates occur. The difference ( $\Delta$ ) between the development of the crop under historic climatic and future climatic conditions is also listed.

Sowing Date	Region	Climate Scenario	Emergence			Harvest		
			Date	DAP	$\Delta$	Date	DAP	$\Delta$
15 Nov.	Northern	1971-2000	25 Nov.	10	N/A	28 Aug.	286	N/A
		2020-2049 RCP 4.5	24 Nov.	9	-1	29 Jun.	227	-59
		2020-2049 RCP 8.5	24 Nov.	9	-1	23 Jun.	221	-65
		2040-2069 RCP 4.5	24 Nov.	9	-1	29 Jun.	227	-59
		2040-2069 RCP 8.5	24 Nov.	9	-1	23 Jun.	221	-65
	Central	1971-2000	24 Nov.	9	N/A	18 Jul.	245	N/A
		2020-2049 RCP 4.5	23 Nov.	8	-1	28 May	195	-50
		2020-2049 RCP 8.5	23 Nov.	8	-1	23 May	190	-55
		2040-2069 RCP 4.5	23 Nov.	8	-1	28 May	195	-50
		2040-2069 RCP 8.5	23 Nov.	8	-1	23 May	190	-55
	Southern	1971-2000	23 Nov.	8	N/A	27 May	193	N/A
		2020-2049 RCP 4.5	22 Nov.	7	-1	01 May	168	-25
		2020-2049 RCP 8.5	22 Nov.	7	-1	29 Apr.	166	-27
		2040-2069 RCP 4.5	22 Nov.	7	-1	01 May	168	-25
		2040-2069 RCP 8.5	22 Nov.	7	-1	29 Apr.	166	-27
10 Dec.	Northern	1971-2000	21 Dec.	11	N/A	26 Sep.	290	N/A
		2020-2049 RCP 4.5	20 Dec.	10	-1	14 Aug.	248	-42
		2020-2049 RCP 8.5	20 Dec.	10	-1	09 Aug.	243	-47
		2040-2069 RCP 4.5	20 Dec.	10	-1	14 Aug.	248	-42
		2040-2069 RCP 8.5	20 Dec.	10	-1	09 Aug.	243	-47
	Central	1971-2000	20 Dec.	10	N/A	01 Sep.	265	N/A
		2020-2049 RCP 4.5	19 Dec.	9	-1	17 Jul.	220	-45
		2020-2049 RCP 8.5	18 Dec.	8	-2	10 Jul.	213	-52
		2040-2069 RCP 4.5	19 Dec.	9	-1	18 Jul.	221	-44


		2040-2069 RCP 8.5	18 Dec.	8	-2	10 Jul.	213	-52
	Southern	1971-2000	18 Dec.	8	N/A	21 Jul.	223	N/A
		2020-2049 RCP 4.5	18 Dec.	8	0	10 Jun.	183	-40
		2020-2049 RCP 8.5	18 Dec.	8	0	06 Jun.	179	-44
		2040-2069 RCP 4.5	18 Dec.	8	0	10 Jun.	183	-40
		2040-2069 RCP 8.5	18 Dec.	8	0	06 Jun.	179	-44
<b>30 Dec.</b>	Northern	1971-2000	10 Jan.	11	N/A	14 Oct.	288	N/A
		2020-2049 RCP 4.5	09 Jan.	10	-1	08 Sep.	253	-35
		2020-2049 RCP 8.5	09 Jan.	10	-1	03 Sep.	248	-40
		2040-2069 RCP 4.5	09 Jan.	10	-1	08 Sep.	253	-35
		2040-2069 RCP 8.5	09 Jan.	10	-1	03 Sep.	248	-40
	Central	1971-2000	09 Jan.	10	N/A	23 Sep.	267	N/A
		2020-2049 RCP 4.5	08 Jan.	9	-1	20 Aug.	234	-33
		2020-2049 RCP 8.5	08 Jan.	9	-1	14 Aug.	228	-39
		2040-2069 RCP 4.5	08 Jan.	9	-1	20 Aug.	234	-33
		2040-2069 RCP 8.5	08 Jan.	9	-1	14 Aug.	228	-39
	Southern	1971-2000	08 Jan.	9	N/A	26 Aug.	239	N/A
		2020-2049 RCP 4.5	07 Jan.	8	-1	21 Jul.	204	-35
		2020-2049 RCP 8.5	07 Jan.	8	-1	14 Jul.	197	-42
		2040-2069 RCP 4.5	07 Jan.	8	-1	21 Jul.	204	-35
		2040-2069 RCP 8.5	07 Jan.	8	-1	14 Jul.	197	-42




**Table 9:** AFI for slow- and fast-developing maize cultivars in each region of Malawi under differing climatic conditions and sowing dates. The bold red text indicates AFI levels which are consistent with exceedances of the EU threshold for AFB<sub>1</sub> contamination in food.

	Sowing Date	Region	Baseline (1971-2000)	2035 RCP 4.5	2035 RCP 8.5	2055 RCP 4.5	2055 RCP 8.5
Slow-Development Cultivar	15 Nov.	North	5.62	5.90	28.06	0.44	32.02
	10 Dec.		21.08	38.02	90.07	45.40	54.15
	30 Dec.		35.75	60.19	<b>101.57</b>	<b>169.69</b>	83.42
	15 Nov.	Central	31.46	60.43	<b>116.33</b>	47.80	93.21
	10 Dec.		<b>97.63</b>	<b>103.05</b>	<b>146.55</b>	94.04	<b>129.55</b>
	30 Dec.		81.98	<b>151.65</b>	<b>165.21</b>	<b>167.04</b>	<b>215.79</b>
	15 Nov.	South	<b>178.01</b>	<b>235.61</b>	<b>247.04</b>	<b>210.21</b>	<b>232.87</b>
	10 Dec.		<b>179.40</b>	<b>251.67</b>	<b>244.28</b>	<b>234.12</b>	<b>197.55</b>
	30 Dec.		<b>147.21</b>	<b>238.50</b>	<b>247.30</b>	<b>265.58</b>	<b>282.81</b>
Fast-Development Cultivar	15 Nov.	North	4.14	3.43	21.85	0.00	45.61
	10 Dec.		17.20	30.68	75.75	30.32	44.53
	30 Dec.		31.57	64.15	<b>96.19</b>	<b>139.86</b>	81.83
	15 Nov.	Central	22.27	55.07	<b>124.70</b>	39.69	<b>110.78</b>
	10 Dec.		84.62	<b>111.70</b>	<b>127.55</b>	74.40	<b>110.26</b>
	30 Dec.		89.86	<b>156.09</b>	<b>168.65</b>	<b>154.72</b>	<b>209.15</b>
	15 Nov.	South	<b>178.60</b>	<b>231.56</b>	<b>240.76</b>	<b>215.12</b>	<b>236.56</b>
	10 Dec.		<b>185.92</b>	<b>251.83</b>	<b>243.54</b>	<b>235.35</b>	<b>180.24</b>
	30 Dec.		<b>156.20</b>	<b>238.02</b>	<b>250.55</b>	<b>268.90</b>	<b>247.19</b>

## Supplementary Figures



**Figure 1:** The top panel shows daily temperatures, the height of the bar indicates the average mean daily temperature during the relevant month over that 30 year period, with the whisker bars indicating the average daily minimum and maximum temperatures. The second, third and fourth panels down show average daily precipitation, average daily reference evapotranspiration and average daily relative humidity respectively. All four panels show this data for the three regions of Malawi in the 1971-2000 period, with the darkest blue representing the Northern Region, the middle blue representing the Central Region and the lightest blue representing the Southern Region.



**Figure 2:** AFI for slow (A) and fast (B) development maize cultivars in each region of Malawi under different climatic conditions and planting dates

## Methodology for Calculating Reference Evapotranspiration for Central Malawi

To calculate evapotranspiration for Central Malawi, the FAO Penman Monteith (FPM) model was applied (Allen et al, 1998a).

$$\text{Equation 1} \quad ETo = \frac{0.408\Delta(R_n - G) + \gamma \frac{900}{T + 273} u_2 (e_s - e_a)}{\Delta + \gamma(1 + 0.34u_2)}$$

Where:

- ETo is the reference evapotranspiration (mm day<sup>-1</sup>)
- R<sub>n</sub> is the net radiation at the crop surface (MJ m<sup>2</sup> day<sup>-1</sup>),
- G is the soil heat flux density (MJ m<sup>-2</sup> day<sup>-1</sup>)
- T is the mean daily air temperature (°C)
- u<sub>2</sub> is wind speed at 2 m height (m s<sup>-1</sup>)
- e<sub>s</sub> is the saturation vapour pressure (kPa)
- e<sub>a</sub> is the actual vapour pressure (kPa), see Equation 10
- e<sub>s</sub> – e<sub>a</sub> is the saturation vapour pressure deficit (kPa)
- Δ is the slope vapour pressure curve (kPa °C<sup>-1</sup>)
- γ is the psychrometric constant (kPa °C<sup>-1</sup>)

It is not possible to get data for all of the above variables for Central Malawi, either from observed data of the past, or from climate models used to hindcast the past or forecast future climates. Therefore, temperature-based calculation methods were applied for climatic variables with no primary data available (Allen et al, 1998b). This methodology has been tested for Malawi by Wang et al. (2011), and for South Malawi by Ngongondo et al. (2012) and deemed to be appropriate for use.

### 2.1. Net Radiation at the Crop Surface

R<sub>n</sub> is the net radiation at the crop surface (MJ m<sup>2</sup> day<sup>-1</sup>) and can be calculated as follows:

$$\text{Equation 2} \quad R_n = R_{ns} - R_{nl}$$

Where:

- R<sub>ns</sub> is the net incoming shortwave radiation (MJm<sup>-2</sup> day<sup>-1</sup>) and can be calculated as follows:

$$\text{Equation 3} \quad R_{ns} = (1 - \alpha) R_s$$

Where:

- α is the albedo or canopy reflection coefficient, which is 0.23 for the hypothetical grass reference crop
- R<sub>s</sub> is the fraction of the solar radiation not reflected from the surface (MJm<sup>-2</sup> day<sup>-1</sup>) and can be calculated as follows:

$$\text{Equation 4} \quad R_s = k_{RS} \sqrt{T_{max} - T_{min}} R_a$$

Where:

- K<sub>RS</sub> is adjustment coefficient. For inland regions not influenced by large bodies of water, K<sub>RS</sub> = 0.16 ; for coastal regions, or regions where the air mass is influenced by a large nearby water body, K<sub>RS</sub> = 0.19. Since Central Malawi is highly influenced by the presence of a large water body (Lake Malawi). K<sub>RS</sub> is considered to be 0.19 in this study.
- T<sub>max</sub> is the maximum air temperature (°C)
- T<sub>min</sub> is the minimum air temperature (°C)

- $R_a$  is extra-terrestrial radiation ( $\text{MJm}^{-2} \text{ day}^{-1}$ ) and can be calculated as follows:

$$\text{Equation 5} \quad R_a = \frac{24(60)}{\pi} G_{sc} d_r [\omega_s \sin(\varphi) \sin(\delta) + \cos(\varphi) \cos(\delta) \sin(\omega_s)]$$

Where:

- $G_{sc}$  is the solar constant =  $0.0820 \text{ MJm}^{-2}\text{min}^{-1}$
- $d_r$  is the inverse relative since earth-Sun (rad) which can be calculated as follows:

$$\text{Equation 6} \quad d_r = 1 + 0.033 \cos\left(\frac{2\pi}{365} J\right)$$

Where:

- $J$  is the number of days in the year between 1 (1 January) and 365 or 266 (31 December).  $J$  at the middle of each month =  $30.4M - 15$  where  $M$  is the month number
- $\omega_s$  is the sunset hour angle (rad) which can be calculated as follows:

$$\text{Equation 7} \quad \omega_s = \arccos[-\tan(\varphi) \tan(\delta)]$$

Where:

- $\varphi$  is the latitude (rad)
- $\delta$  is the solar declination (rad) which can be calculated as follows:

$$\text{Equation 8} \quad \delta = 1 + 0.409 \sin\left(\frac{2\pi}{365} J - 1.39\right)$$

- $R_{nl}$  is in the net outgoing longwave radiation ( $\text{MJm}^{-2} \text{ day}^{-1}$ ) and can be calculated as follows:

$$\text{Equation 9} \quad R_{nl} = \sigma \left[ \frac{T_{\max K^4} + T_{\min K^4}}{2} \right] (0.34 - 0.14\sqrt{e_a}) \left( 1.35 \frac{R_s}{R_{so}} - 0.35 \right)$$

Where:

- $\sigma$  is the Stefan-Boltzmann constant [ $4.903 \times 10^{-9} \text{ MJ K}^{-4} \text{ m}^{-2} \text{ day}^{-1}$ ]
- $T_{\max, K}$  is the maximum absolute temperature during the 24-hour period [ $K = ^\circ\text{C} + 273.16$ ],
- $T_{\min, K}$  minimum absolute temperature during the 24-hour period [ $K = ^\circ\text{C} + 273.16$ ],
- $e_a$  actual vapour pressure [kPa], which can be calculated as follows:

$$\text{Equation 10} \quad e_a = e^o(T_{dew}) = 0.6108 \exp\left(\frac{17.27 T_{dew}}{T_{dew} + 237.3}\right)$$

Where:

- $T_{dew}$  is the dew point temperature.  $T_{dew}$  is near the minimum temperature ( $T_{\min}$ ) when the relative humidity is nearly 100%. In semi-arid regions,  $T_{dew}$  is estimated by subtracting  $2^\circ\text{C}$  from  $T_{\min}$ . As Central Malawi's humidity is typically under 90%, the  $T_{dew}$  can be calculated as follows:

$$\text{Equation 11} \quad T_{dew} = T_{\min} - 2$$

- $R_s$  is the solar radiation [ $\text{MJ m}^{-2} \text{ day}^{-1}$ ], see Equation 4.

- $R_{so}$  is the clear-sky solar radiation [ $\text{MJ m}^{-2} \text{ day}^{-1}$ ], which can be calculated as follows:

$$\text{Equation 12 } R_{so} = (0.75 + 0.00002(h))R_a$$

Where:

- $h$  is the elevation above sea level (m)
- $R_a$  is extra-terrestrial radiation, ( $\text{MJm}^{-2} \text{ day}^{-1}$ ), see Equation 5.

## 2.2. Soil Heat Flux Density

$G$  is the soil heat flux density ( $\text{MJ m}^{-2} \text{ day}^{-1}$ )

- For daily assessment,  $G$  is assumed to be zero (0) as the soil heat flux is relatively small

$$\text{Equation 13: } G_{day} = 0$$

- For monthly assessments,

$$\text{Equation 14: } G = 0.07 (T_{month,i+1} - T_{month,i-1})$$

Where:

- $T_{mon, i-1}$  is the mean air temperature of the previous month ( $^{\circ}\text{C}$ )
- $T_{mon, i+1}$  is the mean air temperature of the next month ( $^{\circ}\text{C}$ )

## 2.3. Mean Temperature

$T$  is the mean daily air temperature ( $^{\circ}\text{C}$ ), which can be calculated as follows:

$$\text{Equation 15: } T_{mean} = \frac{T_{min} + T_{max}}{2}$$

Where:

- $T_{max}$  is the maximum air temperature ( $^{\circ}\text{C}$ )
- $T_{min}$  is the minimum air temperature ( $^{\circ}\text{C}$ )

## 2.4. Wind Speed at 2m height

$u_2$  is wind speed at 2 m height ( $\text{m s}^{-1}$ ). We can use a default value of  $172 \text{ km day}^{-1}$  which is the average value over different weather stations around the globe. This was recommended by Allen et al. (1998). To convert to the correct units for the equation above ( $\text{m s}^{-1}$ ) we can do the following:

$$\text{Equation 16: } \frac{172 \text{ km}}{\text{day}} \times \frac{\text{day}}{24 \text{ hours}} \times \frac{\text{hour}}{60 \text{ minutes}} \times \frac{\text{minute}}{60 \text{ seconds}} \times \frac{1000 \text{ meters}}{\text{km}} = \frac{172,000 \text{ meters}}{86,400 \text{ seconds}}$$

## 2.5. Vapour Pressure

To calculate ETO, various vapour pressure variables are required, including the saturation vapour pressure ( $e_s$ ), the actual vapour pressure ( $e_a$ ) and the slope vapour pressure curve ( $\Delta$ ).

- $e_s$  is the saturation vapour pressure (kPa), it can be calculated as follows:

$$\text{Equation 17: } e_s = \frac{e^0(T_{max}) + e^0(T_{min})}{2}$$

Where:

- $e^0(T_{max})$  is the vapour pressure at maximum temperature, and can be calculated as follows:

$$\text{Equation 18: } e^0(T_{max}) = 0.6108 \exp\left(\frac{17.27T_{max}}{T_{max} + 237.3}\right)$$

Where:

- $T_{\max}$  is the maximum air temperature ( $^{\circ}\text{C}$ )
- $e^0(T_{\min})$  is the vapour pressure at minimum temperature, and can be calculated as follows:

$$\text{Equation 19: } e^0(T_{\min}) = 0.6108 \exp\left(\frac{17.27T_{\min}}{T_{\min} + 237.3}\right)$$

Where:

- $T_{\min}$  is the minimum air temperature ( $^{\circ}\text{C}$ )
- $e_a$  is the actual vapour pressure (kPa), see Equation 10
- $\Delta$  is the slope vapour pressure curve ( $\text{kPa}^{\circ}\text{C}^{-1}$ )

$$\text{Equation 20: } \Delta = \frac{4098 \left[ 0.6108 \exp\left(\frac{17.27T}{T+237.3}\right) \right]}{(T+237.3)^2}$$

Where:

- $T$  is the mean air temperature ( $^{\circ}\text{C}$ ), see Equation 15
- $\exp[\dots]$  2.7183 (base of natural logarithm) raised to the power [...]

## 2.6. Psychrometric Constant

$\gamma$  is the psychrometric constant ( $\text{kPa } ^{\circ}\text{C}^{-1}$ ), it can be calculated as follows:

$$\text{Equation 21: } \gamma = \frac{c_p P}{\varepsilon \lambda}$$

Where:

- $C_p$  is the specific heat at a constant pressure,  $C_p = 1.013 \times 10^{-3} \text{ MJ kg}^{-1} ^{\circ}\text{C}^{-1}$
- $P$  is atmospheric pressure (kPa), which can be calculated as follows:

$$\text{Equation 22: } P = 101.325(293 - 0.0065(h))^{5.25588}$$

Where:

- $h$  is the altitude above sea level in meters (m)
  - For Central Malawi, the average altitude above sea level ( $h$ ) is 948.1944444m (determined using data from JISAO (2014))
- $\varepsilon$  is the ratio molecular weight of water vapour / dry air,  $\varepsilon = 0.622$
- $\lambda$  is the latent heat of vaporization,  $\lambda = 2.45 \text{ MJ kg}^{-1}$

Allen, R. G., Pereira, L. S., Raes, D. & Smith, M. (1998a) FAO Penman-Monteith Equation, *Crop evapotranspiration - Guidelines for computing crop water requirements*. FAO Irrigation and Drainage Paper 56.

Allen, R. G., Pereira, L. S., Raes, D. & Smith, M. (1998b) *Meteorological Data, Crop evapotranspiration - Guidelines for computing crop water requirements*. FAO Irrigation and Drainage Paper 56. Rome: FAO.

Benson, T., Mabiso, A. & Nankhuni, F. (2016) *Detailed crop suitability maps and an agricultural zonation scheme for Malawi: Spatial information for agricultural planning purposes*. East Lansing, MI: (IFPRI), M. S. U. a. I. E. P. R. I.

CCCma (2017) *Canadian Regional Climate Model Output*, 2017. Available online: <http://climate-modelling.canada.ca/climatemodeldata/canrcm/CanRCM4/index.shtml> [Accessed.

Christensen, O. B., Drews, M., Christensen, J. H., Dethloff, K., Katelsen, K., Hebestadt, I. & Rinke, A. (2007) *Technical report 06-17. The HIRHAM Regional Climate Model*

Version 5 (β). Copenhagen: Danish Climate Centre, D.

COSMO (2017) *Core Documentation of the COSMO-model*, 2017. Available online: <http://www.cosmo-model.org/content/model/documentation/core/default.htm#p1> [Accessed.

ESGF (2017) *ESGF@LiU/CORDEX*, 2017. Available online: <https://esg-dn1.nsc.liu.se/projects/cordex/> [Accessed].

Fiwa, L. (2015) *Improving rainfed cereal production and water productivity in Malawi*. PhD KU Leuven. Available online: [height:14.25pt">](#) [Accessed. Available online: [<td nowrap="" style="width:82.9pt](#) [Accessed. Available online: [height:14.25pt" valign="bottom" width="111">](#) [Accessed. Available online:

[https://lirias.kuleuven.be/bitstream/123456789/493765/1/Lameck\\_FIWA.pdf](https://lirias.kuleuven.be/bitstream/123456789/493765/1/Lameck_FIWA.pdf). [Accessed.

Harris, I., Jones, P. D., Osborn, T. J. & Lister, D. H. (2014) Updated high-resolution grids of monthly climatic observations - the CRU TS3.10 Dataset. *International Journal of Climatology*, 34(3), 623-642.

Jacob, D., Elizalde, A., Haensler, A., Hagemann, S., Kumar, P., Podzun, R., Rechid, D., Remedio, A. R., Saeed, F., Sieck, K., Teichmann, C. & Wilhelm, C. (2012) Assessing the

Regional Climate Downscaling Experiment (CORDEX) Regions. *Atmosphere*, 3(1), 181-199.

JISAO (2014) *Elevation data in netCDF. 0.25-degree latitude-longitude resolution elevation (TBASE)*. (05 December 2017). Available online: [http://research.jisao.washington.edu/data\\_sets/elevation/](http://research.jisao.washington.edu/data_sets/elevation/).

Ngongondo, C., Xu, C.-Y., Tallaksen, L. M. & Alemaw, B. (2012) Evolution of the FAO Penman-Monthith, Priestley-Taylor and Hargreaves models for estimating reference evapotranspiration in southern Malawi. *Hydrology Research*, 44(4), 706-722.

Samuelsson, P., Gollvik, S., Jansson, C., Kupiainen, M., Kourzeneva, E. & Jan van de Berg, W. (2015) *The surface processes of the Rossby Centre*

regional atmospheric climate model (RCA4). Norrköping, Sweden: SMHI.

Schneider, U., Becker, A., Finger, P., Meyer-Christoffer, A., Rudolf, B. & Ziese, M. (2015) GPCP Full Data Reanalysis Version 7.0 at 1.0 °: Monthly Land-Surface Precipitation from Rain-Gauges built on GTS-based and Historic Data.

Scinocca, J. F., Kharin, V. V., Jiao, Y., Qian, M. W., Lazare, M., Solheim, L., Flato, G. M., Biner, S., Desgagne, M. & Dugas, B. (2016) Coordinated Global and Regional Climate Modeling. *Journal of Climate*, 29(1), 17-35.

Sutcliffe, C. A. J. (2014) *Adoption of improved maize cultivars for climate vulnerability reduction in Malawi*. PhD University of Leeds. Available online:  
[http://etheses.whiterose.ac.uk/7794/1/Sutcliffe\\_CAJ\\_2014.pdf](http://etheses.whiterose.ac.uk/7794/1/Sutcliffe_CAJ_2014.pdf) [Accessed].

van Meijgaard, E., van Ulft, L. H., van de Berg, W. J., Bosveld, F. C., van den Hurk, B. J. J. M., Lenderink, G. & Siebesma, A. P. (2008) *Technical report ; TR - 302. The KNMI regional atmospheric climate model RACMO version 2.1*. De Bilt: KNMI.



Wang, Y.-M., Namaona, W., Gladden, L. A., Traore, S. & Deng, L.-T. (2011) Comparative study on estimating reference evapotranspiration under limited climate data condition in Malawi.

*International Journal of the Physical Sciences*, 6(9), 2239-2248.

Willmott, C. J. & Matsuura, K. (2001) *Terrestrial Air Temperature and Precipitation: Monthly and Annual Time Series (1950-1999)*, 2001. Available online:

[http://climate.geog.udel.edu/~climate/html\\_pages/README.ghcn\\_ts2.html](http://climate.geog.udel.edu/~climate/html_pages/README.ghcn_ts2.html). [Accessed.

Wiyo, K. A., Kasomekera, Z. M. & Feyen, J. (1999) Variability in ridge and furrow size and shape and maize population density on small subsistence farms in Malawi. *Soil and Tillage Research*, 51(1-2), 113-119.