HL7 FHIR R4 — Chapter 4: Terminology Module — part 5 of 6 (sections 4.4.1.482 – 4.4.2.111)

Source: HL7 FHIR R4 (v4.0.1) specification, https://hl7.org/fhir/R4/ — text-only rendition for AI ingestion; page order follows toc.html.

4.4.1.482 Value Set http://hl7.org/fhir/ValueSet/procedure-not-performed-reason

URL: https://hl7.org/fhir/R4/valueset-procedure-not-performed-reason.html
Patient Care Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined at http://hl7.org/special/committees/CQI .

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/procedure-not-performed-reason
Version: | 4.0.1
Name: | ProcedureNotPerformedReason(SNOMED-CT)
Title: | Procedure Not Performed Reason (SNOMED-CT)
Definition: | Situation codes describing the reason that a procedure, which might otherwise be expected, was not performed, or a procedure that was started and was not completed. Consists of SNOMED CT codes, children of procedure contraindicated (183932001), procedure discontinued (416406003), procedure not done (416237000), procedure not indicated (428119001), procedure not offered (416064006), procedure not wanted (416432009), procedure refused (183944003), and procedure stopped (394908001).
Committee: | Patient Care Work Group
OID: | 2.16.840.1.113883.4.642.3.431 (for OID based terminology systems)
Copyright: | This resource includes content from SNOMED Clinical Terms® (SNOMED CT®) which is copyright of the International Health Terminology Standards Development Organisation (IHTSDO). Implementers of these specifications must have the appropriate SNOMED CT Affiliate license - for more information contact http://www.snomed.org/snomed-ct/get-snomed-ct or info@snomed.org
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: Procedure.statusReason (CodeableConcept / Example)

4.4.1.482.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include codes from http://snomed.info/sct where concept is-a 183932001 (Procedure contraindicated)
• Include codes from http://snomed.info/sct where concept is-a 416406003 (Procedure discontinued)
• Include codes from http://snomed.info/sct where concept is-a 416237000 (Procedure not done)
• Include codes from http://snomed.info/sct where concept is-a 428119001 (Procedure not indicated)
• Include codes from http://snomed.info/sct where concept is-a 416064006 (Procedure not offered)
• Include codes from http://snomed.info/sct where concept is-a 416432009 (Procedure not wanted)
• Include codes from http://snomed.info/sct where concept is-a 183944003 (Procedure refused)
• Include codes from http://snomed.info/sct where concept is-a 394908001 (Procedure stopped)

4.4.1.482.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 485 concepts

All codes from system http://snomed.info/sct

Code | Display | Definition
135809002 | Nitrate contraindicated
135811006 | Diuretic contraindicated
135822005 | Lipid lowering therapy contraindicated
135823000 | Calcium channel blocker contraindicated
169551000 | Contraception contraindicated
170954008 | Hormone replacement therapy (HRT) contraindicated
171277005 | Sensitivity C/I - immunization
171284002 | Pertussis vaccine contraindicated
183934000 | Sensitivity contraindication to procedure
183936003 | Contraindication to live immunization
266757004 | Medical contraindication to procedure
266758009 | Immunization contraindicated
268560002 | Medical C/I - immunization
275935009 | Surgery contraindicated
275974002 | Measles/mumps/rubella vaccine contraindicated

312451002 | Aspirin prophylaxis contra-indicated
315061006 | Warfarin contraindicated
315062004 | Beta blocker contraindicated
315363002 | Statins contraindicated
315364008 | Angiotensin converting enzyme inhibitors contraindicated
315591006 | Fibrates contraindicated
390795005 | Pneumococcal vaccination contraindicated
390796006 | Influenza vaccination contraindicated
390910005 | Thrombolysis contraindicated
394987009 | Angiotensin II receptor antagonists contraindicated
395174005 | Nicotine replacement therapy contraindicated
395175006 | Bupropion contraindicated
395675007 | Oral contraceptive pill contraindicated
407582002 | Clopidogrel contraindicated
413558003 | Anticoagulation contraindicated
414055003 | Drug dependence home detoxification contraindicated
414159007 | Exercise tolerance test contraindicated
414677003 | Metformin contraindicated
415571003 | Spirometry reversibility testing contraindicated
415666001 | Administration of sulfonylurea contraindicated
416091008 | Glitazones contraindicated
416475003 | Combined calcium and vitamin D3 preparation contraindicated
416704001 | DXA scan contraindicated
416759002 | Non-steroidal anti-inflammatory drugs contraindicated
416996000 | Strontium ranelate contraindicated
417101006 | Selective estrogen receptor modulator contraindicated
417128001 | Bisphosphonates contraindicated
438767006 | Termination of pregnancy contraindicated
438833006 | Administration of drug or medicament contraindicated
698468005 | Bisoprolol contraindicated
698521002 | Simvastatin contraindicated
699037006 | Hepatitis B vaccination contraindicated
703997000 | Teriparatide contraindicated
703999002 | Raloxifene contraindicated
707298000 | Phosphodiesterase-5 inhibitor contraindicated
715163003 | Diphtheria vaccination contraindicated
715166006 | Bacillus Calmette-Guerin vaccination contraindicated
170624001 | Oral steroids stopped
170688000 | Lithium stopped
170689008 | Injectable phenothiazine stopped
170919001 | Warfarin therapy stopped
170958006 | Hormonal replacement therapy (HRT) stopped
182840001 | Drug treatment stopped - medical advice
182841002 | Dr stopped drugs - ineffective
182842009 | Doctor stopped drugs - side effect
182843004 | Dr stopped drugs -inconvenient
182844005 | Doctor stopped drugs - patient dislikes
182845006 | Doctor stopped drugs - avoid interaction
182846007 | Doctor stopped drugs - medical aim achieved
274512008 | Drug therapy discontinued
309841001 | Thyroxine treatment stopped
394909009 | Syringe driver discontinued
395006008 | Medication stopped - interaction
395007004 | Medication stopped - ineffective
395008009 | Medication stopped - contra-indication
395009001 | Medication stopped - side effect
412782003 | Inappropriate medication stopped
425519007 | Cardiopulmonary resuscitation discontinued due to medical control order
425921008 | Cardiopulmonary resuscitation discontinued due to do not resuscitate order
426970003 | Cardiopulmonary resuscitation discontinued due to return of spontaneous circulation
426999008 | Cardiopulmonary resuscitation discontinued due to signs of death
427569000 | Cardiopulmonary resuscitation discontinued as per protocol
430279008 | Drug treatment stopped at end of clinical trial
438604008 | Peritoneal dialysis therapy discontinued
439516000 | Hemodialysis therapy discontinued
710765004 | Wound drain discontinued
713246009 | Procedure discontinued by healthcare professional
713247000 | Procedure discontinued by patient
713248005 | Procedure discontinued by doctor
162650008 | Patient not examined
162675007 | General examination not done
162882008 | Respiratory system not examined

162980001 | CVS not examined
163128002 | GIT not examined
163348001 | GU system not examined
163582008 | Nervous system not examined
163628001 | Sensory system not examined
163900009 | O/E - CNS not examined
164176006 | ENT examination not performed
164286000 | O/E - temperature not taken
164321005 | O/E - skin not examined
164445005 | O/E - extremities not examined
164503006 | Orthopedic exam. not done
164713001 | Neurological diagnostic procedure - not done
164725003 | Visual testing not done
164746007 | Auditory/vestibular test not done
164766003 | Special ENT procedure not done
164776000 | Special CVS test not done
164786004 | Special GIT test not done
164795007 | Special urinary test not done
164808009 | Special female test not done
164823004 | Special male test not done
164831009 | Musculoskeletal test not done
164853006 | ECG not done
164964005 | Susceptibility skin test not done
164978006 | Mantoux test not done
164985005 | Kveim test not done
164997006 | Hypersensitivity skin test not done
165008002 | Allergy testing not done
165017002 | Lung function testing not done
165026004 | Lung volume test not done
165035006 | Respiratory flow rate not measured
165074004 | Cardiac function test not done
165081006 | Exercise tolerance test not done
165104002 | Metabolic function not tested
165122004 | Non-surgical biopsy not done
165139002 | Endoscopy not carried out
165343008 | Laboratory test requested - not done
165377003 | Hematology test request - not done
165393007 | Hemoglobin not estimated
167219008 | Urine not examined
167260001 | Urine glucose test not done
167272007 | Urine protein test not done
167286006 | Urine ketone test not done
167296002 | Urine blood test not done
167306007 | Urine pH test not done
167317008 | Urine bacteria test not done
167322008 | Urine urobilinogen not tested
167594003 | Feces not examined
167701000 | CSF: not examined
167728005 | CSF chemistry - not tested
168497006 | Radiographic imaging procedure not carried out
169607004 | No antenatal care
169610006 | Antenatal care: not attended
169611005 | No antenatal care: not known pregnant
169699008 | Antenatal syphilis screening not done
169704000 | Antenatal blood group screening not done
169708002 | Antenatal sickle screening not done
170100005 | Child not examined at birth
170108003 | Child not examined at 10 days
170117003 | Child 6 week examination not attended
170126000 | 8-9 month exam not attended
170135007 | 18 month examination not attended
170144008 | 2.5 year examination not attended
170153001 | 3.5 year examination not attended
170162004 | 4.5 year examination not attended
170171008 | 10 year examination not attended
170180008 | 15 year examination not attended
170257006 | Child 1 year examination not attended
170266005 | Child 6 month examination not attended
170275007 | Child 21 month examination not attended
170284007 | Child 3 year exam not attended
170293008 | Child 39 month examination not attended
170304008 | Child 8 week examination not attended

182849000 | No drug therapy prescribed
183928007 | Residence remote from medical care
183929004 | No home medical services
183930009 | Awaiting admission elsewhere
266756008 | Medical care unavailable
266881002 | No history taken
309846006 | Treatment not available
310352003 | Mammography not attended
314374001 | 360 degree sweep of cervix not performed
371900001 | Medication not administered
373147003 | Medication not administered because contraindicated
373148008 | Thrombolytic agent not administered because contraindicated
390848004 | Retinae not examined
391016008 | Osteoporosis risk assessment defaulted
394965000 | Urine leukocyte test not done
396781004 | No chemotherapy AND/OR radiation therapy prior to lymphadenectomy
397004005 | Immunophenotypic analysis not performed
397399004 | Cell phenotyping not performed
399538001 | Cytogenetic study not performed
401318002 | Shuttle walking test not done
408504000 | Child 7 month examination not attended
408836004 | Sample not obtained
408837008 | Amniocentesis sample not obtained
416128008 | No past history of procedure
418014008 | Procedure not ordered
418731009 | No stain applied
428348004 | Cardiopulmonary resuscitation not attempted
439867009 | Determination of pattern of resistance to antiviral agent not done
441989008 | Delayed hypersensitivity skin test for histoplasmin not done
443788002 | Maternal postnatal examination not attended
698569004 | Postnatal depression not discussed
698753008 | Never had cervical smear
699231000 | Domestic abuse not discussed
703942005 | Never had mammogram
712751006 | Radiation therapy not done
712985002 | Chemotherapy not done
713207007 | Neck dissection not done
715621003 | Heart failure education not done
715622005 | Warfarin not prescribed
4451000175106 | Asthma action plan not done
4461000175108 | Cervical cancer screening not done
4471000175100 | Colorectal cancer screening not done
4501000175108 | Hemoglobin A1C test not done
4541000175105 | Spirometry not done
4551000175107 | Diabetic foot exam not done
4591000175100 | Bone density screening not done
5991000175105 | Peak expiratory flow not done
6021000175100 | Vaccination for diphtheria, pertussis, and tetanus not done
6031000175102 | Varicella vaccination not done
6041000175108 | Hepatitis A vaccination not done
6051000175105 | Hepatitis B vaccination not done
134390006 | Angiotensin converting enzyme inhibitor not indicated
134391005 | Statin not indicated
134392003 | Warfarin not indicated
134394002 | Aspirin not indicated
135805008 | Nitrate not indicated
135806009 | Calcium channel blocker not indicated
135807000 | Lipid lowering therapy not indicated
135808005 | Diuretic not indicated
183964008 | Treatment not indicated
183965009 | Surgery not indicated
183966005 | Drug treatment not indicated
315214003 | Beta blocker not indicated
407571005 | Clopidogrel not indicated
407572003 | Angiotensin II receptor antagonist not indicated
407573008 | Influenza vaccination not indicated
408339001 | Thrombolysis therapy not indicated
408396006 | Diabetic retinopathy screening not indicated
408397002 | Diabetic foot examination not indicated
408398007 | Smoking review not indicated
413167008 | Compression bandaging not indicated
413560001 | Anticoagulation not indicated

413812009 | Cervical smear not indicated
415119001 | Pneumococcal vaccination not indicated
415570002 | Spirometry not indicated
415667005 | Administration of sulfonylurea not indicated
416664005 | Selective estrogen receptor modulator not indicated
416670004 | Glitazones not indicated
416678006 | Combined calcium and vitamin D3 preparation not indicated
416741000 | Strontium ranelate not indicated
417013004 | DXA scan not indicated
417218000 | Carotid ultrasound not indicated
417434004 | Bisphosphonates not indicated
699048004 | Exercise tolerance test not indicated
700384008 | Antibiotic prophylaxis not indicated
703971006 | Hepatitis A vaccination not indicated
703996009 | Teriparatide not indicated
704000000 | Raloxifene not indicated
704033005 | Phosphodiesterase 5 inhibitor not indicated
704052004 | Referral to cardiac rehabilitation program not indicated
704096004 | Referral to heart failure exercise program not indicated
707287001 | Hepatitis B vaccination not indicated
707853004 | Testing for Ebola virus not indicated
709198003 | Education not indicated
712740006 | Medication monitoring not indicated
146922001 | Rubella screening not offered
147022006 | Double test not offered
148324002 | Screening not offered
148477008 | Immunization not offered
169608009 | Antenatal care: not offered
170115006 | Child 6 week examination not offered
170124002 | 8-9 month exam not offered
170133000 | 18 month examination not offered
170142007 | 2.5 year examination not offered
170151004 | 3.5 year examination not offered
170160007 | 4.5 year examination not offered
170169008 | 10 year examination not offered
170178002 | 15 year examination not offered
170255003 | Child 1 year examination not offered
170264008 | Child 6 month examination not offered
170273000 | Child 21 month examination not offered
170282006 | Child 3 year exam not offered
170291005 | Child 39 month examination not offered
170301000 | Child 8 week examination not offered
408506003 | Child 7 month examination not offered
444062004 | Maternal postnatal examination not offered
712790005 | Triple test not offered
712803007 | Administration of Anti-D globulin not offered
712812009 | Antenatal ultrasound scan not offered
712851004 | Alpha-fetoprotein blood test not offered
712852006 | Amniocentesis not offered
712854007 | Antenatal screening for viral hepatitis type B not offered
712869008 | Antenatal HIV (human immunodeficiency virus) screening not offered
712870009 | Antenatal Rhesus antibody screening not offered
712872001 | Antenatal thalassemia screening not offered
712996005 | Parenting education not offered
169609001 | Antenatal care: not wanted
169635000 | Parent craft not wanted
169649009 | A/N amniocentesis - not wanted
169660000 | A/N U/S scan not wanted
169687001 | Alpha-fetoprotein gene (AFP) blood test not wanted
169693009 | Rubella screening not wanted
169795009 | Triple test not wanted
169796005 | Double test not wanted
170116007 | Child 6 week examination not wanted
170125001 | 8-9 month exam not wanted
170134006 | 18 month examination not wanted
170143002 | 2.5 year examination not wanted
170152006 | 3.5 year examination not wanted
170161006 | 4.5 year examination not wanted
170170009 | 10 year examination not wanted
170179005 | 15 year examination not wanted
170256002 | Child 1 year examination not wanted
170265009 | Child 6 month examination not wanted

170274006 | Child 21 month examination not wanted
170283001 | Child 3 year exam not wanted
170292003 | Child 39 month examination not wanted
170302007 | Child 8 week examination not wanted
171034000 | Health education not wanted
171103002 | Screening not wanted
171152003 | Cancer of cervix screening not wanted
184163005 | Child surveillance not wanted
408505004 | Child 7 month examination not wanted
105480006 | Refusal of treatment by patient
134385008 | Referral to dietician declined
134386009 | Referral to chiropodist declined
134396000 | Statin declined
134397009 | Angiotensin converting enzyme inhibitor declined
134398004 | Warfarin declined
135812004 | Nitrate declined
135814003 | Diuretic declined
135825007 | Calcium channel blocker declined
135826008 | Lipid lowering therapy declined
165342003 | Patient refused laboratory test
165376007 | Patient refused hematology test
168499009 | Radiology refused
182895007 | Drug declined by patient
182896008 | Drug declined by patient - dislikes taste
182897004 | Drug declined by patient - side effects
182898009 | Drug declined by patient - inconvenient
182899001 | Drug declined by patient - problem swallowing
182900006 | Drug declined by patient - patient beliefs
182901005 | Drug declined by patient - alternative therapy
182902003 | Drug declined by patient - cannot pay script
182903008 | Drug declined by patient - reason unknown
183945002 | Procedure refused for religious reason
183946001 | Procedure refused - uncooperative
183947005 | Refused procedure - after thought
183948000 | Refused procedure - parent's wish
183949008 | Assessment examination refused
183956002 | Care/help refused by patient
183957006 | Patient refuses day hospital
183958001 | Refuses meals on wheels
183959009 | Social Services help refused
183960004 | Patient refuses hospital admit
183961000 | Part III accommodation refused
183962007 | Patient non-compliant with specific advice
185754009 | Refuses diabetes monitoring
275936005 | Patient noncompliance - general
315020006 | Beta blocker therapy refused
315021005 | Refuses CHD monitoring
315022003 | Nicotine replacement therapy refused
315023008 | Aspirin prophylaxis refused
315640000 | Influenza vaccination declined
371138003 | Refusal of treatment by parents
390914001 | Thrombolytic therapy refused
391015007 | Osteoporosis risk assessment refused
395176007 | Bupropion refused
395703005 | Meningitis C immunization refused
401047000 | Hypertension treatment refused
401084003 | Angiotensin II receptor antagonist declined
401086001 | Pneumococcal vaccination declined
406149000 | Medication refused
407583007 | Clopidogrel declined
408548005 | Magnetic resonance imaging scan declined
408549002 | Angiocardiography declined
408551003 | Exercise tolerance test refused
408558009 | Multidisciplinary team falls assessment declined
408559001 | Primary health care team falls assessment declined
408566000 | Echocardiogram declined
408567009 | CT scan brain declined
408569007 | Diagnostic procedure declined
408572000 | Carotid artery doppler declined
408778004 | Hepatitis B immunization declined
408791003 | BCG vaccination declined
408795007 | Vitamin K prophylaxis declined

412713002 | Coronary arteriography declined
412718006 | Chlamydia screening declined
412725004 | Medication review declined
412752009 | Diabetic foot examination declined
413122001 | Diabetic retinopathy screening refused
413123006 | Blood pressure procedure refused
413310006 | Patient non-compliant - refused access to services
413311005 | Patient non-compliant - refused intervention / support
413312003 | Patient non-compliant - refused service
413559006 | Anticoagulation declined
413756001 | Cardiac rehabilitation declined
415572005 | Spirometry test declined
416126007 | Bisphosphonates declined
416522000 | Strontium ranelate declined
416888009 | DXA scan declined
417045009 | Combined calcium and vitamin D3 preparation declined
417114007 | Refused referral to minor ailments clinic
417512003 | Selective estrogen receptor modulator declined
426544006 | History refused
428042006 | Medium-chain acyl-coenzyme A dehydrogenase deficiency screening test declined
428073005 | Alcohol consumption screening test declined
428299008 | Sickle cell screening declined
428650003 | Congenital hypothyroidism screening declined
428841003 | Cystic fibrosis screening declined
429284000 | CVD risk assessment declined
429688007 | Nurse triage declined
429726000 | Phenylketonuria screening declined
429739001 | Patient refused transportation
438370008 | Vascular disease risk assessment declined
438765003 | Human papillomavirus vaccination declined
439495000 | Counseling declined
440621003 | Referral declined by patient
442324008 | Antenatal screening declined
442392002 | Ambulance transport to hospital declined
442444001 | Referral to specialist alcohol treatment service declined by patient
444020006 | Maternal postnatal examination refused
473149004 | Long acting reversible contraception declined
698357006 | Abdominal aortic aneurysm screening declined
698451006 | Ankle brachial pressure index test declined
698460003 | Antiviral therapy declined
698473004 | Blood glucose test declined
698483000 | Weight management advice declined
698484006 | Measurement of waist circumference declined
698490005 | Urine dipstick test declined
698758004 | Nebivolol therapy refused
698950001 | Influenza A virus subtype H1N1 vaccination declined
699034004 | Home oxygen therapy declined
699042003 | History taking of sexual orientation declined
699053009 | Foot pulse check declined
699054003 | Administration of blood product declined
699128009 | Blood transfusion declined
699137009 | Carvedilol therapy declined
700359005 | Booster meningitis C vaccination declined
700411009 | Routine enquiry about domestic abuse declined
703427001 | Refusal of treatment by patient against dental advice
703989007 | Teriparatide therapy declined
703991004 | Raloxifene therapy declined
703992006 | Breastfeeding support declined
704046000 | Mental health assessment declined
704047009 | Phosphodiesterase 5 inhibitor declined
704049007 | Referral to erectile dysfunction clinic declined
704051006 | Referral to cardiac rehabilitation program declined
704097008 | Referral to heart failure exercise program declined
705140004 | Physical health assessment declined
706890009 | Vaginal birth after previous cesarean section refused
707745004 | Tuberculosis screening declined
707746003 | Screening chest X-ray declined
708129006 | Transfusion of blood product refused for religious reason
713068007 | Over 75 years of age health check declined
713615000 | Advance care planning declined
714747005 | Discussion about advance care planning declined
715508000 | Assessment using Generalized Anxiety Disorder 7 item scale declined

716048005 | Review of advance care plan declined
591000119102 | Vaccine refused by patient
921000119109 | Vaccine refused by parent
116101000119100 | Colonoscopy refused
314553311000087102 | Decided to wait for a specific provider
914933391000087108 | Request procedure deferral
168533005 | Radiotherapy stopped
170585007 | Cardiac disease treatment stopped
170588009 | Treatment for hypertension stopped
170620005 | Respiratory disease treatment stopped
170621009 | Home oxygen supply stopped
170675003 | Gastrointestinal tract treatment stopped
170687005 | Psychiatric treatment stopped
170700002 | Neurological disorder treatment stopped
170712003 | Epilepsy treatment stopped
170725006 | Ophthalmological treatment stopped
170800007 | Treatment of obesity stopped
170816009 | Endocrine disease treatment stopped
170826002 | B12 injections stopped
170834008 | Blood disorder treatment stopped
170843004 | Allergic disorder treatment stopped
170854009 | Rheumatology disorder treatment stopped
170863006 | Skin disorder treatment stopped
170872003 | Gynecological disorder treatment stopped
170882002 | Urinary disorder treatment stopped
170894006 | ENT disease treatment stopped
170904002 | Ostomy - treatment stopped
170912005 | High risk treatment stopped
170929008 | Repeat prescription treatment stopped
180863000 | Operation aborted
268521004 | Thyroid disease treatment stopped
288120003 | Operation abandoned before onset
391021006 | Osteoporosis treatment stopped
397023003 | Operation not completed
414120003 | ERCP not completed due to anatomical derangements from previous surgery
698459008 | Antidepressant drug treatment stopped
700110004 | Bisphosphonate prophylaxis suspended
708000007 | Insulin treatment stopped

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.483 Value Set http://hl7.org/fhir/ValueSet/audit-source-type

URL: https://hl7.org/fhir/R4/valueset-audit-source-type.html
Security Work Group | Maturity Level: 3 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/audit-source-type
Version: | 4.0.1
Name: | AuditEventSourceType
Title: | Audit Event Source Type
Definition: | The type of process where the audit event originated from.
Committee: | Security Work Group
OID: | 2.16.840.1.113883.4.642.3.466 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for AuditEventSourceType
• Resource: AuditEvent.source.type (Coding / Extensible)

4.4.1.483.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/security-source-type

4.4.1.483.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 9 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/security-source-type version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/security-source-type

Code | Display | Definition
1 | User Device | End-user display device, diagnostic device.
2 | Data Interface | Data acquisition device or instrument.
3 | Web Server | Web Server process or thread.
4 | Application Server | Application Server process or thread.
5 | Database Server | Database Server process or thread.
6 | Security Server | Security server, e.g. a domain controller.
7 | Network Device | ISO level 1-3 network component.
8 | Network Router | ISO level 4-6 operating software.
9 | Other | Other kind of device (defined by DICOM, but some other code/system can be used).

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.484 Value Set http://hl7.org/fhir/ValueSet/conditional-delete-status

URL: https://hl7.org/fhir/R4/valueset-conditional-delete-status.html
FHIR Infrastructure Work Group | Maturity Level: N | Normative (from v4.0.0) | Use Context: Any

| This page has been approved as part of an ANSI standard. See the Conformance Package for further details.

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/conditional-delete-status
Version: | 4.0.1
Name: | ConditionalDeleteStatus
Title: | ConditionalDeleteStatus
Definition: | A code that indicates how the server supports conditional delete.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.194 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ConditionalDeleteStatus

• Resource: CapabilityStatement.rest.resource.conditionalDelete (code / Required)

4.4.1.484.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/conditional-delete-status

4.4.1.484.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 3 concepts

Expansion based on http://hl7.org/fhir/conditional-delete-status version 4.0.1

All codes from system http://hl7.org/fhir/conditional-delete-status

Code | Display | Definition
not-supported | Not Supported | No support for conditional deletes.

single | Single Deletes Supported | Conditional deletes are supported, but only single resources at a time.
multiple | Multiple Deletes Supported | Conditional deletes are supported, and multiple resources can be deleted in a single interaction.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.485 Value Set http://hl7.org/fhir/ValueSet/adverse-event-causality-assess

URL: https://hl7.org/fhir/R4/valueset-adverse-event-causality-assess.html
Patient Care Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/adverse-event-causality-assess
Version: | 4.0.1
Name: | AdverseEventCausalityAssessment
Title: | AdverseEventCausalityAssessment
Definition: | Codes for the assessment of whether the entity caused the event.
Committee: | Patient Care Work Group
OID: | 2.16.840.1.113883.4.642.3.840 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for AdverseEventCausalityAssessment

• Resource: AdverseEvent.suspectEntity.causality.assessment (CodeableConcept / Example)

4.4.1.485.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/adverse-event-causality-assess

4.4.1.485.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 6 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/adverse-event-causality-assess version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/adverse-event-causality-assess

Code | Display | Definition
Certain | Certain | i) Event or laboratory test abnormality, with plausible time relationship to drug intake; ii) Cannot be explained by disease or other drugs; iii) Response to withdrawal plausible (pharmacologically, pathologically); iv) Event definitive pharmacologically or phenomenologically (i.e. an objective and specific medical disorder or a recognized pharmacological phenomenon); or v) Re-challenge satisfactory, if necessary.
Probably-Likely | Probably/Likely | i) Event or laboratory test abnormality, with reasonable time relationship to drug intake; ii) Unlikely to be attributed to disease or other drugs; iii) Response to withdrawal clinically reasonable; or iv) Re-challenge not required.
Possible | Possible | i) Event or laboratory test abnormality, with reasonable time relationship to drug intake; ii) Could also be explained by disease or other drugs; or iii) Information on drug withdrawal may be lacking or unclear.
Unlikely | Unlikely | i) Event or laboratory test abnormality, with a time to drug intake that makes a relationship improbable (but not impossible); or ii) Disease or other drugs provide plausible explanations.
Conditional-Classified | Conditional/Classified | i) Event or laboratory test abnormality; ii) More data for proper assessment needed; or iii) Additional data under examination.
Unassessable-Unclassifiable | Unassessable/Unclassifiable | i) Report suggesting an adverse reaction; ii) Cannot be judged because information is insufficient or contradictory; or iii) Data cannot be supplemented or verified.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)

Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.486 Value Set http://hl7.org/fhir/ValueSet/verificationresult-primary-source-type

URL: https://hl7.org/fhir/R4/valueset-verificationresult-primary-source-type.html
Patient Administration Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/verificationresult-primary-source-type
Version: | 4.0.1
Name: | primary-source-type
Title: | Primary-source-type
Definition: | Type of the validation primary source
Committee: | Patient Administration Work Group
OID: | 2.16.840.1.113883.4.642.3.892 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for primary-source-type
• Resource: VerificationResult.primarySource.type (CodeableConcept / Example)

4.4.1.486.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/primary-source-type

4.4.1.486.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 9 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/primary-source-type version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/primary-source-type

Code | Display | Definition
lic-board | License Board
prim | Primary Education
cont-ed | Continuing Education
post-serv | Postal Service
rel-own | Relationship owner
reg-auth | Registration Authority
legal | Legal source
issuer | Issuing source
auth-source | Authoritative source

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.487 Value Set http://hl7.org/fhir/ValueSet/document-relationship-type

URL: https://hl7.org/fhir/R4/valueset-document-relationship-type.html
Structured Documents Work Group | Maturity Level: 3 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/document-relationship-type
Version: | 4.0.1
Name: | DocumentRelationshipType
Title: | DocumentRelationshipType
Definition: | The type of relationship between documents.
Committee: | Structured Documents Work Group
OID: | 2.16.840.1.113883.4.642.3.244 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for DocumentRelationshipType
• Resource: DocumentReference.relatesTo.code (code / Required)
• Resource: Composition.relatesTo.code (code / Required)

4.4.1.487.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/document-relationship-type

4.4.1.487.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 4 concepts

Expansion based on http://hl7.org/fhir/document-relationship-type version 4.0.1

All codes from system http://hl7.org/fhir/document-relationship-type

Code | Display | Definition
replaces | Replaces | This document logically replaces or supersedes the target document.
transforms | Transforms | This document was generated by transforming the target document (e.g. format or language conversion).
signs | Signs | This document is a signature of the target document.
appends | Appends | This document adds additional information to the target document.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.488 Value Set http://hl7.org/fhir/ValueSet/subscription-channel-type

URL: https://hl7.org/fhir/R4/valueset-subscription-channel-type.html
FHIR Infrastructure Work Group | Maturity Level: 3 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/subscription-channel-type
Version: | 4.0.1
Name: | SubscriptionChannelType
Title: | SubscriptionChannelType
Definition: | The type of method used to execute a subscription.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.500 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for SubscriptionChannelType
• Resource: Subscription.channel.type (code / Required)

4.4.1.488.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/subscription-channel-type

4.4.1.488.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 5 concepts

Expansion based on http://hl7.org/fhir/subscription-channel-type version 4.0.1

All codes from system http://hl7.org/fhir/subscription-channel-type

Code | Display | Definition
rest-hook | Rest Hook | The channel is executed by making a post to the URI. If a payload is included, the URL is interpreted as the service base, and an update (PUT) is made.
websocket | Websocket | The channel is executed by sending a packet across a web socket connection maintained by the client. The URL identifies the websocket, and the client binds to this URL.
email | Email | The channel is executed by sending an email to the email addressed in the URI (which must be a mailto:).
sms | SMS | The channel is executed by sending an SMS message to the phone number identified in the URL (tel:).
message | Message | The channel is executed by sending a message (e.g. a Bundle with a MessageHeader resource etc.) to the application identified in the URI.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.489 Value Set http://hl7.org/fhir/ValueSet/medication-codes

URL: https://hl7.org/fhir/R4/valueset-medication-codes.html
Pharmacy Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/medication-codes
Version: | 4.0.1
Name: | SNOMEDCTMedicationCodes
Title: | SNOMED CT Medication Codes
Definition: | This value set includes all drug or medicament substance codes and all pharmaceutical/biologic products from SNOMED CT - provided as an exemplar value set.
Committee: | Pharmacy Work Group
OID: | 2.16.840.1.113883.4.642.3.375 (for OID based terminology systems)
Copyright: | This resource includes content from SNOMED Clinical Terms® (SNOMED CT®) which is copyright of the International Health Terminology Standards Development Organisation (IHTSDO). Implementers of these specifications must have the appropriate SNOMED CT Affiliate license - for more information contact http://www.snomed.org/snomed-ct/get-snomed-ct or info@snomed.org
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: ActivityDefinition.product[x] (Reference(Medication|Substance)|CodeableConcept / Example)
• Profile: ShareableActivityDefinition: ActivityDefinition.product[x] (Reference, CodeableConcept / Example)
• Resource: CarePlan.activity.detail.product[x] (CodeableConcept|Reference(Medication|Substance) / Example)
• Resource: MedicationRequest.medication[x] (CodeableConcept|Reference(Medication) / Example)
• Resource: MedicationAdministration.medication[x] (CodeableConcept|Reference(Medication) / Example)
• Resource: MedicationStatement.medication[x] (CodeableConcept|Reference(Medication) / Example)
• Resource: MedicationDispense.medication[x] (CodeableConcept|Reference(Medication) / Example)
• Resource: MedicationKnowledge.code (CodeableConcept / Example)

4.4.1.489.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include codes from http://snomed.info/sct where concept is-a 410942007 (Drug or medicament)

• Include codes from http://snomed.info/sct where concept is-a 373873005 (Pharmaceutical / biologic product)

• Include codes from http://snomed.info/sct where concept is-a 106181007 (Immunologic substance)

4.4.1.489.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 2993 concepts

All codes from system http://snomed.info/sct

Code	Display	Definition
261000	Codeine phosphate	
336001	Fibrinogen Tokyo II	
363000	Fibrinogen San Juan	
519005	Free protein S	
585007	SP - Substance P	
693002	Trichothecene	
698006	Erythromycin lactobionate	
699003	Coal tar extract	
747006	Oxamniquine	
873008	Urethan	
925002	Sodium iodipamide	
1018001	Nornicotine	
1025008	^93^Molybdenum	
1057009	Phosphate salt	
1091008	Coagulation factor inhibitor	
1171004	Fibrinogen Kawaguchi	
1190007	Mephenoxalone	
1244009	Fibrinogen Madrid I	
1269009	Amikacin sulfate	
1325004	Metocurine iodide	
1336006	Deoxycortone	
1355006	Coagulation factor IX Oxford 3 variant	
1368003	Iodine 131	
1381005	^126^Iodine	
1450002	Methylpentynol	
1476002	Codeine sulfate	
1536005	Pargyline hydrochloride	
1575001	Maltose tetrapalmitate	
1603001	Cobalt isotope	
1668008	Ceforanide	
1914001	von Willebrand factor antibody	
1944003	Coagulation factor X Patient variant	
1956002	Buclizine hydrochloride	
1971003	Loxapine hydrochloride	
2029004	Fibrinogen Oslo II	
2125008	Betazole	
2195004	Tocainide hydrochloride	
2197007	Boric acid topical agent	
2201007	Bacteriopurpurin	
2212007	Fibrinogen Bethesda II	
2215009	Azuresin	
2249001	Gentamicin sulfate	
2264001	Selenium isotope	
2346005	Vascormone	
2537003	Antituberculosis agent	
2660003	Sodium dehydrocholate	
2680002	Factor XIII antibody	
2796008	Methanthelinium	
2799001	Methylbenzethonium	
2869004	Ethanoic acid	
2878005	Pethidine hydrochloride	
2927004	Rescinnamine	
2958003	Zinc caprylate	
2964005	Dimethoxyamphetamine	
2991007	Mecamylamine hydrochloride	
2995003	Arecoline	
3031003	Dihydroxyaluminum sodium carbonate	
3066001	Triiodothyroacetic acid	
3087006	Oil of male fern	
3136005	Cefoperazone sodium	
3142009	Azacyclonol	

3145006 | Penicillic acid
3209002 | Pancuronium sodium
3212004 | Manganese sulfate
3225007 | Fibrinogen Seattle I
3342004 | Copper isotope
3378009 | Imipramine hydrochloride
3379001 | Merthiolate
3501003 | Isoxsuprine hydrochloride
3592004 | Short-acting thyroid stimulator
3597005 | Acebutolol hydrochloride
3617004 | ^79^Selenium
3672002 | Fibrinogen Caracas
3693004 | Fibrinogen Dusart
3793006 | Soluble barium compound
3823007 | Prochlorperazine edisylate
3829006 | Fe - Iron
3874004 | Hydrocodone bitartrate
3906002 | Rectified birch tar oil
3932008 | ^64^Copper
3941003 | Metronidazole hydrochloride
3945007 | Tin isotope
3983008 | N,-N-dimethyltryptamine
4014000 | Sulfisomidine
4025009 | Captodiame
4043008 | Etidocaine hydrochloride
4076007 | PTH - Parathyroid hormone
4188007 | Fibrinogen Sydney II
4203009 | Imipramine pamoate
4334005 | Tar oil
4353000 | Di-n-butylphthalate
4355007 | Coagulation factor IX San Dimas variant
4413004 | Benzypyrinium
4423008 | Fibrinogen New York II
4480008 | Sulfaethidole
4518006 | Buthenal
4616002 | Triclobisonium chloride
4681002 | Potassium permanganate
4700006 | Bovine insulin
4780009 | Secbutabarbital sodium
4814001 | Valethamate
4844003 | 3,3' Diiodothyronine
5007006 | Papain
5142007 | Coagulation factor II Houston variant
5167005 | Zinc chloride fumes
5172001 | Coagulation factor Xa
5220000 | Bacitracin
5312000 | ^65^Zinc
5405008 | ^60^Cobalt
5477001 | Ferric subsulfate
5641004 | Valproate semisodium
5647000 | Griseofulvin ultramicrosize
5657004 | Coal tar topical solution
5681006 | Ceftizoxime sodium
5691000 | Absorbable gelatin sponge
5699003 | Somatomedin C
5739006 | Stramonium
5767002 | Sulfamerazine
5774007 | White petrolatum
5907009 | Quinidine polygalacturonate
5955009 | Diethylene glycol
6054001 | Doxylamine succinate
6088007 | Benzfetamine hydrochloride
6162007 | Meclocycline
6237004 | Bemegride
6260007 | Protokylol
6333002 | Squill extract
6394006 | Phentermine hydrochloride
6495008 | Fibrinogen Montreal II
6513000 | Flumethiazide
6524003 | Distilled spirits
6602005 | Aminoacridine
6612003 | Chloramphenicol sodium succinate

6710000 | Nitric oxide
6717002 | Nifuroxime
6790004 | Aminopterin
6809003 | Phomopsin
6826009 | Sterol hormone
6837005 | Dextropropoxyphene napsylate
6865007 | Theophylline calcium salicylate
6873003 | Cefapirin sodium
6881002 | Magnesium fumes
6910009 | Sodium fluoride
6927001 | ^121^Tin
6952004 | ^121m^Tin
6992002 | Triflupromazine hydrochloride
7034005 | Diclofenac
7084003 | Fibrinogen Buenos Aires II
7156005 | Phosphorus isotope
7179006 | Prekallikrein
7302008 | Ambuphylline
7318002 | Bacteriochlorophyll
7328006 | Red petrolatum
7348004 | Factor II
7451005 | Tobramycin ophthalmic agent
7509000 | Mannitol hexanitrate
7515000 | Hepatotoxic mycotoxin
7549008 | Fibrinogen Bethesda I
7608003 | Triethylene glycol
7737009 | Chlortetracycline hydrochloride
7770004 | ^89^Strontium
7774008 | Neo-b-vitamin A>1<
7816005 | Antazoline hydrochloride
7834009 | Acetyl digitoxin
7904003 | Deanol
7975001 | Amiphenazole
8030004 | PEG - Polyethylene glycol
8143001 | Diosmin
8202008 | ^43^Potassium
8203003 | Human menopausal gonadotropin
8222007 | Coagulation factor II Padua variant
8252004 | Chlorothiazide sodium
8397006 | Nicotine resin complex
8450009 | Lemon grass oil
8534003 | ^110^Tin
8631001 | Potassium chloride
8689007 | Kanamycin sulfate
8701002 | Sulfachlorpyridazine
8767001 | Santonin
8886003 | Flecainide acetate
8919000 | Biotin
8987006 | Cycle-phase specific agent
9174005 | Fibrinogen Poitiers
9234005 | Chlorobutanol
9271007 | Fibrinogen Pontoise
9396007 | ^60^Iron
9457002 | Fibrinogen Almeria
9471005 | Polypropylene glycol
9530002 | Amine hormone
9532005 | Coagulation factor XIIIa
9539001 | Chlorprothixene lactate
9643009 | Chlorphentermine
9663002 | Pecazine
9676008 | Fibrinogen New York III
9680003 | Central depressant
9721008 | Phencyclidine
9765000 | Lithium salt
9797000 | Phosphorus trichloride
9817005 | Mycoplasma pulmonis antibody test kit
9910008 | Oxymetazoline hydrochloride
9974009 | Angiotensin
9992001 | Molybdenum radioisotope
10016008 | Bithionol
10020007 | Biperiden hydrochloride
10109009 | Fibrinogen London III

10133003 | Cyclizine lactate
10174003 | Procarbazine hydrochloride
10192006 | Prostaglandin F>2<
10202007 | Prostaglandin E3
10240005 | Lethanes
10270000 | Erythromycin estolate
10282009 | Betahistidine
10324005 | Demeclocycline hydrochloride
10329000 | Zinc insulin
10333007 | Clobenoside
10342000 | Heparin cofactor II
10354000 | Somantin
10377000 | Sodium nitrite
10424009 | Maprotiline hydrochloride
10466005 | Phosphide compound
10471003 | Fibrinogen Vienna
10500003 | Xanthinol
10534002 | TRH - Thyrotrophin releasing hormone
10595003 | Pseudoephedrine sulfate
10644005 | Black phosphorus
10660009 | Luteoskyrin
10682002 | Fibrinogen Grand Rapids
10730008 | Azlocillin sodium
10750007 | Neurotoxic mycotoxin
10751006 | Netilmicin sulfate
10767000 | Calcium phosphate dibasic
10782005 | PG - Pentagastrin
10790005 | ^53^Manganese
10843002 | Anterior pituitary hormone
10955007 | Anti-factor X
11036001 | Alum
11038000 | Dipentene (limonene)
11115001 | Thromboxane A2
11121002 | ^135^Iodine
11137008 | Chymotrypsin
11170003 | Vanillin
11202003 | Manganese salt
11331001 | ^56^Cobalt
11345007 | Tribromsalan
11420005 | Rhubarb agent
11473005 | Trichlormethiazide
11496005 | Mercuric chloride
11504003 | Edrophonium chloride
11633008 | Flurbiprofen sodium
11644000 | Piperacillin sodium
11652002 | VIP - Vasoactive intestinal polypeptide
11714005 | Strong silver protein
11943009 | Hydroxydione
11984007 | 1, Hydroxy cholecalciferol
11986009 | Penicillin G potassium
12009000 | Coagulation factor IX Chapel Hill variant
12034000 | Coagulation factor II Salatka variant
12177002 | Pseudoephedrine hydrochloride
12208001 | Syrosingopine
12218006 | Diltiazem hydrochloride
12290003 | Emetine hydrochloride
12315006 | Halazone
12391001 | Dextran 70
12525000 | Tybamate
12568004 | Belladonnine
12578001 | Erythromycin ethylsuccinate
12597001 | Sn - Tin
12689001 | Magnesium radioisotope
12716009 | Aluminum carbonate
12801003 | Iodamide meglumine
12821002 | Clemizole
12870003 | Coagulation factor IX Durham variant
12930006 | Calcium phosphate dibasic dihydrate
12970004 | Inositol hexanitrate
13030002 | Piperocaine
13150000 | Animal fat
13188003 | Tobramycin sulfate

13235001 | Vitamin B2
13342004 | Diethyl 2-chlorovinyl phosphate
13502005 | Hydroxychloroquine sulfate
13585009 | Cefotetan
13668001 | Propylene glycol
13708006 | Rectified pine tar oil
13787003 | Protein secretory trypsin inhibitor
13789000 | Coal tar creosote
14013006 | Guanadrel sulfate
14071002 | ^90^Strontium
14146003 | Potassium isotope
14172007 | Intrinsic factor concentrate agent
14193005 | Sulfobromophthalein
14285000 | Coagulation factor XI variant type III
14312008 | Vitamin L>2<
14321009 | Difolatan
14340003 | Verapamil hydrochloride
14349002 | Fibrinogen Seattle II
14399003 | Iodine radioisotope
14409006 | Neocinchophen
14438009 | Carbenicillin disodium
14443002 | Aminoglycoside -class of antibiotic-
14461006 | Aluminum phosphate
14507006 | Arsthinol
14638000 | Thiobarbiturate
14645000 | Zinc phenolsulfonate
14715007 | Dextran 75
14743003 | Cinchonine
14767006 | A-1AT - Alpha-1-antitrypsin
14796007 | Amphechloral
14819006 | Aspidium
14903000 | Antimony sodium thioglycolate
14905007 | Promethazine hydrochloride
15009009 | Meprylcaine
15017001 | Beeswax
15098005 | Alseroxylon
15129007 | Zinc propionate
15322006 | Benzoquinonium
15331006 | Glycine - amino acid
15352003 | Cyproheptadine hydrochloride
15505005 | Preprodynorphin
15571002 | Mezlocillin sodium
15660006 | Bleomycin sulfate
15698006 | Lysergic acid diethylamide
15730005 | Porphyrin
15785009 | Phenazopyridine
15810003 | Tuaminoheptane
15895007 | Fibrinogen London I
15901005 | Fibrinogen Paris III
16106007 | Sulfametoxydiazine
16125005 | Styramate
16130009 | Deoxyribonuclease IV (Phage T>4<-induced)
16214005 | Deslanoside
16257000 | Dopamine hydrochloride
16276003 | Coagulation factor IX Eagle Rock variant
16318005 | Dibenzothiepin
16355005 | Tetracycline hydrochloride
16359004 | Phthalylsulfathiazole
16392005 | Hexylcaine
16395007 | Pituitary gonadotropin
16462002 | Alpha neoendorphin
16477005 | Diagnostic vaccine
16492006 | Cloxacillin sodium
16613008 | Prostaglandin D2
16624005 | Bromide salt
16628008 | GRF - Growth hormone releasing factor
16670003 | Fibrinopeptide B-beta 1-42
16683002 | Progesterone
16717002 | 4-Pregnen-21-ol-3,11,20-trione
16744007 | Lactobacillus acidophilus agent
16745008 | Xenon isotope
16748005 | Zolamine

16808006 | Trichloroethylene
16826009 | Pentamidine isethionate
16915004 | Streptozocin
16923002 | Lupus anticoagulant
16946000 | Triacetin
17008002 | Levallorphan
17062003 | Nafoxidine hydrochloride
17072000 | Cathepsin D
17117004 | Androsterone
17147002 | CA - Cholic acid
17212003 | Bismuth subcarbonate
17240008 | Oil in water agent
17243005 | Uramustine
17244004 | Apraclonidine hydrochloride
17356001 | Pralidoxime chloride
17614005 | Fibrinogen Buenos Aires I
17731005 | Coagulation factor IX London variant
17777001 | Blastomycin
17798001 | Coagulation factor II Cardeza variant
17836006 | Aromatic ammonia spirit
17916009 | Propylene glycol dinitrate
17932007 | Fibrinogen Vicenza
17942009 | Fibrinogen Houston
17990002 | Melarsoprol
17991003 | Fibrinogen Adelaide
18017000 | Phenylglycidyl ether
18039003 | Alkyl mercuric phosphate
18143001 | Fibrinogen Quebec II
18220004 | Thyroid hormone
18288009 | vWF- von Willebrand factor
18321003 | Thiethylperazine maleate
18394004 | Collagenase agent
18414002 | VITD - Cholecalciferol
18449009 | Lincomycin hydrochloride
18462008 | Methdilazine
18535002 | Hypothalamic releasing factor
18550006 | Thioridazine hydrochloride
18600008 | Glucurolactone
18616005 | Lithium hydride
18712002 | Phenacemide
18815007 | Diethylphthalate
18832006 | Butylphenamide
18852007 | Fibrinogen New York IV
18959002 | Dibenzazepine derivative
18970009 | Prolactin releasing factor
19012003 | Fibrinogen Tokyo I
19041007 | Tolazoline hydrochloride
19046002 | Fibrinogen Pamplona
19114000 | Mafenide acetate
19126005 | Merbromin
19163001 | Prohormone
19205004 | Secretin
19421007 | Chloroprocaine hydrochloride
19427006 | ^59^Nickel
19462001 | Carbon dioxide absorbent
19499001 | Carbinoxamine maleate
19510001 | Diphenhydramine hydrochloride
19524002 | Penthienate
19595005 | Phenolphthalein
19622008 | ^48^Chromium
19839007 | Sorbitol
19893005 | Potassium dichromate
19918001 | Dihydroergocornine
19967004 | Viomycin
19978007 | Hexafluorenium
20056006 | Dibromosalicylaldehyde
20170008 | Lung surfactant
20217007 | Trimethaphan camsylate
20229003 | Pyrrobutamine
20231007 | Aminosalicylate sodium
20238001 | Chlorinated lime
20327004 | Sodium caprylate

20340009 | Methysergide maleate
20368008 | Diphacinone
20378006 | Methyldimethoxyamphetamine
20379003 | Neomycin C
20413008 | Levopropoxyphene
20450009 | Ciprofloxacin hydrochloride
20468007 | Isopropamide
20495002 | Fibrinogen Haifa
20686000 | Fibrinogen Christchurg II
20752000 | Prothrombin antibody
20771003 | Congenital dysfibrinogen
20844009 | Oxyprocaine
20847002 | Streptokinase agent
20887007 | Tretamine
21028006 | Fibrinogen Bergamo I
21066009 | Buprenorphine hydrochloride
21075006 | ^61^Cobalt
21094001 | Coal tar ointment
21168008 | Acetosulfone
21175009 | Methanthelinium bromide
21235009 | Piperoxan
21246007 | Fibrinogen Detroit
21289006 | Platelet factor 4
21303006 | Methoxamine hydrochloride
21373005 | Soap enema
21394008 | Adiphenine
21518006 | Naloxone hydrochloride
21556007 | 4-Methylaminophenol sulfate
21559000 | ^69m^Zinc
21566004 | Methyldopate hydrochloride
21568003 | Adrenal cortical hormone
21572004 | I123 - Iodine 123
21611007 | Boric acid
21614004 | Phenelzine sulfate
21706000 | Tetrahydrofolic acid
21891005 | Digestive enzyme
21903000 | Bismuth violet
21919007 | Opium
22038003 | Se - Selenium
22086005 | Sodium antimonyl gluconate
22165008 | Metamizole sodium
22236007 | Acetarsol
22242006 | Glutardialdehyde
22250002 | Fibrinogen Birmingham
22362000 | Calcium hydrogen phosphate anhydrous
22424004 | Ochratoxin
22453003 | Cathepsin G
22496008 | Fibrinogen Cleveland I
22606007 | Vitamin K>2<
22635004 | Factor V antibody
22654004 | Propantheline bromide
22769006 | Penthienate bromide
22779008 | Coagulation factor II Habana variant
22790003 | Physostigmine sulfate
22792006 | Tripelennamine citrate
22827004 | Prochlorperazine maleate
22840009 | Tetraethyl pyrophosphate
22882008 | Coagulation factor II Molise variant
22941009 | Cortodoxone
22976006 | Aluminum acetate
23053002 | Iophenoic acid
23068001 | Caffeine citrate
23077008 | Barbituric acid
23176001 | Bacampicillin hydrochloride
23295004 | Coagulation factor I
23375008 | Polymyxin E sulfate
23433006 | Calciferol
23459009 | Selenium radioisotope
23564005 | Dyclonine
23692005 | Tribasic calcium phosphate
23701001 | Triclocarban
23814005 | Guanethidine monosulfate

23816007 | THF - Tetrahydrocortisol
23861006 | Phenolsulfonphthalein
23862004 | Fibrinogen Bethesda III
23866001 | Fluoroacetic acid
23883005 | Methadone hydrochloride
23959001 | TG - Thyroglobulin
23969007 | Tryparsamide
24022008 | Bupivacaine hydrochloride
24202000 | Ranitidine hydrochloride
24237002 | Prostaglandin F>1<
24261009 | Trimethobenzamide hydrochloride
24336008 | Aminophylline anhydrous
24357000 | CSF-1
24434007 | Sodium tartrate
24435008 | Fibrinogen Versailles
24556008 | n-Butylglycidylether
24570008 | Purgative
24583009 | Terbutaline sulfate
24650007 | Dihydro-alpha-ergocryptine
24686008 | Acaricide
24721007 | Chlorothymol
24751001 | Oxymorphone
24809001 | Spectinomycin hydrochloride
24823004 | Pipobroman
24838003 | Undecylenic acid and zinc undecylenate
24869004 | Lampit
25002001 | Perazine
25013003 | Pyrantel pamoate
25027008 | Glycoprotein hormone
25128000 | ^72^Zinc
25141001 | Tubocurarine chloride
25183006 | Manganese chloride
25204006 | Aluminum soluble salt
25205007 | Fluorine radioisotope
25213008 | Bicyclic antidepressant
25217009 | Pituitary follicle stimulating hormone
25254000 | Procainamide hydrochloride
25305005 | Long-acting insulin
25307002 | Petrolatum
25401000 | Barbiturate analog
25500001 | ^40^Potassium
25525005 | Protein C
25538002 | Tiotixene hydrochloride
25571003 | Clodantoin
25607008 | DDimer
25620006 | Bitter orange oil
25710007 | ^113^Tin
25747007 | ^135m^Xenon
25796002 | Aluminum aspirin
25886000 | Fibrinogen Bergamo III
25911004 | Prostaglandin H2
26120001 | Desipramine hydrochloride
26191006 | Bromazine hydrochloride
26277008 | Dynorphin
26288002 | Mitotane
26327007 | Red phosphorus
26346008 | Ethambutol hydrochloride
26351002 | Prostaglandin
26371006 | Chlorophacinone
26379008 | Quinisocaine
26437003 | Oil of mustard
26469007 | Prepro-opiomelanocortin
26518005 | Coagulation factor XIa
26656004 | Aromatic castor oil
26817007 | Methylated naphthalene
26945002 | Phendimetrazine tartrate
26992003 | Chlorisondamine
27054007 | ^57^Cobalt
27079005 | Meclocycline sulfosalicylate
27081007 | ^127^Xenon
27082000 | Sulfapyridine
27122000 | ^56^Nickel

27184001 | 17-alpha-Hydroxypregnenolone
27192005 | Aminosalicylic acid
27244000 | Lithium isotope
27248002 | Coagulation factor X R.E.D. variant
27345002 | Hemin
27361000 | Barium salt
27499006 | Oxyphencyclimine
27586005 | Undecoylium chloride iodine
27594003 | Ferric pyrophosphate
27656005 | Gitalin amorphous
27730007 | Merodicein
27736001 | Bacitracin A
27766008 | Prothipendyl
27822002 | Phenylpropylmethylamine
27928002 | Flurazepam hydrochloride
27931001 | Dipeptidyl peptidase I
27989007 | Coagulation factor II Segovia variant
28021008 | Ethyl acrylate
28029005 | Metescufylline
28069006 | Refrigerant anesthetic
28121005 | Iofendylate
28223003 | Cycloguanil
28268006 | Pregnanediol
28344001 | Mephenytoin
28421003 | Sorbate
28464005 | Dioxyline
28521000 | Coagulation factor II Denver variant
28580002 | Diprenorphine
28588009 | Cefaloridine
28662003 | Hydralazine hydrochloride
28702005 | Ambutonium
28825003 | Sterigmatocystin
28829009 | Coal tar naphtha
28927005 | Diphenylheptane derivative
28942008 | Coconut oil
28983006 | 6-Ketoprostaglandin F1
28999000 | Fusarin
29011006 | Manganese trioxide
29184007 | Diphemanil metilsulfate
29190006 | Fentanyl citrate
29301006 | Isoprenaline hydrochloride
29327006 | Chloramphenicol palmitate
29406002 | alpha-Amylase agent
29522004 | Toluidine blue stain
29527005 | Benzatropine mesylate
29531004 | Nickel radioisotope
29584000 | Diagnostic antigen
29588002 | Brompheniramine maleate
29666006 | Nortriptyline hydrochloride [enuresis - no agents here]
29671004 | Lithium bromide
29725000 | Heparin calcium
29765001 | Fumagillin
29783009 | Chromocarb
29876006 | ^129m^Xenon
30034004 | Dimethoxanate
30053009 | Brefeldin
30094001 | Riboflavin dinucleotide
30145004 | Activin hormone
30178006 | Vitamin D
30179003 | Carbonyl fluoride
30203009 | Paromomycin sulfate
30205002 | Thymic T lymphocyte factor
30236005 | Tilorone
30324001 | ^132^Iodine
30325000 | Chlorfenvinphos
30326004 | Atrial natriuretic hormone
30329006 | Triflupromazine
30333004 | Mercaptomerin sodium
30424002 | Proparacaine hydrochloride
30589007 | Dibutyl phosphate
30658004 | Turacoporphyrin
30676006 | Metharbital

30745005 | Loxapine succinate
30804005 | Factor VII
30820000 | Phosphorus
30827002 | Azapetine
30848000 | Fibrinogen Oslo III
30863002 | Desiccated whole bile
30990007 | Abnormal fibrinogen
31011004 | 4-hydroxycoumarin
31055005 | GIH - Gastrointestinal hormone
31147000 | Metoclopramide hydrochloride
31178001 | Bethanechol chloride
31422009 | Ox bile extract
31522006 | Mild silver protein
31617001 | Hydrophilic petrolatum
31622001 | ^127m^Xenon
31662002 | Orange flower oil
31706007 | Ketamine hydrochloride
31707003 | Zinc bacitracin
31731008 | Magnesium phosphate
31780003 | Preproenkephalin
31787000 | Coagulation factor IX Alabama variant
31799007 | Mephentermine sulfate
31801005 | Benzonatate
31815007 | Oxybutynin hydrochloride
31818009 | Deoxyribonuclease (pyrimidine dimer)
31827005 | Ristocetin
31895006 | Gonadotrophin
31936008 | Sodium isotope
31990000 | Fibrinogen Cleveland II
32050003 | Pine oil
32083005 | Oxanamide
32120008 | Camphorated oil
32133002 | Microarazide nitrate
32154009 | Inulin
32157002 | Cathepsin B
32281001 | Fibrinogen Oslo IV
32370002 | Diprophylline
32436002 | Phentolamine mesilate
32437006 | Triphenyl phosphate
32498003 | Cortisone
32505007 | Phosphorus 32
32519007 | Charcoal medicinal vegetable
32707001 | Deoxyribonuclease I
32714004 | Magnesium dust
32757009 | Dibenzepin
32789000 | Ferritin
32800009 | Ethionamide
32824001 | Ergot alkaloid
32836007 | Sodium acetrizoate
32852005 | MSH - Beta melanocyte stimulating hormone
32898006 | Fibrinogen San Francisco
32901007 | Prostaglandin A2
33280006 | Medicinal zinc peroxide
33307008 | Sodium meralein
33396006 | Ni - Nickel
33435008 | Capillary active agent
33440000 | Ceftriaxone sodium
33447002 | Bephenium hydroxynaphthoate
33535006 | Renal hormone
33619005 | Plasminogen activator
33635003 | 5-Hydroxytryptamine
33642003 | Fibrinogen Sydney I
33667000 | Mercumatilin
33752008 | Motilin
33837008 | Aluminum glycinate
33922005 | Vitamin L
33963004 | Angiotensin 3
34007009 | Cholelitholytic agent
34070005 | Fibrinogen Nagoya
34074001 | Borate salt
34086003 | Heparin cofactor I
34113002 | Acrisorcin

34120009 | Fibrinogen Amsterdam
34198005 | Folescutol
34211007 | ^83^Strontium
34239008 | Castor oil
34274009 | Iodine pentafluoride
34332002 | Nitrophenol
34425005 | Amolanone
34471003 | ^121^Iodine
34654009 | Iodine solution
34657002 | Isopropamide iodide
34658007 | Met-enkephalin
34737006 | C1-esterase inhibitor
34915005 | Pyridostigmine bromide
34919004 | Potassium tartrate
34953000 | Colocynth
34983009 | Epicillin
35060001 | Testicular hormone
35135004 | Aglycone
35150008 | Glucocorticoid hormone
35236008 | Thenyldiamine
35257000 | Entsulfon
35281007 | Acetophenazine
35318005 | Esmolol hydrochloride
35321007 | Fludeoxyglucose F 18
35343004 | Cefonicid sodium
35431001 | Adenosine
35466004 | Relaxin
35473009 | Fibrinogen St. Louis
35605007 | Wool fat
35733004 | Fat soluble vitamin
35748005 | Wine
35765001 | Sincalide
35864006 | Parathiazine
35883004 | F - Fluorine
35903003 | Potassium bromide
35946000 | Pentolinium
35954003 | Coagulation factor II variant
35966009 | Ouabain
35976007 | PPP - Human pancreatic polypeptide
36020009 | Factor IX antibody
36021008 | Cefadroxil monohydrate
36022001 | Fibrinogen Freiberg
36136008 | Penitrem-A
36167005 | Fibrinogen Torino
36173006 | Tetraiodothyroacetic acid
36176003 | Thrombin
36235004 | Pine needle oil
36378007 | Lithium compound
36380001 | Oxyphencyclimine hydrochloride
36516003 | Mepyramine maleate
36541005 | Mercuric iodide
36651003 | Bismuth salt
36661005 | Tyrothricin
36712003 | Factor XII antibody
36872003 | Tridihexethyl
36887008 | Mineralocorticoid hormone
36953002 | Fibrinogen Nancy
37006008 | Cyclothiazide
37013008 | Dipivefrine hydrochloride
37077000 | ^122^Xenon
37078005 | Nitromersol
37123002 | Copper dust
37225000 | ^52^Manganese
37237003 | VITE - Tocopherol
37262003 | Phenyl p-aminosalicylate
37276002 | Polypeptide hormone
37352007 | Fungus antigenic agent
37365003 | Fibrinogen Rouen
37411004 | Tissue plasminogen activator
37433002 | Polycarbophil
37451001 | Opium tincture
37527009 | Sufentanil citrate

37648000 | Clindamycin phosphate
37656002 | Thiamazole
37691005 | Hetacillin
37716005 | White ointment
37751002 | Beta 2 blocking agent
37756007 | GIP - Gastric inhibitory polypeptide
37758008 | Drug-induced coagulation inhibitor
37765000 | Amfepramone hydrochloride
37951005 | Fibrinogen Paris I
37957009 | Pentoxyverine
37978007 | Nitrofurantoin sodium
37986007 | Tremorgen
37994000 | Fibrinogen Hanover
38044001 | Paromomycin
38122009 | Anisindione
38218009 | Hyaluronic acid
38245005 | Thymic hormone
38344006 | Sodium iodomethamate
38399002 | ^135^Xenon
38408002 | Paregoric
38543004 | Lissamine green B stain
38595007 | Trimethyl phosphite
38622005 | Aluminum oxide ore
38623000 | ^69^Zinc
38648002 | Mephentermine
38686006 | Colistimethate sodium
38714005 | IGF 2 - Insulin-like growth factor 2
38794009 | Molybdenum isotope
38909000 | Glutamic acid hydrochloride
38914001 | Thymol iodide
38937007 | Water in oil agent
39012008 | Thiram
39044007 | Aluminum alkyl
39123009 | Cefaloglycin
39152007 | Erythromycin stearate
39263003 | Amisometradine
39292004 | Iodine trichloride
39294003 | Iron carbohydrate complex
39428005 | Deuteroporphyrin
39514001 | Decamethonium
39546001 | Manganese isotope
39552000 | Scabicide
39808001 | Cinchocaine hydrochloride
39815009 | Clorazepate
39933002 | Pancreatic hormone
39962001 | Coagulation factor II Barcelona variant
39972003 | Na - Sodium
39973008 | C peptide
39979007 | T2 toxin
40036000 | Sulfadimidine
40115000 | Aluminum hydroxychloride
40342009 | Thiamylal sodium
40404004 | Papaverine hydrochloride
40438007 | Antimony sodium tartrate
40534007 | Aflatrem
40545005 | Amphotericin A
40601003 | Chlordiazepoxide hydrochloride
40699008 | Barium fluorosilicate
40710000 | Diodone
40789008 | ACTH - Adrenocorticotrophic hormone
40924008 | Water soluble vitamin
40937006 | ^124^Iodine
40940006 | Beta human chorionic gonadotrophin
41062004 | Methoxsalen
41067005 | Oxiconazole nitrate
41091001 | Mebutamate
41143004 | Ursodesoxycholic acid
41153003 | Amyl nitrate
41199001 | Melatonin
41261002 | Quinethazone
41332001 | Oleandomycin
41395001 | Tamoxifen citrate

41410009 | IF - Intrinsic factor
41433005 | Aluminum compound
41464007 | ^90^Molybdenum
41485007 | Leuc-enkephalin
41492002 | Satratoxins
41503000 | Zinc compound
41509001 | Potassium warfarin
41573001 | Linseed oil
41592004 | ^82^Strontium
41598000 | Estrogen
41722006 | Cefotaxime sodium
41793006 | Calcium cyanamide
41834005 | Olive oil
41903005 | Hexapradol
41945007 | Alkylpiperidine derivative of phenothiazine
41989007 | Lolitrem
42033003 | Karaya gum
42056007 | Placental hormone
42145009 | Fibrinogen Copenhagen
42146005 | I - Iodide
42159002 | Mushroom toxin
42163009 | Methylphenidate hydrochloride
42180008 | Vitamin D>2<, phosphate ester
42193003 | Stannous fluoride
42212002 | Sodium pentachlorophenate
42230009 | Bentonite
42319009 | Lipotropin
42382009 | 4-Ipomeanol
42464005 | 1,4-Benzenediamine
42520004 | Bulrush millet ergot alkaloid
42549007 | A-2M - Alpha-2 macroglobulin
42605004 | Aldosterone
42671007 | Rye ergot alkaloid
42692007 | Naproxen sodium
42710006 | Coagulation factor XI variant type II
42730005 | Periodate salt
42761001 | 1, 2-Benzenediol
42841002 | Zinc oxide
42877009 | Anhydrous borate
43004008 | Glucagon-like peptide 1
43032004 | Anabasine
43048003 | Amfomycin
43136004 | Sr - Strontium
43171004 | Ovarian hormone
43239002 | ^75^Selenium
43289005 | Dihydrofolate
43332008 | Coagulation factor IX Lake Elsinor variant
43356007 | Betaine
43440003 | Melanocyte stimulating hormone releasing factor
43462003 | Para-aminohippuric acid
43538006 | Non radiopaque medium
43541002 | Pentapiperide
43585000 | Sulfonamide diuretic
43592005 | Cactinomycin
43597004 | Chymodenin
43613009 | Phosphorous pentachloride
43621003 | Coagulation factor IX Oxford 2 variant
43688007 | Testosterone
43698001 | Hydroxystilbamidine isethionate
43706004 | VITC - Ascorbic acid
43784004 | Thymic lymphopoietin suppressing factor
43835003 | Zinc gelatin
43848004 | Ancrod
43880002 | Chloroacetamide
43897008 | ^56^Manganese
43909000 | Thiamine triphosphate
43989002 | 3,4-Methylenedioxybenzyl methyl ketone
44044007 | Calcium phosphate product
44068004 | Doxylamine
44159003 | Calcined baryta
44234001 | Strontium isotope
44262008 | Oxidized cellulose

44293009 | Phenoxybenzamine hydrochloride
44330008 | Pyrvinium pamoate
44347009 | Lergotrile
44469001 | Fibrinogen Petoskey
44508008 | Hydromorphone
44520000 | Buphenine hydrochloride
44555003 | Methylenedioxyamphetamine
44588005 | Iodine
44603007 | Iodinated glycerol
44609006 | Calcitonin gene-related peptide
44624002 | Fibrinogen New Orleans I
44644008 | Mycotoxin
44681007 | Hypothalamic inhibiting factor
44711006 | Gastrin II
44776005 | Xenon radioisotope
44908000 | Chlorzoxozone
45084007 | Fibrin degradation agent, D fragment
45119009 | Glycine salt of bile acid
45158004 | Fluorine isotope
45159007 | Azatadine maleate
45207006 | Dexamfetamine sulphate
45266004 | Alpha-2-antiplasmin
45483006 | Psilocin
45524001 | Dextranomer
45555007 | NA - Norepinephrine
45604009 | Tranquilizer
45754009 | Alpha interferon
45807004 | Coagulation factor IX variant
45940009 | Theophylline anhydrous
45946003 | Proglucagon
45960001 | Naepaine
46015007 | MSH - Melanocyte stimulating hormone
46021006 | ^91^Strontium
46058006 | Prostaglandin G2
46097003 | Lactated Ringer solution
46122001 | Antitreponemal agent
46134009 | Prostaglandin A1
46146008 | Cefotetan disodium
46191004 | Cataglykin
46201000 | Piperidolate
46225008 | CCK-PZ - Cholecystokinin-pancreozymin
46250006 | Slaframine
46257009 | Triiodomethane
46293006 | Bromocriptine mesylate
46492003 | Nivalenol
46514003 | Calcium mandelate
46558003 | Imipenem
46572007 | Factor XI
46654009 | THE - Tetrahydrocortisone
46668002 | Homatropine methylbromide
46711008 | Diglycol hydroiodide
46749007 | Medicinal iodine
46861006 | Deoxynivalenol
46887006 | Ambenonium chloride
46921009 | Quinoline dye
46943001 | (3-alpha,5-beta,20S)-3,17,20,21-Tetrahydroxypregnan-11-one
46950002 | Protriptyline hydrochloride
47026005 | Thymol oxide
47176005 | Methdilazine hydrochloride
47180000 | Metisazone
47192000 | Meglumine amidotrizoate
47218005 | Fibrinogen Giessen II
47247003 | Fibrinogen Kyoto
47280005 | Macrocyclic trichothecenes
47336007 | Fibrinogen Manchester
47349002 | beta Neoendorphin
47350002 | Pregnenolone
47355007 | Benzodiazepine nucleus
47383009 | Cephaelin
47389008 | MTBE - Methyl tert-butyl ether
47408003 | Naftifine hydrochloride
47413004 | Fat emulsion

47414005 | Trimethidinium
47500008 | Borate pentahydrate
47663000 | Clindamycin palmitate
169008 | Hypothalamic releasing factor preparation
211009 | Norethandrolone preparation
302007 | Spiramycin
439007 | Therapeutic radiopharmaceuticals
449005 | Procaine penicillin
544002 | Melphalan
669007 | Vaccinia virus vaccine
716000 | Hematological agents
796001 | Digoxin product
847003 | D-thyroxine preparation
902003 | Experimental drug
922004 | Pralidoxime
1039008 | 6MP - Mercaptopurine
1148001 | Ticarcillin
1182007 | Hypotensive agent
1206000 | Alpha 2 adrenergic blocking agent
1222004 | Metronidazole
1389007 | Beclometasone
1434005 | Calamine
1528001 | Folinic acid product
1552008 | Saluretic drug
1594006 | Azatadine
1613009 | Dexbrompheniramine maleate
1756009 | Antipruritic topical anesthetic
1758005 | Motilin preparation
1842003 | Diphemanil
1878008 | Hexachlorophane product
1887004 | Permethrin
1982006 | Bacitracin ophthalmic preparation
2016004 | Dextromethorphan
2037007 | Otic anti-inflammatory preparation
2183004 | Tetryzoline
2190009 | Benzhexol
2497003 | Hexetidine
2571007 | Busulfan
2596005 | Lincomycin
2679000 | Oxandrolone
2944000 | Depilatory agent
2949005 | Diagnostic agents
3037004 | Flumethasone preparation
3046005 | Enalapril maleate
3127006 | 5FU - Fluorouracil
3221003 | Ringer solution
3334000 | Cefotaxime
3361000 | Anti-heparin agent
3526007 | Rabies vaccine, human
3593009 | Otic anti-infective preparation
3814009 | Propylthiouracil product
3822002 | Suxamethonium
4126008 | Fluprednisolone preparation
4194004 | Mazindol
4219002 | Penicillamine
4220008 | Tolazoline
4382004 | Centrally acting antihypertensive agent
4704002 | Iothiouracil
4741001 | Sodium folate preparation
4753002 | Triamcinolone dental paste
4865009 | Prolactin releasing factor preparation
4937008 | Cefaclor
5067008 | Antithyroid drugs
5465006 | Propitocaine hydrochloride
5478006 | Trifluperidol
5487002 | Therapeutic agent
5606003 | Dexamethasone nasal preparation
5720001 | Latrodectus mactans antivenin
5737008 | Demeclocycline
5776009 | Anesthetic
5786005 | Chlorothiazide
5797005 | Clotrimazole

5924003 | Isosorbide dinitrate
5975000 | Niclosamide
6028009 | Triamcinolone preparation
6067003 | Orciprenaline
6071000 | Coal tar preparation
6102009 | Baclofen
6116004 | Oxymetholone preparation
6122008 | Class Ia antiarrhythmic drug
6232005 | Naphazoline
6247001 | Folic acid preparation
6259002 | Hydrogen peroxide solution 3%
6369005 | Penicillin -class of antibiotic-
6425004 | Antihistamine
6517004 | Butyrophenone derivative antipsychotic agent
6526001 | Nalorphine
6625006 | Zinc sulfate
6652001 | Calcium pantothenate preparation
6716006 | Abortient agent
6960003 | Polymyxin B
6985007 | Opium preparation
7092007 | Metoprolol
7140000 | Radiographic contrast media
7168001 | Magnesium carbonate
7230005 | Brucella vaccine
7235000 | Ethylenediamine derivative antihistamine
7292004 | Indocyanine green
7336002 | Trazodone
7561000 | Dexamethasone preparation
7577004 | Ciprofloxacin
7624002 | Sodium perborate product
7836006 | Expectorants
7947003 | Acetylsalicylic acid
7959004 | Teniposide
7982002 | Phenmetrazine hydrochloride
8028001 | Butacaine
8109005 | Alimemazine
8163008 | Nitroprusside
8348002 | Cyclopentolate
8372007 | Promethazine
8416000 | Dicloxacillin
8571001 | Vasoconstrictor
8658008 | Human serum albumin preparation
8661009 | Replacement preparation
8692006 | Metamfetamine
8696009 | Antispasmodic
9190005 | Tropicamide
9268004 | Butabarbitone
9307009 | Animal serum globulin preparation
9500005 | MAOI - Phenelzine
9542007 | Antihepatitis B immunoglobulin
9690006 | Nikethamide
9745007 | Sucrose product
9778000 | Cytomegalovirus immune globulin
9944001 | Chlorphenamine
10099000 | Ketoprofen
10135005 | Cinchona alkaloid
10243007 | Benzoic and salicylic acid ointment
10312003 | Prednisone preparation
10332002 | Vaccine - bacterin - toxoid
10355004 | Injectable vitamin preparation
10356003 | Undecylenic acid and undecylenate salt
10368007 | Colloidal oatmeal powder
10422008 | Nitropentaerythrol
10504007 | Doxycycline
10515002 | Lututrin preparation
10555000 | Tocainide
10632007 | Multivitamin preparation
10668002 | Skin anti-infective
10712001 | Glucagon product
10756001 | Haloperidol
10784006 | Anti-psychotic agent
11185009 | Autogenous vaccine - autogenous bacterin

11260008 | Ophthalmic anti-infective preparation
11402001 | Medicinal enzyme
11430001 | Tetracyclic antidepressant drug
11563006 | Vitamin D preparation
11719000 | Cetylpyridinium
11777007 | Ophthalmic antibiotic
11783005 | Stool softener
11796006 | Methysergide
11841005 | Doxepin
11847009 | Naproxen
11866009 | Plague vaccine
11959009 | Procainamide
12096000 | Nystatin
12236006 | Pancreatin product
12289007 | Whole blood product
12335007 | Diatrizoate
12369008 | Oxytocin product
12425002 | Human white blood cell product
12436009 | Vinblastine
12495006 | Magnesium citrate
12512008 | Triamterene
12559001 | Emetine
12657005 | Fludrocortisone acetate preparation
12839006 | Estradiol product
12968008 | Vitamin products
13132007 | Dextran product
13154009 | Chlormethine
13178004 | Rickettsial vaccine
13252002 | Salsalate
13414000 | Cefadroxil
13432000 | Nortriptyline
13512003 | Minocycline
13525006 | Acetylcholine preparation
13565005 | Bisacodyl
13592004 | Pyrazinamide
13664004 | BAL - British Anti-Lewisite
13790009 | Oral form iron preparation
13800009 | Naftifine
13813003 | Biotin preparation
13929005 | Spironolactone
13936006 | Butorphanol
13965000 | Valproic acid
14054003 | Vitamin B complex preps
14103001 | Opiate antagonist
14170004 | Capreomycin
14542005 | Caffeine and sodium benzoate injection solution vial
14601000 | Anticholinergic agent
14706000 | Pheneticillin
14728000 | Chloroquine
14745005 | Hepatitis A vaccine
14814001 | Trimethobenzamide
14816004 | Cocaine product
15117003 | Coagulant
15222008 | Enalapril
15375005 | Phenanthrene derivative
15383004 | Levodopa
15389000 | Hydantoin derivative anticonvulsant
15432003 | Ethinylestradiol
15772006 | Beta 1 blocking product
15857002 | Ethanolamine derivative antihistamine
16031005 | Dexchlorpheniramine
16037009 | Terfenadine product
16047007 | Benzodiazepine
16131008 | Animal antiserum
16403005 | Non-steroidal anti-inflammatory agent
16602005 | Hydrocortisone preparation
16787005 | Streptococcus equisimilis - suis antiserum
16791000 | Cefradine
16832004 | Conjugated estrogens
16858004 | Urea
16867004 | Sulfathiazole
16970001 | Proguanil

16977003 | Lithium carbonate
17016006 | Animal gamma globulin
17055007 | Systemic acidifier
17308007 | Dapsone
17386008 | Anti-infective agent
17502009 | Gallamine triethiodide
17554004 | Paramethasone preparation
17558001 | Corn oil
17600005 | Diagnostic radiopharmaceuticals
17805003 | Lithium citrate
17859000 | Polyvalent crotalidae antivenin
17893001 | Skeletal muscle relaxant
18002004 | Auranofin
18125000 | Fluocinonide
18335001 | Aureolic acid
18340009 | Oxychlorosene
18381001 | Prindolol
18511007 | Generic drug
18548003 | Methylphenidate
18679008 | Potassium-removing resin
18811003 | L-asparaginase preparation
18914005 | Hydroflumethiazide
18952006 | Econazole
19194001 | ddI
19225000 | Lorazepam
19232009 | Prilocaine
19261005 | Sulfinpyrazone
19315007 | Pyridoxine hydrochloride preparation
19403009 | Flurazepam
19405002 | Netilmicin
19581007 | Parasympathomimetic agent
19583005 | Diclofenamide
19630009 | Silver sulfadiazine
19768003 | Alkylating drugs
19841008 | Ceftriaxone
20091003 | Somatotropin releasing factor preparation
20201001 | Nafoxidine
20237006 | Dihydrotachysterol preparation
20249007 | Progestin preparation
20303001 | Vinblastine sulfate
20320002 | Hydrocodone
20577002 | HCG - Human chorionic gonadotrophin product
20865003 | Diflunisal
20969008 | Polyvalent snake antivenin
21069002 | Lipotropic agent
21159006 | Pargyline
21451004 | Indium-113m chloride
21691008 | Magnesium trisilicate
21701005 | Cromoglicic acid
21767006 | Iron dextran complex
21788002 | Phenazopyridine hydrochloride
21986005 | Erysipelothrix rhusiopathiae antiserum
22091006 | Hormone preparation
22168005 | Glycopyrronium
22198003 | Metolazone
22274004 | Methandriol preparation
22474002 | Aldosterone preparation
22587006 | Depolarizing muscle relaxant
22657006 | Calcitonin product
22672005 | Amfetamine group
22696000 | Hydralazine
22801004 | Sterile water solution
22826008 | Ammonia detoxicant
22969001 | Oxytetracycline
23079006 | Vincristine
23222006 | Antisera
23343005 | Human platelet product
23532003 | Phenmetrazine
23822003 | Human plasma fraction
23827009 | Sulfacetamide
23838009 | Nonspecific alpha-adrenergic blocking agent
23888001 | Cascara

24036003 | Antianemia drug
24450004 | Ethambutolol
24504000 | Methylcellulose
24866006 | Salmonella typhimurium antiserum
24891006 | Sodium amidotrizoate
25014009 | Tripelennamine
25057004 | Mercurial diuretic product
25076002 | Carisoprodol
25085002 | Cholecystokinin preparation
25142008 | Trilostane
25246002 | Allopurinol
25398003 | Ichthammol
25419009 | Barium sulfate
25465007 | Escherichia coli monoclonal antibody
25673006 | Omeprazole
25716001 | Tetanus toxoid, human
25849001 | SNP - Sodium nitroprusside
25860005 | Terconazole
25912006 | Menadiol sodium diphosphate preparation
25995007 | Triprolidine
26122009 | Dimetindene
26124005 | Glipizide product
26244009 | Antimuscarinic
26303005 | Hexestrol preparation
26370007 | Hemostatic agent
26456008 | Levothyroxine sodium preparation
26458009 | Diphenhydramine
26462003 | Cyproheptadine
26503009 | Deserpidine
26523005 | Dobutamine
26548008 | Pancreatic hormone product
26574002 | Droperidol
26580005 | Digestant
26736008 | Ferrous gluconate
26800000 | Midazolam
26842003 | Burbot liver oil preparation
27035007 | Heavy metal antagonist
27085003 | Sodium polystyrene sulphonate
27196008 | Bupivacaine
27242001 | Methylprednisolone preparation
27479000 | AZT - Azidothymidine
27518004 | Parathyroid hormone
27548001 | LA - Local anesthetic drug
27566006 | Drug vehicle preservative
27638005 | T-PA - tissue-type plasminogen activator
27658006 | Amoxicilline
27707001 | Medroxyprogesterone acetate preparation
27754002 | Pyrazinobutazone
27867009 | Antineoplastic
28028002 | Synthetic hormone preparation
28149003 | Pentostatin
28235004 | Doxapram
28240007 | Eye cosmetic
28410007 | alpha-Carboxypenicillin
28415002 | Methscopolamine
28426008 | Fluocinolone
28506006 | Flucytosine
28748001 | Chloral hydrate
28841002 | Anhydrohydroxyprogesterone
28906000 | Percoid liver oil preparation
29051009 | Loop diuretic product
29058003 | Halcinonide
29089004 | Dibromomannitol
29121001 | Mersalyl
29129004 | Oxymetazoline
29156002 | Mechlorethamine
29175007 | Rifampicin product
29237004 | Arachnida antivenin
29439004 | Captopril
29620001 | beta-Tocopherol preparation
29840005 | Amoxapine
29877002 | MAOI - Isocarboxazid

29883004 | Class IV antiarrhythmic
29896003 | Betamethasone
29987004 | Vitamins A and D preparation
30010009 | Cyanocobalamin product
30125007 | Senna product
30306003 | Vitamin B1 preparation
30317002 | Cisapride
30427009 | Erythromycin product
30466001 | Chloramiphene
30492008 | Diuretic
30666008 | Toxoid (human)
30729008 | Iron preparation
30761007 | Mannitol product
30964009 | Methyprylone
30988006 | Dienestrol preparation
31025003 | Dibenzoxazepine derivative antipsychotic agent
31087008 | Ampicillin
31181006 | Injectable testosterone preparation
31231007 | H2O2 - Hydrogen peroxide
31305008 | Streptococcus equisimilis antiserum
31306009 | Quinidine
31352002 | Normal colostral whey
31587003 | Almond oil liquid
31684002 | Buprenorphine
31690003 | Bethanechol
31692006 | Pentamidine
31716004 | Frozen plasma product, human
31865003 | Fluconazole
31872002 | Pramocaine
31970009 | Nitrate vasodilator
32249005 | Antiviral drugs
32313007 | Enflurane
32460003 | Beclometasone dipropionate nasal preparation
32462006 | Melanocyte stimulating hormone releasing factor preparation
32473004 | Spider antivenin
32474005 | Probucol
32583002 | Antiseborrheic drug
32647002 | Ergotamine
32653002 | Ergosterol preparation
32792001 | Trimethoprim
32823007 | Maprotiline
32841004 | NaCl - Sodium chloride solution
32955006 | Domperidone
32960005 | Thiosalicylate
33124007 | Butamidum
33133009 | Ophthalmic anti-inflammatory preparation
33219003 | Tricyclic antidepressant
33231001 | Pentobarbital
33234009 | Var/Vac
33236006 | Cortisone acetate preparation
33252009 | beta-adrenoceptor blocking drug
33378002 | Desipramine
33484000 | Proprietary drug
33588000 | Thioridazine
33589008 | Cardiac glycoside
33664007 | Acetazolamide
33675006 | Carbachol
33682005 | Mydriatic
33815001 | Streptococcus suis antiserum
34012005 | Sulfonylurea product
34217006 | Oxyquinoline
34364009 | Mefenamic acid
34393009 | Tolazamide
34462007 | Natamycin
34598001 | Immunobiologic agent available from the Centers for Disease Control and Prevention
34599009 | Thyroglobulin preparation
34689006 | Hepatitis B vaccine
34693000 | Zalcitabine
34696008 | Sodium ascorbate preparation
34731007 | Carbenicillin
34816007 | Cod liver oil
34833000 | Hydrocortisone ophthalmic preparation

34929006 | Benzethonium
35035001 | Orphenadrine
35063004 | Ribavirin
35097006 | Flea antigen
35282000 | Gemfibrozil
35300007 | Daunorubicin
35324004 | Paraldehyde product
35392005 | Calcium-removing resin
35476001 | Silver nitrate product
35531004 | Hydrocortamate preparation
35736007 | Cholera vaccines
35768004 | Oxybutynin
35792007 | Peritoneal dialysis solution
35967000 | Medazepam
35983000 | Cellular blood product, human
36068003 | Pyrantel
36113004 | Imipramine
36168000 | Anti-infective nasal preparations
36218003 | Thiethylperazine
36236003 | Antidepressant
36391008 | Primaquine
36537006 | Ambenonium
36594001 | Tiabendazole
36621009 | Medroxyprogesterone
36642006 | Propantheline
36763003 | Immune globulin IM
36795002 | Piperazine estrone sulfate
36893000 | Ceftazidime
36909007 | Phenindamine
36940000 | Thioxanthene derivative antipsychotic agent
36944009 | Vitamin B6 preparation
37084008 | 4-quinolones
37146000 | Typhus vaccine
37306000 | Vidarabine
37400007 | Magnesium sulfate
37628007 | Cefalotin
37643009 | Autogenous bacterin
37773009 | Diphenylbutylpiperidine derivative antipsychotic agent
37803001 | d-TC - Tubocurarine
38076006 | Sodium thyroxine
38131009 | Synthetic drug
38166006 | Tolnaftate
38181000 | Antineoplastic antibiotic
38231004 | Polysaccharide iron complex
38268001 | Propionic acid-ibuprofen
38314008 | Isotretinoin
38413003 | Otic dosage form product
38578004 | Megestrol preparation
38828006 | Sodium thiosulfate
38900001 | Proliferant agent
39064002 | Acetohexamide
39124003 | Methohexital
39128000 | Famotidine
39142008 | Phendimetrazine
39252001 | Water balance agent
39359008 | Penicillin V
39432004 | Deodorant
39487003 | Regular insulin product
39516004 | Disulfiram product
39598004 | Marijuana derivative
39608003 | Pentazocine
39707000 | Aminobenzoic acid
39741008 | Drug groups primarily affecting the gastro-intestinal system
39860005 | Fructose product
39939003 | Phenyltoloxamine
40162007 | Bulk-forming laxatives
40232005 | Ketoconazole
40339003 | Calcium lactate
40429005 | Etomidate
40430000 | Bromelains
40556005 | Phenytoin product
40562000 | Methylergonovine preparation

40589005 | Amitriptyline
40606008 | Orphan drug
40648001 | Fentanyl product
40652001 | Cromolyn sodium
40820003 | Carbamazepine product
40877002 | Streptomycin
40901001 | Desferrioxamine mesylate
40905005 | Beractant
40974005 | Phenylpiperone
40999006 | CCNU - Lomustine
41001009 | Dinoprost
41015006 | Metaraminol
41147003 | Perphenazine
41193000 | Aciclovir
41324009 | Propiomazine
41365009 | Fluphenazine
41367001 | Enterogastrone preparation
41399007 | Oxazolidinedione
41470001 | Corbadrine
41493007 | Dicycloverine
41549009 | Angiotensin-converting enzyme inhibitor
41985001 | Bitolterol
42082003 | Vancomycin
42098005 | Dexamethasone ophthalmic preparation
42271003 | Glutamic acid product
42348003 | Methyltestosterone
42374009 | Dextromethorphan hydrobromide
42444000 | Secobarbital
42514000 | Procaine product
42638008 | Methylrosanilium chloride
42714002 | Escherichia coli antiserum
42720001 | Miconazole
42863006 | Desmopressin acetate preparation
43343000 | Magaldrate
43499003 | Injectable interferon
43533002 | Chloramphenicol ophthalmic preparation
43594006 | Skin bleaching preparations
43684009 | Misoprostol product
43747001 | Drug excipient
43753001 | Dydrogesterone
43879000 | Flunisolide
43927002 | Analeptic agent
44175000 | Diperodon
44418001 | Percomorph liver oil preparation
44658005 | Promazine
44731005 | Otic form hydrocortisone
44790008 | Ethosuximide
44798001 | Dinoprostone product
44938006 | Cefoperazone
44990002 | Procyclidine
45218006 | Meclastine
45311002 | Terbutaline
45313004 | Propylpiperazine derivative of phenothiazine
45355008 | Rocky Mountain spotted fever vaccine
45518007 | Thiazide diuretic product
45680002 | Tolmetin
45844004 | Sulfasalazine
45888006 | gamma-Tocopherol preparation
45923000 | E331 - Sodium citrate
46009007 | Chlorambucil
46063005 | Psychotherapeutic agent
46123006 | Ascorbic acid product
46233009 | FLU - Influenza vaccine
46436003 | Haloprogin
46479001 | Encainide
46532003 | Brilliant green
46547007 | Labetalol
46576005 | Flecainide
46709004 | Methylphenobarbital
46741005 | SA - Salicylic acid
46913003 | Edrophonium
47065008 | Quinine

47120002 | Hexamidinum
47124006 | Aminoglutethimide
47140009 | Medrysone
47331002 | CPZ - Chlorpromazine
47527007 | Phenindione
47602007 | Nalidixic acid
47755009 | Potassium sparing diuretic
47772002 | Fenfluramine hydrochloride
47898004 | Verapamil
48028008 | Haemophilus influenzae Type b (Hib)/diphtheria/tetanus/pertussis vaccine
48174005 | Ranitidine
48256008 | Benzyl benzoate
48279009 | Emollient product
48351000 | Butadione
48546005 | Diazepam product
48603004 | Warfarin
48614001 | Clobetasol
48647005 | Pancrelipase
48698004 | Calcium-channel blocker
48836000 | Amikacin
48875009 | Dihydroergotamine product
48899009 | Hyoscyamine
49019002 | Prednisolone ophthalmic preparation
49157004 | Uricosuric product
49192007 | Antiflatulent agent
49267000 | Hydroxyphenylbutazone
49299006 | Protriptyline
49485009 | Norfloxacin
49577002 | Minoxidil
49590009 | Class I antiarrhythmic
49600008 | Normal animal plasma
49617001 | Disodium enoxolone
49663007 | Sunscreening preparations
49669006 | Escherichia coli - pasteurella multocida - salmonella typhimurium antiserum
49688004 | Hexocyclium
49694007 | Mucolytics
49705006 | Idoxuridine
49802003 | Anti-D (Rh) immunoglobulin
49860008 | Gold product
49953001 | Pheniramine
49992008 | Dexamethasone sodium phosphate
50094009 | Hetastarch
50137001 | Glycine solution
50182002 | Hemodialysis fluid
50244000 | Intravenous anesthetic
50256004 | Antitrichomonal agent
50318003 | Luteal hormone
50335004 | Levorphanol
50520001 | Neomycin B
50841004 | Otic form chloramphenicol
50868004 | Dexamfetamine
51013009 | Sulfadimethoxine
51073002 | Phenobarbital product
51103004 | Hypertonic saline
51126006 | Benzestrol preparation
51132001 | Hyaluronidase product
51326002 | BCNU - Carmustine
51334008 | Cycloserine
51361008 | Amantadine
51425004 | Scorpion antivenin
51555009 | Desiccated liver preparation
51752005 | Chologon
51758009 | Amidine
51779009 | Cephalosporin -class of antibiotic-
51908007 | Prenylamine
51992002 | Gastrin preparation
52017007 | Antiemetic
52108005 | Ferrous fumarate
52215008 | Desonide
52388000 | Prednisolone
52412009 | Methylergonovine maleate preparation
52423008 | Tar preparation

52451004 | Dihydroindolone derivative antipsychotic agent
52489009 | Vasopressin tannate preparation
52628005 | Typhoid-paratyphoid vaccine
52761007 | Anticholinesterase parasympathomimetic
52883001 | para-Hydroxyamphetamine
52896000 | Chinoform
53009005 | Analgesic
53480001 | Phentermine
53584007 | Methacholine
53640004 | Fluoxetine
53641000 | Flavoxate
53691001 | Calcium gluconate
53793005 | Escherichia coli antibody
53800008 | Dithranol
53848009 | Metyrapone
53855006 | Neostigmine bromide
54142000 | Domiphen
54344006 | Flurbiprofen
54391004 | Levamisole
54406003 | Methoxamine
54541002 | Ergometrine preparation
54544005 | Pethidine
54577009 | Ceftizoxime
54659006 | Thiamine hydrochloride preparation
54705000 | Temazepam
54765002 | Phenylephrine
54824008 | Isometheptene
54882005 | Amfepramone
54887004 | Cefalexin
54972005 | Tretinoin
54982006 | Promethestrol preparation
55015008 | E325 - Sodium lactate
55217007 | Calcium carbonate
55432002 | Azlocillin
55556000 | Tetracaine
55673009 | Iothalamate sodium I^125^
55745002 | Propranolol
55782006 | Amide-type anesthetic
55784007 | Natural hormone preparation
55830003 | Menotrophin
55867006 | Aminophylline
55889005 | Praziquantel
56011005 | Stinging insects immunotherapy extract
56014002 | Hydroxyprogesterone
56032002 | Androstanolone
56059005 | Mebendazole product
56123002 | Hexamine
56230001 | Bretylium
56234005 | Growth hormone product
56240003 | Paramethasone acetate preparation
56281006 | Anesthetic ether
56480005 | Parabromdylamine
56549003 | Metoclopramide
56602009 | Hydroxycarbamide
56615000 | Polyestradiol phosphate preparation
56656002 | Magnesium-based antacid
56844000 | Yellow fever vaccine product
56928005 | Etoposide product
56934003 | PVP - Polyvinylpyrrolidone
57002000 | Chlorprothixene
57066004 | Cisplatin product
57123008 | Urinary acidifier
57191001 | Chloramphenicol product
57263002 | Oxiconazole
57376006 | Monosodium carbonate
57538001 | Chlortetracycline
57616006 | STD - Sodium tetradecyl sulfate
57670008 | Cefoxitin
57752001 | Gentamicin product
57811004 | Dihydrocodeine
57819002 | Somatotropin inhibiting factor preparation
57845006 | Isoprenaline

57853003 | Clidinium
57893000 | Chlortalidone
57952007 | Lipid lowering drug
58050004 | Antiparkinsonian agent
58098008 | Tetanus toxoid adsorbed
58360000 | Phenazocine
58467001 | Papaverine
58502006 | Propylamine derivative antihistamine
58760003 | Antimetabolites
58805000 | Pituitary hormone preparation
58883005 | Clindamycin
58892008 | F3T - trifluorothymidine
58905004 | Diazoxide
58944007 | Vasodilator
59057006 | Antihemophilic factor preparation
59187003 | Dopamine product
59240002 | MMC-Mitomycin-C
59245007 | Hypophysiostatic factor preparation
59255006 | Sulfonamide
59261009 | Oral contraceptive preparation
59270007 | Loxapine
59456005 | Astemizole
59589008 | Pyrimethamine product
59594008 | Nondepolarizing agent
59751001 | Cough suppressant
59799001 | Osmotic diuretic product
59941008 | Diltiazem
59953007 | Pyridostigmine
59999009 | Tetanus and diphtheria toxoid adsorbed for adult use
60090003 | Soft drug
60093001 | Phenylpropanolamine hydrochloride
60149003 | Indometacin
60169008 | Antacid
60468008 | Magnesium hydroxide
60520004 | Thiamine mononitrate preparation
60533005 | Astringent drug
60541005 | Lanatoside C
60682004 | Echothiophate
60731009 | Diethylcarbamazine
60767002 | Normal animal serum
60881009 | Heroin
60972002 | Irrigating solution
60978003 | Barbiturate
61020000 | Thyroid preparation
61093008 | Prolactin inhibiting factor preparation
61123001 | Amobarbital sodium
61132004 | Gas gangrene antitoxin
61153008 | Measles, mumps and rubella vaccine
61181002 | Meprednisone preparation
61408004 | Molindone
61457001 | Adrenal hormone preparation
61602008 | Pertussis vaccine
61621000 | Laxative
61623002 | Buclizine
61650007 | Niacinamide preparation
61651006 | Cefamandole
61862008 | Meticillin
61939007 | Diphenylmethane laxative
61946003 | Estrogenic preparation
62051009 | Dichlorisone preparation
62063004 | Nasal anti-inflammatory preparation
62191005 | Injectable liver preparation
62288001 | Anthelmintic
62294009 | Varicella-zoster immune globulin
62353003 | Clomifene citrate
62354009 | Bronchitis vaccine
62529008 | Tiotixene
62560008 | Fluorometholone ophthalmic preparation
62782004 | Clonidine
63094006 | Anticonvulsant
63136007 | Phytonadione preparation
63318000 | Benzoic acid

63338004 | Drug flavoring
63470003 | Fluoxymesterone preparation
63639004 | Niacin preparation
63682003 | Halothane
63758001 | Norethindrone preparation
63822004 | Vitamin E preparation
64083002 | Genitourinary smooth muscle relaxant
64115004 | Amodiaquin
64127001 | Dactinomycin
64240003 | Methandrostenolone preparation
64296005 | Staphylococcus toxoid
64349004 | Griseofulvin
64462001 | Terpin
64558005 | Metixene
64851009 | Diiodohydroxyquin
64878006 | Methylthiouracil
65020006 | Benzocaine
65026000 | Ephedrine
65041000 | Biperiden
65092008 | Chloropyrilene
65265006 | Thromboplastin
65461004 | Adrenal cortex extract
65484006 | Prostacycline PGI>2< preparation
65502005 | Epinephrine preparation
65505007 | Gastrointestinal smooth muscle relaxant
65627005 | 4-Amino-2-methyl-1-naphthol hydrate preparation
65628000 | Skin agent
65638005 | Corticotropin zinc hydroxide preparation
65695006 | General inhalation anesthetic
65823007 | Protective agent
65884003 | Antrapurol
65965000 | Micrurus fulvius antivenin
66065008 | Saline laxative
66094001 | Probenecid
66125007 | Flunisolide nasal preparation
66243004 | Norethindrone acetate preparation
66261008 | Tetracycline
66349002 | Androgenic preparation
66441000 | Pantothenic acid preparation
66492008 | Isoflurane
66493003 | Theophylline
66602007 | Stanozolol preparation
66672008 | Ester type local anesthetic
66742008 | Pigmenting agent
66854004 | Betamethasone valerate
66859009 | Dipyridamole
66860004 | Aluminum chloride
66919007 | Methyclothiazide
66971004 | Colestipol
67213005 | Lymphocyte immune globulin
67423003 | Acylaminopenicillin
67440007 | Alpha-adrenergic blocking agent
67477004 | Valproate
67507000 | Antiarrhythmic drug
67585005 | Topical anesthetic
67735003 | Paclitaxel
67891001 | Second generation cephalosporin
67939000 | Apomorphine
68088000 | Acebutolol
68113000 | Hib oligosaccharide conjugate (diphtheria CRM>197< protein conjugate) vaccine
68206008 | Calcitriol preparation
68208009 | Diphtheria and tetanus toxoid adsorbed
68395000 | Calcium chloride
68398003 | Somatomedin preparation
68402007 | Carbonic anhydrase inhibitor
68407001 | Hydrogen peroxide solution 30%
68422006 | Cloxacillin
68424007 | Isoflurophate
68444001 | Doxorubicin
68461003 | Surgical jelly
68490009 | Sodium propionate
68622003 | Secretin preparation

68702006 | Sodium aurothiomalate
68774008 | Isoxsuprine
68864001 | Polyvalent reptile antivenin
68887009 | Methotrexate
68892006 | Penicillinase-resistant penicillin
69107004 | Dantrolene
69204002 | Guanadrel
69227005 | Class II antiarrhythmic drug
69236009 | Amiodarone
69242008 | Miotic
69331001 | Aminocyclohexylpenicillin
69431002 | Immunosuppressant
69509008 | Biological agent
69576000 | Menadione preparation
69708003 | Clonazepam
69801001 | Sodium estrone sulfate preparation
69879000 | Altretamine
69918003 | Aztreonam
69967001 | Sucralfate
70047000 | Sulfamethoxazole
70216000 | Undecylenate salt
70254000 | Sulfamethizole
70343008 | Piperazine derivative antihistamine
70379000 | Sodium chloride
70447008 | Polyvalent pneumococcal vaccine
70460002 | Fish liver oil preparation
70702006 | Deferoxamine
70776005 | Pemoline
70841003 | Chymotrypsin preparation
70864001 | Meprobamate
70934008 | Demecarium
71181003 | Vaccine
71289008 | Snake antivenin
71338005 | Vasopressin preparation
71451001 | Kanamycin
71453003 | Mupirocin
71455005 | Flurandrenolone preparation
71462001 | Podophyllum resin
71516007 | Ergocalciferol preparation
71584004 | Sodium phosphate
71634000 | Chlormezanone
71699007 | Trifluoperazine
71724000 | Ferrous sulfate
71731001 | Medrysone ophthalmic preparation
71759000 | Nitroglycerin
71770007 | Monoamine oxidase inhibitor
71798005 | Fenoprofen
71837009 | Cytotoxic agent
72312007 | Cyclandelate
72398005 | Ophthalmic sulfonamide preparation
72416006 | Metacycline
72623000 | 2-Amino-6-mercaptopurine
72824008 | Cholestyramine
72870001 | Scopolamine
72924009 | Clofazimine
72968006 | Sodium salicylate
72996000 | Drug builder
73074003 | Colistin
73093001 | Neomycin
73133000 | Colchicine
73170009 | Menthol
73212002 | Adipiodone
73228007 | Non-steroidal estrogen preparation
73274006 | Human plasma cryoprecipitate
73277004 | Mecamylamine
73454001 | Desmopressin preparation
73572009 | Morphine
73585007 | Phenytoin sodium
73627008 | Caloric agent
73639000 | Prescription drug
73647000 | Dipivefrine
73756003 | Amobarbital

73763003 | Extended spectrum penicillin
73805002 | Thyrotropin releasing hormone preparation
73949004 | Atropine
73986003 | Cefuroxime
74022005 | Mepenzolate
74065006 | Prazepam
74074008 | Atracurium
74147005 | Edetate disodium
74213004 | Indapamide
74226000 | Vitamin K preparation
74470007 | Cyclophosphamide
74480006 | Potassium supplement
74575000 | Piperacillin
74583006 | Hydroquinone product
74626007 | Drug diluent
74632002 | Succinimide
74674007 | Propofol
74771007 | Phenoxybenzamine
74780007 | Anthraquinone laxatives
74782004 | Naturally occurring alkaloid
74798006 | Pipenzolate
74819009 | Acetohydroxamic acid
75029008 | Desoxycorticosterone preparation
75201000 | Bacterin - toxoid
75203002 | Mometasone
75292001 | Vincristine sulfate
75366006 | Sodium calciumedetate
75429004 | Dexbrompheniramine
75501004 | Bromazine
75661008 | delta-Tocopherol preparation
75748004 | Theophylline sodium glycinate
75770001 | Urea topical preparation
75927008 | Floxuridine
75959001 | Tamoxifen
75969007 | Gonadotropin releasing factor preparation
76058001 | Prazosin
76155001 | Iopanoic acid
76286000 | Gallamine
76289007 | Xylometazoline
76385003 | Alpha 1 adrenergic blocking agent
76390000 | Practolol
76591000 | Bleomycin
76696004 | Noscapine
76759004 | Disopyramide product
76962009 | Iproniazid
77035009 | Clofibrate
77048008 | Dip/ser
77237006 | Emetic
77331005 | Normal animal globulin concentrate
77390008 | Benzatropine
77398001 | Antidiarrhea agent
77549006 | Terpene
77657003 | Illegal drug
77731008 | N-Acetylcysteine
77750008 | Dacarbazine
77856005 | Esmolol
77885004 | Mestranol product
77898008 | Simethicone
78025001 | Ganciclovir product
78057000 | Carbol-fuchsin topical solution
78174002 | Mezlocillin
78379001 | Reserpine
78411008 | Thiopental sodium
78439009 | Citrate salt
78449007 | Nitrazepam
78507004 | Benzylpenicillin
78542000 | Potassium citrate product
186002 | HLA-Cw9 antigen
217008 | Blood group antigen IH
476005 | CD1b - Cluster of differentiation antigen 1b
501001 | Blood group antibody Sf^a^
505005 | Blood group antibody M'

515004 | Blood group antigen Giaigue
576007 | Blood group antibody Duck
584006 | Blood group antibody Wr^b^
593007 | Blood group antibody Holmes
704006 | Blood group antigen Rx
735000 | Blood group antigen Jobbins
819002 | Lytic antibody
876000 | D antigen
923009 | Complement component 2
1097007 | Blood group antigen M^A^
1160000 | Antibody to antigen in ISBT system LU
1223009 | Blood group antigen Marks
1273007 | Blood group antibody Evelyn
1319003 | Blood group antibody K18
1371006 | Blood group antigen Big
1405004 | Blood group antibody M^e^
1408002 | Blood group antibody 1123K
1506001 | CH1 (ISBT symbol)
1517000 | HLA - Human leukocyte antigen B21
1565004 | Blood group antibody Buckalew
1634002 | Blood group antigen Ven
1649005 | Blood group antigen Sul
1676005 | Blood group antibody LW^ab^
1681001 | Blood group antibody BLe^b^
1696002 | 12-Hydroperoxy eicosatetraenoic acid
1975007 | Blood group antibody Niemetz
2038002 | Blood group antibody Bg^b^
2082006 | Vitronectin receptor alpha chain
2147008 | Blood group antigen Paular
2260005 | Human leukocyte antigen DRw18
2329007 | Blood group antibody Vel
2404002 | Blood group antibody St^a^
2431004 | Blood group antibody Friedberg
2444009 | Human leukocyte antigen Dw25
2500009 | Lymphocyte antigen CDw41b
2568004 | Blood group antigen McAuley
2611008 | Blood group antibody La Fave
2676007 | C3(H20)
2706001 | Blood group antigen Vennera
2721007 | Blood group antigen McC^f^
2728001 | Lewis system antigen
2753003 | Blood group antibody M>1<
2765004 | SC3 (ISBT symbol)
2988007 | Human leukocyte antigen Aw
3131000 | Blood group antigen Middel
3187008 | Blood group antigen Nielsen
3555004 | Blood group antigen Morrison
3588006 | Complement enzyme
3602003 | Warm antibody
3730007 | Blood group antigen Lanthois
3807007 | RH4 (ISBT symbol)
3897001 | MNS30 (ISBT symbol)
3961009 | Blood group antigen Ritherford
3976001 | Blood group antigen HEMPAS
4115002 | Blood group antibody Norlander
4167003 | PECAM-1 - Platelet endothelial cell adhesion molecule-1
4169000 | Blood group antibody Le^bH^
4314009 | Blood group antibody Allchurch
4393002 | Blood group antigen Fedor
4401009 | Blood group antibody H>T<
4422003 | Blood group antigen
4425001 | Blood group antibody Binge
4555006 | Blood group antibody Rils
4564001 | Blood group antibody Sisson
4582003 | Blood group antigen N^A^
4591004 | MNS22 (ISBT symbol)
4761007 | CD30 - Cluster of differentiation antigen 30
4762000 | Platelet antigen HPA-3b
4789005 | Blood group antibody Bultar
4878009 | Human leukocyte antigen Dw3
4933007 | Leukocyte M1
5004004 | Blood group antibody Panzar

5064001 | Blood group antibody D 1276
5081005 | RH31 (ISBT symbol)
5094007 | Blood group antigen Rios
5250008 | Thymus-independent antigen
5259009 | Blood group antigen Braden
5307001 | Blood group antigen Hamet
5340005 | Blood group antigen Swietlik
5404007 | Lymphocyte antigen CD45RA
5439007 | Blood group antibody Do^a^
5504009 | Blood group antigen Fuerhart
5533005 | Blood group antibody Kp^a^
5573004 | Blood group antigen Oca
5637003 | HLA-DQw6 antigen
5700002 | Blood group antibody Gomez
5757007 | HLA-Cw8 antigen
5840001 | Blood group antibody Wj
5915007 | Blood group antigen Gladding
5977008 | Blood group antigen Bullock
6056004 | Blood group antibody Wk^a^
6068008 | Blood group antigen Mil
6091007 | Blood group antibody L Harris
6115000 | Blood group antibody Anuszewska
6135004 | Blood group antigen Duck
6138002 | Blood group antigen Le Provost
6170002 | Heat labile antibody
6178009 | Lymphocyte antigen CD63
6197009 | Blood group antigen Zd
6310003 | Particulate antigen
6360009 | Kallidin II
6411000 | Interleukin-12
6422001 | HLA-DRw14 antigen
6507009 | Blood group antigen Much
6529008 | Blood group antigen Cl^a^
6532006 | Macrophage activating factor
6592009 | HLA-Dw12 antigen
6642000 | Opsonin
6671004 | Blood group antigen Caw
6741004 | Anti DNA antibody
6755007 | TL antigen
6945001 | Blood group antigen Le^bH^
6958000 | Blood group antibody Frando
7029006 | Blood group antigen Greenlee
7120007 | Antigen
7158006 | HLA-Dw19 antigen
7161007 | Complement component C2a
7211005 | Blood group antibody Haakestad
7337006 | Blood group antibody Tr^a^
7411007 | Blood group antibody HLA-B8
7489000 | Homocytotropic antibody
7588005 | Blood group antibody Sk^a^
7616007 | Blood group antibody Pruitt
7648006 | HLA-Bw70 antigen
7675004 | Blood group antigen Towey
7685003 | Blood group antibody Bg^c^
7738004 | HLA-B49 antigen
7948008 | Reed-Sternberg antibody
7974002 | Blood group antibody Dalman
8002004 | Blood group antibody Fleming
8025003 | Blood group antibody Gibson
8048008 | Blood group antigen Th
8123007 | Blood group antibody Schuppenhauer
8164002 | Cluster of differentiation antigen 67
8268005 | Blood group antibody Hildebrandt
8343006 | Blood group antibody Re^a^
8362009 | Anti c
8376005 | Duffy blood group antibody
8452001 | Blood group antigen Sisson
8460000 | Blood group antibody Vg^a^
8484008 | Blood group antigen Mur
8486005 | HLA-DRw15 antigen
8491006 | Complement inhibitor
8612007 | Tumor necrosis factor

8660005 | Complement component C3c
8731008 | Blood group antibody Austin
8740007 | C3(H20)Bb
8786009 | Blood group antigen Wd^a^
8818009 | Blood group antibody Tri W
8878003 | Blood group antigen Evelyn
8908003 | Blood group antibody I^T^
8977007 | Blood group antibody Tarplee
9054000 | Human leukocyte antigen B15
9172009 | Blood group antibody Alda
9253000 | HLA-DRw16 antigen
9296005 | Gamma interferon
9319001 | Blood group antibody Vennera
9392009 | Blood group antibody Pollio
9398008 | Blood group antigen Pillsbury
9472003 | Blood group antigen Schneider
9493000 | Homologous antigen
9608008 | Blood group antigen Noble
9672005 | S antigen
9701007 | Blood group antibody Pr>3<
9716005 | Blood group antibody Luke
9921004 | Blood group antibody 'N'
9930007 | Blood group antigen Hartley
9980001 | CDw75 - Cluster of differentiation antigen (workshop) 75
9985006 | Desarginisated complement enzyme
10105003 | Active C3bBbC3b
10158008 | Blood group antigen K13
10189007 | Conglutinin
10228005 | Blood group antibody Mil
10265007 | Blood group antibody Jobbins
10393009 | HLA - Human leukocyte antigen Dw20
10398000 | Blood group antibody iH
10436003 | Blood group antibody Ad
10440007 | HLA class II antigen
10473000 | C3 - Complement component 3
10560001 | Blood group antibody By
10570004 | Blood group antigen Sf^a^
10622000 | Blood group antibody Gilbraith
10889009 | Blood group antigen Cr3
10912008 | Blood group antigen Le^a^
10987005 | Platelet-derived growth factor
11004008 | Blood group antigen Ge3
11022006 | Blood group antibody Cr2
11041009 | Blood group antibody Dr^a^
11064006 | Blood group antigen Lu^b^
11091008 | Blood group antibody Madden
11123004 | Blood group antigen Simpson
11222004 | Blood group antigen Ge1
11233001 | Public blood group antigen
11252007 | Blood group antigen Sa
11267006 | Interleukin-10
11330000 | Platelet antibody HPA-4b
11353004 | Anti-glomerular basement membrane antibody
11447000 | Antibody to hepatitis B core antigen, IgM type
11474004 | Blood group antibody French
11479009 | Blood group antibody Ok^a^
11489008 | Blood group antigen Nickolai
11566003 | Blood group antibody Braden
11594006 | RH19 (ISBT symbol)
11600003 | Blood group antibody Terrell
11684009 | Blood group antigen Kennedy
11725001 | Blood group antigen Gould
11744008 | Blood group antigen Knudsen
11770009 | FY1 (ISBT symbol)
11799004 | Blood group antibody Donaldson
11825009 | Anti endomysial antibody
11863001 | GE6 (ISBT symbol)
11877003 | HLA-DRw10 antigen
11886008 | Blood group antibody Mckeever
12018003 | Trichophyton extract skin test
12103002 | HLA-B45 antigen
12190009 | Blood group antibody Lazicki

12194000 | Leukotriene
12379005 | Blood group antibody Do^b^
12498008 | Blood group antibody Kn^b^
12564002 | HLA class III antigen
12577006 | Blood group antibody Ch^a^
12627001 | Macrophage chemotactic factor
12642003 | Artificial antigen
12685007 | Blood group antigen Wiley
12752008 | Blood group antibody HLA-A7
12753003 | Blood group antibody Fr^a^
12899008 | Anti Lua
12934002 | Human leukocyte antigen Cw7
12998001 | Blood group antibody Mineo
13016007 | Blood group antigen Li^a^
13083005 | Eosinophilic chemotactic factor
13105002 | Hbs Ag/ayr
13377004 | MNS9 (ISBT symbol)
13400000 | HLA-Bw65 antigen
13435003 | Blood group antibody Cost-Stirling
13484006 | Blood group antibody NOR
13492002 | Blood group antibody Di^b^
13539006 | Blood group antibody Sharp
13571004 | Blood group antibody Stevenson
13590007 | Blood group antibody Kosis
13625002 | HLA - Human leukocyte antigen A24(9)
13701000 | Blood group antigen E. Amos
13717006 | Blood group antibody McCall
13723001 | Blood group antigen Man
13772008 | Blood group antibody Middel
13841004 | Leukotriene C
13872000 | Blood group antibody Fuller
14090005 | MNS2 (ISBT symbol)
14104007 | Blood group antigen O'Connor
14190008 | Blood group antibody T
14226002 | Blood group antigen Friedberg
14271005 | Blood group antigen Gon
14279007 | Blood group antibody Epi
14396005 | Blood group antibody Ls^a^
14444008 | Blood group antibody Todd
14464003 | HLA - Human leukocyte antigen Cw3
14517001 | Blood group antibody Jordan
14574003 | Blood group antibody Bovet
14604008 | Blood group antibody Hg^a^
14620003 | Blood group antibody B 9724
14665007 | Blood group antigen Parra
14711003 | A antigen
14726001 | Antibody to antigen in ISBT system 007
14827002 | DI1 (ISBT symbol)
14849008 | HLA-Bw77 antigen
14986005 | Blood group antigen Wilson
15011000 | Blood group antibody Ts
15021008 | Neoantigen
15073009 | Antigen excess immune complex
15275007 | Blood group antibody FR
15286009 | HLA - Human leukocyte antigen Cw2
15313005 | Blood group antibody Gf
15392001 | DO5 (ISBT symbol)
15416001 | Blood group antigen Pruitt
15469000 | Blood group antibody p
15551006 | Alternative complement pathway factor
15620005 | KN5 (ISBT symbol)
15653000 | Cluster of differentiation antigen 76
15683009 | Blood group antigen Robert
15754000 | Interleukin-7
15781000 | KEL20 (ISBT symbol)
15797005 | Blood group antigen A. Owens
15798000 | Blood group antibody Bp^a^
15909007 | Blood group antibody Yk^a^
15942008 | Blood group antibody Lanthois
16019008 | Blood group antibody Fy^x^
16025007 | HLA-DQw8 antigen
16074008 | Immune complex at equivalence

16122008 | Blood group antibody hr^H^
16133006 | Blood group antigen Kamiya
16138002 | Blood group antigen M'
16230002 | Blood group antigen Madden
16519001 | Blood group antibody Ny^a^
16705004 | HLA-Bw47 antigen
16734005 | Blood group antibody S>2<
16752005 | Blood group antigen Pearl
16878007 | Anti E
16951006 | Antigen in Rh blood group system
16996004 | Blood group antibody Gd
17074004 | Blood group antigen Pelletier
17108004 | Blood group antibody En^a^TS
17119001 | Blood group antibody Yh^a^
17152007 | Blood group antibody I^D^
17165003 | Blood group antigen 754
17430007 | Blood group antigen Hey
17462005 | KEL12 (ISBT symbol)
17595001 | Lymphocyte antigen CD32
17627009 | Hepatitis B e antibody
17640004 | Blood group antibody Savery
17659002 | Blood group antigen R.M.
17701004 | Brucella protein nucleate
17707000 | Blood group antibody Ritter
17740009 | Blood group antigen Epi
17853004 | Antibody excess immune complex
18030004 | Blood group antibody Balkin
18082002 | RH10 (ISBT symbol)
18150002 | Anti-A,B
18195009 | HLA - Human leukocyte antigen DR9
18359006 | Blood group antibody Fedor
18380000 | Blood group antibody K^w^
18426007 | Blood group antibody MZ 443
18436004 | Cluster of differentiation antigen 58
18468007 | Blood group antibody M^g^
18486005 | LE6 (ISBT symbol)
18501000 | Human leucocyte antigen B51
18533009 | RH34 (ISBT symbol)
18611000 | RH18 (ISBT symbol)
18627007 | Blood group antigen iP>1<
18659000 | Fungal antibody
18737007 | Blood group antigen Rh38
18761000 | Leukocyte 23
18763002 | Blood group antigen Dropik
18853002 | CD2 - Cluster of differentiation antigen 2
19009001 | Lymphocyte antigen CD18
19011005 | Anti N
19022009 | Blood group antigen Jopson
19035000 | Blood group antibody Hall J
19089003 | Leukocyte 11
19136002 | Blood group antibody S1^a^
19178008 | Blood group antibody U
19182005 | C>5b67< inhibitor
19331004 | DI6 (ISBT symbol)
19395006 | Blood group antigen Pe
19400007 | Blood group antibody Baumler
19530002 | Blood group antibody ISBT P1
19565002 | Blood group antibody Rios
19677004 | LCA - Leukocyte common antigen
19728002 | Lymphocyte antigen CD17
19755006 | Blood group antibody Shannon
19783008 | Blood group antibody Groslouis
19830006 | Blood group antibody
19868008 | Lymphocyte antigen CDw78
19917006 | Hydroperoxy eicosatetraenoic acid
19945000 | MNS1 (ISBT symbol)
20009008 | RG1 (ISBT symbol)
20040001 | DI2 (ISBT symbol)
20057002 | C6 - Complement component 6
20228006 | KEL5 (ISBT symbol)
20304007 | Blood group antibody globoside
20355000 | Granulocyte antibody

20383003 | Blood group antibody Rh38
20493009 | HLA - Human leukocyte antigen Dw24
20576006 | Blood group antigen Santano
20591008 | Blood group antibody Nielsen
20748000 | Blood group antigen VK
20764008 | Leukocyte 7
20777004 | Blood group antibody Margaret
20807009 | Anti nucleolus antibody
20823009 | Complement, thermolabile serum protein complex with cytotoxic effect
20878003 | Blood group antibody Hut
20898008 | Extracellular matrix receptor type III
20907008 | Blood group antibody Cipriano
21023002 | RH42 (ISBT symbol)
21064007 | Blood group antibody Rm
21102006 | Blood group antigen McC^d^
21145004 | Blood group antibody Hr>o<
21149005 | Blood group antibody Pr>1h<
21158003 | Independent high incidence blood group antibody
21166007 | Lymphocyte antigen CD21
21256006 | Human leukocyte antigen Dw23
21315005 | MNS15 (ISBT symbol)
21521008 | Human leucocyte antigen Bw71
21642002 | RH12 (ISBT symbol)
21774001 | Complement component, precursor
21802009 | Blood group antibody HEMPAS
21873000 | Blood group antibody Griffith
21876008 | Blood group antigen NOR
21920001 | LU14 (ISBT symbol)
22023004 | Blood group antigen Le^x^
22269007 | Blood group antibody Sa
22290004 | Hepatitis B surface antigen
22332006 | Blood group antibody McC^e^
22348007 | Human leukocyte antigen DR5
22503007 | Human leukocyte antigen Bw50
22568000 | RH17 (ISBT symbol)
22627002 | Blood group antibody Barrett
22839007 | Blood group antibody Au^a^
22939008 | Blood group antibody Messenger
22971001 | I system antibody
23016008 | HLA - Human leukocyte antigen DPw1
23064004 | DI17 (ISBT symbol)
23101007 | CROM5 (ISBT symbol)
23103005 | Blood group antigen Niemetz
23164001 | Sp40/40
23165000 | Blood group antigen Terrano
23200004 | FY3 (ISBT symbol)
23240005 | Homologous restriction factor
23314002 | Blood group antibody Schwend
23318004 | Neutrophil cytoplasmic antibody
23369004 | Circulating immune complex
23380004 | KEL3 (ISBT symbol)
23385009 | Blood group antibody ALe^b^
23396001 | Blood group antibody Green
23434000 | MNS31 (ISBT symbol)
23603009 | GE4 (ISBT symbol)
23689006 | Platelet antigen HPA-3a
23760003 | Blood group antibody Wb
23774005 | HLA - Human leukocyte antigen Dw9
23775006 | Blood group antibody Rh40
23878002 | Blood group antibody Whittle
23893003 | Blood group antigen La Fave
24002009 | KN2 (ISBT symbol)
24008008 | Blood group antibody Laine
24023003 | Properdin native
24143007 | Platelet antibody HPA-2a
24207006 | Blood group antigen Tri W
24248009 | Saline agglutinin
24304001 | Blood group antibody K11
24391001 | Platelet antigen HPA-4a
24404002 | ABO3 (ISBT symbol)
24479006 | Blood group antibody Kollogo
24487007 | High incidence antigen

24503006 | Blood group antibody Vr
24540003 | Blood group antibody En^a^KT
24574004 | FY2 (ISBT symbol)
24655002 | Leukocyte 3a
24673004 | Human leukocyte antigen Dw11
24710003 | Blood group antibody Pr^a^
24821002 | Blood group antibody Tx
24857007 | Complement fixing antibody
24860000 | Blood group antibody Don E. W.
24898000 | Independent low incidence blood group antibody
24926008 | LW6 (ISBT symbol)
24978006 | Blood group antigen Bert
24995003 | Human leukocyte antigen A28
25095009 | Blood group antigen Ol^a^
25176004 | Mumps skin test antigen
25282007 | Human leukocyte antigen Bw55
25315004 | HLA - Human leukocyte antigen A34(10)
25400004 | Blood group antibody Yt^b^
25426009 | Blood group antigen Bridgewater
25453008 | Antibody to antigen in ISBT system JK
25513007 | Blood group antigen Stewart
25557006 | Blood group antigen Langer
25717005 | Myeloid-macrophage antibody
25773002 | Blood group antigen Elder
25826003 | Platelet antibody HPA-5a
25833003 | LU1 (ISBT symbol)
25867008 | Blood group antigen Haven
25964000 | KEL17 (ISBT symbol)
26005009 | Blood group antigen Tajama
26021004 | Blood group antibody Sd^a^
26066008 | MNS5 (ISBT symbol)
26093006 | Platelet antigen HPA-4b
26274001 | Hydroxyeicosatetraenoic acid
26355006 | Blood group antibody Cameron
26441004 | Blood group antigen Bg^a^
26647007 | Blood group antibody Coates
26651009 | Blood group antigen Rd^a^
26720006 | Blood group antibody McC^c^
26740004 | Eosinophilic derived inhibitor
26749003 | Blood group antibody Kaj
26855002 | KEL14 (ISBT symbol)
26937007 | MNS20 (ISBT symbol)
27024002 | Blood group antigen By
27044008 | Blood group antibody Becker
27047001 | Blood group antigen Schwend
27048006 | Blood group antigen Can
27076003 | Blood group antibody Rich
27089009 | Anti Ce
27130004 | Mac-1 leukocyte adhesive protein
27205008 | Blood group antigen IAB
27273002 | Complement component 1s
27340007 | Human leukocyte antigen A10
27384007 | GLOB3 (ISBT symbol)
27417007 | Blood group antigen Geslin
27425009 | Platelet antigen HPA-2a
27453009 | Blood group antigen John Smith
27459008 | CO2 (ISBT symbol)
27487004 | Blood group antigen Talbert
27489001 | Blood group antigen Don
27607009 | Blood group antigen Ts
27626001 | Anti S
27800009 | Blood group antibody BLe^d^
27862003 | Blocking antibody
27899003 | Blood group antibody Ol^a^
27963007 | Blood group antibody Toms
27999002 | Blood group antigen Hands
28145009 | Blood group antibody Cr3
28262007 | Blood group antibody Robert
28298004 | Pan-leukocyte antibody
28361003 | Blood group antibody Mathison
28408005 | LW7 (ISBT symbol)
28424006 | CD62 - Cluster of differentiation antigen 62

28427004 | Human leukocyte antigen DQw9
28495003 | Blood group antibody El
28564007 | Blood group antibody Reiter
28577003 | Blood group antibody Chr^a^
28585007 | Platelet-specific antigen
28673005 | Ribosomal antibody
28679009 | CD28 - Cluster of differentiation antigen 28
28708009 | Blood group antigen Bovet
28716000 | Cluster of differentiation antigen (workshop) 65
28731009 | Blood group antibody Morrison
28779002 | Blood group antibody Savior
28808000 | Blood group antigen Stevenson
28819002 | Blood group antibody K12
28881009 | Blood group antibody B 9208
29244008 | Blood group antibody Lu4
29266001 | Blood group antigen Sadler
29275004 | Blood group antibody Tollefsen-Oyen
29468003 | DI8 (ISBT symbol)
29502003 | Blood group antigen IBH
29573007 | Blood group antigen Wade
29652001 | Blood group antibody Noble
29719004 | Blood group antibody Dav
29735006 | Cluster of differentiation antigen 33
29776002 | Complement component 7
29831006 | Blood group antigen Taylor
29988009 | Blood group antibody McC^f^
30030008 | Interleukin-9
30137009 | RH22 (ISBT symbol)
30209008 | Blood group antibody Gladding
30245006 | Blood group antibody Kelly
30357004 | Blood group antibody Santano
30413004 | Blood group antigen Cad
30498007 | Blood group antigen Emma
30540002 | Blood group antibody Simpson
30594007 | Leukocyte 1
30604008 | Platelet antigen HPA-2b
30607001 | LU3 (ISBT symbol)
30609003 | Blood group antibody Terrano
30621004 | Abs - Autoantibodies
30702003 | Blood group antibody D^w^
30723009 | Blood group antigen Payer
30797003 | CROM4 (ISBT symbol)
30954000 | Blood group antigen Charles
30965005 | Interleukin-6
30987001 | Blood group antibody Rh35
31001006 | CD68 - Cluster of differentiation antigen 68
31008000 | Blood group antibody Talbert
31024004 | Blood group antigen Good
31036005 | Blood group antigen Peacock
31245001 | Blood group antibody Oca
31317005 | RH8 (ISBT symbol)
31357008 | Blood group antibody Sc3
31400002 | Human leucocyte antigen Bw63
31854008 | Blood group antibody Terschurr
31897003 | MNS26 (ISBT symbol)
31960007 | La antibody
32073006 | Human leukocyte antigen Bw60
32079005 | Blood group antigen Ramskin
32089009 | Blood group antibody VS
32147003 | Blood group antigen Suhany
32167007 | Blood group antibody Nickolai
32233008 | Blood group antibody Kasamatsuo
32235001 | Blood group antibody A 8306
32237009 | Blood group antibody IBH
32241008 | DI4 (ISBT symbol)
32261002 | Blood group antibody Lu6
32277001 | Soluble immune complex
32305006 | Blood group antibody Rd^a^
32314001 | Blood group antibody Marriott
32324009 | Blood group antibody BR 726750
32329004 | Blood group antigen I fetal
32351007 | Thymus-dependent antigen

32364008 | Blood group antigen Tm
32396000 | Blood group antibody Lu5
32403003 | Blood group antibody Pr>a<
32481003 | Blood group antibody Mackin
32609007 | Hepatitis A antibody
32616008 | Blood group antibody Zim
32699000 | Blood group antigen R>2<R>2<-202
32842006 | Blood group antibody Rh42
32860006 | HLA - Human leukocyte antigen A9
32943000 | Cluster of differentiation antigen 24
32974003 | Blood group antigen Banks
32990003 | HF - Factor H
33029004 | Blood group antibody Bowyer
33037007 | Blood group antigen Austin
33053005 | Blood group antigen Bruno
33057006 | Macrophage antibody
33063002 | LU13 (ISBT symbol)
33188008 | Blood group antigen Christiansen
33210004 | Blood group antibody trihexosylceramide
33309006 | Human leukocyte antigen DQw5
33355008 | Blood group antibody Banks
33509005 | Blood group antibody Mur
33550002 | Blood group antigen Kirkpatrick
33604009 | Blood group antigen Finlay
33680002 | HLA - Human leukocyte antigen B12
33691009 | Blood group antibody Co^b^
33825006 | JK2 (ISBT symbol)
33865005 | Blood group antibody Baltzer
33987002 | Public blood group antibody
34049004 | Blood group antibody Lu9
34161009 | Blood group antibody Ku
34180006 | Blood group antibody Min
34204008 | Blood group antibody Warren
34316000 | Blood group antibody Ge1
34355004 | Inactivated complement enzyme
34388006 | Blood group antibody Fuerhart
34400001 | Blood group antibody Teremok
34453005 | HLA - Human leukocyte antigen B27
34465009 | Human leukocyte antigen DQw7
34510007 | Clonal inhibitory factor
34569000 | Blood group antibody Jn^a^
34745001 | Slow reacting substance-A of anaphylaxis
34750007 | Blood group antigen Panzar
34829007 | Complement component
34832005 | Blood group antibody I^s^
34868000 | Human leukocyte antigen DQ3
34900009 | ABO2 (ISBT symbol)
34912008 | Blood group antibody Ramskin
34968004 | Blood group antigen Lee
34970008 | RH48 (ISBT symbol)
35012004 | Blood group antibody HLA-A9
35040009 | Blood group antibody Rh29
35068008 | Anti C
35072007 | HLA - Human leukocyte antigen B16
35092000 | Lymphocyte antigen CD70
35118003 | Blood group antibody Fy5
35213004 | Blood group antibody Wallin
35215006 | Scarlet fever streptococcus toxin
35310003 | Polyclonal antibody
35367007 | Blood group antigen McC^e^
35410004 | Blood group antibody Kp^c^
35556005 | Sessile antibody
35614002 | LU17 (ISBT symbol)
35645003 | Blood group antigen French
35760006 | Myeloid antibody
35825008 | Cat scratch disease antigen
35841009 | Macrophage migration inhibitory factor
35906006 | Blood group antibody MPD
35922007 | Blood group antibody Black
35952004 | Blood group antibody Block
36005003 | Blood group antibody Tofts
36016005 | Blood group antibody Haase

36100005 | DO2 (ISBT symbol)
36260004 | Blood group antibody Raison
36396003 | Blood group antigen Van Buggenhout
36403001 | Blood group antibody ELO
36418000 | Blood group antigen McC^b^
36562006 | Hemolysin
36613003 | Blood group antigen Pr>1h<
36652005 | Blood group antigen H>T<
36713008 | Blood group antibody McC^d^
36744004 | RH3 (ISBT symbol)
36757007 | Blood group antigen Raison
36759005 | Human leukocyte antigen Bw6
36804003 | Blood group antigen Tasich
36842009 | HLA - Human leukocyte antigen Dw16
36853003 | Blood group antigen Vienna
36907009 | Blood group antibody Kennedy
36933001 | Antibody to antigen in ISBT system RH
37067002 | Blood group antibody Shier
37100000 | Blood group antigen Bradford
37282004 | Blood group antibody B 7358
37287005 | HLA - Human leukocyte antigen A1
37300006 | Blood group antibody h
37583005 | Blood group antigen Buckalew
37654004 | KEL19 (ISBT symbol)
37713002 | Blood group antigen Dautriche
37861002 | KEL7 (ISBT symbol)
37902007 | Blood group antigen A.M.
37984005 | Blood group antibody Don
38227005 | MNS6 (ISBT symbol)
38262000 | Active C5b678
38269009 | CD1c - Cluster of differentiation antigen 1c
38367008 | Blood group antigen Hoalzel
38445008 | Blood group antigen Rils
38476002 | Interleukin
38499003 | Blood group antibody Naz
38519002 | Blood group antigen Donaldson
38553003 | Blood group antigen Schuppenhauer
38554009 | HLA - Human leukocyte antigen B5
38558007 | Blood group antibody Ghawiler
38771008 | HLA - Human leukocyte antigen DPw6
38779005 | Blood group antibody Ht^a^
38874005 | Blood group antigen V.G.
38908008 | Blood group antigen Jankowski
39024001 | Blood group antibody Yt^a^
39053000 | Factor D
39082004 | Hepatitis B core antigen
39118009 | High incidence antibody
39138005 | Blood group antibody Milano
39241007 | Human leukocyte antigen Dw1
39327001 | Blood group antibody Rh43
39337006 | Blood group antibody Es^a^
39442002 | Paratope
39515000 | Blood group antigen Ht^a^
39525005 | Tumor necrosis factor alpha
39650003 | Human leukocyte antigen HLA-Bw54 antigen
39665002 | Blood group antigen Pr>2<
39678002 | Blood group antigen Kominarek
39840002 | Blood group antibody Di^a^
39988003 | Skin reactive factor
39999001 | RH21 (ISBT symbol)
40018000 | Blood group antibody Oliver
40030006 | MNS8 (ISBT symbol)
40044000 | Human leukocyte antigen DRw11
40048002 | Blood group antigen Englund
40065006 | Human leucocyte antigen Bw73
40140007 | Blood group antibody Kirkpatrick
40154004 | Blood group antibody Singleton
40270009 | Blood group antibody Truax
40364000 | LE5 (ISBT symbol)
40447004 | Blood group antibody Hy
40456007 | Blood group antigen IB
40558006 | Blood group antigen VA

40621002 | MNS12 (ISBT symbol)
40706003 | Blood group antigen Toms
40856000 | Leukotriene A
40992002 | CD - Cluster of differentiation antigen marker
41044008 | Blood group antigen Woit
41093003 | Blood group antibody E^w^
41096006 | Blood group antibody Y. Bern
41285005 | Blood group antigen Jones
41301002 | H-2 locus
41318003 | Blood group antibody Js^b^
41403003 | MNS14 (ISBT symbol)
41562008 | Blood group antibody Tm
41576009 | RH26 (ISBT symbol)
41644009 | Blood group antigen Baltzer
41667006 | Blood group antigen Begovitch
41761003 | Blood group antibody Stewart
41771001 | Blood group antigen Gallner
41858009 | Blood group antigen Wetz
41886001 | Blood group antigen Kenneddy
41930000 | Blood group antigen McDermott
41978000 | Blood group antibody V.G.
41999002 | Blood group antibody Joslin
42038007 | Human leukocyte antigen Bw62
42048009 | Blood group antibody Terry
42078000 | Blood group antigen Kursteiner
42092006 | Blood group antigen Allchurch
42107008 | Human leucocyte antigen Cw
42122003 | Blood group antibody M^v^
42166001 | XK1 (ISBT symbol)
42255003 | Blood group antibody Zaw
42333009 | Blood group antibody LW^b^
42371001 | Heterocytotropic antibody
42473002 | Blood group antibody Cross
42490008 | Blood group antibody Tn
42664002 | Human leukocyte antigen Dw17
42732002 | Leukocyte 6
42768007 | Blood group antigen En^a^ trypsin-sensitive
42784008 | Blood group antibody Wd^a^
42799008 | Idiotype
42804003 | Lymphocyte antigen CD13
42830004 | Blood group antibody Wilson
42848008 | Blood group antigen MPD
42888003 | Blood group antigen Cipriano
42891003 | Leukocyte 19
42897004 | Blood group antigen Donati
42926001 | Blood group antigen Seymour
43042002 | Platelet antibody HPA-5b
43076006 | Blood group antibody Rh37
43095004 | Lymphocyte antigen CD35
43181000 | Blood group antibody Cl^a^
43241001 | Blood group antibody Pelletier
43278003 | Platelet activating factor
43347004 | Blood group antigen A>1< Le^d^
43397000 | Idiotope
43417002 | Blood group antibody IH
43425000 | Blood group antigen Dahl
43576000 | Blood group antibody N^A^
43596008 | Human leukocyte antigen Bw64
43632003 | Blood group antibody K14
43678006 | Blood group antigen Pr>3<
43788001 | Blood group antibody Davis
43807002 | IN2 (ISBT symbol)
43827001 | Blood group antigen Mineo
43864007 | Blood group antigen Ull
43871002 | Human leukocyte antigen Dw7
44040003 | Human leucocyte antigen Bw57
44045008 | Blood group antibody Tasich
44046009 | Blood group antibody Paular
44049002 | Blood group antigen Lindsay
44063008 | Blood group antigen Pt^a^
44093002 | Blood group antibody KL
44112005 | Blood group antigen Reynolds

44139002 | Blood group antigen Don E. W.
44158006 | CD64 - Cluster of differentiation antigen 64
44212003 | Ro antibody
44239006 | Platelet antibody HPA-1
44331007 | Blood group antibody In^b^
44369002 | Chemotaxin
44439008 | Blood group antigen Smith
44447008 | Blood group antigen Fleming
44459007 | Interleukin-8
44472008 | Blood group antibody Begovitch
44675004 | Blood group antibody Nou
44680008 | Factor VIII antigen
44686002 | Blood group antigen Lud
44706009 | T3 - T-cell lineage 3
44719008 | Mediator of immune response
44728009 | Complement component C1
44824000 | Blood group antibody Pearl
45026006 | MNS21 (ISBT symbol)
45044003 | Blood group antibody Lud
45047005 | Lymphokine
45095001 | KEL18 (ISBT symbol)
45193006 | Blood group antibody Horw
45246008 | C4BP - Complement component 4 binding protein
45273009 | RH28 (ISBT symbol)
45321005 | Anti M
45333000 | KN4 (ISBT symbol)
45388001 | Blood group antigen Laine
45396006 | ACLA - Anti-cardiolipin antibody
45407009 | Blood group antigen Ghawiler
45428000 | Blood group antibody Perry
45441001 | Blood group antigen Tk
45510000 | Blood group antibody Jopson
45528003 | Blood group antibody Dugstad
45530001 | ANF - Anti-nuclear factor
45601001 | Blood group antibody A.M.
45637006 | Blood group antibody Bonde
45656001 | Human leukocyte antigen Bw22
45668005 | Blood group antigen Bouteille
45672009 | Blood group antibody Reynolds
45857007 | Antilysosomal antibody
45932003 | Jo-1 antibody
45934002 | MNS38 (ISBT symbol)
45969000 | Blood group antibody i
45992000 | MNS4 (ISBT symbol)
46096007 | Blood group antibody Knudsen
46150001 | HLA - Human leukocyte antigen Bw4
46158008 | Human leukocyte antigen Dw14
46195008 | Smith antibody
46234003 | MNS27 (ISBT symbol)
46261003 | RH9 (ISBT symbol)
46331009 | Blood group antibody K13
46336004 | Complement component C2b
46469003 | Properdin convertase, complement component
46519008 | Blood group antibody ILe^bH^
46539007 | C1r - Complement component 1r
46548002 | Leukotriene B
46613001 | Human leucocyte antigen Bw58
46663006 | Blood group antibody Lee
46730008 | Blood group antigen Bio-5
46755002 | Blood group antigen Schor
46841000 | DI15 (ISBT symbol)
46959001 | Leu-M5 antigen
47019005 | Blood group antigen Hildebrandt
47023002 | CD66 - Cluster of differentiation antigen 66
47038001 | Human leukocyte antigen (HLA) antigen
47068005 | Blood group antibody Jr^a^
47104007 | CO3 (ISBT symbol)
47136000 | Blood group antigen Manley
47151009 | Blood group antibody Win
47201006 | 5-Hydroperoxy-6,8,11,14-eicosatetraenoic acid
47215008 | SC2 (ISBT symbol)
47341004 | Blood group antigen Driver

47364002 | Blood group antigen Ryan
47464003 | Blood group antibody Woit
47469008 | Blood group antibody Seymour
47553004 | Blood group antibody Sul
47565000 | I region, major histocompatibility complex
47581005 | Type 1 chain precursor structure (lacto-N-tetraosylceramide)
47601000 | Blood group antigen Savery
47603002 | Blood group antibody Pillsbury
47626009 | Blood group antibody Kemma
47662005 | Blood group antigen h
47674009 | Blood group antigen Pr>1d<
47692001 | Blood group antigen Rm
47733001 | Blood group antibody Bradford
47769009 | Platelet antibody HPA-5
47784000 | Blood group antibody IP
47851007 | Human leukocyte antigen A69(28)
47860004 | Human leukocyte antigen A3
47894002 | Lymphotoxin
47974007 | Blood group antibody Tg^a^
48060000 | Blood group antigen Ritter
48109004 | KEL6 (ISBT symbol)
48116003 | Blood group antigen Paris
48131007 | Blood group antibody Neut
48140006 | Blood group antibody Whittaker
48154003 | Blood group antibody Zwal
48170001 | HLA - Human leukocyte antigen Cw1
48217002 | Complement regulator
48270008 | Lymphocyte antigen CDw49f
48284003 | Kell system antigen
48323009 | Blood group antibody Schneider
48366002 | RH39 (ISBT symbol)
48401006 | I1 (ISBT symbol)
48416009 | Blood group antigen Green
48417000 | Human leukocyte antigen Dw26
48444005 | Freund adjuvant
48464000 | Blood group antibody Sw^a^
48699007 | Blood group antigen Carson
48727007 | Interleukin-4
48751002 | Blood group antibody Can
48779008 | Blood group antibody Hamet
48798005 | Antigen in Lutheran (LU) blood group system
48815002 | Blood group antigen Shannon
48821003 | Leukocyte 12
48869000 | Blood group antigen Jordan
48919009 | Blood group antigen Block
49028001 | Blood group antibody K16
49106003 | Human leukocyte antigen DR1
49119004 | Blood group antigen Bryant
49148005 | Human leukocyte antigen Cw11
49163008 | Blood group antigen Sd^a^
49212001 | Blood group antigen D 1276
49327004 | Interferon
49344000 | Blood group antibody VK
49352002 | Mediator of inflammation
49377001 | Blood group antigen Davis
49419007 | Active C4b
49449009 | Blood group antibody Wimberly
49495002 | Human leukocyte antigen A
49568009 | Blood group antigen Terrell
49582009 | MNS25 (ISBT symbol)
49599005 | Private blood group antibody
49616005 | Monoclonal antibody
49653004 | Blood group antigen Taur
49677005 | Human leukocyte antigen Dw22
49744003 | Blood group antibody M1^a^
49772005 | Blood group antigen Simon
49837000 | Blood group antigen Horn
49858006 | CDw52 - Cluster of differentiation antigen (workshop) 52
49912009 | RH23 (ISBT symbol)
49991001 | Blood group antigen Meteja
49997002 | Human leukocyte antigen Bw59
50028004 | Blood group antibody Boston

50156006 | LE2 (ISBT symbol)
50272007 | Blood group antibody hr^s^
50383001 | Human leukocyte antigen Bw76
50480002 | MNS16 (ISBT symbol)
50526007 | Blood group antibody S^D^
50700004 | Blood group antigen Heibel
50752003 | Blood group antibody Wiley
50762005 | Interleukin-1
50864002 | Blood group antibody CE
50899003 | Blood group antibody Je^a^
50988004 | c-ALL - Common acute lymphoblastic leukemia antigen
51222003 | Blood group antigen IP>1<
51238009 | Lymphocyte antigen CD2R
51242007 | RH45 (ISBT symbol)
51262004 | Blood group antibody Donati
51276003 | Blood group antibody Bothrops
51293003 | Blood group antigen rr-35
51311004 | Blood group antigen B 9208
51372003 | Blood group antibody An^a^
51414008 | Human leukocyte antigen DR2
51415009 | Blood group antibody Kn^a^
51447004 | Blood group antibody Kam
51468003 | Blood group antibody Tajama
51511003 | Blood group antigen Kosis
51515007 | Human leukocyte antigen DRw
51645003 | Blood group antibody Good
51718007 | Human leukocyte antigen Bw46
51809003 | Blood group antibody Pantaysh
51810008 | Proliferative inhibitory factor
51873003 | Blood group antibody Lagay
51941005 | Anti-B
51950007 | Mitochondrial antibody
52078008 | Epitope
52081003 | Blood group antigen Griffith
52160001 | Cluster of differentiation antigen 9
52169000 | Platelet antibody HPA-2
52193005 | Blood group antibody Lindsay
52210003 | Blood group antibody Manley
52227006 | Platelet antibody HPA-3b
52258007 | Blood group antibody C^x^
52310000 | Blood group antibody Dp
52313003 | Complement component 8
52419000 | Human leukocyte antigen Bw72
52467005 | Blood group antigen Krog
52555006 | Blood group antigen Shier
52573009 | Blood group antibody PP>1<P^k^
52680001 | Hbs Ag/adr
52764004 | Blood group antibody Cad
52912003 | Blood group antibody Marks
52935000 | Human leucocyte antigen Bw42
52991006 | Blood group antibody Bryant
53023007 | Leukotriene D

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.490 Value Set http://hl7.org/fhir/ValueSet/specimen-collection-priority

URL: https://hl7.org/fhir/R4/valueset-specimen-collection-priority.html
Orders and Observations Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/specimen-collection-priority
Version: | 4.0.1
Name: | SpecimenCollectionPriority
Title: | Specimen collection priority
Definition: | This example value set defines a set of codes that can be used to indicate the priority of collection of a specimen.
Committee: | Orders and Observations Work Group
OID: | 2.16.840.1.113883.4.642.3.939 (for OID based terminology systems)
Copyright: | TBD
Source Resource | XML / JSON

This value set is used in the following places:

• Extension: http://hl7.org/fhir/StructureDefinition/specimen-collectionPriority: collectionPriority (CodeableConcept / Example)

4.4.1.490.1 Content Logical Definition

This value set includes codes from the following code systems:

http://example.com

Code | Display
1 | STAT
2 | ASAP
3 | ASAP-ED
4 | AM
5 | ROUTINE
6 | NURSE COLLECT
7 | CALL OR FAX

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.491 Value Set http://hl7.org/fhir/ValueSet/concept-property-type

URL: https://hl7.org/fhir/R4/valueset-concept-property-type.html
Vocabulary Work Group | Maturity Level: N | Normative (from v4.0.0) | Use Context: Any

| This page has been approved as part of an ANSI standard. See the Conformance Package for further details.

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/concept-property-type
Version: | 4.0.1
Name: | PropertyType
Title: | PropertyType
Definition: | The type of a property value.
Committee: | Vocabulary Work Group
OID: | 2.16.840.1.113883.4.642.3.780 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for PropertyType

• Resource: CodeSystem.property.type (code / Required)

4.4.1.491.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/concept-property-type

4.4.1.491.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 7 concepts

Expansion based on http://hl7.org/fhir/concept-property-type version 4.0.1

All codes from system http://hl7.org/fhir/concept-property-type

Code | Display | Definition
code | code (internal reference) | The property value is a code that identifies a concept defined in the code system.
Coding | Coding (external reference) | The property value is a code defined in an external code system. This may be used for translations, but is not the intent.
string | string | The property value is a string.
integer | integer | The property value is a string (often used to assign ranking values to concepts for supporting score assessments).
boolean | boolean | The property value is a boolean true | false.
dateTime | dateTime | The property is a date or a date + time.
decimal | decimal | The property value is a decimal number.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.492 Value Set http://hl7.org/fhir/ValueSet/operation-outcome

URL: https://hl7.org/fhir/R4/valueset-operation-outcome.html
FHIR Infrastructure Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/operation-outcome
Version: | 4.0.1
Name: | OperationOutcomeCodes
Title: | Operation Outcome Codes
Definition: | Operation Outcome codes used by FHIR test servers (see Implementation file translations.xml)
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.412 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: OperationOutcome.issue.details (CodeableConcept / Example)

4.4.1.492.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/operation-outcome

4.4.1.492.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 50 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/operation-outcome version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/operation-outcome

Code | Display | Definition
DELETE_MULTIPLE_MATCHES | Error: Multiple matches exist for the conditional delete
MSG_AUTH_REQUIRED | You must authenticate before you can use this service
MSG_BAD_FORMAT | Bad Syntax: "%s" must be a %s'
MSG_BAD_SYNTAX | Bad Syntax in %s
MSG_CANT_PARSE_CONTENT | Unable to parse feed (entry content type = "%s")
MSG_CANT_PARSE_ROOT | Unable to parse feed (root element name = "%s")
MSG_CREATED | New resource created
MSG_DATE_FORMAT | The Date value %s is not in the correct format (Xml Date Format required)
MSG_DELETED | This resource has been deleted
MSG_DELETED_DONE | Resource deleted
MSG_DELETED_ID | The resource "%s" has been deleted
MSG_DUPLICATE_ID | Duplicate Id %s for resource type %s

MSG_ERROR_PARSING | Error parsing resource Xml (%s)
MSG_ID_INVALID | Id "%s" has an invalid character "%s"
MSG_ID_TOO_LONG | Id "%s" too long (length limit 36)
MSG_INVALID_ID | Id not accepted
MSG_JSON_OBJECT | Json Source for a resource should start with an object
MSG_LOCAL_FAIL | Unable to resolve local reference to resource %s
MSG_NO_EXIST | Resource Id "%s" does not exist
MSG_NO_MATCH | No Resource found matching the query "%s"
MSG_NO_MODULE | No module could be found to handle the request "%s"
MSG_NO_SUMMARY | No Summary for this resource
MSG_OP_NOT_ALLOWED | Operation %s not allowed for resource %s (due to local configuration)
MSG_PARAM_CHAINED | Unknown chained parameter name "%s"
MSG_PARAM_INVALID | Parameter "%s" content is invalid
MSG_PARAM_MODIFIER_INVALID | Parameter "%s" modifier is invalid
MSG_PARAM_NO_REPEAT | Parameter "%s" is not allowed to repeat
MSG_PARAM_UNKNOWN | Parameter "%s" not understood
MSG_RESOURCE_EXAMPLE_PROTECTED | Resources with identity "example" cannot be deleted (for testing/training purposes)
MSG_RESOURCE_ID_FAIL | unable to allocate resource id
MSG_RESOURCE_ID_MISMATCH | Resource Id Mismatch
MSG_RESOURCE_ID_MISSING | Resource Id Missing
MSG_RESOURCE_NOT_ALLOWED | Not allowed to submit a resource for this operation
MSG_RESOURCE_REQUIRED | A resource is required
MSG_RESOURCE_TYPE_MISMATCH | Resource Type Mismatch
MSG_SORT_UNKNOWN | Unknown sort parameter name "%s"
MSG_TRANSACTION_DUPLICATE_ID | Duplicate Identifier in transaction: %s
MSG_TRANSACTION_MISSING_ID | Missing Identifier in transaction - an entry.id must be provided
MSG_UNHANDLED_NODE_TYPE | Unhandled xml node type "%s"
MSG_UNKNOWN_CONTENT | Unknown Content (%s) at %s
MSG_UNKNOWN_OPERATION | unknown FHIR http operation
MSG_UNKNOWN_TYPE | Resource Type "%s" not recognised
MSG_UPDATED | existing resource updated
MSG_VERSION_AWARE | Version aware updates are required for this resource
MSG_VERSION_AWARE_CONFLICT | Update Conflict (server current version = "%s", client version referenced = "%s")
MSG_VERSION_AWARE_URL | Version specific URL not recognised
MSG_WRONG_NS | This does not appear to be a FHIR element or resource (wrong namespace "%s")
SEARCH_MULTIPLE | Error: Multiple matches exist for %s search parameters "%s"
SEARCH_NONE | Error: no processable search found for %s search parameters "%s"
UPDATE_MULTIPLE_MATCHES | Error: Multiple matches exist for the conditional update

Additional Language Displays

Code | Spanish (es) | French (fr) | Italian (it) | Nederlands (Dutch, nl) | Polskie (Polish, pl) | Русский (Russian, ru) | □□ (Chinese, zh)
DELETE_MULTIPLE_MATCHES | | | | Fout: er is meer dan één resultaat voor de conditionele delete
MSG_AUTH_REQUIRED | Debe autenticarse antes de poder usar este servicio | Vous devez être authentifié avant de pouvoir utiliser ce service | Autenticazione richiesta prima di usare questo servizio | U moet zich authenticeren voor gebruik van deze service | Wymagana autentykacja przed użyciem tego serwisu | | □□□□□□□□□□
MSG_BAD_FORMAT | Sintaxis Incorrecta: "%s" debe de ser un %s' | Erreur de Syntaxe : "%s" doit être un %s | Sintassi Errata: "%s" deve essere un %s' | Verkeerde syntax: "%s" moet een %s zijn | Błąd składni: "%s" powinno być %s' | Неверный синтакс: "%s" должен быть %s | □□□□: "%s" □□□□□ %s'
MSG_BAD_SYNTAX | Sintaxis Incorrecta en %s | Erreur de Syntaxe dans %s | Sintassi errata in %s | Verkeerde syntax in %s | Błąd składni w %s | Неверный синтакс: %s | %s □□□□□
MSG_CANT_PARSE_CONTENT | No se pudo parsear el feed (el tipo de contenido de la entry = "%s") | Impossible d'analyser le flux (type de contenu de l'entrée = "%s") | Impossibile effettuare il parsing del feed (tipo del contenuto della entry = "%s") | Kan feed niet verwerken (contenttype inhoud = "%s") | Błąd parsowania (typ zawartości wejściowej = "%s") | | □□□□feed (□□□□□□□ = "%s")
MSG_CANT_PARSE_ROOT | No se pudo parsear el feed (nombre del elemento raiz = "%s") | Impossible d'analyser le flux (nom de l'élément racine = "%s") | Impossibile effettuare il parsing del feed (nome elemento root = "%s") | Kan feed niet verwerken (rootelementnaam = "%s") | Błąd parsowania (nazwa elementu root = "%s") | Не удалось разобрать данные (корневой элемент = "%s") | □□□□feed (□□□□ = "%s")
MSG_CREATED | | | | Nieuwe resource gemaakt
MSG_DATE_FORMAT | El valor de la fecha %s no está en el formato correcto (se requiere un formato de fecha Xml) | Le format de la date %s est incorrect (format Date Xml attendu) | Il valore %s per la data non è nel formato corretto (richiesto il Formato Data Xml) | De Datum-waarde %s heeft niet de juiste structuur (Xml Date vereist) | Niepoprawny format wartości daty %s (wymagany format XML) | Значение Date %s в неверном формате (требуется Xml Date формат) | □□□□ %s □□□□□ (□□□Xml Date□□)
MSG_DELETED | Este recurso ha sido borrado | La ressource a été supprimée | Questa risorsa è stata cancellata | Deze resource is verwijderd | Ten zasób został usunięty | Данный ресурс был удалён | □□□□□□
MSG_DELETED_DONE | Recurso borrado | Ressource supprimée | Risorsa cancellata | Resource verwijderd | Zasób usunięto | Ресурс удалён | □□□□□
MSG_DELETED_ID | El recurso "%s" ha sido borrado | La ressource "%s" a été supprimée | La risorsa "%s" è stata eliminata | De resource "%s" is verwijderd | Zasób "%s" został usunięty | Ресурс "%s" был удалён | □□ "%s" □□□□
MSG_DUPLICATE_ID | Id %s duplicada para el recurso de tipo %s | Id %s en double pour le type de ressource %s | Id %s duplicato per il tipo di risorsa %s | Duplicaat-id %s voor resourcetype %s | Zdublowany identyfikator %s dla zasobu typu %s | Дублирующий Id %s для типа ресурса %s | ID %s □□□□□□□ %s□
MSG_ERROR_PARSING | Error parseando el recurso Xml (%s) | Erreur d'analyse de la ressource Xml (%s) | Errore nel parsing della risorsa Xml (%s) | Fout in verwerking resource Xml (%s) | Błąd w trakcie parsowania zasobu XML (%s) | Ошибка синтаксического разбора ресурса Xml (%s) | □□□□Xml□□□ (%s)

MSG_ID_INVALID | El Id "%s" contiene el caracter inválido "%s" | Id "%s" possède un caractère invalide "%s" | L''Id "%s" ha un carattere non valido: "%s" | Id "%s" heeft een ongeldig teken "%s" | Identyfikator "%s" zawiera niepoprawny znak "%s" | Id "%s" содержит недопустимые символы "%s" | ID "%s" □□□□□□: "%s"

MSG_ID_TOO_LONG | El Id "%s" es demasiado largo (limite de longitud: 36) | Id "%s" trop long (la longueur limite est 36) | Id "%s" troppo lunga (limite di lunghezza: 36) | Id "%s" te lang (max lengte 36) | Identyfikator "%s" jest zbyt długi (limit długości 36) | Недопустимая длина Id "%s" (ограничение 36) | Id "%s" □□ (□□□□: 36)

MSG_INVALID_ID | Id no aceptada | Id non accepté | Id non accettato | Id niet geaccepteerd | Identyfikator nie zaakceptowany | Id не принято | Id□□□□

MSG_JSON_OBJECT | La fuente Json para un recurso debería empezar con un objeto | La source Json pour une ressource doit commencer par un objet | Il sorgente Json di una risorsa dovrebbe iniziare con un oggetto | Json Source van een resource moeten beginnen met een object | Źródło json dla zasobu powinno rozpoczynać się od obiektu | Json Source для ресурса должен начинаться с объекта | □□□Json□□□□□object□□

MSG_LOCAL_FAIL | Imposible resolver la referencia al recurso %s | Impossible de résourdre la référence locale à la ressource %s | Impossibile risolvere il riferimento locale alla risorsa %s | De resource met lokale referentie %s is niet gevonden | Zasób wskazywany przez loklaną referencję %s nie został odnaleziony | Невозможно определить локальную ссылку на ресурс %s | □□□□□□□ %s □□□□□

MSG_NO_EXIST | El recurso con Id "%s" no existe | La ressource avec l'Id "%s" n'existe pas | La risorsa con Id "%s" non esiste | Resource-id "%s" bestaat niet | Zasób o identyfikatorze "%s" nie istnieje | Ресурс Id "%s" не существует | □□Id "%s"□□□

MSG_NO_MATCH | | | Nessuna Risorsa soddisfa la query "%s" | Geen resource gevonden met query "%s"

MSG_NO_MODULE | No se encontró un modulo que gestione la petición "%s" | Aucun module disponible pour traiter la requête "%s" | Impossibile trovare un modulo per gestire la richiesta "%s" | Er kon geen module worden gevonden om verzoek "%s" te verwerken | Nie można odnaleźć modułu, aby obsłużyć żądanie "%s" | Модуль для обработки запроса "%s" не найден | □□□□□□□□"%s"□□□

MSG_NO_SUMMARY | No existe un resumen para este recurso | Aucun résumé disponible pour cette ressource | Nessun riepilogo per questa risorsa | Geen samenvatting voor deze resource | Brak podsumowania (Summary) dla tego zasobu | Отсутствует Summary для данного ресурса | □□□□summary

MSG_OP_NOT_ALLOWED | Operación %s no permitida para el recurso %s (debido a la configuración local) | L'opération %s n'est pas permise pour la ressource %s (à cause de la configuration locale) | Operazione %s non consentita per la risorsa %s (a causa di configurazioni locali) | Bewerking %s niet toegestaan voor resource %s (vanwege lokale configuratie) | Niedozwolona operacja %s dla zasobu %s (ze względu na lokalną konfiguracje) | Операция %s недопустима для ресурса %s (по причине локальной конфигурации) | □□ %s □□□□□□□□ %s (□□□□□□)

MSG_PARAM_CHAINED | Nombre de parametro encadenado desconocido: "%s" | Nom du paramètre chainé inconnu : "%s" | Nome di parametro concatenato sconosciuto: "%s" | Onbekende geschakelde parameternaam "%s" | Nieznana nazwa parametru powiązanego "%s" | Неизвестое вложенное наименование параметра "%s" | □□□□□□□□: "%s"

MSG_PARAM_INVALID | | | Il contenuto del Parametro "%s" non è valido | Waarde van parameter "%s" is ongeldig

MSG_PARAM_MODIFIER_INVALID | | | | Modifier van parameter "%s" is ongeldig

MSG_PARAM_NO_REPEAT | No se permite la repetición del parámetro "%s" | Le paramètre "%s" ne peut pas être répété | Non consentito ripetere il parametro "%s" | Parameter "%s" mag niet herhalen | Parametr "%s" nie może zostać powtórzony | Параметр "%s" не может быть повторён | □□"%s"□□□□

MSG_PARAM_UNKNOWN | Parámetro "%s" no reconocido | Paramètre "%s" non reconnu | Parametro "%s" non riconosciuto | Parameter "%s" onbekend | Niezrozumiały parametr "%s" | Параметр "%s" не понят | □□□□□□"%s"

MSG_RESOURCE_EXAMPLE_PROTECTED | Recursos con la identidad "example" no pueden ser borrados (son usados para pruebas/entrenamiento) | Les ressources ayant l'identité "example" ne peuvent pas être supprimées (utilisées pour les tests/formations) | Le Risorse aventi l'identità "example" non possono essere cancellate (per finalità di test/formazione) | Resources met identiteit "voorbeeld" kunnen niet worden verwijderd (ten behoeve van testen/training) | Zasoby oznaczone jako "example" nie mogą zostać usunięte (dla celów testów/szkoleń) | Ресурс с идентификатором "example" не может быть удалён (для случаев тестирования/обучения) | □"example" □ID□□□□□□□□ (□□□□/□□)

MSG_RESOURCE_ID_FAIL | imposible encontrar el id del recurso | impossible d'allouer l'id de la ressource | impossibile allocare l''id della risorsa | kan geen resource-id reserveren | nie można nadać identyfikatora zasobu | невозможно выделить идентификатор ресурса | □□□□□□ID

MSG_RESOURCE_ID_MISMATCH | | Problème de correspondance d'Id de la Ressource | | Resource ID's komen niet overeen

MSG_RESOURCE_ID_MISSING | | Id de la Ressource manquante | Id della Risorsa mancante | Resource ID ontbreekt

MSG_RESOURCE_NOT_ALLOWED | No se permite el envío de un recurso para esta operación | Non autorisé à soumettre une ressource pour cette opération | Non è consentito sottomettere una risorsa per questa operazione | Niet toegestaan om een resource in te dienen voor deze bewerking | Nie można zgłosić zasobu dla tej operacji | Для данной операции отправка ресурса недопустима | □□□□□□□□□□

MSG_RESOURCE_REQUIRED | Se requiere un recurso | Une ressource est requise | E'' richiesta una risorsa | Een resource is verplicht | Zasób jest wymagany | Требуется ресурс | □□□□□□□□

MSG_RESOURCE_TYPE_MISMATCH | Los Tipos de los recursos no coinciden | Type de ressource incorrect | Tipo Risorsa non corrispondente | Verkeerd resourcetype | Niepoprawny typ zasobu | Несоответствие типа ресурса | □□□□□□□

MSG_SORT_UNKNOWN | Nombre del parámetro de ordenación "%s" desconocido | Nom du paramètre de tri inconnu "%s" | Nome del parametro di ordinamento "%s" non riconosciuto | Onbekende parameternaam "%s" voor sortering | Nieznany parametr sortowania "%s" | Неизвестное имя параметра сортировки "%s" | □□□□□□□□□"%s"

MSG_TRANSACTION_DUPLICATE_ID | Identificador duplicado en la transacción: %s | Identifiant en double dans la transaction : %s | | Dubbele identificatie in transactie: %s | Zdublowany identyfikator w transakcji: %s | Дублирующий идентификатор в транзакции: %s | □□□□□□□Id: %s

MSG_TRANSACTION_MISSING_ID | Identificador de la transacción no encontrado - se debe proporcionar un entry.id | Identifiant manquant dans la transaction - un élément entry.id doit être fourni | | Ontbrekende identificatie in transactie - entry.id is verplicht | Brak identyfikatora w transakcji - należy podać entry.id | Отсутствует идентификатор в транзакции - требуется entry.id | □□□□□Id - □□□□□□entry.id

MSG_UNHANDLED_NODE_TYPE | Tipo de nodo Xml no soportado "%s" | Type de noeud xml "%s" non traité | Tipo di nodo Xml non gestito "%s" | Kan xml nodetype "%s" niet verwerken | Nieobsługiwany typ węzła XML "%s" | Не обработанный xml узел "%s" | □□□□XML□□□□"%s"

MSG_UNKNOWN_CONTENT | Contenido desconocido (%s) en %s | Contenu inconnu (%s) à %s | Contenuto Sconosciuto (%s) at %s | Onbekende content (%s) at %s | Nieznana zawartość (%s) dla %s | Неизвестный контент (%s) в %s | □□□□ (%s) □□ %s

MSG_UNKNOWN_OPERATION | Operación http FHIR desconocida | operation http FHIR inconnue | operazione http FHIR sconosciuta | onbekende FHIR http operation | nieznana operacja FHIR http | неизвестная операция FHIR http | □□□FHIR HTTP□□

MSG_UNKNOWN_TYPE | Tipo de Recurso "%s" no reconocido | Type de ressource "%s" non reconnu | Tipo di Risorsa "%s" non riconosciuto | Resourcetype "%s" niet herkend | Nie rozpoznany typ zasobu: "%s" | Тип ресурса "%s" не распознан | □□□□"%s"□□□
MSG_UPDATED | Recurso existente actualizado | ressource existante mise à jour | risorsa esistente aggiornata | bestaande resource updated | uaktualniono istniejący zasób | существующий ресурс обновлён | □□□□□□□
MSG_VERSION_AWARE | Este recurso requiere actualizaciones en base a versiones | Des mises à jour en relation avec la version sont requises pour cette ressource | Questa risorsa richiede aggiornamenti per versione | Versie-bewuste updates zijn vereist voor deze resource | Uaktualnienia zakładające wersjonowanie są wymagane dla tego zasobu | Для данного ресурса необходимы обновления с учётом версии | □□□□□□□□□□□□
MSG_VERSION_AWARE_CONFLICT | Conflicto de actualizaciones (versión actual del servidor = "%s", versión del cliente referenciada = "%s") | Conflit de mise à jour (version courante du serveur = "%s", version référencée du client = "%s") | Conflitto nell''aggiornamento (attuale = "%s", quotato = "%s") | Updateconflict (huidige serverversie = "%s", opgegeven clientversie = "%s") | Konflikt podczas uaktualnienia (obecna wersja na serwerze = "%s", wersja wskazana przez klienta = "%s") | Конфликт обновления (текущая версия сервера = "%s", указанная версия клиента = "%s") | □□□□ (□□□□□□□ = "%s", □□□□□□□□ = "%s")
MSG_VERSION_AWARE_URL | URL especifica de la versión no reconocida | URL spécifique à une version non reconnue | URL specifico alla versione non riconosciuto | Versie-specifieke URL niet herkend | Nie rozpoznany URL specyficzny dla wersji | URL для указанной версии не распознан | □□□□□□□□URL
MSG_WRONG_NS | | | Dit lijkt geen FHIR element of resource te zijn (verkeerde namespace "%s")
SEARCH_MULTIPLE | Error: Multiples ocurrencias existen para %s parametros de búsqueda "%s" | Erreur : Plusieurs correspondances existent pour ce paramètre de recherche %s | Errore: Trovate corrispondenze multiple per %s parametri di ricerca "%s" | Fout: er is meer dan één resultaat voor %s zoekparameters "%s" | Błąd: Istnieją wielokrotne dopasowania dla %s parametrów wyszukiwania "%s" | Ошибка: множественные совпадения для %s с параметрами поиска "%s" | □□: □□ %s □□□□□ "%s"□□□□□□
SEARCH_NONE | Error: no se encontro una búsqueda apropiada para %s parametros de búsqueda "%s" | Erreur : aucune recherche trouvée pour les paramètres %s "%s" | Errore: non è stato trovato alcun parametro di ricerca processabile per %s parametri di ricerca "%s" | Fout: geen verwerkbare zoekactie gevonden voor %s zoekparameters "%s" | Błąd: Niewykonalne wyszukiwanie dla %s parametrów wyszukiwania "%s" | Ошибка: обрабатываемых результатов поиска для %s с параметрами поиска "%s" не найдено | □□: □%s□□□□"%s"□□□□□□□□□
UPDATE_MULTIPLE_MATCHES | | | Fout: er is meer dan één resultaat voor de conditionele update

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.493 Value Set http://hl7.org/fhir/ValueSet/allergy-intolerance-criticality

URL: https://hl7.org/fhir/R4/valueset-allergy-intolerance-criticality.html
Patient Care Work Group | Maturity Level: 3 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/allergy-intolerance-criticality
Version: | 4.0.1
Name: | AllergyIntoleranceCriticality
Title: | AllergyIntoleranceCriticality
Definition: | Estimate of the potential clinical harm, or seriousness, of a reaction to an identified substance.
Committee: | Patient Care Work Group
OID: | 2.16.840.1.113883.4.642.3.129 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for AllergyIntoleranceCriticality
• Resource: AllergyIntolerance.criticality (code / Required)

4.4.1.493.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/allergy-intolerance-criticality

4.4.1.493.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 3 concepts

Expansion based on http://hl7.org/fhir/allergy-intolerance-criticality version 4.0.1

All codes from system http://hl7.org/fhir/allergy-intolerance-criticality

Code | Display | Definition
low | Low Risk | Worst case result of a future exposure is not assessed to be life-threatening or having high potential for organ system failure.
high | High Risk | Worst case result of a future exposure is assessed to be life-threatening or having high potential for organ system failure.
unable-to-assess | Unable to Assess Risk | Unable to assess the worst case result of a future exposure.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.494 Value Set http://hl7.org/fhir/ValueSet/type-restful-interaction

URL: https://hl7.org/fhir/R4/valueset-type-restful-interaction.html
FHIR Infrastructure Work Group | Maturity Level: N | Normative (from v4.0.0) | Use Context: Any

| This page has been approved as part of an ANSI standard. See the Conformance Package for further details.

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/type-restful-interaction
Version: | 4.0.1
Name: | TypeRestfulInteraction
Title: | TypeRestfulInteraction
Definition: | Operations supported by REST at the type or instance level.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.180 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: CapabilityStatement.rest.resource.interaction.code (code / Required)

4.4.1.494.1 Content Logical Definition

This value set includes codes from the following code systems:

http://hl7.org/fhir/restful-interaction

Code | Display
read | read | Read the current state of the resource.
vread | vread | Read the state of a specific version of the resource.
update | update | Update an existing resource by its id (or create it if it is new).
patch | patch | Update an existing resource by posting a set of changes to it.
delete | delete | Delete a resource.
history-instance | history-instance | Retrieve the change history for a particular resource.
history-type | history-type | Retrieve the change history for all resources of a particular type.
create | create | Create a new resource with a server assigned id.
search-type | search-type | Search all resources of the specified type based on some filter criteria.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.495 Value Set http://hl7.org/fhir/ValueSet/immunization-reason

URL: https://hl7.org/fhir/R4/valueset-immunization-reason.html
Public Health and Emergency Response Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/immunization-reason
Version: | 4.0.1
Name: | ImmunizationReasonCodes
Title: | Immunization Reason Codes
Definition: | The value set to instantiate this attribute should be drawn from a terminologically robust code system that consists of or contains concepts to support describing the reason why a dose of vaccine was administered. This value set is provided as a suggestive example.
Committee: | Public Health and Emergency Response Work Group
OID: | 2.16.840.1.113883.4.642.3.290 (for OID based terminology systems)
Copyright: | This resource includes content from SNOMED Clinical Terms® (SNOMED CT®) which is copyright of the International Health Terminology Standards Development Organisation (IHTSDO). Implementers of these specifications must have the appropriate SNOMED CT Affiliate license - for more information contact http://www.snomed.org/snomed-ct/get-snomed-ct or info@snomed.org
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: Immunization.reasonCode (CodeableConcept / Example)

4.4.1.495.1 Content Logical Definition

This value set includes codes from the following code systems:

http://snomed.info/sct

Code | Display
429060002 | Procedure to meet occupational requirement
281657000 | Travel vaccinations

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.496 Value Set http://hl7.org/fhir/ValueSet/clinical-findings

URL: https://hl7.org/fhir/R4/valueset-clinical-findings.html
Orders and Observations Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/clinical-findings
Version: | 4.0.1
Name: | SNOMEDCTClinicalFindings
Title: | SNOMED CT Clinical Findings
Definition: | This value set includes all the "Clinical finding" SNOMED CT codes - concepts where concept is-a 404684003 (Clinical finding (finding)).
Committee: | Orders and Observations Work Group
OID: | 2.16.840.1.113883.4.642.3.232 (for OID based terminology systems)
Copyright: | This resource includes content from SNOMED Clinical Terms® (SNOMED CT®) which is copyright of the International Health Terminology Standards Development Organisation (IHTSDO). Implementers of these specifications must have the appropriate SNOMED CT Affiliate license - for more information contact http://www.snomed.org/snomed-ct/get-snomed-ct or info@snomed.org
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: Goal.description (CodeableConcept / Example)
• Resource: Goal.outcomeCode (CodeableConcept / Example)

- Resource: CareTeam.reasonCode (CodeableConcept / Example)
- Resource: FamilyMemberHistory.reasonCode (CodeableConcept / Example)
- Profile: FamilyMemberHistory-Genetic: FamilyMemberHistory.reasonCode (CodeableConcept / Example)
- Resource: Communication.reasonCode (CodeableConcept / Example)
- Resource: AllergyIntolerance.reaction.manifestation (CodeableConcept / Example)
- Resource: CarePlan.activity.detail.reasonCode (CodeableConcept / Example)
- Resource: DiagnosticReport.conclusionCode (CodeableConcept / Example)
- Profile: LipidProfile: DiagnosticReport.conclusionCode (CodeableConcept / Example)
- Profile: DiagnosticReport-Genetics: DiagnosticReport.conclusionCode (CodeableConcept / Example)
- Profile: HLAResult: DiagnosticReport.conclusionCode (CodeableConcept / Example)
- Resource: PlanDefinition.goal.description (CodeableConcept / Example)
- Profile: ShareablePlanDefinition: PlanDefinition.goal.description (CodeableConcept / Example)
- Profile: ComputablePlanDefinition: PlanDefinition.goal.description (CodeableConcept / Example)
- Profile: CDSHooksServicePlanDefinition: PlanDefinition.goal.description (CodeableConcept / Example)

4.4.1.496.1 Content Logical Definition

This value set includes codes from the following code systems:

- Include codes from http://snomed.info/sct where concept is-a 404684003 (Clinical finding)

4.4.1.496.2 Expansion

This expansion generated 01 Nov 2019

This value set has >1000 codes in it. In order to keep the publication size manageable, only a selection (1000 codes) of the whole set of codes is shown

All codes from system http://snomed.info/sct

Code	Display	Definition
109006	Anxiety disorder of childhood OR adolescence	
122003	Choroidal hemorrhage	
127009	Spontaneous abortion with laceration of cervix	
129007	Homoiothermia	
134006	Decreased hair growth	
140004	Chronic pharyngitis	
144008	Normal peripheral vision	
147001	Superficial foreign body of scrotum without major open wound but with infection	
150003	Abnormal bladder continence	
151004	Meningitis due to gonococcus	
162004	Severe manic bipolar I disorder without psychotic features	
165002	Accident-prone	
168000	Typhlolithiasis	
171008	Injury of ascending right colon without open wound into abdominal cavity	
172001	Endometritis following molar AND/OR ectopic pregnancy	
175004	Supraorbital neuralgia	
177007	Poisoning by sawfly larvae	
179005	Apraxia of dressing	
181007	Hemorrhagic bronchopneumonia	
183005	Autoimmune pancytopenia	
184004	Withdrawal arrhythmia	
188001	Intercostal artery injury	
192008	Congenital syphilitic hepatomegaly	
193003	Benign hypertensive renal disease	
195005	Illegal abortion with endometritis	
198007	Disease due to Filoviridae	
199004	Decreased lactation	
208008	Neurocutaneous melanosis sequence	
216004	Delusion of persecution	
219006	Alcohol user	
222008	Acute epiglottitis with obstruction	
223003	Tumor of body of uterus affecting pregnancy	
228007	Lucio phenomenon	
241006	Motor simple partial status	
242004	Noninfectious jejunitis	
253005	Sycosis	
257006	Acne rosacea, erythematous telangiectatic type	
258001	Pseudoknuckle pad	
264008	Blind hypertensive eye	

276008 | Oxytocin poisoning
279001 | Senile myocarditis
281004 | Alcoholic dementia
282006 | Acute myocardial infarction of basal-lateral wall
290006 | Melnick-Fraser syndrome
292003 | EEG finding
297009 | Acute myringitis
299007 | Paraffinoma of skin
303002 | Apoplectic pancreatitis
308006 | Pearly penile papules
310008 | Penile boil
313005 | Déjà vu
317006 | Reactive hypoglycemia
320003 | Cervical dilatation, 1cm
324007 | Plaster ulcer
330007 | Occipital headache
335002 | Pylorospasm
341009 | ABO incompatibility reaction
349006 | Absent tendon reflex
355001 | Hemorrhagic shock
357009 | Closed fracture trapezoid
358004 | Smallpox vaccine poisoning
359007 | Kernicterus due to isoimmunization
360002 | Acute radiation disease
364006 | Acute left-sided heart failure
366008 | Hidromeiosis
368009 | Heart valve disorder
369001 | Normal jugular venous pressure
378007 | Morquio syndrome
382009 | Legal history finding relating to child
383004 | Finding of passive range of hip extension
385006 | Secondary peripheral neuropathy
387003 | Melanuria
398002 | Left axis deviation greater than -90 degrees by EKG
407000 | Congenital hepatomegaly
408005 | Tooth chattering
409002 | Food allergy diet
426008 | Superficial injury of ankle without infection
431005 | Hypertrophy of scrotum
437009 | Abnormal composition of urine
440009 | Persistent hyperphenylalaninemia
442001 | Secondary hypopituitarism
443006 | Cystocele affecting pregnancy
447007 | Coach in sports activity accident
450005 | Ulcerative stomatitis
452002 | Blister of groin without infection
460001 | Squamous metaplasia of prostate gland
467003 | Old laceration of pelvic floor muscle
470004 | Vitreous touch syndrome
479003 | Graves' disease with pretibial myxedema AND with thyrotoxic crisis
486006 | Acute vascular insufficiency
488007 | Fibroid myocarditis
490008 | Upper respiratory tract hypersensitivity reaction
496002 | Closed traumatic dislocation of third cervical vertebra
504009 | Androgen-dependent hirsutism
517007 | Foreign body in hypopharynx
518002 | Multiple aggregation
520004 | Congenital bent nose
527001 | Spontaneous fetal evolution, Roederer's method
536002 | Glissonian cirrhosis
539009 | Conjunctival argyrosis
547009 | Hypersecretion of calcitonin
548004 | 13p partial trisomy syndrome
554003 | 2p partial trisomy syndrome
555002 | Dicentra species poisoning
563001 | Nystagmus
568005 | Habit disorder
586008 | Contact dermatitis due to primrose
590005 | Congenital aneurysm of anterior communicating artery
596004 | Premenstrual dysphoric disorder
599006 | Persistent pneumothorax
600009 | Pyromania

602001 | Ross river fever
607007 | Decreased vital capacity
610000 | Spastic aphonia
613003 | FRAXA - Fragile X syndrome
615005 | Obstruction due to foreign body accidentally left in operative wound AND/OR body cavity during a procedure
616006 | Sensorimotor disorder of eyelid
626004 | Hypercortisolism due to nonpituitary tumor
631002 | Transfusion reaction due to minor incompatibility
634005 | Saddle boil
640003 | Injury of pneumogastric nerve
643001 | Hypertrophy of lip
646009 | Idiopathic cyst of anterior chamber
649002 | Open fracture of distal end of ulna
651003 | Root work
652005 | Gangrenous tonsillitis
655007 | Abnormal fetal heart beat noted before labor in liveborn infant
658009 | Injury of colon without open wound into abdominal cavity
663008 | Pulmonary embolism following molar AND/OR ectopic pregnancy
664002 | Delayed ovulation
666000 | Poisoning by antivaricose drug AND/OR sclerosing agent
675003 | Torsion of intestine
682004 | Thrombosis complicating pregnancy AND/OR puerperium
685002 | Acquired telangiectasia of small AND/OR large intestines
701003 | Adult osteochondritis of spine
703000 | Congenital adhesion of tongue
714002 | Abrasion AND/OR friction burn of toe with infection
715001 | Nontraumatic rupture of urethra
718004 | Acute bronchiolitis with obstruction
733007 | Superficial foreign body of groin without major open wound but with infection
734001 | Opocephalus
736004 | Abscess of hip
750009 | Schistosoma mansoni infection
755004 | Postgastrectomy phytobezoar
756003 | Chronic rheumatic myopericarditis
758002 | Cyst of uterus
775008 | Open wound of head with complication
776009 | Partial arterial retinal occlusion
781000 | Cestrum diurnum poisoning
786005 | Clinical stage I B
787001 | Rheumatic mitral stenosis with regurgitation
788006 | Disease-related diet
792004 | CJD - Creutzfeldt-Jakob disease
799008 | Sigmoid colon ulcer
801006 | Insect bite, nonvenomous, of foot, infected
805002 | Pneumoconiosis due to silica
811004 | Flail motion
813001 | Ankle instability
815008 | Episcleritis
816009 | Genetic recombination
818005 | Third degree burn of multiple sites of lower limb
825003 | Superficial injury of axilla with infection
827006 | Late congenital syphilis, latent (+ sero., - C.S.F., 2 years OR more)
832007 | Moderate major depression
834008 | Chair-seated facing coital position
841002 | Congenital absence of skull bone
842009 | Consanguinity
843004 | Poliomyelomalacia
844005 | Finding relating to behavior
845006 | Inferior mesenteric artery injury
849000 | Total cataract
857002 | Erythema simplex
862001 | Anemia due to chlorate
865004 | Hyperalimentation formula for ileus
871005 | Contracted pelvis
874002 | Therapeutic diuresis
875001 | Chalcosis of eye
888003 | Foetal or neonatal effect of maternal blood loss
890002 | Deep third degree burn of elbow
899001 | Axis I diagnosis
903008 | Chorioretinal infarction
904002 | Pinard's sign
908004 | Superficial injury of interscapular region without infection

919001 | Pseudohomosexual state
928000 | Musculoskeletal disorder
931004 | Gestation period, 9 weeks
932006 | Flat affect
934007 | Thalassemia intermedia
943003 | Congenital retinal aneurysm
954008 | Renon-Delille syndrome
961007 | Erythema nodosum, acute form
962000 | Disability evaluation, disability 6%
964004 | Open wound of pharynx without complication
965003 | Toxic amblyopia
975000 | Anorectal agenesis
978003 | Chronic infantile eczema
981008 | Hemorrhagic proctitis
984000 | Perirectal cellulitis
987007 | Cellulitis of temple region
991002 | Wide QRS complex
998008 | Chagas' disease with heart involvement
1003002 | Religious discrimination
1020003 | Disease due to Nairovirus
1023001 | Apneic
1027000 | Biliary esophagitis
1031006 | Open wound of trachea without complication
1033009 | Thoracic arthritis
1034003 | Mesenteric-portal fistula
1038000 | Disaccharidurie
1045000 | Colonospasm
1046004 | Ureteritis glandularis
1051005 | Hyperplasia of islet alpha cells with gastrin excess
1055001 | Stenosis of precerebral artery
1059007 | Opisthorchiasis
1070000 | Facial myokymia
1073003 | Xeroderma pigmentosum group B
1074009 | Glucocorticoid-responsive primary hyperaldosteronism
1077002 | Septal infarction by EKG
1079004 | Macular retinal cyst
1085006 | Vulval candidiasis
1089000 | Congenital sepsis
1102005 | Intraerythrocytic parasitosis by Nuttallia
1107004 | Early latent syphilis, positive serology, negative cerebrospinal fluid, with relapse after treatment
1108009 | Female pattern alopecia
1111005 | Normal sebaceous gland activity
1112003 | Degenerative disorder of eyelid
1116000 | Chronic aggressive type B viral hepatitis
1124005 | Postpartum period, 6 days
1125006 | Septicemia during labor
1126007 | Knee locking
1131009 | Congenital valvular insufficiency
1134001 | Muehrcke lines
1135000 | Solar retinitis
1139006 | Confrontation (visual) test
1140008 | Thermal hypesthesia
1141007 | Circumoral paresthesia
1145003 | DSD - Developmental speech disorder
1150009 | Congenital microcheilia
1151008 | Constricted visual field
1152001 | Skin reaction negative
1155004 | Myocardial hypertrophy, determined by electrocardiogram
1156003 | Cavitary prostatitis
1168007 | Allotype
1184008 | Glasgow coma scale, 10
1192004 | Familial amyloid neuropathy, Finnish type
1194003 | Disease condition determination, well controlled
1196001 | Chronic bipolar II disorder, most recent episode major depressive
1197005 | Carbuncle of heel
1201005 | Benign essential hypertension
1203008 | Deep third degree burn of forehead AND/OR cheek with loss of body part
1207009 | Optic disc glaucomatous atrophy
1208004 | Gastroptosis
1212005 | Juvenile dermatomyositis
1214006 | Infection by Strongyloides
1230003 | No diagnosis on Axis 1

1232006 | Congenital articular rigidity with myopathy
1239002 | Congenital anteversion of femoral neck
1240000 | Lying prone
1259003 | Schistosis
1261007 | Multiple fractures of ribs
1264004 | Injury of descending left colon without open wound into abdominal cavity
1271009 | Knuckle pads, leuconychia and deafness
1280009 | Isologous chimera
1282001 | Laryngeal perichondritis
1283006 | Visual acuity less than .02 (1/60, count fingers 1 meter) or visual field less than 5 degrees, but better than 5.
1284000 | Abnormal jaw closure
1286003 | Vitamin K deficiency coagulation disorder
1287007 | Congenital absence of bile duct
1297003 | Infection by Cladosporium carrionii
1308001 | Complication of reimplant
1310004 | Impaired glucose tolerance associated with genetic syndrome
1317001 | Injury of ovary without open wound into abdominal cavity
1318006 | Post-translational genetic protein processing
1323006 | Kanamycin poisoning
1332008 | Conjugated visual deviation
1335005 | Peyronies disease
1343000 | DTA - Deep transverse arrest
1345007 | Hang nail
1351002 | Iliac artery injury
1356007 | Calculus of common duct with obstruction
1361009 | Leucocoria
1363007 | Fetal or neonatal effect of chronic maternal respiratory disease
1367008 | Injury of superior mesenteric artery
1370007 | Open fracture of metacarpal bone(s)
1372004 | Unicornate uterus
1376001 | Obsessive compulsive personality disorder
1378000 | Supination-eversion injury of ankle
1380006 | Agoraphobia without history of panic disorder with limited symptom attacks
1383008 | Hallucinogen induced mood disorder
1384002 | Diffuse cholesteatosis of middle ear
1386000 | Intracranial hemorrhage
1387009 | Solanum nigrum poisoning
1388004 | Metabolic alkalosis
1393001 | Lenz-Majewski dysplasia
1395008 | Complication of ultrasound therapy
1402001 | Frightened
1412008 | Anterior subcapsular polar cataract
1415005 | Inflammation of lymphatics
1418007 | Hypoplastic chondrodystrophy
1419004 | Injury of prostate without open wound into abdominal cavity
1426004 | Necrotizing glomerulonephritis
1427008 | Intraspinal abscess
1430001 | Intracranial hemorrhage following injury without open intracranial wound AND with prolonged loss of consciousness (more than 24 hours) without return to pre-existing level
1447000 | Icthyoparasitism
1469007 | Miscarriage with urinary tract infection
1474004 | Hypertensive heart AND renal disease complicating AND/OR reason for care during childbirth
1475003 | Herpes labialis
1478001 | Obliteration of lymphatic vessel
1479009 | 20q partial trisomy syndrome
1482004 | Chalazion
1486001 | Orbital congestion
1488000 | PONV - Postoperative nausea and vomiting
1489008 | External hordeolum
1492007 | Congenital anomaly of large intestine
1493002 | Acute endophthalmitis
1499003 | Bipolar I disorder, single manic episode with postpartum onset
1512006 | Congenital stricture of bile duct
1515008 | Gorham disease
1518005 | Splenitis
1519002 | Congenital phlebectasia
1521007 | Blister of buttock without infection
1523005 | Clinical stage IV B
1525003 | Blister of foot without infection
1531000 | Nitrofuran derivative poisoning
1532007 | Viral pharyngitis
1534008 | Palsy of conjugate gaze

1538006 | Central nervous system malformation in foetus affecting obstetrical care
1539003 | Nodular tendinous disease of finger
1542009 | Omphalocele with obstruction
1544005 | Open dislocation of knee
1551001 | Osteomyelitis of femur
1556006 | Clark melanoma level 4
1563006 | Protein S deficiency
1567007 | Chronic gastric ulcer without hemorrhage, without perforation AND without obstruction
1588003 | Heterosexual precocious puberty
1592005 | Failed attempted termination of pregnancy with uremia
1593000 | Infantile hemiplegia
1606009 | Infection caused by Macracanthorhynchus hirudinaceus
1608005 | Increased capillary permeability
1639007 | Abnormality of organs AND/OR soft tissues of pelvis affecting pregnancy
1647007 | Primaquine poisoning
1648002 | Lymphocytic pseudotumor of lung
1654001 | Steroid-induced glaucoma
1657008 | Toxic effect of phosdrin
1658003 | Closed fracture clavicle, lateral end
1663004 | Tumor grade G2
1667003 | Early fontanel closure
1670004 | Cerebral hemiparesis
1671000 | Sago spleen
1674008 | Meesman's epithelial corneal dystrophy
1679003 | Arthritis associated with another disorder
1682008 | Transitory amino acid metabolic disorder
1685005 | Rat-bite fever
1686006 | Sedative, hypnotic AND/OR anxiolytic-induced anxiety disorder
1694004 | Accessory lobe of lung
1698001 | Ulcer of bile duct
1703007 | Increased leg circumference
1705000 | Closed fracture of base of neck of femur
1708003 | Open dislocation of clavicle
1714005 | Photokeratitis
1717003 | Guttate hypomelanosis
1723008 | Urethral stricture due to schistosomiasis
1724002 | Infection caused by Crenosoma
1734006 | Fracture of vertebral column with spinal cord injury
1735007 | Thrill
1739001 | Occipital fracture
1742007 | Female hypererotism
1744008 | Connation of teeth
1748006 | Thrombophlebitis of deep femoral vein
1755008 | Healed coronary
1761006 | Biliary cirrhosis
1763009 | Stromal keratitis
1767005 | Fisher syndrome
1769008 | Thoracodidymus
1771008 | Insulin biosynthesis defect
1776003 | RTA - Renal tubular acidosis
1777007 | Increased molecular dissociation
1778002 | Malocclusion due to abnormal swallowing
1779005 | OFD II - Orofacial-digital syndrome II
1794009 | Idiopathic corneal edema
1816003 | Panic disorder with agoraphobia, severe agoraphobic avoidance AND mild panic attacks
1821000 | Chemoreceptor apnea
1822007 | Impaired glucose tolerance associated with pancreatic disease
1824008 | Allergic gastritis
1826005 | Granuloma of lip
1828006 | Infestation caused by Gasterophilus hemorrhoidalis
1829003 | Microcephalus
1833005 | Phacoanaphylactic endophthalmitis
1835003 | Necrosis of pancreas
1837006 | Orciprenaline poisoning
1845001 | Paraparesis
1847009 | Endophthalmitis
1848004 | Poisoning caused by gaseous anesthetic
1852004 | Traumatic injury of sixth cranial nerve
1855002 | Developmental academic disorder
1856001 | Accessory nose
1857005 | Congenital rubella syndrome
1858000 | Infection caused by Stilesia globipunctata

1860003 | Fluid volume disorder
1865008 | Impaired intestinal protein absorption
1869002 | Rupture of iris sphincter
1881003 | Increased nutritional requirement
1892002 | Star figure at the macula
1896004 | Ectopic breast tissue
1897008 | Amsinckia species poisoning
1899006 | Autosomal hereditary disorder
1903004 | Infestation caused by Psorergates ovis
1908008 | von Willebrand disease, type IIC
1909000 | Impairment level: better eye: severe impairment: lesser eye: near-total impairment
1922008 | Congenital absence of urethra
1926006 | Osteopetrosis
1938002 | Emotional AND/OR mental disease in mother complicating pregnancy, childbirth AND/OR puerperium
1939005 | Abnormal vascular flow
1943009 | Left-right confusion
1953005 | Congenital deficiency of pigment of skin
1954004 | Dilated cardiomyopathy secondary to toxic reaction
1955003 | Preauricular pit
1959009 | Encephalartos species poisoning
1961000 | Chronic polyarticular juvenile rheumatoid arthritis
1963002 | PNH - Paroxysmal nocturnal hemoglobinuria
1965009 | Normal skin pH
1967001 | Congenital absence of forearm only
1973000 | Sedative, hypnotic AND/OR anxiolytic-induced psychotic disorder with delusions
1977004 | Oxymetholone poisoning
1979001 | Focal choroiditis
1980003 | Seromucinous otitis media
1981004 | Urhidrosis
1988005 | Late effect of injury to nerve roots, spinal plexus AND/OR other nerves of trunk
1989002 | Burn of vagina AND/OR uterus
2004005 | Normotensive
2012002 | Fracture of lunate
2024009 | Dilated cardiomyopathy secondary to metazoal myocarditis
2028007 | Erythema induratum
2032001 | Cerebral edema
2036003 | Acquired factor VII deficiency disease
2040007 | Neurogenic thoracic outlet syndrome
2041006 | Eunuchoid gigantism
2043009 | Alcoholic gastritis
2053005 | Late effect of injury to blood vessels of thorax, abdomen AND/OR pelvis
2055003 | Recurrent erosion syndrome
2058001 | Bilateral loss of labyrinthine reactivity
2061000 | Conductive hearing loss of combined sites
2065009 | Autosomal dominant optic atrophy
2066005 | Gastric ulcer with hemorrhage AND perforation but without obstruction
2070002 | Burning sensation in eye
2073000 | Delusions
2087000 | Pulmonary nocardiosis
2089002 | Pagets disease of bone
2091005 | Pharyngeal diverticulitis
2094002 | Carbon disulfide causing toxic effect
2102007 | Deep corneal vascularization
2103002 | Reflex sympathetic dystrophy of upper extremity
2107001 | Anisomelia
2109003 | Isolated somatotropin deficiency
2114004 | Infection caused by Cysticercus pisiformis
2116002 | Intramembranous bone formation
2120003 | Weak cry
2121004 | Ethopropazine poisoning
2128005 | Disorder of adenoid
2129002 | Edema of pharynx
2132004 | Meconium in amniotic fluid noted before labor in liveborn infant
2134003 | Diffuse pain
2136001 | Open wound of jaw with complication
2138000 | LSP - Left sacroposterior position
2145000 | Anal intercourse
2149006 | Decreased hormone production
2158004 | Infection caused by Contracaecum
2167004 | Retinal hemangioblastomatosis
2169001 | Thoracic radiculitis
2170000 | Gallop rhythm

2176006 | Halogen acne
2177002 | PHN - Post-herpetic neuralgia
2186007 | Compensated metabolic alkalosis
2198002 | Visceral epilepsy
2202000 | Open posterior dislocation of distal end of femur
2204004 | Acquired deformity of pinna
2213002 | Congenital anomaly of vena cava
2216005 | Nocturnal emission
2217001 | Superficial injury of perineum without infection
2219003 | Disability evaluation, disability 100%
2224000 | Selenium poisoning
2228002 | Scintillating scotoma
2229005 | Chimera
2231001 | Nerve plexus disorder
2237002 | Painful breathing -pleurodynia
2239004 | Previous pregnancies 6
2241003 | X-linked absence of thyroxine-binding globulin
2243000 | Hypercalcemia due to hyperthyroidism
2245007 | Foreign body in nasopharynx
2251002 | Primary hypotony of eye
2256007 | Monovular twins
2261009 | Obstetrical pulmonary fat embolism
2268003 | Victim of homosexual aggression
2284002 | Pulsating exophthalmos
2295008 | Closed fracture of upper end of forearm
2296009 | Iron dextran toxicity
2298005 | Focal facial dermal dysplasia
2301009 | Psychosomatic factor in physical condition, psychological component of unknown degree
2303007 | Inguinal hernia with gangrene
2304001 | Intervertebral discitis
2307008 | Peripancreatic fat necrosis
2308003 | Silent alleles
2312009 | Reactive attachment disorder of infancy OR early childhood, inhibited type
2314005 | Unprotected intercourse
2326000 | Marriage annulment
2339001 | Sexual overexposure
2341000 | Infection caused by Moniliformis
2351004 | Genetic transduction
2355008 | Rud syndrome
2359002 | Hyper-beta-alaninemia
2365002 | Simple chronic pharyngitis
2366001 | Late effect of dislocation
2367005 | Acute hemorrhagic gastritis
2374000 | Monofascicular block
2385003 | Cellulitis of pectoral region
2388001 | Normal variation in translucency
2390000 | Acute gonococcal vulvovaginitis
2391001 | Achondrogenesis
2396006 | Malignant pyoderma
2398007 | Quinidine toxicity by electrocardiogram
2403008 | Dependence syndrome
2415007 | Lumbosacral root lesion
2418009 | Polyester fume causing toxic effect
2419001 | Open wound of forehead with complication
2420007 | Third degree burn of multiple sites of upper limb
2432006 | Cerebrospinal fluid circulation disorder
2435008 | Ascaridiasis
2437000 | Placenta circumvallata
2438005 | Iniencephaly
2439002 | Purulent endocarditis
2443003 | Hydrogen sulfide poisoning
2452007 | Fetal rotation
2463005 | Acquired heterochromia of iris
2469009 | Onychomalacia
2470005 | Brain damage
2471009 | Intra-abdominal abscess postprocedure
2472002 | Passes no urine
2473007 | Intermittent vertical squint
2477008 | Superficial phlebitis
2492009 | Disorder of nutrition
2495006 | Congenital cerebral arteriovenous aneurysm
2496007 | Acalculia

2506003 | Early onset dysthymia
2513003 | Tinea capitis caused by Trichophyton
2518007 | Cryptogenic sexual precocity
2521009 | Bone conduction better than air
2523007 | Salmonella pneumonia
2526004 | Noninflammatory disorder of the female genital organs
2528003 | Viremia
2532009 | Choroidal rupture
2534005 | Congenital absence of vena cava
2538008 | Ketosis
2541004 | Compulsive buying
2554006 | Acute purulent pericarditis
2556008 | Disease of supporting structures of teeth
2560006 | Complex syndactyly of fingers
2562003 | Athanasia trifurcata poisoning
2576002 | Trachoma
2581006 | Clasp knife rigidity
2582004 | Deep third degree burn of multiple sites of lower limb
2583009 | Filigreed network of venous valves
2584003 | Cerebral degeneration in childhood
2585002 | Pneumococcal pleurisy
2589008 | Acute dacryoadenitis
2591000 | Crush injury of shoulder region
2593002 | Dubowitz syndrome
2602008 | Hemarthrosis of shoulder
2606006 | Boil of perineum
2615004 | Graafian follicle cyst
2618002 | Chronic recurrent major depressive disorder
2622007 | Infected ulcer of skin
2624008 | Prepubertal periodontitis
2625009 | Senter syndrome
2630008 | Open wound of finger without complication
2634004 | Decreased blood erythrocyte volume
2638001 | Hypercalcemia caused by a drug
2640006 | Clinical stage 4
2651006 | Closed traumatic dislocation of elbow joint
2655002 | Invalidism
2657005 | Overflow proteinuria
2663001 | Palpatory proteinuria
2665008 | Coordinate convulsion
2683000 | Nonvenomous insect bite of axilla without infection
2689001 | Dominant dystrophic epidermolysis bullosa with absence of skin
2694001 | Myelophthisic anemia
2704003 | Acute disease
2707005 | Necrotizing enterocolitis
2713001 | Closed pneumothorax
2724004 | Auditory recruitment
2725003 | Previous abnormality of glucose tolerance
2733002 | Heel pain
2736005 | Honeycomb atrophy of face
2740001 | Gouty proteinuria
2749000 | Congenital deformity of hip
2751001 | Fibrocalculous pancreatic diabetes
2761008 | Decreased stool caliber
2764000 | Joint crackle
2770006 | Fetal or neonatal effect of antibiotic transmitted via placenta and/or breast milk
2772003 | Epidermolysis bullosa acquisita
2775001 | Intra-articular loose body
2776000 | Organic brain syndrome
2781009 | Miscarriage complicated by delayed and/or excessive hemorrhage
2782002 | Temporomandibular dysplasia
2783007 | Gastrojejunal ulcer without hemorrhage AND without perforation
2786004 | Epithelial ovarian tumor, International Federation of Gynecology and Obstetrics stage III
2790002 | Impairment level: one eye: total impairment: other eye: not specified
2805007 | Phosmet poisoning
2806008 | Impaired psychomotor development
2807004 | Chronic gastrojejunal ulcer with perforation
2808009 | Infection caused by Prosthenorchis elegans
2815001 | Sexual pyromania
2816000 | Dilated cardiomyopathy secondary to myotonic dystrophy
2818004 | Congenital vascular anomaly of eye
2819007 | Magnesium sulfate poisoning

2825006 | Abrasion and/or friction burn of gum without infection
2828008 | Congenital stenosis of nares
2829000 | Uhl disease
2831009 | Pyloric antral vascular ectasia
2835000 | Hemolytic anemia due to cardiac trauma
2836004 | Butane causing toxic effect
2838003 | Piblokto
2840008 | Open fracture of vault of skull with cerebral laceration AND/OR contusion
2850009 | Infection caused by Schistosoma incognitum
2853006 | Macular keratitis
2856003 | Vitamin A-responsive dermatosis
2858002 | Postpartum sepsis
2884008 | Spherophakia-brachymorphia syndrome
2893009 | Anomaly of chromosome pair 10
2897005 | Immune thrombocytopenia
2899008 | Thought blocking
2900003 | Fibromuscular dysplasia of renal artery
2901004 | Altered blood passed per rectum
2902006 | Decreased lymphocyte life span
2904007 | Male infertility
2910007 | Discharge from penis
2912004 | Cystic-bullous disease of the lung
2917005 | Transient hypothyroidism
2918000 | Infection caused by Bacteroides
2919008 | Nausea, vomiting and diarrhea
2929001 | Arterial occlusion
2935001 | Antiasthmatic poisoning
2940009 | Intrabasal vesicular dermatitis
2946003 | Osmotic diarrhea
2951009 | Atopic cataract
2955000 | Chronic ulcerative pulpitis
2965006 | Nevoid congenital alopecia
2967003 | Non-comitant strabismus
2972007 | Occlusion of anterior spinal artery
2973002 | Pelvic organ injury without open wound into abdominal cavity
2978006 | Aneurysm of conjunctiva
2981001 | Pulsatile mass of abdomen
2989004 | Complication following molar AND/OR ectopic pregnancy
2990008 | Lymphocytic leukemoid reaction
2992000 | Pigmentary pallidal degeneration
2994004 | Brain fag
2999009 | Injury of ear region
3002002 | Thyroid hemorrhage
3004001 | Congenital dilatation of esophagus
3006004 | Altered consciousness
3009006 | Solanum malacoxylon poisoning
3013004 | Open wound of ear drum without complication
3014005 | Autoeczematization
3018008 | Penetration of eyeball with magnetic foreign body
3019000 | Closed anterior dislocation of elbow
3021005 | Normal gastric acidity
3023008 | Acute peptic ulcer without hemorrhage, without perforation AND without obstruction
3032005 | Nonvenomous insect bite of cheek without infection
3033000 | Bone AND/OR joint disorder of pelvis in mother complicating pregnancy, childbirth AND/OR puerperium
3038009 | Acute lymphangitis of umbilicus
3044008 | Vitreous prolapse
3053001 | Poisoning caused by nitroglycerin
3059002 | Acute lymphangitis of thigh
3067005 | Weak C phenotype
3071008 | Widow
3072001 | Hormone-induced hypopituitarism
3073006 | Ruvalcaba syndrome
3084004 | Nonvenomous insect bite of gum without infection
3089009 | Disability evaluation, impairment, class 7
3094009 | Vomiting in infants AND/OR children
3095005 | Induced malaria
3097002 | Superficial injury of lip with infection
3098007 | Ventricular septal rupture
3105002 | Intron
3109008 | Secondary dysthymia early onset
3110003 | AOM - Acute otitis media
3119002 | Brain stem laceration with open intracranial wound AND loss of consciousness

3129009 | Infarction of ovary
3135009 | OE - Otitis externa
3140001 | Citrullinemia, subacute type
3144005 | Staphylococcal pleurisy
3158007 | Panic disorder with agoraphobia, agoraphobic avoidance in partial remission AND panic attacks in partial remission
3160009 | Infertility of cervical origin
3163006 | Acute adenoviral follicular conjunctivitis
3168002 | Thrombophlebitis of intracranial venous sinus
3185000 | Mood-congruent delusion
3199001 | Sprain of shoulder joint
3200003 | Sacrocoxalgia
3208005 | Open wound of ossicles without complication
3214003 | Invasive pulmonary aspergillosis
3217005 | Open dislocation of sixth cervical vertebra
3218000 | Mycotic disease
3219008 | Disease type AND/OR category unknown
3228009 | Closed fracture of the radial shaft
3229001 | Tracheal ulcer
3230006 | Illegal termination of pregnancy with afibrinogenemia
3238004 | Pericarditis
3239007 | Lymphocyte disorder
3253007 | Dyschromia
3254001 | Infection caused by Strongyloides westeri
3259006 | Homeria species poisoning
3261002 | Migratory osteolysis
3263004 | Verumontanitis
3272007 | Stomatocytosis
3274008 | Flat chest
3275009 | Behcet syndrome, vascular type
3276005 | Toad poisoning
3277001 | Terminal mood insomnia
3282008 | Arc eye
3283003 | Feeling of sand or foreign body in eye
3286006 | Patient status determination, greatly improved
3289004 | Anisometropia
3291007 | Closed fracture of two ribs
3298001 | Amnestic syndrome
3303004 | Disease caused by Arenavirus
3304005 | Bronchial compression
3305006 | Disorder of lymphatic vessel
3308008 | Atrophic-hyperplastic gastritis
3310005 | Foreign body granuloma of skin
3321001 | Renal abscess
3323003 | Leukoplakia of penis
3327002 | Acquired jerk nystagmus
3331008 | Open fracture of neck of metacarpal bone
3344003 | Toxic labyrinthitis
3345002 | Idiopathic osteoporosis
3355003 | Anti-common cold drug poisoning
3358001 | Lichen ruber moniliformis
3368006 | Dull chest pain
3376008 | Pseudoptyalism
3381004 | Open fracture of astragalus
3387000 | Auditory discrimination aphasia
3391005 | Negative for tumor cells
3393008 | Phlebitis following infusion, perfusion AND/OR transfusion
3398004 | Cadmium poisoning
3401001 | Cercopithecus herpesvirus 1 disease
3415004 | Cyanosis
3419005 | Faucial diphtheria
3421000 | Open blow-out fracture orbit
3424008 | Heart rate fast
3426005 | Retained magnetic intraocular foreign body
3427001 | Nonglucosuric melituria
3434004 | Myotonia
3439009 | Severe combined immunodeficiency (SCID) due to absent peripheral T cell maturation
3441005 | Disorder of sebaceous gland
3446000 | Open fracture of T7-T12 level with spinal cord injury
3449007 | Finger agnosia
3456001 | Chronic progressive non-hereditary chorea
3458000 | Myositis ossificans associated with dermato / polymyositis
3461004 | Deep third degree burn of thumb

3464007 | Infection caused by Oesophagostomum dentatum
3468005 | Neonatal infective mastitis
3469002 | Partial thickness burn of thumb
3472009 | Spondylolisthesis, grade 4
3474005 | Glycine max poisoning
3480002 | Burn of wrist
3482005 | Postoperative esophagitis
3483000 | Chronic peptic ulcer with perforation
3487004 | Pulmonary candidiasis
3500002 | Open wound of ossicles with complication
3502005 | Cervical lymphadenitis
3503000 | Gender identity disorder of adolescence, previously asexual
3505007 | Nonallopathic lesion of the arm
3506008 | Stenosis of retinal artery
3507004 | Abscess of thigh
3511005 | Infectious thyroiditis
3514002 | Peribronchial fibrosis of lung
3519007 | Disorder of synovium
3528008 | Restricted carbohydrate fat controlled diet
3529000 | Infection caused by Sanguinicola
3530005 | Bipolar 1 disorder, single manic episode, full remission
3531009 | Intrapsychic conflict
3533007 | Acute palmoplantar pustular psoriasis
3539006 | Enteromenia
3542000 | Laceration extending into parenchyma of spleen with open wound into abdominal cavity
3544004 | Hair-splitting
3545003 | Diastolic dysfunction
3548001 | Brachial plexus disorder
3549009 | Pancreatic acinar atrophy
3558002 | Mesenteric infarction
3560000 | Bilateral recurrent inguinal hernia
3570003 | Increased blood erythrocyte volume
3571004 | Megaloblastic anemia due to pancreatic insufficiency
3577000 | Lattice retinal degeneration
3585009 | Blinking
3586005 | Psychogenic fugue
3589003 | Syphilitic pericarditis
3590007 | Enteroenteric fistula
3591006 | Metabolic acidosis, normal anion gap, bicarbonate losses
3598000 | Partial recent retinal detachment with single defect
3611003 | Demeton poisoning
3633001 | Abscess of hand
3634007 | Legal termination of pregnancy complicated by metabolic disorder
3639002 | Glossoptosis
3640000 | Late effect of traumatic amputation
3641001 | Infection caused by Coenurosis serialis
3642008 | Steryl-sulfate sulfohydrolase deficiency
3644009 | Macerated skin
3649004 | Contusion, multiple sites of trunk
3650004 | Congenital absence of liver,total
3652007 | Overproduction of growth hormone
3657001 | Osteospermum species poisoning
3660008 | Lethal glossopharyngeal defect
3662000 | Rolling hiatus hernia
3677008 | Academic problem
3680009 | Monocephalus tripus dibrachius
3681008 | Thrombophlebitis of torcular Herophili
3696007 | Functional dyspepsia
3699000 | Transverse deficiency of arm
3703002 | Ischiatic hernia with gangrene
3704008 | Diffuse endocapillary proliferative glomerulonephritis
3705009 | Congenital malformation of anterior chamber of eye
3712000 | Degenerated eye
3716002 | Thyroid goiter
3720003 | Abnormal presence of hemoglobin
3723001 | Joint inflammation
3733009 | Congenital eventration of right crus of diaphragm
3736001 | Open wound of thumbnail with tendon involvement
3738000 | VH - Viral hepatitis
3744001 | Hyperlipoproteinemia
3745000 | Sleep rhythm problem
3747008 | EC - Ejection click

3750006 | Arteriospasm
3751005 | Contusion of labium
3752003 | Infection by Trichuris
3754002 | Dysplasia of vagina
3755001 | PRP - Pityriasis rubra pilaris
3756000 | Static ataxia
3759007 | Injury of heart with open wound into thorax
3760002 | Familial multiple factor deficiency syndrome, type V
3762005 | Bilateral recurrent femoral hernia with gangrene
3763000 | Expected bereavement due to life event
3783004 | Enamel pearls
3797007 | Periodontal cyst
3798002 | Premature birth of identical twins, both stillborn
3815005 | Crohn disease of rectum
3820005 | Inner ear conductive hearing loss
3827008 | Aneurysm of artery of neck
3830001 | Subcutaneous emphysema
3841004 | Blister of cheek with infection
3845008 | Duplication of intestine
3855007 | Disorder of pancreas
3859001 | Late effect of open wound of extremities without tendon injury
3873005 | Failed attempted termination of pregnancy with acute necrosis of liver
3885002 | ABO isoimmunization in pregnancy
3886001 | Congenital fecaliths
3899003 | Neutropenic typhlitis
3900008 | Mixed sensory-motor polyneuropathy
3902000 | Non dose-related drug-induced neutropenia
3903005 | Closed traumatic pneumothorax
3908001 | Infestation caused by Haematopinus
3909009 | Coeur en sabot
3913002 | Injury of gastrointestinal tract with open wound into abdominal cavity
3914008 | Mental disorder in childhood
3928002 | Zika virus disease
3939004 | Bacterial colony density, transparent
3944006 | X-linked placental steryl-sulfatase deficiency
3947004 | High oxygen affinity hemoglobin polycythemia
3950001 | Birth
3951002 | Proctitis
3972004 | Idiopathic insomnia
3975002 | Deep third degree burn of lower limb
3978000 | AIHA - Warm autoimmune hemolytic anemia
3987009 | Congenital absence of trachea
3993001 | Infection caused by Muellerius
3999002 | Acute pyelitis without renal medullary necrosis
4003003 | Alphavirus disease
4004009 | Monster with cranial anomalies
4006006 | Foetal tachycardia affecting management of mother
4009004 | Lower urinary tract infection
4016003 | Empyema of mastoid
4017007 | Increased stratum corneum adhesiveness
4022007 | Vulvitis circumscripta plasmacellularis
4026005 | Interstitial mastitis associated with childbirth
4030008 | Le Dantec virus disease
4038001 | Myrotheciotoxicosis
4039009 | Multiple vitamin deficiency disease
4040006 | Hassall-Henle bodies
4041005 | Congenital anomaly of macula
4046000 | Degenerative spondylolisthesis
4062006 | Lumbosacral plexus lesion
4063001 | Achillodynia
4069002 | Anoxic brain damage during AND/OR resulting from a procedure
4070001 | Palinphrasia
4075006 | Peganum harmala poisoning
4082005 | Syphilitic myocarditis
4088009 | Acquired hydrocephalus
4089001 | Meningococcemia
4092002 | Nonallopathic lesion of costovertebral region
4103001 | Complex partial seizure
4106009 | Rotator cuff rupture
4107000 | Infertile male syndrome
4113009 | Arrested hydrocephalus
4120002 | Bronchiolitis

4124006 | Insect bite, nonvenomous, of vagina, infected
4127004 | Prostatic obstruction
4129001 | Argyll-Robertson pupil
4135001 | 11p partial monosomy syndrome
4136000 | Macrodactylia of toes
4142001 | Muscular asthenopia
4152002 | Acquired hypoprothrombinemia
4160001 | Congenital anomaly of upper respiratory system
4168008 | Tibial plateau chondromalacia
4170004 | Ehlers-Danlos syndrome, procollagen proteinase resistant
4174008 | Tripartite placenta
4175009 | Infestation by Estrus
4178006 | Partial recent retinal detachment with multiple defects
4181001 | Normal peak expiratory flow rate
4183003 | Charcot-Marie-Tooth disease, type IC
4184009 | Congenital malformation of the endocrine glands
4191007 | Scaphoid head
4195003 | Duplication of anus
4197006 | Disability evaluation, impairment, class 5
4199009 | 18p partial trisomy syndrome
4208000 | Closed multiple fractures of both lower limbs
4210003 | OH - Ocular hypertension
4223005 | Parkinsonism caused by drug
4224004 | Complication of infusion
4225003 | Nasal tuberculosis
4229009 | Phthisical eye
4232007 | Chronic vulvitis
4237001 | Suppurative pulpitis
4240001 | Rupture of aorta
4241002 | Listeria infection
4242009 | 18q partial monosomy syndrome
4244005 | Urticaria neonatorum
4248008 | Synovitis AND/OR tenosynovitis associated with another disease
4249000 | Poor peripheral circulation
4251001 | Internal eye sign
4260009 | Sacral spinal cord injury without bone injury
4262001 | Phlebitis of superior sagittal sinus
4264000 | Chronic pericoronitis
4269005 | Chronic gastrojejunal ulcer without hemorrhage AND without perforation
4273008 | Closed posterior dislocation of elbow
4275001 | Conjugate gaze spasm
4278004 | Superficial foreign body of axilla without major open wound but with infection
4283007 | Mirizzi syndrome
4287008 | Chordee of penis
4294006 | Isosexual precocious puberty
4300009 | Deep third degree burn of forearm
4301008 | Autoimmune state
4306003 | Cluster B personality disorder
4307007 | Pregestational diabetes mellitus AND/OR impaired glucose tolerance, modified White class F
4308002 | RSIS - Repetitive strain injury syndrome
4310000 | Third degree burn of wrist AND/OR hand
4313003 | Acardiacus anceps
4316006 | Myometritis
4320005 | Factor V deficiency
4324001 | Subacute cystitis
4325000 | 11q partial monosomy syndrome
4332009 | Subarachnoid hemorrhage following injury without open intracranial wound AND with concussion
4338008 | Arnold nerve reflex cough syndrome
4340003 | Acrodermatitis chronica atrophicans
4349002 | Open fracture of multiple sites of metacarpus
4354006 | Open dislocation of scapula
4356008 | Gingival soft tissue recession
4359001 | Early congenital syphilis
4364002 | Structure of associations
4367009 | Hoover sign
4373005 | Clubbing of nail
4374004 | TV - Congenital tricuspid valve abnormality
4381006 | Verbal paraphasia
4386001 | Bronchospasm
4390004 | Chronic lithium nephrotoxicity
4397001 | Partial congenital duodenal obstruction
4399003 | Acute hemorrhagic pancreatitis

4403007 | Exclamation point hair
4406004 | Congenital anomaly of male genital system
4409006 | Combined methylmalonic acidemia and homocystinuria due to defects in adenosylcobalamin and methylcobalamin synthesis
4410001 | Retroperitoneal hernia with obstruction
4412009 | Digital nerve injury
4414005 | Infection caused by Setaria
4416007 | Heerfordt syndrome
4418008 | Gangrenous ergotism
4426000 | Ten previous induced terminations of pregnancy
4434006 | BS - Bloom syndrome
4439001 | Axenfeld-Schurenberg syndrome
4441000 | Severe bipolar disorder with psychotic features
4445009 | TB - Urogenital tuberculosis
4448006 | Allergic headache
4451004 | Illegal termination of pregnancy with renal tubular necrosis
4461006 | Complication of administrative procedure
4463009 | Indiana-Maryland type amyloid polyneuropathy
4464003 | Rocio virus disease
4465002 | Spherophakia
4468000 | Oppenheim gait
4470009 | Blanching of skin
4473006 | Migraine with aura
4477007 | Juvenile myopathy AND lactate acidosis
4478002 | Multiple fractures of upper AND lower limbs
4481007 | Abnormal gastric secretion regulation
4483005 | Syphilitic punched out ulcer

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.497 Value Set http://hl7.org/fhir/ValueSet/immunization-subpotent-reason

URL: https://hl7.org/fhir/R4/valueset-immunization-subpotent-reason.html
Public Health and Emergency Response Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined at http://www.hl7.org/Special/committees/pher/index.cfm .

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/immunization-subpotent-reason
Version: | 4.0.1
Name: | ImmunizationSubpotentReason
Title: | Immunization Subpotent Reason
Definition: | The value set to instantiate this attribute should be drawn from a terminologically robust code system that consists of or contains concepts to support describing the reason why a dose is considered to be subpotent. This value set is provided as a suggestive example.
Committee: | Public Health and Emergency Response Work Group
OID: | 2.16.840.1.113883.4.642.3.993 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ImmunizationSubpotentReason

• Resource: Immunization.subpotentReason (CodeableConcept / Example)

4.4.1.497.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/immunization-subpotent-reason

4.4.1.497.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 3 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/immunization-subpotent-reason version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/immunization-subpotent-reason

Code | Display | Definition
partial | Partial Dose | The full volume of the dose was not administered to the patient.
coldchainbreak | Cold Chain Break | The vaccine experienced a cold chain break.
recall | Manufacturer Recall | The vaccine was recalled by the manufacturer.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.498 Value Set http://hl7.org/fhir/ValueSet/organization-type

URL: https://hl7.org/fhir/R4/valueset-organization-type.html
Patient Administration Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/organization-type
Version: | 4.0.1
Name: | OrganizationType
Title: | Organization type
Definition: | This example value set defines a set of codes that can be used to indicate a type of organization.
Committee: | Patient Administration Work Group
OID: | 2.16.840.1.113883.4.642.3.414 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for OrganizationType
• Resource: Organization.type (CodeableConcept / Example)

4.4.1.498.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/organization-type

4.4.1.498.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 12 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/organization-type version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/organization-type

Code | Display | Definition
prov | Healthcare Provider | An organization that provides healthcare services.
dept | Hospital Department | A department or ward within a hospital (Generally is not applicable to top level organizations)
team | Organizational team | An organizational team is usually a grouping of practitioners that perform a specific function within an organization (which could be a top level organization, or a department).
govt | Government | A political body, often used when including organization records for government bodies such as a Federal Government, State or Local Government.
ins | Insurance Company | A company that provides insurance to its subscribers that may include healthcare related policies.
pay | Payer | A company, charity, or governmental organization, which processes claims and/or issues payments to providers on behalf of patients or groups of patients.
edu | Educational Institute | An educational institution that provides education or research facilities.
reli | Religious Institution | An organization that is identified as a part of a religious institution.
crs | Clinical Research Sponsor | An organization that is identified as a Pharmaceutical/Clinical Research Sponsor.
cg | Community Group | An un-incorporated community group.
bus | Non-Healthcare Business or Corporation | An organization that is a registered business or corporation but not identified by other types.
other | Other | Other type of organization not already specified.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.499 Value Set http://hl7.org/fhir/ValueSet/defined-types

URL: https://hl7.org/fhir/R4/valueset-defined-types.html
FHIR Infrastructure Work Group | Maturity Level: N | Normative (from v4.0.0) | Use Context: Any

| This page has been approved as part of an ANSI standard. See the Conformance Package for further details.

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/defined-types
Version: | 4.0.1
Name: | FHIRDefinedType
Title: | FHIRDefinedType
Definition: | A list of all the concrete types defined in this version of the FHIR specification - Data Types and Resource Types.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.24 (for OID based terminology systems)
Source Resource | XML / JSON

This Value Set is normative - it is generated based on the information defined in this specification. The definition will remain fixed across versions, but the actual contents will change from version to version

This value set is used in the following places:

• Resource: StructureDefinition.type (uri / Extensible)
• Resource: TestScript.setup.action.operation.resource (code / Required)
• Resource: TestScript.setup.action.assert.resource (code / Required)
• Resource: ElementDefinition.type.code (uri / Extensible)

4.4.1.499.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/data-types
• Include all codes defined in http://hl7.org/fhir/resource-types

4.4.1.499.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 211 concepts

Expansion based on http://hl7.org/fhir/resource-types version 4.0.1, http://hl7.org/fhir/data-types version 4.0.1

Code | System | Display | Definition
Address | http://hl7.org/fhir/data-types | Address | An address expressed using postal conventions (as opposed to GPS or other location definition formats). This data type may be used to convey addresses for use in delivering mail as well as for visiting locations which might not be valid for mail delivery. There are a variety of postal address formats defined around the world.
Age | http://hl7.org/fhir/data-types | Age | A duration of time during which an organism (or a process) has existed.
Annotation | http://hl7.org/fhir/data-types | Annotation | A text note which also contains information about who made the statement and when.
Attachment | http://hl7.org/fhir/data-types | Attachment | For referring to data content defined in other formats.
BackboneElement | http://hl7.org/fhir/data-types | BackboneElement | Base definition for all elements that are defined inside a resource - but not those in a data type.
CodeableConcept | http://hl7.org/fhir/data-types | CodeableConcept | A concept that may be defined by a formal reference to a terminology or ontology or may be provided by text.
Coding | http://hl7.org/fhir/data-types | Coding | A reference to a code defined by a terminology system.
ContactDetail | http://hl7.org/fhir/data-types | ContactDetail | Specifies contact information for a person or organization.
ContactPoint | http://hl7.org/fhir/data-types | ContactPoint | Details for all kinds of technology mediated contact points for a person or organization, including telephone, email, etc.
Contributor | http://hl7.org/fhir/data-types | Contributor | A contributor to the content of a knowledge asset, including authors, editors, reviewers, and endorsers.
Count | http://hl7.org/fhir/data-types | Count | A measured amount (or an amount that can potentially be measured). Note that measured amounts include amounts that are not precisely quantified, including amounts involving arbitrary units and floating currencies.

DataRequirement | http://hl7.org/fhir/data-types | DataRequirement | Describes a required data item for evaluation in terms of the type of data, and optional code or date-based filters of the data.
Distance | http://hl7.org/fhir/data-types | Distance | A length - a value with a unit that is a physical distance.
Dosage | http://hl7.org/fhir/data-types | Dosage | Indicates how the medication is/was taken or should be taken by the patient.
Duration | http://hl7.org/fhir/data-types | Duration | A length of time.
Element | http://hl7.org/fhir/data-types | Element | Base definition for all elements in a resource.
ElementDefinition | http://hl7.org/fhir/data-types | ElementDefinition | Captures constraints on each element within the resource, profile, or extension.
Expression | http://hl7.org/fhir/data-types | Expression | A expression that is evaluated in a specified context and returns a value. The context of use of the expression must specify the context in which the expression is evaluated, and how the result of the expression is used.
Extension | http://hl7.org/fhir/data-types | Extension | Optional Extension Element - found in all resources.
HumanName | http://hl7.org/fhir/data-types | HumanName | A human's name with the ability to identify parts and usage.
Identifier | http://hl7.org/fhir/data-types | Identifier | An identifier - identifies some entity uniquely and unambiguously. Typically this is used for business identifiers.
MarketingStatus | http://hl7.org/fhir/data-types | MarketingStatus | The marketing status describes the date when a medicinal product is actually put on the market or the date as of which it is no longer available.
Meta | http://hl7.org/fhir/data-types | Meta | The metadata about a resource. This is content in the resource that is maintained by the infrastructure. Changes to the content might not always be associated with version changes to the resource.
Money | http://hl7.org/fhir/data-types | Money | An amount of economic utility in some recognized currency.
MoneyQuantity | http://hl7.org/fhir/data-types | MoneyQuantity
Narrative | http://hl7.org/fhir/data-types | Narrative | A human-readable summary of the resource conveying the essential clinical and business information for the resource.
ParameterDefinition | http://hl7.org/fhir/data-types | ParameterDefinition | The parameters to the module. This collection specifies both the input and output parameters. Input parameters are provided by the caller as part of the $evaluate operation. Output parameters are included in the GuidanceResponse.
Period | http://hl7.org/fhir/data-types | Period | A time period defined by a start and end date and optionally time.
Population | http://hl7.org/fhir/data-types | Population | A populatioof people with some set of grouping criteria.
ProdCharacteristic | http://hl7.org/fhir/data-types | ProdCharacteristic | The marketing status describes the date when a medicinal product is actually put on the market or the date as of which it is no longer available.
ProductShelfLife | http://hl7.org/fhir/data-types | ProductShelfLife | The shelf-life and storage information for a medicinal product item or container can be described using this class.
Quantity | http://hl7.org/fhir/data-types | Quantity | A measured amount (or an amount that can potentially be measured). Note that measured amounts include amounts that are not precisely quantified, including amounts involving arbitrary units and floating currencies.
Range | http://hl7.org/fhir/data-types | Range | A set of ordered Quantities defined by a low and high limit.
Ratio | http://hl7.org/fhir/data-types | Ratio | A relationship of two Quantity values - expressed as a numerator and a denominator.
Reference | http://hl7.org/fhir/data-types | Reference | A reference from one resource to another.
RelatedArtifact | http://hl7.org/fhir/data-types | RelatedArtifact | Related artifacts such as additional documentation, justification, or bibliographic references.
SampledData | http://hl7.org/fhir/data-types | SampledData | A series of measurements taken by a device, with upper and lower limits. There may be more than one dimension in the data.
Signature | http://hl7.org/fhir/data-types | Signature | A signature along with supporting context. The signature may be a digital signature that is cryptographic in nature, or some other signature acceptable to the domain. This other signature may be as simple as a graphical image representing a hand-written signature, or a signature ceremony Different signature approaches have different utilities.
SimpleQuantity | http://hl7.org/fhir/data-types | SimpleQuantity
SubstanceAmount | http://hl7.org/fhir/data-types | SubstanceAmount | Chemical substances are a single substance type whose primary defining element is the molecular structure. Chemical substances shall be defined on the basis of their complete covalent molecular structure; the presence of a salt (counter-ion) and/or solvates (water, alcohols) is also captured. Purity, grade, physical form or particle size are not taken into account in the definition of a chemical substance or in the assignment of a Substance ID.
Timing | http://hl7.org/fhir/data-types | Timing | Specifies an event that may occur multiple times. Timing schedules are used to record when things are planned, expected or requested to occur. The most common usage is in dosage instructions for medications. They are also used when planning care of various kinds, and may be used for reporting the schedule to which past regular activities were carried out.
TriggerDefinition | http://hl7.org/fhir/data-types | TriggerDefinition | A description of a triggering event. Triggering events can be named events, data events, or periodic, as determined by the type element.
UsageContext | http://hl7.org/fhir/data-types | UsageContext | Specifies clinical/business/etc. metadata that can be used to retrieve, index and/or categorize an artifact. This metadata can either be specific to the applicable population (e.g., age category, DRG) or the specific context of care (e.g., venue, care setting, provider of care).
base64Binary | http://hl7.org/fhir/data-types | base64Binary | A stream of bytes
boolean | http://hl7.org/fhir/data-types | boolean | Value of "true" or "false"
canonical | http://hl7.org/fhir/data-types | canonical | A URI that is a reference to a canonical URL on a FHIR resource
code | http://hl7.org/fhir/data-types | code | A string which has at least one character and no leading or trailing whitespace and where there is no whitespace other than single spaces in the contents
date | http://hl7.org/fhir/data-types | date | A date or partial date (e.g. just year or year + month). There is no time zone. The format is a union of the schema types gYear, gYearMonth and date. Dates SHALL be valid dates.
dateTime | http://hl7.org/fhir/data-types | dateTime | A date, date-time or partial date (e.g. just year or year + month). If hours and minutes are specified, a time zone SHALL be populated. The format is a union of the schema types gYear, gYearMonth, date and dateTime. Seconds must be provided due to schema type constraints but may be zero-filled and may be ignored. Dates SHALL be valid dates.
decimal | http://hl7.org/fhir/data-types | decimal | A rational number with implicit precision
id | http://hl7.org/fhir/data-types | id | Any combination of letters, numerals, "-" and ".", with a length limit of 64 characters. (This might be an integer, an unprefixed OID, UUID or any other identifier pattern that meets these constraints.) Ids are case-insensitive.

instant | http://hl7.org/fhir/data-types | instant | An instant in time - known at least to the second
integer | http://hl7.org/fhir/data-types | integer | A whole number
markdown | http://hl7.org/fhir/data-types | markdown | A string that may contain Github Flavored Markdown syntax for optional processing by a mark down presentation engine
oid | http://hl7.org/fhir/data-types | oid | An OID represented as a URI
positiveInt | http://hl7.org/fhir/data-types | positiveInt | An integer with a value that is positive (e.g. >0)
string | http://hl7.org/fhir/data-types | string | A sequence of Unicode characters
time | http://hl7.org/fhir/data-types | time | A time during the day, with no date specified
unsignedInt | http://hl7.org/fhir/data-types | unsignedInt | An integer with a value that is not negative (e.g. >= 0)
uri | http://hl7.org/fhir/data-types | uri | String of characters used to identify a name or a resource
url | http://hl7.org/fhir/data-types | url | A URI that is a literal reference
uuid | http://hl7.org/fhir/data-types | uuid | A UUID, represented as a URI
xhtml | http://hl7.org/fhir/data-types | XHTML | XHTML format, as defined by W3C, but restricted usage (mainly, no active content)
Account | http://hl7.org/fhir/resource-types | Account | A financial tool for tracking value accrued for a particular purpose. In the healthcare field, used to track charges for a patient, cost centers, etc.
ActivityDefinition | http://hl7.org/fhir/resource-types | ActivityDefinition | This resource allows for the definition of some activity to be performed, independent of a particular patient, practitioner, or other performance context.
AdverseEvent | http://hl7.org/fhir/resource-types | AdverseEvent | Actual or potential/avoided event causing unintended physical injury resulting from or contributed to by medical care, a research study or other healthcare setting factors that requires additional monitoring, treatment, or hospitalization, or that results in death.
AllergyIntolerance | http://hl7.org/fhir/resource-types | AllergyIntolerance | Risk of harmful or undesirable, physiological response which is unique to an individual and associated with exposure to a substance.
Appointment | http://hl7.org/fhir/resource-types | Appointment | A booking of a healthcare event among patient(s), practitioner(s), related person(s) and/or device(s) for a specific date/time. This may result in one or more Encounter(s).
AppointmentResponse | http://hl7.org/fhir/resource-types | AppointmentResponse | A reply to an appointment request for a patient and/or practitioner(s), such as a confirmation or rejection.
AuditEvent | http://hl7.org/fhir/resource-types | AuditEvent | A record of an event made for purposes of maintaining a security log. Typical uses include detection of intrusion attempts and monitoring for inappropriate usage.
Basic | http://hl7.org/fhir/resource-types | Basic | Basic is used for handling concepts not yet defined in FHIR, narrative-only resources that don't map to an existing resource, and custom resources not appropriate for inclusion in the FHIR specification.
Binary | http://hl7.org/fhir/resource-types | Binary | A resource that represents the data of a single raw artifact as digital content accessible in its native format. A Binary resource can contain any content, whether text, image, pdf, zip archive, etc.
BiologicallyDerivedProduct | http://hl7.org/fhir/resource-types | BiologicallyDerivedProduct | A material substance originating from a biological entity intended to be transplanted or infused into another (possibly the same) biological entity.
BodyStructure | http://hl7.org/fhir/resource-types | BodyStructure | Record details about an anatomical structure. This resource may be used when a coded concept does not provide the necessary detail needed for the use case.
Bundle | http://hl7.org/fhir/resource-types | Bundle | A container for a collection of resources.
CapabilityStatement | http://hl7.org/fhir/resource-types | CapabilityStatement | A Capability Statement documents a set of capabilities (behaviors) of a FHIR Server for a particular version of FHIR that may be used as a statement of actual server functionality or a statement of required or desired server implementation.
CarePlan | http://hl7.org/fhir/resource-types | CarePlan | Describes the intention of how one or more practitioners intend to deliver care for a particular patient, group or community for a period of time, possibly limited to care for a specific condition or set of conditions.
CareTeam | http://hl7.org/fhir/resource-types | CareTeam | The Care Team includes all the people and organizations who plan to participate in the coordination and delivery of care for a patient.
CatalogEntry | http://hl7.org/fhir/resource-types | CatalogEntry | Catalog entries are wrappers that contextualize items included in a catalog.
ChargeItem | http://hl7.org/fhir/resource-types | ChargeItem | The resource ChargeItem describes the provision of healthcare provider products for a certain patient, therefore referring not only to the product, but containing in addition details of the provision, like date, time, amounts and participating organizations and persons. Main Usage of the ChargeItem is to enable the billing process and internal cost allocation.
ChargeItemDefinition | http://hl7.org/fhir/resource-types | ChargeItemDefinition | The ChargeItemDefinition resource provides the properties that apply to the (billing) codes necessary to calculate costs and prices. The properties may differ largely depending on type and realm, therefore this resource gives only a rough structure and requires profiling for each type of billing code system.
Claim | http://hl7.org/fhir/resource-types | Claim | A provider issued list of professional services and products which have been provided, or are to be provided, to a patient which is sent to an insurer for reimbursement.
ClaimResponse | http://hl7.org/fhir/resource-types | ClaimResponse | This resource provides the adjudication details from the processing of a Claim resource.
ClinicalImpression | http://hl7.org/fhir/resource-types | ClinicalImpression | A record of a clinical assessment performed to determine what problem(s) may affect the patient and before planning the treatments or management strategies that are best to manage a patient's condition. Assessments are often 1:1 with a clinical consultation / encounter, but this varies greatly depending on the clinical workflow. This resource is called "ClinicalImpression" rather than "ClinicalAssessment" to avoid confusion with the recording of assessment tools such as Apgar score.
CodeSystem | http://hl7.org/fhir/resource-types | CodeSystem | The CodeSystem resource is used to declare the existence of and describe a code system or code system supplement and its key properties, and optionally define a part or all of its content.
Communication | http://hl7.org/fhir/resource-types | Communication | An occurrence of information being transmitted; e.g. an alert that was sent to a responsible provider, a public health agency that was notified about a reportable condition.
CommunicationRequest | http://hl7.org/fhir/resource-types | CommunicationRequest | A request to convey information; e.g. the CDS system proposes that an alert be sent to a responsible provider, the CDS system proposes that the public health agency be notified about a reportable condition.
CompartmentDefinition | http://hl7.org/fhir/resource-types | CompartmentDefinition | A compartment definition that defines how resources are accessed on a server.
Composition | http://hl7.org/fhir/resource-types | Composition | A set of healthcare-related information that is assembled together into a single logical package that provides a single coherent statement of meaning, establishes its own context and that has clinical attestation with regard to who is making the statement. A Composition defines the structure and narrative content necessary for a document. However, a Composition alone does not constitute a document. Rather, the Composition must be the

first entry in a Bundle where Bundle.type=document, and any other resources referenced from Composition must be included as subsequent entries in the Bundle (for example Patient, Practitioner, Encounter, etc.).

ConceptMap | http://hl7.org/fhir/resource-types | ConceptMap | A statement of relationships from one set of concepts to one or more other concepts - either concepts in code systems, or data element/data element concepts, or classes in class models.

Condition | http://hl7.org/fhir/resource-types | Condition | A clinical condition, problem, diagnosis, or other event, situation, issue, or clinical concept that has risen to a level of concern.

Consent | http://hl7.org/fhir/resource-types | Consent | A record of a healthcare consumer’s choices, which permits or denies identified recipient(s) or recipient role(s) to perform one or more actions within a given policy context, for specific purposes and periods of time.

Contract | http://hl7.org/fhir/resource-types | Contract | Legally enforceable, formally recorded unilateral or bilateral directive i.e., a policy or agreement.

Coverage | http://hl7.org/fhir/resource-types | Coverage | Financial instrument which may be used to reimburse or pay for health care products and services. Includes both insurance and self-payment.

CoverageEligibilityRequest | http://hl7.org/fhir/resource-types | CoverageEligibilityRequest | The CoverageEligibilityRequest provides patient and insurance coverage information to an insurer for them to respond, in the form of an CoverageEligibilityResponse, with information regarding whether the stated coverage is valid and in-force and optionally to provide the insurance details of the policy.

CoverageEligibilityResponse | http://hl7.org/fhir/resource-types | CoverageEligibilityResponse | This resource provides eligibility and plan details from the processing of an CoverageEligibilityRequest resource.

DetectedIssue | http://hl7.org/fhir/resource-types | DetectedIssue | Indicates an actual or potential clinical issue with or between one or more active or proposed clinical actions for a patient; e.g. Drug-drug interaction, Ineffective treatment frequency, Procedure-condition conflict, etc.

Device | http://hl7.org/fhir/resource-types | Device | A type of a manufactured item that is used in the provision of healthcare without being substantially changed through that activity. The device may be a medical or non-medical device.

DeviceDefinition | http://hl7.org/fhir/resource-types | DeviceDefinition | The characteristics, operational status and capabilities of a medical-related component of a medical device.

DeviceMetric | http://hl7.org/fhir/resource-types | DeviceMetric | Describes a measurement, calculation or setting capability of a medical device.

DeviceRequest | http://hl7.org/fhir/resource-types | DeviceRequest | Represents a request for a patient to employ a medical device. The device may be an implantable device, or an external assistive device, such as a walker.

DeviceUseStatement | http://hl7.org/fhir/resource-types | DeviceUseStatement | A record of a device being used by a patient where the record is the result of a report from the patient or another clinician.

DiagnosticReport | http://hl7.org/fhir/resource-types | DiagnosticReport | The findings and interpretation of diagnostic tests performed on patients, groups of patients, devices, and locations, and/or specimens derived from these. The report includes clinical context such as requesting and provider information, and some mix of atomic results, images, textual and coded interpretations, and formatted representation of diagnostic reports.

DocumentManifest | http://hl7.org/fhir/resource-types | DocumentManifest | A collection of documents compiled for a purpose together with metadata that applies to the collection.

DocumentReference | http://hl7.org/fhir/resource-types | DocumentReference | A reference to a document of any kind for any purpose. Provides metadata about the document so that the document can be discovered and managed. The scope of a document is any seralized object with a mime-type, so includes formal patient centric documents (CDA), cliical notes, scanned paper, and non-patient specific documents like policy text.

DomainResource | http://hl7.org/fhir/resource-types | DomainResource | A resource that includes narrative, extensions, and contained resources.

EffectEvidenceSynthesis | http://hl7.org/fhir/resource-types | EffectEvidenceSynthesis | The EffectEvidenceSynthesis resource describes the difference in an outcome between exposures states in a population where the effect estimate is derived from a combination of research studies.

Encounter | http://hl7.org/fhir/resource-types | Encounter | An interaction between a patient and healthcare provider(s) for the purpose of providing healthcare service(s) or assessing the health status of a patient.

Endpoint | http://hl7.org/fhir/resource-types | Endpoint | The technical details of an endpoint that can be used for electronic services, such as for web services providing XDS.b or a REST endpoint for another FHIR server. This may include any security context information.

EnrollmentRequest | http://hl7.org/fhir/resource-types | EnrollmentRequest | This resource provides the insurance enrollment details to the insurer regarding a specified coverage.

EnrollmentResponse | http://hl7.org/fhir/resource-types | EnrollmentResponse | This resource provides enrollment and plan details from the processing of an EnrollmentRequest resource.

EpisodeOfCare | http://hl7.org/fhir/resource-types | EpisodeOfCare | An association between a patient and an organization / healthcare provider(s) during which time encounters may occur. The managing organization assumes a level of responsibility for the patient during this time.

EventDefinition | http://hl7.org/fhir/resource-types | EventDefinition | The EventDefinition resource provides a reusable description of when a particular event can occur.

Evidence | http://hl7.org/fhir/resource-types | Evidence | The Evidence resource describes the conditional state (population and any exposures being compared within the population) and outcome (if specified) that the knowledge (evidence, assertion, recommendation) is about.

EvidenceVariable | http://hl7.org/fhir/resource-types | EvidenceVariable | The EvidenceVariable resource describes a "PICO" element that knowledge (evidence, assertion, recommendation) is about.

ExampleScenario | http://hl7.org/fhir/resource-types | ExampleScenario | Example of workflow instance.

ExplanationOfBenefit | http://hl7.org/fhir/resource-types | ExplanationOfBenefit | This resource provides: the claim details; adjudication details from the processing of a Claim; and optionally account balance information, for informing the subscriber of the benefits provided.

FamilyMemberHistory | http://hl7.org/fhir/resource-types | FamilyMemberHistory | Significant health conditions for a person related to the patient relevant in the context of care for the patient.

Flag | http://hl7.org/fhir/resource-types | Flag | Prospective warnings of potential issues when providing care to the patient.

Goal | http://hl7.org/fhir/resource-types | Goal | Describes the intended objective(s) for a patient, group or organization care, for example, weight loss, restoring an activity of daily living, obtaining herd immunity via immunization, meeting a process improvement objective, etc.

GraphDefinition | http://hl7.org/fhir/resource-types | GraphDefinition | A formal computable definition of a graph of resources - that is, a coherent set of resources that form a graph by following references. The Graph Definition resource defines a set and makes rules about the set.

Group | http://hl7.org/fhir/resource-types | Group | Represents a defined collection of entities that may be discussed or acted upon collectively but which are not expected to act collectively, and are not formally or legally recognized; i.e. a collection of entities that isn't an Organization.

GuidanceResponse | http://hl7.org/fhir/resource-types | GuidanceResponse | A guidance response is the formal response to a guidance request, including any output parameters returned by the evaluation, as well as the description of any proposed actions to be taken.

HealthcareService | http://hl7.org/fhir/resource-types | HealthcareService | The details of a healthcare service available at a location.

ImagingStudy | http://hl7.org/fhir/resource-types | ImagingStudy | Representation of the content produced in a DICOM imaging study. A study comprises a set of series, each of which includes a set of Service-Object Pair Instances (SOP Instances - images or other data) acquired or produced in a common context. A series is of only one modality (e.g. X-ray, CT, MR, ultrasound), but a study may have multiple series of different modalities.

Immunization | http://hl7.org/fhir/resource-types | Immunization | Describes the event of a patient being administered a vaccine or a record of an immunization as reported by a patient, a clinician or another party.

ImmunizationEvaluation | http://hl7.org/fhir/resource-types | ImmunizationEvaluation | Describes a comparison of an immunization event against published recommendations to determine if the administration is "valid" in relation to those recommendations.

ImmunizationRecommendation | http://hl7.org/fhir/resource-types | ImmunizationRecommendation | A patient's point-in-time set of recommendations (i.e. forecasting) according to a published schedule with optional supporting justification.

ImplementationGuide | http://hl7.org/fhir/resource-types | ImplementationGuide | A set of rules of how a particular interoperability or standards problem is solved - typically through the use of FHIR resources. This resource is used to gather all the parts of an implementation guide into a logical whole and to publish a computable definition of all the parts.

InsurancePlan | http://hl7.org/fhir/resource-types | InsurancePlan | Details of a Health Insurance product/plan provided by an organization.

Invoice | http://hl7.org/fhir/resource-types | Invoice | Invoice containing collected ChargeItems from an Account with calculated individual and total price for Billing purpose.

Library | http://hl7.org/fhir/resource-types | Library | The Library resource is a general-purpose container for knowledge asset definitions. It can be used to describe and expose existing knowledge assets such as logic libraries and information model descriptions, as well as to describe a collection of knowledge assets.

Linkage | http://hl7.org/fhir/resource-types | Linkage | Identifies two or more records (resource instances) that refer to the same real-world "occurrence".

List | http://hl7.org/fhir/resource-types | List | A list is a curated collection of resources.

Location | http://hl7.org/fhir/resource-types | Location | Details and position information for a physical place where services are provided and resources and participants may be stored, found, contained, or accommodated.

Measure | http://hl7.org/fhir/resource-types | Measure | The Measure resource provides the definition of a quality measure.

MeasureReport | http://hl7.org/fhir/resource-types | MeasureReport | The MeasureReport resource contains the results of the calculation of a measure; and optionally a reference to the resources involved in that calculation.

Media | http://hl7.org/fhir/resource-types | Media | A photo, video, or audio recording acquired or used in healthcare. The actual content may be inline or provided by direct reference.

Medication | http://hl7.org/fhir/resource-types | Medication | This resource is primarily used for the identification and definition of a medication for the purposes of prescribing, dispensing, and administering a medication as well as for making statements about medication use.

MedicationAdministration | http://hl7.org/fhir/resource-types | MedicationAdministration | Describes the event of a patient consuming or otherwise being administered a medication. This may be as simple as swallowing a tablet or it may be a long running infusion. Related resources tie this event to the authorizing prescription, and the specific encounter between patient and health care practitioner.

MedicationDispense | http://hl7.org/fhir/resource-types | MedicationDispense | Indicates that a medication product is to be or has been dispensed for a named person/patient. This includes a description of the medication product (supply) provided and the instructions for administering the medication. The medication dispense is the result of a pharmacy system responding to a medication order.

MedicationKnowledge | http://hl7.org/fhir/resource-types | MedicationKnowledge | Information about a medication that is used to support knowledge.

MedicationRequest | http://hl7.org/fhir/resource-types | MedicationRequest | An order or request for both supply of the medication and the instructions for administration of the medication to a patient. The resource is called "MedicationRequest" rather than "MedicationPrescription" or "MedicationOrder" to generalize the use across inpatient and outpatient settings, including care plans, etc., and to harmonize with workflow patterns.

MedicationStatement | http://hl7.org/fhir/resource-types | MedicationStatement | A record of a medication that is being consumed by a patient. A MedicationStatement may indicate that the patient may be taking the medication now or has taken the medication in the past or will be taking the medication in the future. The source of this information can be the patient, significant other (such as a family member or spouse), or a clinician. A common scenario where this information is captured is during the history taking process during a patient visit or stay. The medication information may come from sources such as the patient's memory, from a prescription bottle, or from a list of medications the patient, clinician or other party maintains. The primary difference between a medication statement and a medication administration is that the medication administration has complete administration information and is based on actual administration information from the person who administered the medication. A medication statement is often, if not always, less specific. There is no required date/time when the medication was administered, in fact we only know that a source has reported the patient is taking this medication, where details such as time, quantity, or rate or even medication product may be incomplete or missing or less precise. As stated earlier, the medication statement information may come from the patient's memory, from a prescription bottle or from a list of medications the patient, clinician or other party maintains. Medication administration is more formal and is not missing detailed information.

MedicinalProduct | http://hl7.org/fhir/resource-types | MedicinalProduct | Detailed definition of a medicinal product, typically for uses other than direct patient care (e.g. regulatory use).

MedicinalProductAuthorization | http://hl7.org/fhir/resource-types | MedicinalProductAuthorization | The regulatory authorization of a medicinal product.

MedicinalProductContraindication | http://hl7.org/fhir/resource-types | MedicinalProductContraindication | The clinical particulars - indications, contraindications etc. of a medicinal product, including for regulatory purposes.

MedicinalProductIndication | http://hl7.org/fhir/resource-types | MedicinalProductIndication | Indication for the Medicinal Product.
MedicinalProductIngredient | http://hl7.org/fhir/resource-types | MedicinalProductIngredient | An ingredient of a manufactured item or pharmaceutical product.
MedicinalProductInteraction | http://hl7.org/fhir/resource-types | MedicinalProductInteraction | The interactions of the medicinal product with other medicinal products, or other forms of interactions.
MedicinalProductManufactured | http://hl7.org/fhir/resource-types | MedicinalProductManufactured | The manufactured item as contained in the packaged medicinal product.
MedicinalProductPackaged | http://hl7.org/fhir/resource-types | MedicinalProductPackaged | A medicinal product in a container or package.
MedicinalProductPharmaceutical | http://hl7.org/fhir/resource-types | MedicinalProductPharmaceutical | A pharmaceutical product described in terms of its composition and dose form.
MedicinalProductUndesirableEffect | http://hl7.org/fhir/resource-types | MedicinalProductUndesirableEffect | Describe the undesirable effects of the medicinal product.
MessageDefinition | http://hl7.org/fhir/resource-types | MessageDefinition | Defines the characteristics of a message that can be shared between systems, including the type of event that initiates the message, the content to be transmitted and what response(s), if any, are permitted.
MessageHeader | http://hl7.org/fhir/resource-types | MessageHeader | The header for a message exchange that is either requesting or responding to an action. The reference(s) that are the subject of the action as well as other information related to the action are typically transmitted in a bundle in which the MessageHeader resource instance is the first resource in the bundle.
MolecularSequence | http://hl7.org/fhir/resource-types | MolecularSequence | Raw data describing a biological sequence.
NamingSystem | http://hl7.org/fhir/resource-types | NamingSystem | A curated namespace that issues unique symbols within that namespace for the identification of concepts, people, devices, etc. Represents a "System" used within the Identifier and Coding data types.
NutritionOrder | http://hl7.org/fhir/resource-types | NutritionOrder | A request to supply a diet, formula feeding (enteral) or oral nutritional supplement to a patient/resident.
Observation | http://hl7.org/fhir/resource-types | Observation | Measurements and simple assertions made about a patient, device or other subject.
ObservationDefinition | http://hl7.org/fhir/resource-types | ObservationDefinition | Set of definitional characteristics for a kind of observation or measurement produced or consumed by an orderable health care service.
OperationDefinition | http://hl7.org/fhir/resource-types | OperationDefinition | A formal computable definition of an operation (on the RESTful interface) or a named query (using the search interaction).
OperationOutcome | http://hl7.org/fhir/resource-types | OperationOutcome | A collection of error, warning, or information messages that result from a system action.
Organization | http://hl7.org/fhir/resource-types | Organization | A formally or informally recognized grouping of people or organizations formed for the purpose of achieving some form of collective action. Includes companies, institutions, corporations, departments, community groups, healthcare practice groups, payer/insurer, etc.
OrganizationAffiliation | http://hl7.org/fhir/resource-types | OrganizationAffiliation | Defines an affiliation/assotiation/relationship between 2 distinct oganizations, that is not a part-of relationship/sub-division relationship.
Parameters | http://hl7.org/fhir/resource-types | Parameters | This resource is a non-persisted resource used to pass information into and back from an [operation](operations.html). It has no other use, and there is no RESTful endpoint associated with it.
Patient | http://hl7.org/fhir/resource-types | Patient | Demographics and other administrative information about an individual or animal receiving care or other health-related services.
PaymentNotice | http://hl7.org/fhir/resource-types | PaymentNotice | This resource provides the status of the payment for goods and services rendered, and the request and response resource references.
PaymentReconciliation | http://hl7.org/fhir/resource-types | PaymentReconciliation | This resource provides the details including amount of a payment and allocates the payment items being paid.
Person | http://hl7.org/fhir/resource-types | Person | Demographics and administrative information about a person independent of a specific health-related context.
PlanDefinition | http://hl7.org/fhir/resource-types | PlanDefinition | This resource allows for the definition of various types of plans as a sharable, consumable, and executable artifact. The resource is general enough to support the description of a broad range of clinical artifacts such as clinical decision support rules, order sets and protocols.
Practitioner | http://hl7.org/fhir/resource-types | Practitioner | A person who is directly or indirectly involved in the provisioning of healthcare.
PractitionerRole | http://hl7.org/fhir/resource-types | PractitionerRole | A specific set of Roles/Locations/specialties/services that a practitioner may perform at an organization for a period of time.
Procedure | http://hl7.org/fhir/resource-types | Procedure | An action that is or was performed on or for a patient. This can be a physical intervention like an operation, or less invasive like long term services, counseling, or hypnotherapy.
Provenance | http://hl7.org/fhir/resource-types | Provenance | Provenance of a resource is a record that describes entities and processes involved in producing and delivering or otherwise influencing that resource. Provenance provides a critical foundation for assessing authenticity, enabling trust, and allowing reproducibility. Provenance assertions are a form of contextual metadata and can themselves become important records with their own provenance. Provenance statement indicates clinical significance in terms of confidence in authenticity, reliability, and trustworthiness, integrity, and stage in lifecycle (e.g. Document Completion - has the artifact been legally authenticated), all of which may impact security, privacy, and trust policies.
Questionnaire | http://hl7.org/fhir/resource-types | Questionnaire | A structured set of questions intended to guide the collection of answers from end-users. Questionnaires provide detailed control over order, presentation, phraseology and grouping to allow coherent, consistent data collection.
QuestionnaireResponse | http://hl7.org/fhir/resource-types | QuestionnaireResponse | A structured set of questions and their answers. The questions are ordered and grouped into coherent subsets, corresponding to the structure of the grouping of the questionnaire being responded to.
RelatedPerson | http://hl7.org/fhir/resource-types | RelatedPerson | Information about a person that is involved in the care for a patient, but who is not the target of healthcare, nor has a formal responsibility in the care process.
RequestGroup | http://hl7.org/fhir/resource-types | RequestGroup | A group of related requests that can be used to capture intended activities that have inter-dependencies such as "give this medication after that one".
ResearchDefinition | http://hl7.org/fhir/resource-types | ResearchDefinition | The ResearchDefinition resource describes the conditional state (population and any exposures being compared within the population) and outcome (if specified) that the knowledge (evidence, assertion, recommendation) is about.

ResearchElementDefinition | http://hl7.org/fhir/resource-types | ResearchElementDefinition | The ResearchElementDefinition resource describes a "PICO" element that knowledge (evidence, assertion, recommendation) is about.

ResearchStudy | http://hl7.org/fhir/resource-types | ResearchStudy | A process where a researcher or organization plans and then executes a series of steps intended to increase the field of healthcare-related knowledge. This includes studies of safety, efficacy, comparative effectiveness and other information about medications, devices, therapies and other interventional and investigative techniques. A ResearchStudy involves the gathering of information about human or animal subjects.

ResearchSubject | http://hl7.org/fhir/resource-types | ResearchSubject | A physical entity which is the primary unit of operational and/or administrative interest in a study.

Resource | http://hl7.org/fhir/resource-types | Resource | This is the base resource type for everything.

RiskAssessment | http://hl7.org/fhir/resource-types | RiskAssessment | An assessment of the likely outcome(s) for a patient or other subject as well as the likelihood of each outcome.

RiskEvidenceSynthesis | http://hl7.org/fhir/resource-types | RiskEvidenceSynthesis | The RiskEvidenceSynthesis resource describes the likelihood of an outcome in a population plus exposure state where the risk estimate is derived from a combination of research studies.

Schedule | http://hl7.org/fhir/resource-types | Schedule | A container for slots of time that may be available for booking appointments.

SearchParameter | http://hl7.org/fhir/resource-types | SearchParameter | A search parameter that defines a named search item that can be used to search/filter on a resource.

ServiceRequest | http://hl7.org/fhir/resource-types | ServiceRequest | A record of a request for service such as diagnostic investigations, treatments, or operations to be performed.

Slot | http://hl7.org/fhir/resource-types | Slot | A slot of time on a schedule that may be available for booking appointments.

Specimen | http://hl7.org/fhir/resource-types | Specimen | A sample to be used for analysis.

SpecimenDefinition | http://hl7.org/fhir/resource-types | SpecimenDefinition | A kind of specimen with associated set of requirements.

StructureDefinition | http://hl7.org/fhir/resource-types | StructureDefinition | A definition of a FHIR structure. This resource is used to describe the underlying resources, data types defined in FHIR, and also for describing extensions and constraints on resources and data types.

StructureMap | http://hl7.org/fhir/resource-types | StructureMap | A Map of relationships between 2 structures that can be used to transform data.

Subscription | http://hl7.org/fhir/resource-types | Subscription | The subscription resource is used to define a push-based subscription from a server to another system. Once a subscription is registered with the server, the server checks every resource that is created or updated, and if the resource matches the given criteria, it sends a message on the defined "channel" so that another system can take an appropriate action.

Substance | http://hl7.org/fhir/resource-types | Substance | A homogeneous material with a definite composition.

SubstanceNucleicAcid | http://hl7.org/fhir/resource-types | SubstanceNucleicAcid | Nucleic acids are defined by three distinct elements: the base, sugar and linkage. Individual substance/moiety IDs will be created for each of these elements. The nucleotide sequence will be always entered in the 5’-3’ direction.

SubstancePolymer | http://hl7.org/fhir/resource-types | SubstancePolymer | Todo.

SubstanceProtein | http://hl7.org/fhir/resource-types | SubstanceProtein | A SubstanceProtein is defined as a single unit of a linear amino acid sequence, or a combination of subunits that are either covalently linked or have a defined invariant stoichiometric relationship. This includes all synthetic, recombinant and purified SubstanceProteins of defined sequence, whether the use is therapeutic or prophylactic. This set of elements will be used to describe albumins, coagulation factors, cytokines, growth factors, peptide/SubstanceProtein hormones, enzymes, toxins, toxoids, recombinant vaccines, and immunomodulators.

SubstanceReferenceInformation | http://hl7.org/fhir/resource-types | SubstanceReferenceInformation | Todo.

SubstanceSourceMaterial | http://hl7.org/fhir/resource-types | SubstanceSourceMaterial | Source material shall capture information on the taxonomic and anatomical origins as well as the fraction of a material that can result in or can be modified to form a substance. This set of data elements shall be used to define polymer substances isolated from biological matrices. Taxonomic and anatomical origins shall be described using a controlled vocabulary as required. This information is captured for naturally derived polymers (. starch) and structurally diverse substances. For Organisms belonging to the Kingdom Plantae the Substance level defines the fresh material of a single species or infraspecies, the Herbal Drug and the Herbal preparation. For Herbal preparations, the fraction information will be captured at the Substance information level and additional information for herbal extracts will be captured at the Specified Substance Group 1 information level. See for further explanation the Substance Class: Structurally Diverse and the herbal annex.

SubstanceSpecification | http://hl7.org/fhir/resource-types | SubstanceSpecification | The detailed description of a substance, typically at a level beyond what is used for prescribing.

SupplyDelivery | http://hl7.org/fhir/resource-types | SupplyDelivery | Record of delivery of what is supplied.

SupplyRequest | http://hl7.org/fhir/resource-types | SupplyRequest | A record of a request for a medication, substance or device used in the healthcare setting.

Task | http://hl7.org/fhir/resource-types | Task | A task to be performed.

TerminologyCapabilities | http://hl7.org/fhir/resource-types | TerminologyCapabilities | A TerminologyCapabilities resource documents a set of capabilities (behaviors) of a FHIR Terminology Server that may be used as a statement of actual server functionality or a statement of required or desired server implementation.

TestReport | http://hl7.org/fhir/resource-types | TestReport | A summary of information based on the results of executing a TestScript.

TestScript | http://hl7.org/fhir/resource-types | TestScript | A structured set of tests against a FHIR server or client implementation to determine compliance against the FHIR specification.

ValueSet | http://hl7.org/fhir/resource-types | ValueSet | A ValueSet resource instance specifies a set of codes drawn from one or more code systems, intended for use in a particular context. Value sets link between [[[CodeSystem]]] definitions and their use in [coded elements](terminologies.html).

VerificationResult | http://hl7.org/fhir/resource-types | VerificationResult | Describes validation requirements, source(s), status and dates for one or more elements.

VisionPrescription | http://hl7.org/fhir/resource-types | VisionPrescription | An authorization for the provision of glasses and/or contact lenses to a patient.

Additional Language Displays

Code | English (English, en) | Spanish (es) | French (fr) | Italian (it) | □□ (Chinese, zh)

Address
Age
Annotation
Attachment
BackboneElement
CodeableConcept
Coding
ContactDetail
ContactPoint
Contributor
Count
DataRequirement
Distance
Dosage
Duration
Element
ElementDefinition
Expression
Extension
HumanName
Identifier
MarketingStatus
Meta
Money
MoneyQuantity
Narrative
ParameterDefinition
Period
Population
ProdCharacteristic
ProductShelfLife
Quantity
Range
Ratio
Reference
RelatedArtifact
SampledData
Signature
SimpleQuantity
SubstanceAmount
Timing
TriggerDefinition
UsageContext
base64Binary
boolean
canonical
code
date
dateTime
decimal
id
instant
integer
markdown
oid
positiveInt
string
time
unsignedInt
uri
url
uuid
xhtml
Account | Account | Cuenta | | | □□
ActivityDefinition | ActivityDefinition | DefinicionDeActividad | | DefinizioneAttivita | □□□□
AdverseEvent | AdverseEvent | EventoAdverso | | EventoAvverso | □□□□
AllergyIntolerance | AllergyIntolerance | AllergiaIntolerancia | IntoléranceAllergique | AllergiaIntolleranza | □□□□□□□□
Appointment | Appointment | Cita | RendezVous | Appuntamento | □□
AppointmentResponse | AppointmentResponse | CitaRespuesta | RéponseRendezVous | RispostaAppuntamento | □□□□
AuditEvent | AuditEvent | EventoSeguridad | ÉvènementSécurité | | □□□□
Basic | Basic | Basico | Basique | | □□□□
Binary | Binary | Binario | Binaire | Binario | □□□□□

BiologicallyDerivedProduct | BiologicallyDerivedProduct
BodyStructure | BodyStructure | EstructuraDelCuerpo | MorphologieDeCorps | StrutturaDelCorpo | □□□□
Bundle | Bundle | Paquete | Paquet | | □□
CapabilityStatement | CapabilityStatement | DeclaracionDeCapacidad | | | □□□□
CarePlan | CarePlan | PlanDeCuidado | PlanDeSoins | PianoDiCura | □□□□
CareTeam | CareTeam | EquipoDeCuidado | | | □□□□
CatalogEntry | CatalogEntry | EntradaDeCatalogo | | | □□□□
ChargeItem | ChargeItem | CargoDeItem | | | □□□□
ChargeItemDefinition | ChargeItemDefinition
Claim | Claim | Reclamación / Factura | Réclamation | | □□
ClaimResponse | ClaimResponse | | RéponseARéclamation | | □□□□
ClinicalImpression | ClinicalImpression | HallazgoClinico | ImpressionClinique | ImpressioneClinica | □□□□
CodeSystem | CodeSystem | SistemaDeCodigos | | SistemaDiCodifica | □□□□
Communication | Communication | Comunicación | Communication | Comunicazione | □□
CommunicationRequest | CommunicationRequest | ComunicaciónRequerimiento | DemandeDeCommunication | RichiestaDiComunicazione | □□□□
CompartmentDefinition | CompartmentDefinition | DefinicionDeCompartimento | | | □□□□□□
Composition | Composition | Composición | Composition | Composizione | □□□□□
ConceptMap | ConceptMap | MapaDeConceptos | CarteDeConcepts | MappaDiConcetti | □□□□
Condition | Condition | Condición | Condition | Condizione | □□
Consent | Consent | Consentimiento | | Consenso | □□□
Contract | Contract | Contato | Contrat | Contratto | □□
Coverage | Coverage | Cobertura | Couverture | Copertura | □□□□
CoverageEligibilityRequest | CoverageEligibilityRequest | | CouvertureDemandeEligibilité | RichiestaEleggibilitaCopertura | □□□□
CoverageEligibilityResponse | CoverageEligibilityResponse | | RéponseEligibilitéCouverture | RispostaEleggibilitaCopertura | □□□□
DetectedIssue | DetectedIssue | Problema-Detectado /ProblemaDetectado | Problème Détecté | ProblemaRilevato | □□□□□
Device | Device | Dispositivo | Dispositif | Dispositivo | □□
DeviceDefinition | DeviceDefinition | DefiniciónDeDispositivo | DéfinitionDeDispositif | | □□□□
DeviceMetric | DeviceMetric | MetricaDeDispositivo | MétriqueDispositif | | □□□□
DeviceRequest | DeviceRequest | SolicitudDeDispositivo | DemandeUtilisationDispositif | RichiestaDispositivo | □□□□
DeviceUseStatement | DeviceUseStatement | | | | □□□□□□
DiagnosticReport | DiagnosticReport | InformeDiagnostico | RapportDiagnostique | RefertoDiagnostico | □□□□
DocumentManifest | DocumentManifest | ManifestoDocumento | Manifeste | | □□□□
DocumentReference | DocumentReference | ReferenciaDocumento | RéférenceDocument | RiferimentoDocumento | □□□□
DomainResource | DomainResource | RecursoDeDominio | | | □□□□
EffectEvidenceSynthesis | EffectEvidenceSynthesis
Encounter | Encounter | Encuentro | Venue | | □□□□
Endpoint | Endpoint | Endpoint | | | □□
EnrollmentRequest | EnrollmentRequest | SolicitudDeEnrolamiento | DemandeInscription | RichiestaIscrizione | □□□□□□
EnrollmentResponse | EnrollmentResponse | RespuestaDeEnrolamiento | RéponseInscription | RispostaIscrizione | □□□□□□
EpisodeOfCare | EpisodeOfCare | EpisodioDeCuidado | ÉpisodeDeSoins | EpisodioDiCura | □□□□□□
EventDefinition | EventDefinition | DefinitionDeEvento | | DefinizioneEvento | □□□□
Evidence | Evidence
EvidenceVariable | EvidenceVariable
ExampleScenario | ExampleScenario | EjemploDeEscenario | | ScenarioDiEsempio | □□□□
ExplanationOfBenefit | ExplanationOfBenefit | | ExplicationDuBénéfice | | □□□□
FamilyMemberHistory | FamilyMemberHistory | HistorialMiembroFamiliar | HistoireMembreFamille | | □□□
Flag | Flag | Bandera | Drapeau | | □□
Goal | Goal | Objetivo | But | | □□
GraphDefinition | GraphDefinition | DefinitionGrafico | | | □□□□
Group | Group | Grupo | Groupe | Gruppo | □□
GuidanceResponse | GuidanceResponse | RespuestaDeOrientacion | | | □□□□□□
HealthcareService | HealthcareService | ServicioDeCuidado | ServiceDeSanté | ServizioSanitario | □□□□□□□□□
ImagingStudy | ImagingStudy | EstudioImagen / EstudioImagen | EtudeImagerie | | □□□□
Immunization | Immunization | Inmunización | Immunisation | Immunizzazione | □□□□
ImmunizationEvaluation | ImmunizationEvaluation
ImmunizationRecommendation | ImmunizationRecommendation | RecomendaciónInmunización / | RecommendationImmunisation | | □□□□□□
ImplementationGuide | ImplementationGuide | GuiaDeImplementacion | | | □□□□
InsurancePlan | InsurancePlan
Invoice | Invoice
Library | Library | Librería | | | □
Linkage | Linkage | Enlace / Conexión / Vinculo / Acoplamiento | | | □□□□
List | List | Lista | Liste | Lista | □□
Location | Location | Locacion | Localisation | | □□
Measure | Measure | Medida | | Misura | □□
MeasureReport | MeasureReport | ReporteMedida | | | □□□□
Media | Media | Medio | Média | Media | □□
Medication | Medication | Medicación /Medicamento | Médication | | □□
MedicationAdministration | MedicationAdministration | AdministraciónMedicación / AdministracionMedicamento | AdministrationMédicaments | | □□□□
MedicationDispense | MedicationDispense | DispensaciónMedicación /DispensacionMedicamento | DispensationMédicaments | | □□□□

MedicationKnowledge | MedicationKnowledge
MedicationRequest | MedicationRequest | PrescripciónMedicaciónTODO /PrescripcionMedicamento | PrescriptionMédicamenteuseTODO | | □□□□
MedicationStatement | MedicationStatement | ResumenMedicación /ResumenMedicamento | ÉtatMédication | | □□□□
MedicinalProduct | MedicinalProduct | | | ProdottoMedicinale
MedicinalProductAuthorization | MedicinalProductAuthorization | | | AutorizzazioneProdottoMedicinale
MedicinalProductContraindication | MedicinalProductContraindication
MedicinalProductIndication | MedicinalProductInteraction
MedicinalProductIngredient | MedicinalProductIngredient | | | IngredienteProdottoMedicinale
MedicinalProductInteraction | MedicinalProductInteraction
MedicinalProductManufactured | MedicinalProductManufactured
MedicinalProductPackaged | MedicinalProductPackaged
MedicinalProductPharmaceutical | MedicinalProductPharmaceutical
MedicinalProductUndesirableEffect | MedicinalProductUndesirableEffect
MessageDefinition | MessageDefinition | DefinicionMensaje | | DefinizioneMessaggio | □□□□
MessageHeader | MessageHeader | CabeceraMensaje | EntêteMessage | | □□□□
MolecularSequence | MolecularSequence | SecuenciaMolecular | | | □□□□
NamingSystem | NamingSystem | SistemaDeNombres | SystèmeDeNommage | | □□□□
NutritionOrder | NutritionOrder | OrdenNutrición | OrdreNutrition | | □□□□
Observation | Observation | Observación | Observation | Osservazione | □□
ObservationDefinition | ObservationDefinition | DefinicionDeEspecimen | | DefinizioneOsservazione | □□□□
OperationDefinition | OperationDefinition | DefinicionDeOperacion | DéfinitionOpération | DefinizioneOperazione | □□□□
OperationOutcome | OperationOutcome | ResultadoOperación | RésultatOpération | RisultatoOperazione | □□□□
Organization | Organization | Organización | Organisation | Organizzazione | □□□□
OrganizationAffiliation | OrganizationAffiliation
Parameters | Parameters | Parametros | | | □□
Patient | Patient | Paciente | Patient | Paziente | □□
PaymentNotice | PaymentNotice | AvisoDePago | AvisPaiement | AvvisoDiPagamento | □□□□
PaymentReconciliation | PaymentReconciliation | ConciliacionDePago | RéconciliationPaiement | RiconciliazionePagamento | □□□□
Person | Person | Persona | Personne | Persona | □□
PlanDefinition | PlanDefinition | DefinicionDePlan | | | □□□□
Practitioner | Practitioner | Practicante / Profesional | Praticien | | □□□□
PractitionerRole | PractitionerRole | RolProfesional | | | □□□□□□
Procedure | Procedure | Procedimiento | Procédure | Procedura | □□□□
Provenance | Provenance | Procedencia | Provenance | Provenienza | □□
Questionnaire | Questionnaire | Cuestionario | Questionnaire | Questionario | □□□□
QuestionnaireResponse | QuestionnaireResponse | RespuestaAlCuestionario | RéponseQuestionnaire | RispostaQuestionario | □□□□□□
RelatedPerson | RelatedPerson | PersonaRelacionada | PersonneEnRelation | PersonaCorrelata | □□□□
RequestGroup | RequestGroup | GrupoDeSolicitudes | | GruppoDiRichieste | □□□□
ResearchDefinition | ResearchDefinition
ResearchElementDefinition | ResearchElementDefinition
ResearchStudy | ResearchStudy | EstudioDeInvestigacion | | | □□□□
ResearchSubject | ResearchSubject | SujetoDeInvestigacion | | | □□□□
Resource | Resource | Recurso | | | □□
RiskAssessment | RiskAssessment | EvaluacionDeRiesgo | ÉvaluationRisques | | □□□□
RiskEvidenceSynthesis | RiskEvidenceSynthesis
Schedule | Schedule | Agenda | | | □□□□
SearchParameter | SearchParameter | ParametroDeBusqueda | ParamètreRecherche | ParametroDiRicerca | □□□□
ServiceRequest | ServiceRequest | PeticiónServicio | DemandeService | RichiestaDiServizio | □□□□□□
Slot | Slot | Hueco / Zocalo / Espacio | Slot | | □□
Specimen | Specimen | Especimen | Spécimen | Campione | □□
SpecimenDefinition | SpecimenDefinition | DefinicionDeEspecimen | | DefinizioneCampione | □□□□
StructureDefinition | StructureDefinition | DefinicionDeEstructura | DéfinitionStructure | DefinizioneStruttura | □□□□
StructureMap | StructureMap | MapaDeEstructura | | | □□□□
Subscription | Subscription | Suscripción | Souscription | Sottoscrizione | □□
Substance | Substance | Sustancia | Substance | Sostanza | □□
SubstanceNucleicAcid | SubstanceNucleicAcid
SubstancePolymer | SubstancePolymer
SubstanceProtein | SubstanceProtein
SubstanceReferenceInformation | SubstanceReferenceInformation
SubstanceSourceMaterial | SubstanceSourceMaterial
SubstanceSpecification | SubstanceSpecification
SupplyDelivery | SupplyDelivery | Entrega de Suministro | Supply Livraison | | □□□□
SupplyRequest | SupplyRequest | Solicitud de Suministro | Demande d'approvisionnement | | □□□□
Task | Task | Tarea | | | □□
TerminologyCapabilities | TerminologyCapabilities | CapacidadTerminologica | | | □□□□□□
TestReport | TestReport | ReporteDePrueba | RapportTest | | □□□□
TestScript | TestScript | ScriptDePrueba | ScriptTest | | □□□□
ValueSet | ValueSet | ConjuntoValores / ConjuntoDeValores | EnsembleValeurs | | □□□□
VerificationResult | VerificationResult
VisionPrescription | VisionPrescription | PrescripcionDeVision | PrescriptionVision | | □□□□

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.500 Value Set http://hl7.org/fhir/ValueSet/reference-handling-policy

URL: https://hl7.org/fhir/R4/valueset-reference-handling-policy.html
FHIR Infrastructure Work Group | Maturity Level: N | Normative (from v4.0.0) | Use Context: Any

| This page has been approved as part of an ANSI standard. See the Conformance Package for further details.

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/reference-handling-policy
Version: | 4.0.1
Name: | ReferenceHandlingPolicy
Title: | ReferenceHandlingPolicy
Definition: | A set of flags that defines how references are supported.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.202 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ReferenceHandlingPolicy
• Resource: CapabilityStatement.rest.resource.referencePolicy (code / Required)

4.4.1.500.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/reference-handling-policy

4.4.1.500.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 5 concepts

Expansion based on http://hl7.org/fhir/reference-handling-policy version 4.0.1

All codes from system http://hl7.org/fhir/reference-handling-policy

Code | Display | Definition
literal | Literal References | The server supports and populates Literal references (i.e. using Reference.reference) where they are known (this code does not guarantee that all references are literal; see 'enforced').
logical | Logical References | The server allows logical references (i.e. using Reference.identifier).
resolves | Resolves References | The server will attempt to resolve logical references to literal references - i.e. converting Reference.identifier to Reference.reference (if resolution fails, the server may still accept resources; see logical).
enforced | Reference Integrity Enforced | The server enforces that references have integrity - e.g. it ensures that references can always be resolved. This is typically the case for clinical record systems, but often not the case for middleware/proxy systems.
local | Local References Only | The server does not support references that point to other servers.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.501 Value Set http://hl7.org/fhir/ValueSet/medicationrequest-course-of-therapy

URL: https://hl7.org/fhir/R4/valueset-medicationrequest-course-of-therapy.html
Pharmacy Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/medicationrequest-course-of-therapy
Version: | 4.0.1
Name: | medicationRequest Course of Therapy Codes
Title: | Medication request course of therapy codes
Definition: | MedicationRequest Course of Therapy Codes
Committee: | Pharmacy Work Group
OID: | 2.16.840.1.113883.4.642.3.1326 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for medicationRequest Course of Therapy Codes

• Resource: MedicationRequest.courseOfTherapyType (CodeableConcept / Example)

4.4.1.501.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/medicationrequest-course-of-therapy

4.4.1.501.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 3 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/medicationrequest-course-of-therapy version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/medicationrequest-course-of-therapy

Code | Display | Definition
continuous | Continuous long term therapy | A medication which is expected to be continued beyond the present order and which the patient should be assumed to be taking unless explicitly stopped.
acute | Short course (acute) therapy | A medication which the patient is only expected to consume for the duration of the current order and which is not expected to be renewed.
seasonal | Seasonal | A medication which is expected to be used on a part time basis at certain times of the year

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.502 Value Set http://hl7.org/fhir/ValueSet/report-action-result-codes

URL: https://hl7.org/fhir/R4/valueset-report-action-result-codes.html
FHIR Infrastructure Work Group | Maturity Level: 0 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/report-action-result-codes
Version: | 4.0.1
Name: | TestReportActionResult
Title: | TestReportActionResult
Definition: | The results of executing an action.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.720 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for TestReportActionResult

• Resource: TestReport.setup.action.operation.result (code / Required)

• Resource: TestReport.setup.action.assert.result (code / Required)

4.4.1.502.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/report-action-result-codes

4.4.1.502.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 5 concepts

Expansion based on http://hl7.org/fhir/report-action-result-codes version 4.0.1

All codes from system http://hl7.org/fhir/report-action-result-codes

Code	Display	Definition
pass	Pass	The action was successful.
skip	Skip	The action was skipped.
fail	Fail	The action failed.
warning	Warning	The action passed but with warnings.
error	Error	The action encountered a fatal error and the engine was unable to process.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.503 Value Set http://hl7.org/fhir/ValueSet/specimen-contained-preference

URL: https://hl7.org/fhir/R4/valueset-specimen-contained-preference.html
Orders and Observations Work Group | Maturity Level: 0 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/specimen-contained-preference
Version: | 4.0.1
Name: | SpecimenContainedPreference
Title: | SpecimenContainedPreference
Definition: | Degree of preference of a type of conditioned specimen.
Committee: | Orders and Observations Work Group
OID: | 2.16.840.1.113883.4.642.3.852 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for SpecimenContainedPreference

• Resource: SpecimenDefinition.typeTested.preference (code / Required)

4.4.1.503.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/specimen-contained-preference

4.4.1.503.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 2 concepts

Expansion based on http://hl7.org/fhir/specimen-contained-preference version 4.0.1

All codes from system http://hl7.org/fhir/specimen-contained-preference

Code | Display | Definition
preferred | Preferred | This type of contained specimen is preferred to collect this kind of specimen.
alternate | Alternate | This type of conditioned specimen is an alternate.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.504 Value Set http://hl7.org/fhir/ValueSet/coverage-selfpay

URL: https://hl7.org/fhir/R4/valueset-coverage-selfpay.html
Financial Management Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/coverage-selfpay
Version: | 4.0.1
Name: | CoverageSelfPayCodes
Title: | Coverage SelfPay Codes
Definition: | This value set includes Coverage SelfPay codes.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.525 (for OID based terminology systems)
Copyright: | HL7 International.
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for CoverageSelfPayCodes

4.4.1.504.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/coverage-selfpay

4.4.1.504.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 1 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/coverage-selfpay version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/coverage-selfpay

Code | Display | Definition
pay | Pay | An individual or organization is paying directly for goods and services.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.505 Value Set http://hl7.org/fhir/ValueSet/action-participant-role

URL: https://hl7.org/fhir/R4/valueset-action-participant-role.html
Clinical Decision Support Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/action-participant-role
Version: | 4.0.1
Name: | ActionParticipantRole
Title: | Action participant role
Definition: | Either a practitioner role or a relationship type.
Committee: | Clinical Decision Support Work Group
OID: | 2.16.840.1.113883.4.642.3.796 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: ActivityDefinition.participant.role (CodeableConcept / Example)
• Profile: ShareableActivityDefinition: ActivityDefinition.participant.role (CodeableConcept / Example)
• Resource: PlanDefinition.action.participant.role (CodeableConcept / Example)
• Profile: ShareablePlanDefinition: PlanDefinition.action.participant.role (CodeableConcept / Example)
• Profile: ComputablePlanDefinition: PlanDefinition.action.participant.role (CodeableConcept / Example)
• Profile: CDSHooksServicePlanDefinition: PlanDefinition.action.participant.role (CodeableConcept / Example)

4.4.1.505.1 Content Logical Definition

This value set includes codes from the following code systems:

• Import all the codes that are contained in http://hl7.org/fhir/ValueSet/practitioner-role, http://hl7.org/fhir/ValueSet/relatedperson-relationshiptype

4.4.1.505.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 0 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/v3-RoleCode version 2018-08-12, http://terminology.hl7.org/CodeSystem/v2-0131 version 2.9, http://hl7.org/fhir/ValueSet/relatedperson-relationshiptype version 4.0.1, http://terminology.hl7.org/CodeSystem/practitioner-role version 4.0.1, http://hl7.org/fhir/ValueSet/practitioner-role version 4.0.1

Code | System | Display

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.506 Value Set http://hl7.org/fhir/ValueSet/jurisdiction

URL: https://hl7.org/fhir/R4/valueset-jurisdiction.html

This value set could not be expanded by the publication tooling: Error from server: Unable to provide support for code system urn:iso:std:iso:3166:-2

Vocabulary Work Group | Maturity Level: N | Normative (from v4.0.0) | Use Context: Any

| This page has been approved as part of an ANSI standard. See the Conformance Package for further details.

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/jurisdiction
Version: | 4.0.1
Name: | Jurisdiction ValueSet
Title: | Jurisdiction
Definition: | This value set defines a base set of codes for country, country subdivision and region for indicating where a resource is intended to be used. Note: The codes for countries and country subdivisions are taken from ISO 3166 while the codes for "supra-national" regions are from UN Standard country or area codes for statistical use (M49) .
Committee: | Vocabulary Work Group
OID: | 2.16.840.1.113883.4.642.3.48 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

- Resource: EventDefinition.jurisdiction (CodeableConcept / Extensible)
- Resource: MessageDefinition.jurisdiction (CodeableConcept / Extensible)
- Resource: CapabilityStatement.jurisdiction (CodeableConcept / Extensible)
- Resource: Measure.jurisdiction (CodeableConcept / Extensible)
- Profile: ShareableMeasure: Measure.jurisdiction (CodeableConcept / Extensible)
- Resource: GraphDefinition.jurisdiction (CodeableConcept / Extensible)
- Resource: ResearchDefinition.jurisdiction (CodeableConcept / Extensible)
- Resource: ImplementationGuide.jurisdiction (CodeableConcept / Extensible)
- Resource: ResearchElementDefinition.jurisdiction (CodeableConcept / Extensible)
- Resource: SearchParameter.jurisdiction (CodeableConcept / Extensible)
- Resource: ActivityDefinition.jurisdiction (CodeableConcept / Extensible)
- Profile: ShareableActivityDefinition: ActivityDefinition.jurisdiction (CodeableConcept / Extensible)
- Resource: Questionnaire.jurisdiction (CodeableConcept / Extensible)
- Profile: CQF-Questionnaire: Questionnaire.jurisdiction (CodeableConcept / Extensible)
- Resource: ResearchStudy.location (CodeableConcept / Extensible)
- Resource: StructureDefinition.jurisdiction (CodeableConcept / Extensible)
- Resource: ChargeItemDefinition.jurisdiction (CodeableConcept / Extensible)
- Resource: ConceptMap.jurisdiction (CodeableConcept / Extensible)
- Resource: ValueSet.jurisdiction (CodeableConcept / Extensible)
- Profile: ShareableValueSet: ValueSet.jurisdiction (CodeableConcept / Extensible)
- Resource: OperationDefinition.jurisdiction (CodeableConcept / Extensible)
- Resource: EffectEvidenceSynthesis.jurisdiction (CodeableConcept / Extensible)
- Resource: EvidenceVariable.jurisdiction (CodeableConcept / Extensible)
- Profile: PICOElement: EvidenceVariable.jurisdiction (CodeableConcept / Extensible)
- Resource: CodeSystem.jurisdiction (CodeableConcept / Extensible)
- Profile: ShareableCodeSystem: CodeSystem.jurisdiction (CodeableConcept / Extensible)
- Resource: RiskEvidenceSynthesis.jurisdiction (CodeableConcept / Extensible)
- Resource: StructureMap.jurisdiction (CodeableConcept / Extensible)
- Resource: Library.jurisdiction (CodeableConcept / Extensible)
- Profile: ShareableLibrary: Library.jurisdiction (CodeableConcept / Extensible)
- Profile: CQLLibrary: Library.jurisdiction (CodeableConcept / Extensible)
- Resource: TestScript.jurisdiction (CodeableConcept / Extensible)
- Resource: TerminologyCapabilities.jurisdiction (CodeableConcept / Extensible)
- Resource: Evidence.jurisdiction (CodeableConcept / Extensible)
- Profile: Synthesis: Evidence.jurisdiction (CodeableConcept / Extensible)
- Resource: NamingSystem.jurisdiction (CodeableConcept / Extensible)
- Resource: ExampleScenario.jurisdiction (CodeableConcept / Extensible)
- Resource: PlanDefinition.jurisdiction (CodeableConcept / Extensible)
- Profile: ShareablePlanDefinition: PlanDefinition.jurisdiction (CodeableConcept / Extensible)
- Profile: ComputablePlanDefinition: PlanDefinition.jurisdiction (CodeableConcept / Extensible)
- Profile: CDSHooksServicePlanDefinition: PlanDefinition.jurisdiction (CodeableConcept / Extensible)

4.4.1.506.1 Content Logical Definition

This value set includes codes from the following code systems:

- Include all codes defined in urn:iso:std:iso:3166
- Include all codes defined in urn:iso:std:iso:3166:-2
- Include codes from http://unstats.un.org/unsd/methods/m49/m49.htm where class = region

4.4.1.506.2 Expansion

This expansion generated 01 Nov 2019

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information

Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.507 Value Set http://hl7.org/fhir/ValueSet/immunization-route

URL: https://hl7.org/fhir/R4/valueset-immunization-route.html
Public Health and Emergency Response Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/immunization-route
Version: | 4.0.1
Name: | ImmunizationRouteCodes
Title: | Immunization Route Codes
Definition: | The value set to instantiate this attribute should be drawn from a terminologically robust code system that consists of or contains concepts to support describing the administrative routes used during vaccination. This value set is provided as a suggestive example.
Committee: | Public Health and Emergency Response Work Group
OID: | 2.16.840.1.113883.4.642.3.289 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: Immunization.route (CodeableConcept / Example)

4.4.1.507.1 Content Logical Definition

This value set includes codes from the following code systems:

http://terminology.hl7.org/CodeSystem/v3-RouteOfAdministration

Code | Display
IDINJ | Injection, intradermal | Injection, intradermal
IM | Injection, intramuscular | Injection, intramuscular
NASINHLC | Inhalation, nasal | Inhalation, nasal, prongs
IVINJ | Injection, intravenous | Injection, intravenous
PO | Swallow, oral | Swallow, oral
SQ | Injection, subcutaneous | Injection, subcutaneous
TRNSDERM | Transdermal | Transdermal

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.508 Value Set http://hl7.org/fhir/ValueSet/metric-category

URL: https://hl7.org/fhir/R4/valueset-metric-category.html
Health Care Devices Work Group | Maturity Level: 1 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/metric-category
Version: | 4.0.1
Name: | DeviceMetricCategory
Title: | DeviceMetricCategory
Definition: | Describes the category of the metric.
Committee: | Health Care Devices Work Group
OID: | 2.16.840.1.113883.4.642.3.648 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for DeviceMetricCategory

• Resource: DeviceMetric.category (code / Required)

4.4.1.508.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/metric-category

4.4.1.508.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 4 concepts

Expansion based on http://hl7.org/fhir/metric-category version 4.0.1

All codes from system http://hl7.org/fhir/metric-category

Code	Display	Definition
measurement	Measurement	DeviceObservations generated for this DeviceMetric are measured.
setting	Setting	DeviceObservations generated for this DeviceMetric is a setting that will influence the behavior of the Device.
calculation	Calculation	DeviceObservations generated for this DeviceMetric are calculated.
unspecified	Unspecified	The category of this DeviceMetric is unspecified.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.509 Value Set http://hl7.org/fhir/ValueSet/research-study-objective-type

URL: https://hl7.org/fhir/R4/valueset-research-study-objective-type.html
Biomedical Research and Regulation Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/research-study-objective-type
Version: | 4.0.1
Name: | ResearchStudyObjectiveType
Title: | ResearchStudyObjectiveType
Definition: | Codes for the kind of study objective.
Committee: | Biomedical Research and Regulation Work Group
OID: | 2.16.840.1.113883.4.642.3.823 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ResearchStudyObjectiveType

• Resource: ResearchStudy.objective.type (CodeableConcept / Preferred)

4.4.1.509.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/research-study-objective-type

4.4.1.509.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 3 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/research-study-objective-type version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/research-study-objective-type

Code	Display	Definition

primary | Primary | The main question to be answered, and the one that drives any statistical planning for the study—e.g., calculation of the sample size to provide the appropriate power for statistical testing.
secondary | Secondary | Question to be answered in the study that is of lesser importance than the primary objective.
exploratory | Exploratory | Exploratory questions to be answered in the study.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.510 Value Set http://hl7.org/fhir/ValueSet/binding-strength

URL: https://hl7.org/fhir/R4/valueset-binding-strength.html
FHIR Infrastructure Work Group | Maturity Level: N | Normative (from v4.0.0) | Use Context: Any

| This page has been approved as part of an ANSI standard. See the Conformance Package for further details.

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/binding-strength
Version: | 4.0.1
Name: | BindingStrength
Title: | BindingStrength
Definition: | Indication of the degree of conformance expectations associated with a binding.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.43 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for BindingStrength
• Resource: OperationDefinition.parameter.binding.strength (code / Required)
• Resource: ElementDefinition.binding.strength (code / Required)

4.4.1.510.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/binding-strength

4.4.1.510.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 4 concepts

Expansion based on http://hl7.org/fhir/binding-strength version 4.0.1

All codes from system http://hl7.org/fhir/binding-strength

Code | Display | Definition
required | Required | To be conformant, the concept in this element SHALL be from the specified value set.
extensible | Extensible | To be conformant, the concept in this element SHALL be from the specified value set if any of the codes within the value set can apply to the concept being communicated. If the value set does not cover the concept (based on human review), alternate codings (or, data type allowing, text) may be included instead.
preferred | Preferred | Instances are encouraged to draw from the specified codes for interoperability purposes but are not required to do so to be considered conformant.
example | Example | Instances are not expected or even encouraged to draw from the specified value set. The value set merely provides examples of the types of concepts intended to be included.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application

Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.511 Value Set http://hl7.org/fhir/ValueSet/guide-parameter-code

URL: https://hl7.org/fhir/R4/valueset-guide-parameter-code.html
FHIR Infrastructure Work Group | Maturity Level: 1 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/guide-parameter-code
Version: | 4.0.1
Name: | GuideParameterCode
Title: | GuideParameterCode
Definition: | Code of parameter that is input to the guide.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.996 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for GuideParameterCode

• Resource: ImplementationGuide.definition.parameter.code (code / Required)

4.4.1.511.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/guide-parameter-code

4.4.1.511.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 10 concepts

Expansion based on http://hl7.org/fhir/guide-parameter-code version 4.0.1

All codes from system http://hl7.org/fhir/guide-parameter-code

Code | Display | Definition
apply | Apply Metadata Value | If the value of this string 0..* parameter is one of the metadata fields then all conformance resources will have any specified [Resource].[field] overwritten with the ImplementationGuide.[field], where field is one of: version, date, status, publisher, contact, copyright, experimental, jurisdiction, useContext.
path-resource | Resource Path | The value of this string 0..* parameter is a subfolder of the build context's location that is to be scanned to load resources. Scope is (if present) a particular resource type.
path-pages | Pages Path | The value of this string 0..1 parameter is a subfolder of the build context's location that contains files that are part of the html content processed by the builder.
path-tx-cache | Terminology Cache Path | The value of this string 0..1 parameter is a subfolder of the build context's location that is used as the terminology cache. If this is not present, the terminology cache is on the local system, not under version control.
expansion-parameter | Expansion Profile | The value of this string 0..* parameter is a parameter (name=value) when expanding value sets for this implementation guide. This is particularly used to specify the versions of published terminologies such as SNOMED CT.
rule-broken-links | Broken Links Rule | The value of this string 0..1 parameter is either "warning" or "error" (default = "error"). If the value is "warning" then IG build tools allow the IG to be considered successfully build even when there is no internal broken links.
generate-xml | Generate XML | The value of this boolean 0..1 parameter specifies whether the IG publisher creates examples in XML format. If not present, the Publication Tool decides whether to generate XML.
generate-json | Generate JSON | The value of this boolean 0..1 parameter specifies whether the IG publisher creates examples in JSON format. If not present, the Publication Tool decides whether to generate JSON.
generate-turtle | Generate Turtle | The value of this boolean 0..1 parameter specifies whether the IG publisher creates examples in Turtle format. If not present, the Publication Tool decides whether to generate Turtle.
html-template | HTML Template | The value of this string singleton parameter is the name of the file to use as the builder template for each generated page (see templating).

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.512 Value Set http://hl7.org/fhir/ValueSet/search-modifier-code

URL: https://hl7.org/fhir/R4/valueset-search-modifier-code.html
FHIR Infrastructure Work Group | Maturity Level: 3 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/search-modifier-code
Version: | 4.0.1
Name: | SearchModifierCode
Title: | SearchModifierCode
Definition: | A supported modifier for a search parameter.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.639 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for SearchModifierCode

• Resource: SearchParameter.modifier (code / Required)

4.4.1.512.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/search-modifier-code

4.4.1.512.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 12 concepts

Expansion based on http://hl7.org/fhir/search-modifier-code version 4.0.1

All codes from system http://hl7.org/fhir/search-modifier-code

Code | Display | Definition
missing | Missing | The search parameter returns resources that have a value or not.
exact | Exact | The search parameter returns resources that have a value that exactly matches the supplied parameter (the whole string, including casing and accents).
contains | Contains | The search parameter returns resources that include the supplied parameter value anywhere within the field being searched.
not | Not | The search parameter returns resources that do not contain a match.
text | Text | The search parameter is processed as a string that searches text associated with the code/value - either CodeableConcept.text, Coding.display, or Identifier.type.text.
in | In | The search parameter is a URI (relative or absolute) that identifies a value set, and the search parameter tests whether the coding is in the specified value set.
not-in | Not In | The search parameter is a URI (relative or absolute) that identifies a value set, and the search parameter tests whether the coding is not in the specified value set.
below | Below | The search parameter tests whether the value in a resource is subsumed by the specified value (is-a, or hierarchical relationships).
above | Above | The search parameter tests whether the value in a resource subsumes the specified value (is-a, or hierarchical relationships).
type | Type | The search parameter only applies to the Resource Type specified as a modifier (e.g. the modifier is not actually :type, but :Patient etc.).
identifier | Identifier | The search parameter applies to the identifier on the resource, not the reference.
ofType | Of Type | The search parameter has the format system|code|value, where the system and code refer to an Identifier.type.coding.system and .code, and match if any of the type codes match. All 3 parts must be present.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.513 Value Set http://hl7.org/fhir/ValueSet/device-status-reason

URL: https://hl7.org/fhir/R4/valueset-device-status-reason.html
Orders and Observations Work Group | Maturity Level: 2 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/device-status-reason
Version: | 4.0.1
Name: | FHIRDeviceStatusReason
Title: | FHIRDeviceStatusReason
Definition: | The availability status reason of the device.
Committee: | Orders and Observations Work Group
OID: | 2.16.840.1.113883.4.642.3.1081 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for FHIRDeviceStatusReason

• Resource: Device.statusReason (CodeableConcept / Extensible)

4.4.1.513.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/device-status-reason

4.4.1.513.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 8 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/device-status-reason version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/device-status-reason

Code | Display | Definition
online | Online | The device is off.
paused | Paused | The device is paused.
standby | Standby | The device is ready but not actively operating.
offline | Offline | The device is offline.
not-ready | Not Ready | The device is not ready.
transduc-discon | Transducer Disconnected | The device transducer is disconnected.
hw-discon | Hardware Disconnected | The device hardware is disconnected.
off | Off | The device is off.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.514 Value Set http://hl7.org/fhir/ValueSet/question-max-occurs

URL: https://hl7.org/fhir/R4/valueset-question-max-occurs.html
FHIR Infrastructure Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/question-max-occurs
Version: | 4.0.1
Name: | MaxOccurs
Title: | MaxOccurs
Definition: | Flags an element as having unlimited repetitions.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.929 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for MaxOccurs

4.4.1.514.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/question-max-occurs

4.4.1.514.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 1 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/question-max-occurs version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/question-max-occurs

Code	Display	Definition
*	Repeating	Element can repeat an unlimited number of times.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.515 Value Set http://hl7.org/fhir/ValueSet/operation-kind

URL: https://hl7.org/fhir/R4/valueset-operation-kind.html
FHIR Infrastructure Work Group | Maturity Level: N | Normative (from v4.0.0) | Use Context: Any

| This page has been approved as part of an ANSI standard. See the Conformance Package for further details.

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/operation-kind
Version: | 4.0.1
Name: | OperationKind
Title: | OperationKind
Definition: | Whether an operation is a normal operation or a query.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.506 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for OperationKind

• Resource: OperationDefinition.kind (code / Required)

4.4.1.515.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/operation-kind

4.4.1.515.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 2 concepts

Expansion based on http://hl7.org/fhir/operation-kind version 4.0.1

All codes from system http://hl7.org/fhir/operation-kind

Code	Display	Definition
operation	Operation	This operation is invoked as an operation.
query	Query	This operation is a named query, invoked using the search mechanism.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')

Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.516 Value Set http://hl7.org/fhir/ValueSet/contract-assetsubtype

URL: https://hl7.org/fhir/R4/valueset-contract-assetsubtype.html
Financial Management Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/contract-assetsubtype
Version: | 4.0.1
Name: | ContractResourceAssetSub-TypeCodes
Title: | Contract Resource Asset Sub-Type codes
Definition: | This value set contract specific codes for asset subtype.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.1301 (for OID based terminology systems)
Copyright: | HL7 International.
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ContractResourceAssetSub-TypeCodes

• Resource: Contract.term.asset.subtype (CodeableConcept / Example)

4.4.1.516.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/contract-asset-subtype

4.4.1.516.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 1 concepts

Expansion based on http://hl7.org/fhir/contract-asset-subtype version 4.0.1

All codes from system http://hl7.org/fhir/contract-asset-subtype

Code | Display | Definition
participation | Participation | To be completed

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.517 Value Set http://hl7.org/fhir/ValueSet/secondary-finding

URL: https://hl7.org/fhir/R4/valueset-secondary-finding.html
Orders and Observations Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/secondary-finding
Version: | 4.0.1
Name: | ObservationCategoryCodes
Title: | Observation Category Codes
Definition: | Codes to denote a guideline or policy statement.when a genetic test result is being shared as a secondary finding.
Committee: | Orders and Observations Work Group
OID: | 2.16.840.1.113883.4.642.3.1285 (for OID based terminology systems)

Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ObservationCategoryCodes

• Extension: http://hl7.org/fhir/StructureDefinition/observation-secondaryFinding: secondaryFinding (CodeableConcept / Extensible)

4.4.1.517.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/secondary-finding

4.4.1.517.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 2 concepts

Expansion based on http://hl7.org/fhir/secondary-finding version 4.0.1

All codes from system http://hl7.org/fhir/secondary-finding

Code | Display | Definition
acmg-version1 | ACMG Version 1 | First release (2013): ACMG Recommendations for Reporting of Incidental Findings in Clinical Exome and Genome Sequencing. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3727274/
acmg-version2 | ACMG Version 2 | Second release (2016): Recommendations for reporting of secondary findings in clinical exome and genome sequencing, 2016 update (ACMG SF v2.0): a policy statement of the American College of Medical Genetics and Genomics. https://www.ncbi.nlm.nih.gov/pubmed/27854360

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.518 Value Set http://hl7.org/fhir/ValueSet/fundsreserve

URL: https://hl7.org/fhir/R4/valueset-fundsreserve.html
Financial Management Work Group | Maturity Level: 1 | Draft | Use Context: Not Intended for Production use

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/fundsreserve
Version: | 4.0.1
Name: | Funds Reservation Codes
Title: | FundsReserve
Definition: | This value set includes sample funds reservation type codes.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.33 (for OID based terminology systems)
Copyright: | This is an example set.
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for Funds Reservation Codes

• Resource: ExplanationOfBenefit.fundsReserveRequested (CodeableConcept / Example)

• Resource: ExplanationOfBenefit.fundsReserve (CodeableConcept / Example)

• Resource: ClaimResponse.fundsReserve (CodeableConcept / Example)

• Resource: Claim.fundsReserve (CodeableConcept / Example)

4.4.1.518.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/fundsreserve

4.4.1.518.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 3 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/fundsreserve version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/fundsreserve

Code | Display | Definition
patient | Patient | The payor is requested to reserve funds for the provision of the named services by any provider for settlement of future claims related to this request.
provider | Provider | The payor is requested to reserve funds solely for the named provider for settlement of future claims related to this request.
none | None | The payor is not being requested to reserve any funds for the settlement of future claims.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.519 Value Set http://hl7.org/fhir/ValueSet/contract-party-role

URL: https://hl7.org/fhir/R4/valueset-contract-party-role.html
Financial Management Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/contract-party-role
Version: | 4.0.1
Name: | ContractResourcePartyRoleCodes
Title: | Contract Resource Party Role codes
Definition: | This value set contract specific codes for offer party participation.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.1224 (for OID based terminology systems)
Copyright: | HL7 International.
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ContractResourcePartyRoleCodes

• Resource: Contract.term.offer.party.role (CodeableConcept / Example)

4.4.1.519.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/contract-party-role

4.4.1.519.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 1 concepts

Expansion based on http://hl7.org/fhir/contract-party-role version 4.0.1

All codes from system http://hl7.org/fhir/contract-party-role

Code | Display | Definition
flunky | FLunky | To be completed

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.520 Value Set http://hl7.org/fhir/ValueSet/special-values

URL: https://hl7.org/fhir/R4/valueset-special-values.html
FHIR Infrastructure Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/special-values
Version: | 4.0.1
Name: | SpecialValues
Title: | SpecialValues
Definition: | A set of generally useful codes defined so they can be included in value sets.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.9 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for SpecialValues

• This value set is used implicitly when searching boolean values

4.4.1.520.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/special-values

4.4.1.520.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 6 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/special-values version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/special-values

Code | Display | Definition
true | true | Boolean true.
false | false | Boolean false.
trace | Trace Amount Detected | The content is greater than zero, but too small to be quantified.
sufficient | Sufficient Quantity | The specific quantity is not known, but is known to be non-zero and is not specified because it makes up the bulk of the material.
withdrawn | Value Withdrawn | The value is no longer available.
nil-known | Nil Known | The are no known applicable values in this context.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.521 Value Set http://hl7.org/fhir/ValueSet/observation-codes

URL: https://hl7.org/fhir/R4/valueset-observation-codes.html
FHIR Infrastructure Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/observation-codes
Version: | 4.0.1
Name: | LOINCCodes
Title: | LOINC Codes
Definition: | This value set includes all LOINC codes
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.396 (for OID based terminology systems)
Copyright: | This content from LOINC® is copyright © 1995 Regenstrief Institute, Inc. and the LOINC Committee, and available at no cost under the license at http://loinc.org/terms-of-use.
Source Resource | XML / JSON

This value set is used in the following places:

- Resource: Goal.target.measure (CodeableConcept / Example)
- Resource: ObservationDefinition.code (CodeableConcept / Example)
- Profile: DeviceMetricObservation: Observation.code (CodeableConcept / Example)
- Profile: DeviceMetricObservation: Observation.component.code (CodeableConcept / Example)
- Profile: Observation-Genetics: Observation.code (CodeableConcept / Example)
- Profile: Observation-Genetics: Observation.component.code (CodeableConcept / Example)
- Profile: Cholesterol: Observation.code (CodeableConcept / Example)
- Profile: Cholesterol: Observation.component.code (CodeableConcept / Example)
- Profile: Triglyceride: Observation.code (CodeableConcept / Example)
- Profile: Triglyceride: Observation.component.code (CodeableConcept / Example)
- Profile: HDLCholesterol: Observation.code (CodeableConcept / Example)
- Profile: HDLCholesterol: Observation.component.code (CodeableConcept / Example)
- Profile: LDLCholesterol: Observation.component.code (CodeableConcept / Example)
- Resource: PlanDefinition.goal.target.measure (CodeableConcept / Example)
- Profile: ShareablePlanDefinition: PlanDefinition.goal.target.measure (CodeableConcept / Example)
- Profile: ComputablePlanDefinition: PlanDefinition.goal.target.measure (CodeableConcept / Example)
- Profile: CDSHooksServicePlanDefinition: PlanDefinition.goal.target.measure (CodeableConcept / Example)
- Resource: ElementDefinition.code (Coding / Example)

4.4.1.521.1 Content Logical Definition

This value set includes codes from the following code systems:

- Include all codes defined in http://loinc.org

4.4.1.521.2 Expansion

This expansion generated 01 Nov 2019

This value set has >1000 codes in it. In order to keep the publication size manageable, only a selection (1000 codes) of the whole set of codes is shown

All codes from system http://loinc.org

Code	Display
1-8	Acyclovir [Susceptibility]
10-9	Amdinocillin [Susceptibility] by Serum bactericidal titer
100-8	Cefoperazone [Susceptibility] by Minimum inhibitory concentration (MIC)
1000-9	DBG Ab [Presence] in Serum or Plasma from Blood product unit
10000-8	R wave duration in lead AVR
10001-6	R wave duration in lead I
10002-4	R wave duration in lead II
10003-2	R wave duration in lead III
10004-0	R wave duration in lead V1
10005-7	R wave duration in lead V2
10006-5	R wave duration in lead V3
10007-3	R wave duration in lead V4
10008-1	R wave duration in lead V5
10009-9	R wave duration in lead V6
1001-7	DBG Ab [Presence] in Serum or Plasma from Donor
10010-7	R' wave amplitude in lead AVF
10011-5	R' wave amplitude in lead AVL
10012-3	R' wave amplitude in lead AVR
10013-1	R' wave amplitude in lead I
10014-9	R' wave amplitude in lead II
10015-6	R' wave amplitude in lead III
10016-4	R' wave amplitude in lead V1
10017-2	R' wave amplitude in lead V2
10018-0	R' wave amplitude in lead V3
10019-8	R' wave amplitude in lead V4
1002-5	DBG Ab [Presence] in Serum or Plasma
10020-6	R' wave amplitude in lead V5
10021-4	R' wave amplitude in lead V6
10022-2	R' wave duration in lead AVF
10023-0	R' wave duration in lead AVL
10024-8	R' wave duration in lead AVR
10025-5	R' wave duration in lead I
10026-3	R' wave duration in lead II

10027-1 | R' wave duration in lead III
10028-9 | R' wave duration in lead V1
10029-7 | R' wave duration in lead V2
1003-3 | Indirect antiglobulin test.complement specific reagent [Presence] in Serum or Plasma
10030-5 | R' wave duration in lead V3
10031-3 | R' wave duration in lead V4
10032-1 | R' wave duration in lead V5
10033-9 | R' wave duration in lead V6
10034-7 | S wave amplitude in lead AVF
10035-4 | S wave amplitude in lead AVL
10036-2 | S wave amplitude in lead AVR
10037-0 | S wave amplitude in lead I
10038-8 | S wave amplitude in lead II
10039-6 | S wave amplitude in lead III
1004-1 | Direct antiglobulin test.complement specific reagent [Presence] on Red Blood Cells
10040-4 | S wave amplitude in lead V1
10041-2 | S wave amplitude in lead V2
10042-0 | S wave amplitude in lead V3
10043-8 | S wave amplitude in lead V4
10044-6 | S wave amplitude in lead V5
10045-3 | S wave amplitude in lead V6
10046-1 | S wave duration in lead AVF
10047-9 | S wave duration in lead AVL
10048-7 | S wave duration in lead AVR
10049-5 | S wave duration in lead I
1005-8 | Indirect antiglobulin test.IgG specific reagent [Presence] in Serum or Plasma
10050-3 | S wave duration in lead II
10051-1 | S wave duration in lead III
10052-9 | S wave duration in lead V1
10053-7 | S wave duration in lead V2
10054-5 | S wave duration in lead V3
10055-2 | S wave duration in lead V4
10056-0 | S wave duration in lead V5
10057-8 | S wave duration in lead V6
10058-6 | S' wave amplitude in lead AVF
10059-4 | S' wave amplitude in lead AVL
1006-6 | Direct antiglobulin test.IgG specific reagent [Interpretation] on Red Blood Cells
10060-2 | S' wave amplitude in lead AVR
10061-0 | S' wave amplitude in lead I
10062-8 | S' wave amplitude in lead II
10063-6 | S' wave amplitude in lead III
10064-4 | S' wave amplitude in lead V1
10065-1 | S' wave amplitude in lead V2
10066-9 | S' wave amplitude in lead V3
10067-7 | S' wave amplitude in lead V4
10068-5 | S' wave amplitude in lead V5
10069-3 | S' wave amplitude in lead V6
1007-4 | Direct antiglobulin test.poly specific reagent [Presence] on Red Blood Cells
10070-1 | S' wave duration in lead AVF
10071-9 | S' wave duration in lead AVL
10072-7 | S' wave duration in lead AVR
10073-5 | S' wave duration in lead I
10074-3 | S' wave duration in lead II
10075-0 | S' wave duration in lead III
10076-8 | S' wave duration in lead V1
10077-6 | S' wave duration in lead V2
10078-4 | S' wave duration in lead V3
10079-2 | S' wave duration in lead V4
1008-2 | Indirect antiglobulin test.poly specific reagent [Presence] in Serum or Plasma
10080-0 | S' wave duration in lead V5
10081-8 | S' wave duration in lead V6
10082-6 | ST initial amplitude 6 ms in lead AVF
10083-4 | ST initial amplitude 6 ms in lead AVL
10084-2 | ST initial amplitude 6 ms in lead AVR
10085-9 | ST initial amplitude 6 ms in lead I
10086-7 | ST initial amplitude 6 ms in lead II
10087-5 | ST initial amplitude 6 ms in lead III
10088-3 | ST initial amplitude 6 ms in lead V1
10089-1 | ST initial amplitude 6 ms in lead V2
1009-0 | Deprecated Direct antiglobulin test.poly specific reagent [Presence] on Red Blood Cells
10090-9 | ST initial amplitude 6 ms in lead V3
10091-7 | ST initial amplitude 6 ms in lead V4

10092-5 | ST initial amplitude 6 ms in lead V5
10093-3 | ST initial amplitude 6 ms in lead V6
10094-1 | ST slope in lead AVF
10095-8 | ST slope in lead AVL
10096-6 | ST slope in lead AVR
10097-4 | ST slope in lead I
10098-2 | ST slope in lead II
10099-0 | ST slope in lead III
101-6 | Cefoperazone [Susceptibility] by Disk diffusion (KB)
1010-8 | E sup(w) Ab [Presence] in Serum or Plasma from Blood product unit
10100-6 | ST slope in lead V1
10101-4 | ST slope in lead V2
10102-2 | ST slope in lead V3
10103-0 | ST slope in lead V4
10104-8 | ST slope in lead V5
10105-5 | ST slope in lead V6
10106-3 | ST wave end displacement in lead AVF
10107-1 | ST wave end displacement in lead AVL
10108-9 | ST wave end displacement in lead AVR
10109-7 | ST wave end displacement in lead I
1011-6 | E sup(w) Ab [Presence] in Serum or Plasma from Donor
10110-5 | ST wave end displacement in lead II
10111-3 | ST wave end displacement in lead III
10112-1 | ST wave end displacement in lead V1
10113-9 | ST wave end displacement in lead V2
10114-7 | ST wave end displacement in lead V3
10115-4 | ST wave end displacement in lead V4
10116-2 | ST wave end displacement in lead V5
10117-0 | ST wave end displacement in lead V6
10118-8 | ST wave mid displacement in lead AVF
10119-6 | ST wave mid displacement in lead AVL
1012-4 | E sup(w) Ab [Presence] in Serum or Plasma
10120-4 | ST wave mid displacement in lead AVR
10121-2 | ST wave mid displacement in lead I
10122-0 | ST wave mid displacement in lead II
10123-8 | ST wave mid displacement in lead III
10124-6 | ST wave mid displacement in lead V1
10125-3 | ST wave mid displacement in lead V2
10126-1 | ST wave mid displacement in lead V3
10127-9 | ST wave mid displacement in lead V4
10128-7 | ST wave mid displacement in lead V5
10129-5 | ST wave mid displacement in lead V6
1013-2 | E sup(w) Ag [Presence] on Red Blood Cells from Blood product unit
10130-3 | T' wave amplitude in lead AVF
10131-1 | T' wave amplitude in lead AVL
10132-9 | T' wave amplitude in lead AVR
10133-7 | T' wave amplitude in lead I
10134-5 | T' wave amplitude in lead II
10135-2 | T' wave amplitude in lead III
10136-0 | T' wave amplitude in lead V1
10137-8 | T' wave amplitude in lead V2
10138-6 | T' wave amplitude in lead V3
10139-4 | T' wave amplitude in lead V4
1014-0 | E sup(w) Ag [Presence] on Red Blood Cells from Donor
10140-2 | T' wave amplitude in lead V5
10141-0 | T' wave amplitude in lead V6
10142-8 | T wave amplitude in lead AVF
10143-6 | T wave amplitude in lead AVL
10144-4 | T wave amplitude in lead AVR
10145-1 | T wave amplitude in lead I
10146-9 | T wave amplitude in lead II
10147-7 | T wave amplitude in lead III
10148-5 | T wave amplitude in lead V1
10149-3 | T wave amplitude in lead V2
1015-7 | E sup(w) Ag [Presence] on Red Blood Cells
10150-1 | T wave amplitude in lead V3
10151-9 | T wave amplitude in lead V4
10152-7 | T wave amplitude in lead V5
10153-5 | T wave amplitude in lead V6
10154-3 | Chief complaint Narrative - Reported
10155-0 | History of allergies, reported
10156-8 | History of Childhood diseases Narrative

10157-6 | History of family member diseases Narrative
10158-4 | History of Functional status Narrative
10159-2 | History of Industrial exposure Narrative
1016-5 | E Ab [Presence] in Serum or Plasma from Blood product unit
10160-0 | History of Medication use Narrative
10161-8 | History of Occupational exposure Narrative
10162-6 | History of pregnancies Narrative
10163-4 | History of pregnancies
10164-2 | History of Present illness Narrative
10165-9 | Deprecated History of psychiatric symptoms and diseases Narrative
10166-7 | History of Social function Narrative
10167-5 | History of Surgical procedures Narrative
10168-3 | History of Cardiovascular system disorders Narrative
10169-1 | History of Ear disorders Narrative
1017-3 | E Ab [Presence] in Serum or Plasma from Donor
10170-9 | History of Endocrine system disorders Narrative
10171-7 | History of Eyes disorders Narrative
10172-5 | History of Hematologic system disorders Narrative
10173-3 | History of Musculoskeletal system disorders Narrative
10174-1 | History of Nose disorders Narrative
10175-8 | History of Oral cavity disorders Narrative
10176-6 | History of Reproductive system disorders Narrative
10177-4 | History of Respiratory system disorders Narrative
10178-2 | History of Skin disorders Narrative
10179-0 | History of Throat and Neck disorders Narrative
1018-1 | E Ab [Presence] in Serum or Plasma
10180-8 | History of Throat and Neck disorders
10181-6 | History of Urinary tract disorders Narrative
10182-4 | History of Travel Narrative
10183-2 | Hospital discharge medications Narrative
10184-0 | Hospital discharge physical findings Narrative
10185-7 | Hospital discharge procedures Narrative
10186-5 | Identifying information Narrative Observed
10187-3 | Review of systems Narrative - Reported
10188-1 | Review of systems overview Narrative - Reported
10189-9 | Review of systems overview - Reported
1019-9 | E Ag [Presence] on Red Blood Cells from Blood product unit
10190-7 | Mental status Narrative
10191-5 | Physical findings of Abdomen Narrative
10192-3 | Physical findings of Back Narrative
10193-1 | Physical findings of Breasts Narrative
10194-9 | Physical findings of Neurologic deep tendon reflexes Narrative
10195-6 | Physical findings of Ear Narrative
10196-4 | Physical findings of Extremities Narrative
10197-2 | Physical findings of Eye Narrative
10198-0 | Physical findings of Genitourinary tract Narrative
10199-8 | Physical findings of Head Narrative
102-4 | Cefoperazone [Susceptibility] by Serum bactericidal titer
1020-7 | E Ag [Presence] on Red Blood Cells from Donor
10200-4 | Physical findings of Heart Narrative
10201-2 | Physical findings of Mouth and Throat and Teeth Narrative
10202-0 | Physical findings of Nervous system Narrative
10203-8 | Physical findings of Nose Narrative
10204-6 | Physical findings of Pelvis Narrative
10205-3 | Physical findings of Rectum Narrative
10206-1 | Physical findings of Skin Narrative
10207-9 | Physical findings of Thorax and Lungs Narrative
10208-7 | Physical findings of Vessels Narrative
10209-5 | Physical findings of Neurologic balance and Coordination Narrative
1021-5 | E Ag [Presence] on Red Blood Cells
10210-3 | Physical findings of General status Narrative
10211-1 | Physical findings of Sense of touch Narrative
10212-9 | Physical findings of Strength Narrative
10213-7 | Surgical operation note anesthesia Narrative
10214-5 | Surgical operation note anesthesia duration
10215-2 | Surgical operation note findings Narrative
10216-0 | Surgical operation note fluids Narrative
10217-8 | Surgical operation note indications [Interpretation] Narrative
10218-6 | Surgical operation note postoperative diagnosis Narrative
10219-4 | Surgical operation note preoperative diagnosis Narrative
1022-3 | Fy sup(a) Ab [Presence] in Serum or Plasma from Blood product unit
10220-2 | Surgical operation note prep time duration

10221-0 | Surgical operation note specimens taken Narrative
10222-8 | Surgical operation note surgical complications [Interpretation] Narrative
10223-6 | Surgical operation note surgical procedure Narrative
10224-4 | Hemodynamic method special circumstances
10225-1 | Cardiac measurement device Institution inventory number
10226-9 | Oxygen content in Intravascular space
10227-7 | Cardiac measurement device Vendor model number
10228-5 | Cardiac measurement device Vendor serial number
10229-3 | Hemodynamic method Type of
1023-1 | Fy sup(a) Ab [Presence] in Serum or Plasma from Donor
10230-1 | Left ventricular Ejection fraction
10231-9 | Right ventricular Ejection fraction
10232-7 | Oxygen content in Aorta root
10233-5 | Oxygen content in Left atrium
10234-3 | Oxygen content in Right atrium
10235-0 | Oxygen content in High right atrium
10236-8 | Oxygen content in Low right atrium
10237-6 | Oxygen content in Mid right atrium
10238-4 | Oxygen content in Left ventricle
10239-2 | Oxygen content in Right ventricular outflow tract
1024-9 | Fy sup(a) Ab [Presence] in Serum or Plasma
10240-0 | Oxygen content in Right ventricle
10241-8 | Oxygen content in Coronary sinus
10242-6 | Oxygen content in Ductus arteriosus
10243-4 | Oxygen content in Inferior vena cava
10244-2 | Oxygen content in Left pulmonary artery
10245-9 | Oxygen content in Main pulmonary artery
10246-7 | Oxygen content in Right pulmonary artery
10247-5 | Oxygen content in Pulmonary wedge
10248-3 | Oxygen content in Superior vena cava
10249-1 | Heart rate device Institution inventory number
1025-6 | Fy sup(a) Ag [Presence] on Red Blood Cells from Blood product unit
10250-9 | Heart rate device Vendor model number
10251-7 | Heart rate device Vendor serial number
10252-5 | Biliary drain site
10253-3 | Type of Biliary drain
10254-1 | Type of Peritoneal drain
10255-8 | Type of Subarachnoid drain
10256-6 | Biliary drain Institution inventory number
10257-4 | Chest tube Institution inventory number
10258-2 | Enteral tube Institution inventory number
10259-0 | Gastrostomy tube Institution inventory number
1026-4 | Fy sup(a) Ag [Presence] on Red Blood Cells from Donor
10260-8 | GI tube Institution inventory number
10261-6 | Intravascular tube Institution inventory number
10262-4 | IO tube Institution inventory number
10263-2 | Nasogastric tube Institution inventory number
10264-0 | Oral tube Institution inventory number
10265-7 | Peritoneal drain Institution inventory number
10266-5 | Peritoneal tube Institution inventory number
10267-3 | Rate control device Institution inventory number
10268-1 | Stool collection device Institution inventory number
10269-9 | Subarachnoid drain Institution inventory number
1027-2 | Fy sup(a) Ag [Presence] on Red Blood Cells
10270-7 | Synovial drain Institution inventory number
10271-5 | Tube or drain Institution inventory number
10272-3 | Upper GI tube Institution inventory number
10273-1 | Bladder irrigation tube Institution inventory number
10274-9 | Urine tube Institution inventory number
10275-6 | Wound drain device Institution inventory number
10276-4 | Peritoneal drain site
10277-2 | Stool collection site
10278-0 | Subarachnoid drain site
10279-8 | Biliary drain Vendor model number
1028-0 | Fy sup(b) Ab [Presence] in Serum or Plasma from Blood product unit
10280-6 | Chest tube Vendor model number
10281-4 | Enteral tube Vendor model number
10282-2 | Gastrostomy tube Vendor model number
10283-0 | GI tube Vendor model number
10284-8 | Intravascular tube Vendor model number
10285-5 | IO tube Vendor model number
10286-3 | Nasogastric tube Vendor model number

10287-1 | Oral tube Vendor model number
10288-9 | Peritoneal drain Vendor model number
10289-7 | Peritoneal tube Vendor model number
1029-8 | Fy sup(b) Ab [Presence] in Serum or Plasma from Donor
10290-5 | Rate control device Vendor model number
10291-3 | Stool collection device Vendor model number
10292-1 | Subarachnoid drain Vendor model number
10293-9 | Synovial drain Vendor model number
10294-7 | Tube or drain Vendor model number
10295-4 | Upper GI tube Vendor model number
10296-2 | Bladder irrigation tube Vendor model number
10297-0 | Urine tube Vendor model number
10298-8 | Wound drain device Vendor model number
10299-6 | Biliary drain Vendor serial number
103-2 | Ceforanide [Susceptibility] by Minimum lethal concentration (MLC)
1030-6 | Fy sup(b) Ab [Presence] in Serum or Plasma
10300-2 | Chest tube Vendor serial number
10301-0 | Enteral tube Vendor serial number
10302-8 | Gastrostomy tube Vendor serial number
10303-6 | GI tube Vendor serial number
10304-4 | Intravascular tube Vendor serial number
10305-1 | IO tube Vendor serial number
10306-9 | Nasogastric tube Vendor serial number
10307-7 | Oral tube Vendor serial number
10308-5 | Peritoneal drain Vendor serial number
10309-3 | Peritoneal tube Vendor serial number
1031-4 | Fy sup(b) Ag [Presence] on Red Blood Cells from Blood product unit
10310-1 | Rate control device Vendor serial number
10311-9 | Stool collection device Vendor serial number
10312-7 | Subarachnoid drain Vendor serial number
10313-5 | Synovial drain Vendor serial number
10314-3 | Tube or drain Vendor serial number
10315-0 | Upper GI tube Vendor serial number
10316-8 | Bladder irrigation tube Vendor serial number
10317-6 | Urine tube Vendor serial number
10318-4 | Wound drain device Vendor serial number
10319-2 | Enema device Institution inventory number
1032-2 | Fy sup(b) Ag [Presence] on Red Blood Cells from Donor
10320-0 | Enema device Vendor model number
10321-8 | Enema device Vendor serial number
10322-6 | Potassium intake 24 hour
10323-4 | Wound drain fluid [Appearance] Lower GI tract
10324-2 | Breath rate device Institution inventory number
10325-9 | Breath rate device Vendor model number
10326-7 | Breath rate device Vendor serial number
10327-5 | Eosinophils/100 leukocytes in Sputum by Manual count
10328-3 | Lymphocytes/100 leukocytes in Cerebral spinal fluid by Manual count
10329-1 | Monocytes/100 leukocytes in Cerebral spinal fluid by Manual count
1033-0 | Fy sup(b) Ag [Presence] on Red Blood Cells
10330-9 | Monocytes/100 leukocytes in Body fluid by Manual count
10331-7 | Rh [Type] in Blood
10332-5 | Cortisol [Mass/volume] in Serum or Plasma --pre 250 ug corticotropin IM
10333-3 | Appearance of Cerebral spinal fluid
10334-1 | Cancer Ag 125 [Units/volume] in Serum or Plasma
10335-8 | Color of Cerebral spinal fluid
10336-6 | Gonadotropin peptide [Mass/volume] in Urine
10337-4 | Procollagen type I [Mass/volume] in Serum
10338-2 | Barbiturates [Mass/volume] in Serum or Plasma
10339-0 | Fluoxetine+Norfluoxetine [Mass/volume] in Serum or Plasma
1034-8 | Fetal cell screen [Interpretation] in Blood
10340-8 | Molindone [Mass/volume] in Serum or Plasma
10341-6 | Norpropoxyphene [Mass/volume] in Serum or Plasma
10342-4 | Sulfamethoxazole [Mass/volume] in Serum or Plasma
10343-2 | Temazepam [Mass/volume] in Serum or Plasma
10344-0 | Tranylcypromine [Mass/volume] in Serum or Plasma
10345-7 | Trihexyphenidyl [Mass/volume] in Urine
10346-5 | Hemoglobin A [Units/volume] in Blood by Electrophoresis
10347-3 | Babesia microti identified in Blood by Light microscopy
10348-1 | Bordetella parapertussis Ab [Presence] in Serum
10349-9 | Brucella sp Ab [Units/volume] in Serum
1035-5 | Fresh frozen plasma given [Volume]
10350-7 | Herpes simplex virus IgM Ab [Titer] in Serum by Immunoassay

10351-5 | HIV 1 RNA [Units/volume] (viral load) in Serum or Plasma by Probe with amplification
10352-3 | Bacteria identified in Genital specimen by Aerobe culture
10353-1 | Bacteria identified in Nose by Aerobe culture
10354-9 | Bacteria identified in Urethra by Culture
10355-6 | Microscopic observation [Identifier] in Bone marrow by Wright Giemsa stain
10356-4 | Deprecated Microscopic observation [Identifier] in Stool by Trichrome stain
10357-2 | Microscopic observation [Identifier] in Wound by Gram stain
10358-0 | Teichoate Ab [Titer] in Serum by Immune diffusion (ID)
10359-8 | Asialoganglioside GM1 IgM Ab [Titer] in Serum
1036-3 | G Ab [Presence] in Serum or Plasma from Blood product unit
10360-6 | DNA single strand IgG Ab [Units/volume] in Serum
10361-4 | DNA single strand IgM Ab [Units/volume] in Serum
10362-2 | Endomysium IgA Ab [Presence] in Serum
10363-0 | Barbiturates [Presence] in Unspecified specimen
10364-8 | Cotinine [Mass/mass] in Hair
10365-5 | Cotinine [Mass/volume] in Serum or Plasma
10366-3 | Cotinine [Mass/volume] in Urine
10367-1 | Ethanol [Mass/volume] in Gastric fluid
10368-9 | Lead [Mass/volume] in Capillary blood
10369-7 | Opiates [Mass/mass] in Hair
1037-1 | G Ab [Presence] in Serum or Plasma from Donor
10370-5 | Phencyclidine [Mass/mass] in Hair
10371-3 | Bite cells [Presence] in Blood by Light microscopy
10372-1 | Blister cells [Presence] in Blood by Light microscopy
10373-9 | Fragments [Presence] in Blood by Light microscopy
10374-7 | Helmet cells [Presence] in Blood by Light microscopy
10375-4 | Irregularly contracted cells [Presence] in Blood by Light microscopy
10376-2 | Oval macrocytes [Presence] in Blood by Light microscopy
10377-0 | Pencil cells [Presence] in Blood by Light microscopy
10378-8 | Polychromasia [Presence] in Blood by Light microscopy
10379-6 | Erythrocytes.dual population [Presence] in Blood by Light microscopy
1038-9 | G Ab [Presence] in Serum or Plasma
10380-4 | Stomatocytes [Presence] in Blood by Light microscopy
10381-2 | Target cells [Presence] in Blood by Light microscopy
10382-0 | A,B variant NOS Ag [Presence] on Red Blood Cells from Blood product unit
10383-8 | A,B variant NOS Ag [Presence] on Red Blood Cells from Donor
10384-6 | A,B variant NOS Ag [Presence] on Red Blood Cells
10385-3 | Albumin concentration given
10386-1 | Albumin given [Volume]
10387-9 | Autologous erythrocytes given [Volume]
10388-7 | Autologous whole blood given [Volume]
10389-5 | Blood product.other [Type]
1039-7 | G Ag [Presence] on Red Blood Cells from Blood product unit
10390-3 | Blood product special preparation [Type]
10391-1 | Cytomegalovirus immune globulin given [Volume]
10392-9 | Cryoprecipitate given [Volume]
10393-7 | Factor IX given [Type]
10394-5 | Factor IX given [Volume]
10395-2 | Factor VIII given [Type]
10396-0 | Factor VIII given [Volume]
10397-8 | Hepatitis B immune globulin given [Volume]
10398-6 | I Ag [Presence] on Red Blood Cells from Blood product unit
10399-4 | I Ag [Presence] on Red Blood Cells from Donor
104-0 | Ceforanide [Susceptibility] by Minimum inhibitory concentration (MIC)
1040-5 | G Ag [Presence] on Red Blood Cells from Donor
10400-0 | I Ag [Presence] on Red Blood Cells
10401-8 | Immune serum globulin given [Type]
10402-6 | Immune serum globulin given [Volume]
10403-4 | Inject immune serum globulin [Volume]
10404-2 | Inject Rh immune globulin [Volume]
10405-9 | Inject varicella zoster virus immune globulin [Volume]
10406-7 | little i Ag [Presence] on Red Blood Cells from Blood product unit
10407-5 | little i Ag [Presence] on Red Blood Cells from Donor
10408-3 | little i Ag [Presence] on Red Blood Cells
10409-1 | Pentaspan given [Volume]
1041-3 | G Ag [Presence] on Red Blood Cells
10410-9 | Plasma given [Type]
10411-7 | Plasma given [Volume]
10412-5 | Platelets given [Type]
10413-3 | Rh immune globulin given [Volume]
10414-1 | Transfuse albumin [Volume]
10415-8 | Transfuse blood exchange transfusion [Volume]

10416-6 | Transfuse blood product other [Volume]
10417-4 | Transfuse cryoprecipitate [Volume]
10418-2 | Transfuse factor IX [Volume]
10419-0 | Transfuse factor VIII [Volume]
1042-1 | H Ab [Presence] in Serum or Plasma from Blood product unit
10420-8 | Transfuse immune serum globulin [Volume]
10421-6 | Transfuse Pentaspan [Volume]
10422-4 | Transfuse plasma [Volume]
10423-2 | Transfuse platelets [Volume]
10424-0 | Transfuse erythrocytes [Volume]
10425-7 | Transfuse Rh immune globulin [Volume]
10426-5 | Transfuse whole blood [Volume]
10427-3 | Transfuse whole blood autologous [Volume]
10428-1 | Varicella zoster virus immune globulin given [Volume]
10429-9 | AE 1 Ag [Presence] in Tissue by Immune stain
1043-9 | H Ab [Presence] in Serum or Plasma from Donor
10430-7 | AE 3 Ag [Presence] in Tissue by Immune stain
10431-5 | B-cell Ag [Presence] in Tissue by Immune stain
10432-3 | CD30 Ag [Presence] in Tissue by Immune stain
10433-1 | BR-2 Ag [Presence] in Tissue by Immune stain
10434-9 | Complement C3 Ag [Presence] in Tissue by Immune stain
10435-6 | C5B-9 Ag [Presence] in Tissue by Immune stain
10436-4 | CD15 Ag [Presence] in Tissue by Immune stain
10437-2 | CD16 Ag [Presence] in Tissue by Immune stain
10438-0 | CD20 Ag [Presence] in Tissue by Immune stain
10439-8 | CD3 Ag [Presence] in Tissue by Immune stain
1044-7 | H Ab [Presence] in Serum
10440-6 | Deprecated CD30 Ag [Presence] in Tissue by Immune stain
10441-4 | CD34 Ag [Presence] in Tissue by Immune stain
10442-2 | CD56 Ag [Presence] in Tissue by Immune stain
10443-0 | CD43 T-cell monocyte+myeloid cell Ag [Presence] in Tissue by Immune stain
10444-8 | CD57 Ag [Presence] in Tissue by Immune stain
10445-5 | CD11c Ag [Presence] in Tissue by Immune stain
10446-3 | Leukocyte common Ag [Presence] in Tissue by Immune stain
10447-1 | M-5 Ag [Presence] in Tissue by Immune stain
10448-9 | T-cell Ag [Presence] in Tissue by Immune stain
10449-7 | Glucose [Mass/volume] in Serum or Plasma --1 hour post meal
1045-4 | H NOS Ab [Presence] in Serum or Plasma from Blood product unit
10450-5 | Glucose [Mass/volume] in Serum or Plasma --10 hours fasting
10451-3 | Proinsulin [Moles/volume] in Serum or Plasma --12 hours fasting
10452-1 | Xylose [Mass/volume] in Serum or Plasma --1 hour post 25 g xylose PO
10453-9 | Xylose [Mass/volume] in Serum or Plasma --1 hour post dose xylose PO
10454-7 | Xylose [Mass/volume] in Serum or Plasma --2 hours post 25 g xylose PO
10455-4 | Xylose [Mass/volume] in Serum or Plasma --30 minutes post 25 g xylose PO
10456-2 | Xylose [Mass/volume] in Serum or Plasma --6 hours fasting
10457-0 | Actin Ag [Presence] in Tissue by Immune stain
10458-8 | Alkaline phosphatase.placental Ag [Presence] in Tissue by Immune stain
10459-6 | Alpha-1-Fetoprotein Ag [Presence] in Tissue by Immune stain
1046-2 | H NOS Ab [Presence] in Serum or Plasma from Donor
10460-4 | Lactalbumin alpha Ag [Presence] in Tissue by Immune stain
10461-2 | Alpha-1-Antichymotrypsin Ag [Presence] in Tissue by Immune stain
10462-0 | Alpha 1 antitrypsin Ag [Presence] in Tissue by Immune stain
10463-8 | Amyloid A component Ag [Presence] in Tissue by Immune stain
10464-6 | Amyloid P component Ag [Presence] in Tissue by Immune stain
10465-3 | Amyloid.prealbumin Ag [Presence] in Tissue by Immune stain
10466-1 | Anion gap 3 in Serum or Plasma
10467-9 | Beta-2-Microglobulin amyloid Ag [Presence] in Tissue by Immune stain
10468-7 | Calcitonin Ag [Presence] in Tissue by Immune stain
10469-5 | Carcinoembryonic Ag [Presence] in Tissue by Immune stain
1047-0 | H NOS Ab [Presence] in Serum or Plasma
10470-3 | Choriogonadotropin Ag [Presence] in Tissue by Immune stain
10471-1 | Chromogranin A Ag [Presence] in Tissue by Immune stain
10472-9 | Chromogranin Ag [Presence] in Tissue by Immune stain
10473-7 | Chymotrypsin Ag [Presence] in Tissue by Immune stain
10474-5 | Collagen type 4 Ag [Presence] in Tissue by Immune stain
10475-2 | Corticotropin Ag [Presence] in Tissue by Immune stain
10476-0 | Desmin Ag [Presence] in Tissue by Immune stain
10477-8 | Enolase.neuron specific Ag [Presence] in Tissue by Immune stain
10478-6 | Eosinophil major basic protein Ag [Presence] in Tissue by Immune stain
10479-4 | CD227 Ag [Presence] in Tissue by Immune stain
1048-8 | H NOS Ag [Presence] on Red Blood Cells from Blood product unit
10480-2 | Estrogen+Progesterone receptor Ag [Presence] in Tissue by Immune stain

10481-0 | Follitropin.alpha subunit Ag [Presence] in Tissue by Immune stain
10482-8 | Follitropin.beta subunit Ag [Presence] in Tissue by Immune stain
10483-6 | Gastrin Ag [Presence] in Tissue by Immune stain
10484-4 | Glial fibrillary acidic protein Ag [Presence] in Tissue by Immune stain
10485-1 | Glucagon Ag [Presence] in Tissue by Immune stain
10486-9 | Hemoglobin Ag [Presence] in Tissue by Immune stain
10487-7 | HMB-45 Ag [Presence] in Tissue by Immune stain
10488-5 | IgA Ag [Presence] in Tissue by Immune stain
10489-3 | IgA.heavy chain Ag [Presence] in Tissue by Immune stain
1049-6 | H NOS Ag [Presence] on Red Blood Cells from Donor
10490-1 | IgE Ag [Presence] in Tissue by Immune stain
10491-9 | IgG Ag [Presence] in Tissue by Immune stain
10492-7 | IgG.heavy chain Ag [Presence] in Tissue by Immune stain
10493-5 | IgM Ag [Presence] in Tissue by Immune stain
10494-3 | IgM.heavy chain Ag [Presence] in Tissue by Immune stain
10495-0 | Insulin Ag [Presence] in Tissue by Immune stain
10496-8 | Kappa light chains Ag [Presence] in Tissue by Immune stain
10497-6 | Immunoglobulin light chains.kappa amyloid Ag [Presence] in Tissue by Immune stain
10498-4 | Keratin Ag [Presence] in Tissue by Immune stain
10499-2 | Lambda light chains Ag [Presence] in Tissue by Immune stain
105-7 | Ceforanide [Susceptibility] by Disk diffusion (KB)
1050-4 | H NOS Ag [Presence] on Red Blood Cells
10500-7 | Immunoglobulin light chains.lambda amyloid Ag [Presence] in Tissue by Immune stain
10501-5 | Lutropin [Units/volume] in Serum or Plasma
10502-3 | Lutropin Ag [Presence] in Tissue by Immune stain
10503-1 | Lysozyme Ag [Presence] in Tissue by Immune stain
10504-9 | Myelin basic protein Ag [Presence] in Tissue by Immune stain
10505-6 | Myoglobin Ag [Presence] in Tissue by Immune stain
10506-4 | Peanut agglutinin Ag [Presence] in Tissue by Immune stain
10507-2 | Prolactin Ag [Presence] in Tissue by Immune stain
10508-0 | Prostate specific Ag [Presence] in Tissue by Immune stain
10509-8 | Prostatic acid phosphatase Ag [Presence] in Tissue by Immune stain
1051-2 | Hemolytic disease of newborn screen [Interpretation] in Blood
10510-6 | S-100 Ag Ag [Presence] in Tissue by Immune stain
10511-4 | Serotonin Ag [Presence] in Tissue by Immune stain
10512-2 | Somatostatin Ag [Presence] in Tissue by Immune stain
10513-0 | Somatotropin Ag [Presence] in Tissue by Immune stain
10514-8 | Synaptophysin Ag [Presence] in Tissue by Immune stain
10515-5 | Thyroglobulin Ag [Presence] in Tissue by Immune stain
10516-3 | Thyrotropin Ag [Presence] in Tissue by Immune stain
10517-1 | Trypsin Ag [Presence] in Tissue by Immune stain
10518-9 | Ulex europaeus I lectin Ag [Presence] in Tissue by Immune stain
10519-7 | Vimentin Ag [Presence] in Tissue by Immune stain
1052-0 | Deprecated I (intermediate) subtype [Presence] in Serum or Plasma from Blood product unit
10520-5 | Coagulation factor VI Ag [Presence] in Tissue by Immune stain
10521-3 | Coagulation factor VIII Ag [Presence] in Tissue by Immune stain
10522-1 | Coagulation factor XIII Ag [Presence] in Tissue by Immune stain
10523-9 | Fibrinogen Ag [Presence] in Tissue by Immune stain
10524-7 | Microscopic observation [Identifier] in Cervix by Cyto stain
10525-4 | Microscopic observation [Identifier] in Unspecified specimen by Cyto stain
10526-2 | Microscopic observation [Identifier] in Sputum by Cyto stain
10527-0 | Microscopic observation [Identifier] in Tissue by Cyto stain
10528-8 | Acetophenazine [Mass/volume] in Serum or Plasma
10529-6 | Amoxapine metabolite [Mass/volume] in Serum or Plasma
1053-8 | Deprecated I (intermediate) subtype [Presence] in serum or plasma from donor
10530-4 | Aprobarbital [Mass/volume] in Serum or Plasma
10531-2 | Bretylium [Mass/volume] in Serum or Plasma
10532-0 | Deprecated Desethylamiodarone [Mass/volume] in Serum or Plasma
10533-8 | Propoxyphene+Acetaminophen [Mass/volume] in Serum or Plasma
10534-6 | Diazoxide [Mass/volume] in Serum or Plasma
10535-3 | Digoxin [Mass/volume] in Serum or Plasma
10536-1 | Dipyridamole [Mass/volume] in Serum or Plasma
10537-9 | Deprecated Doxepin+Nordoxepin [Mass/volume] in Serum or Plasma
10538-7 | Deprecated Fluoxetine+Norfluoxetine [Mass/volume] in Serum or Plasma
10539-5 | Glipizide [Mass/volume] in Serum or Plasma
1054-6 | Deprecated I (intermediate) subtype [Presence] in srum
10540-3 | Glyburide [Mass/volume] in Serum or Plasma
10541-1 | Mepivacaine [Mass/volume] in Serum or Plasma
10542-9 | Metharbital [Mass/volume] in Serum or Plasma
10543-7 | Methsuximide+Normethsuximide [Mass/volume] in Serum or Plasma
10544-5 | Normeperidine [Presence] in Serum or Plasma
10545-2 | Normephenytoin [Mass/volume] in Serum or Plasma

10546-0 | Normethsuximide [Mass/volume] in Serum or Plasma
10547-8 | Primidone+Phenobarbital [Mass/volume] in Serum or Plasma
10548-6 | Phenytoin Free/Phenytoin.total in Serum or Plasma
10549-4 | Pirmenol [Mass/volume] in Serum or Plasma
1055-3 | Deprecated I (intermediate) subtype [Presence] on Red Blood Cells from Blood product unit
10550-2 | Deprecated Temazepam [Mass/volume] in Serum or Plasma
10551-0 | Triamterene [Mass/volume] in Serum or Plasma
10552-8 | Tricyclic antidepressants [Mass/volume] in Serum or Plasma
10553-6 | Prostatic acid phosphatase [Enzymatic activity/volume] in Genital fluid
10554-4 | Deprecated Prostatic acid phosphatase [Enzymatic activity/volume] in Seminal plasma
10555-1 | Acrosin [Entitic Catalytic Activity] in Spermatozoa
10556-9 | Deprecated Adenosine triphosphatase [Enzymatic activity/volume] in Seminal plasma
10557-7 | Adenosine triphosphate [Moles/volume] in Semen
10558-5 | Albumin [Moles/volume] in Semen
10559-3 | Deprecated Calcium [Molecules/volume] in Semen
1056-1 | Deprecated I (intermediate) subtype [Presence] on Red Blood Cells from donor
10560-1 | Carcinoembryonic Ag [Units/volume] in Semen
10561-9 | Carnitine [Moles/volume] in Semen
10562-7 | Cells [#/volume] in Cervical mucus
10563-5 | Cells other than spermatozoa [#/volume] in Semen
10564-3 | Cervical mucus [Volume]
10565-0 | Choriogonadotropin [Units/volume] in Semen
10566-8 | Deprecated Choriogonadotropin [Molecules/volume] in Semen
10567-6 | Citrate [Moles/volume] in Semen
10568-4 | Clarity of Semen
10569-2 | Color of Semen
1057-9 | Deprecated I (intermediate) subtype [Presence] on Red Blood Cells
10570-0 | Consistency of Cervical mucus
10571-8 | Deprecated Consistency of Semen
10572-6 | Deprecated Duration^post ejaculation
10573-4 | Ferning [Type] in Cervical mucus
10574-2 | Fructose [Moles/volume] in Semen
10575-9 | Deprecated Gamma glutamyl transferase [Enzymatic activity/volume] in Semen
10576-7 | Germ cells.immature [#/volume] in Semen
10577-5 | Glucosidase [Enzymatic activity/volume] in Seminal plasma
10578-3 | Glycerophosphocholine [Moles/volume] in Seminal plasma
10579-1 | Leukocytes [#/volume] in Semen
1058-7 | I Ab [Presence] in Serum or Plasma from Blood product unit
10580-9 | Liquefaction [Time] in Semen
10581-7 | Deprecated Number of entities [#] in Spermatozoa
10582-5 | pH of Cervical mucus
10583-3 | Prostaglandin F1 alpha [Moles/volume] in Semen
10584-1 | Deprecated Protein [Mass/volume] in Semen
10585-8 | Round cells [#/volume] in Semen
10586-6 | Deprecated Volume of Semen
10587-4 | Sexual abstinence duration
10588-2 | Spermatogonia [#/volume] in Semen
10589-0 | Spermatids [#/volume] in Semen by Streptoavidin-biotin method (SAB)
1059-5 | I Ab [Presence] in Serum or Plasma from Donor
10590-8 | Spermatids [#/volume] in Semen by Sertoli cell barrier method (SCB)
10591-6 | Primary Spermatocytes [#/volume] in Semen
10592-4 | Secondary Spermatocytes [#/volume] in Semen
10593-2 | Spermatozoa Pyriform Head/100 spermatozoa in Semen
10594-0 | Spermatozoa Tapering Head/100 spermatozoa in Semen
10595-7 | Deprecated Spermatozoa [#/volume] in Semen
10596-5 | Deprecated Spermatozoa Ab in cervical mucosa
10597-3 | Deprecated Spermatozoa Ab in semen
10598-1 | Deprecated Spermatozoa Ab in serum
10599-9 | Spermatozoa penetration [Presence] in Seminal fluid and Cervical mucosa by Kremer test
106-5 | Ceforanide [Susceptibility] by Serum bactericidal titer
1060-3 | I Ab [Presence] in Serum or Plasma
10600-5 | Spermatozoa penetration [Presence] in Seminal fluid and Cervical mucosa
10601-3 | Spermatozoa penetration [Presence] in Seminal fluid and Cervical mucosa --post coitus
10602-1 | Spermatozoa Abnormal Head/100 spermatozoa in Semen
10603-9 | Spermatozoa Abnormal Midpiece/100 spermatozoa in Semen
10604-7 | Spermatozoa Abnormal Tail/100 spermatozoa in Semen
10605-4 | Spermatozoa Agglutinated/100 spermatozoa in Semen
10606-2 | Spermatozoa Amorphous Head/100 spermatozoa in Semen
10607-0 | Spermatozoa Coiled Tail/100 spermatozoa in Semen
10608-8 | Spermatozoa Cytoplasmic Droplet/100 spermatozoa in Semen
10609-6 | Spermatozoa Duplicate Head/100 spermatozoa in Semen
1061-1 | I NOS Ab [Presence] in Serum or Plasma from Blood product unit

10610-4 | Spermatozoa Duplicate Tail/100 spermatozoa in Semen
10611-2 | Spermatozoa Immotile/100 spermatozoa in Semen
10612-0 | Spermatozoa Large Oval Head/100 spermatozoa in Semen
10613-8 | Spermatozoa Viable/100 spermatozoa in Semen
10614-6 | Spermatozoa Motile w IgA/100 spermatozoa in Semen by Immunobead
10615-3 | Spermatozoa Motile w IgA/100 spermatozoa in Semen by Mixed antiglobulin reaction
10616-1 | Spermatozoa Motile w IgG/100 spermatozoa in Semen by Immunobead
10617-9 | Spermatozoa Motile w IgG/100 spermatozoa in Semen by Mixed antiglobulin reaction
10618-7 | Spermatozoa Motile w IgM/100 spermatozoa in Semen by Immunobead
10619-5 | Spermatozoa Motile w IgM/100 spermatozoa in Semen by Mixed antiglobulin reaction
1062-9 | I NOS Ab [Presence] in Serum or Plasma from Donor
10620-3 | Spermatozoa Nonprogressive/100 spermatozoa in Semen
10621-1 | Spermatozoa Normal Head/100 spermatozoa in Semen
10622-9 | Spermatozoa Normal/100 spermatozoa in Semen
10623-7 | Spermatozoa Pin Head/100 spermatozoa in Semen
10624-5 | Spermatozoa Rapid/100 spermatozoa in Semen
10625-2 | Spermatozoa Round Head/100 spermatozoa in Semen
10626-0 | Spermatozoa Slow/100 spermatozoa in Semen
10627-8 | Spermatozoa Small Oval Head/100 spermatozoa in Semen
10628-6 | Spermatozoa Tail Swelling/100 spermatozoa in Semen
10629-4 | Spermatozoa Vacuolated Head/100 spermatozoa in Semen
1063-7 | I NOS Ab [Presence] in Serum or Plasma
10630-2 | Spinnbarkeit [Length] in Cervical mucus
10631-0 | Testosterone [Moles/volume] in Semen
10632-8 | Time until next menstrual period
10633-6 | Zinc [Moles/volume] in Semen
10634-4 | Complement C1 esterase inhibitor.functional/Complement C1 esterase inhibitor.total in Serum or Plasma
10635-1 | Acanthamoeba sp identified in Eye by Organism specific culture
10636-9 | Acanthamoeba sp identified in Eye by Wet preparation
10637-7 | Microscopic observation [Identifier] in Tissue by Night blue stain
10638-5 | Thermophilic Actinomycetes colony count [#/volume] in Unspecified specimen by Organism specific culture
10639-3 | Thermophilic Actinomycetes identified in Unspecified specimen by Organism specific culture
1064-5 | I NOS Ag [Presence] on Red Blood Cells from Blood product unit
10640-1 | Deprecated Adenovirus 40+41 [Identifier] in Stool by Electron microscopy
10641-9 | Amoeba identified in Aspirate by Immune stain
10642-7 | Amoeba identified in Aspirate by Wet preparation
10643-5 | Amoeba identified in Stool by Organism specific culture
10644-3 | Arthropod identified in Unspecified specimen
10645-0 | Aspergillus fumigatus Ag [Presence] in Tissue by Immunofluorescence
10646-8 | Astrovirus [Identifier] in Stool by Electron microscopy
10647-6 | Babesia sp identified in Blood by Thick film
10648-4 | Babesia sp identified in Blood by Thin film
10649-2 | Calicivirus [Identifier] in Stool by Electron microscopy
1065-2 | I NOS Ag [Presence] on Red Blood Cells from Donor
10650-0 | Candida sp DNA [Presence] in Blood by Probe with amplification
10651-8 | Chlamydophila pneumoniae Ag [Presence] in Sputum or Bronchial
10652-6 | Chlamydophila pneumoniae DNA [Presence] in Sputum or Bronchial by Probe with amplification
10653-4 | Clotrimazole [Susceptibility] by Minimum inhibitory concentration (MIC)
10654-2 | Clotrimazole [Susceptibility] by Minimum lethal concentration (MLC)
10655-9 | Coccidia identified in Duodenal fluid by Acid fast stain
10656-7 | Coccidia identified in Stool by Acid fast stain
10657-5 | Deprecated Cryptococcus neoformans Ag [Presence] in Tissue by Immunofluorescence
10658-3 | Cyanobacterium identified in Water by Light microscopy
10659-1 | Cyclospora sp identified in Stool by Acid fast stain
1066-0 | I NOS Ag [Presence] on Red Blood Cells
10660-9 | Cytomegalovirus Ag [Presence] in Tissue by Immune stain
10661-7 | Dinoflagellate identified in Water by Light microscopy
10662-5 | Filaria identified in Blood by Concentration
10663-3 | Filaria identified in Blood by Thick film
10664-1 | Filaria identified in Blood by Thin film
10665-8 | Fungus colony count [#/volume] in Unspecified specimen by Environmental culture
10666-6 | Fungus identified in Tissue by Fontana-Masson stain
10667-4 | Fungus identified in Unspecified specimen by Animal inoculation
10668-2 | Fungus identified in Unspecified specimen by Environmental culture
10669-0 | Fungus identified in Unspecified specimen by Sticky tape for environmental fungus
1067-8 | Jk sup(a) Ab [Presence] in Serum or Plasma from Blood product unit
10670-8 | Giardia lamblia [Presence] in Stool by Organism specific culture
10671-6 | Helminth identified in Unspecified specimen
10672-4 | Helminth+Arthropod identified in Unspecified specimen
10673-2 | Hepatitis B virus core Ag [Presence] in Tissue by Immune stain
10674-0 | Hepatitis B virus surface Ag [Presence] in Tissue by Immune stain
10675-7 | Hepatitis B virus surface Ag [Presence] in Tissue by Orcein stain

10676-5 | Hepatitis C virus RNA [Units/volume] (viral load) in Serum or Plasma by Probe with amplification
10677-3 | Herpes simplex virus 1 Ag [Presence] in Tissue by Immune stain
10678-1 | Herpes simplex virus 1+2 Ag [Presence] in Tissue by Immune stain
10679-9 | Herpes simplex virus 2 Ag [Presence] in Tissue by Immune stain
1068-6 | Jk sup(a) Ab [Presence] in Serum or Plasma from Donor
10680-7 | Herpes simplex virus identified in Cerebral spinal fluid by Electron microscopy
10681-5 | Herpes simplex virus identified in Tissue by Electron microscopy
10682-3 | Deprecated HIV 1 RNA [Units/volume] in Serum or Plasma by Probe with amplification
10683-1 | Hydatid cyst identified in Aspirate by Immune stain
10684-9 | Hydatid cyst identified in Liver by Wet preparation
10685-6 | Hydatid cyst identified in Lung tissue by Wet preparation
10686-4 | Leishmania sp identified in Tissue by Giemsa stain
10687-2 | Leishmania sp identified in Tissue by Organism specific culture
10688-0 | Microscopic observation [Identifier] in Hair by KOH preparation
10689-8 | Microscopic observation [Identifier] in Nail by KOH preparation
1069-4 | Jk sup(a) Ab [Presence] in Serum or Plasma
10690-6 | Microsporidia identified in Duodenal fluid by Trichrome stain modified
10691-4 | Mushroom.toxic identified by Inspection
10692-2 | Mushroom.toxic identified in Food by Light microscopy
10693-0 | Mushroom.toxic identified in Vomitus by Light microscopy
10694-8 | Naegleria sp identified in Tissue by Organism specific culture
10695-5 | Naegleria sp identified in Tissue by Wet preparation
10696-3 | Norovirus [Identifier] in Stool by Electron microscopy
10697-1 | Nystatin [Susceptibility] by Minimum inhibitory concentration (MIC)
10698-9 | Nystatin [Susceptibility] by Minimum lethal concentration (MLC)
10699-7 | Onchocerca sp identified in Tissue by Wet preparation
107-3 | Cefotaxime [Susceptibility] by Minimum lethal concentration (MLC)
1070-2 | Jk sup(a) Ag [Presence] on Red Blood Cells from Blood product unit
10700-3 | Orthopoxvirus [Identifier] in Skin by Electron microscopy
10701-1 | Ova and parasites identified in Stool by Concentration
10702-9 | Ova and parasites identified in Stool by Immune stain
10703-7 | Ova and parasites identified in Stool by Kinyoun iron hematoxylin stain
10704-5 | Ova and parasites identified in Stool by Light microscopy
10705-2 | Human papilloma virus Ag [Presence] in Tissue by Immune stain
10706-0 | Picornavirus [Identifier] in Stool by Electron microscopy
10707-8 | Toxic Plant identified in Plant specimen by Inspection
10708-6 | Toxic Plant identified in Vomitus by Light microscopy
10709-4 | Plasmodium falciparum Ag [Units/volume] in Blood by Immunofluorescence
1071-0 | Jk sup(a) Ag [Presence] on Red Blood Cells from Donor
10710-2 | Plasmodium sp identified in Blood by Thin film
10711-0 | Plasmodium vivax Ag [Units/volume] in Blood by Immunofluorescence
10712-8 | Pneumocystis sp identified in Lung tissue
10713-6 | Prototheca identified in Unspecified specimen by Culture
10714-4 | Deprecated Rotavirus [Identifier] in Stool by Electron microscopy
10715-1 | Schistosoma sp identified in Urine sediment by Light microscopy
10716-9 | Schistosoma sp identified in Unspecified specimen by Organism specific culture
10717-7 | Streptococcus pneumoniae Ab [Units/volume] in Serum by Latex agglutination
10718-5 | Strongyloides sp Ab [Units/volume] in Serum by Immunoassay
10719-3 | Taenia solium adult Ab [Units/volume] in Serum by Immunoblot
1072-8 | Jk sup(a) Ag [Presence] on Red Blood Cells
10720-1 | Terbinafine [Susceptibility] by Minimum inhibitory concentration (MIC)
10721-9 | Terbinafine [Susceptibility] by Minimum lethal concentration (MLC)
10722-7 | Torovirus [Identifier] in Stool by Electron microscopy
10723-5 | Toxoplasma gondii IgA Ab [Units/volume] in Serum by Immunoassay
10724-3 | Toxoplasma gondii IgE Ab [Units/volume] in Serum by Immunoassay
10725-0 | Toxoplasma gondii DNA [Units/volume] in Body fluid by Probe with amplification
10726-8 | Toxoplasma gondii [Presence] in Tissue by Giemsa stain
10727-6 | Toxoplasma gondii identified in Tissue
10728-4 | Trichomonas sp identified in Genital specimen by Organism specific culture
10729-2 | Trypanosoma sp identified in Blood by Light microscopy
1073-6 | Jk sup(b) Ab [Presence] in Serum or Plasma from Blood product unit
10730-0 | Trypanosoma sp identified in Blood by Organism specific culture
10731-8 | Trypanosoma sp identified in Blood by Thick film
10732-6 | Trypanosoma sp identified in Blood by Thin film
10733-4 | Trypanosoma sp identified in Blood by Wet preparation
10734-2 | Varicella zoster virus identified in Skin by Electron microscopy
10735-9 | Viral sequencing [Identifier] in Serum by Sequencing
10736-7 | Virus identified in Cerebral spinal fluid by Electron microscopy
10737-5 | Virus identified in Stool by Electron microscopy
10738-3 | Virus identified in Tissue by Electron microscopy
10739-1 | Virus identified in Unspecified specimen by Electron microscopy
1074-4 | Jk sup(b) Ab [Presence] in Serum or Plasma from Donor

10740-9 | Aluminum.microscopic observation [Identifier] in Bone by Histomorphometry stain
10741-7 | Amyloid.microscopic observation [Identifier] in Brain by Thioflavine-S stain
10742-5 | Amyloid.microscopic observation [Identifier] in Tissue by Bennhold stain.Putchler modified
10743-3 | Amyloid.microscopic observation [Identifier] in Tissue by Highman stain
10744-1 | Amyloid.microscopic observation [Identifier] in Tissue by Vassar-culling stain
10745-8 | Bile.microscopic observation [Identifier] in Tissue by Fouchet stain
10746-6 | Calcium.microscopic observation [Identifier] in Tissue by Von Kossa stain
10747-4 | Collagen fibers+Elastic fibers.microscopic observation [Identifier] in Tissue by Lawson-Van Gieson stain
10748-2 | Collagen fibers+Elastic fibers.microscopic observation [Identifier] in Tissue by Verhoeff-Van Gieson stain
10749-0 | Collagen fibers.microscopic observation [Identifier] in Tissue by Van Gieson stain
1075-1 | Jk sup(b) Ab [Presence] in Serum or Plasma
10750-8 | Connective tissue.microscopic observation [Identifier] in Tissue by Trichrome stain.Masson
10751-6 | Copper.microscopic observation [Identifier] in Tissue by Rhodamine stain
10752-4 | Fat.microscopic observation [Identifier] in Milk by Sudan IV stain
10753-2 | Fat.microscopic observation [Identifier] in Stool by Sudan IV stain
10754-0 | Fat.microscopic observation [Identifier] in Tissue by Sudan IV stain
10755-7 | Fungus.microscopic observation [Identifier] in Tissue by Methenamine silver stain.Grocott
10756-5 | Glial fibers.microscopic observation [Identifier] in Tissue by Holzer stain
10757-3 | Hematologic+Nuclear elements.microscopic observation [Identifier] in Tissue by Giemsa stain.May-Grunwald
10758-1 | Iron.microscopic observation [Identifier] in Bone by Histomorphometry stain
10759-9 | Iron.microscopic observation [Identifier] in Sputum by Gomori stain
1076-9 | Jk sup(b) Ag [Presence] on Red Blood Cells from Blood product unit
10760-7 | Iron.microscopic observation [Identifier] in Tissue by Gomori stain
10761-5 | Iron.microscopic observation [Identifier] in Tissue by Other stain
10762-3 | Microscopic observation [Identifier] in Blood by Hemosiderin stain
10763-1 | Microscopic observation [Identifier] in Body fluid by Sudan black stain
10764-9 | Microscopic observation [Identifier] in Sputum by Silver stain
10765-6 | Microscopic observation [Identifier] in Tissue by Acetate esterase stain
10766-4 | Microscopic observation [Identifier] in Tissue by Alcian blue stain
10767-2 | Microscopic observation [Identifier] in Tissue by Alcian blue stain.sulfated
10768-0 | Microscopic observation [Identifier] in Tissue by Alcian blue stain.with periodic acid-Schiff
10769-8 | Microscopic observation [Identifier] in Tissue by Alizarin red S stain
1077-7 | Jk sup(b) Ag [Presence] on Red Blood Cells from Donor
10770-6 | Microscopic observation [Identifier] in Tissue by Argentaffin stain
10771-4 | Deprecated Microscopic observation [Identifier] in Tissue by Rhodamine-auramine fluorochrome stain
10772-2 | Microscopic observation [Identifier] in Tissue by Azure-eosin stain
10773-0 | Microscopic observation [Identifier] in Tissue by Basic fuchsin stain
10774-8 | Microscopic observation [Identifier] in Tissue by Bielschowsky stain
10775-5 | Microscopic observation [Identifier] in Tissue by Bleach stain
10776-3 | Microscopic observation [Identifier] in Tissue by Bodian stain
10777-1 | Microscopic observation [Identifier] in Tissue by Brown and Brenn stain
10778-9 | Microscopic observation [Identifier] in Tissue by Butyrate esterase stain
10779-7 | Microscopic observation [Identifier] in Tissue by Carmine stain.Best
1078-5 | Jk sup(b) Ag [Presence] on Red Blood Cells
10780-5 | Microscopic observation [Identifier] in Tissue by Chloracetate esterase stain
10781-3 | Microscopic observation [Identifier] in Tissue by Churukian-Schenk stain
10782-1 | Microscopic observation [Identifier] in Tissue by Congo red stain
10783-9 | Microscopic observation [Identifier] in Tissue by Crystal violet stain
10784-7 | Microscopic observation [Identifier] in Tissue by Esterase stain.non-specific
10785-4 | Microscopic observation [Identifier] in Tissue by Fite-Faraco stain
10786-2 | Deprecated Microscopic observation [Identifier] in Tissue by Giemsa stain.3 micron
10787-0 | Microscopic observation [Identifier] in Tissue by Gridley stain
10788-8 | Microscopic observation [Identifier] in Tissue by Hansel stain
10789-6 | Microscopic observation [Identifier] in Tissue by Hematoxylin-eosin-Harris regressive stain
1079-3 | Js sup(a) Ab [Presence] in Serum or Plasma from Blood product unit
10790-4 | Microscopic observation [Identifier] in Tissue by Hematoxylin-eosin-Mayers progressive stain
10791-2 | Microscopic observation [Identifier] in Tissue by Mallory-Heidenhain stain
10792-0 | Microscopic observation [Identifier] in Tissue by Methenamine silver stain.Jones
10793-8 | Microscopic observation [Identifier] in Tissue by Methyl green stain
10794-6 | Microscopic observation [Identifier] in Tissue by Methyl green-pyronine Y stain
10795-3 | Microscopic observation [Identifier] in Tissue by Methyl violet stain
10796-1 | Microscopic observation [Identifier] in Tissue by Mucicarmine stain
10797-9 | Microscopic observation [Identifier] in Tissue by Neutral red stain
10798-7 | Microscopic observation [Identifier] in Tissue by Oil red O stain
10799-5 | Microscopic observation [Identifier] in Tissue by Pentachrome stain.Movat
108-1 | Cefotaxime [Susceptibility] by Minimum inhibitory concentration (MIC)
1080-1 | Js sup(a) Ab [Presence] in Serum or Plasma from Donor
10800-1 | Microscopic observation [Identifier] in Tissue by Periodic acid-Schiff stain with diatase digestion
10801-9 | Microscopic observation [Identifier] in Tissue by Phosphotungstic acid Hematoxylin (PTAH) Stain
10802-7 | Microscopic observation [Identifier] in Tissue by Prussian blue stain
10803-5 | Microscopic observation [Identifier] in Tissue by Quinacrine fluorescent stain
10804-3 | Microscopic observation [Identifier] in Tissue by Reticulin stain

10805-0 | Microscopic observation [Identifier] in Tissue by Safranin stain
10806-8 | Microscopic observation [Identifier] in Tissue by Schmorl stain
10807-6 | Microscopic observation [Identifier] in Tissue by Sevier-Munger stain
10808-4 | Microscopic observation [Identifier] in Tissue by Silver impregnation stain.Dieterle
10809-2 | Microscopic observation [Identifier] in Tissue by Silver nitrate stain
1081-9 | Js sup(a) Ab [Presence] in Serum or Plasma
10810-0 | Microscopic observation [Identifier] in Tissue by Silver stain.Fontana-Masson
10811-8 | Microscopic observation [Identifier] in Tissue by Silver stain.Grimelius
10812-6 | Microscopic observation [Identifier] in Tissue by Steiner stain
10813-4 | Microscopic observation [Identifier] in Tissue by Sudan black stain
10814-2 | Microscopic observation [Identifier] in Tissue by Supravital stain
10815-9 | Microscopic observation [Identifier] in Tissue by Tetrachrome stain
10816-7 | Deprecated Microscopic observation [Identifier] in Tissue by Toluidine blue O stain
10817-5 | Microscopic observation [Identifier] in Tissue by Trichrome stain.Gomori-Wheatley
10818-3 | Microscopic observation [Identifier] in Tissue by Trichrome stain.Masson modified
10819-1 | Microscopic observation [Identifier] in Tissue by Wade stain
1082-7 | Js sup(a) Ag [Presence] on Red Blood Cells from Blood product unit
10820-9 | Deprecated Microscopic observation [Identifier] in Tissue by Warthin-Starry stain
10821-7 | Deprecated Microscopic observation [Identifier] in Tissue by Wright Giemsa stain
10822-5 | Mucin.microscopic observation [Identifier] in Tissue by Mucicarmine stain.Mayer
10823-3 | Mucopolysaccharides.microscopic observation [Identifier] in Tissue by Colloidal ferric oxide stain.Hale
10824-1 | Myelin+Myelin breakdown products.microscopic observation [Identifier] in Tissue by Luxol fast blue/Periodic acid-Schiff stain
10825-8 | Myelin+Nerve cells.microscopic observation [Identifier] in Tissue by Luxol fast blue/Cresyl violet stain
10826-6 | Nissel.microscopic observation [Identifier] in Tissue by Cresyl echt violet stain
10827-4 | Reticulum.microscopic observation [Identifier] in Tissue by Gomori stain
10828-2 | Urate crystals.microscopic observation [type] in Tissue by De Galantha stain
10829-0 | Silicon [Mass/volume] in Serum or Plasma
1083-5 | Js sup(a) Ag [Presence] on Red Blood Cells from Donor
10830-8 | Deprecated Surgical operation note complications
10831-6 | Surgical operation note complications [Interpretation]
10832-4 | Glucose [Mass/volume] in Serum or Plasma --15 minutes post 50 g lactose PO
10833-2 | Insulin [Mass/volume] in Serum or Plasma --7 hours post 75 g glucose PO
10834-0 | Globulin [Mass/volume] in Serum by calculation
10835-7 | Lipoprotein a [Mass/volume] in Serum or Plasma
10836-5 | Niacin [Mass/volume] in Blood
10837-3 | Organic acids [Presence] in Serum or Plasma
10838-1 | Phosphoserine [Mass/volume] in Serum or Plasma
10839-9 | Troponin I.cardiac [Mass/volume] in Serum or Plasma
1084-3 | Js sup(a) Ag [Presence] on Red Blood Cells
10840-7 | Atropine [Mass/volume] in Urine
10841-5 | Methazolamide [Mass/volume] in Serum or Plasma
10842-3 | Deprecated HLA-DQ1 [Presence]
10843-1 | Deprecated HLA-DQ2 [Presence]
10844-9 | Deprecated HLA-DQ3 [Presence]
10845-6 | Deprecated HLA-DQ4 [Presence]
10846-4 | Borrelia burgdorferi DNA [Presence] in Blood by Probe with amplification
10847-2 | Borrelia burgdorferi DNA [Presence] in Body fluid by Probe with amplification
10848-0 | Chlamydia sp IgG Ab [Titer] in Serum by Immunofluorescence
10849-8 | Chlamydia sp IgM Ab [Titer] in Serum by Immunofluorescence
1085-0 | Js sup(b) Ab [Presence] in Serum or Plasma from Blood product unit
10850-6 | Cyclospora cayetanensis [Presence] in Unspecified specimen
10851-4 | Escherichia coli O157:H7 [Presence] in Stool by Organism specific culture
10852-2 | Deprecated Fungus identified in Blood by Culture
10853-0 | Isospora belli [Presence] in Unspecified specimen by Acid fast stain.Kinyoun modified
10854-8 | Deprecated Midrofilaria identified in Blood by Concentration
10855-5 | Ova and parasites identified in Duodenal fluid or Gastric fluid by Light microscopy
10856-3 | Microscopic observation [Identifier] in Genital mucus by Gram stain
10857-1 | Microsporidia identified in Unspecified specimen by Light microscopy
10858-9 | Teichoate Ab [Presence] in Serum
10859-7 | Trypanosoma sp [Identifier] in Blood by Acridine Orange + Giemsa Stain
1086-8 | Js sup(b) Ab [Presence] in Serum or Plasma from Donor
10860-5 | Varicella zoster virus [Presence] in Unspecified specimen by Organism specific culture
10861-3 | Progesterone receptor [Mass/mass] in Tissue
10862-1 | Basement membrane Ab [Titer] in Serum
10863-9 | Endomysium IgA Ab [Titer] in Serum
10864-7 | Immune complex [Units/volume] in Serum or Plasma by Raji cell assay
10865-4 | Intercellular substance Ab [Presence] in Serum
10866-2 | Sulfatide IgG Ab [Titer] in Serum
10867-0 | Sulfatide IgM Ab [Titer] in Serum
10868-8 | Bacitracin [Susceptibility] by Disk diffusion (KB)
10869-6 | Xylose [Mass/volume] in Blood --3 hours post 25 g xylose PO

1087-6 | Js sup(b) Ab [Presence] in Serum or Plasma
10870-4 | Xylose [Mass/volume] in Blood --4 hours post 25 g xylose PO
10871-2 | Xylose [Mass/volume] in Blood --5 hours post 25 g xylose PO
10872-0 | Xylose [Mass/volume] in Blood --baseline
10873-8 | Beta-2-Microglobulin [Mass/time] in 24 hour Urine
10874-6 | Bombesin [Mass/volume] in Plasma
10875-3 | Carnitine esters [Mass/volume] in Urine
10876-1 | Carnitine esters [Mass/volume] in Serum or Plasma
10877-9 | Carnitine free (C0) [Mass/volume] in Urine
10878-7 | Iodine.protein bound [Mass/volume] in Serum or Plasma
10879-5 | Isovalerylglycine [Mass/volume] in Urine
1088-4 | Js sup(b) Ag [Presence] on Red Blood Cells from Blood product unit
10880-3 | Magnesium [Mass/mass] in Stool
10881-1 | Pentacarboxylporphyrins [Mass/volume] in Red Blood Cells
10882-9 | Pentacarboxylporphyrins [Mass/volume] in Serum or Plasma
10883-7 | Phenolphthalein [Mass/mass] in Stool
10884-5 | Phosphate [Mass/mass] in Stool
10885-2 | Porphyrins [Mass/time] in 24 hour Urine
10886-0 | Prostate Specific Ag Free [Mass/volume] in Serum or Plasma
10887-8 | Pyridinoline [Mass/time] in 24 hour Urine
10888-6 | Lipase [Enzymatic activity/volume] in Urine
10889-4 | Bisacodyl [Mass/mass] in Stool
1089-2 | Js sup(b) Ag [Presence] on Red Blood Cells from Donor
10890-2 | Deprecated Normethsuximide [Mass/volume] in Serum or Plasma
10891-0 | Oxyphenisatin [Mass/mass] in Stool
10892-8 | Pentoxifylline [Mass/volume] in Serum or Plasma
10893-6 | Trenbolone [Mass/volume] in Urine
10894-4 | Aspergillus niger Ab [Presence] in Serum by Immune diffusion (ID)
10895-1 | Clostridium difficile toxin B [Presence] in Stool
10896-9 | Eastern equine encephalitis virus IgG Ab [Titer] in Serum by Immunofluorescence
10897-7 | Eastern equine encephalitis virus IgG Ab [Titer] in Cerebral spinal fluid by Immunofluorescence
10898-5 | Eastern equine encephalitis virus IgM Ab [Titer] in Serum by Immunofluorescence

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.522 Value Set http://hl7.org/fhir/ValueSet/research-subject-status

URL: https://hl7.org/fhir/R4/valueset-research-subject-status.html
Biomedical Research and Regulation Work Group | Maturity Level: 1 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/research-subject-status
Version: | 4.0.1
Name: | ResearchSubjectStatus
Title: | ResearchSubjectStatus
Definition: | Indicates the progression of a study subject through a study.
Committee: | Biomedical Research and Regulation Work Group
OID: | 2.16.840.1.113883.4.642.3.829 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ResearchSubjectStatus
• ConceptMap: Translation source in Canonical Mapping for "ResearchSubjectStatus"
• Resource: ResearchSubject.status (code / Required)

4.4.1.522.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/research-subject-status

4.4.1.522.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 13 concepts

Expansion based on http://hl7.org/fhir/research-subject-status version 4.0.1

All codes from system http://hl7.org/fhir/research-subject-status

Code | Display | Definition | Canonical Status
candidate | Candidate | An identified person that can be considered for inclusion in a study.
eligible | Eligible | A person that has met the eligibility criteria for inclusion in a study.
follow-up | Follow-up | A person is no longer receiving study intervention and/or being evaluated with tests and procedures according to the protocol, but they are being monitored on a protocol-prescribed schedule. | ~replaced
ineligible | Ineligible | A person who did not meet one or more criteria required for participation in a study is considered to have failed screening or is ineligible for the study. | ~declined
not-registered | Not Registered | A person for whom registration was not completed. | ~abandoned
off-study | Off-study | A person that has ended their participation on a study either because their treatment/observation is complete or through not responding, withdrawal, non-compliance and/or adverse event. | ~complete
on-study | On-study | A person that is enrolled or registered on a study.
on-study-intervention | On-study-intervention | The person is receiving the treatment or participating in an activity (e.g. yoga, diet, etc.) that the study is evaluating.
on-study-observation | On-study-observation | The subject is being evaluated via tests and assessments according to the study calendar, but is not receiving any intervention. Note that this state is study-dependent and might not exist in all studies. A synonym for this is "short-term follow-up".
pending-on-study | Pending on-study | A person is pre-registered for a study. | ~suspended
potential-candidate | Potential Candidate | A person that is potentially eligible for participation in the study.
screening | Screening | A person who is being evaluated for eligibility for a study.
withdrawn | Withdrawn | The person has withdrawn their participation in the study before registration. | ~failed

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.523 Value Set http://hl7.org/fhir/ValueSet/explanationofbenefit-status

URL: https://hl7.org/fhir/R4/valueset-explanationofbenefit-status.html
Financial Management Work Group | Maturity Level: 2 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/explanationofbenefit-status
Version: | 4.0.1
Name: | ExplanationOfBenefitStatus
Title: | ExplanationOfBenefitStatus
Definition: | A code specifying the state of the resource instance.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.618 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

- CodeSystem: This value set is the designated 'entire code system' value set for ExplanationOfBenefitStatus
- ConceptMap: Translation source in Canonical Mapping for "ExplanationOfBenefitStatus"
- Resource: ExplanationOfBenefit.status (code / Required)

4.4.1.523.1 Content Logical Definition

This value set includes codes from the following code systems:

- Include all codes defined in http://hl7.org/fhir/explanationofbenefit-status

4.4.1.523.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 4 concepts

Expansion based on http://hl7.org/fhir/explanationofbenefit-status version 4.0.1

All codes from system http://hl7.org/fhir/explanationofbenefit-status

Code | Display | Definition | Canonical Status
active | Active | The resource instance is currently in-force. | ~active
cancelled | Cancelled | The resource instance is withdrawn, rescinded or reversed. | ~abandoned
draft | Draft | A new resource instance the contents of which is not complete. | ~draft
entered-in-error | Entered In Error | The resource instance was entered in error. | ~error

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.524 Value Set http://hl7.org/fhir/ValueSet/audit-event-sub-type

URL: https://hl7.org/fhir/R4/valueset-audit-event-sub-type.html
Security Work Group | Maturity Level: 3 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/audit-event-sub-type
Version: | 4.0.1
Name: | AuditEventSub-Type
Title: | Audit Event Sub-Type
Definition: | More detailed code concerning the type of the audit event - defined by DICOM with some FHIR specific additions.
Committee: | Security Work Group
OID: | 2.16.840.1.113883.4.642.3.464 (for OID based terminology systems)
Copyright: | These codes are excerpted from Digital Imaging and Communications in Medicine (DICOM) Standard, Part 16: Content Mapping Resource, Copyright © 2011 by the National Electrical Manufacturers Association.
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: AuditEvent.subtype (Coding / Extensible)

4.4.1.524.1 Content Logical Definition

This value set includes codes from the following code systems:

http://dicom.nema.org/resources/ontology/DCM

Code | Display
110120 | Application Start | Audit event: Application Entity has started
110121 | Application Stop | Audit event: Application Entity has stopped
110122 | Login | Audit event: User login has been attempted
110123 | Logout | Audit event: User logout has been attempted
110124 | Attach | Audit event: Node has been attached
110125 | Detach | Audit event: Node has been detached
110126 | Node Authentication | Audit event: Node Authentication has been attempted
110127 | Emergency Override Started | Audit event: Emergency Override has started
110128 | Network Configuration | Audit event: Network configuration has been changed
110129 | Security Configuration | Audit event: Security configuration has been changed
110130 | Hardware Configuration | Audit event: Hardware configuration has been changed
110131 | Software Configuration | Audit event: Software configuration has been changed
110132 | Use of Restricted Function | Audit event: A use of a restricted function has been attempted
110133 | Audit Recording Stopped | Audit event: Audit recording has been stopped
110134 | Audit Recording Started | Audit event: Audit recording has been started
110135 | Object Security Attributes Changed | Audit event: Security attributes of an object have been changed
110136 | Security Roles Changed | Audit event: Security roles have been changed
110137 | User security Attributes Changed | Audit event: Security attributes of a user have been changed
110138 | Emergency Override Stopped | Audit event: Emergency Override has Stopped
110139 | Remote Service Operation Started | Audit event: Remote Service Operation has Begun
110140 | Remote Service Operation Stopped | Audit event: Remote Service Operation has Stopped
110141 | Local Service Operation Started | Audit event: Local Service Operation has Begun
110142 | Local Service Operation Stopped | Audit event: Local Service Operation Stopped

• Include all codes defined in http://hl7.org/fhir/restful-interaction

4.4.1.524.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 40 concepts

Expansion based on DICOM version 01, http://hl7.org/fhir/restful-interaction version 4.0.1

Code | System | Display | Definition
110120 | http://dicom.nema.org/resources/ontology/DCM | Application Start | Audit event: Application Entity has started
110121 | http://dicom.nema.org/resources/ontology/DCM | Application Stop | Audit event: Application Entity has stopped
110122 | http://dicom.nema.org/resources/ontology/DCM | Login | Audit event: User login has been attempted
110123 | http://dicom.nema.org/resources/ontology/DCM | Logout | Audit event: User logout has been attempted
110124 | http://dicom.nema.org/resources/ontology/DCM | Attach | Audit event: Node has been attached
110125 | http://dicom.nema.org/resources/ontology/DCM | Detach | Audit event: Node has been detached
110126 | http://dicom.nema.org/resources/ontology/DCM | Node Authentication | Audit event: Node Authentication has been attempted
110127 | http://dicom.nema.org/resources/ontology/DCM | Emergency Override Started | Audit event: Emergency Override has started
110128 | http://dicom.nema.org/resources/ontology/DCM | Network Configuration | Audit event: Network configuration has been changed
110129 | http://dicom.nema.org/resources/ontology/DCM | Security Configuration | Audit event: Security configuration has been changed
110130 | http://dicom.nema.org/resources/ontology/DCM | Hardware Configuration | Audit event: Hardware configuration has been changed
110131 | http://dicom.nema.org/resources/ontology/DCM | Software Configuration | Audit event: Software configuration has been changed
110132 | http://dicom.nema.org/resources/ontology/DCM | Use of Restricted Function | Audit event: A use of a restricted function has been attempted
110133 | http://dicom.nema.org/resources/ontology/DCM | Audit Recording Stopped | Audit event: Audit recording has been stopped
110134 | http://dicom.nema.org/resources/ontology/DCM | Audit Recording Started | Audit event: Audit recording has been started
110135 | http://dicom.nema.org/resources/ontology/DCM | Object Security Attributes Changed | Audit event: Security attributes of an object have been changed
110136 | http://dicom.nema.org/resources/ontology/DCM | Security Roles Changed | Audit event: Security roles have been changed
110137 | http://dicom.nema.org/resources/ontology/DCM | User security Attributes Changed | Audit event: Security attributes of a user have been changed
110138 | http://dicom.nema.org/resources/ontology/DCM | Emergency Override Stopped | Audit event: Emergency Override has Stopped
110139 | http://dicom.nema.org/resources/ontology/DCM | Remote Service Operation Started | Audit event: Remote Service Operation has Begun
110140 | http://dicom.nema.org/resources/ontology/DCM | Remote Service Operation Stopped | Audit event: Remote Service Operation has Stopped
110141 | http://dicom.nema.org/resources/ontology/DCM | Local Service Operation Started | Audit event: Local Service Operation has Begun
110142 | http://dicom.nema.org/resources/ontology/DCM | Local Service Operation Stopped | Audit event: Local Service Operation Stopped
read | http://hl7.org/fhir/restful-interaction | read | Read the current state of the resource.
vread | http://hl7.org/fhir/restful-interaction | vread | Read the state of a specific version of the resource.
update | http://hl7.org/fhir/restful-interaction | update | Update an existing resource by its id (or create it if it is new).
patch | http://hl7.org/fhir/restful-interaction | patch | Update an existing resource by posting a set of changes to it.
delete | http://hl7.org/fhir/restful-interaction | delete | Delete a resource.
history | http://hl7.org/fhir/restful-interaction | history | Retrieve the change history for a particular resource, type of resource, or the entire system.
history-instance | http://hl7.org/fhir/restful-interaction | history-instance | Retrieve the change history for a particular resource.
history-type | http://hl7.org/fhir/restful-interaction | history-type | Retrieve the change history for all resources of a particular type.
history-system | http://hl7.org/fhir/restful-interaction | history-system | Retrieve the change history for all resources on a system.
create | http://hl7.org/fhir/restful-interaction | create | Create a new resource with a server assigned id.
search | http://hl7.org/fhir/restful-interaction | search | Search a resource type or all resources based on some filter criteria.
search-type | http://hl7.org/fhir/restful-interaction | search-type | Search all resources of the specified type based on some filter criteria.
search-system | http://hl7.org/fhir/restful-interaction | search-system | Search all resources based on some filter criteria.
capabilities | http://hl7.org/fhir/restful-interaction | capabilities | Get a Capability Statement for the system.
transaction | http://hl7.org/fhir/restful-interaction | transaction | Update, create or delete a set of resources as a single transaction.
batch | http://hl7.org/fhir/restful-interaction | batch | perform a set of a separate interactions in a single http operation
operation | http://hl7.org/fhir/restful-interaction | operation | Perform an operation as defined by an OperationDefinition.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)

Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.525 Value Set http://hl7.org/fhir/ValueSet/namingsystem-identifier-type

URL: https://hl7.org/fhir/R4/valueset-namingsystem-identifier-type.html
FHIR Infrastructure Work Group | Maturity Level: 1 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/namingsystem-identifier-type
Version: | 4.0.1
Name: | NamingSystemIdentifierType
Title: | NamingSystemIdentifierType
Definition: | Identifies the style of unique identifier used to identify a namespace.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.492 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for NamingSystemIdentifierType

• Resource: NamingSystem.uniqueId.type (code / Required)

• Operation: Parameter $preferred-id.type (code /: (Required))

4.4.1.525.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/namingsystem-identifier-type

4.4.1.525.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 4 concepts

Expansion based on http://hl7.org/fhir/namingsystem-identifier-type version 4.0.1

All codes from system http://hl7.org/fhir/namingsystem-identifier-type

Code | Display | Definition
oid | OID | An ISO object identifier; e.g. 1.2.3.4.5.
uuid | UUID | A universally unique identifier of the form a5afddf4-e880-459b-876e-e4591b0acc11.
uri | URI | A uniform resource identifier (ideally a URL - uniform resource locator); e.g. http://unitsofmeasure.org.
other | Other | Some other type of unique identifier; e.g. HL7-assigned reserved string such as LN for LOINC.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.526 Value Set http://hl7.org/fhir/ValueSet/contract-data-meaning

URL: https://hl7.org/fhir/R4/valueset-contract-data-meaning.html
Financial Management Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/contract-data-meaning
Version: | 4.0.1

Name: | ContractDataMeaning
Title: | ContractDataMeaning
Definition: | How a resource reference is interpreted when evaluating contract offers.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.749 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ContractDataMeaning

4.4.1.526.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/contract-data-meaning

4.4.1.526.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 4 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/contract-data-meaning version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/contract-data-meaning

Code | Display | Definition
instance | Instance | The consent applies directly to the instance of the resource.
related | Related | The consent applies directly to the instance of the resource and instances it refers to.
dependents | Dependents | The consent applies directly to the instance of the resource and instances that refer to it.
authoredby | AuthoredBy | The consent applies to instances of resources that are authored by.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.527 Value Set http://hl7.org/fhir/ValueSet/verificationresult-need

URL: https://hl7.org/fhir/R4/valueset-verificationresult-need.html
Patient Administration Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/verificationresult-need
Version: | 4.0.1
Name: | need
Title: | Need
Definition: | The frequency with which the target must be validated
Committee: | Patient Administration Work Group
OID: | 2.16.840.1.113883.4.642.3.882 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for need

• Resource: VerificationResult.need (CodeableConcept / Preferred)

4.4.1.527.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/need

4.4.1.527.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 3 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/need version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/need

Code | Display | Definition
none | None | ***TODO***
initial | Initial | ***TODO***
periodic | Periodic | ***TODO***

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.528 Value Set http://hl7.org/fhir/ValueSet/allerg-intol-substance-exp-risk

URL: https://hl7.org/fhir/R4/valueset-allerg-intol-substance-exp-risk.html
Patient Care Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/allerg-intol-substance-exp-risk
Version: | 4.0.1
Name: | AllergyIntoleranceSubstanceExposureRisk
Title: | AllergyIntoleranceSubstanceExposureRisk
Definition: | The risk of an adverse reaction (allergy or intolerance) for this patient upon exposure to the substance (including pharmaceutical products).
Committee: | Patient Care Work Group
OID: | 2.16.840.1.113883.4.642.3.940 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for AllergyIntoleranceSubstanceExposureRisk

4.4.1.528.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/allerg-intol-substance-exp-risk

4.4.1.528.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 2 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/allerg-intol-substance-exp-risk version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/allerg-intol-substance-exp-risk

Code | Display | Definition
known-reaction-risk | Known Reaction Risk | Known risk of allergy or intolerance reaction upon exposure to the specified substance.
no-known-reaction-risk | No Known Reaction Risk | No known risk of allergy or intolerance reaction upon exposure to the specified substance.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.529 Value Set http://hl7.org/fhir/ValueSet/questionnaire-enable-behavior

URL: https://hl7.org/fhir/R4/valueset-questionnaire-enable-behavior.html
FHIR Infrastructure Work Group | Maturity Level: 3 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/questionnaire-enable-behavior
Version: | 4.0.1
Name: | EnableWhenBehavior
Title: | EnableWhenBehavior
Definition: | Controls how multiple enableWhen values are interpreted - whether all or any must be true.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.1007 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for EnableWhenBehavior

• Resource: Questionnaire.item.enableBehavior (code / Required)

4.4.1.529.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/questionnaire-enable-behavior

4.4.1.529.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 2 concepts

Expansion based on http://hl7.org/fhir/questionnaire-enable-behavior version 4.0.1

All codes from system http://hl7.org/fhir/questionnaire-enable-behavior

Code | Display | Definition
all | All | Enable the question when all the enableWhen criteria are satisfied.
any | Any | Enable the question when any of the enableWhen criteria are satisfied.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.530 Value Set http://hl7.org/fhir/ValueSet/contract-assettype

URL: https://hl7.org/fhir/R4/valueset-contract-assettype.html
Financial Management Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/contract-assettype
Version: | 4.0.1
Name: | ContractResourceAssetTypeCodes
Title: | Contract Resource Asset Type codes
Definition: | This value set contract specific codes for asset type.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.1299 (for OID based terminology systems)
Copyright: | HL7 International.
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ContractResourceAssetTypeCodes

• Resource: Contract.term.asset.type (CodeableConcept / Example)

4.4.1.530.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/contract-asset-type

4.4.1.530.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 1 concepts

Expansion based on http://hl7.org/fhir/contract-asset-type version 4.0.1

All codes from system http://hl7.org/fhir/contract-asset-type

Code | Display | Definition
participation | Participation | To be completed

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.531 Value Set http://hl7.org/fhir/ValueSet/coverage-copay-type

URL: https://hl7.org/fhir/R4/valueset-coverage-copay-type.html
Financial Management Work Group | Maturity Level: 2 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/coverage-copay-type
Version: | 4.0.1
Name: | CoverageCopayTypeCodes
Title: | Coverage Copay Type Codes
Definition: | This value set includes sample Coverage Copayment Type codes.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.527 (for OID based terminology systems)
Copyright: | This is an example set.
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for CoverageCopayTypeCodes

• Resource: Coverage.costToBeneficiary.type (CodeableConcept / Extensible)

4.4.1.531.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/coverage-copay-type

4.4.1.531.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 10 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/coverage-copay-type version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/coverage-copay-type

Code | Display | Definition
gpvisit | GP Office Visit | An office visit for a general practitioner of a discipline.
spvisit | Specialist Office Visit | An office visit for a specialist practitioner of a discipline
emergency | Emergency | An episode in an emergency department.
inpthosp | Inpatient Hospital | An episode of an Inpatient hospital stay.
televisit | Tele-visit | A visit held where the patient is remote relative to the practitioner, e.g. by phone, computer or video conference.
urgentcare | Urgent Care | A visit to an urgent care facility - typically a community care clinic.

copaypct | Copay Percentage | A standard percentage applied to all classes or service or product not otherwise specified.
copay | Copay Amount | A standard fixed currency amount applied to all classes or service or product not otherwise specified.
deductible | Deductible | The accumulated amount of patient payment before the coverage begins to pay for services.
maxoutofpocket | Maximum out of pocket | The maximum amout of payment for services which a patient, or family, is expected to incur - typically annually.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.532 Value Set http://hl7.org/fhir/ValueSet/administration-method-codes

URL: https://hl7.org/fhir/R4/valueset-administration-method-codes.html
Pharmacy Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/administration-method-codes
Version: | 4.0.1
Name: | SNOMEDCTAdministrationMethodCodes
Title: | SNOMED CT Administration Method Codes
Definition: | This value set includes some method codes from SNOMED CT - provided as an exemplar
Committee: | Pharmacy Work Group
OID: | 2.16.840.1.113883.4.642.3.97 (for OID based terminology systems)
Copyright: | This value set includes content from SNOMED CT, which is copyright © 2002+ International Health Terminology Standards Development Organisation (IHTSDO), and distributed by agreement between IHTSDO and HL7. Implementer use of SNOMED CT is not covered by this agreement
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: MedicationAdministration.dosage.method (CodeableConcept / Example)

4.4.1.532.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include codes from http://snomed.info/sct where concept is-a 422096002 (Dosing instruction fragment)

4.4.1.532.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 58 concepts

All codes from system http://snomed.info/sct

Code | Display | Definition
417924000 | Apply
418283001 | Administer
419385000 | Use
419582001 | Give
419652001 | Take
419747000 | Chew
420045007 | Suck
420246001 | At
420247005 | Dosing instruction imperative
420295001 | Only
420341009 | Constant
420360002 | Sniff
420484009 | Subtract - dosing instruction fragment
420503003 | As
420561004 | Or
420604000 | Finish
420606003 | Shampoo
420620005 | Push
420652005 | Until gone

420771004 | Upon
420806001 | Per
420883007 | Sparingly
420942008 | Call
420974001 | When
421035004 | To
421066005 | Place
421067001 | Then
421134003 | Inhale
421139008 | Hold
421206002 | Multiply
421257003 | Insert
421286000 | Discontinue
421298005 | Swish and swallow
421399004 | Dilute
421463005 | With
421484000 | Then discontinue
421521009 | Swallow
421538008 | Instill
421548005 | Until
421612001 | Every
421682005 | Dissolve
421718005 | Before
421723005 | Now
421769005 | Follow directions
421803000 | If
421805007 | Swish
421829000 | And
421832002 | Twice
421939007 | Follow
421984009 | Until finished
421994004 | During
422033008 | Divide
422106007 | Add
422114001 | Once
422145002 | Inject
422152000 | Wash
422219000 | Sprinkle
422327006 | Then stop

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.533 Value Set http://hl7.org/fhir/ValueSet/participationstatus

URL: https://hl7.org/fhir/R4/valueset-participationstatus.html
Patient Administration Work Group | Maturity Level: 3 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/participationstatus
Version: | 4.0.1
Name: | ParticipationStatus
Title: | ParticipationStatus
Definition: | The Participation status of an appointment.
Committee: | Patient Administration Work Group
OID: | 2.16.840.1.113883.4.642.3.486 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ParticipationStatus

• ConceptMap: Translation source in Canonical Mapping for "ParticipationStatus"
• Resource: Appointment.participant.status (code / Required)
• Resource: AppointmentResponse.participantStatus (code / Required)

4.4.1.533.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/participationstatus

4.4.1.533.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 4 concepts

Expansion based on http://hl7.org/fhir/participationstatus version 4.0.1

All codes from system http://hl7.org/fhir/participationstatus

Code | Display | Definition | Canonical Status
accepted | Accepted | The participant has accepted the appointment. | ~accepted
declined | Declined | The participant has declined the appointment and will not participate in the appointment. | ~declined
tentative | Tentative | The participant has tentatively accepted the appointment. This could be automatically created by a system and requires further processing before it can be accepted. There is no commitment that attendance will occur. | ~draft
needs-action | Needs Action | The participant needs to indicate if they accept the appointment by changing this status to one of the other statuses. | ~failed

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.534 Value Set http://hl7.org/fhir/ValueSet/research-study-reason-stopped

URL: https://hl7.org/fhir/R4/valueset-research-study-reason-stopped.html
Biomedical Research and Regulation Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/research-study-reason-stopped
Version: | 4.0.1
Name: | ResearchStudyReasonStopped
Title: | ResearchStudyReasonStopped
Definition: | Codes for why the study ended prematurely.
Committee: | Biomedical Research and Regulation Work Group
OID: | 2.16.840.1.113883.4.642.3.825 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ResearchStudyReasonStopped
• Resource: ResearchStudy.reasonStopped (CodeableConcept / Example)

4.4.1.534.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/research-study-reason-stopped

4.4.1.534.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 4 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/research-study-reason-stopped version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/research-study-reason-stopped

Code | Display | Definition
accrual-goal-met | Accrual Goal Met | The study prematurely ended because the accrual goal was met.
closed-due-to-toxicity | Closed due to toxicity | The study prematurely ended due to toxicity.
closed-due-to-lack-of-study-progress | Closed due to lack of study progress | The study prematurely ended due to lack of study progress.
temporarily-closed-per-study-design | Temporarily closed per study design | The study prematurely ended temporarily per study design.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.535 Value Set http://hl7.org/fhir/ValueSet/conformance-expectation

URL: https://hl7.org/fhir/R4/valueset-conformance-expectation.html
FHIR Infrastructure Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/conformance-expectation
Version: | 4.0.1
Name: | ConformanceExpectation
Title: | ConformanceExpectation
Definition: | Indicates the degree of adherence to a specified behavior or capability expected for a system to be deemed conformant with a specification.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.919 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ConformanceExpectation

4.4.1.535.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/conformance-expectation

4.4.1.535.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 4 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/conformance-expectation version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/conformance-expectation

Code | Display | Definition
SHALL | SHALL | Support for the specified capability is required to be considered conformant.
SHOULD | SHOULD | Support for the specified capability is strongly encouraged, and failure to support it should only occur after careful consideration.
MAY | MAY | Support for the specified capability is not necessary to be considered conformant, and the requirement should be considered strictly optional.
SHOULD-NOT | SHOULD-NOT | Support for the specified capability is strongly discouraged and should occur only after careful consideration.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept

Comments | Additional notes about how to use the code

4.4.1.536 Value Set http://hl7.org/fhir/ValueSet/languages

URL: https://hl7.org/fhir/R4/valueset-languages.html
FHIR Infrastructure Work Group | Maturity Level: 3 | Trial Use | Use Context: Not yet ready for Production use

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/languages
Version: | 4.0.1
Name: | CommonLanguages
Title: | Common Languages
Definition: | This value set includes common codes from BCP-47 (http://tools.ietf.org/html/bcp47)
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.20 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

- Resource: Resource.language (code / Preferred)
- Profile: ShareableMeasure: Measure.language (code / Preferred)
- Profile: ServiceRequest-Genetics: ServiceRequest.language (code / Preferred)
- Resource: RelatedPerson.communication.language (CodeableConcept / Preferred)
- Resource: Practitioner.communication (CodeableConcept / Preferred)
- Profile: GroupDefinition: Group.language (code / Preferred)
- Profile: ActualGroup: Group.language (code / Preferred)
- Profile: FamilyMemberHistory-Genetic: FamilyMemberHistory.language (code / Preferred)
- Profile: ShareableActivityDefinition: ActivityDefinition.language (code / Preferred)
- Profile: CDSHooksRequestGroup: RequestGroup.language (code / Preferred)
- Profile: provenance-relevant-history: Provenance.language (code / Preferred)
- Profile: CQF-Questionnaire: Questionnaire.language (code / Preferred)
- Resource: ExplanationOfBenefit.processNote.language (CodeableConcept / Preferred)
- Resource: ValueSet.compose.include.concept.designation.language (code / Preferred)
- Profile: ShareableValueSet: ValueSet.language (code / Preferred)
- Profile: ShareableValueSet: ValueSet.compose.include.concept.designation.language (code / Preferred)
- Profile: PICOElement: EvidenceVariable.language (code / Preferred)
- Resource: CodeSystem.concept.designation.language (code / Preferred)
- Profile: ShareableCodeSystem: CodeSystem.language (code / Preferred)
- Profile: ShareableCodeSystem: CodeSystem.concept.designation.language (code / Preferred)
- Profile: CDSHooksGuidanceResponse: GuidanceResponse.language (code / Preferred)
- Profile: DeviceMetricObservation: Observation.language (code / Preferred)
- Profile: Observation-Genetics: Observation.language (code / Preferred)
- Profile: VitalSigns: Observation.language (code / Preferred)
- Profile: BodyWeight: Observation.language (code / Preferred)
- Profile: VitalsPanel: Observation.language (code / Preferred)
- Profile: BodyHeight: Observation.language (code / Preferred)
- Profile: RespRate: Observation.language (code / Preferred)
- Profile: HeartRate: Observation.language (code / Preferred)
- Profile: BodyTemp: Observation.language (code / Preferred)
- Profile: HeadCircum: Observation.language (code / Preferred)
- Profile: OxygenSat: Observation.language (code / Preferred)
- Profile: BMI: Observation.language (code / Preferred)
- Profile: BP: Observation.language (code / Preferred)
- Profile: ShareableLibrary: Library.language (code / Preferred)
- Profile: CQLLibrary: Library.language (code / Preferred)
- Resource: ClaimResponse.processNote.language (CodeableConcept / Preferred)
- Profile: LipidProfile: DiagnosticReport.language (code / Preferred)

• Profile: Cholesterol: Observation.language (code / Preferred)
• Profile: Triglyceride: Observation.language (code / Preferred)
• Profile: HDLCholesterol: Observation.language (code / Preferred)
• Profile: LDLCholesterol: Observation.language (code / Preferred)
• Profile: DiagnosticReport-Genetics: DiagnosticReport.language (code / Preferred)
• Profile: HLAResult: DiagnosticReport.language (code / Preferred)
• Resource: HealthcareService.communication (CodeableConcept / Preferred)
• Profile: Synthesis: Evidence.language (code / Preferred)
• Profile: ClinicalDocument: Composition.language (code / Preferred)
• Profile: Catalog: Composition.language (code / Preferred)
• Resource: Patient.communication.language (CodeableConcept / Preferred)
• Profile: ShareablePlanDefinition: PlanDefinition.language (code / Preferred)
• Profile: ComputablePlanDefinition: PlanDefinition.language (code / Preferred)
• Profile: CDSHooksServicePlanDefinition: PlanDefinition.language (code / Preferred)
• Resource: Attachment.language (code / Preferred)
• Extension: http://hl7.org/fhir/StructureDefinition/language: Human Language (code / Preferred)
• Extension: http://hl7.org/fhir/StructureDefinition/cqf-recipientLanguage: recipientLanguage (CodeableConcept / Preferred)
• Extension: http://hl7.org/fhir/StructureDefinition/devicerequest-patientInstruction: patientInstruction (code / Preferred)
• Extension: http://hl7.org/fhir/StructureDefinition/cqf-systemUserLanguage: systemUserLanguage (CodeableConcept / Preferred)
• Extension: http://hl7.org/fhir/StructureDefinition/translation: Translation (code / Preferred)

4.4.1.536.1 Content Logical Definition

This value set includes codes from the following code systems:

urn:ietf:bcp:47

Code	Display
ar	Arabic
bn	Bengali
cs	Czech
da	Danish
de	German
de-AT	German (Austria)
de-CH	German (Switzerland)
de-DE	German (Germany)
el	Greek
en	English
en-AU	English (Australia)
en-CA	English (Canada)
en-GB	English (Great Britain)
en-IN	English (India)
en-NZ	English (New Zeland)
en-SG	English (Singapore)
en-US	English (United States)
es	Spanish
es-AR	Spanish (Argentina)
es-ES	Spanish (Spain)
es-UY	Spanish (Uruguay)
fi	Finnish
fr	French
fr-BE	French (Belgium)
fr-CH	French (Switzerland)
fr-FR	French (France)
fy	Frysian
fy-NL	Frysian (Netherlands)
hi	Hindi
hr	Croatian
it	Italian
it-CH	Italian (Switzerland)
it-IT	Italian (Italy)
ja	Japanese
ko	Korean

nl | Dutch
nl-BE | Dutch (Belgium)
nl-NL | Dutch (Netherlands)
no | Norwegian
no-NO | Norwegian (Norway)
pa | Punjabi
pl | Polish
pt | Portuguese
pt-BR | Portuguese (Brazil)
ru | Russian
ru-RU | Russian (Russia)
sr | Serbian
sr-RS | Serbian (Serbia)
sv | Swedish
sv-SE | Swedish (Sweden)
te | Telegu
zh | Chinese
zh-CN | Chinese (China)
zh-HK | Chinese (Hong Kong)
zh-SG | Chinese (Singapore)
zh-TW | Chinese (Taiwan)

Additional Language Displays

Code | Dansk (Danish, da) | Deutsch (German, de) | English (English, en) | Nederlands (Dutch, nl) | Polskie (Polish, pl) | Русский (Russian, ru) | □□ (Chinese, zh)
ar | Arabisk | | Arabic | Arabisch
bn | Bengalsk | Bengalisch | Bengali | Bengaals | | Бенгальский | □□□□
cs | Tjekkisk | Tschechisch | Czech | Tsjechisch | | Чешский | □□□
da | Dansk | Dänisch | Danish | Deens | | Датский | □□□
de | Tysk | Deutsch | German | Duits | | Немецкий | □□
de-AT | Tysk (Østrig | Deutsch (Österreich) | German (Austria) | Duits (Oostenrijk) | | Немецкий (Австрия) | □□ □□□□□
de-CH | Tysk (Schweiz) | Deutsch (Schweiz) | German (Switzerland) | Duits (Zwitserland) | | Немецкий (Швейцария) | □□ □□□□
de-DE | Tysk (Tyskland) | Deutsch (Deutschland) | German (Germany) | Duits (Duitsland) | | Немецкий (Германия) | □□ □□□□
el | Græsk | Griechisch | Greek | Grieks | | Греческий | □□□
en | Engelsk | Englisch | English | Engels | | Английский | □□
en-AU | Engelsk (Australien) | Englisch (Australien) | English (Australia) | Engels (Australië) | | Английский (Австралия) | □□ □□□□□□
en-CA | Engelsk (Canada) | Englisch (Kanada) | English (Canada) | Engels (Canada) | | Английский (Канада) | □□ □□□□□
en-GB | Engelsk (Storbritannien) | Englisch (Großbritannien) | English (Great Britain) | Engels (Groot Brittannië) | | Английский (Великобритания) | □□ □□□□
en-IN | Engelsk (Indien) | Englisch (Indien) | English (India) | Engels (India) | | Английский (Индия) | □□ □□□□
en-NZ | Engelsk (New Zeeland) | Englisch (Neuseeland) | English (New Zeland) | Engels (Nieuw Zeeland) | | Английский (Новая Зеландия) | □□ □□□□□
en-SG | Engelsk (Singapore) | Englisch (Singapur) | English (Singapore) | Engels (Singapore) | | Английский (Сингапур) | □□ □□□□□
en-US | Engelsk (Amerikansk) | Englisch (USA) | English (United States) | Engels (Verenigde Staten) | | Английский (США) | □□ □□□□
es | Spansk | Spanisch | Spanish | Spaans | | Испанский | □□□□
es-AR | Spansk (Argentina) | Spanisch (Argentinien) | Spanish (Argentina) | Spaans (Argentinië) | | Испанский (Аргентина) | □□□□ □□□□□
es-ES | Spansk (Spanien) | Spanisch (Spanien) | Spanish (Spain) | Spaans (Spanje) | | Испанский (Испания) | □□□□ □□□□□
es-UY | Spansk (Uruguay) | Spanisch (Uruguay) | Spanish (Uruguay) | Spaans (Uruguay) | | Испанский (Уругвай) | □□□□ □□□□□
fi | Finsk | Finnisch | Finnish | Fins | | Финский | □□□
fr | Fransk | Französisch | French | Frans | | Французский | □□
fr-BE | Finsk (Belgien) | Französisch (Belgien) | French (Belgium) | Frans (België)
fr-CH | Fransk (Schweiz) | Französisch (Schweiz) | French (Switzerland) | Frans (Zwitserland) | | Французский (Швейцария) | □□ □□□□□
fr-FR | Fransk (Frankrig) | Französisch (Frankreich) | French (France) | Frans (Frankrijk) | | Французский (Франция) | □□ □□□□
fy | Frisisk | Friesisch | Frysian | Fries | | | □□□□□
fy-NL | Frisisk (Holland) | Friesisch (Niederlande) | Frysian (Netherlands) | Fries (Nederland) | | | □□□□□□□□□□
hi | Hindi | | Hindi | Hindi
hr | Kroatisk | Kroatisch | Croatian | Kroatisch | | Хорватский | □□□□□
it | Italiensk | Italienisch | Italian | Italiaans | | Итальянский | □□□□
it-CH | Italiensk (Schweiz) | Italienisch (Schweiz) | Italian (Switzerland) | Italiaans (Zwitserland) | | Итальянский (Швейцария) | □□□□ □□□□
it-IT | Italiensk (Italien) | Italienisch (Italien) | Italian (Italy) | Italiaans (Italië) | | Итальянский (Италия) | □□□□ □□□□□
ja | Japansk | Japanisch | Japanese | Japans | | Японский | □□
ko | Koreansk | Koreanisch | Korean | Koreaans | | Корейский | □□□
nl | Hollandsk | Niederländisch | Dutch | Nederlands | | Нидерландский | □□□
nl-BE | Hollandsk (Belgien) | Niederländisch (Belgien) | Dutch (Belgium) | Nederlands (België) | | | □□□ □□□□□
nl-NL | Hollandsk (Holland) | Niederländisch (Niederlande) | Dutch (Netherlands) | Nederlands (Nederland) | | Нидерландский (Нидерланды) | □□□ □□□□
no | Norsk | Norwegisch | Norwegian | Noors | | Норвежский | □□□
no-NO | Norsk (Norge) | Norwegisch (Norwegen) | Norwegian (Norway) | Noors (Noorwegen) | | Норвежский (Норвегия) | □□□ □□□□
pa | Punjabi | | Punjabi | Punjabi

pl | | | | | Polskie
pt | Portugisisk | Portugiesisch | Portuguese | Portugees | | Португальский | □□□□
pt-BR | Portugisisk (Brasilien) | Portugiesisch (Brasilien) | Portuguese (Brazil) | Portugees (Brazilië) | | Португальский (Бразилия) | □□□□ □□□□
ru | Russisk | Russisch | Russian | Russisch | | Русский | □□
ru-RU | Russisk (Rusland) | Russisch (Russland) | Russian (Russia) | Russisch (Rusland) | | Русский (Россия) | □□ □□□□□
sr | Serbisk | Serbisch | Serbian | Servisch | | Сербский | □□□□□
sr-RS | Serbisk (Serbien) | Serbisch (Serbien) | Serbian (Serbia) | Servisch (Servië) | | Сербский (Сербия) | □□□□□ □□□□□□
sv | Svensk | Schwedisch | Swedish | Zweeds | | Шведский | □□□
sv-SE | Svensk (Sverige) | Schwedisch (Schweden) | Swedish (Sweden) | Zweeds (Zweden) | | Шведский (Швеция) | □□□ □□□□
te | Telugu | Telugu | Telegu | Teloegoe | | Телугу | □□□□
zh | Kinesisk | Chinesisch | Chinese | Chinees | | Китайский | □□
zh-CN | Kinesisk (Kina) | Chinesisch (China) | Chinese (China) | Chinees (China) | | Китайский (Китай) | □□ □□□□
zh-HK | Kinesisk (Hong Kong) | Chinesisch (Hong Kong) | Chinese (Hong Kong) | Chinees (Hong Kong) | | Китайский (Гонконг) | □□ □□□□□□□□□
zh-SG | Kinesisk (Singapore) | Chinesisch (Singapur) | Chinese (Singapore) | Chinees (Singapore) | | Китайский (Сингапур) | □□ □□□□□
zh-TW | Kinesisk (Taiwan) | Chinesisch (Taiwan) | Chinese (Taiwan) | Chinees (Taiwan) | | Китайский (Тайвань) | □□ □□□□

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.537 Value Set http://hl7.org/fhir/ValueSet/missing-tooth-reason

URL: https://hl7.org/fhir/R4/valueset-missing-tooth-reason.html
Financial Management Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/missing-tooth-reason
Version: | 4.0.1
Name: | MissingToothReasonCodes
Title: | Missing Tooth Reason Codes
Definition: | This value set includes sample Missing Tooth Reason codes.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.534 (for OID based terminology systems)
Copyright: | This is an example set.
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for MissingToothReasonCodes
• Resource: ExplanationOfBenefit.supportingInfo.reason (Coding / Example)

4.4.1.537.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/missingtoothreason

4.4.1.537.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 4 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/missingtoothreason version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/missingtoothreason

Code | Display | Definition
e | E | Extraction
c | C | Congenital
u | U | Unknown
o | O | Other

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.538 Value Set http://hl7.org/fhir/ValueSet/precision-estimate-type

URL: https://hl7.org/fhir/R4/valueset-precision-estimate-type.html
Clinical Decision Support Work Group | Maturity Level: 0 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/precision-estimate-type
Version: | 4.0.1
Name: | PrecisionEstimateType
Title: | PrecisionEstimateType
Definition: | Method of reporting variability of estimates, such as confidence intervals, interquartile range or standard deviation.
Committee: | Clinical Decision Support Work Group
OID: | 2.16.840.1.113883.4.642.3.1357 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for PrecisionEstimateType
• Resource: EffectEvidenceSynthesis.effectEstimate.precisionEstimate.type (CodeableConcept / Extensible)
• Resource: RiskEvidenceSynthesis.riskEstimate.precisionEstimate.type (CodeableConcept / Extensible)

4.4.1.538.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/precision-estimate-type

4.4.1.538.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 4 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/precision-estimate-type version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/precision-estimate-type

Code | Display | Definition
CI | confidence interval | confidence interval.
IQR | interquartile range | interquartile range.
SD | standard deviation | standard deviation.
SE | standard error | standard error.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.539 Value Set http://hl7.org/fhir/ValueSet/additionalmaterials

URL: https://hl7.org/fhir/R4/valueset-additionalmaterials.html
Financial Management Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/additionalmaterials
Version: | 4.0.1
Name: | AdditionalMaterialCodes
Title: | Additional Material Codes
Definition: | This value set includes sample additional material type codes.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.529 (for OID based terminology systems)
Copyright: | This is an example set.
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for AdditionalMaterialCodes

4.4.1.539.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/additionalmaterials

4.4.1.539.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 6 concepts

Expansion based on http://hl7.org/fhir/additionalmaterials version 4.0.1

All codes from system http://hl7.org/fhir/additionalmaterials

Code	Display	Definition
xray	XRay	XRay
image	Image	Image
email	Email	Email
model	Model	Model
document	Document	Document
other	Other	Other

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.540 Value Set http://hl7.org/fhir/ValueSet/medicationknowledge-package-type

URL: https://hl7.org/fhir/R4/valueset-medicationknowledge-package-type.html
Pharmacy Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/medicationknowledge-package-type
Version: | 4.0.1
Name: | medicationKnowledge Package Type Codes
Title: | Medication knowledge package type codes
Definition: | MedicationKnowledge Package Type Codes
Committee: | Pharmacy Work Group
OID: | 2.16.840.1.113883.4.642.3.1339 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for medicationKnowledge Package Type Codes

• Resource: MedicationKnowledge.packaging.type (CodeableConcept / Example)

4.4.1.540.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/medicationknowledge-package-type

4.4.1.540.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 22 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/medicationknowledge-package-type version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/medicationknowledge-package-type

Code | Display | Definition
amp | Ampule
bag | Bag
blstrpk | Blister Pack
bot | Bottle
box | Box
can | Can
cart | Cartridge
disk | Disk
doset | Dosette
jar | Jar
jug | Jug
minim | Minim
nebamp | Nebule Amp
ovul | Ovule
pch | Pouch
pkt | Packet
sash | Sashet
strip | Strip
tin | Tin
tub | Tub
tube | Tube
vial | Vial

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.541 Value Set http://hl7.org/fhir/ValueSet/care-plan-activity-outcome

URL: https://hl7.org/fhir/R4/valueset-care-plan-activity-outcome.html
Patient Care Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined at http://www.hl7.org/Special/committees/patientcare/ .

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/care-plan-activity-outcome
Version: | 4.0.1
Name: | CarePlanActivityOutcome
Title: | Care Plan Activity Outcome
Definition: | Example codes indicating the outcome of a care plan activity. Note that these are in no way complete and might not even be appropriate for some uses.
Committee: | Patient Care Work Group
OID: | 2.16.840.1.113883.4.642.3.149 (for OID based terminology systems)
Copyright: | This resource includes content from SNOMED Clinical Terms® (SNOMED CT®) which is copyright of the International Health Terminology Standards Development Organisation (IHTSDO). Implementers of these specifications must have the appropriate SNOMED CT Affiliate license - for more information contact http://www.snomed.org/snomed-ct/get-snomed-ct or info@snomed.org
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: CarePlan.activity.outcomeCodeableConcept (CodeableConcept / Example)

4.4.1.541.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include codes from http://snomed.info/sct where concept is-a 397640006 (Healthcare knowledge finding)

4.4.1.541.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 42 concepts

All codes from system http://snomed.info/sct

Code	Display	Definition
54777007	Deficient knowledge	
129862009	Deficient knowledge: diagnostic test	
129863004	Deficient knowledge of dietary regimen	
129864005	Deficient knowledge: disease process	
129865006	Deficient knowledge of fluid volume management	
129866007	Deficient knowledge: medication regimen	
129867003	Deficient knowledge of safety precautions	
129868008	Deficient knowledge: therapeutic regimen	
182891003	Patient misunderstood treatment instructions	
397588000	Demonstrates knowledge of expected responses to operative procedure	
397598006	Demonstrates knowledge of nutritional requirements related to operative procedure	
397675006	Demonstrates knowledge of wound healing process	
397699006	Demonstrates knowledge of medication management	
397724004	Demonstrates knowledge of pain management	
408364003	Patient does not understand why taking all medication	
423875000	Deficient knowledge of family planning methods	
424537004	Deficient knowledge of preconception health practices	
704375003	Deficient knowledge of breast feeding	
704377006	Deficient knowledge of peritoneal dialysis	
704379009	Deficient knowledge of sexual behavior	
704381006	Deficient knowledge of oral hygiene	
704383009	Deficient knowledge of child development	
704400005	Deficient knowledge of community services	
704402002	Deficient knowledge of physical therapy	
704403007	Deficient knowledge of fall prevention	
704404001	Deficient knowledge of pain management	
704406004	Deficient knowledge of symptom	
704446005	Deficient knowledge of exercise	
704491001	Deficient knowledge of traditional therapy	
704680001	Deficient knowledge of behavior change process	
705086000	Deficient knowledge of patient controlled analgesia	
706869001	Deficient knowledge of medication	
708571001	Demonstrates knowledge of who to contact in medical emergency	
713398008	Deficient knowledge of travel health	
713447003	Deficient knowledge of parenting	
713448008	Deficient knowledge of infant feeding	
713449000	Deficient knowledge of infant care	
713451001	Deficient knowledge of childbirth	
713452008	Deficient knowledge of foetal development	
714670007	Deficient knowledge of pregnancy	
716391009	Able to adapt to sensory changes in body part	
716438000	Able to recognize need for pressure area care	

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information

Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)

Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')

Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application

Definition | An explanation of the meaning of the concept

Comments | Additional notes about how to use the code

4.4.1.542 Value Set http://hl7.org/fhir/ValueSet/codesystem-content-mode

URL: https://hl7.org/fhir/R4/valueset-codesystem-content-mode.html

Vocabulary Work Group | Maturity Level: N | Normative (from v4.0.0) | Use Context: Any

| This page has been approved as part of an ANSI standard. See the Conformance Package for further details.

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/codesystem-content-mode
Version: | 4.0.1
Name: | CodeSystemContentMode
Title: | CodeSystemContentMode
Definition: | The extent of the content of the code system (the concepts and codes it defines) are represented in a code system resource.
Committee: | Vocabulary Work Group
OID: | 2.16.840.1.113883.4.642.3.782 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for CodeSystemContentMode
• Resource: CodeSystem.content (code / Required)

4.4.1.542.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/codesystem-content-mode

4.4.1.542.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 5 concepts

Expansion based on http://hl7.org/fhir/codesystem-content-mode version 4.0.1

All codes from system http://hl7.org/fhir/codesystem-content-mode

Code | Display | Definition
not-present | Not Present | None of the concepts defined by the code system are included in the code system resource.
example | Example | A few representative concepts are included in the code system resource. There is no useful intent in the subset choice and there's no process to make it workable: it's not intended to be workable.
fragment | Fragment | A subset of the code system concepts are included in the code system resource. This is a curated subset released for a specific purpose under the governance of the code system steward, and that the intent, bounds and consequences of the fragmentation are clearly defined in the fragment or the code system documentation. Fragments are also known as partitions.
complete | Complete | All the concepts defined by the code system are included in the code system resource.
supplement | Supplement | The resource doesn't define any new concepts; it just provides additional designations and properties to another code system.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.543 Value Set http://hl7.org/fhir/ValueSet/benefit-unit

URL: https://hl7.org/fhir/R4/valueset-benefit-unit.html
Financial Management Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/benefit-unit
Version: | 4.0.1
Name: | UnitTypeCodes
Title: | Unit Type Codes
Definition: | This value set includes a smattering of Unit type codes.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.610 (for OID based terminology systems)
Copyright: | This is an example set.
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for UnitTypeCodes

• Resource: CoverageEligibilityResponse.insurance.item.unit (CodeableConcept / Example)
• Resource: ExplanationOfBenefit.benefitBalance.unit (CodeableConcept / Example)

4.4.1.543.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/benefit-unit

4.4.1.543.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 2 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/benefit-unit version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/benefit-unit

Code	Display	Definition
individual	Individual	A single individual
family	Family	A family, typically includes self, spouse(s) and children to a defined age

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl	A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source	The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code	The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display	The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition	An explanation of the meaning of the concept
Comments	Additional notes about how to use the code

4.4.1.544 Value Set http://hl7.org/fhir/ValueSet/device-type

URL: https://hl7.org/fhir/R4/valueset-device-type.html

Orders and Observations Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL:	http://hl7.org/fhir/ValueSet/device-type
Version:	4.0.1
Name:	DeviceType
Title:	Device Type
Definition:	Codes used to identify medical devices. Includes concepts from SNOMED CT (http://www.snomed.org/) where concept is-a 49062001 (Device) and is provided as a suggestive example.
Committee:	Orders and Observations Work Group
OID:	2.16.840.1.113883.4.642.3.1376 (for OID based terminology systems)
Copyright:	This resource includes content from SNOMED Clinical Terms® (SNOMED CT®) which is copyright of the International Health Terminology Standards Development Organisation (IHTSDO). Implementers of these specifications must have the appropriate SNOMED CT Affiliate license - for more information contact http://www.snomed.org/snomed-ct/get-snomed-ct or info@snomed.org
Source Resource	XML / JSON

This value set is used in the following places:

• Resource: Device.type (CodeableConcept / Example)

4.4.1.544.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include codes from http://snomed.info/sct where concept is-a 49062001 (Device)

4.4.1.544.2 Expansion

This expansion generated 01 Nov 2019

This value set has >1000 codes in it. In order to keep the publication size manageable, only a selection (1000 codes) of the whole set of codes is shown

All codes from system http://snomed.info/sct

Code	Display	Definition
156009	Spine board	
271003	Bone plate	
287000	Air receiver	
291005	Atomizer	

678001 | Epilator
739006 | Bicycle ergometer
793009 | Mechanical power press
882002 | Diagnostic implant
972002 | Air filter, device
989005 | Linen cloth
994005 | Brush, device
1211003 | Treadmill, device
1333003 | Emesis basin, device
1422002 | Plastic mold, device
1579007 | Surgical drill, device
1766001 | Toboggan, device
1941006 | Silk cloth
1962007 | Dike, device
2248009 | Tracheal tube cuff
2282003 | Breast implant
2287009 | Nail file, device
2468001 | Breath analyzer, device
2478003 | Ocular prosthesis
2491002 | Intra-aortic balloon catheter, device
3201004 | Tent, device
3319006 | Artificial liver, device
4408003 | Endoscopic camera, device
4632004 | Hair cloth
4816004 | Metal device
5041003 | Adhesive strip, device
5042005 | Patient scale, device
5159002 | Physiologic monitoring system, device
5679009 | Bed sheet
6012004 | Hearing aid
6097006 | T-tube
6822006 | Microwave oven
6919005 | Protective clothing material
6972009 | Lithotripter
7007007 | Radiographic-fluoroscopic unit
7402007 | Probe
7406005 | Crib
7704007 | Stabilizing appliance
7733008 | Hydrocephalic shunt catheter
7968002 | Three-wheeled all-terrain vehicle
7971005 | Fogarty catheter
8060009 | Denture
8118007 | Crane
8170008 | Adhesive
8384009 | Band saw
8407004 | Bile collection bag
8434001 | Gaol
8451008 | Intramedullary nail
8615009 | Blood electrolyte analyzer
8643000 | Mortising machine
8682003 | Protective shield
9017009 | Ventricular intracranial catheter
9096001 | Support
9129003 | Feeding catheter
9419002 | Bobsled
9458007 | Elastic bandage
9611009 | Dermatotome
9883003 | Cargo handling gear
10244001 | Needle guide
10371004 | Electrostimulating analgesia unit
10507000 | Toeboard
10826000 | Industrial saw
10850003 | Radiographic-therapeutic unit
10906003 | Vein stripper
11141007 | Bone growth stimulator
11158002 | Electromyographic monitor and recorder
11358008 | Prosthetic valve
11987000 | Clinical chemistry analyzer
12150006 | Cannula
12183004 | Upper limb prosthesis
12198002 | Ice skate
12953007 | File

13118005 | Wool cloth
13219008 | Gastroscope
13288007 | Monitors
13459008 | Temporary artificial heart prosthesis
13764006 | Uterine sound
13855007 | Pillow
13905003 | Tennis ball
14106009 | Cardiac pacemaker implant
14108005 | Cage
14116001 | Analgesia unit
14208000 | Oil well
14288003 | Nasal septum button
14339000 | Button
14364002 | Camera
14423008 | Adhesive bandage
14519003 | Aspirator
14548009 | Harrington rod
14762000 | Alloy steel chain sling
14789005 | Prosthetic implant
15000008 | Air-conditioner
15340005 | Wood's light
15447007 | Arthroplasty prosthesis
15644007 | Anesthesia unit
15869005 | Dosimeter, device
15873008 | Boiler, device
15922004 | Gown, device
16056004 | Boots
16349000 | Orthopedic equipment
16417001 | Commercial breathing supply hoses diving operation, device
16470007 | Electrode, device
16497000 | Electric clipper, device
16524003 | Cotton cloth
16540000 | Umbrella catheter, device
16650009 | Splint, device
17102003 | NG - Nasogastric tube
17107009 | Prosthetic mitral valve
17207004 | Mattress, device
17306006 | Hernia belt, device
17404008 | Cardiac compression board, device
17472008 | Knife, device
18151003 | Punch, device
18153000 | Fluorescence immunoassay analyzer, device
18411005 | Chisel, device
18666004 | Finespun glass
19257004 | Defibrillator, device
19328000 | Blanket, device
19443004 | Radioactive implant, device
19627002 | Leather
19762002 | Leather belt
19817005 | Fan blade, device
19892000 | Scale, device
19923001 | Catheter, device
20195009 | Leg prosthesis, device
20235003 | Toothbrush, device
20273004 | Industrial machine, device
20359006 | Contraceptive diaphragm, device
20406008 | Back rests, device
20428008 | Oxygen tent, device
20513005 | Power tool, device
20568009 | Urinary catheter, device
20613002 | Cystoscope, device
20861007 | Plug pack, device
20867006 | Experimental implant, device
20873007 | Plastic cloth-like material
20997002 | Hand tool, device
21079000 | Carbon monoxide analyzer, device
21546008 | Icebox
21870002 | Transluminal extraction catheter, device
21944004 | Abdominal binder, device
22251003 | Timer, device
22283009 | Artificial membrane, device
22566001 | Cytology brush, device

22662007 | Retaining harness, device
22679001 | Handcuffs, device
22744006 | Artificial hair wig, device
22852002 | Therapeutic implant, device
23228005 | Arthroscope, device
23366006 | Motorized wheelchair device
23699001 | Baseball, device
23785007 | Arthroscopic irrigation/distension pump, device
23973005 | Indwelling urinary catheter, device
24073000 | Mechanical cardiac valve prosthesis
24110008 | Anoscope, device
24174009 | Bronchoscope, device
24230000 | Vibrator, device
24290003 | Cardiac valve bioprosthesis
24402003 | Stepladder, device
24470005 | Wrench, device
24513003 | Plastic boots
24697008 | Ostomy belt, device
24767007 | Eustachian tube prosthesis, device
25005004 | Snare, device
25062003 | Feeding tube, device
25152007 | Squeeze cage, device
25510005 | Heart valve prosthesis
25632005 | Hockey puck, device
25680008 | Scaffold, device
25742001 | Orthodontic appliance, device
25937001 | Neurostimulation device
26128008 | Bougie, device
26239002 | Soccer ball, device
26334009 | Dockboard, device
26397000 | Reservoir bag
26412008 | ET - Endotracheal tube
26579007 | Holter valve, device
26719000 | Celestin tube, device
26882005 | Rongeur, device
27042007 | Needle adapter, device
27065002 | Suture
27091001 | Dumbwaiter, device
27126002 | Power belt, device
27229001 | Spray booth, device
27606000 | Dental prosthesis, device
27785006 | Athletic supporter, device
27812008 | Electric heating pad, device
27976001 | Woodworking machinery, device
27991004 | Thermometer, device
28026003 | Hairbrush, device
29292008 | Fur garment
29319002 | Forceps, device
29396008 | Resuscitator, device
30012001 | Elevator, device
30070001 | Multistage suspension scaffolding, device
30115002 | Shield, device
30176005 | Baseball bat, device
30234008 | Medical laboratory analyzer, device
30610008 | Epidural catheter, device
30661003 | Cosmetic prosthesis, device
30929000 | Ligator, device
30968007 | Drainage bag, device
31030004 | Peritoneal catheter, device
31031000 | Internal fixator
31174004 | Lumbosacral belt, device
31791005 | Traction belt, device
31878003 | Surgical scissors, device
32033000 | Arterial pressure monitor, device
32356002 | Machine guarding, device
32504006 | Screwdriver, device
32634007 | Fixed ladder, device
32667006 | Oral airway
32711007 | Ostomy collection bag, device
32712000 | Drain, device
32871007 | Tweezer, device
33194000 | Welding equipment, device

33352006 | Ax, device
33388001 | Carbon dioxide analyzer, device
33482001 | Rubber boots
33686008 | Stylet, device
33690005 | Sharp instrument, device
33802005 | Enema bag, device
33894003 | Experimental device
33918000 | Rubberized cloth
34164001 | POP - Plaster of Paris cast
34188004 | Straightjacket, device
34234003 | Plastic tube, device
34263000 | Medical balloon, device
34362008 | Vascular device
34759008 | Urethral catheter, device
35398009 | Ostomy appliance, device
35593004 | Wire ligature, device
35870000 | Cerebrospinal catheter, device
36365007 | Ice-pick, device
36370000 | Aspirator trap bottle, device
36645008 | Stimulator, device
36761001 | Natural hair wig, device
36965003 | Hemodialysis machine, device
36977008 | Peripheral nerve stimulator
37189001 | Magnetic detector, device
37270008 | Endoscope, device
37284003 | Bag, device
37311003 | Stone retrieval basket, device
37347002 | Dildo, device
37360008 | Patient isolator, device
37503007 | Protective blind, device
37759000 | Surgical instrument, device
37874008 | Continuing positive airway pressure unit, device
37953008 | Bedside rails, device
38126007 | Protective lenses
38141007 | Tourniquet, device
38277008 | Protective device
38806006 | Hockey stick, device
38862006 | Sheet metal bending equipment
38871002 | Metallic cloth
39590006 | Air compressor, device
39690000 | Sphygmomanometer, device
39768008 | Rasp, device
39790008 | Non-electric heating pad, device
39802000 | Tongue blade, device
39821008 | Positron emission tomography unit, device
39849001 | Nasal oxygen cannula
39869006 | Alarm, device
40388003 | Biomedical implant
40519001 | Binder, device
41157002 | Orthopedic immobilizer
41323003 | Urinary collection bag, device
41525006 | Artificial structure, device
41684000 | Industrial tool, device
42152006 | Metal tube, device
42305009 | Ambulation device
42380001 | Ear plug, device
42451009 | Blood warmer, device
42716000 | Wool
42882002 | Hypodermic spray, device
43001000 | Phlebotomy kit, device
43192004 | Bone pencil, device
43252007 | Cochlear implant
43725001 | Airway equipment
43734006 | Blood administration set, device
43770009 | Doppler device
43983001 | Shoes
44056008 | Caliper, device
44396004 | Magnet, device
44492001 | Industrial robot, device
44668000 | Pump, device
44738004 | Laryngoscope, device
44806002 | Esophageal bougie, device

44845007 | Golf ball, device
44907005 | Four-wheeled all-terrain vehicle, device
44959004 | Angioplasty balloon catheter, device
45633005 | Peritoneal dialyzer, device
45901004 | Penrose drain, device
46181005 | Automatic fire extinguisher system, device
46265007 | Artificial lashes, device
46299005 | Sanitary belt, device
46364009 | Clamp, device
46440007 | Basketball, device
46625003 | Suppository
46666003 | Chain, device
46949002 | Deck machinery, device
47162009 | Mirror, device
47326004 | Electrical utilization equipment, device
47424002 | Apgar scoring timer, device
47528002 | Ureteric catheter
47731004 | Birthing chair, device
47734007 | Chromic catgut suture
47776004 | Mittens
47942000 | Proctoscope, device
48066006 | Circular portable saw, device
48096001 | Bathtub rails, device
48240003 | Training equipment, device
48246009 | Studgun, device
48295009 | Vascular filter, device
48473008 | Protective body armor, device
48822005 | Bilirubin light, device
48990009 | Strap, device
49448001 | Razor, device
49890001 | Hospital cart, device
50121007 | Glasses
50457005 | Workover rig service to oil well, device
50483000 | Oil rig
50851003 | Penile tumescence monitor, device
51016001 | Hammer, device
51086006 | Shower curtain, device
51324004 | Stripper, device
51685009 | Roller skate, device
51791000 | Measuring tape, device
51832004 | Valved tube, device
51883004 | Sling, device
52124006 | Central line
52161002 | Molten lava
52291003 | Gloves
52520009 | Ladder, device
52537002 | Aspirator collection canister, device
52624007 | Radiofrequency generator, device
52773007 | Ski, device
52809000 | Nasopharyngeal catheter, device
52893008 | Blood gas/pH analyzer, device
53167006 | Platform suspended boom, device
53177008 | Nasal balloon, device
53217009 | Artificial lung, device
53350007 | Prosthesis, device
53535004 | Retractor, device
53639001 | Stethoscope, device
53671008 | Gastric balloon, device
53996008 | Penile prosthesis, device
54234007 | Cryogenic analgesia unit, device
54638004 | Towel, device
54953005 | Computerized axial tomography scanner, device
55091003 | Blood coagulation analyzer, device
55206006 | Suture button, device
55216003 | Amnioscope, device
55337009 | Auscultoscope, device
55567004 | Bassinet, device
55658008 | Hot object
55986002 | Tong, device
56144002 | Back braces, device
56353002 | Staple, device
56547001 | Trephine, device

56896002 | Pacemaker catheter, device
56961003 | Cardiac transvenous pacemaker, device
57118008 | Perfusion pump, device
57134006 | Clinical instrument
57368009 | Contact lens
57395004 | Physical restraint equipment, device
57730005 | Abrasive grinding, device
58153004 | Android, device
58253008 | Suction catheter, device
58514003 | Infant scale, device
58878002 | Protective vest, device
58938008 | WC - Wheelchair
59102007 | Ice bag, device
59127000 | Apnea alarm, device
59153008 | Barge, device
59160002 | Chipguard, device
59181002 | Oxygen analyzer, device
59432006 | Ligature, device
59746007 | Needle holder, device
59772003 | Culdoscope, device
59782002 | Speculum, device
59833007 | Collapsible balloon, device
60054005 | SB - Seat belt
60110001 | Wig, device
60150003 | Skipole, device
60161006 | Acupuncture needle, device
60185003 | Carbon dioxide absorber, device
60311007 | Leather boots
60773001 | Injector
60806001 | Whirlpool bath
60957001 | Otoscope
61330002 | Nasopharyngeal airway device
61512008 | Tennis racket
61835000 | Dilator
61968008 | Syringe
61979003 | Antiembolic device
62336005 | Electric cable
62495008 | Gamma counter
62614002 | Overhead and gantry crane
62980002 | Tubular bandage
63112008 | Bone wire
63173005 | Hat band
63289001 | Metal nail
63336000 | Bone plug
63548003 | Derrick
63562005 | Cervical collar
63619003 | Fiberoptic cable
63653004 | Medical device
63797009 | Traction unit
63839002 | Electroejaculator
63995005 | Bandage
64174005 | Snowmobile
64255007 | Esophageal balloon
64565002 | Air tool
64571008 | Hair clipper
64883003 | Inhalation analgesia unit
64973003 | Scissors
64989000 | Escalator
65053001 | Electrical battery
65105002 | Surgical drapes
65268008 | Chart recorder
65473004 | Microscope
65577000 | X-ray shield
65818007 | Stent
66222000 | Hospital robot
66415006 | Audiometric testing equipment
66435007 | Electric bed
66494009 | Face cloth
67270000 | Hip prosthesis
67387001 | Coronary perfusion catheter
67670006 | Radiographic-tomographic unit
67777003 | Moving walk

67829007 | Esophagoscope
67920005 | Aerial lift
67966000 | Enema tube
68080007 | Radiographic unit
68181008 | Vibrating electric heating pad
68183006 | Bone screw
68276009 | Bottle
68325009 | Sound
68597009 | Support belt
68685003 | Household robot
68842005 | Gastroduodenoscope
69670004 | Patient utensil kit
69805005 | Insulin pump
69922008 | Tracheostomy button
70080007 | Bayonet
70300000 | Skull tongs
70303003 | Freezer
70453008 | Sled
70665002 | Blood pressure cuff
70793005 | Recreation equipment
70872004 | Wash basin
71384000 | Warmer
71483007 | Diving stage
71545009 | Humidifier
71601002 | Proctosigmoidoscope
71667001 | Bone wax
71948003 | Autoclave
72070000 | Ring
72188006 | Tissue expander
72302000 | Lead cable
72506001 | Implantable defibrillator
72742007 | Aspirator collection bottle
73027007 | Infant incubator
73534004 | Artificial skin
73562006 | Transilluminator
73571002 | Intravenous analgesia unit
73618007 | Power saw
73878004 | Hand saw
73985004 | Face protection in construction industry
74094004 | Belt
74108008 | Recorder
74300007 | Sanitary pad
74444006 | AL - Artificial limb
74566002 | Crutch
74884005 | Boatswain's chair
75075000 | Shoring and bracing (masonry and woodwork)
75187009 | Local anesthesia kit
75192006 | Arterial cannula
75751006 | Manual respirator
75780002 | Artificial kidney
75963008 | Skateboard
76091005 | Stainless steel wire suture
76123001 | Glass tube
76428000 | Elbow joint prosthesis
76433001 | Apron
76664007 | Artificial pancreas
76705002 | Applicator stick
76825006 | Abrasive wheel machinery
76937009 | Guillotine
77444004 | Pins
77541009 | Band
77720000 | Clips
77755003 | Chemical fiber cloth
77777004 | Bone staple
78279003 | Nail clipper
78498003 | Testicular prosthesis
78641001 | Nylon suture
78886001 | Electronic monitor
79051006 | Greenfield filter
79068005 | Needle
79218005 | Vehicle-mounted work platform
79287008 | Tampon

79401009 | Chute
79438009 | Foot protection
79481007 | Swing or sliding cut-off saw
79593001 | Transvenous electrode
79618001 | Storage tank
79811009 | Electric blanket
79834000 | Hickman line
79952001 | Swan-Ganz catheter, device
80278003 | Pediatric bed
80404000 | Chain fall
80617005 | Analysers
80664005 | Motor home
80853009 | Tendon hammer
80950008 | Oven
81293006 | Textile material
81317009 | Socket wrench
81320001 | Enzyme immunoassay analyzer
81826000 | All-terrain vehicle
81892008 | Radial saw
82379000 | Hemostat
82449006 | Peripheral intravenous catheter
82657000 | Bony tissue forceps
82830000 | Robotic arm
82879008 | Safety belt
83059008 | Tube
83315005 | Audio analgesia unit
83320005 | Dip tank
83369007 | Plastic shoes
83517001 | Robot
83903000 | Man lift
84023008 | Ski tow
84330009 | Pliers
84444002 | Chain saw
84546002 | Barricade
84599008 | Detonating cord
84610002 | Implantable dental prosthesis
84683006 | Aortic valve prosthesis
84756000 | Adhesive tape
85106006 | Boring machine
85329008 | Abortion pump
85455005 | Cart
85684007 | Engraving press
86056006 | Golf club
86174004 | Laparoscope
86184003 | Electrocardiographic monitor and recorder
86407004 | Table
86453006 | Defibrillator paddle
86572008 | Arteriovenous shunt catheter
86768006 | Balloon pump
86816008 | Diving ladder
86967005 | Tool
87088005 | Soldering iron
87140005 | Clothing material
87405001 | Cane
87710003 | Physical restraint structure
87717000 | Tester
87851008 | Blood cell counter and analyzer
88063004 | Footwear
88168006 | Maximum security cell
88208003 | Intravenous anesthesia administration set
88765001 | Artificial tissue
88959008 | Hypodermic needle
89149003 | Stretcher
89236003 | Leather shoes
89509004 | Blood culture analyzer
90003000 | Magnetic resonance imaging unit
90035000 | Alcohol sponge
90082007 | Cast cutter
90134004 | Metal periosteal implant
90412006 | Colonoscope
90504001 | Auricular prosthesis
90913005 | Rubber shoes

90948003 | Abrasive blast by cleaning nozzles
91294003 | Thomas collar
91318002 | Hyperbaric chamber
91535004 | Basin
91537007 | Hospital bed
102303004 | Vascular prosthesis
102304005 | Measuring ruler
102305006 | Intramedullary reamer
102306007 | Reamer
102307003 | Surgical knife
102308008 | Scalpel
102309000 | Surgical saw
102310005 | Gigli's wire saw
102311009 | Starck dilator
102312002 | Atherectomy device
102313007 | Rotational atherectomy device
102314001 | Embolization coil
102315000 | Embolization ball
102316004 | Embolization particulate
102317008 | Guiding catheter
102318003 | Implantable venous catheter
102319006 | Percutaneous transluminal angioplasty balloon
102320000 | Detachable balloon
102321001 | Operating microscope
102322008 | External prosthesis for sonographic procedure
102323003 | Water bag prosthesis for imaging procedure
102324009 | Saline bag prosthesis for imaging procedure
102325005 | Gel prosthesis for imaging procedure
102326006 | Dagger
102327002 | Dirk
102328007 | Sword
102384007 | Motor vehicle airbag
102385008 | Front airbag
102386009 | Front driver airbag
102387000 | Front passenger airbag
102388005 | Side airbag
102402008 | Snowboard
102403003 | Water ski
105784003 | Life support equipment
105785002 | Adhesive, bandage AND/OR suture
105787005 | Belt AND/OR binder
105788000 | Probe, sound, bougie AND/OR airway
105789008 | Cannula, tube AND/OR catheter
105790004 | Bag, balloon AND/OR bottle
105791000 | Pump, injector AND/OR aspirator
105792007 | Analgesia AND/OR anesthesia unit
105793002 | Monitor, alarm AND/OR stimulator
105794008 | Scope AND/OR camera
105809003 | Physical restraint equipment AND/OR structure
108874005 | Silicone plug
109184000 | Pregnancy testing kit
109226007 | Dental pin
109227003 | Hand joint prosthesis
109228008 | Knee joint prosthesis
111041008 | Artificial nails
111042001 | Artificial organ
111043006 | Medical tuning fork
111044000 | Bone tap
111045004 | Exerciser
111047007 | Urethral bougie
111048002 | Rhinoscope
111052002 | Protective breast plate
111060001 | Industrial sewing machine
111062009 | Food waste disposal equipment
115961006 | Soft Cast
115962004 | Fiberglass cast
116146000 | Blood product unit
116204000 | Catheter tip
116205004 | Blood bag
116206003 | Plasma bag
116250002 | Filter
116251003 | Wick

118294000 | Solid-state laser
118295004 | Gas laser
118296003 | Chemical laser
118297007 | Excimer laser
118298002 | Dye laser
118299005 | Diode laser
118301003 | Nd:YVO>4< laser
118302005 | Nd:YLF laser
118303000 | Nd:Glass laser
118304006 | Chromium sapphire laser device
118305007 | Er:Glass laser
118306008 | Erbium-YAG laser
118307004 | Ho:YLF laser
118308009 | Holmium-YAG laser
118309001 | Ti:sapphire laser device
118310006 | Alexandrite laser
118311005 | Argon laser
118312003 | CO2 laser
118313008 | He laser
118314002 | Helium cadmium laser
118315001 | HeNe laser
118316000 | Krypton laser
118317009 | Neon gas laser
118318004 | Nitrogen gas laser
118319007 | Xenon gas laser
118320001 | Copper vapor laser
118321002 | Gold vapor laser
118322009 | DF laser
118323004 | DF-CO>2< laser device
118324005 | HF laser
118325006 | ArF laser
118326007 | KrF laser
118327003 | KrCl laser
118328008 | XeCl laser
118329000 | XeFl laser
118330005 | Free electron laser
118331009 | Tunable dye laser
118332002 | Tunable dye argon laser
118333007 | Gallium arsenide laser
118334001 | Gallium aluminum arsenide laser
118335000 | Lead-salt laser
118336004 | Rhodamine 6G dye laser
118337008 | Coumarin C30 dye laser
118338003 | Coumarin 102 dye laser
118342000 | Diode pumped laser
118343005 | Flashlamp pumped laser device
118346002 | Pulsed dye laser
118347006 | QS laser
118348001 | Flashlamp pulsed dye laser
118349009 | CW CO>2< laser
118350009 | High energy pulsed CO>2< laser
118351008 | Frequency doubled Nd:YAG laser
118354000 | Continuous wave laser
118355004 | Pulsed laser
118356003 | Metal vapor laser
118357007 | KTP laser
118371004 | Ion laser
118372006 | Plastic implant
118373001 | Silastic implant
118374007 | Silicone implant
118375008 | Cardiac septum prosthesis
118376009 | Thermocouple
118377000 | Biopsy needle
118378005 | Pacemaker pulse generator
118379002 | Automatic implantable cardioverter sensing electrodes
118380004 | Implantable defibrillator leads
118381000 | Implantable cardioverter leads
118382007 | Neuropacemaker device
118383002 | External fixation device
118384008 | Long arm splint
118385009 | Short arm splint
118386005 | Figure of eight plaster cast

118387001 | Halo jacket
118388006 | Body cast, shoulder to hips
118389003 | Body cast, shoulder to hips including head, Minerva type
118390007 | Body cast, shoulder to hips including one thigh
118391006 | Body cast, shoulder to hips including both thighs
118392004 | Shoulder cast
118393009 | Long arm cylinder
118394003 | Forearm cylinder
118396001 | Cylinder cast, thigh to ankle
118397005 | Long leg cast
118398000 | Long leg cast, walker or ambulatory type
118399008 | Long leg cast, brace type
118400001 | Short leg cast below knee to toes
118401002 | Short leg cast below knee to toes, walking or ambulatory type
118402009 | Clubfoot cast
118403004 | Clubfoot cast, short leg
118404005 | Clubfoot cast, long leg
118405006 | Spica cast
118406007 | Hip spica cast, both legs
118407003 | Hip spica cast, one leg
118408008 | Hip spica cast, one and one-half spica
118409000 | Patellar tendon bearing cast
118410005 | Boot cast
118411009 | Sugar tong cast
118412002 | Gauntlet cast
118413007 | Complete cast
118414001 | Pressure dressing
118415000 | Packing material
118416004 | Wound packing material
118418003 | Trocar
118419006 | Umbrella device
118420000 | Atrial septal umbrella
118421001 | King-Mills umbrella device
118422008 | Mobitz-Uddin umbrella device
118423003 | Rashkind umbrella device
118424009 | Reservoir device
118425005 | Ventricular reservoir
118426006 | Ommaya reservoir
118427002 | Rickham reservoir
118428007 | Flexible fiberoptic endoscope
118429004 | Flexible fiberoptic laryngoscope with strobe
118643004 | Cast
122456005 | Laser
123636009 | SS - Silk suture
126064005 | Gastrostomy tube, device
126065006 | Jejunostomy tube, device
128981007 | Baffle
129113006 | Intra-aortic balloon pump
129121000 | Tracheostomy tube
129247000 | Fine biopsy needle
129248005 | Core biopsy needle
129460009 | Compression paddle
129462001 | Catheter guide wire
129463006 | J wire
129464000 | Medical administrative equipment
129465004 | Medical record
129466003 | Patient chart
129467007 | Identification plate
134823007 | Sterile absorbent dressing pad
134963007 | Wound drainage pouch dressing
170615005 | Home nebulizer
182562006 | Plaster jacket
182563001 | Shoulder spica
182564007 | Humeral U-slab
182565008 | Long arm slab
182566009 | Humeral hanging slab
182567000 | Forearm slab
182568005 | Scaphoid cast
182569002 | Bennett cast
182570001 | Hip spica
182571002 | Long leg spica
182572009 | Below knee non-weight-bearing cast

182573004 | Below knee weight-bearing cast
182574005 | Plaster stripper
182576007 | Humeral brace
182577003 | Functional elbow brace
182578008 | Forearm brace
182579000 | Hip brace
182580002 | Femoral brace
182581003 | Tibial brace
182587004 | Body support
182588009 | Spinal frame
182589001 | Corset support
182590005 | Cardiac bed
182591009 | Water bed
182592002 | High air loss bed
182839003 | Automated drug microinjector
183116000 | Dental aid
183125006 | Ear fitting hearing aid
183135000 | Mobility aid
183141007 | Inshoe orthosis
183143005 | Surgical stockings
183144004 | Antiembolic stockings
183146002 | ZF - Zimmer frame
183147006 | Tripod
183148001 | RGO - Reciprocating gait orthosis
183149009 | Hip guidance orthosis
183150009 | Standing frame
183152001 | Hip abduction orthosis
183153006 | Hip-knee-ankle-foot orthosis
183154000 | Knee-ankle-foot orthosis
183155004 | Flexible knee support
183156003 | Collateral ligament brace
183157007 | Anterior cruciate ligament brace
183158002 | Posterior cruciate ligament brace
183159005 | Ground reaction orthosis
183160000 | Rigid ankle-foot orthosis
183161001 | Flexible ankle-foot orthosis
183162008 | Double below-knee iron
183164009 | Inside iron
183165005 | Outside iron
183166006 | Inside T-strap
183170003 | Hindquarter prosthesis
183171004 | Hip disarticulation prosthesis
183172006 | Above knee prosthesis
183173001 | Through knee prosthesis
183174007 | Below knee prosthesis
183175008 | Syme's prosthesis
183176009 | Midfoot amputation prosthesis
183177000 | Shoe filler
183183002 | Milwaukee brace
183184008 | Boston brace
183185009 | Jewett brace
183187001 | Halo device
183188006 | Four poster brace
183189003 | Rigid collar
183190007 | Flexible collar
183192004 | Shoulder abduction brace
183193009 | Elbow brace
183194003 | Passive wrist extension splint
183195002 | Active wrist extension splint
183196001 | Passive finger extension splint
183197005 | Active finger extension splint
183198000 | Kleinert traction
183199008 | Passive thumb splint
183200006 | Active thumb splint
183202003 | Shin splint
183204002 | Excretory control aid
183235008 | Facial non-surgical prosthesis
183236009 | Breast non-surgical prosthesis
183240000 | Patient-propelled wheelchair
183241001 | Pedal powered wheelchair
183248007 | Attendant powered wheelchair
183249004 | Wheelchair seating

183250004 | Molded wheelchair seat
183251000 | Matrix seat
201706006 | Intracranial pressure transducer
223394001 | Equipment for positioning
224684009 | Top security prison
224685005 | Category B prison
224686006 | Low security prison
224823002 | Street lighting
224824008 | Sign posting
224825009 | Street name sign
224826005 | Building name sign
224827001 | Pedestrian direction sign
224828006 | Traffic sign
224898003 | Orthotic device
224899006 | Walking aid
224900001 | Communication aid
228167008 | Corset
228235002 | Slippers
228236001 | Mules
228237005 | Slippersox
228239008 | Trainers
228240005 | Plimsolls
228241009 | Sandals
228242002 | Gum boots
228243007 | Chappel
228259007 | Fastening
228260002 | Velcro
228261003 | Buckle
228262005 | Zipper
228264006 | Small button
228265007 | Medium button
228266008 | Large button
228267004 | Press stud
228268009 | Hook and eye
228270000 | Laces
228271001 | Shoe laces
228731007 | Radiotherapy equipment and appliances
228732000 | Beam direction shell
228733005 | Head and neck beam direction shell
228734004 | Body beam direction shell
228735003 | Beam modifier
228736002 | Surface bolus
228737006 | Surface compensator
228738001 | Cutout
228739009 | Shielding block
228740006 | Lung block
228741005 | Humerus block
228742003 | Scrotal block
228743008 | Kidney block
228744002 | Eye block
228745001 | Bite block
228746000 | Wedge filter
228747009 | Kilovoltage grid
228748004 | Brachytherapy implant
228749007 | Single plane implant
228750007 | Two plane implant
228751006 | Semicircular implant
228752004 | Regular volume implant
228753009 | Irregular volume implant
228754003 | Brachytherapy surface mold
228755002 | Two plane mold
228756001 | Single plane mold
228757005 | Cylinder mold
228759008 | Adhesive felt mold
228760003 | Elastoplast mold
228761004 | Collimator
228762006 | Multileaf collimator
228763001 | Asymmetric jaws collimator
228765008 | Standard collimator
228766009 | Form of brachytherapy source
228767000 | Wire source
228768005 | Seeds source

228770001 | Hairpins source
228771002 | Needles source
228772009 | Pellets source
228773004 | Capsules source
228774005 | Chains source
228775006 | Tubes source
228776007 | Rods source
228777003 | Grains source
228778008 | Plaque source
228869008 | Manual wheelchair
229772003 | Bed
229839006 | Functional foot orthosis
229840008 | Non-functional foot orthosis
229841007 | Detachable pad for the foot
229842000 | Detachable toe prop
229843005 | Detachable horseshoe pad
243135003 | Spacer
243719003 | Near low vision aid - integral eyeglass magnifier
243720009 | Near low vision aid - clip-on eyeglass magnifier
243722001 | Near low vision aid - integral eyeglass telescope
243723006 | Near low vision aid - clip-on eyeglass telescope
255296002 | Wedge
255712000 | Television
255716002 | Latex rubber gloves
256245006 | Textiles
256246007 | Cotton - textile
256247003 | Flax
256562002 | Cotton wool
256563007 | Cotton wool roll
256564001 | Cotton wool pledget
256589007 | Dental rubber dam
256590003 | Endodontic sponge
256593001 | Orthodontic elastic
256641009 | Ribbon gauze
256642002 | Wet ribbon gauze
256643007 | Dry ribbon gauze
257192006 | Aid to vision
257193001 | Telescopic eyeglasses
257194007 | Video
257211007 | Cylinder cutter
257212000 | Rotary cutter
257213005 | Rotary cutter with steel blades
257214004 | Rotary cutter with plastic blades
257215003 | Nail instrument
257216002 | Flexible endoscope
257217006 | Rigid endoscope
257218001 | Flexible cystoscope
257219009 | Rigid cystoscope
257220003 | Hysteroscope
257221004 | Flexible hysteroscope
257222006 | Rigid hysteroscope
257223001 | Contact hysteroscope
257224007 | Panoramic hysteroscope
257225008 | Flexible bronchoscope
257226009 | Rigid bronchoscope
257227000 | Standard laryngoscope
257228005 | Fiberlight anesthetic laryngoscope
257229002 | Pharyngeal mirror
257230007 | Obstetric forceps
257231006 | Barnes forceps

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information

Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)

Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')

Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application

Definition | An explanation of the meaning of the concept

Comments | Additional notes about how to use the code

4.4.1.545 Value Set http://hl7.org/fhir/ValueSet/device-status

URL: https://hl7.org/fhir/R4/valueset-device-status.html
Orders and Observations Work Group | Maturity Level: 2 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/device-status
Version: | 4.0.1
Name: | FHIRDeviceStatus
Title: | FHIRDeviceStatus
Definition: | The availability status of the device.
Committee: | Orders and Observations Work Group
OID: | 2.16.840.1.113883.4.642.3.209 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for FHIRDeviceStatus
• ConceptMap: Translation source in Canonical Mapping for "FHIRDeviceStatus"
• Resource: Device.status (code / Required)

4.4.1.545.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/device-status

4.4.1.545.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 4 concepts

Expansion based on http://hl7.org/fhir/device-status version 4.0.1

All codes from system http://hl7.org/fhir/device-status

Code | Display | Definition | Canonical Status
active | Active | The device is available for use. Note: For *implanted devices* this means that the device is implanted in the patient. | ~active
inactive | Inactive | The device is no longer available for use (e.g. lost, expired, damaged). Note: For *implanted devices* this means that the device has been removed from the patient. | ~inactive
entered-in-error | Entered in Error | The device was entered in error and voided. | ~error
unknown | Unknown | The status of the device has not been determined. | ~unknown

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.546 Value Set http://hl7.org/fhir/ValueSet/event-capability-mode

URL: https://hl7.org/fhir/R4/valueset-event-capability-mode.html
FHIR Infrastructure Work Group | Maturity Level: N | Normative (from v4.0.0) | Use Context: Any

| This page has been approved as part of an ANSI standard. See the Conformance Package for further details.

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/event-capability-mode
Version: | 4.0.1
Name: | EventCapabilityMode
Title: | EventCapabilityMode
Definition: | The mode of a message capability statement.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.182 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

- CodeSystem: This value set is the designated 'entire code system' value set for EventCapabilityMode
- Resource: CapabilityStatement.messaging.supportedMessage.mode (code / Required)

4.4.1.546.1 Content Logical Definition

This value set includes codes from the following code systems:

- Include all codes defined in http://hl7.org/fhir/event-capability-mode

4.4.1.546.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 2 concepts

Expansion based on http://hl7.org/fhir/event-capability-mode version 4.0.1

All codes from system http://hl7.org/fhir/event-capability-mode

Code | Display | Definition
sender | Sender | The application sends requests and receives responses.
receiver | Receiver | The application receives requests and sends responses.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.547 Value Set http://hl7.org/fhir/ValueSet/subject-type

URL: https://hl7.org/fhir/R4/valueset-subject-type.html
FHIR Infrastructure Work Group | Maturity Level: 5 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/subject-type
Version: | 4.0.1
Name: | SubjectType
Title: | Subject type
Definition: | Possible types of subjects.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.775 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

- Resource: EventDefinition.subject[x] (CodeableConcept|Reference(Group) / Extensible)
- Resource: Measure.subject[x] (CodeableConcept|Reference(Group) / Extensible)
- Profile: ShareableMeasure: Measure.subject[x] (CodeableConcept, Reference / Extensible)
- Resource: ResearchDefinition.subject[x] (CodeableConcept|Reference(Group) / Extensible)
- Resource: ResearchElementDefinition.subject[x] (CodeableConcept|Reference(Group) / Extensible)
- Resource: ActivityDefinition.subject[x] (CodeableConcept|Reference(Group) / Extensible)
- Profile: ShareableActivityDefinition: ActivityDefinition.subject[x] (CodeableConcept, Reference / Extensible)
- Resource: Library.subject[x] (CodeableConcept|Reference(Group) / Extensible)
- Profile: ShareableLibrary: Library.subject[x] (CodeableConcept, Reference / Extensible)
- Profile: CQLLibrary: Library.subject[x] (CodeableConcept, Reference / Extensible)
- Resource: PlanDefinition.subject[x] (CodeableConcept|Reference(Group) / Extensible)
- Resource: PlanDefinition.action.subject[x] (CodeableConcept|Reference(Group) / Extensible)
- Profile: ShareablePlanDefinition: PlanDefinition.subject[x] (CodeableConcept, Reference / Extensible)
- Profile: ShareablePlanDefinition: PlanDefinition.action.subject[x] (CodeableConcept, Reference / Extensible)
- Profile: ComputablePlanDefinition: PlanDefinition.subject[x] (CodeableConcept, Reference / Extensible)

• Profile: ComputablePlanDefinition: PlanDefinition.action.subject[x] (CodeableConcept, Reference / Extensible)

• Profile: CDSHooksServicePlanDefinition: PlanDefinition.subject[x] (CodeableConcept, Reference / Extensible)

• Profile: CDSHooksServicePlanDefinition: PlanDefinition.action.subject[x] (CodeableConcept, Reference / Extensible)

• Resource: DataRequirement.subject[x] (CodeableConcept|Reference(Group) / Extensible)

4.4.1.547.1 Content Logical Definition

This value set includes codes from the following code systems:

http://hl7.org/fhir/resource-types

Code | Display
Patient | Patient | Demographics and other administrative information about an individual or animal receiving care or other health-related services.
Practitioner | Practitioner | A person who is directly or indirectly involved in the provisioning of healthcare.
Organization | Organization | A formally or informally recognized grouping of people or organizations formed for the purpose of achieving some form of collective action. Includes companies, institutions, corporations, departments, community groups, healthcare practice groups, payer/insurer, etc.
Location | Location | Details and position information for a physical place where services are provided and resources and participants may be stored, found, contained, or accommodated.
Device | Device | A type of a manufactured item that is used in the provision of healthcare without being substantially changed through that activity. The device may be a medical or non-medical device.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.548 Value Set http://hl7.org/fhir/ValueSet/remittance-outcome

URL: https://hl7.org/fhir/R4/valueset-remittance-outcome.html
Financial Management Work Group | Maturity Level: 2 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/remittance-outcome
Version: | 4.0.1
Name: | ClaimProcessingCodes
Title: | Claim Processing Codes
Definition: | This value set includes Claim Processing Outcome codes.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.13 (for OID based terminology systems)
Copyright: | HL7 Inc.
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ClaimProcessingCodes

• Resource: ExplanationOfBenefit.outcome (code / Required)

• Resource: ClaimResponse.outcome (code / Required)

4.4.1.548.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/remittance-outcome

4.4.1.548.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 4 concepts

Expansion based on http://hl7.org/fhir/remittance-outcome version 4.0.1

All codes from system http://hl7.org/fhir/remittance-outcome

Code | Display | Definition
queued | Queued | The Claim/Pre-authorization/Pre-determination has been received but processing has not begun.
complete | Processing Complete | The processing has completed without errors
error | Error | One or more errors have been detected in the Claim
partial | Partial Processing | No errors have been detected in the Claim and some of the adjudication has been performed.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.549 Value Set http://hl7.org/fhir/ValueSet/issue-severity

URL: https://hl7.org/fhir/R4/valueset-issue-severity.html
FHIR Infrastructure Work Group | Maturity Level: N | Normative (from v4.0.0) | Use Context: Any

| This page has been approved as part of an ANSI standard. See the Infrastructure Package for further details.

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/issue-severity
Version: | 4.0.1
Name: | IssueSeverity
Title: | IssueSeverity
Definition: | How the issue affects the success of the action.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.408 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for IssueSeverity
• Resource: OperationOutcome.issue.severity (code / Required)

4.4.1.549.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/issue-severity

4.4.1.549.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 4 concepts

Expansion based on http://hl7.org/fhir/issue-severity version 4.0.1

All codes from system http://hl7.org/fhir/issue-severity

Code | Display | Definition
fatal | Fatal | The issue caused the action to fail and no further checking could be performed.
error | Error | The issue is sufficiently important to cause the action to fail.
warning | Warning | The issue is not important enough to cause the action to fail but may cause it to be performed suboptimally or in a way that is not as desired.
information | Information | The issue has no relation to the degree of success of the action.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.550 Value Set http://hl7.org/fhir/ValueSet/doc-typecodes

URL: https://hl7.org/fhir/R4/valueset-doc-typecodes.html
Structured Documents Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined at http://hl7.org .

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/doc-typecodes
Version: | 4.0.1
Name: | FHIRDocumentTypeCodes
Title: | FHIR Document Type Codes
Definition: | FHIR Document Codes - all LOINC codes where scale type = 'DOC'.
Committee: | Structured Documents Work Group
OID: | 2.16.840.1.113883.4.642.3.240 (for OID based terminology systems)
Copyright: | This content from LOINC® is copyright © 1995 Regenstrief Institute, Inc. and the LOINC Committee, and available at no cost under the license at http://loinc.org/terms-of-use.
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: Composition.type (CodeableConcept / Preferred)

• Profile: ClinicalDocument: Composition.type (CodeableConcept / Preferred)

• Profile: Catalog: Composition.type (CodeableConcept / Preferred)

4.4.1.550.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include codes from http://loinc.org where SCALE_TYP = Doc

4.4.1.550.2 Expansion

This expansion generated 01 Nov 2019

This value set has >1000 codes in it. In order to keep the publication size manageable, only a selection (1000 codes) of the whole set of codes is shown

All codes from system http://loinc.org

Code | Display
11485-0 | Anesthesia records
11486-8 | Chemotherapy records
11488-4 | Consult note
11490-0 | Physician Discharge summary
11492-6 | Provider-unspecifed, History and physical note
11494-2 | Deprecated Physician Initial assessment note at First encounter
11495-9 | Deprecated Physical therapy Initial assessment note at First encounter
11496-7 | Deprecated Podiatry Initial assessment note at First encounter
11497-5 | Deprecated Psychology Initial assessment note at First encounter
11498-3 | Deprecated Social work Initial assessment note at First encounter
11499-1 | Deprecated Speech therapy Initial assessment note at First encounter
11500-6 | Deprecated Occupational therapy Initial assessment note at First encounter
11502-2 | Laboratory report
11503-0 | Medical records
11504-8 | Surgical operation note
11505-5 | Physician procedure note
11506-3 | Progress note
11507-1 | Occupational therapy Progress note
11508-9 | Physical therapy Progress note
11509-7 | Podiatry Progress note
11510-5 | Psychology Progress note
11512-1 | Speech-language pathology Progress note
11514-7 | Chiropractic Records total Encounter
11515-4 | Physical therapy Records total Encounter
11516-2 | Physician Records total Encounter
11517-0 | Podiatry Records total Encounter
11518-8 | Psychology Records total Encounter
11519-6 | Social service Records total Encounter
11520-4 | Speech therapy Records total Encounter
11521-2 | Occupational therapy Records total Encounter
11522-0 | Deprecated Cardiac echo study
11523-8 | EEG study
11524-6 | EKG Study
11525-3 | US Pelvis Fetus for pregnancy
11526-1 | Pathology study
11527-9 | Psychiatry study
11528-7 | Deprecated Radiology Unspecified site and modality Study
11529-5 | Surgical pathology study

11534-5 | Temperature charts
11536-0 | Nurse Notes
11538-6 | Deprecated CT Chest study
11539-4 | Deprecated CT Head study
11540-2 | Deprecated CT Abdomen study
11541-0 | Deprecated MRI Brain study
11542-8 | Deprecated Provider-unspecified visit note
11543-6 | Nursery records
15507-7 | Emergency department Progress note
15508-5 | Labor and delivery records
17787-3 | Deprecated NM Thyroid gland Study report
18762-5 | Deprecated Chiropractor Progress note
18763-3 | Physician consulting Initial evaluation note
18764-1 | Deprecated Nurse practitioner Progress note
18765-8 | Deprecated Podiatry visit note
18766-6 | Deprecated Psychology visit note
18841-7 | Hospital consultations Document
18842-5 | Discharge summary
18594-2 | Psychiatric service attachment
18733-6 | Physician attending Progress note
18734-4 | Occupational therapy Initial evaluation note
18735-1 | Physical therapy Initial evaluation note
18736-9 | Physician Initial evaluation note
18737-7 | Podiatry Initial evaluation note
18738-5 | Psychology Initial evaluation note
18739-3 | Social work Initial evaluation note
18740-1 | Speech-language pathology Initial evaluation note
18741-9 | Deprecated Attending physician progress note
18742-7 | Arthroscopy study
18743-5 | Autopsy report
18744-3 | Bronchoscopy study
18745-0 | Cardiac catheterization study
18746-8 | Colonoscopy Study
18747-6 | Deprecated CT Unspecified system Study
18748-4 | Diagnostic imaging study
18749-2 | Electromyogram study
18750-0 | Electrophysiology study
18751-8 | Endoscopy study
18752-6 | Exercise stress test study
18753-4 | Flexible sigmoidoscopy study
18754-2 | Holter monitor study
18755-9 | Deprecated MRI Unspecified site study
18756-7 | MR Spine study
18757-5 | Deprecated Nuclear medicine study
18758-3 | Deprecated PET scan Unspecified system Study
18759-1 | Spirometry study
18760-9 | Deprecated US Unspecified system Study
18761-7 | Transfer summary note
18776-5 | Plan of care note
18823-5 | Alcohol and/or substance abuse service attachment
18824-3 | Cardiac service attachment
18825-0 | Medical social services attachment
18826-8 | Occupational therapy service attachment
18836-7 | Cardiac stress study Procedure
19002-5 | Physical therapy service attachment
19003-3 | Respiratory therapy service attachment
19004-1 | Skilled nursing service attachment
24600-9 | US Guidance for needle localization of Breast
24601-7 | US Breast
24602-5 | MG Guidance for biopsy of Breast
24603-3 | MG stereo Guidance for biopsy of Breast
24604-1 | MG Breast Diagnostic Limited Views
25054-8 | CT Guidance for radiosurgery of Unspecified body region-- W contrast IV
25055-5 | Deprecated Unspecified body region MR additional sequence
25056-3 | MR Unspecified body region
25057-1 | MR Unspecified body region W conscious sedation
24673-6 | Deprecated Duodenum Radio fluoroscopy W contrast PO and hypotonic agent per ng
24674-4 | MR Elbow
24675-1 | MR Elbow WO and W contrast IV
24676-9 | XR Elbow Views
24677-7 | US Pelvis transvaginal
24678-5 | RF Esophagus Views W contrast PO

24787-4 | XR tomography Kidney - bilateral WO contrast and 10M post contrast IV
24788-2 | XR Kidney - bilateral Views W contrast IV
24789-0 | XR tomography Kidney - bilateral
24790-8 | XR tomography Kidney - bilateral W contrast IV
24840-1 | MR Neck WO and W contrast IV
24841-9 | MR Neck W contrast IV
24842-7 | US Neck
24843-5 | XR Neck Lateral
24844-3 | MRA Neck vessels W contrast IV
24965-6 | CT Lumbar spine W contrast IT
24967-2 | MR Lumbar spine WO and W contrast IV
24968-0 | MR Lumbar spine
24969-8 | XR Lumbar spine Lateral
25022-5 | RF Uterus and Fallopian tubes Views W contrast IU
25023-3 | RFA Vein Views W contrast IV
25024-1 | RFA Guidance for placement of peripherally-inserted central venous catheter in Vein
24531-6 | US Retroperitoneum
24532-4 | US Abdomen RUQ
24533-2 | MRA Abdominal vessels W contrast IV
24534-0 | US.doppler Abdominal vessels
24535-7 | XR Acetabulum Views
24536-5 | XR Acromioclavicular Joint Views
24537-3 | US Guidance for aspiration of amniotic fluid of Uterus
24538-1 | MR Ankle
24539-9 | MR Ankle WO and W contrast IV
24540-7 | XR Ankle 2 Views
24541-5 | XR Ankle Views
24542-3 | US Anus
24543-1 | RFA Guidance for angioplasty of Thoracic and abdominal aorta-- W contrast IA
24544-9 | CT Thoracic Aorta
24545-6 | CT Thoracic Aorta W contrast IV
24546-4 | Deprecated RFA Aortic arch W contrast IA
24547-2 | US Thoracic and abdominal aorta
24548-0 | US Appendix
24549-8 | MRA Upper extremity vessels W contrast IV
24550-6 | RFA Upper extremity veins Views W contrast IV
24551-4 | RFA AV fistula Views W contrast IA
24552-2 | RF Stent Views W contrast intra stent
24553-0 | RFA Guidance for embolectomy of Vessel intracranial-- W contrast IV
24554-8 | RFA Guidance for embolization of Artery-- W contrast IA
24555-5 | RFA Guidance for placement of stent in Artery
24556-3 | MR Abdomen
24557-1 | MR Abdomen WO and W contrast IV
24558-9 | US Abdomen
24559-7 | US Guidance for drainage of Abdomen
24560-5 | Portable XR Abdomen AP left lateral-decubitus
24561-3 | XR Abdomen AP left lateral-decubitus
24562-1 | XR Abdomen Left lateral-decubitus and Right lateral-decubitus
24563-9 | XR Abdomen AP right lateral-decubitus
24564-7 | Portable XR Abdomen AP upright
24566-2 | CT Retroperitoneum
24567-0 | Deprecated Abdomen>Retroperitoneum CT W contrast
24568-8 | RFA Guidance for atherectomy of AV fistula-- W contrast IV
24569-6 | RFA AV shunt Views W contrast IV
24570-4 | RF Guidance for removal of calculus from Biliary duct common-- W contrast retrograde intra biliary
24571-2 | NM Biliary ducts and Gallbladder Views for patency of biliary structures and ejection fraction W sincalide and W radionuclide IV
24572-0 | NM Biliary ducts and Gallbladder Views for patency of biliary structures W Tc-99m IV
24573-8 | XR Biliary ducts and Gallbladder Views W contrast IV
24574-6 | RF Biliary ducts and Gallbladder Views during surgery W contrast biliary duct
24575-3 | RF Biliary ducts and Gallbladder Views W contrast percutaneous transhepatic
24576-1 | RFA Urinary bladder arteries Views W contrast IA
24577-9 | XR Bone Views during surgery
24578-7 | SPECT Bones
24579-5 | XR Long bones Survey Views
24580-3 | RFA Guidance for angioplasty of Brachiocephalic artery-- W contrast IA
24581-1 | RFA Subclavian artery and Brachial artery Views W contrast IA
24582-9 | MR Brachial plexus
24583-7 | MR Brachial plexus WO and W contrast IV
24584-5 | MRA Thoracic inlet vessels W contrast IV
24585-2 | CT Guidance for stereotactic biopsy of Head-- W contrast IV
24586-0 | MR Brain W anesthesia

24587-8 | MR Brain WO and W contrast IV
24588-6 | MR Brain WO and W contrast IV and W anesthesia
24589-4 | MR Brain W contrast IV
24590-2 | MR Brain
24591-0 | NM Brain Brain death protocol Views W Tc-99m HMPAO IV
24593-6 | MRA Head vessels W contrast IV
24761-9 | XR Hip Single view
24762-7 | XR Hip Views
24764-3 | RF Hip Arthrogram
24765-0 | XR Humerus 2 Views
24766-8 | RFA Guidance for angioplasty of Iliac artery-- W contrast IA
24767-6 | XR tomography Internal auditory canal
24769-2 | CT Guidance for injection of Joint space
24770-0 | NM Joint Views W In-111 Intrajoint
24943-3 | XR Cervical spine Lateral
24944-1 | XR Cervical spine Swimmers
24945-8 | XR Cervical spine Views W flexion and W extension
24946-6 | XR Cervical spine Views
24947-4 | RF Cervical spine Views W contrast IT
24948-2 | XR Spine Cervical Odontoid and Cervical axis AP
24963-1 | CT Lumbar spine
24964-9 | CT Lumbar spine W contrast IV
25061-3 | US Unspecified body region
25062-1 | XR Unspecified body region Comparison view
25063-9 | RFA Vessel Single view W contrast IA
25064-7 | RFA Vessel Views for angioplasty W contrast IA
25065-4 | RF 15 minutes
25066-2 | RF 30 minutes
25067-0 | RF 45 minutes
25068-8 | RF 1 hour
25069-6 | RF Guidance for biopsy of Unspecified body region
24594-4 | MG Guidance for aspiration of cyst of Breast
24595-1 | MG Guidance for needle localization of mass of Breast
24596-9 | US Breast specimen
24597-7 | MG Breast specimen Views
24598-5 | MG Guidance for fluid aspiration of Breast
24599-3 | US Breast limited
24605-8 | MG Breast Diagnostic
24606-6 | MG Breast Screening
24609-0 | MG Guidance for percutaneous biopsy.core needle of Breast
24610-8 | MG Breast Limited Views
24611-6 | Outpatient Consultation 2nd opinion
24612-4 | XR Calcaneus Views
24613-2 | Deprecated RFA Guidance for placement of catheter in artery in Central cardiovascular artery
24614-0 | RFA Guidance for angioplasty of Carotid artery extracranial-- W contrast IA
24615-7 | RFA Guidance for angioplasty of Carotid artery.intracranial-- W contrast IA
24616-5 | US Carotid arteries
24617-3 | RFA Carotid arteries Views W contrast IA
24619-9 | XR Wrist Views
24620-7 | RF Catheter Views for patency check W contrast via catheter
24621-5 | RF Guidance for percutaneous drainage of Unspecified body region
24622-3 | RFA Celiac artery Views W contrast IA
24623-1 | CT Guidance for nerve block of Celiac plexus
24624-9 | RFA Guidance for change of CV catheter in Vein-- W contrast IV
24625-6 | RFA Guidance for placement of CV catheter in Vein-- W contrast IV
24626-4 | RFA Guidance for reposition of CV catheter in Vein-- W contrast IV
24627-2 | CT Chest
24628-0 | CT Chest W contrast IV
24629-8 | MR Chest
24630-6 | US Chest
24631-4 | RF Unspecified body region Views for central venous catheter placement check
24632-2 | Portable XR Chest Views AP
24634-8 | Portable XR Chest Views W inspiration and expiration
24635-5 | XR Chest PA upright Views W inspiration and expiration
24636-3 | Portable XR Chest AP left lateral-decubitus
24637-1 | XR Chest AP left lateral-decubitus
24638-9 | Portable XR Chest Left lateral upright
24639-7 | XR Chest Left lateral upright
24640-5 | XR Chest Apical lordotic
24641-3 | Portable XR Chest Left oblique
24642-1 | XR Chest AP and PA upright
24643-9 | XR Chest PA and Lateral and Oblique upright

24644-7 | Portable XR Chest PA and Lateral upright
24645-4 | Portable XR Chest PA and Right lateral and Right oblique and Left oblique upright
24646-2 | XR Chest PA and Right lateral and Right oblique and Left oblique upright
24647-0 | XR Chest PA and Lateral upright
24648-8 | XR Chest PA upright
24649-6 | Portable XR Chest Right lateral-decubitus and Left lateral-decubitus
24650-4 | XR Chest Right lateral-decubitus and Left lateral-decubitus
24651-2 | XR Chest Right oblique and Left oblique upright
24652-0 | Portable XR Chest AP right lateral-decubitus
24653-8 | XR Chest AP and AP right lateral-decubitus
24654-6 | Portable XR Chest AP and AP right lateral-decubitus
24655-3 | RF Chest Image intensifier during surgery
24656-1 | RF Chest Single view during surgery
24657-9 | XR tomography Chest
24658-7 | RFA Thoracic and abdominal aorta Views W contrast IA
24659-5 | MRA Chest vessels W contrast IV
24660-3 | MRA Thoracic Aorta
24661-1 | RF Pleural space Views W contrast intra pleural space
24662-9 | US Guidance for fluid aspiration of Pleural space
24663-7 | NM Cerebral cisterns Views W radionuclide IT
24664-5 | XR Clavicle Views
24665-2 | XR Sacrum and Coccyx Views
24666-0 | RF Colon Views W air and barium contrast PR
24667-8 | RF Colon Views W contrast PR
24668-6 | RF Colon Single view for transit Post solid contrast
24669-4 | RF Colon Views W water soluble contrast PR
24670-2 | US Guidance for biopsy of cyst of Unspecified body region
24671-0 | RF Guidance for aspiration of cyst of Unspecified body region
24672-8 | US Diaphragm for motion
24679-3 | RF Esophagus Views W Gastrografin PO
24680-1 | RF Guidance for dilation of Esophagus
24681-9 | RF videography Hypopharynx and Esophagus Views
24682-7 | RF videography Hypopharynx and Esophagus Views W liquid and paste contrast PO during swallowing
24683-5 | NM Esophagus+Stomach Views W Tc-99m SC PO
24684-3 | RFA Guidance for embolectomy of Extracranial vessels-- W contrast IA
24685-0 | RFA Peripheral veins Views W contrast IV
24686-8 | XR Lower extremity Views
24687-6 | MR Lower Extremity Joint
24688-4 | MR Upper extremity
24689-2 | XR Upper extremity Views
24690-0 | CT Extremity
24691-8 | CT Extremity W contrast IV
24692-6 | US Guidance for drainage of Extremity
24693-4 | US Extremity
24694-2 | MR Face WO and W contrast IV
24695-9 | XR Facial bones Views
24696-7 | CT Facial bones
24697-5 | CT Facial bones W contrast IV
24698-3 | RFA Guidance for angioplasty of Femoral artery-- W contrast IA
24699-1 | RFA Femoral artery Runoff W contrast IA
24700-7 | XR Femur and Tibia Views for leg length
24702-3 | MR Thigh
24703-1 | MR Thigh WO and W contrast IV
24704-9 | XR Femur Views
24705-6 | MR Finger
24706-4 | XR Finger Views
24707-2 | MR Foot
24708-0 | XR Foot Views W standing
24709-8 | XR Foot Views
24710-6 | MR Forearm
24711-4 | US Gallbladder
24712-2 | XR Gallbladder Views W contrast PO
24713-0 | XR Gallbladder Views 48H post contrast PO
24714-8 | NM Gastrointestinal tract Views for gastrointestinal bleeding W Tc-99m tagged RBC IV
24715-5 | RF Gastrointestinal tract upper Single view W contrast PO
24716-3 | RF Guidance for placement of decompression tube in Gastrointestinal tract
24717-1 | Deprecated XR Ileal conduit Loopogram
24718-9 | RF Guidance for transjugular biopsy of Liver-- W contrast IV
24719-7 | US Groin
24720-5 | MR Hand
24721-3 | XR Hand 2 Views
24722-1 | XR Hand 3 Views

24723-9 | XR Hand Arthritis
24724-7 | XR Wrist and Hand Bone age Views
24725-4 | CT Head
24726-2 | CT Head WO and W contrast IV
24727-0 | CT Head W contrast IV
24728-8 | CT perfusion Head
24729-6 | CT perfusion Head WO and W contrast IV
24730-4 | NM Brain Views
24731-2 | US Head
24732-0 | US Head during surgery
24733-8 | US.doppler Head vessels
24734-6 | CT Cerebral cisterns W contrast IT
24735-3 | MR Internal auditory canal and Posterior fossa
24740-3 | MR Internal auditory canal and Posterior fossa WO and W contrast IV
24745-2 | XR Petrous part of temporal bone Views
24746-0 | Deprecated Head Sagittal Sinus MR
24747-8 | Deprecated Head Sagittal Sinus MR angiogram W contrast IV
24748-6 | MR Heart
24750-2 | NM Heart Views at rest and W Tl-201 IV
24751-0 | NM Parathyroid gland Views W Tl-201 subtraction Tc-99m IV
24752-8 | RF videography Heart Views
24753-6 | CT Unspecified body region W contrast IV
24754-4 | Administration of vasodilator into catheter Vein
24755-1 | RFA Guidance for atherectomy of Vein-- W contrast IV
24756-9 | RFA Guidance for placement of stent in Vein
24757-7 | Deprecated Coronary arteries CT fast
24760-1 | US Hip
24771-8 | RF Guidance for fluid aspiration of Joint space
24772-6 | US Guidance for biopsy of Kidney
24773-4 | NM Kidney Views W radionuclide transplant scan
24776-7 | NM Kidney Views
24778-3 | XR Kidney - bilateral 3 Serial Views WO and W contrast IV
24779-1 | RF Guidance for percutaneous placement of nephrostomy tube of Kidney - bilateral-- W contrast via tube
24780-9 | RF Kidney - bilateral Views W contrast via nephrostomy tube
24781-7 | RF Guidance for exchange of nephrostomy tube of Kidney - bilateral-- W contrast
24782-5 | RF Guidance for percutaneous placement of nephroureteral stent of Kidney - bilateral-- W contrast via stent
24783-3 | RF Kidney - bilateral Views for urodynamics
24784-1 | XR tomography Kidney - bilateral WO and W contrast IV
24792-4 | Portable XR Abdomen AP and AP left lateral-decubitus
24793-2 | Portable XR Abdomen AP and Lateral
24794-0 | XR Abdomen AP and Lateral
24795-7 | Portable XR Abdomen Supine and Upright
24796-5 | XR Abdomen AP and AP left lateral-decubitus
24797-3 | XR Abdomen AP and Oblique prone
24798-1 | XR Abdomen Supine and Upright
24799-9 | XR Abdomen AP
24800-5 | RF Knee Arthrogram
24801-3 | XR Knee Merchants
24802-1 | MR Knee
24803-9 | MR Knee WO and W contrast IV
24804-7 | NM Knee Views
24805-4 | XR Knee AP and Lateral W standing
24806-2 | XR Knee 2 Views
24807-0 | XR Knee AP W standing
24808-8 | XR Knee AP and PA W standing
24809-6 | XR Knee Views W standing
24811-2 | CT Guidance for fluid aspiration of Liver
24812-0 | CT Guidance for biopsy of Liver
24813-8 | CT Guidance for core needle biopsy of Liver
24814-6 | CT Liver
24815-3 | CT Liver W contrast IV
24816-1 | US Guidance for biopsy of Liver
24817-9 | SPECT Liver W Tc-99m IV
24818-7 | US Diaphragm and Liver
24819-5 | Deprecated Liver and Spleen NM W Tc-99m calcium colloid IV
24820-3 | MRA Lower leg vessels W contrast IV
24821-1 | MR Lower leg
24822-9 | CT Guidance for fluid aspiration of Lung
24823-7 | CT Guidance for biopsy of Lung
24824-5 | NM Lung Portable Views
24825-2 | XR Lung Views W contrast intrabronchial
24826-0 | NM Lymphatic vessels Views W radionuclide intra lymphatic

24827-8 | RFA Lymphatic vessels Views W contrast intra lymphatic
24828-6 | XR tomography Mandible Panoramic
24829-4 | XR Mandible Views
24830-2 | XR Mastoid Views
24831-0 | NM Small bowel Views for Meckel's diverticulum W Tc-99m M04 IV
24832-8 | RFA Guidance for angioplasty of Mesenteric artery-- W contrast IA
24833-6 | RFA Mesenteric artery Views W contrast IA
24834-4 | XR Nasal bones Views
24835-1 | CT Nasopharynx and Neck
24836-9 | CT Nasopharynx and Neck W contrast IV
24837-7 | CT Guidance for fluid aspiration of Neck
24838-5 | CT Guidance for biopsy of Neck
24839-3 | MR Neck
24845-0 | RF Neck Views W contrast intra larynx
24846-8 | XR Optic foramen Views
24848-4 | CT Orbit - bilateral
24849-2 | CT Orbit - bilateral WO and W contrast IV
24850-0 | CT Orbit - bilateral W contrast IV
24851-8 | MR Orbit - bilateral WO and W contrast IV
24852-6 | MR Orbit - bilateral W contrast IV
24853-4 | US Eye+Orbit - bilateral
24854-2 | XR Orbit - bilateral Views
24855-9 | RF videography Oropharynx Views
24856-7 | CT Guidance for fluid aspiration of Pancreas
24857-5 | CT Pancreas
24858-3 | CT Pancreas W contrast IV
24859-1 | US Pancreas
24860-9 | RFA Pancreatic artery Views W contrast IA
24861-7 | XR Patella 2 Views
24862-5 | RFA Iliac artery Internal Views W contrast IA
24863-3 | CT Guidance for fluid aspiration of Pelvis
24864-1 | CT Guidance for biopsy of Pelvis
24865-8 | CT Pelvis
24866-6 | CT Pelvis W contrast IV
24867-4 | MR Pelvis
24868-2 | US Guidance for drainage of Pelvis
24869-0 | US Pelvis
24870-8 | US.doppler Pelvis vessels
24871-6 | XR Pelvis Pelvimetry
24872-4 | MR Pelvis and Hip
24873-2 | MRA Pelvis vessels W contrast IV
24874-0 | RFA Peripheral arteries Views W contrast IA
24875-7 | Deprecated Peripheral vessel US.doppler Peripheral plane
24876-5 | NM Head to Pelvis Views for shunt patency W Tc-99m DTPA IT
24877-3 | CT Petrous part of temporal bone
24878-1 | CT Petrous part of temporal bone W contrast IV
24879-9 | MR Pituitary and Sella turcica WO and W contrast IV
24880-7 | MR Pituitary and Sella turcica
24881-5 | US Popliteal space
24882-3 | RFA Guidance for percutaneous transluminal angioplasty of Popliteal artery-- W contrast IA
24883-1 | US Guidance for percutaneous biopsy of Prostate
24884-9 | US Prostate transrectal
24885-6 | Deprecated US Guidance for repair of Pseudoaneurysm/AV fistula
24887-2 | RFA Guidance for embolectomy of Pulmonary arteries-- W contrast IA
24888-0 | NM Pulmonary system Ventilation and Perfusion W Xe-133 IH and W Tc-99m MAA IV
24889-8 | US Pylorus for Pyloric Stenosis
24891-4 | XR Radius and Ulna Views
24892-2 | US Rectum
24893-0 | RF Rectum Single view post contrast PR during defecation
24894-8 | RF Rectum and Urinary bladder Views W contrast PR and intra bladder during defecation and voiding
24896-3 | US Guidance for drainage of Kidney
24899-7 | XR Ribs Views
24900-3 | XR Sacroiliac Joint Views
24901-1 | CT Guidance for injection of Sacroiliac Joint
24902-9 | RF Salivary gland Views W contrast intra salivary duct
24903-7 | XR Scapula Views
24904-5 | CT Pituitary and Sella turcica WO and W contrast IV
24905-2 | MR Shoulder
24906-0 | MR Shoulder WO and W contrast IV
24907-8 | US Shoulder
24908-6 | XR Shoulder 3 Views
24909-4 | XR Shoulder Views

24910-2 | RF Shoulder Arthrogram
24911-0 | RF Shunt Views
24912-8 | RF Sinus tract Views W contrast intra sinus tract
24913-6 | CT Sinuses limited
24914-4 | MR Sinuses
24915-1 | MR Sinuses W contrast IV
24916-9 | XR Sinuses Views
24917-7 | XR Skull Single view
24918-5 | XR Skull 3 Views
24919-3 | XR Skull AP and Lateral
24920-1 | XR Skull Lateral
24921-9 | XR Skull Waters
24922-7 | XR Skull 5 Views
24923-5 | RF Small bowel Views W positive contrast via enteroclysis tube
24924-3 | RF Small bowel Views W contrast PO
24925-0 | RFA Spinal artery Views W contrast IA
24926-8 | US Spine
24927-6 | RF Spine Views W contrast intradisc
24928-4 | XR Spine AP and Lateral
24929-2 | XR Thoracic and lumbar spine Views for scoliosis W flexion and W extension
24930-0 | XR Thoracic and lumbar spine Views for scoliosis
24931-8 | RF Guidance for injection of Spine facet joint
24932-6 | CT Cervical spine
24933-4 | CT Cervical spine W contrast IV
24934-2 | CT Cervical spine W contrast IT
24935-9 | MR Cervical spine
24936-7 | MR Cervical spine W anesthesia
24937-5 | MR Cervical spine WO and W contrast IV
24938-3 | MR Cervical spine W contrast IV
24939-1 | XR Cervical spine 5 Views
24940-9 | XR Cervical spine Single view
24941-7 | XR Cervical spine 3 Views
24942-5 | XR Cervical spine AP and Lateral
24970-6 | XR Lumbar spine AP and Lateral
24971-4 | XR Lumbar spine Views W flexion and W extension
24972-2 | XR Lumbar spine Views
24973-0 | RF Guidance for fluid aspiration of Spine.lumbar space
24974-8 | RF Lumbar spine Views W contrast IT
24975-5 | XR Spine.lumbar and Sacroiliac joint - bilateral Views
24977-1 | MR Lumbar spine W anesthesia
24978-9 | CT Thoracic spine
24979-7 | CT Thoracic spine W contrast IV
24980-5 | MR Thoracic spine
24981-3 | MR Thoracic spine WO and W contrast IV
24982-1 | MR Thoracic spine W contrast IV
24983-9 | XR Thoracic spine Views
24984-7 | XR Thoracic and lumbar spine 2 Views
24985-4 | RF Thoracic spine Views W contrast IT
24986-2 | CT Guidance for biopsy of Spine
24987-0 | CT Spine W contrast IV
24988-8 | CT Spleen
24989-6 | CT Spleen WO and W contrast IV
24990-4 | US Spleen
24991-2 | RFA Splenic vein and Portal vein Views W contrast IA
24992-0 | RFA Splenic artery Views W contrast IA
24993-8 | Deprecated X-ray
24994-6 | XR Sternum Views
24995-3 | RF Guidance for placement of tube in Stomach
24996-1 | RF Guidance for percutaneous replacement of gastrostomy of Stomach
24997-9 | NM Stomach Views for gastric emptying solid phase W Tc-99m SC PO
24998-7 | RF Placement check of gastrostomy tube W contrast via GI tube
24999-5 | MR Temporomandibular joint
25000-1 | XR Temporomandibular joint Views
25001-9 | NM Scrotum and testicle Views W Tc-99m pertechnetate IV
25002-7 | US Scrotum and testicle
25003-5 | MRA Thigh vessels W contrast IV
25005-0 | RFA Three vessels Views W contrast
25006-8 | XR Thumb Views
25007-6 | NM Thyroid gland Views W I-131 IV
25008-4 | NM Thyroid gland Views and Views uptake W I-131 IV
25009-2 | US Guidance for biopsy of Thyroid gland
25010-0 | US Thyroid gland

25011-8 | XR Tibia and Fibula Views
25012-6 | Deprecated RFA Guidance for angioplasty of Tibial artery-- W contrast IA
25013-4 | XR Toes Views
25014-2 | RFA Two vessels Views W contrast
25015-9 | Replacement of percutaneous gastrojejunostomy Upper GI tract Document
25016-7 | RF Urethra Views W contrast intra urethra
25017-5 | RF Urinary bladder and Urethra Views W contrast intra bladder
25018-3 | NM Urinary bladder Views
25019-1 | US Urinary bladder
25020-9 | RF Urinary bladder and Urethra Views W contrast retrograde via urethra
25025-8 | RFA Vena cava Views W contrast IV
25026-6 | RFA Guidance for placement of venous filter in Inferior vena cava-- W contrast IV
25027-4 | Guidance for placement of large bore CV catheter in Vein
25028-2 | RFA Guidance for placement of catheter for infusion of thrombolytic in Vessel-- W contrast intravascular
25029-0 | RFA Guidance for placement of catheter for vasoconstrictor infusion in Vessels
25030-8 | RFA Abdominal arteries Views W contrast IA
25031-6 | NM Bone Views
25032-4 | NM Bone Views W In-111 tagged WBC IV
25033-2 | MR Wrist
25034-0 | RF Wrist Arthrogram
25035-7 | MR Wrist WO and W contrast IV
25036-5 | US Wrist
25038-1 | Courtesy consultation Unspecified body region Document
25039-9 | CT Unspecified body region limited
25040-7 | Deprecated Unspecified body region CT 3D
25041-5 | CT Guidance for aspiration or biopsy of Unspecified body region-- W contrast IV
25042-3 | CT Guidance for aspiration or biopsy of Unspecified body region
25043-1 | CT Guidance for fluid aspiration of Unspecified body region
25044-9 | CT Guidance for biopsy of Unspecified body region
25045-6 | CT Unspecified body region
25046-4 | CT Unspecified body region W anesthesia
25047-2 | CT Unspecified body region W conscious sedation
25050-6 | Deprecated Unspecified body region CT 3D sagittal and coronal disarticulation
25051-4 | Deprecated Unspecified body region CT Multisectional sagittal
25052-2 | Deprecated Unspecified body region CT sagittal and coronal
25053-0 | CT Guidance for radiosurgery of Unspecified body region
25058-9 | MRA Unspecified body region W contrast IV
25059-7 | US Guidance for biopsy of Unspecified body region
25060-5 | US Unspecified body region No charge
25070-4 | RF Unspecified body region Views during surgery
25071-2 | XR tomography Unspecified body region
25072-0 | Guidance for placement of infusion port
25073-8 | RFA Guidance for removal of foreign body from Vessel
25074-6 | XR Zygomatic arch Views
25076-1 | RFA Hepatic artery Views W contrast IA
25077-9 | RFA Guidance for placement of catheter in Hepatic artery-- W contrast IA
25078-7 | RF Guidance for placement of stent in Intrahepatic portal system
25079-5 | RFA Renal arteries Views W contrast IA
25080-3 | RFA Renal vein - bilateral Views for renin sampling W contrast IV
25081-1 | RFA Guidance for angioplasty of Renal vessel-- W contrast IA
26132-1 | XR Hip - right Views
26133-9 | XR Acetabulum - bilateral Views
26134-7 | XR Acetabulum - left Views
26135-4 | XR Acetabulum - right Views
26136-2 | XR Acromioclavicular joint - bilateral Views
26137-0 | XR Acromioclavicular joint - left Views
26238-6 | MR Finger - bilateral
26239-4 | MR Finger - left
26240-2 | MR Finger - right
26241-0 | MR Foot - bilateral
26242-8 | MR Foot - left
26243-6 | MR Foot - right
26296-4 | RFA Guidance for reposition of CV catheter in Vein - left-- W contrast IV
26297-2 | RFA Guidance for reposition of CV catheter in Vein - right-- W contrast IV
26298-0 | RFA Guidance for atherectomy of Vein - bilateral-- W contrast IV
26299-8 | RFA Guidance for atherectomy of Vein - left-- W contrast IV
26300-4 | RFA Guidance for atherectomy of Vein - right-- W contrast IV
26342-6 | US Guidance for biopsy of Kidney - right
26343-4 | MG Guidance for aspiration of cyst of Breast - bilateral
26344-2 | MG Guidance for aspiration of cyst of Breast - left
26345-9 | MG Guidance for aspiration of cyst of Breast - right
26346-7 | MG Breast - bilateral Diagnostic

26400-2 | XR Hip - bilateral Single view
26401-0 | XR Hip - left Single view
26402-8 | XR Hip - right Single view
26129-7 | XR Foot - right Views
26130-5 | XR Hip - bilateral Views
26131-3 | XR Hip - left Views
26138-8 | XR Acromioclavicular joint - right Views
26139-6 | XR Mastoid - bilateral Views
26140-4 | XR Mastoid - left Views
26141-2 | XR Mastoid - right Views
26142-0 | XR Optic foramen - bilateral Views
26064-6 | RFA Vein - bilateral Views W contrast IV
26065-3 | RFA Vein - left Views W contrast IV
26066-1 | RFA Vein - right Views W contrast IV
26067-9 | RF Salivary gland - bilateral Views W contrast intra salivary duct
26068-7 | RF Salivary gland - left Views W contrast intra salivary duct
26069-5 | RF Salivary gland - right Views W contrast intra salivary duct
26070-3 | RF Hip - bilateral Arthrogram
26071-1 | RF Hip - left Arthrogram
26072-9 | RF Hip - right Arthrogram
26073-7 | RF Knee - bilateral Arthrogram
26074-5 | RF Knee - left Arthrogram
26075-2 | RF Knee - right Arthrogram
26076-0 | RF Shoulder - bilateral Arthrogram
26077-8 | RF Shoulder - left Arthrogram
26078-6 | RF Shoulder - right Arthrogram
26079-4 | RFA Carotid arteries - bilateral Views W contrast IA
26080-2 | RFA Carotid arteries - left Views W contrast IA
26081-0 | RFA Carotid arteries -right Views W contrast IA
26082-8 | RFA Spinal artery - bilateral Views W contrast IA
26083-6 | RFA Spinal artery - left Views W contrast IA
26084-4 | RFA Spinal artery - right Views W contrast IA
26085-1 | XR Knee - bilateral Views W standing
26086-9 | XR Knee - left Views W standing
26087-7 | XR Knee - right Views W standing
26088-5 | NM Knee - bilateral Views
26089-3 | NM Knee - left Views
26090-1 | NM Knee - right Views
26091-9 | NM Scrotum and Testicle - bilateral Views W Tc-99m pertechnetate IV
26092-7 | NM Scrotum and Testicle - left Views W Tc-99m pertechnetate IV
26093-5 | NM Scrotum and Testicle - right Views W Tc-99m pertechnetate IV
26094-3 | XR Foot - bilateral Views W standing
26095-0 | XR Foot - left Views W standing
26096-8 | XR Foot - right Views W standing
26097-6 | XR Ankle - bilateral Views
26098-4 | XR Ankle - left Views
26099-2 | XR Ankle - right Views
26100-8 | XR Calcaneus - bilateral Views
26101-6 | XR Calcaneus - left Views
26102-4 | XR Calcaneus - right Views
26106-5 | XR Clavicle - bilateral Views
26107-3 | XR Clavicle - left Views
26108-1 | XR Clavicle - right Views
26109-9 | XR Elbow - bilateral Views
26110-7 | XR Elbow - left Views
26111-5 | XR Elbow - right Views
26112-3 | XR Lower extremity - bilateral Views
26113-1 | XR Lower extremity - left Views
26114-9 | XR Lower extremity - right Views
26115-6 | XR Upper extremity - bilateral Views
26116-4 | XR Upper extremity - left Views
26117-2 | XR Upper extremity - right Views
26118-0 | XR Femur - bilateral Views
26120-6 | XR Femur - left Views
26122-2 | XR Femur - right Views
26124-8 | XR Finger - bilateral Views
26125-5 | XR Finger - left Views
26126-3 | XR Finger - right Views
26127-1 | XR Foot - bilateral Views
26128-9 | XR Foot - left Views
26143-8 | XR Optic foramen - left Views
26144-6 | XR Optic foramen - right Views

26146-1 | XR Radius and Ulna - bilateral Views
26148-7 | XR Radius and Ulna - left Views
26150-3 | XR Radius and Ulna - right Views
26151-1 | XR Ribs - bilateral Views
26152-9 | XR Ribs - left Views
26153-7 | XR Ribs - right Views
26154-5 | XR Scapula - bilateral Views
26155-2 | XR Scapula - left Views
26156-0 | XR Scapula - right Views
26157-8 | XR Shoulder - bilateral Views
26158-6 | XR Shoulder - left Views
26159-4 | XR Shoulder - right Views
26160-2 | XR Thumb - bilateral Views
26161-0 | XR Thumb - left Views
26162-8 | XR Thumb - right Views
26163-6 | XR Tibia and Fibula - bilateral Views
26164-4 | XR Tibia and Fibula - left Views
26165-1 | XR Tibia and Fibula - right Views
26166-9 | XR Toes - bilateral Views
26167-7 | XR Toes - left Views
26168-5 | XR Toes - right Views
26169-3 | XR Wrist - bilateral Views
26170-1 | XR Wrist - left Views
26171-9 | XR Wrist - right Views
26172-7 | XR Zygomatic arch - bilateral Views
26173-5 | XR Zygomatic arch - left Views
26174-3 | XR Zygomatic arch - right Views
26175-0 | MG Breast - bilateral Screening
26176-8 | MG Breast - left Screening
26177-6 | MG Breast - right Screening
26178-4 | RFA Femoral artery - bilateral Runoff W contrast IA
26179-2 | RFA Femoral artery - left Runoff W contrast IA
26180-0 | RFA Femoral artery - right Runoff W contrast IA
26181-8 | MRA Thoracic inlet vessels - bilateral W contrast IV
26182-6 | MRA Thoracic inlet vessels - left W contrast IV
26183-4 | MRA Thoracic inlet vessels - right W contrast IV
26184-2 | CT Extremity - bilateral W contrast IV
26185-9 | CT Extremity - left W contrast IV
26186-7 | CT Extremity - right W contrast IV
26187-5 | MR Ankle - bilateral WO and W contrast IV
26188-3 | MR Ankle - left WO and W contrast IV
26189-1 | MR Ankle - right WO and W contrast IV
26190-9 | MR Brachial plexus - bilateral WO and W contrast IV
26191-7 | MR Brachial plexus - left WO and W contrast IV
26192-5 | MR Brachial plexus - right WO and W contrast IV
26193-3 | MR Elbow - bilateral WO and W contrast IV
26194-1 | MR Elbow - left WO and W contrast IV
26195-8 | MR Elbow - right WO and W contrast IV
26196-6 | MR Thigh - bilateral WO and W contrast IV
26197-4 | MR Thigh - left WO and W contrast IV
26198-2 | MR Thigh - right WO and W contrast IV
26199-0 | MR Knee - bilateral WO and W contrast IV
26200-6 | MR Knee - left WO and W contrast IV
26201-4 | MR Knee - right WO and W contrast IV
26202-2 | MR Shoulder - bilateral WO and W contrast IV
26203-0 | MR Shoulder - left WO and W contrast IV
26204-8 | MR Shoulder - right WO and W contrast IV
26205-5 | MR Wrist - bilateral WO and W contrast IV
26206-3 | MR Wrist - left WO and W contrast IV
26207-1 | MR Wrist - right WO and W contrast IV
26208-9 | MR Ankle - bilateral
26209-7 | MR Ankle - left
26210-5 | MR Ankle - right
26211-3 | MR Brachial plexus - bilateral
26212-1 | MR Brachial plexus - left
26213-9 | MR Brachial plexus - right
26214-7 | US Breast - bilateral
26215-4 | US Breast - left
26216-2 | US Breast - right
26217-0 | US Carotid arteries - bilateral
26218-8 | US Carotid arteries - left
26219-6 | US Carotid arteries -right

26220-4 | MR Elbow - bilateral
26221-2 | MR Elbow - left
26222-0 | MR Elbow - right
26223-8 | US Extremity - bilateral
26224-6 | CT Extremity - bilateral
26225-3 | US Extremity - left
26226-1 | CT Extremity - left
26227-9 | MR Lower extremity joint - bilateral
26228-7 | MR Lower extremity joint - left
26229-5 | MR Lower extremity joint - right
26230-3 | US Extremity - right
26231-1 | CT Extremity - right
26232-9 | MR Upper extremity - bilateral
26233-7 | MR Upper extremity - left
26234-5 | MR Upper extremity - right
26235-2 | MR Thigh - bilateral
26236-0 | MR Thigh - left
26237-8 | MR Thigh - right
26244-4 | MR Forearm - bilateral
26245-1 | MR Forearm - left
26246-9 | MR Forearm - right
26247-7 | MR Hand - bilateral
26248-5 | MR Hand - left
26249-3 | MR Hand - right
26250-1 | US Hip - bilateral
26251-9 | US Hip - left
26252-7 | US Hip - right
26253-5 | XR tomography Internal auditory canal - bilateral
26254-3 | XR tomography Internal auditory canal - left
26255-0 | XR tomography Internal auditory canal - right
26256-8 | MR Knee - bilateral
26257-6 | MR Knee - left
26258-4 | MR Knee - right
26259-2 | MR Pelvis and Hip - bilateral
26260-0 | MR Pelvis and Hip - left
26261-8 | MR Pelvis and Hip - right
26262-6 | US Popliteal space - bilateral
26263-4 | US Popliteal space - left
26264-2 | US Popliteal space - right
26265-9 | US Shoulder - bilateral
26266-7 | MR Shoulder - bilateral
26267-5 | US Shoulder - left
26268-3 | MR Shoulder - left
26269-1 | US Shoulder - right
26270-9 | MR Shoulder - right
26271-7 | US Scrotum and Testicle - bilateral
26272-5 | US Scrotum and Testicle - left
26273-3 | US Scrotum and Testicle - right
26277-4 | MR Wrist - bilateral
26278-2 | US Wrist - bilateral
26279-0 | MR Wrist - left
26280-8 | US Wrist - left
26281-6 | MR Wrist - right
26282-4 | US Wrist - right
26283-2 | XR Knee - bilateral Merchants
26284-0 | XR Knee - left Merchants
26285-7 | XR Knee - right Merchants
26286-5 | US Breast - bilateral limited
26287-3 | MG Breast - bilateral Limited Views
26288-1 | US Breast - left limited
26289-9 | MG Breast - left Limited Views
26290-7 | US Breast - right limited
26291-5 | MG Breast - right Limited Views
26292-3 | MG stereo Guidance for biopsy of Breast - bilateral
26293-1 | MG stereo Guidance for biopsy of Breast - left
26294-9 | MG stereo Guidance for biopsy of Breast - right
26295-6 | RFA Guidance for reposition of CV catheter in Vein - bilateral-- W contrast IV
26301-2 | RFA Guidance for placement of stent in Vein - bilateral
26302-0 | RFA Guidance for placement of stent in Vein - left
26303-8 | RFA Guidance for placement of stent in Vein - right
26304-6 | RFA Guidance for placement of peripherally-inserted central venous catheter in Vein - bilateral
26305-3 | RFA Guidance for placement of peripherally-inserted central venous catheter in Vein - left

26306-1 | RFA Guidance for placement of peripherally-inserted central venous catheter in Vein - right
26307-9 | Guidance for placement of large bore CV catheter in Vein - bilateral
26308-7 | Guidance for placement of large bore CV catheter in Vein - left
26309-5 | Guidance for placement of large bore CV catheter in Vein - right
26310-3 | RFA Guidance for placement of CV catheter in Vein - bilateral-- W contrast IV
26311-1 | RFA Guidance for placement of CV catheter in Vein - left-- W contrast IV
26312-9 | RFA Guidance for placement of CV catheter in Vein - right-- W contrast IV
26313-7 | US Guidance for needle localization of Breast - bilateral
26314-5 | US Guidance for needle localization of Breast - left
26315-2 | MG Guidance for needle localization of mass of Breast - bilateral
26316-0 | MG Guidance for needle localization of mass of Breast - left
26317-8 | MG Guidance for needle localization of mass of Breast - right
26318-6 | US Guidance for needle localization of Breast - right
26319-4 | CT Guidance for injection of Sacroiliac joint - bilateral
26320-2 | CT Guidance for injection of Sacroiliac joint - left
26321-0 | CT Guidance for injection of Sacroiliac joint - right
26322-8 | RF Guidance for injection of Spine facet joint - bilateral
26323-6 | RF Guidance for injection of Spine facet joint - left
26324-4 | RF Guidance for injection of Spine facet joint - right
26325-1 | US Guidance for drainage of Extremity - bilateral
26326-9 | US Guidance for drainage of Extremity - left
26327-7 | US Guidance for drainage of Extremity - right
26328-5 | US Guidance for drainage of Kidney - bilateral
26329-3 | US Guidance for drainage of Kidney - left
26330-1 | US Guidance for drainage of Kidney - right
26331-9 | RFA Guidance for change of CV catheter in Vein - bilateral-- W contrast IV
26332-7 | RFA Guidance for change of CV catheter in Vein - left-- W contrast IV
26333-5 | RFA Guidance for change of CV catheter in Vein - right-- W contrast IV
26334-3 | MG Guidance for percutaneous biopsy.core needle of Breast - bilateral
26335-0 | MG Guidance for percutaneous biopsy.core needle of Breast - left
26336-8 | MG Guidance for percutaneous biopsy.core needle of Breast - right
26337-6 | MG Guidance for biopsy of Breast - bilateral
26338-4 | MG Guidance for biopsy of Breast - left
26339-2 | MG Guidance for biopsy of Breast - right
26340-0 | US Guidance for biopsy of Kidney - bilateral
26341-8 | US Guidance for biopsy of Kidney - left
26347-5 | MG Breast - left Diagnostic
26348-3 | MG Breast - right Diagnostic
26349-1 | MG Breast - bilateral Diagnostic Limited Views
26350-9 | MG Breast - left Diagnostic Limited Views
26351-7 | MG Breast - right Diagnostic Limited Views
26352-5 | XR Wrist - bilateral and Hand - bilateral Bone age Views
26353-3 | XR Wrist - left and Hand - left Bone age Views
26354-1 | XR Wrist - right and Hand - right Bone age Views
26355-8 | XR Hand - bilateral Arthritis
26356-6 | XR Hand - left Arthritis
26357-4 | XR Hand - right Arthritis
26358-2 | XR Knee - bilateral AP W standing
26359-0 | XR Knee - left AP W standing
26360-8 | XR Knee - right AP W standing
26361-6 | XR Knee - bilateral AP and PA W standing
26362-4 | XR Knee - left AP and PA W standing
26363-2 | XR Knee - right AP and PA W standing
26364-0 | XR Knee - bilateral AP and Lateral W standing
26365-7 | XR Knee - left AP and Lateral W standing
26366-5 | XR Knee - right AP and Lateral W standing
26368-1 | Deprecated Brachiocephalic artery - left Fluoroscopic angiogram Angioplasty W contrast IA
26369-9 | Deprecated Brachiocephalic artery - right Fluoroscopic angiogram Angioplasty W contrast IA
26370-7 | RFA Guidance for angioplasty of Iliac artery - bilateral-- W contrast IA
26371-5 | RFA Guidance for angioplasty of Iliac artery - left-- W contrast IA
26372-3 | RFA Guidance for angioplasty of Iliac artery - right-- W contrast IA
26373-1 | Deprecated RFA Guidance for angioplasty of Tibial artery - bilateral-- W contrast IA
26374-9 | Deprecated RFA Guidance for angioplasty of Tibial artery - left-- W contrast IA
26375-6 | Deprecated RFA Guidance for angioplasty of Tibial artery - right-- W contrast IA
26376-4 | Administration of vasodilator into catheter Vein - bilateral
26377-2 | Administration of vasodilator into catheter Vein - left
26378-0 | Administration of vasodilator into catheter Vein - right
26379-8 | XR Hand - bilateral 3 Views
26380-6 | XR Hand - left 3 Views
26381-4 | XR Hand - right 3 Views
26382-2 | XR Shoulder - bilateral 3 Views
26383-0 | XR Shoulder - left 3 Views

26384-8 | XR Shoulder - right 3 Views
26385-5 | XR Ankle - bilateral 2 Views
26386-3 | XR Ankle - left 2 Views
26387-1 | XR Ankle - right 2 Views
26388-9 | XR Hand - bilateral 2 Views
26389-7 | XR Hand - left 2 Views
26390-5 | XR Hand - right 2 Views
26391-3 | XR Humerus - bilateral 2 Views
26392-1 | XR Humerus - left 2 Views
26393-9 | XR Humerus - right 2 Views
26394-7 | XR Knee - bilateral 2 Views
26395-4 | XR Knee - left 2 Views
26396-2 | XR Knee - right 2 Views
26397-0 | XR Patella - bilateral 2 Views
26398-8 | XR Patella - left 2 Views
26399-6 | XR Patella - right 2 Views
28032-1 | Deprecated Cardiac echo, study
28561-9 | XR Pelvis Views
28564-3 | XR Skull Views
28565-0 | XR Knee Views
28566-8 | CT Spine
28567-6 | XR Humerus Views
28568-4 | Physician Emergency department Note
28569-2 | Physician consulting Progress note
28570-0 | Procedure note
28571-8 | Speech-language pathology Note
28572-6 | Dentistry Initial evaluation note
28573-4 | Physician, Operation note
28574-2 | Deprecated Discharge note
28575-9 | Nurse practitioner Progress note
28576-7 | MR Joint
28577-5 | Dentistry Procedure note
28578-3 | Occupational therapy Note
28579-1 | Physical therapy Note
28580-9 | Chiropractic medicine Progress note
28581-7 | Chiropractic medicine Initial evaluation note
28582-5 | XR Hand Views
28583-3 | Dentist Operation note
28613-8 | XR Spine Views
28614-6 | US Liver
28615-3 | Audiology study
28616-1 | Physician Transfer note
28617-9 | Dentistry Progress note
28618-7 | Dentistry Note
28621-1 | Nurse practitioner Initial evaluation note
28622-9 | Nurse Discharge assessment
28623-7 | Nurse Progress note
28624-5 | Podiatry Operation note
28625-2 | Podiatry Procedure note
28626-0 | Physician History and physical note
28627-8 | Psychiatry Progress note
28628-6 | Psychiatry Note
28629-4 | Perimetry study
28630-2 | Tonometry study
28631-0 | Visual acuity study
28632-8 | Heterophoria study
28633-6 | Polysomnography (sleep) study
28635-1 | Psychiatry Initial evaluation note
28636-9 | Initial evaluation note
28651-8 | Nurse Transfer note
28653-4 | Social work Note
28654-2 | Physician attending Initial evaluation note
28655-9 | Physician attending Discharge summary
28656-7 | Social work Progress note
29272-2 | Eye ultrasound study
29206-0 | Speech therapy service attachment
29252-4 | CT Chest WO contrast
29749-9 | Dialysis records
29750-7 | Neonatal intensive care records
29751-5 | Critical care records
29752-3 | Perioperative records
29753-1 | Nurse Initial evaluation note

29754-9 | Nystagmogram study
29755-6 | Nerve conduction study
29756-4 | Peritoneoscopy study
29757-2 | Colposcopy study
29761-4 | Dentistry Discharge summary
30816-3 | RFA Peritoneum Views W contrast percutaneous intraperitoneal
30817-1 | RF Guidance for injection of Spine Lumbar Facet Joint
30818-9 | RF Guidance for placement of catheter in Fallopian tubes-- transcervical
30819-7 | RFA Epidural veins Views W contrast IV
30604-3 | CT Guidance for biopsy of Pancreas
30605-0 | CT Guidance for fine needle aspiration of Pancreas
30606-8 | CT Guidance for fine needle aspiration of Pelvis
30607-6 | CT Guidance for biopsy of Kidney - bilateral
30608-4 | CT Guidance for fine needle aspiration of Kidney - bilateral
30578-9 | CT Guidance for drainage of abscess of Unspecified body region
30579-7 | CT Guidance for injection of Spine facet joint
30580-5 | CT Guidance for fine needle aspiration of Unspecified body region
30581-3 | CT Guidance for radiation treatment of Unspecified body region-- W contrast IV
30582-1 | CT Guidance for radiation treatment of Unspecified body region-- WO contrast
30583-9 | CT Internal auditory canal W contrast IV
30584-7 | CT Internal auditory canal WO contrast
30585-4 | CT Nasopharynx and Neck WO contrast
30586-2 | CT Neck WO and W contrast IV
30587-0 | CT Orbit - bilateral WO contrast
30588-8 | CT Sinuses
30589-6 | CT Petrous part of temporal bone WO contrast
30590-4 | CT Pituitary and Sella turcica W contrast IV

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.551 Value Set http://hl7.org/fhir/ValueSet/metric-calibration-state

URL: https://hl7.org/fhir/R4/valueset-metric-calibration-state.html
Health Care Devices Work Group | Maturity Level: 1 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/metric-calibration-state
Version: | 4.0.1
Name: | DeviceMetricCalibrationState
Title: | DeviceMetricCalibrationState
Definition: | Describes the state of a metric calibration.
Committee: | Health Care Devices Work Group
OID: | 2.16.840.1.113883.4.642.3.652 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for DeviceMetricCalibrationState

• Resource: DeviceMetric.calibration.state (code / Required)

4.4.1.551.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/metric-calibration-state

4.4.1.551.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 4 concepts

Expansion based on http://hl7.org/fhir/metric-calibration-state version 4.0.1

All codes from system http://hl7.org/fhir/metric-calibration-state

Code | Display | Definition
not-calibrated | Not Calibrated | The metric has not been calibrated.
calibration-required | Calibration Required | The metric needs to be calibrated.
calibrated | Calibrated | The metric has been calibrated.
unspecified | Unspecified | The state of calibration of this metric is unspecified.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.552 Value Set http://hl7.org/fhir/ValueSet/data-absent-reason

URL: https://hl7.org/fhir/R4/valueset-data-absent-reason.html
FHIR Infrastructure Work Group | Maturity Level: N | Normative (from v4.0.0) | Use Context: Any

| This page has been approved as part of an ANSI standard. See the Observation Package for further details.

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/data-absent-reason
Version: | 4.0.1
Name: | DataAbsentReason
Title: | DataAbsentReason
Definition: | Used to specify why the normally expected content of the data element is missing.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.5 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for DataAbsentReason
• ConceptMap: Translation source in v3 map for DataAbsentReason
• Resource: Observation.dataAbsentReason (CodeableConcept / Extensible)
• Resource: Observation.component.dataAbsentReason (CodeableConcept / Extensible)
• Profile: DeviceMetricObservation: Observation.dataAbsentReason (CodeableConcept / Extensible)
• Profile: DeviceMetricObservation: Observation.component.dataAbsentReason (CodeableConcept / Extensible)
• Profile: Observation-Genetics: Observation.dataAbsentReason (CodeableConcept / Extensible)
• Profile: Observation-Genetics: Observation.component.dataAbsentReason (CodeableConcept / Extensible)
• Profile: VitalSigns: Observation.dataAbsentReason (CodeableConcept / Extensible)
• Profile: VitalSigns: Observation.component.dataAbsentReason (CodeableConcept / Extensible)
• Profile: BodyWeight: Observation.dataAbsentReason (CodeableConcept / Extensible)
• Profile: BodyWeight: Observation.component.dataAbsentReason (CodeableConcept / Extensible)
• Profile: VitalsPanel: Observation.dataAbsentReason (CodeableConcept / Extensible)
• Profile: VitalsPanel: Observation.component.dataAbsentReason (CodeableConcept / Extensible)
• Profile: BodyHeight: Observation.dataAbsentReason (CodeableConcept / Extensible)
• Profile: BodyHeight: Observation.component.dataAbsentReason (CodeableConcept / Extensible)
• Profile: RespRate: Observation.dataAbsentReason (CodeableConcept / Extensible)
• Profile: RespRate: Observation.component.dataAbsentReason (CodeableConcept / Extensible)
• Profile: HeartRate: Observation.dataAbsentReason (CodeableConcept / Extensible)
• Profile: HeartRate: Observation.component.dataAbsentReason (CodeableConcept / Extensible)
• Profile: BodyTemp: Observation.dataAbsentReason (CodeableConcept / Extensible)
• Profile: BodyTemp: Observation.component.dataAbsentReason (CodeableConcept / Extensible)

- Profile: HeadCircum: Observation.dataAbsentReason (CodeableConcept / Extensible)
- Profile: HeadCircum: Observation.component.dataAbsentReason (CodeableConcept / Extensible)
- Profile: OxygenSat: Observation.dataAbsentReason (CodeableConcept / Extensible)
- Profile: OxygenSat: Observation.component.dataAbsentReason (CodeableConcept / Extensible)
- Profile: BMI: Observation.dataAbsentReason (CodeableConcept / Extensible)
- Profile: BMI: Observation.component.dataAbsentReason (CodeableConcept / Extensible)
- Profile: BP: Observation.dataAbsentReason (CodeableConcept / Extensible)
- Profile: BP: Observation.component.dataAbsentReason (CodeableConcept / Extensible)
- Profile: Cholesterol: Observation.dataAbsentReason (CodeableConcept / Extensible)
- Profile: Cholesterol: Observation.component.dataAbsentReason (CodeableConcept / Extensible)
- Profile: Triglyceride: Observation.dataAbsentReason (CodeableConcept / Extensible)
- Profile: Triglyceride: Observation.component.dataAbsentReason (CodeableConcept / Extensible)
- Profile: HDLCholesterol: Observation.dataAbsentReason (CodeableConcept / Extensible)
- Profile: HDLCholesterol: Observation.component.dataAbsentReason (CodeableConcept / Extensible)
- Profile: LDLCholesterol: Observation.dataAbsentReason (CodeableConcept / Extensible)
- Profile: LDLCholesterol: Observation.component.dataAbsentReason (CodeableConcept / Extensible)

4.4.1.552.1 Content Logical Definition

This value set includes codes from the following code systems:

- Include all codes defined in http://terminology.hl7.org/CodeSystem/data-absent-reason

4.4.1.552.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 15 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/data-absent-reason version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/data-absent-reason

Lvl | Code | Display | Definition
0 | unknown | Unknown | The value is expected to exist but is not known.
1 | asked-unknown | Asked But Unknown | The source was asked but does not know the value.
1 | temp-unknown | Temporarily Unknown | There is reason to expect (from the workflow) that the value may become known.
1 | not-asked | Not Asked | The workflow didn't lead to this value being known.
1 | asked-declined | Asked But Declined | The source was asked but declined to answer.
0 | masked | Masked | The information is not available due to security, privacy or related reasons.
0 | not-applicable | Not Applicable | There is no proper value for this element (e.g. last menstrual period for a male).
0 | unsupported | Unsupported | The source system wasn't capable of supporting this element.
0 | as-text | As Text | The content of the data is represented in the resource narrative.
0 | error | Error | Some system or workflow process error means that the information is not available.
1 | not-a-number | Not a Number (NaN) | The numeric value is undefined or unrepresentable due to a floating point processing error.
1 | negative-infinity | Negative Infinity (NINF) | The numeric value is excessively low and unrepresentable due to a floating point processing error.
1 | positive-infinity | Positive Infinity (PINF) | The numeric value is excessively high and unrepresentable due to a floating point processing error.
0 | not-performed | Not Performed | The value is not available because the observation procedure (test, etc.) was not performed.
0 | not-permitted | Not Permitted | The value is not permitted in this context (e.g. due to profiles, or the base data types).

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.553 Value Set http://hl7.org/fhir/ValueSet/product-storage-scale

URL: https://hl7.org/fhir/R4/valueset-product-storage-scale.html
Orders and Observations Work Group | Maturity Level: 0 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/product-storage-scale
Version: | 4.0.1
Name: | BiologicallyDerivedProductStorageScale
Title: | BiologicallyDerivedProductStorageScale
Definition: | BiologicallyDerived Product Storage Scale.
Committee: | Orders and Observations Work Group
OID: | 2.16.840.1.113883.4.642.3.904 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for BiologicallyDerivedProductStorageScale

• Resource: BiologicallyDerivedProduct.storage.scale (code / Required)

4.4.1.553.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/product-storage-scale

4.4.1.553.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 3 concepts

Expansion based on http://hl7.org/fhir/product-storage-scale version 4.0.1

All codes from system http://hl7.org/fhir/product-storage-scale

Code | Display | Definition
farenheit | Fahrenheit | Fahrenheit temperature scale.
celsius | Celsius | Celsius or centigrade temperature scale.
kelvin | Kelvin | Kelvin absolute thermodynamic temperature scale.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.554 Value Set http://hl7.org/fhir/ValueSet/related-artifact-type

URL: https://hl7.org/fhir/R4/valueset-related-artifact-type.html
FHIR Infrastructure Work Group | Maturity Level: 5 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/related-artifact-type
Version: | 4.0.1
Name: | RelatedArtifactType
Title: | RelatedArtifactType
Definition: | The type of relationship to the related artifact.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.99 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for RelatedArtifactType

• Resource: RelatedArtifact.type (code / Required)

4.4.1.554.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/related-artifact-type

4.4.1.554.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 8 concepts

Expansion based on http://hl7.org/fhir/related-artifact-type version 4.0.1

All codes from system http://hl7.org/fhir/related-artifact-type

Code | Display | Definition
documentation | Documentation | Additional documentation for the knowledge resource. This would include additional instructions on usage as well as additional information on clinical context or appropriateness.
justification | Justification | A summary of the justification for the knowledge resource including supporting evidence, relevant guidelines, or other clinically important information. This information is intended to provide a way to make the justification for the knowledge resource available to the consumer of interventions or results produced by the knowledge resource.
citation | Citation | Bibliographic citation for papers, references, or other relevant material for the knowledge resource. This is intended to allow for citation of related material, but that was not necessarily specifically prepared in connection with this knowledge resource.
predecessor | Predecessor | The previous version of the knowledge resource.
successor | Successor | The next version of the knowledge resource.
derived-from | Derived From | The knowledge resource is derived from the related artifact. This is intended to capture the relationship in which a particular knowledge resource is based on the content of another artifact, but is modified to capture either a different set of overall requirements, or a more specific set of requirements such as those involved in a particular institution or clinical setting.
depends-on | Depends On | The knowledge resource depends on the given related artifact.
composed-of | Composed Of | The knowledge resource is composed of the given related artifact.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.555 Value Set http://hl7.org/fhir/ValueSet/research-study-prim-purp-type

URL: https://hl7.org/fhir/R4/valueset-research-study-prim-purp-type.html
Biomedical Research and Regulation Work Group | Maturity Level: 1 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/research-study-prim-purp-type
Version: | 4.0.1
Name: | ResearchStudyPrimaryPurposeType
Title: | ResearchStudyPrimaryPurposeType
Definition: | Codes for the main intent of the study.
Committee: | Biomedical Research and Regulation Work Group
OID: | 2.16.840.1.113883.4.642.3.827 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ResearchStudyPrimaryPurposeType

• Resource: ResearchStudy.primaryPurposeType (CodeableConcept / Extensible)

4.4.1.555.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/research-study-prim-purp-type

4.4.1.555.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 8 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/research-study-prim-purp-type version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/research-study-prim-purp-type

Code | Display | Definition

treatment | Treatment | One or more interventions are being evaluated for treating a disease, syndrome, or condition.
prevention | Prevention | One or more interventions are being assessed for preventing the development of a specific disease or health condition.
diagnostic | Diagnostic | One or more interventions are being evaluated for identifying a disease or health condition.
supportive-care | Supportive Care | One or more interventions are evaluated for maximizing comfort, minimizing side effects, or mitigating against a decline in the participant's health or function.
screening | Screening | One or more interventions are assessed or examined for identifying a condition, or risk factors for a condition, in people who are not yet known to have the condition or risk factor.
health-services-research | Health Services Research | One or more interventions for evaluating the delivery, processes, management, organization, or financing of healthcare.
basic-science | Basic Science | One or more interventions for examining the basic mechanism of action (for example, physiology or biomechanics of an intervention).
device-feasibility | Device Feasibility | An intervention of a device product is being evaluated to determine the feasibility of the product or to test a prototype device and not health outcomes. Such studies are conducted to confirm the design and operating specifications of a device before beginning a full clinical trial.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.556 Value Set http://hl7.org/fhir/ValueSet/consent-provision-type

URL: https://hl7.org/fhir/R4/valueset-consent-provision-type.html
Community Based Collaborative Care Work Group | Maturity Level: 2 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/consent-provision-type
Version: | 4.0.1
Name: | ConsentProvisionType
Title: | ConsentProvisionType
Definition: | How a rule statement is applied, such as adding additional consent or removing consent.
Committee: | Community Based Collaborative Care Work Group
OID: | 2.16.840.1.113883.4.642.3.757 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ConsentProvisionType
• Resource: Consent.provision.type (code / Required)

4.4.1.556.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/consent-provision-type

4.4.1.556.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 2 concepts

Expansion based on http://hl7.org/fhir/consent-provision-type version 4.0.1

All codes from system http://hl7.org/fhir/consent-provision-type

Code | Display | Definition
deny | Opt Out | Consent is denied for actions meeting these rules.
permit | Opt In | Consent is provided for actions meeting these rules.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')

Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.557 Value Set http://hl7.org/fhir/ValueSet/contract-assetcontext

URL: https://hl7.org/fhir/R4/valueset-contract-assetcontext.html
Financial Management Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/contract-assetcontext
Version: | 4.0.1
Name: | ContractResourceAssetContextCodes
Title: | Contract Resource Asset Context codes
Definition: | This value set contract specific codes for asset context.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.1297 (for OID based terminology systems)
Copyright: | HL7 International.
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ContractResourceAssetContextCodes

• Resource: Contract.term.asset.context.code (CodeableConcept / Example)

4.4.1.557.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/contract-asset-context

4.4.1.557.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 1 concepts

Expansion based on http://hl7.org/fhir/contract-asset-context version 4.0.1

All codes from system http://hl7.org/fhir/contract-asset-context

Code | Display | Definition
custodian | Custodian | To be completed

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.558 Value Set http://hl7.org/fhir/ValueSet/message-significance-category

URL: https://hl7.org/fhir/R4/valueset-message-significance-category.html
Infrastructure And Messaging Work Group | Maturity Level: 1 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/message-significance-category
Version: | 4.0.1
Name: | MessageSignificanceCategory
Title: | MessageSignificanceCategory
Definition: | The impact of the content of a message.
Committee: | Infrastructure And Messaging Work Group
OID: | 2.16.840.1.113883.4.642.3.188 (for OID based terminology systems)

Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for MessageSignificanceCategory

• Resource: MessageDefinition.category (code / Required)

4.4.1.558.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/message-significance-category

4.4.1.558.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 3 concepts

Expansion based on http://hl7.org/fhir/message-significance-category version 4.0.1

All codes from system http://hl7.org/fhir/message-significance-category

Code | Display | Definition
consequence | Consequence | The message represents/requests a change that should not be processed more than once; e.g., making a booking for an appointment.
currency | Currency | The message represents a response to query for current information. Retrospective processing is wrong and/or wasteful.
notification | Notification | The content is not necessarily intended to be current, and it can be reprocessed, though there may be version issues created by processing old notifications.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.559 Value Set http://hl7.org/fhir/ValueSet/sort-direction

URL: https://hl7.org/fhir/R4/valueset-sort-direction.html
FHIR Infrastructure Work Group | Maturity Level: 5 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/sort-direction
Version: | 4.0.1
Name: | SortDirection
Title: | SortDirection
Definition: | The possible sort directions, ascending or descending.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.979 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for SortDirection

• Resource: DataRequirement.sort.direction (code / Required)

4.4.1.559.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/sort-direction

4.4.1.559.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 2 concepts

Expansion based on http://hl7.org/fhir/sort-direction version 4.0.1

All codes from system http://hl7.org/fhir/sort-direction

Code | Display | Definition

ascending | Ascending | Sort by the value ascending, so that lower values appear first.
descending | Descending | Sort by the value descending, so that lower values appear last.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.560 Value Set http://hl7.org/fhir/ValueSet/list-mode

URL: https://hl7.org/fhir/R4/valueset-list-mode.html
FHIR Infrastructure Work Group | Maturity Level: 2 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/list-mode
Version: | 4.0.1
Name: | ListMode
Title: | ListMode
Definition: | The processing mode that applies to this list.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.318 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ListMode
• Resource: List.mode (code / Required)
• Resource: Composition.section.mode (code / Required)

4.4.1.560.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/list-mode

4.4.1.560.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 3 concepts

Expansion based on http://hl7.org/fhir/list-mode version 4.0.1

All codes from system http://hl7.org/fhir/list-mode

Code | Display | Definition
working | Working List | This list is the master list, maintained in an ongoing fashion with regular updates as the real world list it is tracking changes.
snapshot | Snapshot List | This list was prepared as a snapshot. It should not be assumed to be current.
changes | Change List | A point-in-time list that shows what changes have been made or recommended. E.g. a discharge medication list showing what was added and removed during an encounter.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.561 Value Set http://hl7.org/fhir/ValueSet/exposure-state

URL: https://hl7.org/fhir/R4/valueset-exposure-state.html
Clinical Decision Support Work Group | Maturity Level: 0 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/exposure-state
Version: | 4.0.1
Name: | ExposureState
Title: | ExposureState
Definition: | Whether the results by exposure is describing the results for the primary exposure of interest (exposure) or the alternative state (exposureAlternative).
Committee: | Clinical Decision Support Work Group
OID: | 2.16.840.1.113883.4.642.3.1351 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ExposureState

• Resource: EffectEvidenceSynthesis.resultsByExposure.exposureState (code / Required)

4.4.1.561.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/exposure-state

4.4.1.561.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 2 concepts

Expansion based on http://hl7.org/fhir/exposure-state version 4.0.1

All codes from system http://hl7.org/fhir/exposure-state

Code | Display | Definition
exposure | Exposure | used when the results by exposure is describing the results for the primary exposure of interest.
exposure-alternative | Exposure Alternative | used when the results by exposure is describing the results for the alternative exposure state, control state or comparator state.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.562 Value Set http://hl7.org/fhir/ValueSet/medication-form-codes

URL: https://hl7.org/fhir/R4/valueset-medication-form-codes.html
Pharmacy Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/medication-form-codes
Version: | 4.0.1
Name: | SNOMEDCTFormCodes
Title: | SNOMED CT Form Codes
Definition: | This value set includes all dose form codes from SNOMED CT - provided as an exemplar.
Committee: | Pharmacy Work Group
OID: | 2.16.840.1.113883.4.642.3.374 (for OID based terminology systems)
Copyright: | This value set includes content from SNOMED CT, which is copyright © 2002 International Health Terminology Standards Development Organisation (IHTSDO), and distributed by agreement between IHTSDO and HL7. Implementer use of SNOMED CT is not covered by this agreement.
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: MedicationKnowledge.doseForm (CodeableConcept / Example)

4.4.1.562.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include codes from http://snomed.info/sct where concept is-a 421967003 (Drug dose form)

4.4.1.562.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 352 concepts

All codes from system http://snomed.info/sct

Code	Display	Definition
7946007	Drug suspension	
11190007	Drug stick	
17519006	Lotion	
30843009	Drug aerosol foam	
36875001	Drug patch	
37937005	Drug paste	
46992007	Pill	
48582000	Caplet	
52262001	Drug aerosol	
63316001	Liniment	
64241004	Drug pledget	
66076007	Chewable tablet	
77899000	Drug solution	
85581007	Drug powder	
385018001	Oral drops	
385019009	Oral drops solution	
385020003	Oral drops suspension	
385021004	Oral drops emulsion	
385022006	Oral liquid	
385023001	Oral solution	
385024007	Oral suspension	
385025008	Oral emulsion	
385026009	Powder for oral solution	
385027000	Powder for oral suspension	
385028005	Granules for oral solution	
385029002	Granules for oral suspension	
385032004	Syrup	
385033009	Powder for syrup	
385034003	Granules for syrup	
385035002	Soluble tablet	
385036001	Dispersible tablet	
385038000	Oral gel	
385039008	Oral paste	
385041009	Oral powder	
385042002	Effervescent powder	
385043007	Granules	
385044001	Effervescent granules	
385045000	Gastro-resistant granules	
385046004	Prolonged-release granules	
385047008	Modified-release granules	
385048003	Cachet	
385049006	Capsule	
385050006	Hard capsule	
385051005	Soft capsule	
385052003	Gastro-resistant capsule	
385053008	Prolonged-release capsule	
385054002	Modified-release capsule	
385055001	Tablet	
385057009	Film-coated tablet	
385058004	Effervescent tablet	
385059007	Gastro-resistant tablet	
385060002	Prolonged-release tablet	
385061003	Modified-release tablet	
385062005	Oral lyophilisate	
385063000	Oral gum	
385064006	Pillule	
385069001	Oromucosal liquid	
385070000	Oromucosal solution	
385071001	Oromucosal suspension	
385073003	Oromucosal spray	
385074009	Sublingual spray	
385077002	Gingival solution	
385078007	Oromucosal gel	

385079004 | Oromucosal paste
385080001 | Medicated chewing-gum
385081002 | Gingival gel
385082009 | Gingival paste
385083004 | Oromucosal capsule
385084005 | Sublingual tablet
385085006 | Buccal tablet
385086007 | Muco-adhesive buccal tablet
385087003 | Lozenge
385088008 | Dental gel
385089000 | Dental stick
385090009 | Dental insert
385091008 | Dental powder
385092001 | Dental liquid
385094000 | Dental solution
385095004 | Dental suspension
385096003 | Dental emulsion
385098002 | Bath additive
385099005 | Cream
385100002 | Gel
385101003 | Ointment
385102005 | Cutaneous paste
385103000 | Cutaneous foam
385104006 | Shampoo
385105007 | Cutaneous spray
385106008 | Cutaneous solution spray
385107004 | Cutaneous liquid
385108009 | Cutaneous solution
385110006 | Cutaneous suspension
385111005 | Cutaneous emulsion
385112003 | Cutaneous powder
385113008 | Solution for iontophoresis
385114002 | Transdermal patch
385115001 | Collodion
385116000 | Medicated nail laquer
385117009 | Poultice
385118004 | Cutaneous stick
385119007 | Cutaneous sponge
385121002 | Eye cream
385122009 | Eye gel
385123004 | Eye ointment
385124005 | Eye drops
385125006 | Eye drops solution
385128008 | Prolonged-release eye drops
385130005 | Eye lotion
385132002 | Ophthalmic insert
385133007 | Ear cream
385134001 | Ear gel
385135000 | Ear ointment
385136004 | Ear drops
385137008 | Ear drops solution
385138003 | Ear drops emulsion
385139006 | Ear powder
385140008 | Ear spray
385141007 | Ear spray solution
385142000 | Ear spray suspension
385143005 | Ear spray emulsion
385147006 | Ear tampon
385148001 | Ear stick
385149009 | Nasal cream
385150009 | Nasal gel
385151008 | Nasal ointment
385152001 | Nasal drops
385153006 | Nasal drops solution
385154000 | Nasal drops suspension
385155004 | Nasal drops emulsion
385156003 | Nasal powder
385157007 | Nasal spray
385158002 | Nasal spray solution
385159005 | Nasal spray suspension
385160000 | Nasal spray emulsion
385162008 | Nasal stick

385163003 | Eye/ear/nose drops
385164009 | Eye/ear/nose ointment
385165005 | Vaginal cream
385166006 | Vaginal gel
385167002 | Vaginal ointment
385168007 | Vaginal foam
385169004 | Vaginal liquid
385170003 | Vaginal solution
385171004 | Vaginal suspension
385172006 | Vaginal emulsion
385173001 | Tablet for vaginal solution
385174007 | Pessary
385175008 | Vaginal capsule
385176009 | Hard vaginal capsule
385177000 | Soft vaginal capsule
385178005 | Vaginal tablet
385179002 | Effervescent vaginal tablet
385180004 | Medicated vaginal tampon
385182007 | Rectal cream
385183002 | Rectal gel
385184008 | Rectal ointment
385185009 | Rectal foam
385186005 | Enema
385187001 | Rectal solution
385188006 | Rectal suspension
385189003 | Rectal emulsion
385191006 | Powder for rectal suspension
385192004 | Tablet for rectal solution
385193009 | Tablet for rectal suspension
385194003 | Suppository
385195002 | Rectal capsule
385196001 | Rectal tampon
385197005 | Nebulizer liquid
385198000 | Nebulizer solution
385199008 | Nebulizer suspension
385200006 | Powder for nebulizer suspension
385201005 | Powder for nebulizer solution
385202003 | Nebulizer emulsion
385203008 | Pressurised inhalation
385204002 | Pressurised inhalation solution
385205001 | Pressurised inhalation suspension
385206000 | Pressurised inhalation emulsion
385207009 | Inhalation powder
385208004 | Hard capsule inhalation powder
385210002 | Inhalation vapor
385211003 | Inhalation vapor powder
385212005 | Inhalation vapor capsule
385213000 | Inhalation vapor solution
385214006 | Inhalation vapor tablet
385215007 | Inhalation vapor ointment
385216008 | Inhalation vapor liquid
385217004 | Inhalation gas
385219001 | Injection solution
385220007 | Injection suspension
385221006 | Injection emulsion
385222004 | Injection powder
385223009 | Powder for injection solution
385224003 | Powder for injection suspension
385229008 | Infusion solution
385230003 | Infusion powder
385231004 | Powder for infusion solution
385236009 | Implantation tablet
385237000 | Implantation chain
385242008 | Intravesical solution
385245005 | Urethral gel
385246006 | Urethral stick
385247002 | Endotracheopulmonary instillation solution
385248007 | Powder for endotracheopulmonary instillation solution
385250004 | Endotracheopulmonary instillation suspension
385251000 | Endocervical gel
385257001 | Gastroenteral liquid
385258006 | Gastroenteral solution

385259003 | Gastroenteral suspension
385260008 | Gastroenteral emulsion
385261007 | Wound stick
385262000 | Organ preservation solution
385278003 | Cutaneous powder spray
385279006 | Cutaneous suspension spray
385286003 | Implant dosage form
414951009 | Oral elixir
420243009 | Tampon dose form
420253005 | Oil injection
420275007 | Semi-solid dose form
420283001 | Nasal emulsion
420292003 | Rectal powder
420293008 | Coated pellets capsule
420305009 | Powder for oral liquid
420317006 | Inhaler
420378007 | Extended-release film coated tablet
420385006 | Extended-release insert
420386007 | Emulsion
420407000 | Inhalation aerosol solution
420430006 | Lyophilized powder for injectable extended release liposomal suspension
420450005 | Pressurized nebulizer suspension
420460001 | Film
420509004 | Oral granules
420536002 | Vaginal powder
420540006 | Cutaneous oil
420610000 | Nasal aerosol
420627008 | Extended-release tablet
420631002 | Vaginal insert
420634005 | Microspheres for injectable suspension
420636007 | Eye/ear drops
420641004 | Solution for inhalation
420656008 | Lyophilized powder for injectable liposomal suspension
420692007 | Oral capsule
420699003 | Liquid dose form
420705007 | Inhalation aerosol suspension
420736004 | Eye suspension
420757007 | Delayed-release granules
420761001 | Urethral suppository
420767002 | Delayed-release pellets capsule
420768007 | Pellet
420802004 | Extended-release liquid
420828001 | Powder for reconstitution for drug product
420847003 | Metered dose aerosol
420873008 | Extended release injectable suspension
420887008 | Intrathecal suspension
420891003 | Ear drops suspension
420901005 | Ear emulsion
420927005 | Metered powder
420929008 | Rectal suppository
420955009 | Powder for solution
420956005 | Ultramicronized tablet
420992009 | Implantable pellet
421026006 | Oral tablet
421027002 | Delayed-release capsule
421034000 | Extended-release suppository
421043009 | Extended-release film
421051007 | Sonicated injectable suspension
421056002 | Emulsion for inhalation
421079001 | Pastille
421080003 | Powder for suspension
421104008 | Aerosol powder
421131006 | Gaseous dose form
421155001 | Extended-release enteric coated tablet
421166008 | Foam
421195004 | Vaginal suppository
421221008 | Suspension for inhalation
421271006 | Extended-release bead implant
421288004 | Sponge
421300005 | Extended-release film coated capsule
421316007 | Powder for eye drops
421338009 | Extended-release coated capsule

421340004 | Powder spray
421343002 | Powder for cutaneous solution
421347001 | Cutaneous aerosol
421366001 | Tablet for oral suspension
421374000 | Delayed-release tablet
421378002 | Solid dose form
421382000 | Ear suspension
421410002 | Intravenous solution
421425002 | Powder for inhalation solution
421427005 | Eye/ear ointment
421428000 | Injectable oil suspension
421446006 | Extended-release granules for suspension
421504000 | Cone
421522002 | Liposomal injectable suspension
421532009 | Insert
421535006 | Delayed-release particles tablet
421575003 | Powder for ear drops
421606006 | Aerosol spray
421607002 | Powder for nasal drops
421618002 | Extended-release capsule
421620004 | Sustained-release buccal tablet
421628006 | Cutaneous cream
421637006 | Lyophilized powder for injectable solution
421669002 | Metered gel
421701006 | Tablet for oral solution
421713001 | Ear solution
421716009 | Transdermal drug delivery system
421720008 | Spray dose form
421721007 | Coated particles tablet
421752008 | Extended-release enteric coated capsule
421759004 | Metered dose aerosol inhaler
421765004 | Intraperitoneal solution
421857007 | Powder for ophthalmic solution
421873001 | Oral cream
421890007 | Oil
421932003 | Multilayer tablet
421937009 | Medicated toothpaste
421943006 | Lyophilized powder for injectable suspension
421949005 | Cutaneous gel
421987002 | Intraocular solution
421999009 | Extended-release coated pellets
422054001 | Metered dose powder inhaler
422059006 | Metered dose inhaler
422060001 | Ophthalmic solution
422068008 | Eye drops suspension
422080000 | Nasal suspension
422085005 | Powder for ophthalmic suspension
422151007 | Breath activated powder inhaler
422186009 | Tincture
422197009 | Breath activated inhaler
422199007 | Dental cone
422201009 | Tablet for cutaneous solution
422202002 | Colloidal suspension for injection
422259002 | Spirit
422264003 | Powder for injectable extended release suspension
422301006 | Modified-release pessary
422336005 | Nasal solution
422353003 | Extended-release suspension
424179000 | Aerosol generator
424552006 | Vaginal sponge
425753008 | Topical ointment
425965000 | Rectal spray
426210003 | Gum
426684005 | Drops dose form
426823003 | Vaginal spray
426969004 | Metered spray
427129005 | Coated capsule
427564005 | Pressurized spray
427609008 | Modified release drops dose form
429885007 | Bar
443424002 | Buccal film
447050008 | Orodispersible film

447079001 | Orodispersible tablet

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.563 Value Set http://hl7.org/fhir/ValueSet/all-types

URL: https://hl7.org/fhir/R4/valueset-all-types.html
FHIR Infrastructure Work Group | Maturity Level: N | Normative (from v4.0.0) | Use Context: Any

| This page has been approved as part of an ANSI standard. See the Conformance Package for further details.

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/all-types
Version: | 4.0.1
Name: | FHIRAllTypes
Title: | FHIRAllTypes
Definition: | A list of all the concrete types defined in this version of the FHIR specification - Abstract Types, Data Types and Resource Types.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.25 (for OID based terminology systems)
Source Resource | XML / JSON

This Value Set is normative - it is generated based on the information defined in this specification. The definition will remain fixed across versions, but the actual contents will change from version to version

This value set is used in the following places:

• Resource: OperationDefinition.parameter.type (code / Required)
• Resource: DataRequirement.type (code / Required)
• Resource: ParameterDefinition.type (code / Required)

4.4.1.563.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/data-types
• Include all codes defined in http://hl7.org/fhir/resource-types
• Include all codes defined in http://hl7.org/fhir/abstract-types

4.4.1.563.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 213 concepts

Expansion based on http://hl7.org/fhir/resource-types version 4.0.1, http://hl7.org/fhir/abstract-types version 4.0.1, http://hl7.org/fhir/data-types version 4.0.1

Code | System | Display | Definition
Address | http://hl7.org/fhir/data-types | Address | An address expressed using postal conventions (as opposed to GPS or other location definition formats). This data type may be used to convey addresses for use in delivering mail as well as for visiting locations which might not be valid for mail delivery. There are a variety of postal address formats defined around the world.
Age | http://hl7.org/fhir/data-types | Age | A duration of time during which an organism (or a process) has existed.
Annotation | http://hl7.org/fhir/data-types | Annotation | A text note which also contains information about who made the statement and when.
Attachment | http://hl7.org/fhir/data-types | Attachment | For referring to data content defined in other formats.
BackboneElement | http://hl7.org/fhir/data-types | BackboneElement | Base definition for all elements that are defined inside a resource - but not those in a data type.
CodeableConcept | http://hl7.org/fhir/data-types | CodeableConcept | A concept that may be defined by a formal reference to a terminology or ontology or may be provided by text.
Coding | http://hl7.org/fhir/data-types | Coding | A reference to a code defined by a terminology system.
ContactDetail | http://hl7.org/fhir/data-types | ContactDetail | Specifies contact information for a person or organization.
ContactPoint | http://hl7.org/fhir/data-types | ContactPoint | Details for all kinds of technology mediated contact points for a person or organization, including telephone, email, etc.
Contributor | http://hl7.org/fhir/data-types | Contributor | A contributor to the content of a knowledge asset, including authors, editors, reviewers, and endorsers.

Count | http://hl7.org/fhir/data-types | Count | A measured amount (or an amount that can potentially be measured). Note that measured amounts include amounts that are not precisely quantified, including amounts involving arbitrary units and floating currencies.

DataRequirement | http://hl7.org/fhir/data-types | DataRequirement | Describes a required data item for evaluation in terms of the type of data, and optional code or date-based filters of the data.

Distance | http://hl7.org/fhir/data-types | Distance | A length - a value with a unit that is a physical distance.

Dosage | http://hl7.org/fhir/data-types | Dosage | Indicates how the medication is/was taken or should be taken by the patient.

Duration | http://hl7.org/fhir/data-types | Duration | A length of time.

Element | http://hl7.org/fhir/data-types | Element | Base definition for all elements in a resource.

ElementDefinition | http://hl7.org/fhir/data-types | ElementDefinition | Captures constraints on each element within the resource, profile, or extension.

Expression | http://hl7.org/fhir/data-types | Expression | A expression that is evaluated in a specified context and returns a value. The context of use of the expression must specify the context in which the expression is evaluated, and how the result of the expression is used.

Extension | http://hl7.org/fhir/data-types | Extension | Optional Extension Element - found in all resources.

HumanName | http://hl7.org/fhir/data-types | HumanName | A human's name with the ability to identify parts and usage.

Identifier | http://hl7.org/fhir/data-types | Identifier | An identifier - identifies some entity uniquely and unambiguously. Typically this is used for business identifiers.

MarketingStatus | http://hl7.org/fhir/data-types | MarketingStatus | The marketing status describes the date when a medicinal product is actually put on the market or the date as of which it is no longer available.

Meta | http://hl7.org/fhir/data-types | Meta | The metadata about a resource. This is content in the resource that is maintained by the infrastructure. Changes to the content might not always be associated with version changes to the resource.

Money | http://hl7.org/fhir/data-types | Money | An amount of economic utility in some recognized currency.

MoneyQuantity | http://hl7.org/fhir/data-types | MoneyQuantity

Narrative | http://hl7.org/fhir/data-types | Narrative | A human-readable summary of the resource conveying the essential clinical and business information for the resource.

ParameterDefinition | http://hl7.org/fhir/data-types | ParameterDefinition | The parameters to the module. This collection specifies both the input and output parameters. Input parameters are provided by the caller as part of the $evaluate operation. Output parameters are included in the GuidanceResponse.

Period | http://hl7.org/fhir/data-types | Period | A time period defined by a start and end date and optionally time.

Population | http://hl7.org/fhir/data-types | Population | A populatioof people with some set of grouping criteria.

ProdCharacteristic | http://hl7.org/fhir/data-types | ProdCharacteristic | The marketing status describes the date when a medicinal product is actually put on the market or the date as of which it is no longer available.

ProductShelfLife | http://hl7.org/fhir/data-types | ProductShelfLife | The shelf-life and storage information for a medicinal product item or container can be described using this class.

Quantity | http://hl7.org/fhir/data-types | Quantity | A measured amount (or an amount that can potentially be measured). Note that measured amounts include amounts that are not precisely quantified, including amounts involving arbitrary units and floating currencies.

Range | http://hl7.org/fhir/data-types | Range | A set of ordered Quantities defined by a low and high limit.

Ratio | http://hl7.org/fhir/data-types | Ratio | A relationship of two Quantity values - expressed as a numerator and a denominator.

Reference | http://hl7.org/fhir/data-types | Reference | A reference from one resource to another.

RelatedArtifact | http://hl7.org/fhir/data-types | RelatedArtifact | Related artifacts such as additional documentation, justification, or bibliographic references.

SampledData | http://hl7.org/fhir/data-types | SampledData | A series of measurements taken by a device, with upper and lower limits. There may be more than one dimension in the data.

Signature | http://hl7.org/fhir/data-types | Signature | A signature along with supporting context. The signature may be a digital signature that is cryptographic in nature, or some other signature acceptable to the domain. This other signature may be as simple as a graphical image representing a hand-written signature, or a signature ceremony Different signature approaches have different utilities.

SimpleQuantity | http://hl7.org/fhir/data-types | SimpleQuantity

SubstanceAmount | http://hl7.org/fhir/data-types | SubstanceAmount | Chemical substances are a single substance type whose primary defining element is the molecular structure. Chemical substances shall be defined on the basis of their complete covalent molecular structure; the presence of a salt (counter-ion) and/or solvates (water, alcohols) is also captured. Purity, grade, physical form or particle size are not taken into account in the definition of a chemical substance or in the assignment of a Substance ID.

Timing | http://hl7.org/fhir/data-types | Timing | Specifies an event that may occur multiple times. Timing schedules are used to record when things are planned, expected or requested to occur. The most common usage is in dosage instructions for medications. They are also used when planning care of various kinds, and may be used for reporting the schedule to which past regular activities were carried out.

TriggerDefinition | http://hl7.org/fhir/data-types | TriggerDefinition | A description of a triggering event. Triggering events can be named events, data events, or periodic, as determined by the type element.

UsageContext | http://hl7.org/fhir/data-types | UsageContext | Specifies clinical/business/etc. metadata that can be used to retrieve, index and/or categorize an artifact. This metadata can either be specific to the applicable population (e.g., age category, DRG) or the specific context of care (e.g., venue, care setting, provider of care).

base64Binary | http://hl7.org/fhir/data-types | base64Binary | A stream of bytes

boolean | http://hl7.org/fhir/data-types | boolean | Value of "true" or "false"

canonical | http://hl7.org/fhir/data-types | canonical | A URI that is a reference to a canonical URL on a FHIR resource

code | http://hl7.org/fhir/data-types | code | A string which has at least one character and no leading or trailing whitespace and where there is no whitespace other than single spaces in the contents

date | http://hl7.org/fhir/data-types | date | A date or partial date (e.g. just year or year + month). There is no time zone. The format is a union of the schema types gYear, gYearMonth and date. Dates SHALL be valid dates.

dateTime | http://hl7.org/fhir/data-types | dateTime | A date, date-time or partial date (e.g. just year or year + month). If hours and minutes are specified, a time zone SHALL be populated. The format is a union of the schema types gYear, gYearMonth, date and dateTime. Seconds must be provided due to schema type constraints but may be zero-filled and may be ignored. Dates SHALL be valid dates.

decimal | http://hl7.org/fhir/data-types | decimal | A rational number with implicit precision

id | http://hl7.org/fhir/data-types | id | Any combination of letters, numerals, "-" and ".", with a length limit of 64 characters. (This might be an integer, an unprefixed OID, UUID or any other identifier pattern that meets these constraints.) Ids are case-insensitive.
instant | http://hl7.org/fhir/data-types | instant | An instant in time - known at least to the second
integer | http://hl7.org/fhir/data-types | integer | A whole number
markdown | http://hl7.org/fhir/data-types | markdown | A string that may contain Github Flavored Markdown syntax for optional processing by a mark down presentation engine
oid | http://hl7.org/fhir/data-types | oid | An OID represented as a URI
positiveInt | http://hl7.org/fhir/data-types | positiveInt | An integer with a value that is positive (e.g. >0)
string | http://hl7.org/fhir/data-types | string | A sequence of Unicode characters
time | http://hl7.org/fhir/data-types | time | A time during the day, with no date specified
unsignedInt | http://hl7.org/fhir/data-types | unsignedInt | An integer with a value that is not negative (e.g. >= 0)
uri | http://hl7.org/fhir/data-types | uri | String of characters used to identify a name or a resource
url | http://hl7.org/fhir/data-types | url | A URI that is a literal reference
uuid | http://hl7.org/fhir/data-types | uuid | A UUID, represented as a URI
xhtml | http://hl7.org/fhir/data-types | XHTML | XHTML format, as defined by W3C, but restricted usage (mainly, no active content)
Account | http://hl7.org/fhir/resource-types | Account | A financial tool for tracking value accrued for a particular purpose. In the healthcare field, used to track charges for a patient, cost centers, etc.
ActivityDefinition | http://hl7.org/fhir/resource-types | ActivityDefinition | This resource allows for the definition of some activity to be performed, independent of a particular patient, practitioner, or other performance context.
AdverseEvent | http://hl7.org/fhir/resource-types | AdverseEvent | Actual or potential/avoided event causing unintended physical injury resulting from or contributed to by medical care, a research study or other healthcare setting factors that requires additional monitoring, treatment, or hospitalization, or that results in death.
AllergyIntolerance | http://hl7.org/fhir/resource-types | AllergyIntolerance | Risk of harmful or undesirable, physiological response which is unique to an individual and associated with exposure to a substance.
Appointment | http://hl7.org/fhir/resource-types | Appointment | A booking of a healthcare event among patient(s), practitioner(s), related person(s) and/or device(s) for a specific date/time. This may result in one or more Encounter(s).
AppointmentResponse | http://hl7.org/fhir/resource-types | AppointmentResponse | A reply to an appointment request for a patient and/or practitioner(s), such as a confirmation or rejection.
AuditEvent | http://hl7.org/fhir/resource-types | AuditEvent | A record of an event made for purposes of maintaining a security log. Typical uses include detection of intrusion attempts and monitoring for inappropriate usage.
Basic | http://hl7.org/fhir/resource-types | Basic | Basic is used for handling concepts not yet defined in FHIR, narrative-only resources that don't map to an existing resource, and custom resources not appropriate for inclusion in the FHIR specification.
Binary | http://hl7.org/fhir/resource-types | Binary | A resource that represents the data of a single raw artifact as digital content accessible in its native format. A Binary resource can contain any content, whether text, image, pdf, zip archive, etc.
BiologicallyDerivedProduct | http://hl7.org/fhir/resource-types | BiologicallyDerivedProduct | A material substance originating from a biological entity intended to be transplanted or infused into another (possibly the same) biological entity.
BodyStructure | http://hl7.org/fhir/resource-types | BodyStructure | Record details about an anatomical structure. This resource may be used when a coded concept does not provide the necessary detail needed for the use case.
Bundle | http://hl7.org/fhir/resource-types | Bundle | A container for a collection of resources.
CapabilityStatement | http://hl7.org/fhir/resource-types | CapabilityStatement | A Capability Statement documents a set of capabilities (behaviors) of a FHIR Server for a particular version of FHIR that may be used as a statement of actual server functionality or a statement of required or desired server implementation.
CarePlan | http://hl7.org/fhir/resource-types | CarePlan | Describes the intention of how one or more practitioners intend to deliver care for a particular patient, group or community for a period of time, possibly limited to care for a specific condition or set of conditions.
CareTeam | http://hl7.org/fhir/resource-types | CareTeam | The Care Team includes all the people and organizations who plan to participate in the coordination and delivery of care for a patient.
CatalogEntry | http://hl7.org/fhir/resource-types | CatalogEntry | Catalog entries are wrappers that contextualize items included in a catalog.
ChargeItem | http://hl7.org/fhir/resource-types | ChargeItem | The resource ChargeItem describes the provision of healthcare provider products for a certain patient, therefore referring not only to the product, but containing in addition details of the provision, like date, time, amounts and participating organizations and persons. Main Usage of the ChargeItem is to enable the billing process and internal cost allocation.
ChargeItemDefinition | http://hl7.org/fhir/resource-types | ChargeItemDefinition | The ChargeItemDefinition resource provides the properties that apply to the (billing) codes necessary to calculate costs and prices. The properties may differ largely depending on type and realm, therefore this resource gives only a rough structure and requires profiling for each type of billing code system.
Claim | http://hl7.org/fhir/resource-types | Claim | A provider issued list of professional services and products which have been provided, or are to be provided, to a patient which is sent to an insurer for reimbursement.
ClaimResponse | http://hl7.org/fhir/resource-types | ClaimResponse | This resource provides the adjudication details from the processing of a Claim resource.
ClinicalImpression | http://hl7.org/fhir/resource-types | ClinicalImpression | A record of a clinical assessment performed to determine what problem(s) may affect the patient and before planning the treatments or management strategies that are best to manage a patient's condition. Assessments are often 1:1 with a clinical consultation / encounter, but this varies greatly depending on the clinical workflow. This resource is called "ClinicalImpression" rather than "ClinicalAssessment" to avoid confusion with the recording of assessment tools such as Apgar score.
CodeSystem | http://hl7.org/fhir/resource-types | CodeSystem | The CodeSystem resource is used to declare the existence of and describe a code system or code system supplement and its key properties, and optionally define a part or all of its content.
Communication | http://hl7.org/fhir/resource-types | Communication | An occurrence of information being transmitted; e.g. an alert that was sent to a responsible provider, a public health agency that was notified about a reportable condition.
CommunicationRequest | http://hl7.org/fhir/resource-types | CommunicationRequest | A request to convey information; e.g. the CDS system proposes that an alert be sent to a responsible provider, the CDS system proposes that the public health agency be notified about a reportable condition.
CompartmentDefinition | http://hl7.org/fhir/resource-types | CompartmentDefinition | A compartment definition that defines how resources are accessed on a server.

Composition | http://hl7.org/fhir/resource-types | Composition | A set of healthcare-related information that is assembled together into a single logical package that provides a single coherent statement of meaning, establishes its own context and that has clinical attestation with regard to who is making the statement. A Composition defines the structure and narrative content necessary for a document. However, a Composition alone does not constitute a document. Rather, the Composition must be the first entry in a Bundle where Bundle.type=document, and any other resources referenced from Composition must be included as subsequent entries in the Bundle (for example Patient, Practitioner, Encounter, etc.).

ConceptMap | http://hl7.org/fhir/resource-types | ConceptMap | A statement of relationships from one set of concepts to one or more other concepts - either concepts in code systems, or data element/data element concepts, or classes in class models.

Condition | http://hl7.org/fhir/resource-types | Condition | A clinical condition, problem, diagnosis, or other event, situation, issue, or clinical concept that has risen to a level of concern.

Consent | http://hl7.org/fhir/resource-types | Consent | A record of a healthcare consumer’s choices, which permits or denies identified recipient(s) or recipient role(s) to perform one or more actions within a given policy context, for specific purposes and periods of time.

Contract | http://hl7.org/fhir/resource-types | Contract | Legally enforceable, formally recorded unilateral or bilateral directive i.e., a policy or agreement.

Coverage | http://hl7.org/fhir/resource-types | Coverage | Financial instrument which may be used to reimburse or pay for health care products and services. Includes both insurance and self-payment.

CoverageEligibilityRequest | http://hl7.org/fhir/resource-types | CoverageEligibilityRequest | The CoverageEligibilityRequest provides patient and insurance coverage information to an insurer for them to respond, in the form of an CoverageEligibilityResponse, with information regarding whether the stated coverage is valid and in-force and optionally to provide the insurance details of the policy.

CoverageEligibilityResponse | http://hl7.org/fhir/resource-types | CoverageEligibilityResponse | This resource provides eligibility and plan details from the processing of an CoverageEligibilityRequest resource.

DetectedIssue | http://hl7.org/fhir/resource-types | DetectedIssue | Indicates an actual or potential clinical issue with or between one or more active or proposed clinical actions for a patient; e.g. Drug-drug interaction, Ineffective treatment frequency, Procedure-condition conflict, etc.

Device | http://hl7.org/fhir/resource-types | Device | A type of a manufactured item that is used in the provision of healthcare without being substantially changed through that activity. The device may be a medical or non-medical device.

DeviceDefinition | http://hl7.org/fhir/resource-types | DeviceDefinition | The characteristics, operational status and capabilities of a medical-related component of a medical device.

DeviceMetric | http://hl7.org/fhir/resource-types | DeviceMetric | Describes a measurement, calculation or setting capability of a medical device.

DeviceRequest | http://hl7.org/fhir/resource-types | DeviceRequest | Represents a request for a patient to employ a medical device. The device may be an implantable device, or an external assistive device, such as a walker.

DeviceUseStatement | http://hl7.org/fhir/resource-types | DeviceUseStatement | A record of a device being used by a patient where the record is the result of a report from the patient or another clinician.

DiagnosticReport | http://hl7.org/fhir/resource-types | DiagnosticReport | The findings and interpretation of diagnostic tests performed on patients, groups of patients, devices, and locations, and/or specimens derived from these. The report includes clinical context such as requesting and provider information, and some mix of atomic results, images, textual and coded interpretations, and formatted representation of diagnostic reports.

DocumentManifest | http://hl7.org/fhir/resource-types | DocumentManifest | A collection of documents compiled for a purpose together with metadata that applies to the collection.

DocumentReference | http://hl7.org/fhir/resource-types | DocumentReference | A reference to a document of any kind for any purpose. Provides metadata about the document so that the document can be discovered and managed. The scope of a document is any seralized object with a mime-type, so includes formal patient centric documents (CDA), cliical notes, scanned paper, and non-patient specific documents like policy text.

DomainResource | http://hl7.org/fhir/resource-types | DomainResource | A resource that includes narrative, extensions, and contained resources.

EffectEvidenceSynthesis | http://hl7.org/fhir/resource-types | EffectEvidenceSynthesis | The EffectEvidenceSynthesis resource describes the difference in an outcome between exposures states in a population where the effect estimate is derived from a combination of research studies.

Encounter | http://hl7.org/fhir/resource-types | Encounter | An interaction between a patient and healthcare provider(s) for the purpose of providing healthcare service(s) or assessing the health status of a patient.

Endpoint | http://hl7.org/fhir/resource-types | Endpoint | The technical details of an endpoint that can be used for electronic services, such as for web services providing XDS.b or a REST endpoint for another FHIR server. This may include any security context information.

EnrollmentRequest | http://hl7.org/fhir/resource-types | EnrollmentRequest | This resource provides the insurance enrollment details to the insurer regarding a specified coverage.

EnrollmentResponse | http://hl7.org/fhir/resource-types | EnrollmentResponse | This resource provides enrollment and plan details from the processing of an EnrollmentRequest resource.

EpisodeOfCare | http://hl7.org/fhir/resource-types | EpisodeOfCare | An association between a patient and an organization / healthcare provider(s) during which time encounters may occur. The managing organization assumes a level of responsibility for the patient during this time.

EventDefinition | http://hl7.org/fhir/resource-types | EventDefinition | The EventDefinition resource provides a reusable description of when a particular event can occur.

Evidence | http://hl7.org/fhir/resource-types | Evidence | The Evidence resource describes the conditional state (population and any exposures being compared within the population) and outcome (if specified) that the knowledge (evidence, assertion, recommendation) is about.

EvidenceVariable | http://hl7.org/fhir/resource-types | EvidenceVariable | The EvidenceVariable resource describes a "PICO" element that knowledge (evidence, assertion, recommendation) is about.

ExampleScenario | http://hl7.org/fhir/resource-types | ExampleScenario | Example of workflow instance.

ExplanationOfBenefit | http://hl7.org/fhir/resource-types | ExplanationOfBenefit | This resource provides: the claim details; adjudication details from the processing of a Claim; and optionally account balance information, for informing the subscriber of the benefits provided.

FamilyMemberHistory | http://hl7.org/fhir/resource-types | FamilyMemberHistory | Significant health conditions for a person related to the patient relevant in the context of care for the patient.

Flag | http://hl7.org/fhir/resource-types | Flag | Prospective warnings of potential issues when providing care to the patient.

Goal | http://hl7.org/fhir/resource-types | Goal | Describes the intended objective(s) for a patient, group or organization care, for example, weight loss, restoring an activity of daily living, obtaining herd immunity via immunization, meeting a process improvement objective, etc.

GraphDefinition | http://hl7.org/fhir/resource-types | GraphDefinition | A formal computable definition of a graph of resources - that is, a coherent set of resources that form a graph by following references. The Graph Definition resource defines a set and makes rules about the set.

Group | http://hl7.org/fhir/resource-types | Group | Represents a defined collection of entities that may be discussed or acted upon collectively but which are not expected to act collectively, and are not formally or legally recognized; i.e. a collection of entities that isn't an Organization.

GuidanceResponse | http://hl7.org/fhir/resource-types | GuidanceResponse | A guidance response is the formal response to a guidance request, including any output parameters returned by the evaluation, as well as the description of any proposed actions to be taken.

HealthcareService | http://hl7.org/fhir/resource-types | HealthcareService | The details of a healthcare service available at a location.

ImagingStudy | http://hl7.org/fhir/resource-types | ImagingStudy | Representation of the content produced in a DICOM imaging study. A study comprises a set of series, each of which includes a set of Service-Object Pair Instances (SOP Instances - images or other data) acquired or produced in a common context. A series is of only one modality (e.g. X-ray, CT, MR, ultrasound), but a study may have multiple series of different modalities.

Immunization | http://hl7.org/fhir/resource-types | Immunization | Describes the event of a patient being administered a vaccine or a record of an immunization as reported by a patient, a clinician or another party.

ImmunizationEvaluation | http://hl7.org/fhir/resource-types | ImmunizationEvaluation | Describes a comparison of an immunization event against published recommendations to determine if the administration is "valid" in relation to those recommendations.

ImmunizationRecommendation | http://hl7.org/fhir/resource-types | ImmunizationRecommendation | A patient's point-in-time set of recommendations (i.e. forecasting) according to a published schedule with optional supporting justification.

ImplementationGuide | http://hl7.org/fhir/resource-types | ImplementationGuide | A set of rules of how a particular interoperability or standards problem is solved - typically through the use of FHIR resources. This resource is used to gather all the parts of an implementation guide into a logical whole and to publish a computable definition of all the parts.

InsurancePlan | http://hl7.org/fhir/resource-types | InsurancePlan | Details of a Health Insurance product/plan provided by an organization.

Invoice | http://hl7.org/fhir/resource-types | Invoice | Invoice containing collected ChargeItems from an Account with calculated individual and total price for Billing purpose.

Library | http://hl7.org/fhir/resource-types | Library | The Library resource is a general-purpose container for knowledge asset definitions. It can be used to describe and expose existing knowledge assets such as logic libraries and information model descriptions, as well as to describe a collection of knowledge assets.

Linkage | http://hl7.org/fhir/resource-types | Linkage | Identifies two or more records (resource instances) that refer to the same real-world "occurrence".

List | http://hl7.org/fhir/resource-types | List | A list is a curated collection of resources.

Location | http://hl7.org/fhir/resource-types | Location | Details and position information for a physical place where services are provided and resources and participants may be stored, found, contained, or accommodated.

Measure | http://hl7.org/fhir/resource-types | Measure | The Measure resource provides the definition of a quality measure.

MeasureReport | http://hl7.org/fhir/resource-types | MeasureReport | The MeasureReport resource contains the results of the calculation of a measure; and optionally a reference to the resources involved in that calculation.

Media | http://hl7.org/fhir/resource-types | Media | A photo, video, or audio recording acquired or used in healthcare. The actual content may be inline or provided by direct reference.

Medication | http://hl7.org/fhir/resource-types | Medication | This resource is primarily used for the identification and definition of a medication for the purposes of prescribing, dispensing, and administering a medication as well as for making statements about medication use.

MedicationAdministration | http://hl7.org/fhir/resource-types | MedicationAdministration | Describes the event of a patient consuming or otherwise being administered a medication. This may be as simple as swallowing a tablet or it may be a long running infusion. Related resources tie this event to the authorizing prescription, and the specific encounter between patient and health care practitioner.

MedicationDispense | http://hl7.org/fhir/resource-types | MedicationDispense | Indicates that a medication product is to be or has been dispensed for a named person/patient. This includes a description of the medication product (supply) provided and the instructions for administering the medication. The medication dispense is the result of a pharmacy system responding to a medication order.

MedicationKnowledge | http://hl7.org/fhir/resource-types | MedicationKnowledge | Information about a medication that is used to support knowledge.

MedicationRequest | http://hl7.org/fhir/resource-types | MedicationRequest | An order or request for both supply of the medication and the instructions for administration of the medication to a patient. The resource is called "MedicationRequest" rather than "MedicationPrescription" or "MedicationOrder" to generalize the use across inpatient and outpatient settings, including care plans, etc., and to harmonize with workflow patterns.

MedicationStatement | http://hl7.org/fhir/resource-types | MedicationStatement | A record of a medication that is being consumed by a patient. A MedicationStatement may indicate that the patient may be taking the medication now or has taken the medication in the past or will be taking the medication in the future. The source of this information can be the patient, significant other (such as a family member or spouse), or a clinician. A common scenario where this information is captured is during the history taking process during a patient visit or stay. The medication information may come from sources such as the patient's memory, from a prescription bottle, or from a list of medications the patient, clinician or other party maintains. The primary difference between a medication statement and a medication administration is that the medication administration has complete administration information and is based on actual administration information from the person who administered the medication. A medication statement is often, if not always, less specific. There is no required date/time when the medication was administered, in fact we only know that a source has reported the patient is taking this medication, where details such as time, quantity, or rate or even medication product may be incomplete or missing or less precise. As stated earlier, the medication statement information may come from the patient's memory, from a prescription bottle or from a list of medications the patient, clinician or other party maintains. Medication administration is more formal and is not missing detailed information.

MedicinalProduct | http://hl7.org/fhir/resource-types | MedicinalProduct | Detailed definition of a medicinal product, typically for uses other than direct patient care (e.g. regulatory use).

MedicinalProductAuthorization | http://hl7.org/fhir/resource-types | MedicinalProductAuthorization | The regulatory authorization of a medicinal product.
MedicinalProductContraindication | http://hl7.org/fhir/resource-types | MedicinalProductContraindication | The clinical particulars - indications, contraindications etc. of a medicinal product, including for regulatory purposes.
MedicinalProductIndication | http://hl7.org/fhir/resource-types | MedicinalProductIndication | Indication for the Medicinal Product.
MedicinalProductIngredient | http://hl7.org/fhir/resource-types | MedicinalProductIngredient | An ingredient of a manufactured item or pharmaceutical product.
MedicinalProductInteraction | http://hl7.org/fhir/resource-types | MedicinalProductInteraction | The interactions of the medicinal product with other medicinal products, or other forms of interactions.
MedicinalProductManufactured | http://hl7.org/fhir/resource-types | MedicinalProductManufactured | The manufactured item as contained in the packaged medicinal product.
MedicinalProductPackaged | http://hl7.org/fhir/resource-types | MedicinalProductPackaged | A medicinal product in a container or package.
MedicinalProductPharmaceutical | http://hl7.org/fhir/resource-types | MedicinalProductPharmaceutical | A pharmaceutical product described in terms of its composition and dose form.
MedicinalProductUndesirableEffect | http://hl7.org/fhir/resource-types | MedicinalProductUndesirableEffect | Describe the undesirable effects of the medicinal product.
MessageDefinition | http://hl7.org/fhir/resource-types | MessageDefinition | Defines the characteristics of a message that can be shared between systems, including the type of event that initiates the message, the content to be transmitted and what response(s), if any, are permitted.
MessageHeader | http://hl7.org/fhir/resource-types | MessageHeader | The header for a message exchange that is either requesting or responding to an action. The reference(s) that are the subject of the action as well as other information related to the action are typically transmitted in a bundle in which the MessageHeader resource instance is the first resource in the bundle.
MolecularSequence | http://hl7.org/fhir/resource-types | MolecularSequence | Raw data describing a biological sequence.
NamingSystem | http://hl7.org/fhir/resource-types | NamingSystem | A curated namespace that issues unique symbols within that namespace for the identification of concepts, people, devices, etc. Represents a "System" used within the Identifier and Coding data types.
NutritionOrder | http://hl7.org/fhir/resource-types | NutritionOrder | A request to supply a diet, formula feeding (enteral) or oral nutritional supplement to a patient/resident.
Observation | http://hl7.org/fhir/resource-types | Observation | Measurements and simple assertions made about a patient, device or other subject.
ObservationDefinition | http://hl7.org/fhir/resource-types | ObservationDefinition | Set of definitional characteristics for a kind of observation or measurement produced or consumed by an orderable health care service.
OperationDefinition | http://hl7.org/fhir/resource-types | OperationDefinition | A formal computable definition of an operation (on the RESTful interface) or a named query (using the search interaction).
OperationOutcome | http://hl7.org/fhir/resource-types | OperationOutcome | A collection of error, warning, or information messages that result from a system action.
Organization | http://hl7.org/fhir/resource-types | Organization | A formally or informally recognized grouping of people or organizations formed for the purpose of achieving some form of collective action. Includes companies, institutions, corporations, departments, community groups, healthcare practice groups, payer/insurer, etc.
OrganizationAffiliation | http://hl7.org/fhir/resource-types | OrganizationAffiliation | Defines an affiliation/assotiation/relationship between 2 distinct oganizations, that is not a part-of relationship/sub-division relationship.
Parameters | http://hl7.org/fhir/resource-types | Parameters | This resource is a non-persisted resource used to pass information into and back from an [operation](operations.html). It has no other use, and there is no RESTful endpoint associated with it.
Patient | http://hl7.org/fhir/resource-types | Patient | Demographics and other administrative information about an individual or animal receiving care or other health-related services.
PaymentNotice | http://hl7.org/fhir/resource-types | PaymentNotice | This resource provides the status of the payment for goods and services rendered, and the request and response resource references.
PaymentReconciliation | http://hl7.org/fhir/resource-types | PaymentReconciliation | This resource provides the details including amount of a payment and allocates the payment items being paid.
Person | http://hl7.org/fhir/resource-types | Person | Demographics and administrative information about a person independent of a specific health-related context.
PlanDefinition | http://hl7.org/fhir/resource-types | PlanDefinition | This resource allows for the definition of various types of plans as a sharable, consumable, and executable artifact. The resource is general enough to support the description of a broad range of clinical artifacts such as clinical decision support rules, order sets and protocols.
Practitioner | http://hl7.org/fhir/resource-types | Practitioner | A person who is directly or indirectly involved in the provisioning of healthcare.
PractitionerRole | http://hl7.org/fhir/resource-types | PractitionerRole | A specific set of Roles/Locations/specialties/services that a practitioner may perform at an organization for a period of time.
Procedure | http://hl7.org/fhir/resource-types | Procedure | An action that is or was performed on or for a patient. This can be a physical intervention like an operation, or less invasive like long term services, counseling, or hypnotherapy.
Provenance | http://hl7.org/fhir/resource-types | Provenance | Provenance of a resource is a record that describes entities and processes involved in producing and delivering or otherwise influencing that resource. Provenance provides a critical foundation for assessing authenticity, enabling trust, and allowing reproducibility. Provenance assertions are a form of contextual metadata and can themselves become important records with their own provenance. Provenance statement indicates clinical significance in terms of confidence in authenticity, reliability, and trustworthiness, integrity, and stage in lifecycle (e.g. Document Completion - has the artifact been legally authenticated), all of which may impact security, privacy, and trust policies.
Questionnaire | http://hl7.org/fhir/resource-types | Questionnaire | A structured set of questions intended to guide the collection of answers from end-users. Questionnaires provide detailed control over order, presentation, phraseology and grouping to allow coherent, consistent data collection.
QuestionnaireResponse | http://hl7.org/fhir/resource-types | QuestionnaireResponse | A structured set of questions and their answers. The questions are ordered and grouped into coherent subsets, corresponding to the structure of the grouping of the questionnaire being responded to.
RelatedPerson | http://hl7.org/fhir/resource-types | RelatedPerson | Information about a person that is involved in the care for a patient, but who is not the target of healthcare, nor has a formal responsibility in the care process.

RequestGroup | http://hl7.org/fhir/resource-types | RequestGroup | A group of related requests that can be used to capture intended activities that have inter-dependencies such as "give this medication after that one".

ResearchDefinition | http://hl7.org/fhir/resource-types | ResearchDefinition | The ResearchDefinition resource describes the conditional state (population and any exposures being compared within the population) and outcome (if specified) that the knowledge (evidence, assertion, recommendation) is about.

ResearchElementDefinition | http://hl7.org/fhir/resource-types | ResearchElementDefinition | The ResearchElementDefinition resource describes a "PICO" element that knowledge (evidence, assertion, recommendation) is about.

ResearchStudy | http://hl7.org/fhir/resource-types | ResearchStudy | A process where a researcher or organization plans and then executes a series of steps intended to increase the field of healthcare-related knowledge. This includes studies of safety, efficacy, comparative effectiveness and other information about medications, devices, therapies and other interventional and investigative techniques. A ResearchStudy involves the gathering of information about human or animal subjects.

ResearchSubject | http://hl7.org/fhir/resource-types | ResearchSubject | A physical entity which is the primary unit of operational and/or administrative interest in a study.

Resource | http://hl7.org/fhir/resource-types | Resource | This is the base resource type for everything.

RiskAssessment | http://hl7.org/fhir/resource-types | RiskAssessment | An assessment of the likely outcome(s) for a patient or other subject as well as the likelihood of each outcome.

RiskEvidenceSynthesis | http://hl7.org/fhir/resource-types | RiskEvidenceSynthesis | The RiskEvidenceSynthesis resource describes the likelihood of an outcome in a population plus exposure state where the risk estimate is derived from a combination of research studies.

Schedule | http://hl7.org/fhir/resource-types | Schedule | A container for slots of time that may be available for booking appointments.

SearchParameter | http://hl7.org/fhir/resource-types | SearchParameter | A search parameter that defines a named search item that can be used to search/filter on a resource.

ServiceRequest | http://hl7.org/fhir/resource-types | ServiceRequest | A record of a request for service such as diagnostic investigations, treatments, or operations to be performed.

Slot | http://hl7.org/fhir/resource-types | Slot | A slot of time on a schedule that may be available for booking appointments.

Specimen | http://hl7.org/fhir/resource-types | Specimen | A sample to be used for analysis.

SpecimenDefinition | http://hl7.org/fhir/resource-types | SpecimenDefinition | A kind of specimen with associated set of requirements.

StructureDefinition | http://hl7.org/fhir/resource-types | StructureDefinition | A definition of a FHIR structure. This resource is used to describe the underlying resources, data types defined in FHIR, and also for describing extensions and constraints on resources and data types.

StructureMap | http://hl7.org/fhir/resource-types | StructureMap | A Map of relationships between 2 structures that can be used to transform data.

Subscription | http://hl7.org/fhir/resource-types | Subscription | The subscription resource is used to define a push-based subscription from a server to another system. Once a subscription is registered with the server, the server checks every resource that is created or updated, and if the resource matches the given criteria, it sends a message on the defined "channel" so that another system can take an appropriate action.

Substance | http://hl7.org/fhir/resource-types | Substance | A homogeneous material with a definite composition.

SubstanceNucleicAcid | http://hl7.org/fhir/resource-types | SubstanceNucleicAcid | Nucleic acids are defined by three distinct elements: the base, sugar and linkage. Individual substance/moiety IDs will be created for each of these elements. The nucleotide sequence will be always entered in the 5’-3’ direction.

SubstancePolymer | http://hl7.org/fhir/resource-types | SubstancePolymer | Todo.

SubstanceProtein | http://hl7.org/fhir/resource-types | SubstanceProtein | A SubstanceProtein is defined as a single unit of a linear amino acid sequence, or a combination of subunits that are either covalently linked or have a defined invariant stoichiometric relationship. This includes all synthetic, recombinant and purified SubstanceProteins of defined sequence, whether the use is therapeutic or prophylactic. This set of elements will be used to describe albumins, coagulation factors, cytokines, growth factors, peptide/SubstanceProtein hormones, enzymes, toxins, toxoids, recombinant vaccines, and immunomodulators.

SubstanceReferenceInformation | http://hl7.org/fhir/resource-types | SubstanceReferenceInformation | Todo.

SubstanceSourceMaterial | http://hl7.org/fhir/resource-types | SubstanceSourceMaterial | Source material shall capture information on the taxonomic and anatomical origins as well as the fraction of a material that can result in or can be modified to form a substance. This set of data elements shall be used to define polymer substances isolated from biological matrices. Taxonomic and anatomical origins shall be described using a controlled vocabulary as required. This information is captured for naturally derived polymers (. starch) and structurally diverse substances. For Organisms belonging to the Kingdom Plantae the Substance level defines the fresh material of a single species or infraspecies, the Herbal Drug and the Herbal preparation. For Herbal preparations, the fraction information will be captured at the Substance information level and additional information for herbal extracts will be captured at the Specified Substance Group 1 information level. See for further explanation the Substance Class: Structurally Diverse and the herbal annex.

SubstanceSpecification | http://hl7.org/fhir/resource-types | SubstanceSpecification | The detailed description of a substance, typically at a level beyond what is used for prescribing.

SupplyDelivery | http://hl7.org/fhir/resource-types | SupplyDelivery | Record of delivery of what is supplied.

SupplyRequest | http://hl7.org/fhir/resource-types | SupplyRequest | A record of a request for a medication, substance or device used in the healthcare setting.

Task | http://hl7.org/fhir/resource-types | Task | A task to be performed.

TerminologyCapabilities | http://hl7.org/fhir/resource-types | TerminologyCapabilities | A TerminologyCapabilities resource documents a set of capabilities (behaviors) of a FHIR Terminology Server that may be used as a statement of actual server functionality or a statement of required or desired server implementation.

TestReport | http://hl7.org/fhir/resource-types | TestReport | A summary of information based on the results of executing a TestScript.

TestScript | http://hl7.org/fhir/resource-types | TestScript | A structured set of tests against a FHIR server or client implementation to determine compliance against the FHIR specification.

ValueSet | http://hl7.org/fhir/resource-types | ValueSet | A ValueSet resource instance specifies a set of codes drawn from one or more code systems, intended for use in a particular context. Value sets link between [[[CodeSystem]]] definitions and their use in [coded elements](terminologies.html).

VerificationResult | http://hl7.org/fhir/resource-types | VerificationResult | Describes validation requirements, source(s), status and dates for one or more elements.

VisionPrescription | http://hl7.org/fhir/resource-types | VisionPrescription | An authorization for the provision of glasses and/or contact lenses to a patient.
Type | http://hl7.org/fhir/abstract-types | Type | A place holder that means any kind of data type
Any | http://hl7.org/fhir/abstract-types | Any | A place holder that means any kind of resource

Additional Language Displays

Code | English (English, en) | Spanish (es) | French (fr) | Italian (it) | □□ (Chinese, zh)
Address
Age
Annotation
Attachment
BackboneElement
CodeableConcept
Coding
ContactDetail
ContactPoint
Contributor
Count
DataRequirement
Distance
Dosage
Duration
Element
ElementDefinition
Expression
Extension
HumanName
Identifier
MarketingStatus
Meta
Money
MoneyQuantity
Narrative
ParameterDefinition
Period
Population
ProdCharacteristic
ProductShelfLife
Quantity
Range
Ratio
Reference
RelatedArtifact
SampledData
Signature
SimpleQuantity
SubstanceAmount
Timing
TriggerDefinition
UsageContext
base64Binary
boolean
canonical
code
date
dateTime
decimal
id
instant
integer
markdown
oid
positiveInt
string
time
unsignedInt
uri
url
uuid
xhtml
Account | Account | Cuenta | | | □□
ActivityDefinition | ActivityDefinition | DefinicionDeActividad | | DefinizioneAttivita | □□□□
AdverseEvent | AdverseEvent | EventoAdverso | | EventoAvverso | □□□□

AllergyIntolerance | AllergyIntolerance | AllergiaIntolerancia | IntoléranceAllergique | AllergiaIntolleranza | □□□□□□□□
Appointment | Appointment | Cita | RendezVous | Appuntamento | □□
AppointmentResponse | AppointmentResponse | CitaRespuesta | RéponseRendezVous | RispostaAppuntamento | □□□□
AuditEvent | AuditEvent | EventoSeguridad | ÉvènementSécurité | | □□□□
Basic | Basic | Basico | Basique | | □□□□
Binary | Binary | Binario | Binaire | Binario | □□□□□
BiologicallyDerivedProduct | BiologicallyDerivedProduct
BodyStructure | BodyStructure | EstructuraDelCuerpo | MorphologieDeCorps | StrutturaDelCorpo | □□□□
Bundle | Bundle | Paquete | Paquet | | □□
CapabilityStatement | CapabilityStatement | DeclaracionDeCapacidad | | | □□□□
CarePlan | CarePlan | PlanDeCuidado | PlanDeSoins | PianoDiCura | □□□□
CareTeam | CareTeam | EquipoDeCuidado | | | □□□□
CatalogEntry | CatalogEntry | EntradaDeCatalogo | | | □□□□
ChargeItem | ChargeItem | CargoDeItem | | | □□□□
ChargeItemDefinition | ChargeItemDefinition
Claim | Claim | Reclamación / Factura | Réclamation | | □□
ClaimResponse | ClaimResponse | | RéponseARéclamation | | □□□□
ClinicalImpression | ClinicalImpression | HallazgoClinico | ImpressionClinique | ImpressioneClinica | □□□□
CodeSystem | CodeSystem | SistemaDeCodigos | | SistemaDiCodifica | □□□□
Communication | Communication | Comunicación | Communication | Comunicazione | □□
CommunicationRequest | CommunicationRequest | ComunicaciónRequerimiento | DemandeDeCommunication |
RichiestaDiComunicazione | □□□□
CompartmentDefinition | CompartmentDefinition | DefinicionDeCompartimento | | | □□□□□□
Composition | Composition | Composición | Composition | Composizione | □□□□□
ConceptMap | ConceptMap | MapaDeConceptos | CarteDeConcepts | MappaDiConcetti | □□□□
Condition | Condition | Condición | Condition | Condizione | □□
Consent | Consent | Consentimiento | | Consenso | □□□
Contract | Contract | Contato | Contrat | Contratto | □□
Coverage | Coverage | Cobertura | Couverture | Copertura | □□□□
CoverageEligibilityRequest | CoverageEligibilityRequest | | CouvertureDemandeEligibilité | RichiestaEleggibilitaCopertura | □□□□
CoverageEligibilityResponse | CoverageEligibilityResponse | | | RéponseEligibilitéCouverture | RispostaEleggibilitaCopertura | □□□□
DetectedIssue | DetectedIssue | Problema-Detectado /ProblemaDetectado | Problème Détecté | ProblemaRilevato | □□□□□
Device | Device | Dispositivo | Dispositif | Dispositivo | □□
DeviceDefinition | DeviceDefinition | DefiniciónDeDispositivo | DéfinitionDeDispositif | | □□□□
DeviceMetric | DeviceMetric | MetricaDeDispositivo | MétriqueDispositif | | □□□□
DeviceRequest | DeviceRequest | SolicitudDeDispositivo | DemandeUtilisationDispositif | RichiestaDispositivo | □□□□
DeviceUseStatement | DeviceUseStatement | | | □□□□□□
DiagnosticReport | DiagnosticReport | InformeDiagnostico | RapportDiagnostique | RefertoDiagnostico | □□□□
DocumentManifest | DocumentManifest | ManifestoDocumento | Manifeste | | □□□□
DocumentReference | DocumentReference | ReferenciaDocumento | RéférenceDocument | RiferimentoDocumento | □□□□
DomainResource | DomainResource | RecursoDeDominio | | | □□□□
EffectEvidenceSynthesis | EffectEvidenceSynthesis
Encounter | Encounter | Encuentro | Venue | | □□□□
Endpoint | Endpoint | Endpoint | | | □□
EnrollmentRequest | EnrollmentRequest | SolicitudDeEnrolamiento | DemandeInscription | RichiestaIscrizione | □□□□□□
EnrollmentResponse | EnrollmentResponse | RespuestaDeEnrolamiento | RéponseInscription | RispostaIscrizione | □□□□□□
EpisodeOfCare | EpisodeOfCare | EpisodioDeCuidado | ÉpisodeDeSoins | EpisodioDiCura | □□□□□□
EventDefinition | EventDefinition | DefinitionDeEvento | | DefinizioneEvento | □□□□
Evidence | Evidence
EvidenceVariable | EvidenceVariable
ExampleScenario | ExampleScenario | EjemploDeEscenario | | ScenarioDiEsempio | □□□□
ExplanationOfBenefit | ExplanationOfBenefit | | ExplicationDuBénéfice | | □□□□
FamilyMemberHistory | FamilyMemberHistory | HistorialMiembroFamiliar | HistoireMembreFamille | | □□□
Flag | Flag | Bandera | Drapeau | | □□
Goal | Goal | Objetivo | But | | □□
GraphDefinition | GraphDefinition | DefinitionGrafico | | | □□□□
Group | Group | Grupo | Groupe | Gruppo | □□
GuidanceResponse | GuidanceResponse | RespuestaDeOrientacion | | | □□□□□□
HealthcareService | HealthcareService | ServicioDeCuidado | ServiceDeSanté | ServizioSanitario | □□□□□□□□
ImagingStudy | ImagingStudy | EstudioImagen / EstudioImagen | EtudeImagerie | | □□□□
Immunization | Immunization | Inmunización | Immunisation | Immunizzazione | □□□□
ImmunizationEvaluation | ImmunizationEvaluation
ImmunizationRecommendation | ImmunizationRecommendation | RecomendaciónInmunización / | RecommendationImmunisation
| | □□□□□□
ImplementationGuide | ImplementationGuide | GuiaDeImplementacion | | | □□□□
InsurancePlan | InsurancePlan
Invoice | Invoice
Library | Library | Librería | | | □
Linkage | Linkage | Enlace / Conexión / Vinculo / Acoplamiento | | | □□□□
List | List | Lista | Liste | Lista | □□
Location | Location | Locacion | Localisation | | □□
Measure | Measure | Medida | | Misura | □□

MeasureReport | MeasureReport | ReporteMedida | | | □□□□
Media | Media | Medio | Média | Media | □□
Medication | Medication | Medicación /Medicamento | Médication | | □□
MedicationAdministration | MedicationAdministration | AdministraciónMedicación / AdministracionMedicamento | AdministrationMédicaments | | □□□□
MedicationDispense | MedicationDispense | DispensaciónMedicación /DispensacionMedicamento | DispensationMédicaments | | □□□□
MedicationKnowledge | MedicationKnowledge
MedicationRequest | MedicationRequest | PrescripciónMedicaciónTODO /PrescripcionMedicamento | PrescriptionMédicamenteuseTODO | | □□□□
MedicationStatement | MedicationStatement | ResumenMedicación /ResumenMedicamento | ÉtatMédication | | □□□□
MedicinalProduct | MedicinalProduct | | | ProdottoMedicinale
MedicinalProductAuthorization | MedicinalProductAuthorization | | | AutorizzazioneProdottoMedicinale
MedicinalProductContraindication | MedicinalProductContraindication
MedicinalProductIndication | MedicinalProductInteraction
MedicinalProductIngredient | MedicinalProductIngredient | | | IngredienteProdottoMedicinale
MedicinalProductInteraction | MedicinalProductInteraction
MedicinalProductManufactured | MedicinalProductManufactured
MedicinalProductPackaged | MedicinalProductPackaged
MedicinalProductPharmaceutical | MedicinalProductPharmaceutical
MedicinalProductUndesirableEffect | MedicinalProductUndesirableEffect
MessageDefinition | MessageDefinition | DefinicionMensaje | | DefinizioneMessaggio | □□□□
MessageHeader | MessageHeader | CabeceraMensaje | EntêteMessage | | □□□□
MolecularSequence | MolecularSequence | SecuenciaMolecular | | | □□□□
NamingSystem | NamingSystem | SistemaDeNombres | SystèmeDeNommage | | □□□□
NutritionOrder | NutritionOrder | OrdenNutrición | OrdreNutrition | | □□□□
Observation | Observation | Observación | Observation | Osservazione | □□
ObservationDefinition | ObservationDefinition | DefinicionDeEspecimen | | DefinizioneOsservazione | □□□□
OperationDefinition | OperationDefinition | DefinicionDeOperacion | DéfinitionOpération | DefinizioneOperazione | □□□□
OperationOutcome | OperationOutcome | ResultadoOperación | RésultatOpération | RisultatoOperazione | □□□□
Organization | Organization | Organización | Organisation | Organizzazione | □□□□
OrganizationAffiliation | OrganizationAffiliation
Parameters | Parameters | Parametros | | | □□
Patient | Patient | Paciente | Patient | Paziente | □□
PaymentNotice | PaymentNotice | AvisoDePago | AvisPaiement | AvvisoDiPagamento | □□□□
PaymentReconciliation | PaymentReconciliation | ConciliacionDePago | RéconciliationPaiement | RiconciliazionePagamento | □□□□
Person | Person | Persona | Personne | Persona | □□
PlanDefinition | PlanDefinition | DefinicionDePlan | | | □□□□
Practitioner | Practitioner | Practicante / Profesional | Praticien | | □□□□
PractitionerRole | PractitionerRole | RolProfesional | | | □□□□□□
Procedure | Procedure | Procedimiento | Procédure | Procedura | □□□□
Provenance | Provenance | Procedencia | Provenance | Provenienza | □□
Questionnaire | Questionnaire | Cuestionario | Questionnaire | Questionario | □□□□
QuestionnaireResponse | QuestionnaireResponse | RespuestaAlCuestionario | RéponseQuestionnaire | RispostaQuestionario | □□□□□□
RelatedPerson | RelatedPerson | PersonaRelacionada | PersonneEnRelation | PersonaCorrelata | □□□□
RequestGroup | RequestGroup | GrupoDeSolicitudes | | GruppoDiRichieste | □□□□
ResearchDefinition | ResearchDefinition
ResearchElementDefinition | ResearchElementDefinition
ResearchStudy | ResearchStudy | EstudioDeInvestigacion | | | □□□□
ResearchSubject | ResearchSubject | SujetoDeInvestigacion | | | □□□□
Resource | Resource | Recurso | | | □□
RiskAssessment | RiskAssessment | EvaluacionDeRiesgo | ÉvaluationRisques | | □□□□
RiskEvidenceSynthesis | RiskEvidenceSynthesis
Schedule | Schedule | Agenda | | | □□□□
SearchParameter | SearchParameter | ParametroDeBusqueda | ParamètreRecherche | ParametroDiRicerca | □□□□
ServiceRequest | ServiceRequest | PeticiónServicio | DemandeService | RichiestaDiServizio | □□□□□□
Slot | Slot | Hueco / Zocalo / Espacio | Slot | | □□
Specimen | Specimen | Especimen | Spécimen | Campione | □□
SpecimenDefinition | SpecimenDefinition | DefinicionDeEspecimen | | DefinizioneCampione | □□□□
StructureDefinition | StructureDefinition | DefinicionDeEstructura | DéfinitionStructure | DefinizioneStruttura | □□□□
StructureMap | StructureMap | MapaDeEstructura | | | □□□□
Subscription | Subscription | Suscripción | Souscription | Sottoscrizione | □□
Substance | Substance | Sustancia | Substance | Sostanza | □□
SubstanceNucleicAcid | SubstanceNucleicAcid
SubstancePolymer | SubstancePolymer
SubstanceProtein | SubstanceProtein
SubstanceReferenceInformation | SubstanceReferenceInformation
SubstanceSourceMaterial | SubstanceSourceMaterial
SubstanceSpecification | SubstanceSpecification
SupplyDelivery | SupplyDelivery | Entrega de Suministro | Supply Livraison | | □□□□
SupplyRequest | SupplyRequest | Solicitud de Suministro | Demande d'approvisionnement | | □□□□
Task | Task | Tarea | | | □□

TerminologyCapabilities | TerminologyCapabilities | CapacidadTerminologica | | □□□□□□
TestReport | TestReport | ReporteDePrueba | RapportTest | | □□□□
TestScript | TestScript | ScriptDePrueba | ScriptTest | | □□□□
ValueSet | ValueSet | ConjuntoValores / ConjuntoDeValores | EnsembleValeurs | | □□□□
VerificationResult | VerificationResult
VisionPrescription | VisionPrescription | PrescripcionDeVision | PrescriptionVision | | □□□□
Type
Any

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.564 Value Set http://hl7.org/fhir/ValueSet/practitioner-role

URL: https://hl7.org/fhir/R4/valueset-practitioner-role.html
Patient Administration Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/practitioner-role
Version: | 4.0.1
Name: | PractitionerRole
Title: | Practitioner role
Definition: | This example value set defines a set of codes that can be used to indicate the role of a Practitioner.
Committee: | Patient Administration Work Group
OID: | 2.16.840.1.113883.4.642.3.439 (for OID based terminology systems)
Copyright: | This resource includes content from SNOMED Clinical Terms® (SNOMED CT®) which is copyright of the International Health Terminology Standards Development Organisation (IHTSDO). Implementers of these specifications must have the appropriate SNOMED CT Affiliate license - for more information contact http://www.snomed.org/snomed-ct/get-snomed-ct or info@snomed.org
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: PractitionerRole.code (CodeableConcept / Example)

4.4.1.564.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/practitioner-role
• Include codes from http://snomed.info/sct where concept is-a 223366009 (Healthcare professional)

4.4.1.564.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 500 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/practitioner-role version 4.0.1

Code | System | Display | Definition
doctor | http://terminology.hl7.org/CodeSystem/practitioner-role | Doctor | A qualified/registered medical practitioner
nurse | http://terminology.hl7.org/CodeSystem/practitioner-role | Nurse | A practitioner with nursing experience that may be qualified/registered
pharmacist | http://terminology.hl7.org/CodeSystem/practitioner-role | Pharmacist | A qualified/registered/licensed pharmacist
researcher | http://terminology.hl7.org/CodeSystem/practitioner-role | Researcher | A practitioner that may perform research
teacher | http://terminology.hl7.org/CodeSystem/practitioner-role | Teacher/educator | Someone who is able to provide educational services
ict | http://terminology.hl7.org/CodeSystem/practitioner-role | ICT professional | Someone who is qualified in Information and Communication Technologies
1421009 | http://snomed.info/sct | Specialized surgeon
3430008 | http://snomed.info/sct | Radiation therapist
3842006 | http://snomed.info/sct | Chiropractor
4162009 | http://snomed.info/sct | Dental assistant
5275007 | http://snomed.info/sct | NA - Nursing auxiliary

6816002 | http://snomed.info/sct | Specialized nurse
6868009 | http://snomed.info/sct | Hospital administrator
8724009 | http://snomed.info/sct | Plastic surgeon
11661002 | http://snomed.info/sct | Neuropathologist
11911009 | http://snomed.info/sct | Nephrologist
11935004 | http://snomed.info/sct | Obstetrician
13580004 | http://snomed.info/sct | School dental assistant
14698002 | http://snomed.info/sct | Medical microbiologist
17561000 | http://snomed.info/sct | Cardiologist
18803008 | http://snomed.info/sct | Dermatologist
18850004 | http://snomed.info/sct | Laboratory hematologist
19244007 | http://snomed.info/sct | Gerodontist
20145008 | http://snomed.info/sct | Removable prosthodontist
21365001 | http://snomed.info/sct | Specialized dentist
21450003 | http://snomed.info/sct | Neuropsychiatrist
22515006 | http://snomed.info/sct | Medical assistant
22731001 | http://snomed.info/sct | Orthopedic surgeon
22983004 | http://snomed.info/sct | Thoracic surgeon
23278007 | http://snomed.info/sct | Community health physician
24430003 | http://snomed.info/sct | Physical medicine specialist
24590004 | http://snomed.info/sct | Urologist
25961008 | http://snomed.info/sct | Electroencephalography specialist
26042002 | http://snomed.info/sct | Dental hygienist
26369006 | http://snomed.info/sct | Public health nurse
28229004 | http://snomed.info/sct | Optometrist
28411006 | http://snomed.info/sct | Neonatologist
28544002 | http://snomed.info/sct | Chemical pathologist
36682004 | http://snomed.info/sct | PT - Physiotherapist
37154003 | http://snomed.info/sct | Periodontist
37504001 | http://snomed.info/sct | Orthodontist
39677007 | http://snomed.info/sct | Internal medicine specialist
40127002 | http://snomed.info/sct | Dietitian (general)
40204001 | http://snomed.info/sct | Hematologist
40570005 | http://snomed.info/sct | Interpreter
41672002 | http://snomed.info/sct | Respiratory physician
41904004 | http://snomed.info/sct | Medical X-ray technician
43702002 | http://snomed.info/sct | Occupational health nurse
44652006 | http://snomed.info/sct | Pharmaceutical assistant
45419001 | http://snomed.info/sct | Masseur
45440000 | http://snomed.info/sct | Rheumatologist
45544007 | http://snomed.info/sct | Neurosurgeon
45956004 | http://snomed.info/sct | Sanitarian
46255001 | http://snomed.info/sct | Pharmacist
48740002 | http://snomed.info/sct | Philologist
49203003 | http://snomed.info/sct | Dispensing optometrist
49993003 | http://snomed.info/sct | Maxillofacial surgeon
50149000 | http://snomed.info/sct | Endodontist
54503009 | http://snomed.info/sct | Faith healer
56397003 | http://snomed.info/sct | Neurologist
56466003 | http://snomed.info/sct | Community physician
56542007 | http://snomed.info/sct | Medical record administrator
56545009 | http://snomed.info/sct | Cardiovascular surgeon
57654006 | http://snomed.info/sct | Fixed prosthodontist
59058001 | http://snomed.info/sct | General physician
59169001 | http://snomed.info/sct | Orthopedic technician
59944000 | http://snomed.info/sct | Psychologist
60008001 | http://snomed.info/sct | Community-based dietitian
61207006 | http://snomed.info/sct | Medical pathologist
61246008 | http://snomed.info/sct | Laboratory medicine specialist
61345009 | http://snomed.info/sct | Otorhinolaryngologist
61894003 | http://snomed.info/sct | Endocrinologist
62247001 | http://snomed.info/sct | Family medicine specialist
63098009 | http://snomed.info/sct | Clinical immunologist
66476003 | http://snomed.info/sct | Oral pathologist
66862007 | http://snomed.info/sct | Radiologist
68867008 | http://snomed.info/sct | Public health dentist
68950000 | http://snomed.info/sct | Prosthodontist
69280009 | http://snomed.info/sct | Specialized physician
71838004 | http://snomed.info/sct | Gastroenterologist
73265009 | http://snomed.info/sct | Nursing aid
75271001 | http://snomed.info/sct | MW - Midwife
76166008 | http://snomed.info/sct | Practical aid (pharmacy)

76231001 | http://snomed.info/sct | Osteopath
76899008 | http://snomed.info/sct | Infectious diseases physician
78703002 | http://snomed.info/sct | General surgeon
78729002 | http://snomed.info/sct | Diagnostic radiologist
79898004 | http://snomed.info/sct | Auxiliary midwife
80409005 | http://snomed.info/sct | Translator
80546007 | http://snomed.info/sct | OT - Occupational therapist
80584001 | http://snomed.info/sct | Psychiatrist
80933006 | http://snomed.info/sct | Nuclear medicine physician
81464008 | http://snomed.info/sct | Clinical pathologist
82296001 | http://snomed.info/sct | Pediatrician
83189004 | http://snomed.info/sct | Other professional nurse
83273008 | http://snomed.info/sct | Anatomic pathologist
83685006 | http://snomed.info/sct | Gynecologist
85733003 | http://snomed.info/sct | General pathologist
88189002 | http://snomed.info/sct | Anesthesiologist
88475002 | http://snomed.info/sct | Other dietitians and public health nutritionists
90201008 | http://snomed.info/sct | Pediatric dentist
90655003 | http://snomed.info/sct | Care of the elderly physician
106289002 | http://snomed.info/sct | Dental surgeon
106291005 | http://snomed.info/sct | Dietician AND/OR public health nutritionist
106292003 | http://snomed.info/sct | Nurse
106293008 | http://snomed.info/sct | Nursing personnel
106294002 | http://snomed.info/sct | Midwifery personnel
106296000 | http://snomed.info/sct | Physiotherapist AND/OR occupational therapist
106330007 | http://snomed.info/sct | Philologist, translator AND/OR interpreter
112247003 | http://snomed.info/sct | Medical doctor
158965000 | http://snomed.info/sct | Medical practitioner
158966004 | http://snomed.info/sct | Medical administrator - national
158967008 | http://snomed.info/sct | Consultant physician
158968003 | http://snomed.info/sct | Consultant surgeon
158969006 | http://snomed.info/sct | Consultant gynecology and obstetrics
158970007 | http://snomed.info/sct | Anesthetist
158971006 | http://snomed.info/sct | Hospital registrar
158972004 | http://snomed.info/sct | House officer
158973009 | http://snomed.info/sct | Occupational physician
158974003 | http://snomed.info/sct | Clinical medical officer
158975002 | http://snomed.info/sct | Medical practitioner - teaching
158977005 | http://snomed.info/sct | Dental administrator
158978000 | http://snomed.info/sct | Dental consultant
158979008 | http://snomed.info/sct | Dental general practitioner
158980006 | http://snomed.info/sct | Dental practitioner - teaching
158983008 | http://snomed.info/sct | Nurse administrator - national
158984002 | http://snomed.info/sct | Nursing officer - region
158985001 | http://snomed.info/sct | Nursing officer - district
158986000 | http://snomed.info/sct | Nursing administrator - professional body
158987009 | http://snomed.info/sct | Nursing officer - division
158988004 | http://snomed.info/sct | Nurse education director
158989007 | http://snomed.info/sct | Occupational health nursing officer
158990003 | http://snomed.info/sct | Nursing officer
158992006 | http://snomed.info/sct | Midwifery sister
158993001 | http://snomed.info/sct | Nursing sister (theatre)
158994007 | http://snomed.info/sct | Staff nurse
158995008 | http://snomed.info/sct | Staff midwife
158996009 | http://snomed.info/sct | State enrolled nurse
158997000 | http://snomed.info/sct | District nurse
158998005 | http://snomed.info/sct | Private nurse
158999002 | http://snomed.info/sct | Community midwife
159001001 | http://snomed.info/sct | Clinic nurse
159002008 | http://snomed.info/sct | Practice nurse
159003003 | http://snomed.info/sct | School nurse
159004009 | http://snomed.info/sct | Nurse - teaching
159005005 | http://snomed.info/sct | Student nurse
159006006 | http://snomed.info/sct | Dental nurse
159007002 | http://snomed.info/sct | Community pediatric nurse
159010009 | http://snomed.info/sct | Hospital pharmacist
159011008 | http://snomed.info/sct | Retail pharmacist
159012001 | http://snomed.info/sct | Industrial pharmacist
159013006 | http://snomed.info/sct | Pharmaceutical officer H.A.
159014000 | http://snomed.info/sct | Trainee pharmacist
159016003 | http://snomed.info/sct | Medical radiographer
159017007 | http://snomed.info/sct | Diagnostic radiographer

159018002 | http://snomed.info/sct | Therapeutic radiographer
159019005 | http://snomed.info/sct | Trainee radiographer
159021000 | http://snomed.info/sct | Ophthalmic optician
159022007 | http://snomed.info/sct | Trainee optician
159025009 | http://snomed.info/sct | Remedial gymnast
159026005 | http://snomed.info/sct | Speech and language therapist
159027001 | http://snomed.info/sct | Orthoptist
159028006 | http://snomed.info/sct | Trainee remedial therapist
159033005 | http://snomed.info/sct | Dietician
159034004 | http://snomed.info/sct | Podiatrist
159035003 | http://snomed.info/sct | Dental auxiliary
159036002 | http://snomed.info/sct | ECG technician
159037006 | http://snomed.info/sct | EEG technician
159038001 | http://snomed.info/sct | Artificial limb fitter
159039009 | http://snomed.info/sct | AT - Audiology technician
159040006 | http://snomed.info/sct | Pharmacy technician
159041005 | http://snomed.info/sct | Trainee medical technician
159141008 | http://snomed.info/sct | Geneticist
159972006 | http://snomed.info/sct | Surgical corset fitter
160008000 | http://snomed.info/sct | Dental technician
224529009 | http://snomed.info/sct | Clinical assistant
224530004 | http://snomed.info/sct | Senior registrar
224531000 | http://snomed.info/sct | Registrar
224532007 | http://snomed.info/sct | Senior house officer
224533002 | http://snomed.info/sct | MO - Medical officer
224534008 | http://snomed.info/sct | Health visitor, nurse/midwife
224535009 | http://snomed.info/sct | Registered nurse
224536005 | http://snomed.info/sct | Midwifery tutor
224537001 | http://snomed.info/sct | Accident and Emergency nurse
224538006 | http://snomed.info/sct | Triage nurse
224540001 | http://snomed.info/sct | Community nurse
224541002 | http://snomed.info/sct | Nursing continence advisor
224542009 | http://snomed.info/sct | Coronary care nurse
224543004 | http://snomed.info/sct | Diabetic nurse
224544005 | http://snomed.info/sct | Family planning nurse
224545006 | http://snomed.info/sct | Care of the elderly nurse
224546007 | http://snomed.info/sct | ICN - Infection control nurse
224547003 | http://snomed.info/sct | Intensive therapy nurse
224548008 | http://snomed.info/sct | Learning disabilities nurse
224549000 | http://snomed.info/sct | Neonatal nurse
224550000 | http://snomed.info/sct | Neurology nurse
224551001 | http://snomed.info/sct | Industrial nurse
224552008 | http://snomed.info/sct | Oncology nurse
224553003 | http://snomed.info/sct | Macmillan nurse
224554009 | http://snomed.info/sct | Marie Curie nurse
224555005 | http://snomed.info/sct | Pain control nurse
224556006 | http://snomed.info/sct | Palliative care nurse
224557002 | http://snomed.info/sct | Chemotherapy nurse
224558007 | http://snomed.info/sct | Radiotherapy nurse
224559004 | http://snomed.info/sct | PACU nurse
224560009 | http://snomed.info/sct | Stomatherapist
224561008 | http://snomed.info/sct | Theatre nurse
224562001 | http://snomed.info/sct | Pediatric nurse
224563006 | http://snomed.info/sct | Psychiatric nurse
224564000 | http://snomed.info/sct | Community mental health nurse
224565004 | http://snomed.info/sct | Renal nurse
224566003 | http://snomed.info/sct | Hemodialysis nurse
224567007 | http://snomed.info/sct | Wound care nurse
224569005 | http://snomed.info/sct | Nurse grade
224570006 | http://snomed.info/sct | Clinical nurse specialist
224571005 | http://snomed.info/sct | Nurse practitioner
224572003 | http://snomed.info/sct | Nursing sister
224573008 | http://snomed.info/sct | CN - Charge nurse
224574002 | http://snomed.info/sct | Ward manager
224575001 | http://snomed.info/sct | Nursing team leader
224576000 | http://snomed.info/sct | Nursing assistant
224577009 | http://snomed.info/sct | Healthcare assistant
224578004 | http://snomed.info/sct | Nursery nurse
224579007 | http://snomed.info/sct | Healthcare service manager
224580005 | http://snomed.info/sct | Occupational health service manager
224581009 | http://snomed.info/sct | Community nurse manager
224583007 | http://snomed.info/sct | Behavior therapist

224584001 | http://snomed.info/sct | Behavior therapy assistant
224585000 | http://snomed.info/sct | Drama therapist
224586004 | http://snomed.info/sct | Domiciliary occupational therapist
224587008 | http://snomed.info/sct | Occupational therapy helper
224588003 | http://snomed.info/sct | Psychotherapist
224589006 | http://snomed.info/sct | Community-based physiotherapist
224590002 | http://snomed.info/sct | Play therapist
224591003 | http://snomed.info/sct | Play specialist
224592005 | http://snomed.info/sct | Play leader
224593000 | http://snomed.info/sct | Community-based speech/language therapist
224594006 | http://snomed.info/sct | Speech/language assistant
224595007 | http://snomed.info/sct | Professional counselor
224596008 | http://snomed.info/sct | Marriage guidance counselor
224597004 | http://snomed.info/sct | Trained nurse counselor
224598009 | http://snomed.info/sct | Trained social worker counselor
224599001 | http://snomed.info/sct | Trained personnel counselor
224600003 | http://snomed.info/sct | Psychoanalyst
224601004 | http://snomed.info/sct | Assistant psychologist
224602006 | http://snomed.info/sct | Community-based podiatrist
224603001 | http://snomed.info/sct | Foot care worker
224604007 | http://snomed.info/sct | Audiometrician
224605008 | http://snomed.info/sct | Audiometrist
224606009 | http://snomed.info/sct | Technical healthcare occupation
224607000 | http://snomed.info/sct | Occupational therapy technical instructor
224608005 | http://snomed.info/sct | Administrative healthcare staff
224609002 | http://snomed.info/sct | Complementary health worker
224610007 | http://snomed.info/sct | Supporting services personnel
224614003 | http://snomed.info/sct | Research associate
224615002 | http://snomed.info/sct | Research nurse
224620002 | http://snomed.info/sct | Human aid to communication
224621003 | http://snomed.info/sct | Palantypist
224622005 | http://snomed.info/sct | Note taker
224623000 | http://snomed.info/sct | Cuer
224624006 | http://snomed.info/sct | Lipspeaker
224625007 | http://snomed.info/sct | Interpreter for British sign language
224626008 | http://snomed.info/sct | Interpreter for Signs supporting English
224936003 | http://snomed.info/sct | General practitioner locum
225726006 | http://snomed.info/sct | Lactation consultant
225727002 | http://snomed.info/sct | Midwife counselor
265937000 | http://snomed.info/sct | Nursing occupation
265939002 | http://snomed.info/sct | Medical/dental technicians
283875005 | http://snomed.info/sct | Parkinson disease nurse
302211009 | http://snomed.info/sct | Specialist registrar
303124005 | http://snomed.info/sct | Member of mental health review tribunal
303129000 | http://snomed.info/sct | Hospital manager
303133007 | http://snomed.info/sct | Responsible medical officer
303134001 | http://snomed.info/sct | Independent doctor
304291006 | http://snomed.info/sct | Bereavement counselor
304292004 | http://snomed.info/sct | Surgeon
307988006 | http://snomed.info/sct | Medical technician
308002005 | http://snomed.info/sct | Remedial therapist
309294001 | http://snomed.info/sct | Accident and Emergency doctor
309295000 | http://snomed.info/sct | Clinical oncologist
309296004 | http://snomed.info/sct | Family planning doctor
309322005 | http://snomed.info/sct | Associate general practitioner
309323000 | http://snomed.info/sct | Partner of general practitioner
309324006 | http://snomed.info/sct | Assistant GP
309326008 | http://snomed.info/sct | Deputizing general practitioner
309327004 | http://snomed.info/sct | General practitioner registrar
309328009 | http://snomed.info/sct | Ambulatory pediatrician
309329001 | http://snomed.info/sct | Community pediatrician
309330006 | http://snomed.info/sct | Pediatric cardiologist
309331005 | http://snomed.info/sct | Pediatric endocrinologist
309332003 | http://snomed.info/sct | Pediatric gastroenterologist
309333008 | http://snomed.info/sct | Pediatric nephrologist
309334002 | http://snomed.info/sct | Pediatric neurologist
309335001 | http://snomed.info/sct | Pediatric rheumatologist
309336000 | http://snomed.info/sct | Pediatric oncologist
309337009 | http://snomed.info/sct | Pain management specialist
309338004 | http://snomed.info/sct | Intensive care specialist
309339007 | http://snomed.info/sct | Adult intensive care specialist
309340009 | http://snomed.info/sct | Pediatric intensive care specialist

309341008 | http://snomed.info/sct | Blood transfusion doctor
309342001 | http://snomed.info/sct | Histopathologist
309343006 | http://snomed.info/sct | Physician
309345004 | http://snomed.info/sct | Chest physician
309346003 | http://snomed.info/sct | Thoracic physician
309347007 | http://snomed.info/sct | Clinical hematologist
309348002 | http://snomed.info/sct | Clinical neurophysiologist
309349005 | http://snomed.info/sct | Clinical physiologist
309350005 | http://snomed.info/sct | Diabetologist
309351009 | http://snomed.info/sct | Andrologist
309352002 | http://snomed.info/sct | Neuroendocrinologist
309353007 | http://snomed.info/sct | Reproductive endocrinologist
309354001 | http://snomed.info/sct | Thyroidologist
309355000 | http://snomed.info/sct | Clinical geneticist
309356004 | http://snomed.info/sct | Clinical cytogeneticist
309357008 | http://snomed.info/sct | Clinical molecular geneticist
309358003 | http://snomed.info/sct | Genitourinary medicine physician
309359006 | http://snomed.info/sct | Palliative care physician
309360001 | http://snomed.info/sct | Rehabilitation physician
309361002 | http://snomed.info/sct | Child and adolescent psychiatrist
309362009 | http://snomed.info/sct | Forensic psychiatrist
309363004 | http://snomed.info/sct | Liaison psychiatrist
309364005 | http://snomed.info/sct | Psychogeriatrician
309365006 | http://snomed.info/sct | Psychiatrist for mental handicap
309366007 | http://snomed.info/sct | Rehabilitation psychiatrist
309367003 | http://snomed.info/sct | Obstetrician and gynecologist
309368008 | http://snomed.info/sct | Breast surgeon
309369000 | http://snomed.info/sct | Cardiothoracic surgeon
309371000 | http://snomed.info/sct | Cardiac surgeon
309372007 | http://snomed.info/sct | Ear, nose and throat surgeon
309373002 | http://snomed.info/sct | Endocrine surgeon
309374008 | http://snomed.info/sct | Thyroid surgeon
309375009 | http://snomed.info/sct | Pituitary surgeon
309376005 | http://snomed.info/sct | Gastrointestinal surgeon
309377001 | http://snomed.info/sct | General gastrointestinal surgeon
309378006 | http://snomed.info/sct | Upper gastrointestinal surgeon
309379003 | http://snomed.info/sct | Colorectal surgeon
309380000 | http://snomed.info/sct | Hand surgeon
309381001 | http://snomed.info/sct | Hepatobiliary surgeon
309382008 | http://snomed.info/sct | Ophthalmic surgeon
309383003 | http://snomed.info/sct | Pediatric surgeon
309384009 | http://snomed.info/sct | Pancreatic surgeon
309385005 | http://snomed.info/sct | Transplant surgeon
309386006 | http://snomed.info/sct | Trauma surgeon
309388007 | http://snomed.info/sct | Vascular surgeon
309389004 | http://snomed.info/sct | Medical practitioner grade
309390008 | http://snomed.info/sct | Hospital consultant
309391007 | http://snomed.info/sct | Visiting specialist registrar
309392000 | http://snomed.info/sct | Research registrar
309393005 | http://snomed.info/sct | General practitioner grade
309394004 | http://snomed.info/sct | General practitioner principal
309395003 | http://snomed.info/sct | Hospital specialist
309396002 | http://snomed.info/sct | Associate specialist
309397006 | http://snomed.info/sct | Research fellow
309398001 | http://snomed.info/sct | Allied health professional
309399009 | http://snomed.info/sct | Hospital dietitian
309400002 | http://snomed.info/sct | Domiciliary physiotherapist
309401003 | http://snomed.info/sct | General practitioner-based physiotherapist
309402005 | http://snomed.info/sct | Hospital-based physiotherapist
309403000 | http://snomed.info/sct | Private physiotherapist
309404006 | http://snomed.info/sct | Physiotherapy assistant
309409001 | http://snomed.info/sct | Hospital-based speech and language therapist
309410006 | http://snomed.info/sct | Arts therapist
309411005 | http://snomed.info/sct | Dance therapist
309412003 | http://snomed.info/sct | Music therapist
309413008 | http://snomed.info/sct | Renal dietitian
309414002 | http://snomed.info/sct | Liver dietitian
309415001 | http://snomed.info/sct | Oncology dietitian
309416000 | http://snomed.info/sct | Pediatric dietitian
309417009 | http://snomed.info/sct | Diabetes dietitian
309418004 | http://snomed.info/sct | Audiologist
309419007 | http://snomed.info/sct | Hearing therapist

309420001 | http://snomed.info/sct | Audiological scientist
309421002 | http://snomed.info/sct | Hearing aid dispenser
309422009 | http://snomed.info/sct | Community-based occupational therapist
309423004 | http://snomed.info/sct | Hospital occupational therapist
309427003 | http://snomed.info/sct | Social services occupational therapist
309428008 | http://snomed.info/sct | Orthotist
309429000 | http://snomed.info/sct | Surgical fitter
309434001 | http://snomed.info/sct | Hospital-based podiatrist
309435000 | http://snomed.info/sct | Podiatry assistant
309436004 | http://snomed.info/sct | Lymphedema nurse
309437008 | http://snomed.info/sct | Community learning disabilities nurse
309439006 | http://snomed.info/sct | Clinical nurse teacher
309440008 | http://snomed.info/sct | Community practice nurse teacher
309441007 | http://snomed.info/sct | Nurse tutor
309442000 | http://snomed.info/sct | Nurse teacher practitioner
309443005 | http://snomed.info/sct | Nurse lecturer practitioner
309444004 | http://snomed.info/sct | Outreach nurse
309445003 | http://snomed.info/sct | Anesthetic nurse
309446002 | http://snomed.info/sct | Nurse manager
309450009 | http://snomed.info/sct | Nurse administrator
309452001 | http://snomed.info/sct | Midwifery grade
309453006 | http://snomed.info/sct | Midwife
309454000 | http://snomed.info/sct | Student midwife
309455004 | http://snomed.info/sct | Parentcraft sister
309459005 | http://snomed.info/sct | Healthcare professional grade
309460000 | http://snomed.info/sct | Restorative dentist
310170009 | http://snomed.info/sct | Pediatric audiologist
310171008 | http://snomed.info/sct | Immunopathologist
310172001 | http://snomed.info/sct | Audiological physician
310173006 | http://snomed.info/sct | Clinical pharmacologist
310174000 | http://snomed.info/sct | Private doctor
310175004 | http://snomed.info/sct | Agency nurse
310176003 | http://snomed.info/sct | Behavioral therapist nurse
310177007 | http://snomed.info/sct | Cardiac rehabilitation nurse
310178002 | http://snomed.info/sct | Genitourinary nurse
310179005 | http://snomed.info/sct | Rheumatology nurse specialist
310180008 | http://snomed.info/sct | Continence nurse
310181007 | http://snomed.info/sct | Contact tracing nurse
310182000 | http://snomed.info/sct | General nurse
310183005 | http://snomed.info/sct | Nurse for the mentally handicapped
310184004 | http://snomed.info/sct | Liaison nurse
310185003 | http://snomed.info/sct | Diabetic liaison nurse
310186002 | http://snomed.info/sct | Nurse psychotherapist
310187006 | http://snomed.info/sct | Company nurse
310188001 | http://snomed.info/sct | Hospital midwife
310189009 | http://snomed.info/sct | Genetic counselor
310190000 | http://snomed.info/sct | Mental health counselor
310191001 | http://snomed.info/sct | Clinical psychologist
310192008 | http://snomed.info/sct | Educational psychologist
310193003 | http://snomed.info/sct | Coroner
310194009 | http://snomed.info/sct | Appliance officer
310512001 | http://snomed.info/sct | Medical oncologist
311441001 | http://snomed.info/sct | School medical officer
312485001 | http://snomed.info/sct | Integrated midwife
372102007 | http://snomed.info/sct | RN First Assist
387619007 | http://snomed.info/sct | Optician
394572006 | http://snomed.info/sct | Medical secretary
394618009 | http://snomed.info/sct | Hospital nurse
397824005 | http://snomed.info/sct | Consultant anesthetist
397897005 | http://snomed.info/sct | Paramedic
397903001 | http://snomed.info/sct | Staff grade obstetrician
397908005 | http://snomed.info/sct | Staff grade practitioner
398130009 | http://snomed.info/sct | Medical student
398238009 | http://snomed.info/sct | Acting obstetric registrar
404940000 | http://snomed.info/sct | Physiotherapist technical instructor
405277009 | http://snomed.info/sct | Resident physician
405278004 | http://snomed.info/sct | Certified registered nurse anesthetist
405279007 | http://snomed.info/sct | Attending physician
405623001 | http://snomed.info/sct | Assigned practitioner
405684005 | http://snomed.info/sct | Professional initiating surgical case
405685006 | http://snomed.info/sct | Professional providing staff relief during surgical procedure
408798009 | http://snomed.info/sct | Consultant pediatrician

408799001 | http://snomed.info/sct | Consultant neonatologist
409974004 | http://snomed.info/sct | Health educator
409975003 | http://snomed.info/sct | Certified health education specialist
413854007 | http://snomed.info/sct | Circulating nurse
415075003 | http://snomed.info/sct | Perioperative nurse
415506007 | http://snomed.info/sct | Scrub nurse
416160000 | http://snomed.info/sct | Fellow of American Academy of Osteopathy
420409002 | http://snomed.info/sct | Oculoplastic surgeon
420678001 | http://snomed.info/sct | Retinal surgeon
421841007 | http://snomed.info/sct | Admitting physician
422140007 | http://snomed.info/sct | Medical ophthalmologist
422234006 | http://snomed.info/sct | Ophthalmologist
432100008 | http://snomed.info/sct | Health coach
442867008 | http://snomed.info/sct | Respiratory therapist
443090005 | http://snomed.info/sct | Podiatric surgeon
444912007 | http://snomed.info/sct | Hypnotherapist
445313000 | http://snomed.info/sct | Asthma nurse specialist
445451001 | http://snomed.info/sct | Nurse case manager
446050000 | http://snomed.info/sct | PCP - Primary care physician
446701002 | http://snomed.info/sct | Addiction medicine specialist
449161006 | http://snomed.info/sct | PA - physician assistant
471302004 | http://snomed.info/sct | Government midwife
3981000175106 | http://snomed.info/sct | Nurse complex case manager
231189271000087109 | http://snomed.info/sct | Naturopath
236749831000087105 | http://snomed.info/sct | Prosthetist
258508741000087105 | http://snomed.info/sct | Hip and knee surgeon
260767431000087107 | http://snomed.info/sct | Hepatologist
285631911000087106 | http://snomed.info/sct | Shoulder surgeon
291705421000087106 | http://snomed.info/sct | Interventional radiologist
341320851000087105 | http://snomed.info/sct | Pediatric radiologist
368890881000087105 | http://snomed.info/sct | Emergency medicine specialist
398480381000087106 | http://snomed.info/sct | Family medicine specialist - palliative care
416186861000087101 | http://snomed.info/sct | Surgical oncologist
450044741000087104 | http://snomed.info/sct | Acupuncturist
465511991000087105 | http://snomed.info/sct | Pediatric orthopedic surgeon
494782281000087101 | http://snomed.info/sct | Pediatric hematologist
619197631000087102 | http://snomed.info/sct | Neuroradiologist
623630151000087105 | http://snomed.info/sct | Family medicine specialist - anesthetist
666997781000087107 | http://snomed.info/sct | Doula
673825031000087109 | http://snomed.info/sct | Traditional herbal medicine specialist
682131381000087105 | http://snomed.info/sct | Occupational medicine specialist
724111801000087104 | http://snomed.info/sct | Pediatric emergency medicine specialist
747936471000087102 | http://snomed.info/sct | Family medicine specialist - care of the elderly
766788081000087100 | http://snomed.info/sct | Travel medicine specialist
767205061000087108 | http://snomed.info/sct | Spine surgeon
813758161000087106 | http://snomed.info/sct | Maternal or fetal medicine specialist
822410621000087104 | http://snomed.info/sct | Massage therapist
847240411000087102 | http://snomed.info/sct | Hospitalist
853827051000087104 | http://snomed.info/sct | Sports medicine specialist
926871431000087103 | http://snomed.info/sct | Pediatric respirologist
954544641000087107 | http://snomed.info/sct | Homeopath
956387501000087102 | http://snomed.info/sct | Family medicine specialist - emergency medicine
969118571000087109 | http://snomed.info/sct | Pediatric hematologist or oncologist
984095901000087105 | http://snomed.info/sct | Foot and ankle surgeon
990928611000087105 | http://snomed.info/sct | Invasive cardiologist
999480451000087102 | http://snomed.info/sct | Case manager
999480461000087104 | http://snomed.info/sct | Kinesthesiologist

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information

Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)

Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')

Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application

Definition | An explanation of the meaning of the concept

Comments | Additional notes about how to use the code

4.4.1.565 Value Set http://hl7.org/fhir/ValueSet/flag-category

URL: https://hl7.org/fhir/R4/valueset-flag-category.html
Patient Care Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined at http://www.hl7.org/Special/committees/patientcare .

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/flag-category
Version: | 4.0.1
Name: | FlagCategory
Title: | Flag Category
Definition: | Example list of general categories for flagged issues. (Not complete or necessarily appropriate.)
Committee: | Patient Care Work Group
OID: | 2.16.840.1.113883.4.642.3.122 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for FlagCategory

• Resource: Flag.category (CodeableConcept / Example)

4.4.1.565.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/flag-category

4.4.1.565.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 10 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/flag-category version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/flag-category

Code | Display | Definition
diet | Diet | Flags related to the subject's dietary needs.
drug | Drug | Flags related to the subject's medications.
lab | Lab | Flags related to performing laboratory tests and related processes (e.g. phlebotomy).
admin | Administrative | Flags related to administrative and financial processes.
contact | Subject Contact | Flags related to coming into contact with the patient.
clinical | Clinical | Flags related to the subject's clinical data.
behavioral | Behavioral | Flags related to behavior.
research | Research | Flags related to research.
advance-directive | Advance Directive | Flags related to subject's advance directives.
safety | Safety | Flags related to safety precautions.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.566 Value Set http://hl7.org/fhir/ValueSet/clinicalimpression-status

URL: https://hl7.org/fhir/R4/valueset-clinicalimpression-status.html
Patient Care Work Group | Maturity Level: 0 | Trial Use | Use Context: Any

This is a value set defined at http://www.hl7.org/Special/committees/patientcare/ .

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/clinicalimpression-status
Version: | 4.0.1
Name: | ClinicalImpressionStatus
Title: | Clinical Impression Status
Definition: | Codes that reflect the current state of a clinical impression within its overall lifecycle.
Committee: | Patient Care Work Group
OID: | 2.16.840.1.113883.4.642.3.1046 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• ConceptMap: Translation source in Canonical Mapping for "Clinical Impression Status"
• Resource: ClinicalImpression.status (code / Required)

4.4.1.566.1 Content Logical Definition

This value set includes codes from the following code systems:

http://hl7.org/fhir/event-status

Code | Display | Canonical Status
in-progress | In Progress | The event is currently occurring.
completed | Completed | The event has now concluded.
entered-in-error | Entered in Error | This electronic record should never have existed, though it is possible that real-world decisions were based on it. (If real-world activity has occurred, the status should be "stopped" rather than "entered-in-error".).

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.567 Value Set http://hl7.org/fhir/ValueSet/patient-contactrelationship

URL: https://hl7.org/fhir/R4/valueset-patient-contactrelationship.html
Patient Administration Work Group | Maturity Level: N | Normative (from v4.0.0) | Use Context: Any

| This page has been approved as part of an ANSI standard. See the Patient Package for further details.

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/patient-contactrelationship
Version: | 4.0.1
Name: | PatientContactRelationship
Title: | Patient Contact Relationship
Definition: | The nature of the relationship between the patient and the contact person.
Committee: | Patient Administration Work Group
OID: | 2.16.840.1.113883.4.642.3.1130 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: Patient.contact.relationship (CodeableConcept / Extensible)

4.4.1.567.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include codes from http://terminology.hl7.org/CodeSystem/v2-0131 where concept is-not-a O

4.4.1.567.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 7 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/v2-0131 version 2.9

All codes from system http://terminology.hl7.org/CodeSystem/v2-0131

Code | Display | Definition
C | Emergency Contact
E | Employer
F | Federal Agency
I | Insurance Company
N | Next-of-Kin
S | State Agency
U | Unknown

Additional Language Displays

Code | Deutsch (German, de) | Nederlands (Dutch, nl)
C | Ansprechpartner in Notfällen | Contact bij nood
E | Arbeitgeber | Werkgever

F | Bundesbehörde | Federale overheidsinstelling
I | Versicherung | Verzekeraar
N | Kontaktperson | Familielid
S | Landesbehörde | Staat overheidsinstelling
U | unbekannt | Onbekend

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.568 Value Set http://hl7.org/fhir/ValueSet/list-order

URL: https://hl7.org/fhir/R4/valueset-list-order.html
FHIR Infrastructure Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/list-order
Version: | 4.0.1
Name: | ListOrderCodes
Title: | List Order Codes
Definition: | Base values for the order of the items in a list resource.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.324 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ListOrderCodes
• Resource: List.orderedBy (CodeableConcept / Preferred)
• Resource: Composition.section.orderedBy (CodeableConcept / Preferred)
• Profile: ClinicalDocument: Composition.section.orderedBy (CodeableConcept / Preferred)
• Profile: Catalog: Composition.section.orderedBy (CodeableConcept / Preferred)

4.4.1.568.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/list-order

4.4.1.568.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 8 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/list-order version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/list-order

Code | Display | Definition
user | Sorted by User | The list was sorted by a user. The criteria the user used are not specified.
system | Sorted by System | The list was sorted by the system. The criteria the user used are not specified; define additional codes to specify a particular order (or use other defined codes).
event-date | Sorted by Event Date | The list is sorted by the data of the event. This can be used when the list has items which are dates with past or future events.
entry-date | Sorted by Item Date | The list is sorted by the date the item was added to the list. Note that the date added to the list is not explicit in the list itself.
priority | Sorted by Priority | The list is sorted by priority. The exact method in which priority has been determined is not specified.
alphabetic | Sorted Alphabetically | The list is sorted alphabetically by an unspecified property of the items in the list.
category | Sorted by Category | The list is sorted categorically by an unspecified property of the items in the list.
patient | Sorted by Patient | The list is sorted by patient, with items for each patient grouped together.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)

Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.569 Value Set http://hl7.org/fhir/ValueSet/appointment-cancellation-reason

URL: https://hl7.org/fhir/R4/valueset-appointment-cancellation-reason.html
Patient Administration Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/appointment-cancellation-reason
Version: | 4.0.1
Name: | AppointmentCancellationReason
Title: | Appointment cancellation reason
Definition: | This example value set defines a set of reasons for the cancellation of an appointment.
Committee: | Patient Administration Work Group
OID: | 2.16.840.1.113883.4.642.3.1381 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for AppointmentCancellationReason

• Resource: Appointment.cancelationReason (CodeableConcept / Example)

4.4.1.569.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/appointment-cancellation-reason

4.4.1.569.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 32 concepts

All codes from system http://terminology.hl7.org/CodeSystem/appointment-cancellation-reason

Code | Display | Definition
pat | Patient
pat-crs | Patient: Canceled via automated reminder system
pat-cpp | Patient: Canceled via Patient Portal
pat-dec | Patient: Deceased
pat-fb | Patient: Feeling Better
pat-lt | Patient: Lack of Transportation
pat-mt | Patient: Member Terminated
pat-mv | Patient: Moved
pat-preg | Patient: Pregnant
pat-swl | Patient: Scheduled from Wait List
pat-ucp | Patient: Unhappy/Changed Provider
prov | Provider
prov-pers | Provider: Personal
prov-dch | Provider: Discharged
prov-edu | Provider: Edu/Meeting
prov-hosp | Provider: Hospitalized
prov-labs | Provider: Labs Out of Acceptable Range
prov-mri | Provider: MRI Screening Form Marked Do Not Proceed
prov-onc | Provider: Oncology Treatment Plan Changes
maint | Equipment Maintenance/Repair
meds-inc | Prep/Med Incomplete
other | Other
oth-cms | Other: CMS Therapy Cap Service Not Authorized
oth-err | Other: Error
oth-fin | Other: Financial
oth-iv | Other: Improper IV Access/Infiltrate IV
oth-int | Other: No Interpreter Available
oth-mu | Other: Prep/Med/Results Unavailable
oth-room | Other: Room/Resource Maintenance
oth-oerr | Other: Schedule Order Error

oth-swie | Other: Silent Walk In Error
oth-weath | Other: Weather

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.570 Value Set http://hl7.org/fhir/ValueSet/icd-10-procedures

URL: https://hl7.org/fhir/R4/valueset-icd-10-procedures.html
Financial Management Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/icd-10-procedures
Version: | 4.0.1
Name: | ICD-10ProcedureCodes
Title: | ICD-10 Procedure Codes
Definition: | This value set includes sample ICD-10 Procedure codes.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.574 (for OID based terminology systems)
Copyright: | This is an example set.
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ICD-10ProcedureCodes

• Resource: ExplanationOfBenefit.procedure.procedure[x] (CodeableConcept|Reference(Procedure) / Example)

4.4.1.570.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/sid/ex-icd-10-procedures

4.4.1.570.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 3 concepts

Expansion based on http://hl7.org/fhir/sid/ex-icd-10-procedures version 4.0.1

All codes from system http://hl7.org/fhir/sid/ex-icd-10-procedures

Code | Display | Definition
123001 | PROC-1 | Procedure 1
123002 | PROC-2 | Procedure 2
123003 | PROC-3 | Procedure 3

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.571 Value Set http://hl7.org/fhir/ValueSet/basic-resource-type

URL: https://hl7.org/fhir/R4/valueset-basic-resource-type.html
FHIR Infrastructure Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/basic-resource-type
Version: | 4.0.1
Name: | BasicResourceTypes
Title: | Basic Resource Types
Definition: | This value set defines codes for resources not yet supported by (or which will never be supported by) FHIR. Many of the codes listed here will eventually be turned into official resources. However, there is no guarantee that any particular resource will be created nor that the scope will be exactly as defined by the codes presented here. Codes in this set will be deprecated if/when formal resources are defined that encompass these concepts.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.138 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for BasicResourceTypes
• Resource: Basic.code (CodeableConcept / Example)

4.4.1.571.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/basic-resource-type

4.4.1.571.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 16 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/basic-resource-type version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/basic-resource-type

Code | Display | Definition
consent | Consent | An assertion of permission for an activity or set of activities to occur, possibly subject to particular limitations; e.g. surgical consent, information disclosure consent, etc.
referral | Referral | A request that care of a particular type be provided to a patient. Could involve the transfer of care, a consult, etc.
advevent | Adverse Event | An undesired reaction caused by exposure to some agent (e.g. a medication, immunization, food, or environmental agent).
aptmtreq | Appointment Request | A request that a time be scheduled for a type of service for a specified patient, potentially subject to other constraints
transfer | Transfer | The transition of a patient or set of material from one location to another
diet | Diet | The specification of a set of food and/or other nutritional material to be delivered to a patient.
adminact | Administrative Activity | An occurrence of a non-care-related event in the healthcare domain, such as approvals, reviews, etc.
exposure | Exposure | Record of a situation where a subject was exposed to a substance. Usually of interest to public health.
investigation | Investigation | A formalized inquiry into the circumstances surrounding a particular unplanned event or potential event for the purposes of identifying possible causes and contributing factors for the event
account | Account | A financial instrument used to track costs, charges or other amounts.
invoice | Invoice | A request for payment for goods and/or services. Includes the idea of a healthcare insurance claim.
adjudicat | Invoice Adjudication | The determination of what will be paid against a particular invoice based on coverage, plan rules, etc.
predetreq | Pre-determination Request | A request for a pre-determination of the cost that would be paid under an insurance plan for a hypothetical claim for goods or services
predetermine | Predetermination | An adjudication of what would be paid under an insurance plan for a hypothetical claim for goods or services
study | Study | An investigation to determine information about a particular therapy or product
protocol | Protocol | A set of (possibly conditional) steps to be taken to achieve some aim. Includes study protocols, treatment protocols, emergency protocols, etc.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.572 Value Set http://hl7.org/fhir/ValueSet/testscript-operation-codes

URL: https://hl7.org/fhir/R4/valueset-testscript-operation-codes.html
FHIR Infrastructure Work Group | Maturity Level: 2 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/testscript-operation-codes
Version: | 4.0.1
Name: | TestScriptOperationCode
Title: | Test script operation code
Definition: | This value set defines a set of codes that are used to indicate the supported operations of a testing engine or tool.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.704 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for TestScriptOperationCode
• Resource: TestScript.setup.action.operation.type (Coding / Extensible)

4.4.1.572.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/testscript-operation-codes

4.4.1.572.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 45 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/testscript-operation-codes version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/testscript-operation-codes

Code | Display | Definition
read | Read | Read the current state of the resource.
vread | Version Read | Read the state of a specific version of the resource.
update | Update | Update an existing resource by its id.
updateCreate | Create using Update | Update an existing resource by its id (or create it if it is new).
patch | Patch | Patch an existing resource by its id.
delete | Delete | Delete a resource.
deleteCondSingle | Conditional Delete Single | Conditionally delete a single resource based on search parameters.
deleteCondMultiple | Conditional Delete Multiple | Conditionally delete one or more resources based on search parameters.
history | History | Retrieve the change history for a particular resource or resource type.
create | Create | Create a new resource with a server assigned id.
search | Search | Search based on some filter criteria.
batch | Batch | Update, create or delete a set of resources as independent actions.
transaction | Transaction | Update, create or delete a set of resources as a single transaction.
capabilities | Capabilities | Get a capability statement for the system.
apply | $apply | Realizes an ActivityDefinition in a specific context
closure | $closure | Closure Table Maintenance
find-matches | $find-matches | Finding Codes based on supplied properties
conforms | $conforms | Compare two systems CapabilityStatements
data-requirements | $data-requirements | Aggregates and returns the parameters and data requirements for a resource and all its dependencies as a single module definition
document | $document | Generate a Document
evaluate | $evaluate | Request clinical decision support guidance based on a specific decision support module
evaluate-measure | $evaluate-measure | Invoke an eMeasure and obtain the results
everything | $everything | Return all the related information as described in the Encounter or Patient
expand | $expand | Value Set Expansion
find | $find | Find a functional list
graphql | $graphql | Invoke a GraphQL query
implements | $implements | Test if a server implements a client's required operations
lastn | $lastn | Last N Observations Query
lookup | $lookup | Concept Look Up and Decomposition
match | $match | Find patient matches using MPI based logic
meta | $meta | Access a list of profiles, tags, and security labels
meta-add | $meta-add | Add profiles, tags, and security labels to a resource
meta-delete | $meta-delete | Delete profiles, tags, and security labels for a resource
populate | $populate | Populate Questionnaire
populatehtml | $populatehtml | Generate HTML for Questionnaire
populatelink | $populatelink | Generate a link to a Questionnaire completion webpage

process-message | $process-message | Process a message according to the defined event
questionnaire | $questionnaire | Build Questionnaire
stats | $stats | Observation Statistics
subset | $subset | Fetch a subset of the CapabilityStatement resource
subsumes | $subsumes | CodeSystem Subsumption Testing
transform | $transform | Model Instance Transformation
translate | $translate | Concept Translation
validate | $validate | Validate a resource
validate-code | $validate-code | ValueSet based Validation

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.573 Value Set http://hl7.org/fhir/ValueSet/study-type

URL: https://hl7.org/fhir/R4/valueset-study-type.html
Clinical Decision Support Work Group | Maturity Level: 0 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/study-type
Version: | 4.0.1
Name: | StudyType
Title: | StudyType
Definition: | Types of research studies (types of research methods).
Committee: | Clinical Decision Support Work Group
OID: | 2.16.840.1.113883.4.642.3.1349 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for StudyType
• Resource: EffectEvidenceSynthesis.studyType (CodeableConcept / Extensible)
• Resource: RiskEvidenceSynthesis.studyType (CodeableConcept / Extensible)

4.4.1.573.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/study-type

4.4.1.573.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 7 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/study-type version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/study-type

Code | Display | Definition
RCT | randomized trial | randomized controlled trial.
CCT | controlled trial (non-randomized) | controlled (but not randomized) trial.
cohort | comparative cohort study | observational study comparing cohorts.
case-control | case-control study | case-control study.
series | uncontrolled cohort or case series | uncontrolled cohort or case series.
case-report | case report | a single case report.
mixed | mixed methods | a combination of 1 or more types of studies.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')

Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.574 Value Set http://hl7.org/fhir/ValueSet/sequence-quality-standardSequence

URL: https://hl7.org/fhir/R4/valueset-sequence-quality-standardSequence.html

This value set could not be expanded by the publication tooling: Error from server: Unable to provide support for code system https://precision.fda.gov/files/

Clinical Genomics Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/sequence-quality-standardSequence
Version: | 4.0.1
Name: | FDA-StandardSequence
Title: | F d a- standard sequence
Definition: | This value set includes sequence quality standard
Committee: | Clinical Genomics Work Group
OID: | 2.16.840.1.113883.4.642.3.222 (for OID based terminology systems)
Copyright: | Unless otherwise noted, the contents of the FDA website (www.fda.gov)—both text and graphics—are not copyrighted. They are in the public domain and may be republished, reprinted and otherwise used freely by anyone without the need to obtain permission from FDA. Credit to the U.S. Food and Drug Administration as the source is appreciated but not required. FDA appreciates being informed about the use of website materials. Contact FDA, HFI-50, Rockville, MD 20857 or email webmail@oc.fda.gov.
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: MolecularSequence.quality.standardSequence (CodeableConcept / Example)

4.4.1.574.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in https://precision.fda.gov/files/

4.4.1.574.2 Expansion

This expansion generated 01 Nov 2019

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.575 Value Set http://hl7.org/fhir/ValueSet/vision-product

URL: https://hl7.org/fhir/R4/valueset-vision-product.html
Financial Management Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/vision-product
Version: | 4.0.1
Name: | ExampleVisionPrescriptionProductCodes
Title: | Example Vision Prescription Product Codes
Definition: | This value set includes a smattering of Prescription Product codes.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.658 (for OID based terminology systems)
Copyright: | This is an example set.
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ExampleVisionPrescriptionProductCodes

• Resource: VisionPrescription.lensSpecification.product (CodeableConcept / Example)

4.4.1.575.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/ex-visionprescriptionproduct

4.4.1.575.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 2 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/ex-visionprescriptionproduct version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/ex-visionprescriptionproduct

Code	Display	Definition
lens	Lens	A lens to be fitted to a frame to comprise a pair of glasses.
contact	Contact Lens	A lens to be fitted for wearing directly on an eye.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.576 Value Set http://hl7.org/fhir/ValueSet/bodysite-laterality

URL: https://hl7.org/fhir/R4/valueset-bodysite-laterality.html
Imaging Integration Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/bodysite-laterality
Version: | 4.0.1
Name: | Laterality
Title: | Laterality
Definition: | Laterality: SNOMED-CT concepts for 'left', 'right', and 'bilateral'
Committee: | Imaging Integration Work Group
OID: | 2.16.840.1.113883.4.642.3.287 (for OID based terminology systems)
Copyright: | This value set includes content from SNOMED CT, which is copyright © 2002+ International Health Terminology Standards Development Organisation (IHTSDO), and distributed by agreement between IHTSDO and HL7. Implementer use of SNOMED CT is not covered by this agreement
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: ImagingStudy.series.laterality (Coding / Example)

4.4.1.576.1 Content Logical Definition

This value set includes codes from the following code systems:

http://snomed.info/sct

Code	Display
419161000	Unilateral left
419465000	Unilateral right
51440002	Bilateral

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')

Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.577 Value Set http://hl7.org/fhir/ValueSet/reaction-event-severity

URL: https://hl7.org/fhir/R4/valueset-reaction-event-severity.html
Patient Care Work Group | Maturity Level: 3 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/reaction-event-severity
Version: | 4.0.1
Name: | AllergyIntoleranceSeverity
Title: | AllergyIntoleranceSeverity
Definition: | Clinical assessment of the severity of a reaction event as a whole, potentially considering multiple different manifestations.
Committee: | Patient Care Work Group
OID: | 2.16.840.1.113883.4.642.3.135 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for AllergyIntoleranceSeverity

• Resource: AllergyIntolerance.reaction.severity (code / Required)

4.4.1.577.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/reaction-event-severity

4.4.1.577.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 3 concepts

Expansion based on http://hl7.org/fhir/reaction-event-severity version 4.0.1

All codes from system http://hl7.org/fhir/reaction-event-severity

Code | Display | Definition
mild | Mild | Causes mild physiological effects.
moderate | Moderate | Causes moderate physiological effects.
severe | Severe | Causes severe physiological effects.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.578 Value Set http://hl7.org/fhir/ValueSet/provider-qualification

URL: https://hl7.org/fhir/R4/valueset-provider-qualification.html
Financial Management Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/provider-qualification
Version: | 4.0.1
Name: | ExampleProviderQualificationCodes
Title: | Example Provider Qualification Codes
Definition: | This value set includes sample Provider Qualification codes.
Committee: | Financial Management Work Group

OID: | 2.16.840.1.113883.4.642.3.570 (for OID based terminology systems)
Copyright: | This is an example set.
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ExampleProviderQualificationCodes

• Resource: ExplanationOfBenefit.careTeam.qualification (CodeableConcept / Example)

4.4.1.578.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/ex-providerqualification

4.4.1.578.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 3 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/ex-providerqualification version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/ex-providerqualification

Code | Display | Definition
311405 | Dentist | Dentist General Practitioner (DDS, DDM).
604215 | Ophthalmologist | Ophthalmologist.
604210 | Optometrist | Optometrist.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.579 Value Set http://hl7.org/fhir/ValueSet/eligibilityresponse-purpose

URL: https://hl7.org/fhir/R4/valueset-eligibilityresponse-purpose.html
Financial Management Work Group | Maturity Level: 2 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/eligibilityresponse-purpose
Version: | 4.0.1
Name: | EligibilityResponsePurpose
Title: | EligibilityResponsePurpose
Definition: | A code specifying the types of information being requested.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.1184 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for EligibilityResponsePurpose

• Resource: CoverageEligibilityResponse.purpose (code / Required)

4.4.1.579.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/eligibilityresponse-purpose

4.4.1.579.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 4 concepts

Expansion based on http://hl7.org/fhir/eligibilityresponse-purpose version 4.0.1

All codes from system http://hl7.org/fhir/eligibilityresponse-purpose

Code | Display | Definition
auth-requirements | Coverage auth-requirements | The prior authorization requirements for the listed, or discovered if specified, converages for the categories of service and/or specifed biling codes are requested.
benefits | Coverage benefits | The plan benefits and optionally benefits consumed for the listed, or discovered if specified, converages are requested.
discovery | Coverage Discovery | The insurer is requested to report on any coverages which they are aware of in addition to any specifed.
validation | Coverage Validation | A check that the specified coverages are in-force is requested.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.580 Value Set http://hl7.org/fhir/ValueSet/consent-data-meaning

URL: https://hl7.org/fhir/R4/valueset-consent-data-meaning.html
Community Based Collaborative Care Work Group | Maturity Level: 2 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/consent-data-meaning
Version: | 4.0.1
Name: | ConsentDataMeaning
Title: | ConsentDataMeaning
Definition: | How a resource reference is interpreted when testing consent restrictions.
Committee: | Community Based Collaborative Care Work Group
OID: | 2.16.840.1.113883.4.642.3.759 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ConsentDataMeaning
• Resource: Consent.provision.data.meaning (code / Required)

4.4.1.580.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/consent-data-meaning

4.4.1.580.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 4 concepts

Expansion based on http://hl7.org/fhir/consent-data-meaning version 4.0.1

All codes from system http://hl7.org/fhir/consent-data-meaning

Code | Display | Definition
instance | Instance | The consent applies directly to the instance of the resource.
related | Related | The consent applies directly to the instance of the resource and instances it refers to.
dependents | Dependents | The consent applies directly to the instance of the resource and instances that refer to it.
authoredby | AuthoredBy | The consent applies to instances of resources that are authored by.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.581 Value Set http://hl7.org/fhir/ValueSet/encounter-special-courtesy

URL: https://hl7.org/fhir/R4/valueset-encounter-special-courtesy.html
Patient Administration Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/encounter-special-courtesy
Version: | 4.0.1
Name: | SpecialCourtesy
Title: | Special courtesy
Definition: | This value set defines a set of codes that can be used to indicate special courtesies provided to the patient.
Committee: | Patient Administration Work Group
OID: | 2.16.840.1.113883.4.642.3.252 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: Encounter.hospitalization.specialCourtesy (CodeableConcept / Preferred)

4.4.1.581.1 Content Logical Definition

This value set includes codes from the following code systems:

http://terminology.hl7.org/CodeSystem/v3-EncounterSpecialCourtesy

Code | Display
EXT | extended courtesy | extended courtesy
NRM | normal courtesy | normal courtesy
PRF | professional courtesy | professional courtesy
STF | staff | Courtesies extended to the staff of the entity providing service.
VIP | very important person | very important person

http://terminology.hl7.org/CodeSystem/v3-NullFlavor

Code | Display
UNK | unknown | Description:A proper value is applicable, but not known. Usage Notes: This means the actual value is not known. If the only thing that is unknown is how to properly express the value in the necessary constraints (value set, datatype, etc.), then the OTH or UNC flavor should be used. No properties should be included for a datatype with this property unless: Those properties themselves directly translate to a semantic of "unknown". (E.g. a local code sent as a translation that conveys 'unknown') Those properties further qualify the nature of what is unknown. (E.g. specifying a use code of "H" and a URL prefix of "tel:" to convey that it is the home phone number that is unknown.)

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.582 Value Set http://hl7.org/fhir/ValueSet/vision-eye-codes

URL: https://hl7.org/fhir/R4/valueset-vision-eye-codes.html
Financial Management Work Group | Maturity Level: 2 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/vision-eye-codes
Version: | 4.0.1
Name: | VisionEyes
Title: | VisionEyes
Definition: | A coded concept listing the eye codes.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.660 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for VisionEyes

• Resource: VisionPrescription.lensSpecification.eye (code / Required)

4.4.1.582.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/vision-eye-codes

4.4.1.582.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 2 concepts

Expansion based on http://hl7.org/fhir/vision-eye-codes version 4.0.1

All codes from system http://hl7.org/fhir/vision-eye-codes

Code | Display | Definition
right | Right Eye | Right Eye.
left | Left Eye | Left Eye.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.583 Value Set http://hl7.org/fhir/ValueSet/codesystem-altcode-kind

URL: https://hl7.org/fhir/R4/valueset-codesystem-altcode-kind.html
Vocabulary Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/codesystem-altcode-kind
Version: | 4.0.1
Name: | AlternativeCodeKind
Title: | AlternativeCodeKind
Definition: | Indicates the type of use for which the code is defined.
Committee: | Vocabulary Work Group
OID: | 2.16.840.1.113883.4.642.3.952 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for AlternativeCodeKind

• Extension: http://hl7.org/fhir/StructureDefinition/codesystem-alternate: alternate (Coding / Extensible)

4.4.1.583.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/codesystem-altcode-kind

4.4.1.583.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 5 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/codesystem-altcode-kind version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/codesystem-altcode-kind

Code | Display | Definition
alternate | Alternate Code | The code is an alternative code that can be used in any of the circumstances that the primary code can be used.
deprecated | Deprecated | The code should no longer be used, but was used in the past.
case-insensitive | Case Insensitive | The code is an alternative to be used when a case insensitive code is required (when the primary codes are case sensitive).
case-sensitive | Case Sensitive | The code is an alternative to be used when a case sensitive code is required (when the primary codes are case insensitive).

expression | Expression | The code is an alternative for the primary code that is built using the expression grammar defined by the code system.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.584 Value Set http://hl7.org/fhir/ValueSet/care-plan-activity-status

URL: https://hl7.org/fhir/R4/valueset-care-plan-activity-status.html
Patient Care Work Group | Maturity Level: 2 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/care-plan-activity-status
Version: | 4.0.1
Name: | CarePlanActivityStatus
Title: | CarePlanActivityStatus
Definition: | Codes that reflect the current state of a care plan activity within its overall life cycle.
Committee: | Patient Care Work Group
OID: | 2.16.840.1.113883.4.642.3.146 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for CarePlanActivityStatus
• ConceptMap: Translation source in Canonical Mapping for "CarePlanActivityStatus"
• Resource: CarePlan.activity.detail.status (code / Required)

4.4.1.584.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/care-plan-activity-status

4.4.1.584.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 9 concepts

Expansion based on http://hl7.org/fhir/care-plan-activity-status version 4.0.1

All codes from system http://hl7.org/fhir/care-plan-activity-status

Lvl | Code | Display | Definition | Canonical Status
0 | not-started | Not Started | Care plan activity is planned but no action has yet been taken. | ~draft
0 | scheduled | Scheduled | Appointment or other booking has occurred but activity has not yet begun. | ~accepted
0 | in-progress | In Progress | Care plan activity has been started but is not yet complete. | ~active
0 | on-hold | On Hold | Care plan activity was started but has temporarily ceased with an expectation of resumption at a future time. | ~suspended
0 | completed | Completed | Care plan activity has been completed (more or less) as planned. | ~complete
0 | cancelled | Cancelled | The planned care plan activity has been withdrawn. | ~abandoned
1 | stopped | Stopped | The planned care plan activity has been ended prior to completion after the activity was started. | ~failed
0 | unknown | Unknown | The current state of the care plan activity is not known. Note: This concept is not to be used for "other" - one of the listed statuses is presumed to apply, but the authoring/source system does not know which one. | ~unknown
0 | entered-in-error | Entered in Error | Care plan activity was entered in error and voided. | ~error

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application

Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.585 Value Set http://hl7.org/fhir/ValueSet/linkage-type

URL: https://hl7.org/fhir/R4/valueset-linkage-type.html
Patient Care Work Group | Maturity Level: 0 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/linkage-type
Version: | 4.0.1
Name: | LinkageType
Title: | LinkageType
Definition: | Used to distinguish different roles a resource can play within a set of linked resources.
Committee: | Patient Care Work Group
OID: | 2.16.840.1.113883.4.642.3.314 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for LinkageType

• Resource: Linkage.item.type (code / Required)

4.4.1.585.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/linkage-type

4.4.1.585.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 3 concepts

Expansion based on http://hl7.org/fhir/linkage-type version 4.0.1

All codes from system http://hl7.org/fhir/linkage-type

Code | Display | Definition
source | Source of Truth | The resource represents the "source of truth" (from the perspective of this Linkage resource) for the underlying event/condition/etc.
alternate | Alternate Record | The resource represents an alternative view of the underlying event/condition/etc. The resource may still be actively maintained, even though it is not considered to be the source of truth.
historical | Historical/Obsolete Record | The resource represents an obsolete record of the underlying event/condition/etc. It is not expected to be actively maintained.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.586 Value Set http://hl7.org/fhir/ValueSet/search-param-type

URL: https://hl7.org/fhir/R4/valueset-search-param-type.html
FHIR Infrastructure Work Group | Maturity Level: N | Normative (from v4.0.0) | Use Context: Any

| This page has been approved as part of an ANSI standard. See the Conformance Package for further details.

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/search-param-type
Version: | 4.0.1
Name: | SearchParamType
Title: | SearchParamType
Definition: | Data types allowed to be used for search parameters.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.11 (for OID based terminology systems)

Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for SearchParamType

• Resource: CapabilityStatement.rest.resource.searchParam.type (code / Required)

• Resource: SearchParameter.type (code / Required)

• Resource: OperationDefinition.parameter.searchType (code / Required)

4.4.1.586.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/search-param-type

4.4.1.586.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 9 concepts

Expansion based on http://hl7.org/fhir/search-param-type version 4.0.1

All codes from system http://hl7.org/fhir/search-param-type

Code | Display | Definition
number | Number | Search parameter SHALL be a number (a whole number, or a decimal).
date | Date/DateTime | Search parameter is on a date/time. The date format is the standard XML format, though other formats may be supported.
string | String | Search parameter is a simple string, like a name part. Search is case-insensitive and accent-insensitive. May match just the start of a string. String parameters may contain spaces.
token | Token | Search parameter on a coded element or identifier. May be used to search through the text, display, code and code/codesystem (for codes) and label, system and key (for identifier). Its value is either a string or a pair of namespace and value, separated by a "|", depending on the modifier used.
reference | Reference | A reference to another resource (Reference or canonical).
composite | Composite | A composite search parameter that combines a search on two values together.
quantity | Quantity | A search parameter that searches on a quantity.
uri | URI | A search parameter that searches on a URI (RFC 3986).
special | Special | Special logic applies to this parameter per the description of the search parameter.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.587 Value Set http://hl7.org/fhir/ValueSet/product-category

URL: https://hl7.org/fhir/R4/valueset-product-category.html
Orders and Observations Work Group | Maturity Level: 0 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/product-category
Version: | 4.0.1
Name: | BiologicallyDerivedProductCategory
Title: | BiologicallyDerivedProductCategory
Definition: | Biologically Derived Product Category.
Committee: | Orders and Observations Work Group
OID: | 2.16.840.1.113883.4.642.3.900 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for BiologicallyDerivedProductCategory

• Resource: BiologicallyDerivedProduct.productCategory (code / Required)

4.4.1.587.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/product-category

4.4.1.587.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 5 concepts

Expansion based on http://hl7.org/fhir/product-category version 4.0.1

All codes from system http://hl7.org/fhir/product-category

Code | Display | Definition
organ | Organ | A collection of tissues joined in a structural unit to serve a common function.
tissue | Tissue | An ensemble of similar cells and their extracellular matrix from the same origin that together carry out a specific function.
fluid | Fluid | Body fluid.
cells | Cells | Collection of cells.
biologicalAgent | BiologicalAgent | Biological agent of unspecified type.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.588 Value Set http://hl7.org/fhir/ValueSet/goal-start-event

URL: https://hl7.org/fhir/R4/valueset-goal-start-event.html
Patient Care Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined at http://www.hl7.org/Special/committees/patientcare/ .

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/goal-start-event
Version: | 4.0.1
Name: | GoalStartEvent
Title: | Goal start event
Definition: | Identifies types of events that might trigger the start of a goal.
Committee: | Patient Care Work Group
OID: | 2.16.840.1.113883.4.642.3.279 (for OID based terminology systems)
Copyright: | This resource includes content from SNOMED Clinical Terms® (SNOMED CT®) which is copyright of the International Health Terminology Standards Development Organisation (IHTSDO). Implementers of these specifications must have the appropriate SNOMED CT Affiliate license - for more information contact http://www.snomed.org/snomed-ct/get-snomed-ct or info@snomed.org
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: Goal.start[x] (date|CodeableConcept / Example)
• Resource: PlanDefinition.goal.start (CodeableConcept / Example)
• Profile: ShareablePlanDefinition: PlanDefinition.goal.start (CodeableConcept / Example)
• Profile: ComputablePlanDefinition: PlanDefinition.goal.start (CodeableConcept / Example)
• Profile: CDSHooksServicePlanDefinition: PlanDefinition.goal.start (CodeableConcept / Example)

4.4.1.588.1 Content Logical Definition

This value set includes codes from the following code systems:

http://snomed.info/sct

Code | Display
32485007 | Admission to hospital
308283009 | Discharge from hospital
442137000 | Completion time of procedure
386216000 | Childbirth

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.589 Value Set http://hl7.org/fhir/ValueSet/contract-term-subtype

URL: https://hl7.org/fhir/R4/valueset-contract-term-subtype.html
Financial Management Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/contract-term-subtype
Version: | 4.0.1
Name: | ContractTermSubtypeCodes
Title: | Contract Term Subtype Codes
Definition: | This value set includes sample Contract Term SubType codes.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.735 (for OID based terminology systems)
Copyright: | This is an example set.
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ContractTermSubtypeCodes

• Resource: Contract.term.subType (CodeableConcept / Example)

4.4.1.589.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/contracttermsubtypecodes

4.4.1.589.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 3 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/contracttermsubtypecodes version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/contracttermsubtypecodes

Code | Display | Definition
condition | Condition | Terms that go to the very root of a contract.
warranty | Warranty | Less imperative than a condition, so the contract will survive a breach
innominate | Innominate | Breach of which might or might not go to the root of the contract depending upon the nature of the breach

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.590 Value Set http://hl7.org/fhir/ValueSet/approach-site-codes

URL: https://hl7.org/fhir/R4/valueset-approach-site-codes.html
FHIR Infrastructure Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/approach-site-codes
Version: | 4.0.1
Name: | SNOMEDCTAnatomicalStructureForAdministrationSiteCodes
Title: | SNOMED CT Anatomical Structure for Administration Site Codes
Definition: | This value set includes Anatomical Structure codes from SNOMED CT - provided as an exemplar.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.346 (for OID based terminology systems)
Copyright: | This value set includes content from SNOMED CT, which is copyright © 2002 International Health Terminology Standards Development Organisation (IHTSDO), and distributed by agreement between IHTSDO and HL7. Implementer use of SNOMED CT is not covered by this agreement.
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: MedicationAdministration.dosage.site (CodeableConcept / Example)

• Resource: Dosage.site (CodeableConcept / Example)

4.4.1.590.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include codes from http://snomed.info/sct where concept is-a 91723000 (Anatomical structure)

4.4.1.590.2 Expansion

This expansion generated 01 Nov 2019

This value set has >1000 codes in it. In order to keep the publication size manageable, only a selection (1000 codes) of the whole set of codes is shown

All codes from system http://snomed.info/sct

Code | Display | Definition
106004 | Posterior carpal region
107008 | Fetal part of placenta
108003 | Condylar emissary vein
110001 | Visceral layer of Bowman's capsule
111002 | Parathyroid gland
116007 | Subcutaneous tissue of medial surface of index finger
124002 | Coronoid process of mandible
149003 | Central pair of microtubules, cilium or flagellum, not bacterial
155008 | Deep circumflex artery of ilium
167005 | Supraclavicular part of brachial plexus
202009 | Anterior division of renal artery
205006 | Left commissure of aortic valve
206007 | Gluteus maximus muscle
221001 | Articular surface, phalanges, of fourth metacarpal bone
227002 | Canal of Hering
233006 | Hepatocolic ligament
235004 | Superior labial artery
246001 | Lateral vestibular nucleus
247005 | Mesotympanum
251007 | Pectoral region
256002 | Kupffer cell
263002 | Thoracic nerve
266005 | Right lower lobe of lung
272005 | Superior articular process of lumbar vertebra
273000 | Lateral myocardium
283001 | Central axillary lymph node
284007 | Flexor tendon and tendon sheath of fourth toe
289002 | Metacarpophalangeal joint of index finger
301000 | Fifth metatarsal bone
311007 | Plantar surface of great toe
315003 | Skin of umbilicus
318001 | Cardiac impression of liver
344001 | Ankle
345000 | Penetrating atrioventricular bundle
356000 | Reticular corium
393006 | Wall of urinary bladder
402006 | Dental branches of inferior alveolar artery
405008 | Posterior temporal diploic vein
414003 | Gastric fundus
422005 | Inferior surface of tongue
446003 | Trochanteric bursa
457008 | Collateral ligament
461002 | Lateral corticospinal tract
464005 | Basophilic normoblast

465006 | Ascending frontal gyrus
471000 | Flexor hallucis longus in leg
480000 | Cardiopulmonary circulatory system
485005 | TC - Transverse colon
528006 | Costal surface of lung
552004 | Vagus nerve parasympathetic fibers to cardiac plexus
565008 | Intervertebral disc space of fifth lumbar vertebra
582005 | Head of phalanx of great toe
587004 | Capsule of proximal interphalangeal joint of third toe
589001 | Interventricular septum
595000 | Palpebral fissure
608002 | Subcutaneous tissue of philtrum
621009 | Multivesicular body, internal vesicles
635006 | Tuberosity of distal phalanx of little toe
650002 | Superior articular process of seventh thoracic vertebra
660006 | Tracheal mucous membrane
661005 | Jaw region
667009 | Embryological structure
688000 | Fetal hyaloid artery
691000 | Small intestine submucosa
692007 | Body of ischium
723004 | Dense intermediate filament bundles
774007 | Head and neck
790007 | Visceral surface of liver
798000 | Deep temporal veins
808000 | Posterior intercostal artery
809008 | Fetal chondrocranium
823005 | Posterior cervical spinal cord nerve root
830004 | Spinous process of fifth thoracic vertebra
836005 | Oral region of face
885000 | Lamina muscularis of colonic mucous membrane
895007 | Anterior cruciate ligament of knee joint
917004 | Superior laryngeal aperture
921006 | Thyrohyoid branch of ansa cervicalis
947002 | Crus of diaphragm
955009 | Bronchus
976004 | Ovarian vein
996007 | Meningeal branch of occipital artery
1005009 | Entire diaphragmatic lymph node
1012000 | Structure of fibrous portion of pericardium
1015003 | Structure of peritonsillar tissue
1028005 | Sebaceous gland structure
1030007 | Structure of vesicular bursa of sternohyoid muscle
1063000 | Frontozygomatic suture of skull
1075005 | Promonocyte
1076006 | Subcutaneous prepatellar bursa
1086007 | Female
1087003 | Sternothyroid muscle
1092001 | Superior occipital gyrus
1099005 | Thymic cortex
1101003 | Cranial cavity
1106008 | Major calyx
1110006 | Tarsal gland
1122009 | Inferior longitudinal muscle of tongue
1136004 | Aortopulmonary septum
1159005 | Frenulum linguae
1172006 | Odontoid process of axis
1173001 | Mandibular nerve
1174007 | Chromosomes, group E
1193009 | Teres major muscle
1216008 | Synostosis
1231004 | Central nervous system meninges
1236009 | Duodenal serosa
1243003 | Inferior articular process of sixth cervical vertebra
1246006 | Dorsal digital nerves of radial nerve
1263005 | Distinctive arrangement of microtubules
1277008 | Vertebral nerve
1307006 | Glottis
1311000 | Telogen hair
1350001 | Deep flexor tendon of index finger
1353004 | Gastric serosa
1403006 | Vastus lateralis muscle

1425000 | Posterior limb of stapes
1439000 | Paravesicular lymph node
1441004 | Laryngeal saccule
1456008 | Yellow fibrocartilage
1467009 | Parietal branch of superficial temporal artery
1484003 | Structure of metatarsal region of foot
1490004 | Soft tissues of trunk
1502004 | Anterior cecal artery
1511004 | Ejaculatory duct
1516009 | Frontomental diameter of head
1527006 | Lamina of fourth thoracic vertebra
1537001 | Intervertebral disc of eleventh thoracic vertebra
1541002 | Coccygeal plexus
1562001 | Nucleus pulposus of intervertebral disc of third lumbar vertebra
1569005 | Nail of third toe
1580005 | Nucleus ventralis lateralis
1581009 | Ileal artery
1584001 | Symphysis
1600003 | Splenius capitis muscle
1605008 | Histioblast
1610007 | Otoconia
1611006 | Paramammary lymph node
1617005 | Intrinsic larynx
1620002 | Metaphase nucleus
1626008 | Third thoracic vertebra
1627004 | Medial collateral ligament of knee joint
1630006 | Supraorbital vein
1631005 | Foregut
1650005 | Hilum of left lung
1655000 | Transverse peduncular tract nucleus
1659006 | Nucleus medialis dorsalis
1684009 | Ligamentum teres of liver
1706004 | Thymic lobule
1707008 | Ventral nuclear group of thalamus
1711002 | Periorbital region
1716007 | Cupula ampullaris
1721005 | Right tonsil
1729007 | Central tegmental tract
1732005 | TD - Thoracic duct
1765002 | Structure of lymphatic vessel of thorax
1780008 | Premelanosome
1781007 | Sacroiliac region
1797002 | Naris
1818002 | Greater circulus arteriosus of iris
1825009 | Root of nose
1832000 | Scleral conjunctiva
1840006 | Arrector pili muscle
1849007 | Pharyngeal recess
1853009 | Structure of suprahyoid muscle
1874005 | Promontory lymph node
1893007 | Joint of upper extremity
1895000 | Musculophrenic vein
1902009 | Skin of external ear
1910005 | Ear
1918003 | Suprarenal aorta
1927002 | Left elbow
1992003 | Porus acusticus internus
1997009 | Cingulum dentis
2010005 | Clavicular facet of scapula
2020000 | Superior thoracic artery
2031008 | Structure of anterior median fissure of spinal cord
2033006 | Right fallopian tube
2044003 | Vaginal nerves
2048000 | Lingual tonsil
2049008 | Chorionic villi
2059009 | Skin of ear lobule
2071003 | Reticular formation of spinal cord
2076008 | Head of phalanx of hand
2083001 | Nucleus ambiguus
2095001 | Accessory sinus
2123001 | Mammilloinfundibular nucleus
2150006 | Urinary tract transitional epithelial cell

2156000 | Glial cell
2160002 | Ligamentum arteriosum
2175005 | Pharyngeal cavity
2182009 | Endometrial zona basalis
2192001 | Clavicular part of pectoralis major muscle
2205003 | Lamina of fifth thoracic vertebra
2209009 | Cerebral basal surface
2236006 | Lesser osseous pelvis
2246008 | Type I hair cell
2255006 | Subserosa
2285001 | Structure of torcular Herophili
2292006 | Structure of nasopharyngeal gland
2302002 | Vein of the knee
2305000 | Structure of spinous process of cervical vertebra
2306004 | Structure of base of third metacarpal bone
2327009 | Salivary seromucous gland
2330002 | Structure of segmental bronchial branches
2332005 | Metencephalon of foetus
2334006 | Renal calyx
2349003 | Structure of nasal suture of skull
2372001 | Structure of medial surface of toe
2383005 | Structure of papillary muscles of right ventricle
2389009 | Structure of superior margin of adrenal gland
2395005 | Structure of transverse facial artery
2397002 | Structure of first metatarsal facet of medial cuneiform bone
2400006 | Universal designation 21
2402003 | Structure of dorsum of foot
2421006 | Structure of submaxillary ganglion
2433001 | Structure of digital tendon and tendon sheath of foot
2436009 | Tunica intima of vein
2453002 | Subcutaneous tissue structure of posterior surface of forearm
2454008 | Structure of articular surface, third metacarpal, of second metacarpal bone
2484000 | Skin structure of frenulum of clitoris
2489005 | Structure of medial check ligament of eye
2499000 | Entire cisterna pontis
2502001 | Membrane of lysosome
2504000 | Structure of pancreatic plexus
2510000 | Femoral triangle structure
2539000 | Structure of superior rectal artery
2543001 | Structure of cuboid articular facet of fourth metatarsal bone
2550002 | Bone structure of phalanx of thumb
2577006 | Structure of gracilis muscle
2579009 | Plasmablast
2592007 | All extremities
2600000 | Structure of flexor pollicis longus muscle tendon
2620004 | Intervertebral disc structure of third thoracic vertebra
2639009 | Neuroendocrine tissue
2653009 | Structure of posterior thalamic radiation of internal capsule
2666009 | Structure of semispinalis capitis muscle
2672009 | Structure of anterior cutaneous branch of lumbosacral plexus
2675006 | Structure of anterior ethmoidal artery
2681003 | Structure of dorsal nerve of penis
2682005 | Bladder mucosa
2686008 | Structure of medial olfactory gyrus
2687004 | Structure of Bowman space
2695000 | Left maxillary sinus structure
2703009 | Entire calcarine artery
2712006 | Structure of capsule of ankle joint
2718005 | Structure of apical foramen of tooth
2726002 | Structure of fold for stapes
2730004 | Entire vitelline vein of placenta
2739003 | Endometrial structure
2741002 | Structure of medial occipitotemporal gyrus
2746007 | Circular layer of gastric muscularis
2748008 | Spinal cord structure
2759004 | Eccrine gland structure
2771005 | Lamina propria of ureter
2789006 | Apocrine gland structure
2792005 | Structure of pars tensa of tympanic membrane
2803000 | Structure of tendon sheath of popliteus muscle
2810006 | Structure of cremasteric fascia
2812003 | Structure of head of femur

2824005 | Structure of spinous process of fourth thoracic vertebra
2826007 | Structure of lamina of fourth lumbar vertebra
2830005 | Structure of dorsal digital nerves of lateral hallux and medial second toe
2839006 | Structure of perivesicular tissues of seminal vesicles
2841007 | Renal artery structure
2845003 | Structure of respiratory epithelium
2848001 | Structure of superficial epigastric artery
2855004 | Structure of accessory cephalic vein
2861001 | Entire gland (organ)
2894003 | Structure of posterior epiglottis
2905008 | Structure of anterior ligament of uterus
2909002 | Structure of posterior portion of diaphragmatic aspect of liver
2920002 | Structure of facial nerve motor branch
2922005 | Structure of posterior papillary muscle of left ventricle
2923000 | Subcutaneous tissue structure of supraorbital area
2954001 | Supernumerary deciduous tooth
2969000 | Anatomical space structure
2979003 | Bone structure of medial cuneiform
2986006 | Structure of talar facet of navicular bone of foot
2998001 | Entire right margin of uterus
3003007 | Internal capsule anterior limb structure
3008003 | White fibrocartilage
3028004 | Transitional epithelial cell
3039001 | Subcutaneous tissue structure of thigh
3042007 | Structure of glomerular urinary pole
3054007 | Structure of articular surface, metacarpal, of phalanx of thumb
3055008 | Structure of bone marrow of vertebral body
3056009 | Structure of anteroventral nucleus of thalamus
3057000 | Nerve structure
3058005 | PNS - Peripheral nervous system
3062004 | Spinal arachnoid
3068000 | Structure of seminal vesicle lumen
3081007 | Mitochondrion in division
3093003 | Structure of tendinous arch of pelvic fascia
3100007 | Clinical crown of tooth
3113001 | Structure of suprachoroidal space
3117000 | Structure of dorsal surface of index finger
3118005 | Structure of trabecula carnea of left ventricle
3120008 | Pleural membrane structure
3134008 | Structure of head of fourth metatarsal bone
3138006 | Bone tissue
3153003 | Structure of tractus olivocochlearis
3156006 | Structure of obturator artery
3159004 | Structure of costocervical trunk
3169005 | Spinal nerve structure
3178004 | Structure of raphe of soft palate
3194006 | Endocardium of right atrium
3198009 | Monostomatic sublingual gland
3215002 | Subcutaneous tissue structure of nuchal region
3222005 | All large arteries
3227004 | Left coronary artery main stem
3236000 | Structure of posterior segment of right upper lobe of lung
3243006 | Structure of parametrial lymph node
3255000 | Papillary area
3262009 | Structure of root canal of tooth
3279003 | Structure of pedicle of third cervical vertebra
3295003 | Structure of ventral anterior nucleus of thalamus
3301002 | Tectopontine fibers
3302009 | Splenic branch of splenic artery
3315000 | Structure of vestibulospinal tract
3332001 | Occipitofrontal diameter of head
3336003 | Haversian canal
3341006 | Right lung structure
3350008 | Entire right commissure of pulmonic valve
3362007 | Intertragal incisure structure
3366005 | Structure of anterior papillary muscle of left ventricle
3370002 | Structure of supporting tissue of rectum
3374006 | Secondary spermatocyte
3377004 | Structure of agger nasi
3382006 | Structure of rima oris
3383001 | Nonsegmented basophil
3394002 | Suboccipitobregmatic diameter of head

3395001 | Structure of superior palpebral arch
3396000 | Structure of mesogastrium
3400000 | Cell of bone
3409004 | Structure of lateral margin of forearm
3417007 | Structure of rotator muscle
3438001 | Deep lymphatic of upper extremity
3444002 | Thalamostriate vein
3447009 | Penetrated oocyte
3460003 | Structure of anterodorsal nucleus of thalamus
3462006 | Structure of commissure of tricuspid valve
3471002 | Posterior midline of trunk
3478008 | Structure of vastus medialis muscle
3481003 | Structure of embryonic testis
3488009 | Annulate lamella, cisternal lumen
3490005 | Subcutaneous tissue structure of suboccipital region
3524005 | Structure of lateral wall of mastoid antrum
3538003 | Structure of capsule of proximal tibiofibular joint
3541007 | Structure of dorsal metatarsal artery
3553006 | Structure of thyroid capsule
3556003 | Structure of dorsal nucleus of trapezoid body
3563003 | Muscularis of ureter
3572006 | Vertebral body
3578005 | Structure of body of gallbladder
3582007 | Structure of gastrophrenic ligament
3584008 | T10 dorsal arch
3594003 | Structure of straight part of longus colli muscle
3608004 | Ischiococcygeus muscle
3609007 | Structure of occipital branch of posterior auricular artery
3646006 | Lamellipodium
3663005 | Structure of tympanic ostium of Eustachian tube
3665003 | Pelvic wall structure
3670005 | Entire subpyloric lymph node
3711007 | Great vessel
3743007 | Structure of lateral thoracic artery
3761003 | Structure of nucleus pulposus of intervertebral disc of first thoracic vertebra
3766008 | Subcutaneous tissue structure of lower extremity
3785006 | Entire dorsal metacarpal ligament
3788008 | Structure of apical segment of right lower lobe of lung
3789000 | Enteroendocrine cell
3810000 | Septal cartilage structure
3838008 | Structure of apex of coccyx
3860006 | Structure of transplanted liver
3865001 | Structure of interscapular region of back
3867009 | Structure of dorsal surface of great toe
3876002 | Subcutaneous tissue structure of femoral region
3877006 | Structure of common carotid plexus
3910004 | Subcutaneous tissue structure of lateral surface of fourth toe
3916005 | Structure of occipital lymph node
3924000 | Structure of pericardiophrenic artery
3931001 | OW - Oval window
3935005 | Head of tenth rib structure
3937002 | Structure of entorhinal cortex
3954005 | Lacrimal sac structure
3956007 | Structure of fifth metatarsal articular facet of fourth metatarsal bone
3959000 | Structure of rectus capitis muscle
3960005 | Olfactory tract structure
3964001 | Structure of gyrus of brain
3966004 | Entire parietal branch of anterior cerebral artery
3977005 | Subcutaneous tissue structure of concha
3984002 | Deep vein of clitoris
3989007 | Structure of medial globus pallidus
4015004 | Chromosomes, group A
4019005 | Posterior commissure of labium majorum
4029003 | Eosinophilic progranulocyte
4061004 | Lateral orbital wall
4066009 | Structure of capsule of proximal interphalangeal joint of index finger
4072009 | Structure of fourth coccygeal vertebra
4081003 | Entire dorsal lingual vein
4093007 | Structure of vagus nerve bronchial branch
4111006 | Macula of the saccule
4117005 | Lumbosacral spinal cord central canal structure
4121003 | Structure of superior frontal sulcus

4146003 | Structure of artery of extremity
4148002 | Structure of palmar surface of little finger
4150005 | Celiac nervous plexus structure
4158003 | Abdominal aortic plexus structure
4159006 | Hyparterial bronchus
4180000 | Both lower extremities
4193005 | Entire extensor tendon and tendon sheath of fifth toe
4205002 | Türk cell
4212006 | Epithelial cells
4215008 | Head of second rib structure
4247003 | Bone structure of first metacarpal
4258007 | Posterior tibial vein
4268002 | Lateral spinorubral tract
4276000 | Structure of inferior articular process of seventh cervical vertebra
4281009 | Structure of middle portion of ileum
4295007 | Structure of paracortical area of lymph node
4303006 | Cartilage canal
4312008 | Anterior midline of abdomen
4317002 | Structure of spinalis muscle
4328003 | Protoplasmic astrocyte
4335006 | Upper jaw region
4352005 | Structure of subchorionic space
4358009 | Structure of lateral surface of little finger
4360006 | Stratum spinosum structure
4369007 | Small intestine mucous membrane structure
4371007 | Structure of fourth metatarsal facet of lateral cuneiform bone
4375003 | Structure of incisure of cartilaginous portion of auditory canal
4377006 | Structure of parafascicular nucleus of thalamus
4394008 | Scala vestibuli structure
4402002 | Structure of anterior articular surface for talus
4419000 | Tracheal submucosa
4421005 | Cellular structures
4430002 | Structure of clivus ossis sphenoidalis
4432005 | Structure of ductus arteriosus
4442007 | Dental arch structure
4486002 | Structure of accessory sinus gland
4524000 | Structure of subclavian plexus
4527007 | Joint structure of lower extremity
4537002 | Structure of internal medullary lamina of thalamus
4548009 | Lamellated granule, as in surfactant-secreting cell
4549001 | Entire vagus nerve parasympathetic fibers to liver, gallbladder, bile ducts and pancreas
4566004 | Structure of tentorium cerebelli
4573009 | Desmosome
4578000 | Skin structure of posterior surface of thigh
4583008 | Structure of splenius muscle of trunk
4588004 | Structure of middle trunk of brachial plexus
4596009 | Larynx structure
4603002 | Structure of base of phalanx of foot
4606005 | Tubercle of eighth rib structure
4621004 | Structure of lesser tuberosity of humerus
4624007 | Structure of lymphatic cord
4647008 | Lipid droplet, homogeneous
4651005 | Structure of tunica albuginea of corpus spongiosum
4658004 | Skin structure of nuchal region
4677002 | Basal lamina, inclusion in subepithelial location
4703008 | Cardinal vein structure
4717004 | Neutrophilic myelocyte
4718009 | Entire venous plexus of the foramen ovale basis cranii
4743003 | Structure of ventral sacrococcygeal ligament
4755009 | Structure of medial surface of great toe
4759003 | Structure of gemellus muscle
4766002 | Structure of supracardinal vein
4768001 | Structure of perineal nerve
4774001 | Structure of phrenic nerve pericardial branch
4775000 | Structure of ventral posterior inferior nucleus
4799000 | Deiter cell
4810005 | Structure of uterine venous plexus
4812002 | Anterior tibial compartment structure
4828007 | Femoral canal structure
4840007 | Subcutaneous tissue structure of ring finger
4843009 | Ampulla of semicircular duct
4861000 | Structure of tuberculum impar

4866005 | Constrictor muscle of pharynx structure
4870002 | Structure of dorsal tegmental nuclei of midbrain
4871003 | Lamina of modiolus of cochlea
4881004 | Entire sublingual vein
4888005 | Entire interlobular vein of kidney
4897009 | Cell membrane, prokaryotic
4905007 | Structure of uterovaginal plexus
4906008 | Mastoid antrum structure
4924005 | Cerebellar gracile lobule
4942000 | Lower limb lymph node structure
4954000 | Structure of radial notch of ulna
4956003 | Subcutaneous tissue structure of back
4958002 | Amygdaloid structure
5001007 | Structure of superior temporal sulcus
5023006 | Structure of yellow bone marrow
5026003 | Structure of posterior surface of prostate
5046008 | Structure of superficial dorsal veins of clitoris
5068003 | Structure of obturator internus muscle ischial bursa
5069006 | Structure of rugal column
5076001 | Structure of infrasternal angle
5115006 | Structure of posterior auricular vein
5122003 | Entire angle of first rib
5128004 | Lens zonules
5140004 | Permanent upper right 6 tooth
5192008 | Structure of intervertebral foramen of twelfth thoracic vertebra
5194009 | Structure of epithelium of lens
5195005 | Structure of right external carotid artery
5204005 | Superior ileocecal recess
5213007 | Frontal vein
5225005 | Structure of uterine ostium of fallopian tube
5228007 | Right cerebral hemisphere structure
5229004 | Structure of mucosa of gallbladder
5261000 | Structure of thoracic intervertebral disc
5272005 | Skin structure of lateral portion of neck
5279001 | Structure of foramen singulare
5296000 | Structure of anterior mediastinal lymph node
5324007 | Structure of superior pole of kidney
5329002 | Bone structure of C4
5336001 | Structure of inferior frontal gyrus
5347008 | Synaptic specialization, cytoplasmic
5362005 | Structure of median arcuate ligament of diaphragm
5366008 | Hippocampal structure
5379004 | Small intestine muscularis propria
5382009 | Superior fascia of perineum
5394000 | Uterine paracervical lymph node
5398002 | Normal fat pad
5403001 | Articular process of third lumbar vertebra
5421003 | Sex chromosome Y
5427004 | Apocrine intraepidermal duct
5458003 | Deep artery of clitoris
5459006 | Cardiac incisure of stomach
5491007 | Lacrimal part of orbicularis oculi muscle
5493005 | Metacarpophalangeal joint of little finger
5498001 | Superior aberrant ductule of epididymis
5501001 | Hyaloid artery
5520004 | Subcutaneous tissue of chin
5538001 | Tegmental portion of pons
5542003 | Crista marginalis of tooth
5544002 | Longitudinal layer of duodenal muscularis propria
5560003 | Alveolar ridge mucous membrane
5564007 | Singlet
5574005 | Seventh costal cartilage
5580002 | Tendon of supraspinatus muscle
5597008 | Retina of right eye
5611001 | Anulus fibrosus of intervertebral disc of fifth cervical vertebra
5625000 | Navicular facet of intermediate cuneiform bone
5627008 | Right visceral pleura
5633004 | Muscular portion of interventricular septum
5643001 | Canal of stomach
5644007 | Fractured membrane P face
5653000 | Inner surface of seventh rib
5665001 | Retina

5668004 | Lower digestive tract
5677006 | Lenticular fasciculus
5682004 | Subcutaneous tissue of upper extremity
5696005 | Articular part of tubercle of ninth rib
5697001 | Skin of lateral surface of finger
5709001 | Multifidus muscles
5713008 | Submandibular triangle
5717009 | Temporal fossa
5718004 | Tendon and tendon sheath of leg and ankle
5727003 | Anterior cervical lymph node
5742000 | Skin of forearm
5751008 | Subcutaneous tissue of anterior portion of neck
5769004 | Endocervical epithelium
5780004 | Paradidymis
5798000 | Diaphragm
5802004 | Medium sized neuron
5814007 | Angle of seventh rib
5815008 | Superior rectus muscle
5816009 | Duodenal fold
5825003 | Substantia propria of sclera
5828001 | Posterior cord of brachial plexus
5847003 | Superior articular process of seventh cervical vertebra
5854009 | Orbital plate of ethmoid bone
5868002 | Serosa of urinary bladder
5872003 | Subcutaneous tissue of lateral border of sole of foot
5881009 | Tuberosity of distal phalanx of hand
5882002 | Endothelial sieve plate
5889006 | Articular surface, third metacarpal, of fourth metacarpal bone
5890002 | Posterior cells of ethmoid sinus
5893000 | Superior recess of tympanic membrane
5898009 | Myotome
5923009 | Articular process of twelfth thoracic vertebra
5926001 | Bronchial lumen
5928000 | Great cardiac vein
5942008 | Tensor tympani muscle
5943003 | Vestibular vein
5944009 | Posterior palatine arch
5948007 | Capsule of distal interphalangeal joint of third toe
5951000 | Left wrist
5953002 | Eighth rib
5976004 | Subcutaneous tissue of eyelid
5979006 | Episcleral artery
5996007 | Chromosomes, group D
6001004 | Quadratus lumborum muscle
6004007 | Intervertebral disc of second thoracic vertebra
6006009 | Circular layer of duodenal muscularis propria
6009002 | Mesentery of ascending colon
6013009 | Reticuloendothelial system
6014003 | Penicilliary arteries
6023000 | Heterolysosome
6032003 | Columnar epithelial cell
6046003 | Outer surface of third rib
6050005 | Lacrimal vein
6059006 | Metacarpophalangeal joint of middle finger
6062009 | Deciduous mandibular right canine tooth
6073002 | Ligament of left superior vena cava
6074008 | Capsule of temporomandibular joint
6076005 | Gastrointestinal subserosa
6104005 | Subclavian nerve
6105006 | Body of fifth thoracic vertebra
6110005 | Facial nerve parasympathetic fibers
6194002 | Nail of fourth toe
6216007 | Postcapillary venule
6217003 | Piriform recess
6229007 | Os lacrimale
6253001 | Sulcus terminalis cordis
6268000 | Accessory phrenic nerves
6269008 | Subcutaneous tissue of scalp
6279005 | Skin of dorsal surface of finger
6317000 | Posterior basal branch of left pulmonary artery
6325003 | Aryepiglottic muscle
6326002 | Fetal atloid articulation

6335009 | Lymphoid follicle of stomach
6359004 | Hair medulla
6371005 | Lymphatics of thyroid gland
6375001 | Cavernous portion of urethra
6392005 | Coccygeal nerve
6404004 | Ligamentum nuchae
6413002 | Presymphysial lymph node
6417001 | Medial malleolus
6423006 | Supraspinatus muscle
6424000 | Structure of radiating portion of cortical lobule of kidney
6445007 | Mast cell
6448009 | Posterior vagal trunk
6450001 | Cytotrophoblast
6472004 | Medial aspect of ovary
6504002 | Glans clitoridis
6511003 | Distal portion of circumflex branch of left coronary artery
6530003 | Cardiac valve leaflet
6533001 | Colonic haustra
6538005 | Thyrocervical trunk
6541001 | Anterior commissure of mitral valve
6544009 | Gastrohepatic ligament
6550004 | Angular incisure of stomach
6551000 | Pollicis artery
6553002 | Inferior nasal turbinate
6564004 | Medial border of sole
6566002 | Cerebellar hemisphere
6572002 | Base of phalanx of middle finger
6598008 | Lingual nerve
6606008 | Structure of dorsal intercuneiform ligaments
6608009 | Sphenoparietal sinus
6620001 | Cuticle of nail
6623004 | Sternal muscle
6633007 | Right posterior cerebral artery
6643005 | Right anterior cerebral artery
6646002 | Anterior fossa of cranial cavity
6649009 | Uterine subserosa
6651008 | Central lobule of cerebellum
6684008 | Articular facet of head of fibula
6685009 | Right ankle
6711001 | Arch of second lumbar vertebra
6720005 | Femoral nerve lateral muscular branches
6731002 | Pleural recess
6739000 | Chorda tympani
6742006 | Callosomarginal branch of anterior cerebral artery
6750002 | Mitochondrial inclusion
6757004 | Right knee
6787005 | Tendon and tendon sheath of hand
6789008 | Spermatozoa
6799003 | Macula of utricle
6805009 | Interstitial tissue of spleen
6820003 | Obturator nerve anterior branch
6828005 | Ligament of lumbosacral joint
6829002 | Pars ciliaris of retina
6834003 | Axial skeleton
6841009 | Corticomedullary junction of kidney
6844001 | Spore crystal
6850006 | Secondary foot process
6864006 | Leaf of epiglottis
6866008 | Habenular commissure
6871001 | Visceral pericardium
6894000 | Medial surface of arm
6902008 | Popliteal region
6905005 | Subcutaneous tissue of medial surface of third toe
6912001 | Lower alveolar ridge mucosa
6914000 | Perivascular space
6921000 | Right upper extremity
6930008 | Jugular arch
6944002 | Anterior labial veins
6969002 | Lymphocytic tissue
6975006 | Anterior myocardium
6981003 | Posterior hypothalamic nucleus
6987004 | Collateral sulcus

6989001 | Thoracolumbar region of back
6991009 | Subcutaneous tissue of jaw
7035006 | Bile duct mucous membrane
7050002 | Subcutaneous tissue of external genitalia
7067009 | Right colic artery
7076002 | Interstitial tissue of myocardium
7083009 | Middle phalanx of index finger
7090004 | Supraaortic branches
7091000 | Ventral posterolateral nucleus of thalamus
7099003 | Attachment plaque of desmosome or hemidesmosome
7117004 | Fetal implantation site
7121006 | Maxillary right second molar tooth
7148007 | Anulus fibrosus of intervertebral disc of thoracic vertebra
7149004 | False rib
7154008 | Trigeminal ganglion sensory root
7160008 | Base of metacarpal bone
7167006 | Paraduodenal recess
7173007 | Cauda equina
7188002 | Gustatory pore
7192009 | Isthmus tympani posticus
7227003 | Hypoglossal nerve intrinsic tongue muscle branch
7234001 | Inferior choroid vein
7242000 | Appendiceal muscularis propria
7275008 | Lymphatics of appendix and large intestine
7295002 | Muscle of perineum
7296001 | Deep inguinal ring
7311008 | Anterior surface of arm
7344002 | Lingual gyrus
7345001 | Ciliary processes
7347009 | Infratendinous olecranon bursa
7362006 | Lymphatic of head
7376007 | Left margin of uterus
7378008 | Paraventricular nucleus of thalamus
7384006 | Plantar calcaneocuboidal ligament
7404008 | Anterior semicircular duct
7435002 | Ovarian ligament
7471001 | Lateral surface of sublingual gland
7477002 | Lipid, crystalline
7480001 | Iliotibial tract
7494000 | Cerebellar lenticular nucleus
7498002 | Plantar tarsal ligaments
7507003 | Anterior ligament of head of fibula
7524009 | Vasa vasorum
7532001 | Vagus nerve parasympathetic fibers
7554004 | Deep head of flexor pollicis brevis muscle
7566005 | Mitotic cell in anaphase
7569003 | Finger
7591005 | Intervertebral disc space of eleventh thoracic vertebra
7597009 | Subcutaneous tissue of vertex
7605000 | Connexon
7610001 | Tenth thoracic vertebra
7629007 | Thalamoolivary tract
7651004 | Intervenous tubercle of right atrium
7652006 | Frenulum labii
7657000 | Femoral artery
7658005 | Subtendinous bursa of triceps brachii muscle
7697002 | Pontine portion of medial longitudinal fasciculus
7712004 | Subdural space of spinal region
7726008 | Skin of medial surface of fifth toe
7736000 | Posterior choroidal artery
7742001 | Palatine duct
7748002 | Skin appendage
7755000 | Mesovarian margin of ovary
7756004 | Lamina of third thoracic vertebra
7764005 | Striate artery
7769000 | Right foot
7783003 | Sympathetic trunk spinal nerve branch
7820009 | Lateral posterior nucleus of thalamus
7829005 | Anterior surface of manubrium
7832008 | Abdominal aorta
7835005 | Posterior margin of nasal septum
7840002 | Subcutaneous tissue of submental area

7841003 | Macrocytic normochromic erythrocyte
7844006 | Sternoclavicular joint
7851002 | Intracranial subdural space
7854005 | Mandibular canal
7872004 | Myocardium of ventricle
7874003 | Scapular region of back
7880006 | Rhopheocytotic vesicle
7884002 | Corneal corpuscle
7885001 | Rotator cuff including muscles and tendons
7892006 | Submucosa of anal canal
7896009 | Occipital angle of parietal bone
7911004 | Olivocerebellar fibers
7925003 | Proximal phalanx of third toe
7936005 | Ligament of diaphragm
7944005 | Helper cell
7954009 | Lamina propria of ethmoid sinus
7967007 | First left aortic arch
7986004 | Abdominopelvic portion of sympathetic nervous system
7991003 | Skin of glans penis
7999001 | Articulations of auditory ossicles
8001006 | Mucous membrane of tongue
8012006 | Anterior communicating artery
8017000 | Inflow tract of right ventricle
8024004 | Limitans nucleus
8039003 | Subcutaneous acromial bursa
8040001 | Superficial flexor tendon of little finger
8045006 | Membrane-coating granule, amorphous
8057002 | Lateral nuclei of globus pallidus
8059004 | Pancreatic veins
8067007 | Superficial circumflex iliac vein
8068002 | Stratum lemnisci of corpora quadrigemina
8079007 | Radial nerve
8091003 | Intervertebral disc space of twelfth thoracic vertebra
8100009 | Infundibulum of Fallopian tube
8111001 | Intranuclear crystal
8112008 | Hindgut
8119004 | Delphian lymph node
8128003 | Supraaortic valve area
8133004 | Superior anastomotic vein
8157004 | Vein of head
8158009 | Interlobar duct of pancreas
8159001 | Superior colliculus of corpora quadrigemina
8160006 | Lateral striate arteries
8161005 | Infraorbital nerve
8165001 | Superior articular process of fifth thoracic vertebra
8205005 | Wrist
8225009 | Accessory atrioventricular bundle
8242003 | Apical branch of right pulmonary artery
8251006 | Osseous portion of Eustachian tube
8264007 | Tunica interna of eyeball
8265008 | Articular surface, metacarpal, of phalanx of hand
8266009 | Small intestine serosa
8279000 | Pelvic viscus
8289001 | Below knee region
8292002 | Interlobular arteries of liver
8314003 | Mastoid fontanel of skull
8334002 | Lumbar lymph node
8356004 | Colic lymph node
8361002 | Tunica intima
8369000 | Sphincter pupillae muscle
8373002 | Jugum of sphenoid bone
8387002 | Lamina of eighth thoracic vertebra
8389004 | Birth canal
8412003 | Iliac fossa
8415001 | Renal surface of adrenal gland
8454000 | Joint of lumbar vertebra
8464009 | Ligament of sacroiliac joint and pubic symphysis
8482007 | Sinoatrial node branch of right coronary artery
8483002 | Mesial surface of tooth
8496001 | Obliquus capitis muscle
8523001 | Inferior articular process of twelfth thoracic vertebra
8546004 | Posterior intercavernous sinus

8556000 | Lipid droplet
8559007 | Juxtaintestinal lymph node
8560002 | Interclavicular ligament
8568009 | Abdominal lymph nodes
8580001 | Both feet
8595004 | Meissner's plexus
8598002 | Acoustic nerve
8600008 | Cricoid cartilage
8603005 | Adductor hallucis muscle
8604004 | Medulla oblongata fasciculus cuneatus
8608001 | Right margin of heart
8617001 | Zygomatic region of face
8623006 | Transplanted ureter
8629005 | Superior right pulmonary vein
8640002 | Choroidal branches of posterior spinal artery
8668003 | Glycogen vacuole
8671006 | All toes
8677005 | Body of right atrium
8688004 | Lateral olfactory gyrus
8695008 | Intervertebral foramen of second lumbar vertebra
8710005 | Minor sublingual ducts
8711009 | Periodontal tissues
8714001 | Subcutaneous tissue of interdigital space of hand
8752000 | Cavernous portion of internal carotid artery
8770002 | Nail of second toe
8775007 | Tendinous arch
8784007 | Intranuclear body, granular with filamentous capsule
8810002 | Corticomedullary junction of adrenal gland
8814006 | Iliac tuberosity
8815007 | Thenar and hypothenar spaces
8820007 | Pedicle of eleventh thoracic vertebra
8821006 | Peroneal artery
8827005 | Shaft of phalanx of middle finger
8839002 | Agranular endoplasmic reticulum, connection with other organelle
8845005 | Subtendinous prepatellar bursa
8850004 | Proper fasciculus
8854008 | Crista galli
8862000 | Palmar surface of middle finger
8873007 | Mandibular right second premolar tooth
8887007 | Brachiocephalic vein
8892009 | Diaphragmatic surface of lung
8894005 | Gastric cardiac gland
8897003 | Lateral glossoepiglottic fold
8907008 | Left ulnar artery
8910001 | Inferior transverse scapular ligament
8911002 | Endocardium of right ventricle
8928004 | Inguinal lymph node
8931003 | Coracoid process of scapula
8935007 | Cerebral meninges
8942007 | Trapezoid ligament
8965002 | Stratum zonale of corpora quadrigemina
8966001 | Left eye
8983005 | Joint structure of vertebral column
8988001 | Marginal part of orbicularis oris muscle
8993003 | Hepatic vein
9000002 | Cerebellar peduncle
9003000 | Left parietal lobe
9018004 | Middle colic vein
9040008 | Ascending colon
9055004 | Both forearms
9073001 | White matter of insula
9081000 | Splenic sinusoids
9086005 | Superior laryngeal vein
9089003 | Arch of foot
9108007 | Vein of the scala tympani
9127001 | Transverse folds of palate
9156001 | Embryo stage 1
9181003 | Accessory carpal bone
9185007 | Capsule of metatarsophalangeal joint of fifth toe
9186008 | Filaments of contractile apparatus
9188009 | Intervertebral disc of eighth thoracic vertebra
9208002 | Centriole

9212008 | Shaft of fifth metatarsal bone
9229006 | Rotatores lumborum muscles
9231002 | External pudendal veins
9240003 | Niemann-Pick cell
9242006 | Posterior segment of right lobe of liver
9258009 | Gravid uterus
9261005 | Tendon and tendon sheath of second toe
9262003 | Fascia of pelvis
9284003 | Corpus cavernosum of penis
9290004 | Posterior intraoccipital synchondrosis
9305001 | Labial veins
9317004 | Merkel's tactile disc
9320007 | Subtendinous iliac bursa
9321006 | Tail of epididymis
9325002 | Interdental papilla of gingiva
9332006 | Lateral ligament of temporomandibular joint
9348007 | Skin of medial surface of middle finger
9379006 | Permanent teeth
9380009 | Pecten ani
9384000 | Lumbar vein
9390001 | Lymphatics of stomach
9432007 | Plantar surface of fourth toe
9438006 | Structure of deep cervical lymphatic vessel
9454009 | Subclavian vein
9455005 | Medial cartilaginous lamina of Eustachian tube
9475001 | Amacrine cells of retina
9481009 | Afferent glomerular arteriole
9490002 | Pulmonary ligament
9498009 | Head of metacarpal bone
9502002 | Coronal depression of tooth
9512009 | Calcaneocuboidal ligament
9535007 | Pyramid of medulla oblongata
9558005 | Facet for fifth costal cartilage of sternum
9566001 | Duodenal lumen
9568000 | Subcutaneous tissue of areola
9596006 | Deep branch of ulnar nerve
9609000 | Posterior process of nasal septal cartilage
9625005 | Lanugo hair
9642004 | Left superior vena cava
9646001 | Superior transverse scapular ligament
9654004 | Gastric mucous gland
9659009 | Infraclavicular lymph nodes
9662007 | Subcutaneous tissue of lower margin of nasal septum
9668006 | Ciliary muscle
9677004 | Head of second metatarsal bone
9683001 | Melanocyte
9684007 | Posterior scrotal branches of internal pudendal artery
9708001 | Iliac fascia
9732008 | Medial supraclavicular nerves
9736006 | Right wrist
9743000 | Tendon of index finger
9758008 | Submucosa of tonsil
9770007 | Genital tubercle
9775002 | Left carotid sinus
9779008 | Distinctive shape of mitochondrial cristae
9783008 | Superficial lymphatics of thorax
9791004 | Deep venous system of lower extremity
9796009 | Skeletal muscle fiber, type IIb
9813009 | Fascia of upper extremity
9825007 | Proximal phalanx of little toe
9837009 | Perforating branches of internal thoracic artery
9840009 | Biparietal diameter of head
9841008 | Interspinalis thoracis muscles
9846003 | Right kidney
9847007 | Hilum of adrenal gland
9849005 | Fornix of lacrimal sac
9870004 | Carunculae hymenales
9875009 | Thymus
9878006 | Appendicular vein
9880000 | Thyroid tubercle
9881001 | Peripheral nerve myelinated nerve fiber
9891007 | Transverse arytenoid muscle

9898001 | Paracentral lobule
9951005 | Posterior ethmoidal nerve
9968009 | Primary foot process
9970000 | Ileocecal ostium
9976006 | Rhomboideus cervicis muscle
9994000 | Superior articular process of sixth thoracic vertebra
9999005 | Duodenal ampulla
10013000 | Lateral meniscus of knee joint
10024003 | Base of lung
10025002 | Base of phalanx of index finger
10026001 | Ventral spinocerebellar tract of pons
10036009 | Nucleus pulposus of intervertebral disc of eighth thoracic vertebra
10042008 | Intervertebral foramen of fifth thoracic vertebra
10047002 | Transplanted lung
10052007 | Male
10056005 | Ophthalmic nerve
10062000 | Levator labii superioris muscle
10119003 | Deep volar arch of radial artery
10124000 | Deep dorsal sacrococcygeal ligament
10134009 | Medial surface of third toe

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.591 Value Set http://hl7.org/fhir/ValueSet/service-pharmacy

URL: https://hl7.org/fhir/R4/valueset-service-pharmacy.html
Financial Management Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/service-pharmacy
Version: | 4.0.1
Name: | ExamplePharmacyServiceCodes
Title: | Example Pharmacy Service Codes
Definition: | This value set includes a smattering of Pharmacy Service codes.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.562 (for OID based terminology systems)
Copyright: | This is an example set.
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ExamplePharmacyServiceCodes

4.4.1.591.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/ex-pharmaservice

4.4.1.591.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 6 concepts

Expansion based on http://hl7.org/fhir/ex-pharmaservice version 4.0.1

All codes from system http://hl7.org/fhir/ex-pharmaservice

Code | Display | Definition
smokecess | Smoking cessation | Smoking cessation
flushot | Flu Shot | Flu Shot
drugcost | Drug Cost | The wholesale price of the medication.

markup | Markup | The additional cost assessed on the drug.
dispensefee | Dispense Fee | The professional fee charged for dispensing the product or service.
compoundfee | Compounding Fee | The professional fee charged for compounding the medication.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.592 Value Set http://hl7.org/fhir/ValueSet/adverse-event-category

URL: https://hl7.org/fhir/R4/valueset-adverse-event-category.html
Patient Care Work Group | Maturity Level: 0 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/adverse-event-category
Version: | 4.0.1
Name: | AdverseEventCategory
Title: | AdverseEventCategory
Definition: | Overall categorization of the event, e.g. product-related or situational.
Committee: | Patient Care Work Group
OID: | 2.16.840.1.113883.4.642.3.833 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for AdverseEventCategory

• Resource: AdverseEvent.category (CodeableConcept / Extensible)

4.4.1.592.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/adverse-event-category

4.4.1.592.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 14 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/adverse-event-category version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/adverse-event-category

Lvl | Code | Display | Definition
0 | product-problem | Product Problem | The adverse event pertains to a product problem.
0 | product-quality | Product Quality | The adverse event pertains to product quality.
0 | product-use-error | Product Use Error | The adverse event pertains to a product use error.
1 | wrong-dose | Wrong Dose | The adverse event pertains to a wrong dose.
1 | incorrect-prescribing-information | Incorrect Prescribing Information | The adverse event pertains to incorrect perscribing information.
1 | wrong-technique | Wrong Technique | The adverse event pertains to a wrong technique.
1 | wrong-route-of-administration | Wrong Route of Administration | The adverse event pertains to a wrong route of administration.
1 | wrong-rate | Wrong Rate | The adverse event pertains to a wrong rate.
1 | wrong-duration | Wrong Duration | The adverse event pertains to a wrong duration.
1 | wrong-time | Wrong Time | The adverse event pertains to a wrong time.
1 | expired-drug | Expired Drug | The adverse event pertains to an expired drug.
0 | medical-device-use-error | Medical Device Use Error | The adverse event pertains to a medical device use error.
0 | problem-different-manufacturer | Problem with Different Manufacturer of Same Medicine | The adverse event pertains to a problem with a different manufacturer of the same medication.
0 | unsafe-physical-environment | Unsafe Physical Environment | The adverse event pertains to an unsafe physical environment.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information

Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.593 Value Set http://hl7.org/fhir/ValueSet/contact-point-use

URL: https://hl7.org/fhir/R4/valueset-contact-point-use.html
FHIR Infrastructure Work Group | Maturity Level: N | Normative (from v4.0.0) | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/contact-point-use
Version: | 4.0.1
Name: | ContactPointUse
Title: | ContactPointUse
Definition: | Use of contact point.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.73 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ContactPointUse
• ConceptMap: Translation source in v3 map for ContactPointUse
• ConceptMap: Translation source in v2 map for ContactPointUse
• Resource: ContactPoint.use (code / Required)

4.4.1.593.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/contact-point-use

4.4.1.593.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 5 concepts

Expansion based on http://hl7.org/fhir/contact-point-use version 4.0.1

All codes from system http://hl7.org/fhir/contact-point-use

Code | Display | Definition
home | Home | A communication contact point at a home; attempted contacts for business purposes might intrude privacy and chances are one will contact family or other household members instead of the person one wishes to call. Typically used with urgent cases, or if no other contacts are available.
work | Work | An office contact point. First choice for business related contacts during business hours.
temp | Temp | A temporary contact point. The period can provide more detailed information.
old | Old | This contact point is no longer in use (or was never correct, but retained for records).
mobile | Mobile | A telecommunication device that moves and stays with its owner. May have characteristics of all other use codes, suitable for urgent matters, not the first choice for routine business.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.594 Value Set http://hl7.org/fhir/ValueSet/condition-cause

URL: https://hl7.org/fhir/R4/valueset-condition-cause.html
Patient Care Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/condition-cause
Version: | 4.0.1
Name: | ConditionCauseCodes
Title: | Condition Cause Codes
Definition: | Example value set for Cause of Condition codes
Committee: | Patient Care Work Group
OID: | 2.16.840.1.113883.4.642.3.965 (for OID based terminology systems)
Copyright: | This resource includes content from SNOMED Clinical Terms® (SNOMED CT®) which is copyright of the International Health Terminology Standards Development Organisation (IHTSDO). Implementers of these specifications must have the appropriate SNOMED CT Affiliate license - for more information contact http://www.snomed.org/snomed-ct/get-snomed-ct or info@snomed.org
Source Resource | XML / JSON

This value set is used in the following places:

• Extension: http://hl7.org/fhir/StructureDefinition/condition-dueTo: dueTo (CodeableConcept, Reference / Example)

• Extension: http://hl7.org/fhir/StructureDefinition/workflow-reasonCode: reasonCode (CodeableConcept / Example)

4.4.1.594.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://snomed.info/sct

4.4.1.594.2 Expansion

This expansion generated 01 Nov 2019

This value set cannot be expanded because of the way it is defined - it has an infinite number of members

Code | System | Display

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.595 Value Set http://hl7.org/fhir/ValueSet/history-status

URL: https://hl7.org/fhir/R4/valueset-history-status.html
Patient Care Work Group | Maturity Level: 2 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/history-status
Version: | 4.0.1
Name: | FamilyHistoryStatus
Title: | FamilyHistoryStatus
Definition: | A code that identifies the status of the family history record.
Committee: | Patient Care Work Group
OID: | 2.16.840.1.113883.4.642.3.267 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for FamilyHistoryStatus

• ConceptMap: Translation source in Canonical Mapping for "FamilyHistoryStatus"

• Resource: FamilyMemberHistory.status (code / Required)

4.4.1.595.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/history-status

4.4.1.595.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 4 concepts

Expansion based on http://hl7.org/fhir/history-status version 4.0.1

All codes from system http://hl7.org/fhir/history-status

Code | Display | Definition | Canonical Status
partial | Partial | Some health information is known and captured, but not complete - see notes for details. | ~partial
completed | Completed | All available related health information is captured as of the date (and possibly time) when the family member history was taken. | ~complete
entered-in-error | Entered in Error | This instance should not have been part of this patient's medical record. | ~error
health-unknown | Health Unknown | Health information for this family member is unavailable/unknown. | ~unknown

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.596 Value Set http://hl7.org/fhir/ValueSet/med-admin-perform-function

URL: https://hl7.org/fhir/R4/valueset-med-admin-perform-function.html
Pharmacy Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/med-admin-perform-function
Version: | 4.0.1
Name: | MedicationAdministration Performer Function Codes
Title: | Medication administration performer function codes
Definition: | MedicationAdministration Performer Function Codes
Committee: | Pharmacy Work Group
OID: | 2.16.840.1.113883.4.642.3.349 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for MedicationAdministration Performer Function Codes

• Resource: MedicationAdministration.performer.function (CodeableConcept / Example)

4.4.1.596.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/med-admin-perform-function

4.4.1.596.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 3 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/med-admin-perform-function version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/med-admin-perform-function

Code | Display | Definition
performer | Performer | A person, non-person living subject, organization or device that who actually and principally carries out the action
verifier | Verifier | A person who verifies the correctness and appropriateness of the service (plan, order, event, etc.) and hence takes on accountability.
witness | Witness | A person witnessing the action happening without doing anything. A witness is not necessarily aware, much less approves of anything stated in the service event. Example for a witness is students watching an operation or an advanced directive witness.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.597 Value Set http://hl7.org/fhir/ValueSet/chargeitem-status

URL: https://hl7.org/fhir/R4/valueset-chargeitem-status.html
Patient Administration Work Group | Maturity Level: 0 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/chargeitem-status
Version: | 4.0.1
Name: | ChargeItemStatus
Title: | ChargeItemStatus
Definition: | Codes identifying the lifecycle stage of a ChargeItem.
Committee: | Patient Administration Work Group
OID: | 2.16.840.1.113883.4.642.3.846 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ChargeItemStatus
• ConceptMap: Translation source in Canonical Mapping for "ChargeItemStatus"
• Resource: ChargeItem.status (code / Required)

4.4.1.597.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/chargeitem-status

4.4.1.597.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 7 concepts

Expansion based on http://hl7.org/fhir/chargeitem-status version 4.0.1

All codes from system http://hl7.org/fhir/chargeitem-status

Code | Display | Definition | Canonical Status
planned | Planned | The charge item has been entered, but the charged service is not yet complete, so it shall not be billed yet but might be used in the context of pre-authorization. | ~planned
billable | Billable | The charge item is ready for billing. | ~active
not-billable | Not billable | The charge item has been determined to be not billable (e.g. due to rules associated with the billing code). | ~declined
aborted | Aborted | The processing of the charge was aborted. | ~failed
billed | Billed | The charge item has been billed (e.g. a billing engine has generated financial transactions by applying the associated ruled for the charge item to the context of the Encounter, and placed them into Claims/Invoices. | ~complete
entered-in-error | Entered in Error | The charge item has been entered in error and should not be processed for billing. | ~error
unknown | Unknown | The authoring system does not know which of the status values currently applies for this charge item Note: This concept is not to be used for "other" - one of the listed statuses is presumed to apply, it's just not known which one. | ~unknown

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.598 Value Set http://hl7.org/fhir/ValueSet/assert-operator-codes

URL: https://hl7.org/fhir/R4/valueset-assert-operator-codes.html
FHIR Infrastructure Work Group | Maturity Level: 2 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/assert-operator-codes
Version: | 4.0.1
Name: | AssertionOperatorType
Title: | AssertionOperatorType
Definition: | The type of operator to use for assertion.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.708 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for AssertionOperatorType

• Resource: TestScript.setup.action.assert.operator (code / Required)

4.4.1.598.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/assert-operator-codes

4.4.1.598.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 11 concepts

Expansion based on http://hl7.org/fhir/assert-operator-codes version 4.0.1

All codes from system http://hl7.org/fhir/assert-operator-codes

Code | Display | Definition
equals | equals | Default value. Equals comparison.
notEquals | notEquals | Not equals comparison.
in | in | Compare value within a known set of values.
notIn | notIn | Compare value not within a known set of values.
greaterThan | greaterThan | Compare value to be greater than a known value.
lessThan | lessThan | Compare value to be less than a known value.
empty | empty | Compare value is empty.
notEmpty | notEmpty | Compare value is not empty.
contains | contains | Compare value string contains a known value.
notContains | notContains | Compare value string does not contain a known value.
eval | evaluate | Evaluate the FHIRPath expression as a boolean condition.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.599 Value Set http://hl7.org/fhir/ValueSet/provenance-agent-type

URL: https://hl7.org/fhir/R4/valueset-provenance-agent-type.html
Financial Management Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/provenance-agent-type
Version: | 4.0.1
Name: | ProvenanceParticipantType
Title: | Provenance participant type

Definition: | The type of participation a provenance participant.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.1003 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ProvenanceParticipantType
• Resource: Contract.term.action.performerType (CodeableConcept / Example)
• Profile: provenance-relevant-history: Provenance.agent.type (CodeableConcept / Extensible)

4.4.1.599.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/provenance-participant-type

4.4.1.599.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 10 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/provenance-participant-type version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/provenance-participant-type

Lvl | Code | Display | Definition
0 | enterer | Enterer | A person entering the data into the originating system
0 | performer | Performer | A person, animal, organization or device that who actually and principally carries out the activity
0 | author | Author | A party that originates the resource and therefore has responsibility for the information given in the resource and ownership of this resource
0 | verifier | Verifier | A person who verifies the correctness and appropriateness of activity
1 | legal | Legal Authenticator | The person authenticated the content and accepted legal responsibility for its content
0 | attester | Attester | A verifier who attests to the accuracy of the resource
0 | informant | Informant | A person who reported information that contributed to the resource
0 | custodian | Custodian | The entity that is accountable for maintaining a true an accurate copy of the original record
0 | assembler | Assembler | A device that operates independently of an author on custodian's algorithms for data extraction of existing information for purpose of generating a new artifact.
0 | composer | Composer | A device used by an author to record new information, which may also be used by the author to select existing information for aggregation with newly recorded information for the purpose of generating a new artifact.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.600 Value Set http://hl7.org/fhir/ValueSet/service-referral-method

URL: https://hl7.org/fhir/R4/valueset-service-referral-method.html
Patient Administration Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/service-referral-method
Version: | 4.0.1
Name: | ReferralMethod
Title: | ReferralMethod
Definition: | The methods of referral can be used when referring to a specific HealthCareService resource.
Committee: | Patient Administration Work Group
OID: | 2.16.840.1.113883.4.642.3.510 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ReferralMethod
• Resource: HealthcareService.referralMethod (CodeableConcept / Example)

4.4.1.600.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/service-referral-method

4.4.1.600.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 5 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/service-referral-method version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/service-referral-method

Code	Display	Definition
fax	Fax	Referrals may be accepted by fax.
phone	Phone	Referrals may be accepted over the phone from a practitioner.
elec	Secure Messaging	Referrals may be accepted via a secure messaging system. To determine the types of secure messaging systems supported, refer to the identifiers collection. Callers will need to understand the specific identifier system used to know that they are able to transmit messages.
semail	Secure Email	Referrals may be accepted via a secure email. To send please encrypt with the services public key.
mail	Mail	Referrals may be accepted via regular postage (or hand delivered).

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl	A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source	The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code	The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display	The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition	An explanation of the meaning of the concept
Comments	Additional notes about how to use the code

4.4.1.601 Value Set http://hl7.org/fhir/ValueSet/nutrient-code

URL: https://hl7.org/fhir/R4/valueset-nutrient-code.html

Orders and Observations Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL:	http://hl7.org/fhir/ValueSet/nutrient-code
Version:	4.0.1
Name:	NutrientModifierCodes
Title:	Nutrient Modifier Codes
Definition:	NutrientModifier : Codes for types of nutrients that are being modified such as carbohydrate or sodium. This value set includes codes from SNOMED CT where concept is-a 226355009 (Nutrients(substance)), and the concepts for Sodium, Potassium and Fluid. This is provided as a suggestive example.
Committee:	Orders and Observations Work Group
OID:	2.16.840.1.113883.4.642.3.386 (for OID based terminology systems)
Copyright:	This resource includes content from SNOMED Clinical Terms® (SNOMED CT®) which is copyright of the International Health Terminology Standards Development Organisation (IHTSDO). Implementers of these specifications must have the appropriate SNOMED CT Affiliate license - for more information contact http://www.snomed.org/snomed-ct/get-snomed-ct or info@snomed.org
Source Resource	XML / JSON

This value set is used in the following places:

• Resource: NutritionOrder.oralDiet.nutrient.modifier (CodeableConcept / Example)

4.4.1.601.1 Content Logical Definition

This value set includes codes from the following code systems:

http://snomed.info/sct

Code	Display
33463005	Fluid
39972003	Sodium
88480006	Potassium

• Include codes from http://snomed.info/sct where concept is-a 226355009 (Nutrients)

4.4.1.601.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 699 concepts

All codes from system http://snomed.info/sct

Code | Display | Definition
33463005 | Fluid
39972003 | Sodium
88480006 | Potassium
925002 | Sodium iodipamide
1025008 | ^93^Molybdenum
1057009 | Phosphate salt
1368003 | Iodine 131
1381005 | ^126^Iodine
1603001 | Cobalt isotope
2264001 | Selenium isotope
2331003 | Saccharide
2942001 | Carbon^14^ D-xylose
3161008 | 3-Methylhistidine
3212004 | Manganese sulfate
3325005 | Lipopolysaccharide
3342004 | Copper isotope
3617004 | ^79^Selenium
3829006 | Fe - Iron
3932008 | ^64^Copper
3945007 | Tin isotope
5167005 | Zinc chloride fumes
5312000 | ^65^Zinc
5405008 | ^60^Cobalt
6083003 | Hydroxylysine
6881002 | Magnesium fumes
6910009 | Sodium fluoride
6927001 | ^121^Tin
6952004 | ^121m^Tin
7156005 | Phosphorus isotope
7774008 | Neo-b-vitamin A>1<
7791007 | Soy protein
8143001 | Diosmin
8534003 | ^110^Tin
8919000 | Biotin
9396007 | ^60^Iron
9797000 | Phosphorus trichloride
9992001 | Molybdenum radioisotope
10466005 | Phosphide compound
10644005 | Black phosphorus
10767000 | Calcium phosphate dibasic
10790005 | ^53^Manganese
10796004 | Glucose-6-phosphate
11121002 | ^135^Iodine
11202003 | Manganese salt
11320009 | Sucrose
11331001 | ^56^Cobalt
11984007 | 1, Hydroxy cholecalciferol
12597001 | Sn - Tin
12689001 | Magnesium radioisotope
12930006 | Calcium phosphate dibasic dihydrate
12970004 | Inositol hexanitrate
13235001 | Vitamin B2
13342004 | Diethyl 2-chlorovinyl phosphate
13863008 | Arabinose
14125007 | Serine
14139007 | Arginine glutamate
14312008 | Vitamin L>2<
14399003 | Iodine radioisotope
14564001 | Amylopectin
14645000 | Zinc phenolsulfonate
14971004 | Ile - Isoleucine
15331006 | Glycine - amino acid
15472007 | Disaccharide
16850006 | Ribose
18039003 | Alkyl mercuric phosphate
18414002 | VITD - Cholecalciferol
19160003 | D-xylose
19427006 | ^59^Nickel
19622008 | ^48^Chromium
19839007 | Sorbitol
21075006 | ^61^Cobalt

21239003 | beta-D-fructopyranose
21559000 | ^69m^Zinc
21572004 | I123 - Iodine 123
21706000 | Tetrahydrofolic acid
22038003 | Se - Selenium
22362000 | Calcium hydrogen phosphate anhydrous
22555008 | Sorbose
22559002 | Glycoasparagine
22606007 | Vitamin K>2<
23433006 | Calciferol
23459009 | Selenium radioisotope
23692005 | Tribasic calcium phosphate
25128000 | ^72^Zinc
25183006 | Manganese chloride
25205007 | Fluorine radioisotope
25710007 | ^113^Tin
25761002 | Gln - Glutamine
26327007 | Red phosphorus
26645004 | L-Homocystine
26766001 | Starch glycerite
27054007 | ^57^Cobalt
27122000 | ^56^Nickel
27378009 | L-Tyrosine
29011006 | Manganese trioxide
29531004 | Nickel radioisotope
30094001 | Riboflavin dinucleotide
30178006 | Vitamin D
30179003 | Carbonyl fluoride
30324001 | ^132^Iodine
30589007 | Dibutyl phosphate
30820000 | Phosphorus
31731008 | Magnesium phosphate
32505007 | Phosphorus 32
32714004 | Magnesium dust
32741009 | d-Xylulose
33396006 | Ni - Nickel
33922005 | Vitamin L
34274009 | Iodine pentafluoride
34471003 | ^121^Iodine
35321007 | Fludeoxyglucose F 18
35344005 | Ribulose
35733004 | Fat soluble vitamin
35883004 | F - Fluorine
36397007 | Muramic acid
36434002 | 1-Methylhistidine
36816009 | Glucose-1-phosphate
36998008 | Glycogen
37123002 | Copper dust
37225000 | ^52^Manganese
37237003 | VITE - Tocopherol
38182007 | Galactose
38595007 | Trimethyl phosphite
38623000 | ^69^Zinc
38794009 | Molybdenum isotope
38839009 | Glycerol teichoic acid
39292004 | Iodine trichloride
39360003 | Starch
39546001 | Manganese isotope
40479008 | Fructose-1-phosphate
40924008 | Water soluble vitamin
40937006 | ^124^Iodine
41441005 | Erythrose
41464007 | ^90^Molybdenum
41503000 | Zinc compound
42146005 | I - Iodide
42180008 | Vitamin D>2<, phosphate ester
42193003 | Stannous fluoride
42730005 | Periodate salt
42841002 | Zinc oxide
43239002 | ^75^Selenium
43289005 | Dihydrofolate
43431002 | Maltose

43613009 | Phosphorous pentachloride
43706004 | VITC - Ascorbic acid
43728004 | beta-D-fructofuranose
43835003 | Zinc gelatin
43897008 | ^56^Manganese
43909000 | Thiamine triphosphate
44044007 | Calcium phosphate product
44262008 | Oxidized cellulose
44588005 | Iodine
44970006 | L-Aspartic acid
45158004 | Fluorine isotope
45962009 | Collodion
46075000 | Oligosaccharide
46257009 | Triiodomethane
47192000 | Meglumine amidotrizoate
47703008 | Lactose
48736006 | Ribitol teichoic acid
48861002 | ^99^Molybdenum
49506005 | Gluconic acid
50233008 | gamma-Ketovaleric acid
50374006 | ^120^Iodine
50479000 | Soya protein-iron complex
51224002 | Carmellose sodium
51774004 | ^123^Tin
52130006 | Quercetin
52625008 | L-Arginine
52717004 | Iodate salt
52736009 | L-Threonine
52745005 | ^51^Chromium
52860004 | Manganese radioisotope
53499005 | Flavin mononucleotide
53560004 | Zinc isotope
53646005 | Fructose-6-phosphate
53700003 | ^67^Copper
53834009 | Hemicellulose
54045000 | Glyceraldehyde
54083004 | Iodic acid
54517000 | Hydrogen selenide
54808007 | Co - Cobalt
54821000 | L-Tryptophan
55328009 | ^28^Magnesium
55495002 | Acidulated phosphate fluoride
55946005 | Pectin
56158000 | ^58m^Cobalt
56383006 | Leucocyanidin
56740002 | Triiodotyrosine
57272005 | Iodine compound
57273000 | Zinc radioisotope
57279001 | ^57^Nickel
58202007 | Fructose
58279004 | Selenium hexafluoride
58292001 | Selenium sulfide
58753009 | L-Alanine
60244003 | 3-Dehydroretinol
60260004 | Histidine
60764009 | Neovitamin A
60908004 | ^125^Tin
61025005 | Colloidal iodine
61045002 | Sedoheptulose
61244006 | Monobasic potassium phosphate
62174006 | Fructose-1-6-phosphate
62442005 | Chloriodized oil
62754006 | Sodium iodide
62763008 | Teichoic acid
63004003 | Phenylalanine
63089006 | Mannose
63330006 | Nonessential amino acid
63349005 | Zinc cyanide
63383003 | Zinc stearate
63718003 | Folic acid
63793008 | Phosphorus compound
64179000 | Zinc chloride

64538009 | Iodine heptafluoride
65054007 | ^62^Zinc
65183007 | Vitamin K
65586005 | Phosphorus oxychloride
66656000 | Vitamin K>1<
66781008 | Cobalt dust
66925006 | Copper
66945003 | Phosphite salt
66956003 | ^122^Iodine
67060008 | Vitamin D>3<, phosphate ester
67079006 | Glucose
67154005 | ^66^Nickel
67266008 | Chromium isotope
67377000 | Pangamic acid
67517005 | 25-Hydroxy ergocalciferol
67535001 | Thiamine monophosphate
67690002 | Sodium iodide I^123^
67719005 | Iodine isotope
68024001 | Cellulose acetate
68077006 | Quercetin-3-rutinoside
68334004 | Pyrophosphate salt
68580003 | ^59^Iron
68630002 | ^125^Iodine
68992005 | Cellulose
69089000 | ^52^Iron
69169004 | Vitamin K>3<
69324000 | Phosphorus pentasulfide
69750003 | ^63^Nickel
69770009 | Iron radioisotope
69783005 | Meglumine iodipamide
70203000 | Adenine
70288006 | Methionine
70319005 | Cobalt fumes
70496003 | alpha-Lipoic acid
70587000 | beta Alanine
70706009 | Troxerutin
70726008 | Selenium compound
71128006 | Molybdenum
71425003 | ^61^Copper
71544008 | Polysaccharide
72164009 | Inositol
72309009 | Vitamin P complex
72717003 | Magnesium
72835003 | ^119m^Tin
72840006 | Valine
73131003 | Carotene
73353008 | ^54^Manganese
73722009 | Dibasic potassium phosphate
73916008 | 5-Hydroxy tryptophan
74718007 | Zinc arsenate
74801000 | Sugar
75665004 | E621
75799006 | Lysine
75815006 | Zinc undecylenate
75871008 | Argininosuccinic acid
75876003 | Sodium alginate
75951003 | Iodine monobromide
77004003 | ^18^Fluorine
77404006 | Homoserine
77454000 | Mannoheptulose
77923008 | Phosphorus radioisotope
78481003 | Iofetamine I^123^ hydrochloride
78527003 | Fucose
78529000 | ^130^Iodine
78671009 | Zinc arsenite
78686003 | Copper^64^ acetate
78870008 | Cobalt radioisotope
80393001 | Diiodotyrosine
80582002 | 1,2,3-Propanetriol
80916004 | Potassium phosphate
81100008 | Pentose
82224005 | Ethyl mercury phosphate

82481002 | White phosphorus
82622003 | Vitamin A
83797003 | Leucine
83873000 | Sodium monofluorophosphate
83912003 | ^129^Iodine
84553006 | ^93m^Molybdenum
84987009 | Ethyl phosphate
85214009 | Glutamic acid
85369009 | Tin radioisotope
85668006 | Starch powder
85958004 | ^72^Selenium
86399009 | Nickel isotope
86431009 | Pantothenic acid
86739005 | Zinc
86865003 | ^65^Nickel
86953001 | Zinc salt
87116002 | 5,25-Dihydroxy cholecalciferol
87136001 | Asparagine
87371008 | Phosphoribosyl pyrophosphate
87437000 | ^73^Selenium
87445005 | Ipodate
87708000 | Vitamin
87802005 | ^33^Phosphorus
87869004 | Mn - Manganese
88166005 | Copper^64^ versenate
88236008 | Phosphine
88262004 | Phenylphosphine
88546006 | Sarcosine
88617009 | Magnesium isotope
88954003 | ^55^Iron
89272005 | ^58^Cobalt
89351005 | Potassium iodide
89702005 | Selenium salt
89867007 | Triorthocresyl phosphate
90150007 | ^117m^Tin
90544008 | Iodine monochloride
90567005 | Iron isotope
90664008 | ^127^Tin
90733003 | Metrizamide
91166002 | Monoiodotyrosine
91255004 | Copper fumes
95970003 | Trace element
96277001 | Sodium selenite
96335004 | Prednisolone sodium phosphate
102258003 | Gliadin
102658006 | 4-Pyridoxic acid
102661007 | Galactose-1-phosphate
102663005 | Triosephosphate
102747008 | Homocysteine
102750006 | Phosphoserine
103067001 | Retinaldehyde
103068006 | Provitamin A
103069003 | Vitamin B
103070002 | Pyridoxine congeners
103072005 | Ethylnicotinamide
103074006 | N^1^-methyl-3-carboxamide-6-pyridone
105824007 | Iodine and its derivatives
108946001 | Paricalcitol
109064002 | Hydrocortisone sodium phosphate
109224005 | Ethiodized oil
109240008 | Carrageenan
111039007 | Zinc gluconate
111065006 | Iodine pentoxide
111066007 | ^133^Iodine
111069000 | Copper radioisotope
111070004 | Cr - Chromium
111071000 | Chromium radioisotope
111078006 | ^55^Cobalt
112000000 | Essential amino acid
112121003 | Vitamin U
115599005 | Acid calcium phosphate
116094003 | Doxercalciferol

116119005 | Disodium phosphate anhydrous
116120004 | Dipotassium phosphate anhydrous
116123002 | Dibasic sodium phosphate anhydrous
116124008 | Monobasic sodium phosphate
116128006 | Magnesium chloride hexahydrate
116256008 | Tribasic potassium phosphate
116257004 | Monosaccharide
116258009 | Aldose
116259001 | Hexose
116260006 | Ketose
116261005 | Tetrose
116262003 | Triose
116263008 | Heptose
116264002 | Trisaccharide
116265001 | Raffinose
116266000 | Sugar phosphate
116267009 | Sugar alcohol
116268004 | Galactitol
116269007 | Maltitol
116271007 | Ribitol
116272000 | Dietary fiber
116776001 | D-alpha-tocopherol
116777005 | Beta tocopherol
116778000 | Gamma tocopherol
116934003 | Ergosterol
117634005 | Inorganic phosphorus
126076005 | Fish liver oil
126077001 | Burbot liver oil
126078006 | Percomorph liver oil
126079003 | Percoid liver oil
126080000 | Shark liver oil
126081001 | Halibut liver oil
126082008 | Cod liver oil
126224002 | Sodium folate
126225001 | Calcium pantothenate
126226000 | Dexpanthenol
126227009 | Pyridoxine hydrochloride
126228004 | Thiamine hydrochloride
126229007 | Thiamine nitrate
126230002 | Calcium ascorbate
126231003 | Sodium ascorbate
126235007 | Dihydrotachysterol
126237004 | Delta tocopherol
126238009 | Vitamin K>5<
126239001 | Menadiol sodium phosphate
129497004 | Iron sucrose
173196005 | Nicotinamide
226356005 | Protein and protein derivatives - dietary
226358006 | Fat - dietary
226369009 | Tetrasaccharide
226370005 | Stachyose
226371009 | Pentasaccharide
226372002 | Verbascose
226373007 | Amylose
226374001 | Complex carbohydrate
226375000 | Refined carbohydrate
226781005 | Dried soy milk
226793001 | Soy milk
226842001 | Soy cheese
226880000 | Soy milk yoghurt
226911007 | Soybean oil
227365007 | Tofu
227366008 | Steamed tofu
227367004 | Steamed fried tofu
227368009 | Fu-juk tofu
227537006 | Soya sauce
227948007 | Soy mince granules
227991002 | Carbohydrate food
227992009 | High carbohydrate food
227993004 | Low carbohydrate food
227994005 | Complex carbohydrate food
227995006 | Refined carbohydrate food

227996007 | Simple carbohydrate food
229941008 | Soy flour
229949005 | Low fat soy flour
255673007 | Phosphorus sesquisulfide
255954005 | Zinc omadine
256227006 | Agarose
256579008 | HA - Calcium hydroxyapatite
259333003 | Calcitriol
259336006 | 24R-Dihydroxycalcidiol
259337002 | Calcifediol
259382003 | Amylopectin glycogen
259403006 | Fuco-oligosaccharide
259406003 | Manno-oligosaccharide
259411001 | Sialooligosaccharide
259509008 | 5-deoxyadenosylcobalamin
259512006 | Mecobalamin
259651009 | Cellobiose
259655000 | Rhamnose
259656004 | Vitamin / trace element
259659006 | Vitamin B1
259660001 | Thiamine phosphate
259661002 | Cocarboxylase
259662009 | Vitamin B6
259663004 | Pyridoxal 5'-phosphate
259665006 | Vitamin D3 metabolite
259666007 | Alpha-tocopherol
261353004 | Fluoride toothpaste
273943001 | Nicotinic acid
277951002 | Dental fluoride varnish
277953004 | Dental fluoride gel
277954005 | Dental fluoride solution
284460001 | Soy-based baby milk
290131005 | Co - Cobalt- dietary
301862003 | Free glycerol
304051003 | Ionized magnesium
311719002 | Zinc sulfide
312184007 | L-Methionine
312276003 | Trace metal
372601001 | L-Carnitine
372639009 | Fluoride
386927006 | Hydroxypropyl cellulose
386932007 | Hydroxyethylcellose
386989006 | Povidone iodine
387021003 | Zinc supplement
387030006 | Choline bitartrate
387082008 | Calamine
387131008 | Methylcellulose
387206004 | Polysaccharide-iron complex
387367007 | Zinc sulfate
387415009 | Fluoride salt
391734004 | Alginic acid
395750001 | Iodixanol
395751002 | Iohexol
395752009 | Iomazenil[123I]
395753004 | Iomeprol
395754005 | Iopamidol
395756007 | Iopromide
395757003 | Iothalamate meglumine
395758008 | Iotrolan
395759000 | Ioversol
395760005 | Ioxaglate
395766004 | Calcipotriol
395769006 | Levofolinic acid
395775002 | L-Selenomethionine[75Se]
395782003 | Meglumine iotroxinate
395787009 | Metaiodobenzylguanidine[123I]
395789007 | Metaiodobenzylguanidine[131I]
395791004 | Methylcellulose-450
395792006 | Metrizoate
395815002 | Oxerutins
395830006 | Alpha tocopherol acetate
395838004 | Potassium aminobenzoate

395839007 | Potassium ascorbate
395840009 | Potassium iodate
395875009 | Sodium diatrizoate
395877001 | Sodium iopodate
395878006 | Sodium iothalamate
395879003 | Sodium iothalamate [125I]
395894004 | Tauroselcholic acid[75Se]
395925008 | Zinc sulfate monohydrate
395941009 | Chromium[51Cr] edetate
395951005 | Xylitol
395952003 | Zinc acetate
395981002 | Iothalamate
395982009 | Iothalamic acid
396003009 | Calcium iopodate
396020008 | Diatrizoate
396056006 | Ferric citrate[59Fe]
396065004 | Folinic acid
398939009 | Iodipamide
409093004 | Unbound copper
409094005 | Bound copper
409257009 | Vitamin B>12a<
409258004 | Vitamin B>12b<
409871008 | (1-->3)-beta-D-glucan
411332009 | Zinc diethyldithiocarbamate
412135003 | Icodextrin
412203006 | Hesperidin
412227008 | Iopanoic acid
412228003 | Ioxaglate meglumine
412230001 | Ioxaglate sodium
412244007 | Menadione
412250002 | Multivitamin agent
412410003 | Hypromellose
412415008 | Ipodate sodium
412512009 | Pyrophosphate sodium
412595005 | Cadexomer iodine
413483001 | alpha-Carotene
414506003 | Inositol niacinate
415543009 | Sodium glycerophosphate
415562007 | Soy bean product
417955006 | Iodophore
418346004 | Cellulose derivative
418407000 | Cellulose-derived viscosity modifier
418692007 | Iodoform agent
418929008 | Lactitol
418990001 | Tryptophan agent
419297002 | Lipiodol
419382002 | Cyanocobalamin
419432008 | Carboxymethylcellulose
419508004 | Topical selenium
419679002 | Vitamin B12 - category
420075002 | Monosodium L-aspartate
421289007 | Menadiol
421526004 | Zinc glycinate
421621000 | Retinol acetate
421881000 | Zinc citrate
422073002 | Vitamin K>4<
422160004 | Vitamin A palmitate
422934004 | ^62^Copper
423061008 | Fluorobenzothiazole F^18^
423071005 | Fluoromethane F^18^
423249007 | Monoclonal antibody I^124^
423373002 | Fluorouracil F^18^
423388007 | Fluorethyltyrosin F^18^
423432002 | ATSM Cu^64^
423725005 | Nicotinic acid and derivatives
423795001 | Fluoromisonidazole F^18^
423809007 | PTSM Cu^62^
425329004 | Sphalerite
425364008 | ^60^Copper
425702007 | Zinc sulfate heptahydrate
425867009 | Sodium pantothenate
425874004 | Hydroxyapatite

426722004 | Iodinated contrast media
427638006 | Gluconolactone
428732003 | Hexyl nicotinate
429592005 | Polydextrose
429990006 | Levoleucovorin calcium
430469009 | Vitamin B6
430632000 | Pyridoxamine
430633005 | Pyridoxine 5'-phosphate
430634004 | Pyridoxamine 5'-phosphate
430926008 | Pyridoxal
432165000 | L-methyl folic acid
432219009 | Cobalt[57Co] chloride
432884004 | Chlorotoxin [131I]
433468002 | Levoleucovorin calcium pentahydrate
438901003 | Sugar acid
442247006 | Cyclic oligosaccharide
442330008 | Manganese gluconate
442340006 | Sugammadex
442346000 | Riboflavin sodium phosphate
443148002 | Omega-5 gliadin
446531006 | Iodine^131^ methylnorcholestenol
447134003 | Iodine^123^ 15-(4-iodophenyl)-3(R,S)-methylpentadecanoic acid
449236003 | Levomefolate calcium
456992002 | Fluorocholine F^18^
456995000 | Florbetapir F^18^
456997008 | Flutemetamol F^18^
456999006 | Fluciclatide F^18^
457000009 | Fluciclovine F^18^
699182008 | Sulbutiamine
702796008 | Hidrosmin
703391005 | L-alanyl-L-glutamine
703394002 | Ammonium iodide
703395001 | Ascorbyl palmitate
703399007 | Benfotiamine
703402008 | Bisbentiamine
704384003 | Iobitridol
704430002 | Panthenol
706927001 | Free tryptophan
706929003 | Free thiamine
706930008 | Free retinol
706936002 | Free iodine
706938001 | Free inositol
706941005 | Free hydroxylysine
706942003 | Free homocystine
706943008 | Free homocysteine
706978008 | Arabitol
706980002 | Xylulose
707019004 | 1,25-Dihydroxyvitamin D
707020005 | 1,25-Dihydroxyvitamin D2
707053000 | 25-Hydroxyvitamin D
707070008 | 3-epi-25-Hydroxyvitamin D2
707071007 | 3-epi-25-Hydroxyvitamin D3
707184000 | 5-Methyltetrahydrofolate
707244004 | Dehydroascorbate
707614005 | Inorganic iodine
707661006 | Methionine sulfoxide
707677007 | N-methyl valine
708047003 | Sulfated monosaccharide
708053003 | Succinyladenosine
708259008 | Methionine + tryptophan
708264007 | Isoleucine + leucine + valine
708267000 | Isoleucine + leucine
708272009 | Glutamine + histidine
708276007 | Glutamate + glutamine + threonine
708280002 | Glutamate + glutamine
708290005 | Cystine + homocystine
708296004 | Cystine + homocysteine
708303001 | Asparagine + ornithine
708310007 | Anserine + carnosine + cysteine + histidine + ornithine
708316001 | Alanine + histidine + leucine + phenylalanine + tyrosine
708348002 | Alanine + beta alanine + sarcosine
708351009 | Alanine + ethanolamine

708354001 | Alloisoleucine + hydroxyproline + isoleucine + leucine
708356004 | Alloisoleucine + hydroxyproline + isoleucine + leucine + valine
708562005 | Xantinol nicotinate
708565007 | Phenylalanine + tyrosine
708600008 | Menadione sodium bisulfite
708658008 | Zinc amino acid chelate
708665000 | Alanine + cystine + histidine + homocysteine + leucine + phenylalanine + tyrosine
708703004 | Tocotrienol
708704005 | Threitol
708715000 | 1,5-Anhydroglucitol
708748000 | Arginine + argininosuccinate + lysine + serine + taurine
708846006 | Cyanocobalamin + folate
708850004 | N-2-hydroxyethyl valine
708864004 | Ascorbate + dehydroascorbate
709050009 | Aspartate + citrulline + glycine + homocystine + hydroxyproline
709131007 | Guanidinoacetate
709610005 | Protein-bound glucose
709844000 | N-carbamoyl-beta-alanine
710051007 | Selenomethionine
710208001 | Sodium hypophosphite
710238007 | delta Alanine
711264008 | Beta tocopherol + gamma tocopherol
712565006 | Lysine acetate
712587002 | Calcifediol + ergocalciferol
712590008 | Cysteine + homocysteine disulfide
712594004 | Histadinyl-isoleucine
712681009 | Selenious acid
712684001 | Menatetrenone
712686004 | Fursultiamine
712698009 | Ioxitalamate meglumine
712713006 | Fructooligosaccharide
712736002 | Florbetaben [18F]
712800005 | Glycyl-tyrosine
712801009 | Hesperidin complex
712808003 | Flavin adenine dinucleotide sodium
712847001 | Racemethionine
713311003 | Soybean oil refined
713332006 | Sodium dihydrogen phosphate monohydrate
713337000 | N2-L-alanyl-L-glutamine
713357001 | Methiosulfonium chloride
713365003 | Iron peptonate
713366002 | Iodine norcholestenol
713393004 | Chromic chloride
713664008 | Acylglycine
715884006 | Di-oligosaccharide
715885007 | Mono-oligosaccharide
716119001 | Eluxadoline
5811000122108 | Fluorodopa [18F]
669361000168106 | Wheat dextrin
737661000168109 | Sucroferric oxyhydroxide
999486711000087109 | Invert sugar
999487121000087103 | Xanthan gum

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.602 Value Set http://hl7.org/fhir/ValueSet/match-grade

URL: https://hl7.org/fhir/R4/valueset-match-grade.html
FHIR Infrastructure Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/match-grade
Version: | 4.0.1
Name: | MatchGrade
Title: | MatchGrade
Definition: | A Master Patient Index (MPI) assessment of whether a candidate patient record is a match or not.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.970 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for MatchGrade

4.4.1.602.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/match-grade

4.4.1.602.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 4 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/match-grade version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/match-grade

Code | Display | Definition
certain | Certain Match | This record meets the matching criteria to be automatically considered as a full match.
probable | Probable Match | This record is a close match, but not a certain match. Additional review (e.g. by a human) may be required before using this as a match.
possible | Possible Match | This record may be a matching one. Additional review (e.g. by a human) SHOULD be performed before using this as a match.
certainly-not | Certainly Not a Match | This record is known not to be a match. Note that usually non-matching records are not returned, but in some cases records previously or likely considered as a match may specifically be negated by the matching engine.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.603 Value Set http://hl7.org/fhir/ValueSet/knowledge-resource-types

URL: https://hl7.org/fhir/R4/valueset-knowledge-resource-types.html
Vocabulary Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/knowledge-resource-types
Version: | 4.0.1
Name: | KnowledgeResourceType
Title: | KnowledgeResourceType
Definition: | A list of all the knowledge resource types defined in this version of the FHIR specification.
Committee: | Vocabulary Work Group
OID: | 2.16.840.1.113883.4.642.3.1062 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for KnowledgeResourceType

4.4.1.603.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/knowledge-resource-types

4.4.1.603.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 9 concepts

Expansion based on http://hl7.org/fhir/knowledge-resource-types version 4.0.1

All codes from system http://hl7.org/fhir/knowledge-resource-types

Code | Display | Definition
ActivityDefinition | ActivityDefinition | The definition of a specific activity to be taken, independent of any particular patient or context.
CodeSystem | CodeSystem | A set of codes drawn from one or more code systems.
ConceptMap | ConceptMap | A map from one set of concepts to one or more other concepts.
Library | Library | Represents a library of quality improvement components.
Measure | Measure | A quality measure definition.
PlanDefinition | PlanDefinition | The definition of a plan for a series of actions, independent of any specific patient or context.
StructureDefinition | StructureDefinition | Structural Definition.
StructureMap | StructureMap | A Map of relationships between 2 structures that can be used to transform data.
ValueSet | ValueSet | A set of codes drawn from one or more code systems.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.604 Value Set http://hl7.org/fhir/ValueSet/icd-10

URL: https://hl7.org/fhir/R4/valueset-icd-10.html
Financial Management Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/icd-10
Version: | 4.0.1
Name: | ICD-10Codes
Title: | ICD-10 Codes
Definition: | This value set includes sample ICD-10 codes.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.531 (for OID based terminology systems)
Copyright: | This is an example set.
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: ChargeItem.reason (CodeableConcept / Example)

• Resource: ExplanationOfBenefit.diagnosis.diagnosis[x] (CodeableConcept|Reference(Condition) / Example)

• Resource: CoverageEligibilityRequest.item.diagnosis.diagnosis[x] (CodeableConcept|Reference(Condition) / Example)

• Resource: Claim.diagnosis.diagnosis[x] (CodeableConcept|Reference(Condition) / Example)

4.4.1.604.1 Content Logical Definition

This value set includes codes from the following code systems:

http://hl7.org/fhir/sid/icd-10

Code | Display
123456 | DIAG-1
123457 | DIAG-1a
987654 | DIAG-2
123987 | DIAG-3
112233 | DIAG-4
997755 | DIAG-5
321789 | DIAG-6

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information

Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.605 Value Set http://hl7.org/fhir/ValueSet/condition-category

URL: https://hl7.org/fhir/R4/valueset-condition-category.html
Patient Care Work Group | Maturity Level: 3 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/condition-category
Version: | 4.0.1
Name: | ConditionCategoryCodes
Title: | Condition Category Codes
Definition: | Preferred value set for Condition Categories.
Committee: | Patient Care Work Group
OID: | 2.16.840.1.113883.4.642.3.162 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ConditionCategoryCodes
• Resource: Condition.category (CodeableConcept / Extensible)

4.4.1.605.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/condition-category

4.4.1.605.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 2 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/condition-category version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/condition-category

Code | Display | Definition
problem-list-item | Problem List Item | An item on a problem list that can be managed over time and can be expressed by a practitioner (e.g. physician, nurse), patient, or related person.
encounter-diagnosis | Encounter Diagnosis | A point in time diagnosis (e.g. from a physician or nurse) in context of an encounter.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.606 Value Set http://hl7.org/fhir/ValueSet/animal-species

URL: https://hl7.org/fhir/R4/valueset-animal-species.html
Patient Administration Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/animal-species
Version: | 4.0.1
Name: | AnimalSpecies
Title: | Animal species
Definition: | This example value set defines a set of codes that can be used to indicate species of animal patients.
Committee: | Patient Administration Work Group

OID: | 2.16.840.1.113883.4.642.3.420 (for OID based terminology systems)
Copyright: | This resource includes content from SNOMED Clinical Terms® (SNOMED CT®) which is copyright of the International Health Terminology Standards Development Organisation (IHTSDO). Implementers of these specifications must have the appropriate SNOMED CT Affiliate license - for more information contact http://www.snomed.org/snomed-ct/get-snomed-ct or info@snomed.org
Source Resource | XML / JSON

This value set is used in the following places:

• Extension: http://hl7.org/fhir/StructureDefinition/practitioner-animalSpecies: animalSpecies (CodeableConcept / Example)

• Extension: http://hl7.org/fhir/StructureDefinition/patient-animal: animal (CodeableConcept / Example)

4.4.1.606.1 Content Logical Definition

This value set includes codes from the following code systems:

http://snomed.info/sct

Code	Display
125097000	goat
125099002	sheep
34618005	cow
425134008	turkey
47290002	chicken
15778005	goose
396620009	duck
388445009	horse
85626006	donkey
132950000	mule

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.607 Value Set http://hl7.org/fhir/ValueSet/adjudication-reason

URL: https://hl7.org/fhir/R4/valueset-adjudication-reason.html
Financial Management Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/adjudication-reason
Version: | 4.0.1
Name: | AdjudicationReasonCodes
Title: | Adjudication Reason Codes
Definition: | This value set includes smattering of Adjudication Reason codes.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.598 (for OID based terminology systems)
Copyright: | This is an example set.
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for AdjudicationReasonCodes

• Resource: ExplanationOfBenefit.item.adjudication.reason (CodeableConcept / Example)

• Resource: ClaimResponse.item.adjudication.reason (CodeableConcept / Example)

4.4.1.607.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/adjudication-reason

4.4.1.607.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 2 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/adjudication-reason version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/adjudication-reason

Code | Display | Definition
ar001 | Not covered | Not covered
ar002 | Plan Limit Reached | Plan Limit Reached

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.608 Value Set http://hl7.org/fhir/ValueSet/medicationrequest-intent

URL: https://hl7.org/fhir/R4/valueset-medicationrequest-intent.html
Pharmacy Work Group | Maturity Level: 3 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/medicationrequest-intent
Version: | 4.0.1
Name: | medicationRequest Intent
Title: | Medication request intent
Definition: | MedicationRequest Intent Codes
Committee: | Pharmacy Work Group
OID: | 2.16.840.1.113883.4.642.3.1321 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for medicationRequest Intent
• Resource: MedicationRequest.intent (code / Required)

4.4.1.608.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/CodeSystem/medicationrequest-intent

4.4.1.608.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 8 concepts

Expansion based on http://hl7.org/fhir/CodeSystem/medicationrequest-intent version 4.0.1

All codes from system http://hl7.org/fhir/CodeSystem/medicationrequest-intent

Code | Display | Definition
proposal | Proposal | The request is a suggestion made by someone/something that doesn't have an intention to ensure it occurs and without providing an authorization to act
plan | Plan | The request represents an intention to ensure something occurs without providing an authorization for others to act.
order | Order | The request represents a request/demand and authorization for action
original-order | Original Order | The request represents the original authorization for the medication request.
reflex-order | Reflex Order | The request represents an automatically generated supplemental authorization for action based on a parent authorization together with initial results of the action taken against that parent authorization..
filler-order | Filler Order | The request represents the view of an authorization instantiated by a fulfilling system representing the details of the fulfiller's intention to act upon a submitted order.
instance-order | Instance Order | The request represents an instance for the particular order, for example a medication administration record.
option | Option | The request represents a component or option for a RequestGroup that establishes timing, conditionality and/or other constraints among a set of requests.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information

Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.609 Value Set http://hl7.org/fhir/ValueSet/goal-category

URL: https://hl7.org/fhir/R4/valueset-goal-category.html
Patient Care Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined at http://www.hl7.org/Special/committees/patientcare/ .

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/goal-category
Version: | 4.0.1
Name: | GoalCategory
Title: | Goal category
Definition: | Example codes for grouping goals to use for filtering or presentation.
Committee: | Patient Care Work Group
OID: | 2.16.840.1.113883.4.642.3.275 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for GoalCategory
• Resource: Goal.category (CodeableConcept / Example)
• Resource: PlanDefinition.goal.category (CodeableConcept / Example)
• Profile: ShareablePlanDefinition: PlanDefinition.goal.category (CodeableConcept / Example)
• Profile: ComputablePlanDefinition: PlanDefinition.goal.category (CodeableConcept / Example)
• Profile: CDSHooksServicePlanDefinition: PlanDefinition.goal.category (CodeableConcept / Example)

4.4.1.609.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/goal-category

4.4.1.609.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 5 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/goal-category version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/goal-category

Code | Display | Definition
dietary | Dietary | Goals related to the consumption of food and/or beverages.
safety | Safety | Goals related to the personal protection of the subject.
behavioral | Behavioral | Goals related to the manner in which the subject acts.
nursing | Nursing | Goals related to the practice of nursing or established by nurses.
physiotherapy | Physiotherapy | Goals related to the mobility and/or motor capability of the subject.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.610 Value Set http://hl7.org/fhir/ValueSet/focal-subject

URL: https://hl7.org/fhir/R4/valueset-focal-subject.html
Orders and Observations Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/focal-subject
Version: | 4.0.1
Name: | FocalSubjectCodes
Title: | Focal Subject Codes
Definition: | Example value set composed from SNOMED CT and HL7 V3 codes for observation targets such as donor, fetus or spouse. As use cases are discovered, more values may be added.
Committee: | Orders and Observations Work Group
OID: | 2.16.840.1.113883.4.642.3.954 (for OID based terminology systems)
Copyright: | This resource includes content from SNOMED Clinical Terms® (SNOMED CT®) which is copyright of the International Health Terminology Standards Development Organisation (IHTSDO). Implementers of these specifications must have the appropriate SNOMED CT Affiliate license - for more information contact http://www.snomed.org/snomed-ct/get-snomed-ct or info@snomed.org
Source Resource | XML / JSON

This value set is used in the following places:

• Extension: http://hl7.org/fhir/StructureDefinition/observation-focusCode: focusCode (CodeableConcept / Example)

4.4.1.610.1 Content Logical Definition

This value set includes codes from the following code systems:

http://snomed.info/sct

Code | Display
83418008 | Fetus

http://terminology.hl7.org/CodeSystem/v3-ParticipationType

Code | Display
DON | donor | In some organ transplantation services and rarely in transfusion services a donor will be a target participant in the service. However, in most cases transplantation is decomposed in three services: explantation, transport, and implantation. The identity of the donor (recipient) is often irrelevant for the explantation (implantation) service.

http://terminology.hl7.org/CodeSystem/v3-RoleCode

Code | Display
SPS | spouse | The player of the role is a marriage partner of the scoping person.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.611 Value Set http://hl7.org/fhir/ValueSet/trigger-type

URL: https://hl7.org/fhir/R4/valueset-trigger-type.html
FHIR Infrastructure Work Group | Maturity Level: 5 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/trigger-type
Version: | 4.0.1
Name: | TriggerType
Title: | TriggerType
Definition: | The type of trigger.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.103 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for TriggerType

• Resource: TriggerDefinition.type (code / Required)

4.4.1.611.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/trigger-type

4.4.1.611.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 8 concepts

Expansion based on http://hl7.org/fhir/trigger-type version 4.0.1

All codes from system http://hl7.org/fhir/trigger-type

Lvl | Code | Display | Definition

0 | named-event | Named Event | The trigger occurs in response to a specific named event, and no other information about the trigger is specified. Named events are completely pre-coordinated, and the formal semantics of the trigger are not provided.

0 | periodic | Periodic | The trigger occurs at a specific time or periodically as described by a timing or schedule. A periodic event cannot have any data elements, but may have a name assigned as a shorthand for the event.

0 | data-changed | Data Changed | The trigger occurs whenever data of a particular type is changed in any way, either added, modified, or removed.

1 | data-added | Data Added | The trigger occurs whenever data of a particular type is added.

1 | data-modified | Data Updated | The trigger occurs whenever data of a particular type is modified.

1 | data-removed | Data Removed | The trigger occurs whenever data of a particular type is removed.

0 | data-accessed | Data Accessed | The trigger occurs whenever data of a particular type is accessed.

0 | data-access-ended | Data Access Ended | The trigger occurs whenever access to data of a particular type is completed.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information

Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)

Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')

Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application

Definition | An explanation of the meaning of the concept

Comments | Additional notes about how to use the code

4.4.1.612 Value Set http://hl7.org/fhir/ValueSet/event-timing

URL: https://hl7.org/fhir/R4/valueset-event-timing.html

FHIR Infrastructure Work Group | Maturity Level: N | Normative (from v4.0.0) | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/event-timing

Version: | 4.0.1

Name: | EventTiming

Title: | EventTiming

Definition: | Real world event relating to the schedule.

Committee: | FHIR Infrastructure Work Group

OID: | 2.16.840.1.113883.4.642.3.75 (for OID based terminology systems)

Source Resource | XML / JSON

This value set is used in the following places:

• Resource: Timing.repeat.when (code / Required)

4.4.1.612.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/event-timing

http://terminology.hl7.org/CodeSystem/v3-TimingEvent

Code | Display | Definition

HS | HS | event occurs [offset] before the hour of sleep (or trying to)

WAKE | WAKE | event occurs [offset] after waking

C | C | event occurs at a meal (from the Latin cibus)

CM | CM | event occurs at breakfast (from the Latin cibus matutinus)

CD | CD | event occurs at lunch (from the Latin cibus diurnus)

CV | CV | event occurs at dinner (from the Latin ante cibus vespertinus)

AC | AC | event occurs [offset] before a meal (from the Latin ante cibus)

ACM | ACM | event occurs [offset] before breakfast (from the Latin ante cibus matutinus)

ACD | ACD | event occurs [offset] before lunch (from the Latin ante cibus diurnus)

ACV | ACV | event occurs [offset] before dinner (from the Latin ante cibus vespertinus)

PC | PC | event occurs [offset] after a meal (from the Latin post cibus)

PCM | PCM | event occurs [offset] after breakfast (from the Latin post cibus matutinus)

PCD | PCD | event occurs [offset] after lunch (from the Latin post cibus diurnus)

PCV | PCV | event occurs [offset] after dinner (from the Latin post cibus vespertinus)

4.4.1.612.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 26 concepts

Expansion based on http://hl7.org/fhir/event-timing version 4.0.1,
http://terminology.hl7.org/CodeSystem/v3-TimingEvent version 2018-08-12

Code | System | Display | Definition

MORN | http://hl7.org/fhir/event-timing | Morning | Event occurs during the morning. The exact time is unspecified and established by institution convention or patient interpretation.

MORN.early | http://hl7.org/fhir/event-timing | Early Morning | Event occurs during the early morning. The exact time is unspecified and established by institution convention or patient interpretation.

MORN.late | http://hl7.org/fhir/event-timing | Late Morning | Event occurs during the late morning. The exact time is unspecified and established by institution convention or patient interpretation.

NOON | http://hl7.org/fhir/event-timing | Noon | Event occurs around 12:00pm. The exact time is unspecified and established by institution convention or patient interpretation.

AFT | http://hl7.org/fhir/event-timing | Afternoon | Event occurs during the afternoon. The exact time is unspecified and established by institution convention or patient interpretation.

AFT.early | http://hl7.org/fhir/event-timing | Early Afternoon | Event occurs during the early afternoon. The exact time is unspecified and established by institution convention or patient interpretation.

AFT.late | http://hl7.org/fhir/event-timing | Late Afternoon | Event occurs during the late afternoon. The exact time is unspecified and established by institution convention or patient interpretation.

EVE | http://hl7.org/fhir/event-timing | Evening | Event occurs during the evening. The exact time is unspecified and established by institution convention or patient interpretation.

EVE.early | http://hl7.org/fhir/event-timing | Early Evening | Event occurs during the early evening. The exact time is unspecified and established by institution convention or patient interpretation.

EVE.late | http://hl7.org/fhir/event-timing | Late Evening | Event occurs during the late evening. The exact time is unspecified and established by institution convention or patient interpretation.

NIGHT | http://hl7.org/fhir/event-timing | Night | Event occurs during the night. The exact time is unspecified and established by institution convention or patient interpretation.

PHS | http://hl7.org/fhir/event-timing | After Sleep | Event occurs [offset] after subject goes to sleep. The exact time is unspecified and established by institution convention or patient interpretation.

HS | http://terminology.hl7.org/CodeSystem/v3-TimingEvent | HS | Description: Prior to beginning a regular period of extended sleep (this would exclude naps). Note that this might occur at different times of day depending on a person's regular sleep schedule.

WAKE | http://terminology.hl7.org/CodeSystem/v3-TimingEvent | WAKE | Description: Upon waking up from a regular period of sleep, in order to start regular activities (this would exclude waking up from a nap or temporarily waking up during a period of sleep) Usage Notes: e.g. Take pulse rate on waking in management of thyrotoxicosis. Take BP on waking in management of hypertension Take basal body temperature on waking in establishing date of ovulation

C | http://terminology.hl7.org/CodeSystem/v3-TimingEvent | C | Description: meal (from lat. ante cibus)

CM | http://terminology.hl7.org/CodeSystem/v3-TimingEvent | CM | Description: breakfast (from lat. cibus matutinus)

CD | http://terminology.hl7.org/CodeSystem/v3-TimingEvent | CD | Description: lunch (from lat. cibus diurnus)

CV | http://terminology.hl7.org/CodeSystem/v3-TimingEvent | CV | Description: dinner (from lat. cibus vespertinus)

AC | http://terminology.hl7.org/CodeSystem/v3-TimingEvent | AC | before meal (from lat. ante cibus)

ACM | http://terminology.hl7.org/CodeSystem/v3-TimingEvent | ACM | before breakfast (from lat. ante cibus matutinus)

ACD | http://terminology.hl7.org/CodeSystem/v3-TimingEvent | ACD | before lunch (from lat. ante cibus diurnus)

ACV | http://terminology.hl7.org/CodeSystem/v3-TimingEvent | ACV | before dinner (from lat. ante cibus vespertinus)

PC | http://terminology.hl7.org/CodeSystem/v3-TimingEvent | PC | after meal (from lat. post cibus)

PCM | http://terminology.hl7.org/CodeSystem/v3-TimingEvent | PCM | after breakfast (from lat. post cibus matutinus)

PCD | http://terminology.hl7.org/CodeSystem/v3-TimingEvent | PCD | after lunch (from lat. post cibus diurnus)

PCV | http://terminology.hl7.org/CodeSystem/v3-TimingEvent | PCV | after dinner (from lat. post cibus vespertinus)

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information

Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)

Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')

Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application

Definition | An explanation of the meaning of the concept

Comments | Additional notes about how to use the code

4.4.1.613 Value Set http://hl7.org/fhir/ValueSet/variant-state

URL: https://hl7.org/fhir/R4/valueset-variant-state.html

Clinical Genomics Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/variant-state

Version: | 4.0.1

Name: | sequenceStatus

Title: | sequenceStatus

Definition: | Codes providing the status of the variant test result.
Committee: | Clinical Genomics Work Group
OID: | 2.16.840.1.113883.4.642.3.217 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for sequenceStatus

4.4.1.613.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/variant-state

4.4.1.613.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 3 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/variant-state version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/variant-state

Code | Display | Definition
positive | positive | the variant is detected.
negative | negative | no variant is detected.
absent | absent | result of the variant is missing.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.614 Value Set http://hl7.org/fhir/ValueSet/orientation-type

URL: https://hl7.org/fhir/R4/valueset-orientation-type.html
Clinical Genomics Work Group | Maturity Level: 1 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/orientation-type
Version: | 4.0.1
Name: | orientationType
Title: | orientationType
Definition: | Type for orientation.
Committee: | Clinical Genomics Work Group
OID: | 2.16.840.1.113883.4.642.3.987 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for orientationType

• Resource: MolecularSequence.referenceSeq.orientation (code / Required)

4.4.1.614.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/orientation-type

4.4.1.614.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 2 concepts

Expansion based on http://hl7.org/fhir/orientation-type version 4.0.1

All codes from system http://hl7.org/fhir/orientation-type

Code | Display | Definition

sense | Sense orientation of referenceSeq | Sense orientation of reference sequence.
antisense | Antisense orientation of referenceSeq | Antisense orientation of reference sequence.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.615 Value Set http://hl7.org/fhir/ValueSet/security-labels

URL: https://hl7.org/fhir/R4/valueset-security-labels.html
Vocabulary Work Group | Maturity Level: N | Normative (from v4.0.0) | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/security-labels
Version: | 4.0.1
Name: | All Security Labels
Title: | SecurityLabels
Definition: | A single value set for all security labels defined by FHIR.
Committee: | Vocabulary Work Group
OID: | 2.16.840.1.113883.4.642.3.47 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: Consent.provision.securityLabel (Coding / Extensible)
• Resource: DocumentReference.securityLabel (CodeableConcept / Extensible)
• Resource: AuditEvent.entity.securityLabel (Coding / Extensible)
• Resource: Meta.security (Coding / Extensible)

4.4.1.615.1 Content Logical Definition

This value set includes codes from the following code systems:

• Import all the codes that are contained in http://terminology.hl7.org/ValueSet/v3-ConfidentialityClassification
• Import all the codes that are contained in http://terminology.hl7.org/ValueSet/v3-InformationSensitivityPolicy
• Import all the codes that are contained in http://terminology.hl7.org/ValueSet/v3-Compartment
• Import all the codes that are contained in http://terminology.hl7.org/ValueSet/v3-SecurityIntegrityObservationValue
• Import all the codes that are contained in http://terminology.hl7.org/ValueSet/v3-SecurityControlObservationValue
• Import all the codes that are contained in http://terminology.hl7.org/ValueSet/v3-ActUSPrivacyLaw

4.4.1.615.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 200 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/v3-ActUSPrivacyLaw version 2018-08-12, http://terminology.hl7.org/ValueSet/v3-SecurityControlObservationValue version 2014-03-26, http://terminology.hl7.org/ValueSet/v3-GeneralPurposeOfUse version 2014-03-26, http://terminology.hl7.org/ValueSet/v3-SecurityPolicy version 2014-03-26, http://terminology.hl7.org/CodeSystem/v3-ActCode version 2018-08-12, http://terminology.hl7.org/ValueSet/v3-RefrainPolicy version 2014-03-26, http://terminology.hl7.org/CodeSystem/v3-Confidentiality version 2018-08-12, http://terminology.hl7.org/CodeSystem/v3-ActReason version 2018-08-12, http://terminology.hl7.org/ValueSet/v3-SecurityIntegrityObservationValue version 2014-03-26, http://terminology.hl7.org/ValueSet/v3-PurposeOfUse version 2014-03-26, http://terminology.hl7.org/ValueSet/v3-InformationSensitivityPolicy version 2014-03-26, http://terminology.hl7.org/CodeSystem/v3-ObservationValue version 2018-08-12,

http://terminology.hl7.org/ValueSet/v3-Compartment version 2014-03-26,
http://terminology.hl7.org/ValueSet/v3-ActUSPrivacyLaw version 2018-08-12,
http://terminology.hl7.org/ValueSet/v3-ObligationPolicy version 2014-03-26,
http://terminology.hl7.org/ValueSet/v3-ConfidentialityClassification version 2014-03-26

Code | System | Display | Definition

U | http://terminology.hl7.org/CodeSystem/v3-Confidentiality | unrestricted | Definition: Privacy metadata indicating that the information is not classified as sensitive. Examples: Includes publicly available information, e.g., business name, phone, email or physical address. Usage Note: This metadata indicates that the receiver has no obligation to consider additional policies when making access control decisions. Note that in some jurisdictions, personally identifiable information must be protected as confidential, so it would not be appropriate to assign a confidentiality code of "unrestricted" to that information even if it is publicly available.

L | http://terminology.hl7.org/CodeSystem/v3-Confidentiality | low | Definition: Privacy metadata indicating that the information has been de-identified, and there are mitigating circumstances that prevent re-identification, which minimize risk of harm from unauthorized disclosure. The information requires protection to maintain low sensitivity. Examples: Includes anonymized, pseudonymized, or non-personally identifiable information such as HIPAA limited data sets. Map: No clear map to ISO 13606-4 Sensitivity Level (1) Care Management: RECORD_COMPONENTs that might need to be accessed by a wide range of administrative staff to manage the subject of care's access to health services. Usage Note: This metadata indicates the receiver may have an obligation to comply with a data use agreement.

M | http://terminology.hl7.org/CodeSystem/v3-Confidentiality | moderate | Definition: Privacy metadata indicating moderately sensitive information, which presents moderate risk of harm if disclosed without authorization. Examples: Includes allergies of non-sensitive nature used inform food service; health information a patient authorizes to be used for marketing, released to a bank for a health credit card or savings account; or information in personal health record systems that are not governed under health privacy laws. Map: Partial Map to ISO 13606-4 Sensitivity Level (2) Clinical Management: Less sensitive RECORD_COMPONENTs that might need to be accessed by a wider range of personnel not all of whom are actively caring for the patient (e.g. radiology staff). Usage Note: This metadata indicates that the receiver may be obligated to comply with the receiver's terms of use or privacy policies.

N | http://terminology.hl7.org/CodeSystem/v3-Confidentiality | normal | Definition: Privacy metadata indicating that the information is typical, non-stigmatizing health information, which presents typical risk of harm if disclosed without authorization. Examples: In the US, this includes what HIPAA identifies as the minimum necessary protected health information (PHI) given a covered purpose of use (treatment, payment, or operations). Includes typical, non-stigmatizing health information disclosed in an application for health, workers compensation, disability, or life insurance. Map: Partial Map to ISO 13606-4 Sensitivity Level (3) Clinical Care: Default for normal clinical care access (i.e. most clinical staff directly caring for the patient should be able to access nearly all of the EHR). Maps to normal confidentiality for treatment information but not to ancillary care, payment and operations. Usage Note: This metadata indicates that the receiver may be obligated to comply with applicable jurisdictional privacy law or disclosure authorization.

R | http://terminology.hl7.org/CodeSystem/v3-Confidentiality | restricted | Privacy metadata indicating highly sensitive, potentially stigmatizing information, which presents a high risk to the information subject if disclosed without authorization. May be pre-empted by jurisdictional law, e.g., for public health reporting or emergency treatment. Examples: Includes information that is additionally protected such as sensitive conditions mental health, HIV, substance abuse, domestic violence, child abuse, genetic disease, and reproductive health; or sensitive demographic information such as a patient's standing as an employee or a celebrity. May be used to indicate proprietary or classified information that is not related to an individual, e.g., secret ingredients in a therapeutic substance; or the name of a manufacturer. Map: Partial Map to ISO 13606-4 Sensitivity Level (3) Clinical Care: Default for normal clinical care access (i.e. most clinical staff directly caring for the patient should be able to access nearly all of the EHR). Maps to normal confidentiality for treatment information but not to ancillary care, payment and operations.. Usage Note: This metadata indicates that the receiver may be obligated to comply with applicable, prevailing (default) jurisdictional privacy law or disclosure authorization..

V | http://terminology.hl7.org/CodeSystem/v3-Confidentiality | very restricted | . Privacy metadata indicating that the information is extremely sensitive and likely stigmatizing health information that presents a very high risk if disclosed without authorization. This information must be kept in the highest confidence. Examples: Includes information about a victim of abuse, patient requested information sensitivity, and taboo subjects relating to health status that must be discussed with the patient by an attending provider before sharing with the patient. May also include information held under â€œlegal lockâ€? or attorney-client privilege Map: This metadata indicates that the receiver may not disclose this information except as directed by the information custodian, who may be the information subject. Usage Note: This metadata indicates that the receiver may not disclose this information except as directed by the information custodian, who may be the information subject.

ETH | http://terminology.hl7.org/CodeSystem/v3-ActCode | substance abuse information sensitivity | Policy for handling alcohol or drug-abuse information, which will be afforded heightened confidentiality. Information handling protocols based on organizational policies related to alcohol or drug-abuse information that is deemed sensitive. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

GDIS | http://terminology.hl7.org/CodeSystem/v3-ActCode | genetic disease information sensitivity | Policy for handling genetic disease information, which will be afforded heightened confidentiality. Information handling protocols based on organizational policies related to genetic disease information that is deemed sensitive. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

HIV | http://terminology.hl7.org/CodeSystem/v3-ActCode | HIV/AIDS information sensitivity | Policy for handling HIV or AIDS information, which will be afforded heightened confidentiality. Information handling protocols based on organizational policies related to HIV or AIDS information that is deemed sensitive. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

MST | http://terminology.hl7.org/CodeSystem/v3-ActCode | military sexual trauma information sensitivity | Policy for handling information related to sexual assault or repeated, threatening sexual harassment that occurred while the patient was in the military, which is afforded heightened confidentiality. Access control concerns for military sexual trauma is based on the patient being subject to control by a higher ranking military perpetrator and/or censure by others within the military unit. Due to the relatively unfettered access to healthcare information by higher ranking military personnel and those who have command over the patient, there is a need to sequester this information outside of the typical controls on access to military health records. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law in addition to this more generic code.

SCA | http://terminology.hl7.org/CodeSystem/v3-ActCode | sickle cell anemia information sensitivity | Policy for handling sickle cell disease information, which is afforded heightened confidentiality. Information handling protocols are based on organizational policies related to sickle cell disease information, which is deemed sensitive. Usage Note: If there is a jurisdictional mandate, then

the Act valued with this ActCode should be associated with an Act valued with any applicable laws from the ActPrivacyLaw code system.

SDV | http://terminology.hl7.org/CodeSystem/v3-ActCode | sexual assault, abuse, or domestic violence information sensitivity | Policy for handling sexual assault, abuse, or domestic violence information, which will be afforded heightened confidentiality. Information handling protocols based on organizational policies related to sexual assault, abuse, or domestic violence information that is deemed sensitive. SDV code covers violence perpetrated by related and non-related persons. This code should be specific to physical and mental trauma caused by a related person only. The access control concerns are keeping the patient safe from the perpetrator who may have an abusive psychological control over the patient, may be stalking the patient, or may try to manipulate care givers into allowing the perpetrator to make contact with the patient. The definition needs to be clarified. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

SEX | http://terminology.hl7.org/CodeSystem/v3-ActCode | sexuality and reproductive health information sensitivity | Policy for handling sexuality and reproductive health information, which will be afforded heightened confidentiality. Information handling protocols based on organizational policies related to sexuality and reproductive health information that is deemed sensitive. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

SPI | http://terminology.hl7.org/CodeSystem/v3-ActCode | specially protected information sensitivity | Policy for handling information deemed specially protected by law or policy including substance abuse, substance use, psychiatric, mental health, behavioral health, and cognitive disorders, which is afforded heightened confidentiality. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law in addition to this more generic code.

BH | http://terminology.hl7.org/CodeSystem/v3-ActCode | behavioral health information sensitivity | Policy for handling information related to behavioral and emotional disturbances affecting social adjustment and physical health, which is afforded heightened confidentiality. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law in addition to this more generic code.

COGN | http://terminology.hl7.org/CodeSystem/v3-ActCode | cognitive disability information sensitivity | Policy for handling information related to cognitive disability disorders and conditions caused by these disorders, which are afforded heightened confidentiality. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law in addition to this more generic code. Examples may include dementia, traumatic brain injury, attention deficit, hearing and visual disability such as dyslexia and other disorders and related conditions which impair learning and self-sufficiency. However, the cognitive disabilities to which this term may apply versus other behavioral health categories varies by jurisdiction and organizational policy in part due to overlap with other behavioral health conditions. Implementers should constrain to those diagnoses applicable in the domain in which this code is used.

DVD | http://terminology.hl7.org/CodeSystem/v3-ActCode | developmental disability information sensitivity | Policy for handling information related to developmental disability disorders and conditions caused by these disorders, which is afforded heightened confidentiality. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law in addition to this more generic code. A diverse group of chronic conditions that are due to mental or physical impairments impacting activities of daily living, self-care, language acuity, learning, mobility, independent living and economic self-sufficiency. Examples may include Down syndrome and Autism spectrum. However, the developmental disabilities to which this term applies versus other behavioral health categories varies by jurisdiction and organizational policy in part due to overlap with other behavioral health conditions. Implementers should constrain to those diagnoses applicable in the domain in which this code is used.

EMOTDIS | http://terminology.hl7.org/CodeSystem/v3-ActCode | emotional disturbance information sensitivity | Policy for handling information related to emotional disturbance disorders and conditions caused by these disorders, which is afforded heightened confidentiality. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law in addition to this more generic code. Typical used to characterize behavioral and mental health issues of adolescents where the disorder may be temporarily diagnosed in order to avoid the potential and unnecessary stigmatizing diagnoses of disorder long term.

MH | http://terminology.hl7.org/CodeSystem/v3-ActCode | mental health information sensitivity | Policy for handling information related to psychological disorders, which is afforded heightened confidentiality. Mental health information may be deemed specifically sensitive and distinct from physical health, substance use disorders, and behavioral disabilities and disorders in some jurisdictions. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law in addition to this more generic code.

PSY | http://terminology.hl7.org/CodeSystem/v3-ActCode | psychiatry disorder information sensitivity | Policy for handling psychiatry psychiatric disorder information, which is afforded heightened confidentiality. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

PSYTHPN | http://terminology.hl7.org/CodeSystem/v3-ActCode | psychotherapy note information sensitivity | Policy for handling psychotherapy note information, which is afforded heightened confidentiality. Usage Note: In some jurisdiction, disclosure of psychotherapy notes requires patient consent. If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

SUD | http://terminology.hl7.org/CodeSystem/v3-ActCode | substance use disorder information sensitivity | Policy for handling information related to alcohol or drug use disorders and conditions caused by these disorders, which is afforded heightened confidentiality. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law in addition to this more generic code.

ETHUD | http://terminology.hl7.org/CodeSystem/v3-ActCode | alcohol use disorder information sensitivity | Policy for handling information related to alcohol use disorders and conditions caused by these disorders, which is afforded heightened confidentiality. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law in addition to this more generic code.

OPIOIDUD | http://terminology.hl7.org/CodeSystem/v3-ActCode | opioid use disorder information sensitivity | Policy for handling information related to opioid use disorders and conditions caused by these disorders, which is afforded heightened confidentiality. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law in addition to this more generic code.

STD | http://terminology.hl7.org/CodeSystem/v3-ActCode | sexually transmitted disease information sensitivity | Policy for handling sexually transmitted disease information, which will be afforded heightened confidentiality. Information handling protocols based on organizational policies related to sexually transmitted disease information that is deemed sensitive. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

TBOO | http://terminology.hl7.org/CodeSystem/v3-ActCode | taboo | Policy for handling information not to be initially disclosed or discussed with patient except by a physician assigned to patient in this case. Information handling protocols based on organizational policies related to sensitive patient information that must be initially discussed with the patient by an attending physician before being disclosed to the patient. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code. Open Issue: This definition conflates a rule and a characteristic, and there may be a similar issue with ts sibling codes.

VIO | http://terminology.hl7.org/CodeSystem/v3-ActCode | violence information sensitivity | Policy for handling information related to harm by violence, which is afforded heightened confidentiality. Harm by violence is perpetrated by an unrelated person. Access control concerns for information about mental or physical harm resulting from violence caused by an unrelated person may include manipulation of care givers or access to records that enable the perpetrator contact or locate the patient, but the perpetrator will likely not have established abusive psychological control over the patient. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law in addition to this more generic code.

SICKLE | http://terminology.hl7.org/CodeSystem/v3-ActCode | sickle cell | Types of sensitivity policies that apply to Acts. Act.confidentialityCode is defined in the RIM as "constraints around appropriate disclosure of information about this Act, regardless of mood." Usage Note: ActSensitivity codes are used to bind information to an Act.confidentialityCode according to local sensitivity policy so that those confidentiality codes can then govern its handling across enterprises. Internally to a policy domain, however, local policies guide the access control system on how end users in that policy domain are able to use information tagged with these sensitivity values.

DEMO | http://terminology.hl7.org/CodeSystem/v3-ActCode | all demographic information sensitivity | Policy for handling all demographic information about an information subject, which will be afforded heightened confidentiality. Policies may govern sensitivity of information related to all demographic about an information subject, the disclosure of which could impact the privacy, well-being, or safety of that subject. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

DOB | http://terminology.hl7.org/CodeSystem/v3-ActCode | date of birth information sensitivity | Policy for handling information related to an information subject's date of birth, which will be afforded heightened confidentiality.Policies may govern sensitivity of information related to an information subject's date of birth, the disclosure of which could impact the privacy, well-being, or safety of that subject. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

GENDER | http://terminology.hl7.org/CodeSystem/v3-ActCode | gender and sexual orientation information sensitivity | Policy for handling information related to an information subject's gender and sexual orientation, which will be afforded heightened confidentiality. Policies may govern sensitivity of information related to an information subject's gender and sexual orientation, the disclosure of which could impact the privacy, well-being, or safety of that subject. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

LIVARG | http://terminology.hl7.org/CodeSystem/v3-ActCode | living arrangement information sensitivity | Policy for handling information related to an information subject's living arrangement, which will be afforded heightened confidentiality. Policies may govern sensitivity of information related to an information subject's living arrangement, the disclosure of which could impact the privacy, well-being, or safety of that subject. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

MARST | http://terminology.hl7.org/CodeSystem/v3-ActCode | marital status information sensitivity | Policy for handling information related to an information subject's marital status, which will be afforded heightened confidentiality. Policies may govern sensitivity of information related to an information subject's marital status, the disclosure of which could impact the privacy, well-being, or safety of that subject. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

RACE | http://terminology.hl7.org/CodeSystem/v3-ActCode | race information sensitivity | Policy for handling information related to an information subject's race, which will be afforded heightened confidentiality. Policies may govern sensitivity of information related to an information subject's race, the disclosure of which could impact the privacy, well-being, or safety of that subject. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

REL | http://terminology.hl7.org/CodeSystem/v3-ActCode | religion information sensitivity | Policy for handling information related to an information subject's religious affiliation, which will be afforded heightened confidentiality. Policies may govern sensitivity of information related to an information subject's religion, the disclosure of which could impact the privacy, well-being, or safety of that subject. Usage Notes: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

B | http://terminology.hl7.org/CodeSystem/v3-ActCode | business information sensitivity | Policy for handling trade secrets such as financial information or intellectual property, which will be afforded heightened confidentiality. Description: Since the service class can represent knowledge structures that may be considered a trade or business secret, there is sometimes (though rarely) the need to flag those items as of business level confidentiality. Usage Notes: No patient related information may ever be of this confidentiality level. If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

EMPL | http://terminology.hl7.org/CodeSystem/v3-ActCode | employer information sensitivity | Policy for handling information related to an employer which is deemed classified to protect an employee who is the information subject, and which will be afforded heightened confidentiality. Description: Policies may govern sensitivity of information related to an employer, such as law enforcement or national security, the identity of which could impact the privacy, well-being, or safety of an information subject who is an employee. Usage Notes: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

LOCIS | http://terminology.hl7.org/CodeSystem/v3-ActCode | location information sensitivity | Policy for handling information related to the location of the information subject, which will be afforded heightened confidentiality. Description: Policies may govern sensitivity of information related to the location of the information subject, the disclosure of which could impact the privacy, well-being, or safety of that subject. Usage Notes: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

SSP | http://terminology.hl7.org/CodeSystem/v3-ActCode | sensitive service provider information sensitivity | Policy for handling information related to a provider of sensitive services, which will be afforded heightened confidentiality. Description: Policies may govern sensitivity of information related to providers who deliver sensitive healthcare services in order to protect the privacy, well-being, and safety of the provider and of patients receiving sensitive services. Usage Notes: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

ADOL | http://terminology.hl7.org/CodeSystem/v3-ActCode | adolescent information sensitivity | Policy for handling information related to an adolescent, which will be afforded heightened confidentiality per applicable organizational or jurisdictional policy. An

enterprise may have a policy that requires that adolescent patient information be provided heightened confidentiality. Information deemed sensitive typically includes health information and patient role information including patient status, demographics, next of kin, and location. Usage Note: For use within an enterprise in which an adolescent is the information subject. If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

CEL | http://terminology.hl7.org/CodeSystem/v3-ActCode | celebrity information sensitivity | Policy for handling information related to a celebrity (people of public interest (VIP), which will be afforded heightened confidentiality. Celebrities are people of public interest (VIP) about whose information an enterprise may have a policy that requires heightened confidentiality. Information deemed sensitive may include health information and patient role information including patient status, demographics, next of kin, and location. Usage Note: For use within an enterprise in which the information subject is deemed a celebrity or very important person. If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

DIA | http://terminology.hl7.org/CodeSystem/v3-ActCode | diagnosis information sensitivity | Policy for handling information related to a diagnosis, health condition or health problem, which will be afforded heightened confidentiality. Diagnostic, health condition or health problem related information may be deemed sensitive by organizational policy, and require heightened confidentiality. Usage Note: For use within an enterprise that provides heightened confidentiality to diagnostic, health condition or health problem related information deemed sensitive. If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

DRGIS | http://terminology.hl7.org/CodeSystem/v3-ActCode | drug information sensitivity | Policy for handling information related to a drug, which will be afforded heightened confidentiality. Drug information may be deemed sensitive by organizational policy, and require heightened confidentiality. Usage Note: For use within an enterprise that provides heightened confidentiality to drug information deemed sensitive. If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

EMP | http://terminology.hl7.org/CodeSystem/v3-ActCode | employee information sensitivity | Policy for handling information related to an employee, which will be afforded heightened confidentiality. When a patient is an employee, an enterprise may have a policy that requires heightened confidentiality. Information deemed sensitive typically includes health information and patient role information including patient status, demographics, next of kin, and location. Usage Note: Policy for handling information related to an employee, which will be afforded heightened confidentiality. Description: When a patient is an employee, an enterprise may have a policy that requires heightened confidentiality. Information deemed sensitive typically includes health information and patient role information including patient status, demographics, next of kin, and location.

PDS | http://terminology.hl7.org/CodeSystem/v3-ActCode | patient default information sensitivity | Policy for specially protecting information reported by or about a patient, which is deemed sensitive within the enterprise (i.e., by default regardless of whether the patient requested that the information be deemed sensitive for another reason.) For example information reported by the patient about another person, e.g., a family member, may be deemed sensitive by default. Organizational policy may allow the sensitivity tag to be cleared on patient's request. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law in addition to this more generic code. For example, VA deems employee information sensitive by default. Information about a patient who is being stalked or a victim of abuse or violence may be deemed sensitive by default per a provider organization's policies.

PHY | http://terminology.hl7.org/CodeSystem/v3-ActCode | physician requested information sensitivity | Policy for handling information about a patient, which a physician or other licensed healthcare provider deems sensitive. Once tagged by the provider, this may trigger alerts for follow up actions according to organizational policy or jurisdictional law. Usage Note: For use within an enterprise that provides heightened confidentiality to certain types of information designated by a physician as sensitive. If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code. Use cases in which this code could be used are, e.g., in systems that lack the ability to automatically detect sensitive information and must rely on manual tagging; a system that lacks an applicable sensitivity tag, or for ad hoc situations where criticality of the situation requires that the tagging be done immediately by the provider before coding or transcription of consult notes can be completed, e.g., upon detection of a patient with suicidal tendencies or potential for violence.

PRS | http://terminology.hl7.org/CodeSystem/v3-ActCode | patient requested information sensitivity | Policy for specially protecting information reported by or about a patient, which the patient deems sensitive, and the patient requests that collection, access, use, or disclosure of that information be restricted. For example, a minor patient may request that information about reproductive health not be disclosed to the patient's family or to particular providers and payers. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

COMPT | http://terminology.hl7.org/CodeSystem/v3-ActCode | compartment | This is the healthcare analog to the US Intelligence Community's concept of a Special Access Program. Compartment codes may be used in as a field value in an initiator's clearance to indicate permission to access and use an IT Resource with a security label having the same compartment value in security category label field. Map: Aligns with ISO 2382-8 definition of Compartment - "A division of data into isolated blocks with separate security controls for the purpose of reducing risk."

ACOCOMPT | http://terminology.hl7.org/CodeSystem/v3-ActCode | accountable care organization compartment | A group of health care entities, which may include health care providers, care givers, hospitals, facilities, health plans, and other health care constituents who coordinate care for reimbursement based on quality metrics for improving outcomes and lowering costs, and may be authorized to access the consumer's health information because of membership in that group. Security Compartment Labels assigned to a consumer's information use in accountable care workflows should be met or exceeded by the Security Compartment attribute claimed by a participant in a an accountable care workflow who is requesting access to that information

CTCOMPT | http://terminology.hl7.org/CodeSystem/v3-ActCode | care team compartment | Care coordination across participants in a care plan requires sharing of a healthcare consumer's information specific to that workflow. A care team member should only have access to that information while participating in that workflow or for other authorized uses. Security Compartment Labels assigned to a consumer's information use in care coordination workflows should be met or exceeded by the Security Compartment attribute claimed by a participant in a care team member workflow who is requesting access to that information

FMCOMPT | http://terminology.hl7.org/CodeSystem/v3-ActCode | financial management compartment | Financial management department members who have access to healthcare consumer information as part of a patient account, billing and claims workflows. Security Compartment Labels assigned to consumer information used in these workflows should be met or exceeded by the Security Compartment attribute claimed by a participant in a financial management workflow who is requesting access to that information.

HRCOMPT | http://terminology.hl7.org/CodeSystem/v3-ActCode | human resource compartment | A security category label field value, which indicates that access and use of an IT resource is restricted to members of human resources department or workflow.

LRCOMPT | http://terminology.hl7.org/CodeSystem/v3-ActCode | legitimate relationship compartment | Providers and care givers who have an established relationship per criteria determined by policy are considered to have an established care provision relations with a healthcare consumer, and may be authorized to access the consumer's health information because of that relationship. Providers and care givers should only have access to that information while participating in legitimate relationship workflows or for other authorized uses. Security Compartment Labels assigned to a consumer's information use in legitimate relationship workflows should be met or exceeded by the Security Compartment attribute claimed by a participant in a legitimate relationship workflow who is requesting access to that information.
PACOMPT | http://terminology.hl7.org/CodeSystem/v3-ActCode | patient administration compartment | Patient administration members who have access to healthcare consumer information as part of a patient administration workflows. Security Compartment Labels assigned to consumer information used in these workflows should be met or exceeded by the Security Compartment attribute claimed by a participant in a patient administration workflow who is requesting access to that information.
RESCOMPT | http://terminology.hl7.org/CodeSystem/v3-ActCode | research project compartment | A security category label field value, which indicates that access and use of an IT resource is restricted to members of a research project.
RMGTCOMPT | http://terminology.hl7.org/CodeSystem/v3-ActCode | records management compartment | A security category label field value, which indicates that access and use of an IT resource is restricted to members of records management department or workflow.
ABSTRED | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | abstracted | Security metadata observation values used to indicate the use of a more abstract version of the content, e.g., replacing exact value of an age or date field with a range, or remove the left digits of a credit card number or SSN.
AGGRED | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | aggregated | Security metadata observation values used to indicate the use of an algorithmic combination of actual values with the result of an aggregate function, e.g., average, sum, or count in order to limit disclosure of an IT resource (data, information object, service, or system capability) to the minimum necessary.
ANONYED | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | anonymized | Security metadata observation value conveying the alteration integrity of an IT resource (data, information object, service, or system capability) by used to indicate the mechanism by which software systems can strip portions of the resource that could allow the identification of the source of the information or the information subject. No key to relink the data is retained.
MAPPED | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | mapped | Security metadata observation value used to indicate that the IT resource semantic content has been transformed from one encoding to another. Usage Note: "MAP" code does not indicate the semantic fidelity of the transformed content. To indicate semantic fidelity for maps of HL7 to other code systems, this security alteration integrity observation may be further specified using an Act valued with Value Set: MapRelationship (2.16.840.1.113883.1.11.11052). Semantic fidelity of the mapped IT Resource may also be indicated using a SecurityIntegrityConfidenceObservation.
MASKED | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | masked | Security metadata observation value conveying the alteration integrity of an IT resource (data, information object, service, or system capability) by indicating the mechanism by which software systems can make data unintelligible (that is, as unreadable and unusable by algorithmically transforming plaintext into ciphertext) such that it can only be accessed or used by authorized users. An authorized user may be provided a key to decrypt per license or "shared secret". Usage Note: "MASKED" may be used, per applicable policy, as a flag to indicate to a user or receiver that some portion of an IT resource has been further encrypted, and may be accessed only by an authorized user or receiver to which a decryption key is provided.
PSEUDED | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | pseudonymized | Security metadata observation value conveying the alteration integrity of an IT resource (data, information object, service, or system capability), by indicating the mechanism by which software systems can strip portions of the resource that could allow the identification of the source of the information or the information subject. Custodian may retain a key to relink data necessary to reidentify the information subject. Rationale: Personal data which has been processed to make it impossible to know whose data it is. Used particularly for secondary use of health data. In some cases, it may be possible for authorized individuals to restore the identity of the individual, e.g.,for public health case management. Based on ISO/TS 25237:2008 Health informaticsâ€”Pseudonymization
REDACTED | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | redacted | Security metadata observation value used to indicate the mechanism by which software systems can filter an IT resource (data, information object, service, or system capability) to remove any portion of the resource that is not authorized to be access, used, or disclosed. Usage Note: "REDACTED" may be used, per applicable policy, as a flag to indicate to a user or receiver that some portion of an IT resource has filtered and not included in the content accessed or received.
SUBSETTED | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | subsetted | Metadata observation used to indicate that some information has been removed from the source object when the view this object contains was constructed because of configuration options when the view was created. The content may not be suitable for use as the basis of a record update Usage Note: This is not suitable to be used when information is removed for security reasons - see the code REDACTED for this use.
SYNTAC | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | syntactic transform | Security metadata observation value used to indicate that the IT resource syntax has been transformed from one syntactical representation to another. Usage Note: "SYNTAC" code does not indicate the syntactical correctness of the syntactically transformed IT resource.
TRSLT | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | translated | Security metadata observation value used to indicate that the IT resource has been translated from one human language to another. Usage Note: "TRSLT" does not indicate the fidelity of the translation or the languages translated. The fidelity of the IT Resource translation may be indicated using a SecurityIntegrityConfidenceObservation. To indicate languages, use the Value Set:HumanLanguage (2.16.840.1.113883.1.11.11526)
VERSIONED | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | versioned | Security metadata observation value conveying the alteration integrity of an IT resource (data, information object, service, or system capability) which indicates that the resource only retains versions of an IT resource for access and use per applicable policy Usage Note: When this code is used, expectation is that the system has removed historical versions of the data that falls outside the time period deemed to be the effective time of the applicable version.
CRYTOHASH | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | cryptographic hash function | Security metadata observation value used to indicate the mechanism by which software systems can establish that data was not modified in transit. Rationale: This definition is intended to align with the ISO 22600-2 3.3.19 definition of cryptographic checkvalue: Information which is derived by performing a cryptographic transformation (see cryptography) on the data unit. The derivation of the checkvalue may be performed in one or more steps and is a result of a mathematical function of the key and a data unit. It is usually used to check the integrity of a data unit. Examples: SHA-1 SHA-2 (Secure Hash Algorithm)
DIGSIG | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | digital signature | Security metadata observation value used to indicate the mechanism by which software systems use digital signature to establish that data has not been modified. Rationale: This definition is intended to align with the ISO 22600-2 3.3.26 definition of digital signature: Data appended to, or a

cryptographic transformation (see cryptography) of, a data unit that allows a recipient of the data unit to prove the source and integrity of the data unit and protect against forgery e.g., by the recipient.

HRELIABLE | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | highly reliable | Security metadata observation value used to indicate that the veracity or trustworthiness of an IT resource (data, information object, service, or system capability) for a specified purpose of use is perceived to be or deemed by policy to be very high.

RELIABLE | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | reliable | Security metadata observation value used to indicate that the veracity or trustworthiness of an IT resource (data, information object, service, or system capability) for a specified purpose of use is perceived to be or deemed by policy to be adequate.

UNCERTREL | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | uncertain reliability | Security metadata observation value used to indicate that the veracity or trustworthiness of an IT resource (data, information object, service, or system capability) for a specified purpose of use is perceived to be or deemed by policy to be uncertain.

UNRELIABLE | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | unreliable | Security metadata observation value used to indicate that the veracity or trustworthiness of an IT resource (data, information object, service, or system capability) for a specified purpose of use is perceived to be or deemed by policy to be inadequate.

CLINAST | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | clinician asserted | Security provenance metadata observation value used to indicate that an IT resource (data, information object, service, or system capability) was asserted by a clinician.

DEVAST | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | device asserted | Security provenance metadata observation value used to indicate that an IT resource (data, information object, service, or system capability) was asserted by a device.

HCPAST | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | healthcare professional asserted | Security provenance metadata observation value used to indicate that an IT resource (data, information object, service, or system capability) was asserted by a healthcare professional.

PACQAST | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | patient acquaintance asserted | Security provenance metadata observation value used to indicate that an IT resource (data, information object, service, or system capability) was asserted by a patient acquaintance.

PATAST | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | patient asserted | Security provenance metadata observation value used to indicate that an IT resource (data, information object, service, or system capability) was asserted by a patient.

PAYAST | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | payer asserted | Security provenance metadata observation value used to indicate that an IT resource (data, information object, service, or system capability) was asserted by a payer.

PROAST | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | professional asserted | Security provenance metadata observation value used to indicate that an IT resource (data, information object, service, or system capability) was asserted by a professional.

SDMAST | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | substitute decision maker asserted | Security provenance metadata observation value used to indicate that an IT resource (data, information object, service, or system capability) was asserted by a substitute decision maker.

CLINRPT | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | clinician reported | Security provenance metadata observation value used to indicate that an IT resource (data, information object, service, or system capability) was reported by a clinician.

DEVRPT | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | device reported | Security provenance metadata observation value used to indicate that an IT resource (data, information object, service, or system capability) was reported by a device.

HCPRPT | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | healthcare professional reported | Security provenance metadata observation value used to indicate that an IT resource (data, information object, service, or system capability) was reported by a healthcare professional.

PACQRPT | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | patient acquaintance reported | Security provenance metadata observation value used to indicate that an IT resource (data, information object, service, or system capability) was reported by a patient acquaintance.

PATRPT | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | patient reported | Security provenance metadata observation value used to indicate that an IT resource (data, information object, service, or system capability) was reported by a patient.

PAYRPT | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | payer reported | Security provenance metadata observation value used to indicate that an IT resource (data, information object, service, or system capability) was reported by a payer.

PRORPT | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | professional reported | Security provenance metadata observation value used to indicate that an IT resource (data, information object, service, or system capability) was reported by a professional.

SDMRPT | http://terminology.hl7.org/CodeSystem/v3-ObservationValue | substitute decision maker reported | Security provenance metadata observation value used to indicate that an IT resource (data, information object, service, or system capability) was reported by a substitute decision maker.

SecurityPolicy | http://terminology.hl7.org/CodeSystem/v3-ActCode | security policy | Types of security policies that further specify the ActClassPolicy value set. Examples: obligation to encrypt refrain from redisclosure without consent

AUTHPOL | http://terminology.hl7.org/CodeSystem/v3-ActCode | authorization policy | Authorisation policies are essentially security policies related to access-control and specify what activities a subject is permitted or forbidden to do, to a set of target objects. They are designed to protect target objects so are interpreted by access control agents or the run-time systems at the target system. A positive authorisation policy defines the actions that a subject is permitted to perform on a target. A negative authorisation policy specifies the actions that a subject is forbidden to perform on a target. Positive authorisation policies may also include filters to transform the parameters associated with their actions. (Based on PONDERS)

ACCESSCONSCHEME | http://terminology.hl7.org/CodeSystem/v3-ActCode | access control scheme | An access control policy specific to the type of access control scheme, which is used to enforce one or more authorization policies. Usage Note: Access control schemes are the type of access control policy, which is comprised of access control policy rules concerning the provision of the access control service. There are two categories of access control policies, rule-based and identity-based, which are identified in CCITT Rec. X.800 aka ISO 7498-2. Rule-based access control policies are intended to apply to all access requests by any initiator on any target in a security domain. Identity-based access control policies are based on rules specific to an individual initiator, a group of initiators, entities acting on behalf of initiators, or originators acting in a specific role. Context can modify rule-based or

identity-based access control policies. Context rules may define the entire policy in effect. Real systems will usually employ a combination of these policy types; if a rule-based policy is used, then an identity-based policy is usually in effect also. An access control scheme may be based on access control lists, capabilities, labels, and context or a combination of these. An access control scheme is a component of an access control mechanism or "service") along with the supporting mechanisms required by that scheme to provide access control decision information (ADI) supplied by the scheme to the access decision facility (ADF also known as a PDP). (Based on ISO/IEC 10181-3:1996) Examples: Attribute Based Access Control (ABAC) Discretionary Access Control (DAC) History Based Access Control (HBAC) Identity Based Access Control (IBAC) Mandatory Access Control (MAC) Organization Based Access Control (OrBAC) Relationship Based Access Control (RelBac) Responsibility Based Access Control (RespBAC) Risk Adaptable Access Control (RAdAC) >

DELEPOL | http://terminology.hl7.org/CodeSystem/v3-ActCode | delegation policy | Delegation policies specify which actions subjects are allowed to delegate to others. A delegation policy thus specifies an authorisation to delegate. Subjects must already possess the access rights to be delegated. Delegation policies are aimed at subjects delegating rights to servers or third parties to perform actions on their behalf and are not meant to be the means by which security administrators would assign rights to subjects. A negative delegation policy identifies what delegations are forbidden. A Delegation policy specifies the authorisation policy from which delegated rights are derived, the grantors, which are the entities which can delegate these access rights, and the grantees, which are the entities to which the access rights can be delegated. There are two types of delegation policy, positive and negative. (Based on PONDERS)

ObligationPolicy | http://terminology.hl7.org/CodeSystem/v3-ActCode | obligation policy | Conveys the mandated workflow action that an information custodian, receiver, or user must perform. Usage Notes: Per ISO 22600-2, ObligationPolicy instances 'are event-triggered and define actions to be performed by manager agent'. Per HL7 Composite Security and Privacy Domain Analysis Model: This value set refers to the action required to receive the permission specified in the privacy rule. Per OASIS XACML, an obligation is an operation specified in a policy or policy that is performed in conjunction with the enforcement of an access control decision.

ANONY | http://terminology.hl7.org/CodeSystem/v3-ActCode | anonymize | Custodian system must remove any information that could result in identifying the information subject.

AOD | http://terminology.hl7.org/CodeSystem/v3-ActCode | accounting of disclosure | Custodian system must make available to an information subject upon request an accounting of certain disclosures of the individualâ€™s protected health information over a period of time. Policy may dictate that the accounting include information about the information disclosed, the date of disclosure, the identification of the receiver, the purpose of the disclosure, the time in which the disclosing entity must provide a response and the time period for which accountings of disclosure can be requested.

AUDIT | http://terminology.hl7.org/CodeSystem/v3-ActCode | audit | Custodian system must monitor systems to ensure that all users are authorized to operate on information objects.

AUDTR | http://terminology.hl7.org/CodeSystem/v3-ActCode | audit trail | Custodian system must monitor and maintain retrievable log for each user and operation on information.

CPLYCC | http://terminology.hl7.org/CodeSystem/v3-ActCode | comply with confidentiality code | Custodian security system must retrieve, evaluate, and comply with the information handling directions of the Confidentiality Code associated with an information target.

CPLYCD | http://terminology.hl7.org/CodeSystem/v3-ActCode | comply with consent directive | Custodian security system must retrieve, evaluate, and comply with applicable information subject consent directives.

CPLYJPP | http://terminology.hl7.org/CodeSystem/v3-ActCode | comply with jurisdictional privacy policy | Custodian security system must retrieve, evaluate, and comply with applicable jurisdictional privacy policies associated with the target information.

CPLYOPP | http://terminology.hl7.org/CodeSystem/v3-ActCode | comply with organizational privacy policy | Custodian security system must retrieve, evaluate, and comply with applicable organizational privacy policies associated with the target information.

CPLYOSP | http://terminology.hl7.org/CodeSystem/v3-ActCode | comply with organizational security policy | Custodian security system must retrieve, evaluate, and comply with the organizational security policies associated with the target information.

CPLYPOL | http://terminology.hl7.org/CodeSystem/v3-ActCode | comply with policy | Custodian security system must retrieve, evaluate, and comply with applicable policies associated with the target information.

DECLASSIFYLABEL | http://terminology.hl7.org/CodeSystem/v3-ActCode | declassify security label | Custodian security system must declassify information assigned security labels by instantiating a new version of the classified information so as to break the binding of the classifying security label when assigning a new security label that marks the information as unclassified in accordance with applicable jurisdictional privacy policies associated with the target information. The system must retain an immutable record of the previous assignment and binding.

DEID | http://terminology.hl7.org/CodeSystem/v3-ActCode | deidentify | Custodian system must strip information of data that would allow the identification of the source of the information or the information subject.

DELAU | http://terminology.hl7.org/CodeSystem/v3-ActCode | delete after use | Custodian system must remove target information from access after use.

DOWNGRDLABEL | http://terminology.hl7.org/CodeSystem/v3-ActCode | downgrade security label | Custodian security system must downgrade information assigned security labels by instantiating a new version of the classified information so as to break the binding of the classifying security label when assigning a new security label that marks the information as classified at a less protected level in accordance with applicable jurisdictional privacy policies associated with the target information. The system must retain an immutable record of the previous assignment and binding.

DRIVLABEL | http://terminology.hl7.org/CodeSystem/v3-ActCode | derive security label | Custodian security system must assign and bind security labels derived from compilations of information by aggregation or disaggregation in order to classify information compiled in the information systems under its control for collection, access, use and disclosure in accordance with applicable jurisdictional privacy policies associated with the target information. The system must retain an immutable record of the previous assignment and binding.

ENCRYPT | http://terminology.hl7.org/CodeSystem/v3-ActCode | encrypt | Custodian system must render information unreadable by algorithmically transforming plaintext into ciphertext. Usage Notes: A mathematical transposition of a file or data stream so that it cannot be deciphered at the receiving end without the proper key. Encryption is a security feature that assures that only the parties who are supposed to be participating in a videoconference or data transfer are able to do so. It can include a password, public and private keys, or a complex combination of all. (Per Infoway.)

ENCRYPTR | http://terminology.hl7.org/CodeSystem/v3-ActCode | encrypt at rest | Custodian system must render information unreadable and unusable by algorithmically transforming plaintext into ciphertext when "at rest" or in storage.

ENCRYPTT | http://terminology.hl7.org/CodeSystem/v3-ActCode | encrypt in transit | Custodian system must render information unreadable and unusable by algorithmically transforming plaintext into ciphertext while "in transit" or being transported by any means.

ENCRYPTU | http://terminology.hl7.org/CodeSystem/v3-ActCode | encrypt in use | Custodian system must render information unreadable and unusable by algorithmically transforming plaintext into ciphertext while in use such that operations permitted on the target information are limited by the license granted to the end user.

HUAPRV | http://terminology.hl7.org/CodeSystem/v3-ActCode | human approval | Custodian system must require human review and approval for permission requested.

LABEL | http://terminology.hl7.org/CodeSystem/v3-ActCode | assign security label | Custodian security system must assign and bind security labels in order to classify information created in the information systems under its control for collection, access, use and disclosure in accordance with applicable jurisdictional privacy policies associated with the target information. The system must retain an immutable record of the assignment and binding. Usage Note: In security systems, security policy label assignments do not change, they may supersede prior assignments, and such reassignments are always tracked for auditing and other purposes.

MASK | http://terminology.hl7.org/CodeSystem/v3-ActCode | mask | Custodian system must render information unreadable and unusable by algorithmically transforming plaintext into ciphertext. User may be provided a key to decrypt per license or "shared secret".

MINEC | http://terminology.hl7.org/CodeSystem/v3-ActCode | minimum necessary | Custodian must limit access and disclosure to the minimum information required to support an authorized user's purpose of use. Usage Note: Limiting the information available for access and disclosure to that an authorized user or receiver "needs to know" in order to perform permitted workflow or purpose of use.

PERSISTLABEL | http://terminology.hl7.org/CodeSystem/v3-ActCode | persist security label | Custodian security system must persist the binding of security labels to classify information received or imported by information systems under its control for collection, access, use and disclosure in accordance with applicable jurisdictional privacy policies associated with the target information. The system must retain an immutable record of the assignment and binding.

PRIVMARK | http://terminology.hl7.org/CodeSystem/v3-ActCode | privacy mark | Custodian must create and/or maintain human readable security label tags as required by policy. Map: Aligns with ISO 22600-3 Section A.3.4.3 description of privacy mark: "If present, the privacy-mark is not used for access control. The content of the privacy-mark may be defined by the security policy in force (identified by the security-policy-identifier) which may define a list of values to be used. Alternately, the value may be determined by the originator of the security-label."

PSEUD | http://terminology.hl7.org/CodeSystem/v3-ActCode | pseudonymize | Custodian system must strip information of data that would allow the identification of the source of the information or the information subject. Custodian may retain a key to relink data necessary to reidentify the information subject.

REDACT | http://terminology.hl7.org/CodeSystem/v3-ActCode | redact | Custodian system must remove information, which is not authorized to be access, used, or disclosed from records made available to otherwise authorized users.

UPGRDLABEL | http://terminology.hl7.org/CodeSystem/v3-ActCode | upgrade security label | Custodian security system must declassify information assigned security labels by instantiating a new version of the classified information so as to break the binding of the classifying security label when assigning a new security label that marks the information as classified at a more protected level in accordance with applicable jurisdictional privacy policies associated with the target information. The system must retain an immutable record of the previous assignment and binding.

RefrainPolicy | http://terminology.hl7.org/CodeSystem/v3-ActCode | refrain policy | Conveys prohibited actions which an information custodian, receiver, or user is not permitted to perform unless otherwise authorized or permitted under specified circumstances. Usage Notes: ISO 22600-2 species that a Refrain Policy "defines actions the subjects must refrain from performing". Per HL7 Composite Security and Privacy Domain Analysis Model: May be used to indicate that a specific action is prohibited based on specific access control attributes e.g., purpose of use, information type, user role, etc.

NOAUTH | http://terminology.hl7.org/CodeSystem/v3-ActCode | no disclosure without subject authorization | Prohibition on disclosure without information subject's authorization.

NOCOLLECT | http://terminology.hl7.org/CodeSystem/v3-ActCode | no collection | Prohibition on collection or storage of the information.

NODSCLCD | http://terminology.hl7.org/CodeSystem/v3-ActCode | no disclosure without consent directive | Prohibition on disclosure without organizational approved patient restriction.

NODSCLCDS | http://terminology.hl7.org/CodeSystem/v3-ActCode | no disclosure without information subject's consent directive | Prohibition on disclosure without a consent directive from the information subject.

NOINTEGRATE | http://terminology.hl7.org/CodeSystem/v3-ActCode | no integration | Prohibition on Integration into other records.

NOLIST | http://terminology.hl7.org/CodeSystem/v3-ActCode | no unlisted entity disclosure | Prohibition on disclosure except to entities on specific access list.

NOMOU | http://terminology.hl7.org/CodeSystem/v3-ActCode | no disclosure without MOU | Prohibition on disclosure without an interagency service agreement or memorandum of understanding (MOU).

NOORGPOL | http://terminology.hl7.org/CodeSystem/v3-ActCode | no disclosure without organizational authorization | Prohibition on disclosure without organizational authorization.

NOPAT | http://terminology.hl7.org/CodeSystem/v3-ActCode | no disclosure to patient, family or caregivers without attending provider's authorization | Prohibition on disclosing information to patient, family or caregivers without attending provider's authorization. Usage Note: The information may be labeled with the ActInformationSensitivity TBOO code, triggering application of this RefrainPolicy code as a handling caveat controlling access. Maps to FHIR NOPAT: Typically, this is used on an Alert resource, when the alert records information on patient abuse or non-compliance. FHIR print name is "keep information from patient". Maps to the French realm - code: INVISIBLE_PATIENT. displayName: Document non visible par le patient codingScheme: 1.2.250.1.213.1.1.4.13 French use case: A label for documents that the author chose to hide from the patient until the content can be disclose to the patient in a face to face meeting between a healthcare professional and the patient (in French law some results like cancer diagnosis or AIDS diagnosis must be announced to the patient by a healthcare professional and should not be find out by the patient alone).

NOPERSISTP | http://terminology.hl7.org/CodeSystem/v3-ActCode | no collection beyond purpose of use | Prohibition on collection of the information beyond time necessary to accomplish authorized purpose of use is prohibited.

NORDSCLCD | http://terminology.hl7.org/CodeSystem/v3-ActCode | no redisclosure without consent directive | Prohibition on redisclosure without patient consent directive.

NORDSCLCDS | http://terminology.hl7.org/CodeSystem/v3-ActCode | no redisclosure without information subject's consent directive | Prohibition on redisclosure without a consent directive from the information subject.

NORDSCLW | http://terminology.hl7.org/CodeSystem/v3-ActCode | no disclosure without jurisdictional authorization | Prohibition on disclosure without authorization under jurisdictional law.

NORELINK | http://terminology.hl7.org/CodeSystem/v3-ActCode | no relinking | Prohibition on associating de-identified or pseudonymized information with other information in a manner that could or does result in disclosing information intended to be

masked.
NOREUSE | http://terminology.hl7.org/CodeSystem/v3-ActCode | no reuse beyond purpose of use | Prohibition on use of the information beyond the purpose of use initially authorized.
NOVIP | http://terminology.hl7.org/CodeSystem/v3-ActCode | no unauthorized VIP disclosure | Prohibition on disclosure except to principals with access permission to specific VIP information.
ORCON | http://terminology.hl7.org/CodeSystem/v3-ActCode | no disclosure without originator authorization | Prohibition on disclosure except as permitted by the information originator.
PurposeOfUse | http://terminology.hl7.org/CodeSystem/v3-ActReason | purpose of use | Reason for performing one or more operations on information, which may be permitted by source system's security policy in accordance with one or more privacy policies and consent directives. Usage Notes: The rationale or purpose for an act relating to the management of personal health information, such as collecting personal health information for research or public health purposes.
HMARKT | http://terminology.hl7.org/CodeSystem/v3-ActReason | healthcare marketing | To perform one or more operations on information for marketing services and products related to health care.
HOPERAT | http://terminology.hl7.org/CodeSystem/v3-ActReason | healthcare operations | To perform one or more operations on information used for conducting administrative and contractual activities related to the provision of health care.
CAREMGT | http://terminology.hl7.org/CodeSystem/v3-ActReason | care management | To perform analytics, evaluation and other secondary uses of treatment and healthcare related information to manage the quality, efficacy, patient safety, population health, and cost effectiveness of healthcare delivery. Explicitly excludes the use of information to organize the delivery of health care for care coordination and case management, or to provide healthcare treatment. Usage Note: The concept of care management is narrower than the list of activities related to more general organizational objectives such as provider profiling, education of healthcare and non-healthcare professionals; insurance underwriting, premium rating, reinsurance; organizational legal, medical review, auditing, compliance and fraud and abuse detection; business planning, development, and restructuring; fund-raising; and customer service. Map: Maps to ISO 14265 Classification Term "Health service management and quality assurance" described as "To inform persons or processes responsible for determining the availability, quality, safety, equity and cost-effectiveness of health care services." There is a semantic gap in concepts. This classification term is described as activities, i.e., "to inform persons" or "to inform processes" rather than the rationale for performing actions/operations on information related to the activity.
DONAT | http://terminology.hl7.org/CodeSystem/v3-ActReason | donation | To perform one or more operations on information used for cadaveric organ, eye or tissue donation.
FRAUD | http://terminology.hl7.org/CodeSystem/v3-ActReason | fraud | To perform one or more operations on information used for fraud detection and prevention processes.
GOV | http://terminology.hl7.org/CodeSystem/v3-ActReason | government | To perform one or more operations on information used within government processes.
HACCRED | http://terminology.hl7.org/CodeSystem/v3-ActReason | health accreditation | To perform one or more operations on information for conducting activities related to meeting accreditation criteria.
HCOMPL | http://terminology.hl7.org/CodeSystem/v3-ActReason | health compliance | To perform one or more operations on information used for conducting activities required to meet a mandate.
HDECD | http://terminology.hl7.org/CodeSystem/v3-ActReason | decedent | To perform one or more operations on information used for handling deceased patient matters.
HDIRECT | http://terminology.hl7.org/CodeSystem/v3-ActReason | directory | To perform one or more operation operations on information used to manage a patient directory. Examples: facility enterprise payer health information exchange patient directory
HDM | http://terminology.hl7.org/CodeSystem/v3-ActReason | healthcare delivery management | To perform one or more actions on information used for conducting administrative and contractual activities by or on behalf of organizational entities responsible for delivery of an individual's benefits in a healthcare program, health plan or insurance. Explicitly excludes the use of information to organize the delivery of health care for care coordination and case management, or to provide healthcare treatment. Usage Note: Examples of activities conducted under this purpose of use: provider profiling, risk adjustment, underwriting, fraud and abuse, quality improvement population health and care management. Aligns with HIPAA Operation POU minus coordination of care or other treatment related activities. Similar to the description in SAMHSA Confidentiality of Substance Use Disorder Patient Records Supplemental notice of proposed rulemaking. Map: Maps to ISO 14265 Classification Term "Administration of care for an individual subject of care" described as "To inform persons or processes responsible for enabling the availability of resources or funding or permissions for providing health care services to the subject of care." However, this classification term is described as activities, i.e., "to inform persons" or "to inform processes" rather than the rationale for performing actions/operations on information related to the activity.
HLEGAL | http://terminology.hl7.org/CodeSystem/v3-ActReason | legal | To perform one or more operations on information for conducting activities required by legal proceeding.
HOUTCOMS | http://terminology.hl7.org/CodeSystem/v3-ActReason | health outcome measure | To perform one or more operations on information used for assessing results and comparative effectiveness achieved by health care practices and interventions.
HPRGRP | http://terminology.hl7.org/CodeSystem/v3-ActReason | health program reporting | To perform one or more operations on information used for conducting activities to meet program accounting requirements.
HQUALIMP | http://terminology.hl7.org/CodeSystem/v3-ActReason | health quality improvement | To perform one or more operations on information used for conducting administrative activities to improve health care quality.
HSYSADMIN | http://terminology.hl7.org/CodeSystem/v3-ActReason | health system administration | To perform one or more operations on information to administer the electronic systems used for the delivery of health care.
LABELING | http://terminology.hl7.org/CodeSystem/v3-ActReason | labeling | To perform one or more operations on information to assign, persist, and manage labels to healthcare data to characterize various aspects, such as its security classification, sensitivity, compartment, integrity, and provenance; applicable privacy, consent, security, provenance, and trust policies; and handling caveats such as purpose of use, obligations, and refrain policies. Label management includes classification of target data by constructing and binding of a label set per applicable policies, security policy information file semantics, and classification guides. Label management also includes process and procedures for subsequent revision of a label for, e.g., reclassification, downgrading classification, and declassification. Label revisions may be triggered by, e.g., expiry of classification period; changes in applicable policy, e.g., revocation of a consent directive; or changes in the governing policy domain in which the data is relocated or a copy of the data is sent. If a label is revised, an audit log should be kept and the provenance of the label changes should be tracked.
METAMGT | http://terminology.hl7.org/CodeSystem/v3-ActReason | metadata management | To perform one or more operations on information to assign, persist, and manage metadata to healthcare data to characterize various aspects used for its indexing, discovery, retrieval, and processing by systems, applications, and end users. For example, master index identifier, media type, and location.

MEMADMIN | http://terminology.hl7.org/CodeSystem/v3-ActReason | member administration | To perform one or more operations on information to administer health care coverage to an enrollee under a policy or program.
MILCDM | http://terminology.hl7.org/CodeSystem/v3-ActReason | military command | To perform one or more operations on information for conducting activities required by military processes, procedures, policies, or law.
PATADMIN | http://terminology.hl7.org/CodeSystem/v3-ActReason | patient administration | To perform one or more operations on information used for operational activities conducted to administer the delivery of health care to a patient.
PATSFTY | http://terminology.hl7.org/CodeSystem/v3-ActReason | patient safety | To perform one or more operations on information in processes related to ensuring the safety of health care.
PERFMSR | http://terminology.hl7.org/CodeSystem/v3-ActReason | performance measure | To perform one or more operations on information used for monitoring performance of recommended health care practices and interventions.
RECORDMGT | http://terminology.hl7.org/CodeSystem/v3-ActReason | records management | To perform one or more operations on information used within the health records management process.
SYSDEV | http://terminology.hl7.org/CodeSystem/v3-ActReason | system development | To perform one or more operations on information to design, develop, implement, test, or deploy a healthcare system or application.
HTEST | http://terminology.hl7.org/CodeSystem/v3-ActReason | test health data | To perform one or more operations on information that is simulated or synthetic health data used for testing system capabilities outside of a production or operational system environment. Usage Note: Data marked with a HTEST security label enables an access control system to permit interfacing systems or end users provisioned with a clearance, which includes a HTEST purpose of use attribute, to test, verify, or validate that a system or application will operate in production as intended based on design specifications.
TRAIN | http://terminology.hl7.org/CodeSystem/v3-ActReason | training | To perform one or more operations on information used in training and education.
HPAYMT | http://terminology.hl7.org/CodeSystem/v3-ActReason | healthcare payment | To perform one or more operations on information for conducting financial or contractual activities related to payment for provision of health care.
CLMATTCH | http://terminology.hl7.org/CodeSystem/v3-ActReason | claim attachment | To perform one or more operations on information for provision of additional clinical evidence in support of a request for coverage or payment for health services.
COVAUTH | http://terminology.hl7.org/CodeSystem/v3-ActReason | coverage authorization | To perform one or more operations on information for conducting prior authorization or predetermination of coverage for services.
COVERAGE | http://terminology.hl7.org/CodeSystem/v3-ActReason | coverage under policy or program | To perform one or more operations on information for conducting activities related to coverage under a program or policy.
ELIGDTRM | http://terminology.hl7.org/CodeSystem/v3-ActReason | eligibility determination | To perform one or more operations on information used for conducting eligibility determination for coverage in a program or policy. May entail review of financial status or disability assessment.
ELIGVER | http://terminology.hl7.org/CodeSystem/v3-ActReason | eligibility verification | To perform one or more operations on information used for conducting eligibility verification of coverage in a program or policy. May entail provider contacting coverage source (e.g., government health program such as workers compensation or health plan) for confirmation of enrollment, eligibility for specific services, and any applicable copays.
ENROLLM | http://terminology.hl7.org/CodeSystem/v3-ActReason | enrollment | To perform one or more operations on information used for enrolling a covered party in a program or policy. May entail recording of covered party's and any dependent's demographic information and benefit choices.
MILDCRG | http://terminology.hl7.org/CodeSystem/v3-ActReason | military discharge | To perform one or more operations on information for the process of releasing military personnel from their service obligations, which may include determining service merit, discharge benefits, and disability assessment.
REMITADV | http://terminology.hl7.org/CodeSystem/v3-ActReason | remittance advice | To perform one or more operations on information about the amount remitted for a health care claim.
HRESCH | http://terminology.hl7.org/CodeSystem/v3-ActReason | healthcare research | To perform one or more operations on information for conducting scientific investigations to obtain health care knowledge. Use of the data iincludes basic and applied research such as biomedical, population origin or ancestry, translational research, and disease, discipline, specialty specific healthcare research and clinical trial research.
BIORCH | http://terminology.hl7.org/CodeSystem/v3-ActReason | biomedical research | To perform one or more operations on information for conducting scientific investigations to obtain health care knowledge. Use of the data must be related to specified biomedical basic or applied research. For example, research on rare plants to determine whether biologic properties may be useful for pharmaceutical development. May be used in combination with clinical trial and other healthcare research purposes of use.
CLINTRCH | http://terminology.hl7.org/CodeSystem/v3-ActReason | clinical trial research | To perform one or more operations on information for conducting scientific investigations in accordance with clinical trial protocols to obtain health care knowledge.
CLINTRCHNPC | http://terminology.hl7.org/CodeSystem/v3-ActReason | clinical trial research without patient care | To perform one or more operations on information for conducting scientific investigations in accordance with clinical trial protocols to obtain health care knowledge without provision of patient care. May be post-coordinated or used with other purposes of use such as disease, discipline, specialty, population origins or ancestry, translational healthcare research. For example, a clinical trial conducted on laboratory specimens collected from a specified patient population.
CLINTRCHPC | http://terminology.hl7.org/CodeSystem/v3-ActReason | clinical trial research with patient care | To perform one or more operations on information for conducting scientific investigations with patient care in accordance with clinical trial protocols to obtain health care knowledge. May be post-coordinated or used with other purposes of use such as disease, discipline, specialty, population origins or ancestry, translational healthcare research. For example, an "off-label" drug used for cancer therapy administer to a specified patient population.
PRECLINTRCH | http://terminology.hl7.org/CodeSystem/v3-ActReason | preclinical trial research | To perform one or more operations on information in preparation for conducting scientific investigation to obtain health care knowledge, such as research on animals or review of patient health records, to determine the feasibility of a clinical trial study; assist with protocol design; or in preparation for institutional review board or ethics committee approval process. May be post-coordinated or used with other purposes of use such as disease, discipline, specialty, population origins or ancestry, translational healthcare research.
DSRCH | http://terminology.hl7.org/CodeSystem/v3-ActReason | disease specific healthcare research | To perform one or more operations on information for conducting scientific investigations to obtain health care knowledge. Use of the data must be related to specified conditions, diagnosis, or disease healthcare research. For example, conducting cancer research by testing reaction of tumor cells to certain biologics. May be used in combination with clinical trial and other healthcare research purposes of use.
POARCH | http://terminology.hl7.org/CodeSystem/v3-ActReason | population origins or ancestry healthcare research | To perform one or more operations on information, including genealogical pedigrees, historical records, surveys, family health data, health records, and genetic information, for conducting scientific investigations to obtain health care knowledge. Use of the data must be related to population origins and/or ancestry healthcare research. For example, gathering genetic specimens from a specific

population in order to determine the ancestry and population origins of that group. May be used in combination with clinical trial and other healthcare research purposes of use.
TRANSRCH | http://terminology.hl7.org/CodeSystem/v3-ActReason | translational healthcare research | To perform one or more operations on information for conducting scientific investigations to obtain health care knowledge related to evidence based medicine during the course of providing healthcare treatment. Sometimes referred to as "bench to bedside", which is the iterative feedback loop between healthcare research and clinical trials with input from information collected in the course of routine provision of healthcare. For example, by extending a patient encounter to conduct a survey related to a research topic such as attitudes about use of a wellness device that a patient agreed to use. May be used in combination with clinical trial and other healthcare research purposes of use.
PATRQT | http://terminology.hl7.org/CodeSystem/v3-ActReason | patient requested | To perform one or more operations on information in response to a patient's request.
FAMRQT | http://terminology.hl7.org/CodeSystem/v3-ActReason | family requested | To perform one or more operations on information in response to a request by a family member authorized by the patient.
PWATRNY | http://terminology.hl7.org/CodeSystem/v3-ActReason | power of attorney | To perform one or more operations on information in response to a request by a person appointed as the patient's legal representative.
SUPNWK | http://terminology.hl7.org/CodeSystem/v3-ActReason | support network | To perform one or more operations on information in response to a request by a person authorized by the patient.
PUBHLTH | http://terminology.hl7.org/CodeSystem/v3-ActReason | public health | To perform one or more operations on information for conducting public health activities, such as the reporting of notifiable conditions.
DISASTER | http://terminology.hl7.org/CodeSystem/v3-ActReason | disaster | To perform one or more operations on information used for provision of immediately needed health care to a population of living subjects located in a disaster zone.
THREAT | http://terminology.hl7.org/CodeSystem/v3-ActReason | threat | To perform one or more operations on information used to prevent injury or disease to living subjects who may be the target of violence.
TREAT | http://terminology.hl7.org/CodeSystem/v3-ActReason | treatment | To perform one or more operations on information for provision of health care.
CLINTRL | http://terminology.hl7.org/CodeSystem/v3-ActReason | clinical trial | To perform health care as part of the clinical trial protocol.
COC | http://terminology.hl7.org/CodeSystem/v3-ActReason | coordination of care | To perform one or more actions on information in order to organize the provision and case management of an individualâ€™s healthcare, including: Monitoring a person's goals, needs, and preferences; acting as the communication link between two or more participants concerned with a person's health and wellness; organizing and facilitating care activities and promoting self-management by advocating for, empowering, and educating a person; and ensuring safe, appropriate, non-duplicative, and effective integrated care. Usage Note: Use when describing these functions: 1. Monitoring a personâ€™s goals, needs, and preferences. 2. Acting as the communication link between two or more participants concerned with a person's health and wellness. 3. Organizing and facilitating care activities and promoting self-management by advocating for, empowering, and educating a person. 4. Ensuring safe, appropriate, non-duplicative, and effective integrated care. The goal is to clearly differentiate this type of coordination of care from HIPAA Operations by specifying that these actions on information are undertaken in the provision of healthcare treatment. For similar uses of this concept, see SAMHSA Confidentiality of Substance Use Disorder Patient Records Supplemental notice of proposed rulemaking, which differentiates concepts of care coordination and case management for the provision of treatment as specifically distinct from activities related to health care delivery management and the operations of organizational entities involved in the delivery of healthcare. Map: Maps to ISO 14265 Classification Terms: "Support of care activities within the provider organisation for an individual subject of care" described as "To inform persons or processes enabling others to provide health care services to the subject of care." "Subject of Care Uses" described as "To inform the subject of care in support of his or her own interests."
ETREAT | http://terminology.hl7.org/CodeSystem/v3-ActReason | Emergency Treatment | To perform one or more operations on information for provision of immediately needed health care for an emergent condition.
BTG | http://terminology.hl7.org/CodeSystem/v3-ActReason | break the glass | To perform policy override operations on information for provision of immediately needed health care for an emergent condition affecting potential harm, death or patient safety by end users who are not provisioned for this purpose of use. Includes override of organizational provisioning policies and may include override of subject of care consent directive restricting access. Map: Partially Maps to ISO 14265 Classification Term "Emergency care provision to an individual subject of care" described as "To inform persons needing to provide health care services to the subject of care urgently, possibly needing to over-ride the policies and consents pertaining to Purpose 1 above." Purpose 1 is equivalent to HL7 treatment purpose of use: "Clinical care provision to an individual subject of care" described as "To inform persons or processes responsible for providing health care services to the subject of care." The ISO description conflates both of the proposed specializations of HL7 ETREAT: break the glass and the typically broader access to health information normally available to providers who are provisioned for emergency workflows on a regular basis, e.g., Emergency Room providers. Examples of greater access than is normally accessible by providers based on the need to know are access to sensitive information for which access typically requires a patient's consent. This is not an override of a patient's dissent to disclose sensitive information in cases where the applicable policy waives the need for that consent to access this information. In US, Title 38 Section 7332 and 42 CFR Part 2 both permit emergency access without the need to override a patient's consent directive; rather, this access is a limitation to the patient's right to dissent from disclosure.
ERTREAT | http://terminology.hl7.org/CodeSystem/v3-ActReason | emergency room treatment | To perform one or more operations on information for provision of immediately needed health care for an emergent condition in an emergency room or similar emergent care context by end users provisioned for this purpose, which does not constitute as policy override such as in a "Break the Glass" purpose of use. Map:Partially Maps to ISO 14265 Classification Term "Emergency care provision to an individual subject of care" described as "To inform persons needing to provide health care services to the subject of care urgently, possibly needing to over-ride the policies and consents pertaining to Purpose 1 above." Purpose 1 is equivalent to HL7 treatment purpose of use: "Clinical care provision to an individual subject of care" described as "To inform persons or processes responsible for providing health care services to the subject of care." The ISO description conflates both of the proposed specializations of HL7 ETREAT: break the glass and the typically broader access to health information normally available to providers who are provisioned for emergency workflows on a regular basis, e.g., Emergency Room providers. Examples of greater access than is normally accessible by providers based on the need to know are access to sensitive information for which access typically requires a patient's consent. This is not an override of a patient's dissent to disclose sensitive information in cases where the applicable policy waives the need for that consent to access this information. In US, Title 38 Section 7332 and 42 CFR Part 2 both permit emergency access without the need to override a patient's consent directive; rather, this access is a limitation to the patient's right to dissent from disclosure. There is a semantic gap in concepts. This classification term is described as activities â€œto inform personsâ€? rather than the rationale for performing actions/operations on information related to the activity.
POPHLTH | http://terminology.hl7.org/CodeSystem/v3-ActReason | population health | To perform one or more operations on information for provision of health care to a population of living subjects, e.g., needle exchange program.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.616 Value Set http://hl7.org/fhir/ValueSet/participation-role-type

URL: https://hl7.org/fhir/R4/valueset-participation-role-type.html
Vocabulary Work Group | Maturity Level: 3 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/participation-role-type
Version: | 4.0.1
Name: | ParticipationRoleType
Title: | ParticipationRoleType
Definition: | This FHIR value set is comprised of Actor participation Type codes, which can be used to value FHIR agents, actors, and other role elements. The FHIR Actor participation type value set is based on DICOM Audit Message, C402; ASTM Standard, E1762-95 [2013]; selected codes and derived actor roles from HL7 RoleClass OID 2.16.840.1.113883.5.110; HL7 Role Code 2.16.840.1.113883.5.111, including AgentRoleType; HL7 ParticipationType OID: 2.16.840.1.113883.5.90; and HL7 ParticipationFunction codes OID: 2.16.840.1.113883.5.88. This value set includes, by reference, role codes from external code systems: NUCC Health Care Provider Taxonomy OID: 2.16.840.1.113883.6.101; North American Industry Classification System [NAICS]OID: 2.16.840.1.113883.6.85; IndustryClassificationSystem 2.16.840.1.113883.1.11.16039; and US Census Occupation Code OID: 2.16.840.1.113883.6.243 for relevant recipient or custodian codes not included in this value set. If no source is indicated in the definition comments, then these are example FHIR codes.
Committee: | Vocabulary Work Group
OID: | 2.16.840.1.113883.4.642.3.54 (for OID based terminology systems)
Copyright: | This is an example FHIR value set based on ASTM Standard, E1762-95 (2013) HL7 RoleClass OID 2.16.840.1.113883.5.110, HL7 Role Code 2.16.840.1.113883.5.111, HL7 ParticipationType OID: 2.16.840.1.113883.5.90, HL7 ParticipationFunction codes at OID: 2.16.840.1.113883.5.88, and HL7 Security and Privacy Domain Analysis Model roles classes. These codes are excerpted from Digital Imaging and Communications in Medicine (DICOM) Standard, Part 16: Content Mapping Resource, Copyright © 2011 by the National Electrical Manufacturers Association.
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: AuditEvent.agent.type (CodeableConcept / Extensible)

4.4.1.616.1 Content Logical Definition

This value set includes codes from the following code systems:

http://terminology.hl7.org/CodeSystem/contractsignertypecodes

Code | Display
AMENDER | Amender | A person who has corrected, edited, or amended pre-existing information.
COAUTH | Co-Author | The entity that co-authored content. There can be multiple co-authors of content, which may take such forms as a contract, a healthcare record entry or document, a policy, or a consent directive.
CONT | Contact | A person or an organization that provides or receives information regarding another entity. Examples; patient NOK and emergency contacts; guarantor contact; employer contact.
EVTWIT | Event Witness | A person who attests to observing an occurrence. For example, the witness has observed a procedure and is attesting to this fact.
PRIMAUTH | Primary Author | An entity that is the primary or sole author of information content. In the healthcare context, there can be only one primary author of health information content in a record entry or document.
REVIEWER | Reviewer | A person, device, or algorithm that has used approved criteria for filtered data for inclusion into the patient record. Examples: (1) a medical records clerk who scans a document for inclusion in the medical record, enters header information, or catalogues and classifies the data, or a combination thereof; (2) a gateway that receives data from another computer system and interprets that data or changes its format, or both, before entering it into the patient record.
SOURCE | Source | An automated data source that generates a signature along with content. Examples: (1) the signature for an image that is generated by a device for inclusion in the patient record; (2) the signature for an ECG derived by an ECG system for inclusion in the patient record; (3) the data from a biomedical monitoring device or system that is for inclusion in the patient record.
TRANS | Transcriber | An entity entering the data into the originating system. This includes the transcriptionist for dictated text transcribed into electronic form.
VALID | Validator | A person who validates a health information document for inclusion in the patient record. For example, a medical student or resident is credentialed to perform history or physical examinations and to write progress notes. The attending physician signs the history and physical examination to validate the entry for inclusion in the patient's medical record.

VERF | Verifier | A person who asserts the correctness and appropriateness of an act or the recording of the act, and is accountable for the assertion that the act or the recording of the act complies with jurisdictional or organizational policy. For example, a physician is required to countersign a verbal order that has previously been recorded in the medical record by a registered nurse who has carried out the verbal order.

http://terminology.hl7.org/CodeSystem/v3-RoleClass

Code | Display

AFFL | affiliate | Player of the Affiliate role has a business/professional relationship with scoper. Player and scoper may be persons or organization. The Affiliate relationship does not imply membership in a group, nor does it exist for resource scheduling purposes. Example: A healthcare provider is affiliated with another provider as a business associate.

AGNT | agent | An entity (player) that acts or is authorized to act on behalf of another entity (scoper).

ASSIGNED | assigned entity | An agent role in which the agent is an Entity acting in the employ of an organization. The focus is on functional role on behalf of the organization, unlike the Employee role where the focus is on the 'Human Resources' relationship between the employee and the organization.

CLAIM | claimant | Description: A role played by a party making a claim for coverage under a policy or program. A claimant must be either a person or organization, or a group of persons or organizations. A claimant is not a named insured or a program eligible. Discussion: With respect to liability insurance such as property and casualty insurance, a claimant must file a claim requesting indemnification for a loss that the claimant considers covered under the policy of a named insured. The claims adjuster for the policy underwriter will review the claim to determine whether the loss meets the benefit coverage criteria under a policy, and base any indemnification or coverage payment on that review. If a third party is liable in whole or part for the loss, the underwriter may pursue third party liability recovery. A claimant may be involved in civil or criminal legal proceedings involving claims against a defendant party that is indemnified by an insurance policy or to protest the finding of a claims adjustor. With respect to life insurance, a beneficiary designated by a named insured becomes a claimant of the proceeds of coverage, as in the case of a life insurance policy. However, a claimant for coverage under life insurance is not necessarily a designated beneficiary. Note: A claimant is not a named insured. However, a named insured may make a claim under a policy, e.g., an insured driver may make a claim for an injury under his or her comprehensive automobile insurance policy. Similarly, a program eligible may make a claim under program, e.g., an unemployed worker may claim benefits under an unemployment insurance program, but parties playing these covered party role classes are not, for purposes of this vocabulary and in an effort to clearly distinguish role classes, considered claimants. In the case of a named insured making a claim, a role type code INSCLM (insured claimant) subtypes the class to indicate that either a named insured or an individual insured has filed a claim for a loss. In the case of a program eligible, a role type code INJWKR (injured worker) subtypes the class to indicate that the covered party in a workers compensation program is an injured worker, and as such, has filed a "claim" under the program for benefits. Likewise, a covered role type code UNEMP (unemployed worker) subtypes the program eligible class to indicate that the covered party in an unemployment insurance program has filed a claim for unemployment benefits. Example: A claimant under automobile policy that is not the named insured.

COVPTY | covered party | A role class played by a person who receives benefit coverage under the terms of a particular insurance policy. The underwriter of that policy is the scoping entity. The covered party receives coverage because of some contractual or other relationship with the holder of that policy. Discussion:This reason for coverage is captured in 'Role.code' and a relationship link with type code of indirect authority should be included using the policy holder role as the source, and the covered party role as the target. Note that a particular policy may cover several individuals one of whom may be, but need not be, the policy holder. Thus the notion of covered party is a role that is distinct from that of the policy holder.

DEPEN | dependent | Description: A role played by a person covered under a policy or program based on an association with a subscriber, which is recognized by the policy holder. Note: The party playing the role of a dependent is not a claimant in the sense conveyed by the RoleClassCoveredParty CLAIM (claimant). However, a dependent may make a claim under a policy, e.g., a dependent under a health insurance policy may become the claimant for coverage under that policy for wellness examines or if injured and there is no liable third party. In the case of a dependent making a claim, a role type code INSCLM (insured claimant) subtypes the class to indicate that the dependent has filed a claim for services covered under the health insurance policy. Example: The dependent has an association with the subscriber such as a financial dependency or personal relationship such as that of a spouse, or a natural or adopted child. The policy holder may be required by law to recognize certain associations or may have discretion about the associations. For example, a policy holder may dictate the criteria for the dependent status of adult children who are students, such as requiring full time enrollment, or may recognize domestic partners as dependents. Under certain circumstances, the dependent may be under the indirect authority of a responsible party acting as a surrogate for the subscriber, for example, if the subscriber is differently abled or deceased, a guardian ad Lidem or estate executor may be appointed to assume the subscriberaTMs legal standing in the relationship with the dependent.

ECON | emergency contact | An entity to be contacted in the event of an emergency.

EMP | employee | A relationship between a person or organization and a person or organization formed for the purpose of exchanging work for compensation. The purpose of the role is to identify the type of relationship the employee has to the employer, rather than the nature of the work actually performed. (Contrast with AssignedEntity.)

GUARD | guardian | Guardian of a ward

INVSBJ | Investigation Subject | An entity that is the subject of an investigation. This role is scoped by the party responsible for the investigation.

NAMED | named insured | Description: A role played by a party to an insurance policy to which the insurer agrees to indemnify for losses, provides benefits for, or renders services. A named insured may be either a person, non-person living subject, or an organization, or a group of persons, non-person living subjects, or organizations. Discussion: The coded concept NAMED should not be used where a more specific child concept in this Specializable value set applies. In some cases, the named insured may not be the policy holder, e.g., where a policy holder purchases life insurance policy in which another party is the named insured and the policy holder is the beneficiary of the policy. Note: The party playing the role of a named insured is not a claimant in the sense conveyed by the RoleClassCoveredParty CLAIM (claimant). However, a named insured may make a claim under a policy, e.g., e.g., a party that is the named insured and policy holder under a comprehensive automobile insurance policy may become the claimant for coverage under that policy e.g., if injured in an automobile accident and there is no liable third party. In the case of a named insured making a claim, a role type code INSCLM (insured claimant) subtypes the class to indicate that a named insured has filed a claim for a loss. Example: The named insured under a comprehensive automobile, disability, or property and casualty policy that is the named insured and may or may not be the policy holder.

NOK | next of kin | An individual designated for notification as the next of kin for a given entity.

PAT | patient | A Role of a LivingSubject (player) as an actual or potential recipient of health care services from a healthcare provider organization (scoper). Usage Note: Communication about relationships between patients and specific healthcare practitioners (people) is not done via scoper. Instead this is generally done using the CareProvision act. This allows linkage between patient and a particular healthcare practitioner role and also allows description of the type of care involved in the relationship.

PROV | healthcare provider | An Entity (player) that is authorized to provide health care services by some authorizing agency (scoper).
NOT | notary public | notary public

http://terminology.hl7.org/CodeSystem/v3-RoleCode

Code | Display
CLASSIFIER | classifier | An individual authorized to assign an original classification to information, including compilations of unclassified information, based on a determination that the information requires protection against unauthorized disclosure. The individual marks the information with immutable, computable, and human readable security labels in accordance with applicable security labeling policies. The labeling policies provide instructions on whether and if so how the security labels may be later reclassified [i.e., upgraded, downgraded, used in derivative classification, or declassified] in a manner that preserves the overridden original classification binding and provenance.
CONSENTER | consenter | An entity or an entity's delegatee who is the grantee in an agreement such as a consent for services, advanced directive, or a privacy consent directive in accordance with jurisdictional, organizational, or patient policy.
CONSWIT | consent witness | An entity which has witnessed and attests to observing another entity being counseled about an agreement such as a consent for services, advanced directive, or a privacy consent directive.
COPART | co-participant | An entity which participates in the generation of and attest to veracity of content, but is not an author or coauthor. For example a surgeon who is required by institutional, regulatory, or legal rules to sign an operative report, but who was not involved in the authorship of that report.
DECLASSIFIER | declassifier | An individual which is authorized to declassify information based on a determination that the information no longer requires protection against unauthorized disclosure. The individual marks the information being declassified using computable and human readable security labels indicating that this is copy of previously classified information is unclassified in accordance with applicable security labeling policies. The labeling policies provide instructions on whether and if so how the security labels may be later reclassified [i.e., upgraded or used in derivative classification] in a manner that preserves the overridden original classification binding and provenance.
DELEGATEE | delegatee | A party to whom some right or authority is granted by a delegator.
DELEGATOR | delegator | A party that grants all or some portion its right or authority to another party.
DOWNGRDER | downgrader | An individual authorized to lower the classification level of labeled content and provide rationale for doing so as directed by a classification guide.
DPOWATT | durable power of attorney | A relationship between two people in which one person authorizes another, usually a family member or relative, to act for him or her in a manner which is a legally binding upon the person giving such authority as if he or she personally were to do the acts that is often limited in the kinds of powers that can be assigned. Unlike ordinary powers of attorney, durable powers can survive for long periods of time, and again, unlike standard powers of attorney, durable powers can continue after incompetency.
EXCEST | executor of estate | The role played by a person acting as the estate executor for a deceased subscriber or policyholder who was the responsible party
GRANTEE | grantee | An entity which accepts certain rights or authority from a grantor.
GRANTOR | grantor | An entity which agrees to confer certain rights or authority to a grantee.
GT | Guarantor | An individual or organization that makes or gives a promise, assurance, pledge to pay or has paid the healthcare service provider.
GUADLTM | guardian ad lidem | The role played by a person appointed by the court to look out for the best interests of a minor child during the course of legal proceedings.
HPOWATT | healthcare power of attorney | A relationship between two people in which one person authorizes another to act for him or her in a manner which is a legally binding upon the person giving such authority as if he or she personally were to do the acts that continues (by its terms) to be effective even though the grantor has become mentally incompetent after signing the document.
INTPRTER | interpreter | An entity which converts spoken or written language into the language of key participants in an event such as when a provider is obtaining a patient's consent to treatment or permission to disclose information.
POWATT | power of attorney | A relationship between two people in which one person authorizes another to act for him in a manner which is a legally binding upon the person giving such authority as if he or she personally were to do the acts.
RESPRSN | responsible party | The role played by a party who has legal responsibility for another party.
SPOWATT | special power of attorney | A relationship between two people in which one person authorizes another to act for him or her in a manner which is a legally binding upon the person giving such authority as if he or she personally were to do the acts that is often limited in the kinds of powers that can be assigned.

• Include codes from http://terminology.hl7.org/CodeSystem/v3-RoleCode where concept is-a _CitizenRoleType

http://terminology.hl7.org/CodeSystem/v3-ParticipationFunction

Code | Display
AUCG | caregiver information receiver | Description:Caregiver authorized to receive patient health information.
AULR | legitimate relationship information receiver | Description:Provider with legitimate relationship authorized to receive patient health information.
AUTM | care team information receiver | Description:Member of care team authorized to receive patient health information.
AUWA | work area information receiver | Description:Entities within specified work area authorized to receive patient health information.
PROMSK | authorized provider masking author | Definition:Provider authorized to mask information to protect the patient, a third party, or to ensure that the provider has consulted with the patient prior to release of this information.

http://terminology.hl7.org/CodeSystem/v3-ParticipationType

Code | Display
AUT | author (originator) | Definition: A party that originates the Act and therefore has responsibility for the information given in the Act and ownership of this Act. Example: the report writer, the person writing the act definition, the guideline author, the placer of an order, the EKG cart (device) creating a report etc. Every Act should have an author. Authorship is regardless of mood always actual authorship. Examples of such policies might include: The author and anyone they explicitly delegate may update the report; All administrators within the same clinic may cancel and reschedule appointments created by other administrators within that clinic; A party that is neither an author nor a party who is extended authorship maintenance rights by policy, may only amend,

reverse, override, replace, or follow up in other ways on this Act, whereby the Act remains intact and is linked to another Act authored by that other party.

CST | custodian | An entity (person, organization or device) that is in charge of maintaining the information of this act (e.g., who maintains the report or the master service catalog item, etc.).

INF | informant | A source of reported information (e.g., a next of kin who answers questions about the patient's history). For history questions, the patient is logically an informant, yet the informant of history questions is implicitly the subject.

IRCP | information recipient | A party, who may or should receive or who has recieved the Act or subsequent or derivative information of that Act. Information recipient is inert, i.e., independent of mood." Rationale: this is a generalization of a too diverse family that the definition can't be any more specific, and the concept is abstract so one of the specializations should be used.

LA | legal authenticator | A verifier who legally authenticates the accuracy of an act. An example would be a staff physician who sees a patient and dictates a note, then later signs it. Their signature constitutes a legal authentication.

IRCP | information recipient | A party, who may or should receive or who has recieved the Act or subsequent or derivative information of that Act. Information recipient is inert, i.e., independent of mood." Rationale: this is a generalization of a too diverse family that the definition can't be any more specific, and the concept is abstract so one of the specializations should be used.

TRC | tracker | A secondary information recipient, who receives copies (e.g., a primary care provider receiving copies of results as ordered by specialist).

WIT | witness | Only with service events. A person witnessing the action happening without doing anything. A witness is not necessarily aware, much less approves of anything stated in the service event. Example for a witness is students watching an operation or an advanced directive witness.

• Include all codes defined in http://terminology.hl7.org/CodeSystem/extra-security-role-type

http://dicom.nema.org/resources/ontology/DCM

Code | Display

110150 | Application | Audit participant role ID of software application

110151 | Application Launcher | Audit participant role ID of software application launcher, i.e., the entity that started or stopped an application

110152 | Destination Role ID | Audit participant role ID of the receiver of data

110153 | Source Role ID | Audit participant role ID of the sender of data

110154 | Destination Media | Audit participant role ID of media receiving data during an export

110155 | Source Media | Audit participant role ID of media providing data during an import

4.4.1.616.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 76 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/extra-security-role-type version 4.0.1, http://terminology.hl7.org/CodeSystem/v3-RoleCode version 2018-08-12, http://terminology.hl7.org/CodeSystem/v3-RoleClass version 2018-08-12, http://terminology.hl7.org/CodeSystem/v3-ParticipationFunction version 2018-08-12, http://terminology.hl7.org/CodeSystem/contractsignertypecodes version 4.0.1, http://terminology.hl7.org/CodeSystem/v3-ParticipationType version 2018-08-12, DICOM version 01

Code | System | Display | Definition

AMENDER | http://terminology.hl7.org/CodeSystem/contractsignertypecodes | Amender | A person who has corrected, edited, or amended pre-existing information.

COAUTH | http://terminology.hl7.org/CodeSystem/contractsignertypecodes | Co-Author | The entity that co-authored content. There can be multiple co-authors of content, which may take such forms as a contract, a healthcare record entry or document, a policy, or a consent directive.

CONT | http://terminology.hl7.org/CodeSystem/contractsignertypecodes | Contact | A person or an organization that provides or receives information regarding another entity. Examples; patient NOK and emergency contacts; guarantor contact; employer contact.

EVTWIT | http://terminology.hl7.org/CodeSystem/contractsignertypecodes | Event Witness | A person who attests to observing an occurrence. For example, the witness has observed a procedure and is attesting to this fact.

PRIMAUTH | http://terminology.hl7.org/CodeSystem/contractsignertypecodes | Primary Author | An entity that is the primary or sole author of information content. In the healthcare context, there can be only one primary author of health information content in a record entry or document.

REVIEWER | http://terminology.hl7.org/CodeSystem/contractsignertypecodes | Reviewer | A person, device, or algorithm that has used approved criteria for filtered data for inclusion into the patient record. Examples: (1) a medical records clerk who scans a document for inclusion in the medical record, enters header information, or catalogues and classifies the data, or a combination thereof; (2) a gateway that receives data from another computer system and interprets that data or changes its format, or both, before entering it into the patient record.

SOURCE | http://terminology.hl7.org/CodeSystem/contractsignertypecodes | Source | An automated data source that generates a signature along with content. Examples: (1) the signature for an image that is generated by a device for inclusion in the patient record; (2) the signature for an ECG derived by an ECG system for inclusion in the patient record; (3) the data from a biomedical monitoring device or system that is for inclusion in the patient record.

TRANS | http://terminology.hl7.org/CodeSystem/contractsignertypecodes | Transcriber | An entity entering the data into the originating system. This includes the transcriptionist for dictated text transcribed into electronic form.

VALID | http://terminology.hl7.org/CodeSystem/contractsignertypecodes | Validator | A person who validates a health information document for inclusion in the patient record. For example, a medical student or resident is credentialed to perform history or physical examinations and to write progress notes. The attending physician signs the history and physical examination to validate the entry for inclusion in the patient's medical record.

VERF | http://terminology.hl7.org/CodeSystem/contractsignertypecodes | Verifier | A person who asserts the correctness and appropriateness of an act or the recording of the act, and is accountable for the assertion that the act or the recording of the act complies with jurisdictional or organizational policy. For example, a physician is required to countersign a verbal order that has previously been recorded in the medical record by a registered nurse who has carried out the verbal order.

AFFL | http://terminology.hl7.org/CodeSystem/v3-RoleClass | affiliate | Player of the Affiliate role has a business/professional relationship with scoper. Player and scoper may be persons or organization. The Affiliate relationship does not imply membership in a group, nor does it exist for resource scheduling purposes. Example: A healthcare provider is affiliated with another provider as a business associate.

AGNT | http://terminology.hl7.org/CodeSystem/v3-RoleClass | agent | An entity (player) that acts or is authorized to act on behalf of another entity (scoper).

ASSIGNED | http://terminology.hl7.org/CodeSystem/v3-RoleClass | assigned entity | An agent role in which the agent is an Entity acting in the employ of an organization. The focus is on functional role on behalf of the organization, unlike the Employee role where the focus is on the 'Human Resources' relationship between the employee and the organization.

CLAIM | http://terminology.hl7.org/CodeSystem/v3-RoleClass | claimant | Description: A role played by a party making a claim for coverage under a policy or program. A claimant must be either a person or organization, or a group of persons or organizations. A claimant is not a named insured or a program eligible. Discussion: With respect to liability insurance such as property and casualty insurance, a claimant must file a claim requesting indemnification for a loss that the claimant considers covered under the policy of a named insured. The claims adjuster for the policy underwriter will review the claim to determine whether the loss meets the benefit coverage criteria under a policy, and base any indemnification or coverage payment on that review. If a third party is liable in whole or part for the loss, the underwriter may pursue third party liability recovery. A claimant may be involved in civil or criminal legal proceedings involving claims against a defendant party that is indemnified by an insurance policy or to protest the finding of a claims adjustor. With respect to life insurance, a beneficiary designated by a named insured becomes a claimant of the proceeds of coverage, as in the case of a life insurance policy. However, a claimant for coverage under life insurance is not necessarily a designated beneficiary. Note: A claimant is not a named insured. However, a named insured may make a claim under a policy, e.g., an insured driver may make a claim for an injury under his or her comprehensive automobile insurance policy. Similarly, a program eligible may make a claim under program, e.g., an unemployed worker may claim benefits under an unemployment insurance program, but parties playing these covered party role classes are not, for purposes of this vocabulary and in an effort to clearly distinguish role classes, considered claimants. In the case of a named insured making a claim, a role type code INSCLM (insured claimant) subtypes the class to indicate that either a named insured or an individual insured has filed a claim for a loss. In the case of a program eligible, a role type code INJWKR (injured worker) subtypes the class to indicate that the covered party in a workers compensation program is an injured worker, and as such, has filed a "claim" under the program for benefits. Likewise, a covered role type code UNEMP (unemployed worker) subtypes the program eligible class to indicate that the covered party in an unemployment insurance program has filed a claim for unemployment benefits. Example: A claimant under automobile policy that is not the named insured.

COVPTY | http://terminology.hl7.org/CodeSystem/v3-RoleClass | covered party | A role class played by a person who receives benefit coverage under the terms of a particular insurance policy. The underwriter of that policy is the scoping entity. The covered party receives coverage because of some contractual or other relationship with the holder of that policy. Discussion:This reason for coverage is captured in 'Role.code' and a relationship link with type code of indirect authority should be included using the policy holder role as the source, and the covered party role as the target. Note that a particular policy may cover several individuals one of whom may be, but need not be, the policy holder. Thus the notion of covered party is a role that is distinct from that of the policy holder.

DEPEN | http://terminology.hl7.org/CodeSystem/v3-RoleClass | dependent | Description: A role played by a person covered under a policy or program based on an association with a subscriber, which is recognized by the policy holder. Note: The party playing the role of a dependent is not a claimant in the sense conveyed by the RoleClassCoveredParty CLAIM (claimant). However, a dependent may make a claim under a policy, e.g., a dependent under a health insurance policy may become the claimant for coverage under that policy for wellness examines or if injured and there is no liable third party. In the case of a dependent making a claim, a role type code INSCLM (insured claimant) subtypes the class to indicate that the dependent has filed a claim for services covered under the health insurance policy. Example: The dependent has an association with the subscriber such as a financial dependency or personal relationship such as that of a spouse, or a natural or adopted child. The policy holder may be required by law to recognize certain associations or may have discretion about the associations. For example, a policy holder may dictate the criteria for the dependent status of adult children who are students, such as requiring full time enrollment, or may recognize domestic partners as dependents. Under certain circumstances, the dependent may be under the indirect authority of a responsible party acting as a surrogate for the subscriber, for example, if the subscriber is differently abled or deceased, a guardian ad Lidem or estate executor may be appointed to assume the subscriberaTMs legal standing in the relationship with the dependent.

ECON | http://terminology.hl7.org/CodeSystem/v3-RoleClass | emergency contact | An entity to be contacted in the event of an emergency.

EMP | http://terminology.hl7.org/CodeSystem/v3-RoleClass | employee | A relationship between a person or organization and a person or organization formed for the purpose of exchanging work for compensation. The purpose of the role is to identify the type of relationship the employee has to the employer, rather than the nature of the work actually performed. (Contrast with AssignedEntity.)

GUARD | http://terminology.hl7.org/CodeSystem/v3-RoleClass | guardian | Guardian of a ward

INVSBJ | http://terminology.hl7.org/CodeSystem/v3-RoleClass | Investigation Subject | An entity that is the subject of an investigation. This role is scoped by the party responsible for the investigation.

NAMED | http://terminology.hl7.org/CodeSystem/v3-RoleClass | named insured | Description: A role played by a party to an insurance policy to which the insurer agrees to indemnify for losses, provides benefits for, or renders services. A named insured may be either a person, non-person living subject, or an organization, or a group of persons, non-person living subjects, or organizations. Discussion: The coded concept NAMED should not be used where a more specific child concept in this Specializable value set applies. In some cases, the named insured may not be the policy holder, e.g., where a policy holder purchases life insurance policy in which another party is the named insured and the policy holder is the beneficiary of the policy. Note: The party playing the role of a named insured is not a claimant in the sense conveyed by the RoleClassCoveredParty CLAIM (claimant). However, a named insured may make a claim under a policy, e.g., e.g., a party that is the named insured and policy holder under a comprehensive automobile insurance policy may become the claimant for coverage under that policy e.g., if injured in an automobile accident and there is no liable third party. In the case of a named insured making a claim, a role type code INSCLM (insured claimant) subtypes the class to indicate that a named insured has filed a claim for a loss. Example: The named insured under a comprehensive automobile, disability, or property and casualty policy that is the named insured and may or may not be the policy holder.

NOK | http://terminology.hl7.org/CodeSystem/v3-RoleClass | next of kin | An individual designated for notification as the next of kin for a given entity.

PAT | http://terminology.hl7.org/CodeSystem/v3-RoleClass | patient | A Role of a LivingSubject (player) as an actual or potential recipient of health care services from a healthcare provider organization (scoper). Usage Note: Communication about relationships between patients and specific healthcare practitioners (people) is not done via scoper. Instead this is generally done using the

CareProvision act. This allows linkage between patient and a particular healthcare practitioner role and also allows description of the type of care involved in the relationship.

PROV | http://terminology.hl7.org/CodeSystem/v3-RoleClass | healthcare provider | An Entity (player) that is authorized to provide health care services by some authorizing agency (scoper).

NOT | http://terminology.hl7.org/CodeSystem/v3-RoleClass | notary public | notary public

CLASSIFIER | http://terminology.hl7.org/CodeSystem/v3-RoleCode | classifier | An individual authorized to assign an original classification to information, including compilations of unclassified information, based on a determination that the information requires protection against unauthorized disclosure. The individual marks the information with immutable, computable, and human readable security labels in accordance with applicable security labeling policies. The labeling policies provide instructions on whether and if so how the security labels may be later reclassified [i.e., upgraded, downgraded, used in derivative classification, or declassified] in a manner that preserves the overridden original classification binding and provenance.

CONSENTER | http://terminology.hl7.org/CodeSystem/v3-RoleCode | consenter | An entity or an entity's delegatee who is the grantee in an agreement such as a consent for services, advanced directive, or a privacy consent directive in accordance with jurisdictional, organizational, or patient policy.

CONSWIT | http://terminology.hl7.org/CodeSystem/v3-RoleCode | consent witness | An entity which has witnessed and attests to observing another entity being counseled about an agreement such as a consent for services, advanced directive, or a privacy consent directive.

COPART | http://terminology.hl7.org/CodeSystem/v3-RoleCode | co-participant | An entity which participates in the generation of and attest to veracity of content, but is not an author or coauthor. For example a surgeon who is required by institutional, regulatory, or legal rules to sign an operative report, but who was not involved in the authorship of that report.

DECLASSIFIER | http://terminology.hl7.org/CodeSystem/v3-RoleCode | declassifier | An individual which is authorized to declassify information based on a determination that the information no longer requires protection against unauthorized disclosure. The individual marks the information being declassified using computable and human readable security labels indicating that this is copy of previously classified information is unclassified in accordance with applicable security labeling policies. The labeling policies provide instructions on whether and if so how the security labels may be later reclassified [i.e., upgraded or used in derivative classification] in a manner that preserves the overridden original classification binding and provenance.

DELEGATEE | http://terminology.hl7.org/CodeSystem/v3-RoleCode | delegatee | A party to whom some right or authority is granted by a delegator.

DELEGATOR | http://terminology.hl7.org/CodeSystem/v3-RoleCode | delegator | A party that grants all or some portion its right or authority to another party.

DOWNGRDER | http://terminology.hl7.org/CodeSystem/v3-RoleCode | downgrader | An individual authorized to lower the classification level of labeled content and provide rationale for doing so as directed by a classification guide.

DPOWATT | http://terminology.hl7.org/CodeSystem/v3-RoleCode | durable power of attorney | A relationship between two people in which one person authorizes another, usually a family member or relative, to act for him or her in a manner which is a legally binding upon the person giving such authority as if he or she personally were to do the acts that is often limited in the kinds of powers that can be assigned. Unlike ordinary powers of attorney, durable powers can survive for long periods of time, and again, unlike standard powers of attorney, durable powers can continue after incompetency.

EXCEST | http://terminology.hl7.org/CodeSystem/v3-RoleCode | executor of estate | The role played by a person acting as the estate executor for a deceased subscriber or policyholder who was the responsible party

GRANTEE | http://terminology.hl7.org/CodeSystem/v3-RoleCode | grantee | An entity which accepts certain rights or authority from a grantor.

GRANTOR | http://terminology.hl7.org/CodeSystem/v3-RoleCode | grantor | An entity which agrees to confer certain rights or authority to a grantee.

GT | http://terminology.hl7.org/CodeSystem/v3-RoleCode | Guarantor | An individual or organization that makes or gives a promise, assurance, pledge to pay or has paid the healthcare service provider.

GUADLTM | http://terminology.hl7.org/CodeSystem/v3-RoleCode | guardian ad lidem | The role played by a person appointed by the court to look out for the best interests of a minor child during the course of legal proceedings.

HPOWATT | http://terminology.hl7.org/CodeSystem/v3-RoleCode | healthcare power of attorney | A relationship between two people in which one person authorizes another to act for him or her in a manner which is a legally binding upon the person giving such authority as if he or she personally were to do the acts that continues (by its terms) to be effective even though the grantor has become mentally incompetent after signing the document.

INTPRTER | http://terminology.hl7.org/CodeSystem/v3-RoleCode | interpreter | An entity which converts spoken or written language into the language of key participants in an event such as when a provider is obtaining a patient's consent to treatment or permission to disclose information.

POWATT | http://terminology.hl7.org/CodeSystem/v3-RoleCode | power of attorney | A relationship between two people in which one person authorizes another to act for him in a manner which is a legally binding upon the person giving such authority as if he or she personally were to do the acts.

RESPRSN | http://terminology.hl7.org/CodeSystem/v3-RoleCode | responsible party | The role played by a party who has legal responsibility for another party.

SPOWATT | http://terminology.hl7.org/CodeSystem/v3-RoleCode | special power of attorney | A relationship between two people in which one person authorizes another to act for him or her in a manner which is a legally binding upon the person giving such authority as if he or she personally were to do the acts that is often limited in the kinds of powers that can be assigned.

_CitizenRoleType | http://terminology.hl7.org/CodeSystem/v3-RoleCode | CitizenRoleType | A role type used to qualify a person's legal status within a country or nation.

CAS | http://terminology.hl7.org/CodeSystem/v3-RoleCode | asylum seeker | A person who has fled his or her home country to find a safe place elsewhere.

CASM | http://terminology.hl7.org/CodeSystem/v3-RoleCode | single minor asylum seeker | A person who is someone of below legal age who has fled his or her home country, without his or her parents, to find a safe place elsewhere at time of categorization.

CN | http://terminology.hl7.org/CodeSystem/v3-RoleCode | national | A person who is legally recognized as a member of a nation or country, with associated rights and obligations.

CNRP | http://terminology.hl7.org/CodeSystem/v3-RoleCode | non-country member without residence permit | A foreigner who is present in a country (which is foreign to him/her) unlawfully or without the country's authorization (may be called an illegal alien).

CNRPM | http://terminology.hl7.org/CodeSystem/v3-RoleCode | non-country member minor without residence permit | A person who is below legal age present in a country, without his or her parents, (which is foreign to him/her) unlawfully or without the country's authorization.

CPCA | http://terminology.hl7.org/CodeSystem/v3-RoleCode | permit card applicant | A non-country member admitted to the territory of a nation or country as a non-resident for an explicit purpose.

CRP | http://terminology.hl7.org/CodeSystem/v3-RoleCode | non-country member with residence permit | A foreigner who is a resident of the country but does not have citizenship.

CRPM | http://terminology.hl7.org/CodeSystem/v3-RoleCode | non-country member minor with residence permit | A person who is a resident below legal age of the country without his or her parents and does not have citizenship.

AUCG | http://terminology.hl7.org/CodeSystem/v3-ParticipationFunction | caregiver information receiver | Description:Caregiver authorized to receive patient health information.

AULR | http://terminology.hl7.org/CodeSystem/v3-ParticipationFunction | legitimate relationship information receiver | Description:Provider with legitimate relationship authorized to receive patient health information.

AUTM | http://terminology.hl7.org/CodeSystem/v3-ParticipationFunction | care team information receiver | Description:Member of care team authorized to receive patient health information.

AUWA | http://terminology.hl7.org/CodeSystem/v3-ParticipationFunction | work area information receiver | Description:Entities within specified work area authorized to receive patient health information.

PROMSK | http://terminology.hl7.org/CodeSystem/v3-ParticipationFunction | authorized provider masking author | Definition:Provider authorized to mask information to protect the patient, a third party, or to ensure that the provider has consulted with the patient prior to release of this information.

AUT | http://terminology.hl7.org/CodeSystem/v3-ParticipationType | author (originator) | Definition: A party that originates the Act and therefore has responsibility for the information given in the Act and ownership of this Act. Example: the report writer, the person writing the act definition, the guideline author, the placer of an order, the EKG cart (device) creating a report etc. Every Act should have an author. Authorship is regardless of mood always actual authorship. Examples of such policies might include: The author and anyone they explicitly delegate may update the report; All administrators within the same clinic may cancel and reschedule appointments created by other administrators within that clinic; A party that is neither an author nor a party who is extended authorship maintenance rights by policy, may only amend, reverse, override, replace, or follow up in other ways on this Act, whereby the Act remains intact and is linked to another Act authored by that other party.

CST | http://terminology.hl7.org/CodeSystem/v3-ParticipationType | custodian | An entity (person, organization or device) that is in charge of maintaining the information of this act (e.g., who maintains the report or the master service catalog item, etc.).

INF | http://terminology.hl7.org/CodeSystem/v3-ParticipationType | informant | A source of reported information (e.g., a next of kin who answers questions about the patient's history). For history questions, the patient is logically an informant, yet the informant of history questions is implicitly the subject.

IRCP | http://terminology.hl7.org/CodeSystem/v3-ParticipationType | information recipient | A party, who may or should receive or who has recieved the Act or subsequent or derivative information of that Act. Information recipient is inert, i.e., independent of mood." Rationale: this is a generalization of a too diverse family that the definition can't be any more specific, and the concept is abstract so one of the specializations should be used.

LA | http://terminology.hl7.org/CodeSystem/v3-ParticipationType | legal authenticator | A verifier who legally authenticates the accuracy of an act. An example would be a staff physician who sees a patient and dictates a note, then later signs it. Their signature constitutes a legal authentication.

TRC | http://terminology.hl7.org/CodeSystem/v3-ParticipationType | tracker | A secondary information recipient, who receives copies (e.g., a primary care provider receiving copies of results as ordered by specialist).

WIT | http://terminology.hl7.org/CodeSystem/v3-ParticipationType | witness | Only with service events. A person witnessing the action happening without doing anything. A witness is not necessarily aware, much less approves of anything stated in the service event. Example for a witness is students watching an operation or an advanced directive witness.

authserver | http://terminology.hl7.org/CodeSystem/extra-security-role-type | authorization server | An entity providing authorization services to enable the electronic sharing of health-related information based on resource owner's preapproved permissions. For example, an UMA Authorization Server[UMA]

datacollector | http://terminology.hl7.org/CodeSystem/extra-security-role-type | data collector | An entity that collects information over which the data subject may have certain rights under policy or law to control that information's management and distribution by data collectors, including the right to access, retrieve, distribute, or delete that information.

dataprocessor | http://terminology.hl7.org/CodeSystem/extra-security-role-type | data processor | An entity that processes collected information over which the data subject may have certain rights under policy or law to control that information's management and distribution by data processors, including the right to access, retrieve, distribute, or delete that information.

datasubject | http://terminology.hl7.org/CodeSystem/extra-security-role-type | data subject | A person whose personal information is collected or processed, and who may have certain rights under policy or law to control that information's management and distribution by data collectors or processors, including the right to access, retrieve, distribute, or delete that information.

humanuser | http://terminology.hl7.org/CodeSystem/extra-security-role-type | human user | The human user that has participated.

110150 | http://dicom.nema.org/resources/ontology/DCM | Application | Audit participant role ID of software application

110151 | http://dicom.nema.org/resources/ontology/DCM | Application Launcher | Audit participant role ID of software application launcher, i.e., the entity that started or stopped an application

110152 | http://dicom.nema.org/resources/ontology/DCM | Destination Role ID | Audit participant role ID of the receiver of data

110153 | http://dicom.nema.org/resources/ontology/DCM | Source Role ID | Audit participant role ID of the sender of data

110154 | http://dicom.nema.org/resources/ontology/DCM | Destination Media | Audit participant role ID of media receiving data during an export

110155 | http://dicom.nema.org/resources/ontology/DCM | Source Media | Audit participant role ID of media providing data during an import

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information

Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)

Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')

Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application

Definition | An explanation of the meaning of the concept

Comments | Additional notes about how to use the code

4.4.1.617 Value Set http://hl7.org/fhir/ValueSet/payment-type

URL: https://hl7.org/fhir/R4/valueset-payment-type.html
Financial Management Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/payment-type
Version: | 4.0.1
Name: | PaymentTypeCodes
Title: | Payment Type Codes
Definition: | This value set includes sample Payment Type codes.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.641 (for OID based terminology systems)
Copyright: | This is an example set.
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for PaymentTypeCodes

• Resource: PaymentReconciliation.detail.type (CodeableConcept / Example)

4.4.1.617.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/payment-type

4.4.1.617.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 3 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/payment-type version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/payment-type

Code | Display | Definition
payment | Payment | The amount is partial or complete settlement of the amounts due.
adjustment | Adjustment | The amount is an adjustment regarding claims already paid.
advance | Advance | The amount is an advance against future claims.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.618 Value Set http://hl7.org/fhir/ValueSet/endpoint-payload-type

URL: https://hl7.org/fhir/R4/valueset-endpoint-payload-type.html

This value set could not be expanded by the publication tooling: Error from server: Unable to provide support for code system urn:oid:1.3.6.1.4.1.19376.1.2.3

Patient Administration Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/endpoint-payload-type
Version: | 4.0.1
Name: | EndpointPayloadType
Title: | Endpoint Payload Type
Definition: | This is an example value set defined by the FHIR project, that could be used to represent possible payload document types.
Committee: | Patient Administration Work Group
OID: | 2.16.840.1.113883.4.642.3.496 (for OID based terminology systems)
Copyright: | Some content from IHE® Copyright © 2015 IHE International, Inc. This content is from the IHE Technical Frameworks and Supplements, available for free download and use at http://www.ihe.net/Technical_Frameworks/
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: Endpoint.payloadType (CodeableConcept / Example)

4.4.1.618.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/endpoint-payload-type

urn:oid:1.3.6.1.4.1.19376.1.2.3

Code	Display
urn:ihe:pcc:handp:2008	History and Physical Specification
urn:ihe:pcc:xphr:2007	HL7 CCD Document
urn:ihe:pcc:aps:2007	IHE Antepartum Summary
urn:ihe:pcc:xds-ms:2007	XDS Medical Summaries
urn:ihe:pcc:xphr:2007	Personal Health Records
urn:ihe:pcc:edr:2007	Emergency Department Referral (EDR)
urn:ihe:pcc:edes:2007	Emergency Department Encounter Summary (EDES)
urn:ihe:pcc:apr:handp:2008	Antepartum Record (APR) - History and Physical
urn:ihe:pcc:apr:lab:2008	Antepartum Record (APR) - Laboratory
urn:ihe:pcc:apr:edu:2008	Antepartum Record (APR) - Education
urn:ihe:pcc:irc:2008	Immunization Registry Content (IRC)
urn:ihe:pcc:crc:2008	Cancer Registry Content (CRC)
urn:ihe:pcc:cm:2008	Care Management (CM)
urn:ihe:pcc:ic:2009	Immunization Content (IC)
urn:ihe:pcc:tn:2007	PCC TN
urn:ihe:pcc:nn:2007	PCC NN
urn:ihe:pcc:ctn:2007	PCC CTN
urn:ihe:pcc:edpn:2007	PCC EDPN
urn:ihe:pcc:hp:2008	PCC HP
urn:ihe:pcc:ldhp:2009	PCC LDHP
urn:ihe:pcc:lds:2009	PCC LDS
urn:ihe:pcc:mds:2009	PCC MDS
urn:ihe:pcc:nds:2010	PCC NDS
urn:ihe:pcc:ppvs:2010	PCC PPVS
urn:ihe:pcc:trs:2011	PCC TRS
urn:ihe:pcc:ets:2011	PCC ETS
urn:ihe:pcc:its:2011	PCC ITS
urn:ihe:iti:bppc:2007	Basic Patient Privacy Consents
urn:ihe:iti:bppc-sd:2007	Basic Patient Privacy Consents with Scanned Document
urn:ihe:iti:xdw:2011:workflowDoc	XDW Workflow Document
urn:ihe:iti:dsg:detached:2014	DSG Detached Document
urn:ihe:iti:dsg:enveloping:2014	DSG Enveloping Document
urn:ihe:iti:xds-sd:pdf:2008	PDF embedded in CDA per XDS-SD profile
urn:ihe:iti:xds-sd:text:2008	Text embedded in CDA per XDS-SD profile
urn:ihe:lab:xd-lab:2008	CDA Laboratory Report
urn:ihe:rad:TEXT	Radiology XDS-I Text
urn:ihe:rad:PDF	Radiology XDS-I PDF
urn:ihe:rad:CDA:ImagingReportStructuredHeadings:2013	Radiology XDS-I Structured CDA
urn:ihe:card:imaging:2011	Cardiac Imaging Report
urn:ihe:card:CRC:2012	Cardiology CRC
urn:ihe:card:EPRC-IE:2014	Cardiology EPRC-IE
urn:ihe:dent:TEXT	Dental Text
urn:ihe:dent:PDF	Dental PDF
urn:ihe:dent:CDA:ImagingReportStructuredHeadings:2013	Dental CDA
urn:ihe:pat:apsr:all:2010	Anatomic Pathology Structured Report All
urn:ihe:pat:apsr:cancer:all:2010	Anatomic Pathology Structured Report Cancer All
urn:ihe:pat:apsr:cancer:breast:2010	Anatomic Pathology Structured Report Cancer Breast
urn:ihe:pat:apsr:cancer:colon:2010	Anatomic Pathology Structured Report Cancer Colon
urn:ihe:pat:apsr:cancer:prostate:2010	Anatomic Pathology Structured Report Cancer Prostate
urn:ihe:pat:apsr:cancer:thyroid:2010	Anatomic Pathology Structured Report Cancer Thyroid
urn:ihe:pat:apsr:cancer:lung:2010	Anatomic Pathology Structured Report Cancer Lung
urn:ihe:pat:apsr:cancer:skin:2010	Anatomic Pathology Structured Report Cancer Skin
urn:ihe:pat:apsr:cancer:kidney:2010	Anatomic Pathology Structured Report Cancer Kidney
urn:ihe:pat:apsr:cancer:cervix:2010	Anatomic Pathology Structured Report Cancer Cervix
urn:ihe:pat:apsr:cancer:endometrium:2010	Anatomic Pathology Structured Report Cancer Endometrium
urn:ihe:pat:apsr:cancer:ovary:2010	Anatomic Pathology Structured Report Cancer Ovary
urn:ihe:pat:apsr:cancer:esophagus: 2010	Anatomic Pathology Structured Report Cancer Esophagus
urn:ihe:pat:apsr:cancer:stomach: 2010	Anatomic Pathology Structured Report Cancer Stomach
urn:ihe:pat:apsr:cancer:liver:2010	Anatomic Pathology Structured Report Cancer Liver
urn:ihe:pat:apsr:cancer:pancreas: 2010	Anatomic Pathology Structured Report Cancer Pancreas
urn:ihe:pat:apsr:cancer:testis:2010	Anatomic Pathology Structured Report Cancer Testis
urn:ihe:pat:apsr:cancer:urinary_bladder:2010	Anatomic Pathology Structured Report Cancer Urinary Bladder

urn:ihe:pat:apsr:cancer:lip_oral_cavity:2010 | Anatomic Pathology Structured Report Cancer Lip Oral Cavity
urn:ihe:pat:apsr:cancer:pharynx:2010 | Anatomic Pathology Structured Report Cancer Pharynx
urn:ihe:pat:apsr:cancer:salivary_gland:2010 | Anatomic Pathology Structured Report Cancer Salivary Gland
urn:ihe:pat:apsr:cancer:larynx:2010 | Anatomic Pathology Structured Report Cancer Larynx
urn:ihe:pharm:pre:2010 | Pharmacy Pre
urn:ihe:pharm:padv:2010 | Pharmacy PADV
urn:ihe:pharm:dis:2010 | Pharmacy DIS
urn:ihe:pharm:pml:2013 | Pharmacy PML
urn:hl7-org:sdwg:ccda-structuredBody:1.1 | For documents following C-CDA constraints using a structured body.
urn:hl7-org:sdwg:ccda-nonXMLBody:1.1 | For documents following C-CDA constraints using a non structured body.

4.4.1.618.2 Expansion

This expansion generated 01 Nov 2019

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.619 Value Set http://hl7.org/fhir/ValueSet/device-kind

URL: https://hl7.org/fhir/R4/valueset-device-kind.html
Orders and Observations Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/device-kind
Version: | 4.0.1
Name: | FHIRDeviceTypes
Title: | FHIR Device Types
Definition: | Codes used to identify medical devices. Includes concepts from SNOMED CT (http://www.snomed.org/) where concept is-a 49062001 (Device) and is provided as a suggestive example.
Committee: | Orders and Observations Work Group
OID: | 2.16.840.1.113883.4.642.3.208 (for OID based terminology systems)
Copyright: | This resource includes content from SNOMED Clinical Terms® (SNOMED CT®) which is copyright of the International Health Terminology Standards Development Organisation (IHTSDO). Implementers of these specifications must have the appropriate SNOMED CT Affiliate license - for more information contact http://www.snomed.org/snomed-ct/get-snomed-ct or info@snomed.org
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: ChargeItem.product[x] (Reference(Device|Medication|Substance)|CodeableConcept / Example)

• Resource: DeviceRequest.code[x] (Reference(Device)|CodeableConcept / Example)

• Resource: Procedure.usedCode (CodeableConcept / Example)

• Resource: DeviceDefinition.type (CodeableConcept / Example)

• Extension: http://hl7.org/fhir/StructureDefinition/observation-deviceCode: deviceCode (CodeableConcept / Example)

4.4.1.619.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include codes from http://snomed.info/sct where concept is-a 49062001 (Device)

4.4.1.619.2 Expansion

This expansion generated 01 Nov 2019

This value set has >1000 codes in it. In order to keep the publication size manageable, only a selection (1000 codes) of the whole set of codes is shown

All codes from system http://snomed.info/sct

Code | Display | Definition
156009 | Spine board
271003 | Bone plate

287000 | Air receiver
291005 | Atomizer
678001 | Epilator
739006 | Bicycle ergometer
793009 | Mechanical power press
882002 | Diagnostic implant
972002 | Air filter, device
989005 | Linen cloth
994005 | Brush, device
1211003 | Treadmill, device
1333003 | Emesis basin, device
1422002 | Plastic mold, device
1579007 | Surgical drill, device
1766001 | Toboggan, device
1941006 | Silk cloth
1962007 | Dike, device
2248009 | Tracheal tube cuff
2282003 | Breast implant
2287009 | Nail file, device
2468001 | Breath analyzer, device
2478003 | Ocular prosthesis
2491002 | Intra-aortic balloon catheter, device
3201004 | Tent, device
3319006 | Artificial liver, device
4408003 | Endoscopic camera, device
4632004 | Hair cloth
4816004 | Metal device
5041003 | Adhesive strip, device
5042005 | Patient scale, device
5159002 | Physiologic monitoring system, device
5679009 | Bed sheet
6012004 | Hearing aid
6097006 | T-tube
6822006 | Microwave oven
6919005 | Protective clothing material
6972009 | Lithotripter
7007007 | Radiographic-fluoroscopic unit
7402007 | Probe
7406005 | Crib
7704007 | Stabilizing appliance
7733008 | Hydrocephalic shunt catheter
7968002 | Three-wheeled all-terrain vehicle
7971005 | Fogarty catheter
8060009 | Denture
8118007 | Crane
8170008 | Adhesive
8384009 | Band saw
8407004 | Bile collection bag
8434001 | Gaol
8451008 | Intramedullary nail
8615009 | Blood electrolyte analyzer
8643000 | Mortising machine
8682003 | Protective shield
9017009 | Ventricular intracranial catheter
9096001 | Support
9129003 | Feeding catheter
9419002 | Bobsled
9458007 | Elastic bandage
9611009 | Dermatotome
9883003 | Cargo handling gear
10244001 | Needle guide
10371004 | Electrostimulating analgesia unit
10507000 | Toeboard
10826000 | Industrial saw
10850003 | Radiographic-therapeutic unit
10906003 | Vein stripper
11141007 | Bone growth stimulator
11158002 | Electromyographic monitor and recorder
11358008 | Prosthetic valve
11987000 | Clinical chemistry analyzer
12150006 | Cannula
12183004 | Upper limb prosthesis

12198002 | Ice skate
12953007 | File
13118005 | Wool cloth
13219008 | Gastroscope
13288007 | Monitors
13459008 | Temporary artificial heart prosthesis
13764006 | Uterine sound
13855007 | Pillow
13905003 | Tennis ball
14106009 | Cardiac pacemaker implant
14108005 | Cage
14116001 | Analgesia unit
14208000 | Oil well
14288003 | Nasal septum button
14339000 | Button
14364002 | Camera
14423008 | Adhesive bandage
14519003 | Aspirator
14548009 | Harrington rod
14762000 | Alloy steel chain sling
14789005 | Prosthetic implant
15000008 | Air-conditioner
15340005 | Wood's light
15447007 | Arthroplasty prosthesis
15644007 | Anesthesia unit
15869005 | Dosimeter, device
15873008 | Boiler, device
15922004 | Gown, device
16056004 | Boots
16349000 | Orthopedic equipment
16417001 | Commercial breathing supply hoses diving operation, device
16470007 | Electrode, device
16497000 | Electric clipper, device
16524003 | Cotton cloth
16540000 | Umbrella catheter, device
16650009 | Splint, device
17102003 | NG - Nasogastric tube
17107009 | Prosthetic mitral valve
17207004 | Mattress, device
17306006 | Hernia belt, device
17404008 | Cardiac compression board, device
17472008 | Knife, device
18151003 | Punch, device
18153000 | Fluorescence immunoassay analyzer, device
18411005 | Chisel, device
18666004 | Finespun glass
19257004 | Defibrillator, device
19328000 | Blanket, device
19443004 | Radioactive implant, device
19627002 | Leather
19762002 | Leather belt
19817005 | Fan blade, device
19892000 | Scale, device
19923001 | Catheter, device
20195009 | Leg prosthesis, device
20235003 | Toothbrush, device
20273004 | Industrial machine, device
20359006 | Contraceptive diaphragm, device
20406008 | Back rests, device
20428008 | Oxygen tent, device
20513005 | Power tool, device
20568009 | Urinary catheter, device
20613002 | Cystoscope, device
20861007 | Plug pack, device
20867006 | Experimental implant, device
20873007 | Plastic cloth-like material
20997002 | Hand tool, device
21079000 | Carbon monoxide analyzer, device
21546008 | Icebox
21870002 | Transluminal extraction catheter, device
21944004 | Abdominal binder, device
22251003 | Timer, device

22283009 | Artificial membrane, device
22566001 | Cytology brush, device
22662007 | Retaining harness, device
22679001 | Handcuffs, device
22744006 | Artificial hair wig, device
22852002 | Therapeutic implant, device
23228005 | Arthroscope, device
23366006 | Motorized wheelchair device
23699001 | Baseball, device
23785007 | Arthroscopic irrigation/distension pump, device
23973005 | Indwelling urinary catheter, device
24073000 | Mechanical cardiac valve prosthesis
24110008 | Anoscope, device
24174009 | Bronchoscope, device
24230000 | Vibrator, device
24290003 | Cardiac valve bioprosthesis
24402003 | Stepladder, device
24470005 | Wrench, device
24513003 | Plastic boots
24697008 | Ostomy belt, device
24767007 | Eustachian tube prosthesis, device
25005004 | Snare, device
25062003 | Feeding tube, device
25152007 | Squeeze cage, device
25510005 | Heart valve prosthesis
25632005 | Hockey puck, device
25680008 | Scaffold, device
25742001 | Orthodontic appliance, device
25937001 | Neurostimulation device
26128008 | Bougie, device
26239002 | Soccer ball, device
26334009 | Dockboard, device
26397000 | Reservoir bag
26412008 | ET - Endotracheal tube
26579007 | Holter valve, device
26719000 | Celestin tube, device
26882005 | Rongeur, device
27042007 | Needle adapter, device
27065002 | Suture
27091001 | Dumbwaiter, device
27126002 | Power belt, device
27229001 | Spray booth, device
27606000 | Dental prosthesis, device
27785006 | Athletic supporter, device
27812008 | Electric heating pad, device
27976001 | Woodworking machinery, device
27991004 | Thermometer, device
28026003 | Hairbrush, device
29292008 | Fur garment
29319002 | Forceps, device
29396008 | Resuscitator, device
30012001 | Elevator, device
30070001 | Multistage suspension scaffolding, device
30115002 | Shield, device
30176005 | Baseball bat, device
30234008 | Medical laboratory analyzer, device
30610008 | Epidural catheter, device
30661003 | Cosmetic prosthesis, device
30929000 | Ligator, device
30968007 | Drainage bag, device
31030004 | Peritoneal catheter, device
31031000 | Internal fixator
31174004 | Lumbosacral belt, device
31791005 | Traction belt, device
31878003 | Surgical scissors, device
32033000 | Arterial pressure monitor, device
32356002 | Machine guarding, device
32504006 | Screwdriver, device
32634007 | Fixed ladder, device
32667006 | Oral airway
32711007 | Ostomy collection bag, device
32712000 | Drain, device

32871007 | Tweezer, device
33194000 | Welding equipment, device
33352006 | Ax, device
33388001 | Carbon dioxide analyzer, device
33482001 | Rubber boots
33686008 | Stylet, device
33690005 | Sharp instrument, device
33802005 | Enema bag, device
33894003 | Experimental device
33918000 | Rubberized cloth
34164001 | POP - Plaster of Paris cast
34188004 | Straightjacket, device
34234003 | Plastic tube, device
34263000 | Medical balloon, device
34362008 | Vascular device
34759008 | Urethral catheter, device
35398009 | Ostomy appliance, device
35593004 | Wire ligature, device
35870000 | Cerebrospinal catheter, device
36365007 | Ice-pick, device
36370000 | Aspirator trap bottle, device
36645008 | Stimulator, device
36761001 | Natural hair wig, device
36965003 | Hemodialysis machine, device
36977008 | Peripheral nerve stimulator
37189001 | Magnetic detector, device
37270008 | Endoscope, device
37284003 | Bag, device
37311003 | Stone retrieval basket, device
37347002 | Dildo, device
37360008 | Patient isolator, device
37503007 | Protective blind, device
37759000 | Surgical instrument, device
37874008 | Continuing positive airway pressure unit, device
37953008 | Bedside rails, device
38126007 | Protective lenses
38141007 | Tourniquet, device
38277008 | Protective device
38806006 | Hockey stick, device
38862006 | Sheet metal bending equipment
38871002 | Metallic cloth
39590006 | Air compressor, device
39690000 | Sphygmomanometer, device
39768008 | Rasp, device
39790008 | Non-electric heating pad, device
39802000 | Tongue blade, device
39821008 | Positron emission tomography unit, device
39849001 | Nasal oxygen cannula
39869006 | Alarm, device
40388003 | Biomedical implant
40519001 | Binder, device
41157002 | Orthopedic immobilizer
41323003 | Urinary collection bag, device
41525006 | Artificial structure, device
41684000 | Industrial tool, device
42152006 | Metal tube, device
42305009 | Ambulation device
42380001 | Ear plug, device
42451009 | Blood warmer, device
42716000 | Wool
42882002 | Hypodermic spray, device
43001000 | Phlebotomy kit, device
43192004 | Bone pencil, device
43252007 | Cochlear implant
43725001 | Airway equipment
43734006 | Blood administration set, device
43770009 | Doppler device
43983001 | Shoes
44056008 | Caliper, device
44396004 | Magnet, device
44492001 | Industrial robot, device
44668000 | Pump, device

44738004 | Laryngoscope, device
44806002 | Esophageal bougie, device
44845007 | Golf ball, device
44907005 | Four-wheeled all-terrain vehicle, device
44959004 | Angioplasty balloon catheter, device
45633005 | Peritoneal dialyzer, device
45901004 | Penrose drain, device
46181005 | Automatic fire extinguisher system, device
46265007 | Artificial lashes, device
46299005 | Sanitary belt, device
46364009 | Clamp, device
46440007 | Basketball, device
46625003 | Suppository
46666003 | Chain, device
46949002 | Deck machinery, device
47162009 | Mirror, device
47326004 | Electrical utilization equipment, device
47424002 | Apgar scoring timer, device
47528002 | Ureteric catheter
47731004 | Birthing chair, device
47734007 | Chromic catgut suture
47776004 | Mittens
47942000 | Proctoscope, device
48066006 | Circular portable saw, device
48096001 | Bathtub rails, device
48240003 | Training equipment, device
48246009 | Studgun, device
48295009 | Vascular filter, device
48473008 | Protective body armor, device
48822005 | Bilirubin light, device
48990009 | Strap, device
49448001 | Razor, device
49890001 | Hospital cart, device
50121007 | Glasses
50457005 | Workover rig service to oil well, device
50483000 | Oil rig
50851003 | Penile tumescence monitor, device
51016001 | Hammer, device
51086006 | Shower curtain, device
51324004 | Stripper, device
51685009 | Roller skate, device
51791000 | Measuring tape, device
51832004 | Valved tube, device
51883004 | Sling, device
52124006 | Central line
52161002 | Molten lava
52291003 | Gloves
52520009 | Ladder, device
52537002 | Aspirator collection canister, device
52624007 | Radiofrequency generator, device
52773007 | Ski, device
52809000 | Nasopharyngeal catheter, device
52893008 | Blood gas/pH analyzer, device
53167006 | Platform suspended boom, device
53177008 | Nasal balloon, device
53217009 | Artificial lung, device
53350007 | Prosthesis, device
53535004 | Retractor, device
53639001 | Stethoscope, device
53671008 | Gastric balloon, device
53996008 | Penile prosthesis, device
54234007 | Cryogenic analgesia unit, device
54638004 | Towel, device
54953005 | Computerized axial tomography scanner, device
55091003 | Blood coagulation analyzer, device
55206006 | Suture button, device
55216003 | Amnioscope, device
55337009 | Auscultoscope, device
55567004 | Bassinet, device
55658008 | Hot object
55986002 | Tong, device
56144002 | Back braces, device

56353002 | Staple, device
56547001 | Trephine, device
56896002 | Pacemaker catheter, device
56961003 | Cardiac transvenous pacemaker, device
57118008 | Perfusion pump, device
57134006 | Clinical instrument
57368009 | Contact lens
57395004 | Physical restraint equipment, device
57730005 | Abrasive grinding, device
58153004 | Android, device
58253008 | Suction catheter, device
58514003 | Infant scale, device
58878002 | Protective vest, device
58938008 | WC - Wheelchair
59102007 | Ice bag, device
59127000 | Apnea alarm, device
59153008 | Barge, device
59160002 | Chipguard, device
59181002 | Oxygen analyzer, device
59432006 | Ligature, device
59746007 | Needle holder, device
59772003 | Culdoscope, device
59782002 | Speculum, device
59833007 | Collapsible balloon, device
60054005 | SB - Seat belt
60110001 | Wig, device
60150003 | Skipole, device
60161006 | Acupuncture needle, device
60185003 | Carbon dioxide absorber, device
60311007 | Leather boots
60773001 | Injector
60806001 | Whirlpool bath
60957001 | Otoscope
61330002 | Nasopharyngeal airway device
61512008 | Tennis racket
61835000 | Dilator
61968008 | Syringe
61979003 | Antiembolic device
62336005 | Electric cable
62495008 | Gamma counter
62614002 | Overhead and gantry crane
62980002 | Tubular bandage
63112008 | Bone wire
63173005 | Hat band
63289001 | Metal nail
63336000 | Bone plug
63548003 | Derrick
63562005 | Cervical collar
63619003 | Fiberoptic cable
63653004 | Medical device
63797009 | Traction unit
63839002 | Electroejaculator
63995005 | Bandage
64174005 | Snowmobile
64255007 | Esophageal balloon
64565002 | Air tool
64571008 | Hair clipper
64883003 | Inhalation analgesia unit
64973003 | Scissors
64989000 | Escalator
65053001 | Electrical battery
65105002 | Surgical drapes
65268008 | Chart recorder
65473004 | Microscope
65577000 | X-ray shield
65818007 | Stent
66222000 | Hospital robot
66415006 | Audiometric testing equipment
66435007 | Electric bed
66494009 | Face cloth
67270000 | Hip prosthesis
67387001 | Coronary perfusion catheter

67670006 | Radiographic-tomographic unit
67777003 | Moving walk
67829007 | Esophagoscope
67920005 | Aerial lift
67966000 | Enema tube
68080007 | Radiographic unit
68181008 | Vibrating electric heating pad
68183006 | Bone screw
68276009 | Bottle
68325009 | Sound
68597009 | Support belt
68685003 | Household robot
68842005 | Gastroduodenoscope
69670004 | Patient utensil kit
69805005 | Insulin pump
69922008 | Tracheostomy button
70080007 | Bayonet
70300000 | Skull tongs
70303003 | Freezer
70453008 | Sled
70665002 | Blood pressure cuff
70793005 | Recreation equipment
70872004 | Wash basin
71384000 | Warmer
71483007 | Diving stage
71545009 | Humidifier
71601002 | Proctosigmoidoscope
71667001 | Bone wax
71948003 | Autoclave
72070000 | Ring
72188006 | Tissue expander
72302000 | Lead cable
72506001 | Implantable defibrillator
72742007 | Aspirator collection bottle
73027007 | Infant incubator
73534004 | Artificial skin
73562006 | Transilluminator
73571002 | Intravenous analgesia unit
73618007 | Power saw
73878004 | Hand saw
73985004 | Face protection in construction industry
74094004 | Belt
74108008 | Recorder
74300007 | Sanitary pad
74444006 | AL - Artificial limb
74566002 | Crutch
74884005 | Boatswain's chair
75075000 | Shoring and bracing (masonry and woodwork)
75187009 | Local anesthesia kit
75192006 | Arterial cannula
75751006 | Manual respirator
75780002 | Artificial kidney
75963008 | Skateboard
76091005 | Stainless steel wire suture
76123001 | Glass tube
76428000 | Elbow joint prosthesis
76433001 | Apron
76664007 | Artificial pancreas
76705002 | Applicator stick
76825006 | Abrasive wheel machinery
76937009 | Guillotine
77444004 | Pins
77541009 | Band
77720000 | Clips
77755003 | Chemical fiber cloth
77777004 | Bone staple
78279003 | Nail clipper
78498003 | Testicular prosthesis
78641001 | Nylon suture
78886001 | Electronic monitor
79051006 | Greenfield filter
79068005 | Needle

79218005 | Vehicle-mounted work platform
79287008 | Tampon
79401009 | Chute
79438009 | Foot protection
79481007 | Swing or sliding cut-off saw
79593001 | Transvenous electrode
79618001 | Storage tank
79811009 | Electric blanket
79834000 | Hickman line
79952001 | Swan-Ganz catheter, device
80278003 | Pediatric bed
80404000 | Chain fall
80617005 | Analysers
80664005 | Motor home
80853009 | Tendon hammer
80950008 | Oven
81293006 | Textile material
81317009 | Socket wrench
81320001 | Enzyme immunoassay analyzer
81826000 | All-terrain vehicle
81892008 | Radial saw
82379000 | Hemostat
82449006 | Peripheral intravenous catheter
82657000 | Bony tissue forceps
82830000 | Robotic arm
82879008 | Safety belt
83059008 | Tube
83315005 | Audio analgesia unit
83320005 | Dip tank
83369007 | Plastic shoes
83517001 | Robot
83903000 | Man lift
84023008 | Ski tow
84330009 | Pliers
84444002 | Chain saw
84546002 | Barricade
84599008 | Detonating cord
84610002 | Implantable dental prosthesis
84683006 | Aortic valve prosthesis
84756000 | Adhesive tape
85106006 | Boring machine
85329008 | Abortion pump
85455005 | Cart
85684007 | Engraving press
86056006 | Golf club
86174004 | Laparoscope
86184003 | Electrocardiographic monitor and recorder
86407004 | Table
86453006 | Defibrillator paddle
86572008 | Arteriovenous shunt catheter
86768006 | Balloon pump
86816008 | Diving ladder
86967005 | Tool
87088005 | Soldering iron
87140005 | Clothing material
87405001 | Cane
87710003 | Physical restraint structure
87717000 | Tester
87851008 | Blood cell counter and analyzer
88063004 | Footwear
88168006 | Maximum security cell
88208003 | Intravenous anesthesia administration set
88765001 | Artificial tissue
88959008 | Hypodermic needle
89149003 | Stretcher
89236003 | Leather shoes
89509004 | Blood culture analyzer
90003000 | Magnetic resonance imaging unit
90035000 | Alcohol sponge
90082007 | Cast cutter
90134004 | Metal periosteal implant
90412006 | Colonoscope

90504001 | Auricular prosthesis
90913005 | Rubber shoes
90948003 | Abrasive blast by cleaning nozzles
91294003 | Thomas collar
91318002 | Hyperbaric chamber
91535004 | Basin
91537007 | Hospital bed
102303004 | Vascular prosthesis
102304005 | Measuring ruler
102305006 | Intramedullary reamer
102306007 | Reamer
102307003 | Surgical knife
102308008 | Scalpel
102309000 | Surgical saw
102310005 | Gigli's wire saw
102311009 | Starck dilator
102312002 | Atherectomy device
102313007 | Rotational atherectomy device
102314001 | Embolization coil
102315000 | Embolization ball
102316004 | Embolization particulate
102317008 | Guiding catheter
102318003 | Implantable venous catheter
102319006 | Percutaneous transluminal angioplasty balloon
102320000 | Detachable balloon
102321001 | Operating microscope
102322008 | External prosthesis for sonographic procedure
102323003 | Water bag prosthesis for imaging procedure
102324009 | Saline bag prosthesis for imaging procedure
102325005 | Gel prosthesis for imaging procedure
102326006 | Dagger
102327002 | Dirk
102328007 | Sword
102384007 | Motor vehicle airbag
102385008 | Front airbag
102386009 | Front driver airbag
102387000 | Front passenger airbag
102388005 | Side airbag
102402008 | Snowboard
102403003 | Water ski
105784003 | Life support equipment
105785002 | Adhesive, bandage AND/OR suture
105787005 | Belt AND/OR binder
105788000 | Probe, sound, bougie AND/OR airway
105789008 | Cannula, tube AND/OR catheter
105790004 | Bag, balloon AND/OR bottle
105791000 | Pump, injector AND/OR aspirator
105792007 | Analgesia AND/OR anesthesia unit
105793002 | Monitor, alarm AND/OR stimulator
105794008 | Scope AND/OR camera
105809003 | Physical restraint equipment AND/OR structure
108874005 | Silicone plug
109184000 | Pregnancy testing kit
109226007 | Dental pin
109227003 | Hand joint prosthesis
109228008 | Knee joint prosthesis
111041008 | Artificial nails
111042001 | Artificial organ
111043006 | Medical tuning fork
111044000 | Bone tap
111045004 | Exerciser
111047007 | Urethral bougie
111048002 | Rhinoscope
111052002 | Protective breast plate
111060001 | Industrial sewing machine
111062009 | Food waste disposal equipment
115961006 | Soft Cast
115962004 | Fiberglass cast
116146000 | Blood product unit
116204000 | Catheter tip
116205004 | Blood bag
116206003 | Plasma bag

116250002 | Filter
116251003 | Wick
118294000 | Solid-state laser
118295004 | Gas laser
118296003 | Chemical laser
118297007 | Excimer laser
118298002 | Dye laser
118299005 | Diode laser
118301003 | Nd:YVO>4< laser
118302005 | Nd:YLF laser
118303000 | Nd:Glass laser
118304006 | Chromium sapphire laser device
118305007 | Er:Glass laser
118306008 | Erbium-YAG laser
118307004 | Ho:YLF laser
118308009 | Holmium-YAG laser
118309001 | Ti:sapphire laser device
118310006 | Alexandrite laser
118311005 | Argon laser
118312003 | CO2 laser
118313008 | He laser
118314002 | Helium cadmium laser
118315001 | HeNe laser
118316000 | Krypton laser
118317009 | Neon gas laser
118318004 | Nitrogen gas laser
118319007 | Xenon gas laser
118320001 | Copper vapor laser
118321002 | Gold vapor laser
118322009 | DF laser
118323004 | DF-CO>2< laser device
118324005 | HF laser
118325006 | ArF laser
118326007 | KrF laser
118327003 | KrCl laser
118328008 | XeCl laser
118329000 | XeFl laser
118330005 | Free electron laser
118331009 | Tunable dye laser
118332002 | Tunable dye argon laser
118333007 | Gallium arsenide laser
118334001 | Gallium aluminum arsenide laser
118335000 | Lead-salt laser
118336004 | Rhodamine 6G dye laser
118337008 | Coumarin C30 dye laser
118338003 | Coumarin 102 dye laser
118342000 | Diode pumped laser
118343005 | Flashlamp pumped laser device
118346002 | Pulsed dye laser
118347006 | QS laser
118348001 | Flashlamp pulsed dye laser
118349009 | CW CO>2< laser
118350009 | High energy pulsed CO>2< laser
118351008 | Frequency doubled Nd:YAG laser
118354000 | Continuous wave laser
118355004 | Pulsed laser
118356003 | Metal vapor laser
118357007 | KTP laser
118371004 | Ion laser
118372006 | Plastic implant
118373001 | Silastic implant
118374007 | Silicone implant
118375008 | Cardiac septum prosthesis
118376009 | Thermocouple
118377000 | Biopsy needle
118378005 | Pacemaker pulse generator
118379002 | Automatic implantable cardioverter sensing electrodes
118380004 | Implantable defibrillator leads
118381000 | Implantable cardioverter leads
118382007 | Neuropacemaker device
118383002 | External fixation device
118384008 | Long arm splint

118385009 | Short arm splint
118386005 | Figure of eight plaster cast
118387001 | Halo jacket
118388006 | Body cast, shoulder to hips
118389003 | Body cast, shoulder to hips including head, Minerva type
118390007 | Body cast, shoulder to hips including one thigh
118391006 | Body cast, shoulder to hips including both thighs
118392004 | Shoulder cast
118393009 | Long arm cylinder
118394003 | Forearm cylinder
118396001 | Cylinder cast, thigh to ankle
118397005 | Long leg cast
118398000 | Long leg cast, walker or ambulatory type
118399008 | Long leg cast, brace type
118400001 | Short leg cast below knee to toes
118401002 | Short leg cast below knee to toes, walking or ambulatory type
118402009 | Clubfoot cast
118403004 | Clubfoot cast, short leg
118404005 | Clubfoot cast, long leg
118405006 | Spica cast
118406007 | Hip spica cast, both legs
118407003 | Hip spica cast, one leg
118408008 | Hip spica cast, one and one-half spica
118409000 | Patellar tendon bearing cast
118410005 | Boot cast
118411009 | Sugar tong cast
118412002 | Gauntlet cast
118413007 | Complete cast
118414001 | Pressure dressing
118415000 | Packing material
118416004 | Wound packing material
118418003 | Trocar
118419006 | Umbrella device
118420000 | Atrial septal umbrella
118421001 | King-Mills umbrella device
118422008 | Mobitz-Uddin umbrella device
118423003 | Rashkind umbrella device
118424009 | Reservoir device
118425005 | Ventricular reservoir
118426006 | Ommaya reservoir
118427002 | Rickham reservoir
118428007 | Flexible fiberoptic endoscope
118429004 | Flexible fiberoptic laryngoscope with strobe
118643004 | Cast
122456005 | Laser
123636009 | SS - Silk suture
126064005 | Gastrostomy tube, device
126065006 | Jejunostomy tube, device
128981007 | Baffle
129113006 | Intra-aortic balloon pump
129121000 | Tracheostomy tube
129247000 | Fine biopsy needle
129248005 | Core biopsy needle
129460009 | Compression paddle
129462001 | Catheter guide wire
129463006 | J wire
129464000 | Medical administrative equipment
129465004 | Medical record
129466003 | Patient chart
129467007 | Identification plate
134823007 | Sterile absorbent dressing pad
134963007 | Wound drainage pouch dressing
170615005 | Home nebulizer
182562006 | Plaster jacket
182563001 | Shoulder spica
182564007 | Humeral U-slab
182565008 | Long arm slab
182566009 | Humeral hanging slab
182567000 | Forearm slab
182568005 | Scaphoid cast
182569002 | Bennett cast
182570001 | Hip spica

182571002 | Long leg spica
182572009 | Below knee non-weight-bearing cast
182573004 | Below knee weight-bearing cast
182574005 | Plaster stripper
182576007 | Humeral brace
182577003 | Functional elbow brace
182578008 | Forearm brace
182579000 | Hip brace
182580002 | Femoral brace
182581003 | Tibial brace
182587004 | Body support
182588009 | Spinal frame
182589001 | Corset support
182590005 | Cardiac bed
182591009 | Water bed
182592002 | High air loss bed
182839003 | Automated drug microinjector
183116000 | Dental aid
183125006 | Ear fitting hearing aid
183135000 | Mobility aid
183141007 | Inshoe orthosis
183143005 | Surgical stockings
183144004 | Antiembolic stockings
183146002 | ZF - Zimmer frame
183147006 | Tripod
183148001 | RGO - Reciprocating gait orthosis
183149009 | Hip guidance orthosis
183150009 | Standing frame
183152001 | Hip abduction orthosis
183153006 | Hip-knee-ankle-foot orthosis
183154000 | Knee-ankle-foot orthosis
183155004 | Flexible knee support
183156003 | Collateral ligament brace
183157007 | Anterior cruciate ligament brace
183158002 | Posterior cruciate ligament brace
183159005 | Ground reaction orthosis
183160000 | Rigid ankle-foot orthosis
183161001 | Flexible ankle-foot orthosis
183162008 | Double below-knee iron
183164009 | Inside iron
183165005 | Outside iron
183166006 | Inside T-strap
183170003 | Hindquarter prosthesis
183171004 | Hip disarticulation prosthesis
183172006 | Above knee prosthesis
183173001 | Through knee prosthesis
183174007 | Below knee prosthesis
183175008 | Syme's prosthesis
183176009 | Midfoot amputation prosthesis
183177000 | Shoe filler
183183002 | Milwaukee brace
183184008 | Boston brace
183185009 | Jewett brace
183187001 | Halo device
183188006 | Four poster brace
183189003 | Rigid collar
183190007 | Flexible collar
183192004 | Shoulder abduction brace
183193009 | Elbow brace
183194003 | Passive wrist extension splint
183195002 | Active wrist extension splint
183196001 | Passive finger extension splint
183197005 | Active finger extension splint
183198000 | Kleinert traction
183199008 | Passive thumb splint
183200006 | Active thumb splint
183202003 | Shin splint
183204002 | Excretory control aid
183235008 | Facial non-surgical prosthesis
183236009 | Breast non-surgical prosthesis
183240000 | Patient-propelled wheelchair
183241001 | Pedal powered wheelchair

183248007 | Attendant powered wheelchair
183249004 | Wheelchair seating
183250004 | Molded wheelchair seat
183251000 | Matrix seat
201706006 | Intracranial pressure transducer
223394001 | Equipment for positioning
224684009 | Top security prison
224685005 | Category B prison
224686006 | Low security prison
224823002 | Street lighting
224824008 | Sign posting
224825009 | Street name sign
224826005 | Building name sign
224827001 | Pedestrian direction sign
224828006 | Traffic sign
224898003 | Orthotic device
224899006 | Walking aid
224900001 | Communication aid
228167008 | Corset
228235002 | Slippers
228236001 | Mules
228237005 | Slippersox
228239008 | Trainers
228240005 | Plimsolls
228241009 | Sandals
228242002 | Gum boots
228243007 | Chappel
228259007 | Fastening
228260002 | Velcro
228261003 | Buckle
228262005 | Zipper
228264006 | Small button
228265007 | Medium button
228266008 | Large button
228267004 | Press stud
228268009 | Hook and eye
228270000 | Laces
228271001 | Shoe laces
228731007 | Radiotherapy equipment and appliances
228732000 | Beam direction shell
228733005 | Head and neck beam direction shell
228734004 | Body beam direction shell
228735003 | Beam modifier
228736002 | Surface bolus
228737006 | Surface compensator
228738001 | Cutout
228739009 | Shielding block
228740006 | Lung block
228741005 | Humerus block
228742003 | Scrotal block
228743008 | Kidney block
228744002 | Eye block
228745001 | Bite block
228746000 | Wedge filter
228747009 | Kilovoltage grid
228748004 | Brachytherapy implant
228749007 | Single plane implant
228750007 | Two plane implant
228751006 | Semicircular implant
228752004 | Regular volume implant
228753009 | Irregular volume implant
228754003 | Brachytherapy surface mold
228755002 | Two plane mold
228756001 | Single plane mold
228757005 | Cylinder mold
228759008 | Adhesive felt mold
228760003 | Elastoplast mold
228761004 | Collimator
228762006 | Multileaf collimator
228763001 | Asymmetric jaws collimator
228765008 | Standard collimator
228766009 | Form of brachytherapy source

228767000 | Wire source
228768005 | Seeds source
228770001 | Hairpins source
228771002 | Needles source
228772009 | Pellets source
228773004 | Capsules source
228774005 | Chains source
228775006 | Tubes source
228776007 | Rods source
228777003 | Grains source
228778008 | Plaque source
228869008 | Manual wheelchair
229772003 | Bed
229839006 | Functional foot orthosis
229840008 | Non-functional foot orthosis
229841007 | Detachable pad for the foot
229842000 | Detachable toe prop
229843005 | Detachable horseshoe pad
243135003 | Spacer
243719003 | Near low vision aid - integral eyeglass magnifier
243720009 | Near low vision aid - clip-on eyeglass magnifier
243722001 | Near low vision aid - integral eyeglass telescope
243723006 | Near low vision aid - clip-on eyeglass telescope
255296002 | Wedge
255712000 | Television
255716002 | Latex rubber gloves
256245006 | Textiles
256246007 | Cotton - textile
256247003 | Flax
256562002 | Cotton wool
256563007 | Cotton wool roll
256564001 | Cotton wool pledget
256589007 | Dental rubber dam
256590003 | Endodontic sponge
256593001 | Orthodontic elastic
256641009 | Ribbon gauze
256642002 | Wet ribbon gauze
256643007 | Dry ribbon gauze
257192006 | Aid to vision
257193001 | Telescopic eyeglasses
257194007 | Video
257211007 | Cylinder cutter
257212000 | Rotary cutter
257213005 | Rotary cutter with steel blades
257214004 | Rotary cutter with plastic blades
257215003 | Nail instrument
257216002 | Flexible endoscope
257217006 | Rigid endoscope
257218001 | Flexible cystoscope
257219009 | Rigid cystoscope
257220003 | Hysteroscope
257221004 | Flexible hysteroscope
257222006 | Rigid hysteroscope
257223001 | Contact hysteroscope
257224007 | Panoramic hysteroscope
257225008 | Flexible bronchoscope
257226009 | Rigid bronchoscope
257227000 | Standard laryngoscope
257228005 | Fiberlight anesthetic laryngoscope
257229002 | Pharyngeal mirror
257230007 | Obstetric forceps
257231006 | Barnes forceps

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information

Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)

Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')

Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application

Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.620 Value Set http://hl7.org/fhir/ValueSet/reason-medication-status-codes

URL: https://hl7.org/fhir/R4/valueset-reason-medication-status-codes.html
Pharmacy Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/reason-medication-status-codes
Version: | 4.0.1
Name: | SNOMEDCTDrugTherapyStatusCodes
Title: | SNOMED CT Drug Therapy Status codes
Definition: | This value set includes some taken and not taken reason codes from SNOMED CT - provided as an exemplar
Committee: | Pharmacy Work Group
OID: | 2.16.840.1.113883.4.642.3.369 (for OID based terminology systems)
Copyright: | This value set includes content from SNOMED CT, which is copyright © 2002+ International Health Terminology Standards Development Organisation (IHTSDO), and distributed by agreement between IHTSDO and HL7. Implementer use of SNOMED CT is not covered by this agreement
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: MedicationStatement.statusReason (CodeableConcept / Example)

4.4.1.620.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include codes from http://snomed.info/sct where concept is-a 266710000 (Drugs not taken/completed)

• Include codes from http://snomed.info/sct where concept is-a 410684002 (Drug therapy status)

4.4.1.620.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 120 concepts

All codes from system http://snomed.info/sct

Code | Display | Definition
182862001 | Drug not taken - dislike taste
182863006 | Drug not taken - dislike form
182864000 | Drug not taken - side-effects
182865004 | Drug not taken - inconvenient
182867007 | Drug not taken - pat disbelief
182868002 | Treatment stopped - alternative therapy undertaken
182869005 | Drug not taken - patient lost tablets
182870006 | Drug discontinued - reason unknown
182871005 | Drug discontinued - patient fear/risk
182872003 | Drug discontinued - too expensive
182873008 | Drug treatment stopped - patient ran out of tablets
182874002 | Treatment stopped - patient unable to concentrate
266711001 | Drug not taken - problem swallowing
275929009 | Tablets too large to swallow
134390006 | Angiotensin converting enzyme inhibitor not indicated
134397009 | Angiotensin converting enzyme inhibitor declined
135799002 | Angiotensin converting enzyme inhibitor indicated
161646004 | H/O: high risk medication
161647008 | H/O: anticoagulant therapy
161648003 | H/O: steroid therapy
161649006 | H/O: insulin therapy
161650006 | H/O: anticonvulsant therapy
161652003 | H/O: hormone replacement (HRT)
161653008 | H/O: chemotherapy
170622002 | Oral steroids started
170624001 | Oral steroids stopped
170688000 | Lithium stopped
170689008 | Injectable phenothiazine stopped
170821007 | B12 injections started
170826002 | B12 injections stopped
170919001 | Warfarin therapy stopped
170954008 | Hormone replacement therapy (HRT) contraindicated
170955009 | Hormone replacement therapy (HRT) started

170958006 | Hormonal replacement therapy (HRT) stopped
182834008 | Drug course completed
182840001 | Drug treatment stopped - medical advice
182841002 | Dr stopped drugs - ineffective
182842009 | Doctor stopped drugs - side effect
182843004 | Dr stopped drugs -inconvenient
182844005 | Doctor stopped drugs - patient dislikes
182845006 | Doctor stopped drugs - avoid interaction
182846007 | Doctor stopped drugs - medical aim achieved
274512008 | Drug therapy discontinued
309841001 | Thyroxine treatment stopped
312451002 | Aspirin prophylaxis contra-indicated
315023008 | Aspirin prophylaxis refused
315062004 | Beta blocker contraindicated
315364008 | Angiotensin converting enzyme inhibitors contraindicated
390914001 | Thrombolytic therapy refused
391158008 | Antibiotic prophylaxis recommended
391159000 | Antibiotic prophylaxis not recommended
394888000 | Hormone replacement therapy requested
394987009 | Angiotensin II receptor antagonists contraindicated
395006008 | Medication stopped - interaction
395007004 | Medication stopped - ineffective
395008009 | Medication stopped - contra-indication
395009001 | Medication stopped - side effect
401084003 | Angiotensin II receptor antagonist declined
401244004 | H/O: prolonged corticosteroid therapy
407572003 | Angiotensin II receptor antagonist not indicated
408339001 | Thrombolysis therapy not indicated
408354007 | Drug treatment still needed
408355008 | Drug treatment no longer needed
412782003 | Inappropriate medication stopped
413555000 | Antibiotic prophylaxis indicated
413558003 | Anticoagulation contraindicated
413559006 | Anticoagulation declined
413560001 | Anticoagulation not indicated
416404000 | Angiotensin converting enzyme inhibitor target dose not achieved
417576009 | Deferred antibiotic therapy
430279008 | Drug treatment stopped at end of clinical trial
438553004 | H/O: drug therapy
439631004 | H/O: statin therapy
440935004 | H/O: beta blocker therapy
440991005 | H/O: nitrate therapy
441186000 | H/O: antiplatelet agent therapy
441195008 | H/O: angiotensin converting enzyme (ACE) inhibitor therapy
441196009 | H/O: angiotensin II receptor antagonist therapy
441899004 | History of therapy with ototoxic medication
473151000 | History of tuberculosis chemotherapy
698428006 | Angiotensin II receptor antagonist indicated
698449007 | Antibiotic therapy indicated
698459008 | Antidepressant drug treatment stopped
698460003 | Antiviral therapy declined
698468005 | Bisoprolol contraindicated
698758004 | Nebivolol therapy refused
698985001 | Long term dual antiplatelet drug therapy indicated
699008007 | History of fenfluramine therapy
699137009 | Carvedilol therapy declined
700110004 | Bisphosphonate prophylaxis suspended
700384008 | Antibiotic prophylaxis not indicated
702542006 | Incretin mimetic therapy started
703989007 | Teriparatide therapy declined
703991004 | Raloxifene therapy declined
703996009 | Teriparatide not indicated
703997000 | Teriparatide contraindicated
703999002 | Raloxifene contraindicated
704000000 | Raloxifene not indicated
704033005 | Phosphodiesterase 5 inhibitor not indicated
704047009 | Phosphodiesterase 5 inhibitor declined
707298000 | Phosphodiesterase-5 inhibitor contraindicated
708000007 | Insulin treatment stopped
710159007 | Long-term current use of immunosuppressive drug
710814002 | Long-term current use of drug therapy
710815001 | Long-term current use of insulin

710816000 | Long-term current use of steroid
710817009 | Long-term current use of inhaled steroid
711149003 | Long-term current use of antibiotic
711150003 | Long-term current use of anticoagulant
711370000 | Oral steroids used since last encounter
711436000 | Inhaled steroids used daily
712985002 | Chemotherapy not done
106511000119106 | Long-term current use of mycophenolate
106521000119104 | Long-term current use of cyclophosphamide
106631000119102 | Long-term current use of infliximab
106691000119103 | Long-term current use of etanercept
133621000119103 | Long term current use of non-steroidal anti-inflammatory drug
134651000119108 | History of estrogen therapy
134671000119104 | History of systemic steroid therapy
140561000119102 | Long-term current use of azathioprine

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.621 Value Set http://hl7.org/fhir/ValueSet/publication-status

URL: https://hl7.org/fhir/R4/valueset-publication-status.html
FHIR Infrastructure Work Group | Maturity Level: N | Normative (from v4.0.0) | Use Context: Any

| This page has been approved as part of an ANSI standard. See the Conformance Package for further details.

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/publication-status
Version: | 4.0.1
Name: | PublicationStatus
Title: | PublicationStatus
Definition: | The lifecycle status of an artifact.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.3 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

- CodeSystem: This value set is the designated 'entire code system' value set for PublicationStatus
- ConceptMap: Translation source in Canonical Mapping for "PublicationStatus"
- Resource: CatalogEntry.status (code / Required)
- Resource: EventDefinition.status (code / Required)
- Resource: MessageDefinition.status (code / Required)
- Resource: CapabilityStatement.status (code / Required)
- Resource: Measure.status (code / Required)
- Resource: GraphDefinition.status (code / Required)
- Resource: ResearchDefinition.status (code / Required)
- Resource: ImplementationGuide.status (code / Required)
- Resource: ResearchElementDefinition.status (code / Required)
- Resource: SearchParameter.status (code / Required)
- Resource: ActivityDefinition.status (code / Required)
- Resource: InsurancePlan.status (code / Required)
- Resource: Questionnaire.status (code / Required)
- Resource: StructureDefinition.status (code / Required)
- Resource: ChargeItemDefinition.status (code / Required)
- Resource: ConceptMap.status (code / Required)

- Resource: ValueSet.status (code / Required)
- Resource: OperationDefinition.status (code / Required)
- Resource: EffectEvidenceSynthesis.status (code / Required)
- Resource: EvidenceVariable.status (code / Required)
- Resource: CodeSystem.status (code / Required)
- Resource: RiskEvidenceSynthesis.status (code / Required)
- Resource: StructureMap.status (code / Required)
- Resource: Library.status (code / Required)
- Resource: TestScript.status (code / Required)
- Resource: TerminologyCapabilities.status (code / Required)
- Resource: Evidence.status (code / Required)
- Resource: CompartmentDefinition.status (code / Required)
- Resource: NamingSystem.status (code / Required)
- Resource: ExampleScenario.status (code / Required)
- Resource: PlanDefinition.status (code / Required)

4.4.1.621.1 Content Logical Definition

This value set includes codes from the following code systems:

- Include all codes defined in http://hl7.org/fhir/publication-status

4.4.1.621.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 4 concepts

Expansion based on http://hl7.org/fhir/publication-status version 4.0.1

All codes from system http://hl7.org/fhir/publication-status

Code | Display | Definition | Canonical Status
draft | Draft | This resource is still under development and is not yet considered to be ready for normal use. | ~draft
active | Active | This resource is ready for normal use. | ~active
retired | Retired | This resource has been withdrawn or superseded and should no longer be used. | ~inactive
unknown | Unknown | The authoring system does not know which of the status values currently applies for this resource. Note: This concept is not to be used for "other" - one of the listed statuses is presumed to apply, it's just not known which one. | ~unknown

Additional Language Displays

Code | Nederlands (Dutch, nl) | Русский (Russian, ru)
draft | ontwerp | черновик
active | actief | активный
retired | verouderd | удалён
unknown

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.622 Value Set http://hl7.org/fhir/ValueSet/food-type

URL: https://hl7.org/fhir/R4/valueset-food-type.html
Orders and Observations Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/food-type
Version: | 4.0.1
Name: | FoodTypeCodes
Title: | Food Type Codes

Definition: | This value set represents codes for types of foods and is provided as a suggestive example. It include codes from SNOMED CT where concept is-a 255620007 (Foods (substance)).

Committee: | Orders and Observations Work Group

OID: | 2.16.840.1.113883.4.642.3.384 (for OID based terminology systems)

Copyright: | This resource includes content from SNOMED Clinical Terms® (SNOMED CT®) which is copyright of the International Health Terminology Standards Development Organisation (IHTSDO). Implementers of these specifications must have the appropriate SNOMED CT Affiliate license - for more information contact http://www.snomed.org/snomed-ct/get-snomed-ct or info@snomed.org

Source Resource | XML / JSON

This value set is used in the following places:

• Resource: NutritionOrder.excludeFoodModifier (CodeableConcept / Example)

4.4.1.622.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include codes from http://snomed.info/sct where concept is-a 255620007 (Foods)

4.4.1.622.2 Expansion

This expansion generated 01 Nov 2019

This value set has >1000 codes in it. In order to keep the publication size manageable, only a selection (1000 codes) of the whole set of codes is shown

All codes from system http://snomed.info/sct

Code	Display	Definition
3718001	Cow's milk	
4700006	Bovine insulin	
7661006	Fish bone	
7791007	Soy protein	
10249006	Agar	
10827009	Milk protein	
13577000	Nut	
14263006	Prepared fish	
22836000	Vegetable	
23182003	Cereal	
24515005	Spice	
25743006	Skimmed milk	
28230009	White meat	
28647000	Meat	
28942008	Coconut oil	
39102003	Food particle	
41834005	Olive oil	
43708003	Animal feed additive	
44027008	Seafood	
44632005	4,7,10,13,16,19-Docosahexaenoic acid	
46329000	Chocolate milk	
50479000	Soya protein-iron complex	
50593009	Casein	
51905005	Mustard	
53875002	First milk	
54041009	Safflower oil	
63045006	Berry	
63766005	Flour	
67324005	Rice	
70813002	Milk	
72511004	Fruit	
74242007	Food starch	
75476008	Bovine growth hormone	
77722008	Poultry bone	
80743002	Mustard black	
82450006	Cottonseed oil	
82566005	Animal feed	
83235009	Bovine growth hormone recombinant	
83595008	Goat's milk	
85668006	Starch powder	
89707004	Sesame oil	
89811004	Gluten	
90677004	Mustard white	
91606004	Cochineal	
96364008	Sometribove	
102259006	Citrus fruit	
102260001	Peanut butter	

102261002 | Strawberries
102262009 | Chocolate
102263004 | Eggs (edible)
102264005 | Cheese
102270004 | Growth stimulant
102271000 | Efrotomycin
102697003 | Lactalbumin
102698008 | alpha-Lactalbumin
102699000 | beta-Lactalbumin
108743002 | Pegademase bovine
109010000 | Microfibrillar collagen hemostat
116957009 | Bovine shock protein antibody
117633004 | Bovine osteocalcin
126076005 | Fish liver oil
126077001 | Burbot liver oil
126078006 | Percomorph liver oil
126079003 | Percoid liver oil
126080000 | Shark liver oil
126081001 | Halibut liver oil
126082008 | Cod liver oil
127384006 | Salcatonin
226017009 | Bombay mix
226021002 | Cauliflower cheese
226022009 | Cheese pastry
226023004 | Cheesecake
226024005 | Chocolate digestive biscuit
226037008 | Lasagne
226041007 | Mint
226046002 | Plain digestive biscuit
226061001 | Tamarind
226062008 | Trail mix
226358006 | Fat - dietary
226359003 | Fish oil
226367006 | Eicosapentaenoic acid
226368001 | DHA - Docosahexaenoic acid
226491003 | Apple juice
226492005 | Grape juice
226493000 | Grapefruit juice
226495007 | Lemon juice
226496008 | Orange juice
226497004 | Pineapple juice
226498009 | Tomato juice
226529007 | Starchy food
226530002 | Baby cereal
226531003 | Rusk
226532005 | Flavored rusk
226533000 | Low sugar rusk
226534006 | Plain rusk
226535007 | Wholemeal rusk
226536008 | Rice-based baby cereal
226537004 | Mixed cereal-based baby cereal
226538009 | Wheat-based baby cereal
226539001 | Cookie and/or cracker
226540004 | Chocolate biscuit
226541000 | Full coated chocolate biscuit
226542007 | Half-coated chocolate biscuit
226543002 | Cracker
226544008 | Cream cracker
226546005 | Wholemeal cracker
226547001 | Crispbread
226548006 | Digestive biscuit
226549003 | Gingernut biscuit
226550003 | Home-made biscuit
226551004 | Matzo
226552006 | Oatcake
226553001 | Sandwich biscuit
226554007 | Semi-sweet biscuit
226555008 | Short-sweet biscuit
226556009 | Shortbread biscuit
226557000 | Wafer biscuit
226558005 | Water biscuit
226559002 | Bread

226560007 | Brown bread
226561006 | Brown bread roll
226562004 | Breadcrumbs
226563009 | Chapati
226564003 | White chapati
226565002 | White chapati made with fat
226566001 | White chapati made without fat
226567005 | Wholemeal chapati
226568000 | Wholemeal chapati made with fat
226569008 | Wholemeal chapati made without fat
226570009 | Currant bread
226571008 | Granary bread
226572001 | Granary bread roll
226574000 | Malt bread
226575004 | Naan bread
226576003 | Papadum
226577007 | Fried papadum
226578002 | Grilled papadum
226579005 | Paratha
226580008 | Pitta bread
226581007 | White pitta bread
226582000 | Wholemeal pitta bread
226583005 | Milk bread
226584004 | Rye bread
226585003 | Soda bread
226587006 | Tortilla
226589009 | Wheatgerm bread
226590000 | White bread
226592008 | Fried white bread
226593003 | West Indian white bread
226595005 | White bread roll
226596006 | Wholemeal bread
226597002 | Wholemeal bread roll
226598007 | Breakfast cereal
226601002 | Bran flakes
226604005 | Cornflakes
226605006 | Crunchy nut cornflakes
226611009 | Muesli
226612002 | Muesli Swiss style
226613007 | Muesli with extra fruit
226614001 | Muesli with no added sugar
226616004 | Oat and wheat bran
226617008 | Porridge
226618003 | Porridge made with milk
226619006 | Porridge made with semi-skimmed milk
226620000 | Porridge made with skimmed milk
226621001 | Porridge made with whole milk
226622008 | Porridge made with water
226639005 | Cake
226640007 | All bran loaf
226641006 | Battenburg cake
226642004 | Welsh cheesecake
226643009 | Cherry cake
226644003 | Chinese cake
226645002 | Chinese glutinous rice flour cake
226647005 | Chocolate cake
226648000 | Chocolate cake with butter icing
226649008 | Coconut cake
226650008 | Crispie cake
226651007 | Fancy iced cake
226652000 | Fruit cake
226653005 | Plain fruit cake
226654004 | Rich fruit cake
226655003 | Iced fruit cake
226656002 | Wholemeal fruit cake
226657006 | Gateau
226658001 | Gingerbread
226659009 | Lardy cake
226660004 | Madeira cake
226661000 | Rock cake
226662007 | Sevyiann
226663002 | Sponge cake

226664008 | Fatless sponge cake
226665009 | Frozen sponge cake
226666005 | Jam-filled sponge cake
226667001 | Fresh cream-filled sponge cake
226668006 | Sponge cake with butter icing
226669003 | Sponge cake made with cake mix
226670002 | Swiss roll
226671003 | Chocolate filled swiss roll
226672005 | Jam filled swiss roll
226673000 | Individual swiss roll
226674006 | Welsh cake
226675007 | Individual cakes and pastries
226676008 | Asian pastry
226677004 | Brandy snap
226678009 | Burfi
226679001 | Chelsea bun
226680003 | Choux bun
226681004 | Cream horn
226682006 | Crumpet
226683001 | Currant bun
226684007 | Custard tart
226685008 | Danish pastry
226686009 | Doughnut
226687000 | Jam doughnut
226688005 | Ring doughnut
226689002 | Fresh cream doughnut
226690006 | Eccles cake
226691005 | Eclair
226692003 | Flapjack
226693008 | Greek pastry
226694002 | Gulab jamen
226695001 | Halwa
226697009 | Hot cross bun
226698004 | Jaffa cake
226699007 | Jam tart
226700008 | Jellabi
226701007 | Melting moment
226702000 | Mince pie
226703005 | Mincemeat tart
226704004 | Muffin
226705003 | Pinni
226706002 | Rum baba
226707006 | Scone
226708001 | Cheese scone
226709009 | Fruit scone
226710004 | Plain scone
226711000 | Potato scone
226712007 | Wholemeal scone
226713002 | Wholemeal fruit scone
226714008 | Scotch pancake
226715009 | Shortbread
226716005 | Strawberry tartlet
226717001 | Teacake
226718006 | Vanilla slice
226719003 | Waffle
226720009 | Arrowroot
226721008 | Pearl barley
226722001 | Whole grain barley
226723006 | Buckwheat - cereal
226724000 | Bulgur wheat
226725004 | Oatmeal
226726003 | Popcorn
226727007 | Plain popcorn
226728002 | Sweet popcorn
226729005 | Salted popcorn
226730000 | Buttered popcorn
226731001 | Sago
226732008 | Semolina
226733003 | Tapioca
226734009 | Wheatgerm
226735005 | Noodles
226736006 | Egg noodles

226737002 | Fried noodles
226738007 | Plain noodles
226739004 | Pot noodles
226740002 | Pasta
226741003 | Cannelloni
226742005 | Macaroni
226743000 | Ravioli
226744006 | Spaghetti
226745007 | White spaghetti
226746008 | Wholemeal spaghetti
226747004 | Tinned spaghetti
226748009 | Vermicelli
226749001 | Pastry
226750001 | Choux pastry
226751002 | Flaky pastry
226752009 | Flan case pastry
226753004 | Puff pastry
226754005 | Shortcrust pastry
226755006 | Wholemeal pastry
226756007 | Brown rice
226757003 | Red rice
226758008 | Savory rice
226759000 | White rice
226760005 | Dairy products
226761009 | Gold top milk
226762002 | Flavored cow's milk
226763007 | Semi-skimmed milk
226764001 | Calcium fortified semi-skimmed milk
226765000 | Calcium fortified skimmed milk
226766004 | Sterilized milk
226767008 | Whole milk
226768003 | Calcium fortified whole milk
226769006 | Buttermilk
226770007 | Whey
226771006 | Dried milk
226774003 | Buttermilk powder
226775002 | Dried goat's milk
226776001 | Dried cow's milk
226777005 | Dried semi-skimmed cow's milk
226778000 | Skimmed milk powder
226779008 | Dried skimmed cow's milk with vegetable fat
226780006 | Dried whole milk
226781005 | Dried soy milk
226782003 | Dried whey
226783008 | Tinned milk
226784002 | Condensed milk
226785001 | Skimmed sweetened condensed milk
226786000 | Whole sweetened condensed milk
226787009 | Condensed milk
226788004 | Low fat evaporated milk
226789007 | Mother's milk
226790003 | EBM - Expressed breast milk
226791004 | Sheep milk
226793001 | Soy milk
226795008 | Junior milk
226796009 | Fresh cream
226797000 | Half-cream
226798005 | Double cream
226799002 | Clotted cream
226800003 | Single cream
226801004 | Soured cream
226802006 | Whipping cream
226803001 | Tinned cream
226804007 | Tinned tip-top cream
226805008 | Imitation cream
226806009 | Dessert top cream
226807000 | Elmlea cream
226808005 | Smatana
226809002 | Cheese spread
226810007 | Plain cheese spread
226811006 | Flavored cheese spread
226812004 | Lactic cheese spread

226813009 | Low fat cheese spread
226814003 | Cottage cheese
226815002 | Plain cottage cheese
226816001 | Very low fat cottage cheese
226817005 | Cottage cheese with additions
226818000 | Hard cheese
226819008 | Caerphilly cheese
226820002 | Cheddar cheese
226821003 | Reduced fat cheddar cheese
226822005 | Vegetarian cheddar cheese
226823000 | Cheshire cheese
226824006 | Reduced fat Cheshire cheese
226825007 | Derby cheese
226826008 | Danish blue cheese
226827004 | Double Gloucester cheese
226828009 | Edam cheese
226829001 | Edam-type cheese reduced fat
226830006 | Emmental cheese
226831005 | Feta cheese
226832003 | Gouda cheese
226834002 | Gruyere cheese
226835001 | Lancashire cheese
226836000 | Leicester cheese
226837009 | Mozzarella cheese
226838004 | Parmesan cheese
226839007 | Red Windsor cheese
226840009 | Roquefort cheese
226841008 | Sage Derby cheese
226842001 | Soy cheese
226843006 | Stilton - white cheese
226844000 | Stilton - blue cheese
226845004 | Wensleydale cheese
226846003 | Processed cheese
226847007 | Smoked processed cheese
226848002 | Soft cheese
226849005 | Cream cheese
226850005 | Medium fat cream cheese
226851009 | Low fat cream cheese
226852002 | Goats milk soft cheese
226853007 | Ricotta cheese
226854001 | Soft mold ripened cheese
226855000 | Brie cheese
226856004 | Camembert cheese
226857008 | Fromage frais
226858003 | Plain fromage frais
226859006 | Flavored fromage frais
226860001 | Fruit fromage frais
226861002 | Very low fat fromage frais
226862009 | Quark
226863004 | Yoghurt
226864005 | Whole milk yogurt
226865006 | Whole milk plain yogurt
226866007 | Whole milk fruit yogurt
226867003 | Whole milk flavored yoghurt
226868008 | Live yogurt
226869000 | Organic yogurt
226870004 | Goat's milk yogurt
226871000 | Low fat yogurt
226872007 | Plain low fat yogurt
226873002 | Flavored low fat yoghurt
226874008 | Fruit low fat yogurt
226875009 | Muesli-nut low fat yogurt
226876005 | Low calorie yogurt
226877001 | Drinking yogurt
226878006 | Greek yogurt
226879003 | Sheep's milk yogurt
226880000 | Soy milk yoghurt
226881001 | Dried egg
226882008 | BE - Boiled egg
226883003 | Fried egg
226884009 | PE - Poached egg
226885005 | Raw egg

226886006 | Scrambled egg
226887002 | Dietary fats and oils
226888007 | Butter
226889004 | Blended spread
226890008 | Margarine
226891007 | Hard vegetable fat margarine
226892000 | Hard animal and vegetable fat margarine
226894004 | Soft polyunsaturated fat margarine
226895003 | Olive oil margarine
226896002 | Low fat spread
226897006 | VLF - Very low fat spread
226898001 | Dripping
226899009 | Lard
226900004 | Suet
226901000 | Ghee
226902007 | Butter ghee
226903002 | Palm oil ghee
226904008 | Vegetable oil ghee
226905009 | Oil - dietary
226906005 | Maize oil
226907001 | Palm oil
226908006 | Rapeseed oil
226909003 | Rapeseed oil high erucic acid
226910008 | Rapeseed oil low erucic acid
226911007 | Soybean oil
226912000 | Sunflowerseed oil
226913005 | Blended vegetable oil
226914004 | Polyunsaturated vegetable oil
226915003 | Red meat
226916002 | Beef
226917006 | Beef steak
226918001 | Grilled beef steak
226919009 | Fried beef steak
226920003 | Boiled beef
226921004 | Boiled salt beef
226922006 | Minced beef
226923001 | Stewed minced beef
226924007 | Stewed extra-lean minced beef
226925008 | Roast beef
226926009 | Lean roast beef
226927000 | Stewed beef
226928005 | Bacon
226929002 | Boiled bacon
226930007 | Fried bacon
226931006 | Grilled bacon
226932004 | Gammon
226933009 | Roast ham
226934003 | Pork
226935002 | Fried pork chop
226936001 | Grilled pork chop
226937005 | Minced pork
226938000 | Roast pork
226939008 | Lean roast pork
226940005 | Roast belly pork
226941009 | Boiled ham
226942002 | Lamb - meat
226943007 | Fried lamb chop
226944001 | Grilled lamb chop
226945000 | Minced lamb
226946004 | Roast lamb
226947008 | Lean roast lamb
226948003 | Roast leg of lamb
226949006 | Roast shoulder of lamb
226950006 | Roast breast of lamb
226951005 | Stewed lamb
226952003 | Veal
226953008 | Fried veal
226954002 | Grilled veal
226955001 | Chicken
226956000 | Boiled chicken
226957009 | Fried chicken
226958004 | Fried chicken in batter

226959007 | Fried chicken in breadcrumb
226960002 | Fried chicken nuggets
226961003 | Roast chicken
226962005 | Minced chicken
226963000 | Duck - meat
226964006 | Roast duck
226965007 | Goose - meat
226966008 | Roast goose
226967004 | Turkey - meat
226968009 | Roast turkey
226969001 | Stewed turkey
226970000 | Minced turkey
226971001 | Game - meat
226972008 | Grouse - meat
226973003 | Roast grouse
226974009 | Partridge - meat
226975005 | Roast partridge
226976006 | Pheasant
226977002 | Roast pheasant
226978007 | Roast pigeon
226979004 | Roast venison
226980001 | Stewed hare
226981002 | Stewed rabbit
226982009 | Offal
226983004 | Heart - meat
226984005 | Stewed heart
226985006 | Roast heart
226987003 | Kidney - meat
226988008 | Fried kidney
226989000 | Grilled kidney
226990009 | Stewed kidney
226991008 | Liver - meat
226992001 | Fried liver
226993006 | Grilled liver
226994000 | Stewed liver
226995004 | Oxtail - food
226996003 | Stewed oxtail
226997007 | Sweetbread
226998002 | Fried sweetbread
226999005 | Tongue - meat
227000008 | Tripe
227001007 | Dressed tripe
227002000 | Dressed stewed tripe
227003005 | Meat agents
227004004 | Meat burger
227005003 | Beef burger
227006002 | Pork burger
227007006 | Lamb burger
227008001 | Chicken burger
227009009 | Turkey burger
227010004 | Black pudding
227011000 | Brawn - meat
227012007 | Faggot - meat
227013002 | Haggis
227014008 | Liver pate
227015009 | LF - Low fat liver pate
227016005 | Meat paste
227017001 | Polony - meat agent
227018006 | Salami
227019003 | Saveloy
227020009 | Sausage
227021008 | Frankfurter
227022001 | LF - Low fat sausage
227023006 | Beef sausage
227024000 | LF - Low fat beef sausage
227025004 | Pork sausage
227026003 | LF - Low fat pork sausage
227027007 | Turkey sausage
227028002 | Low fat turkey sausage
227029005 | Liver sausage
227030000 | White pudding
227031001 | Processed meat

227032008 | Chicken roll
227033003 | Chopped ham and pork
227034009 | Corned beef
227036006 | Luncheon meat
227037002 | Fish
227038007 | White fish
227040002 | Bele
227041003 | Boal
227042005 | Bream
227043000 | Catfish - dietary
227044006 | Chinese salted fish
227045007 | Chital
227046008 | Baked cod
227047004 | Poached cod
227048009 | Grilled cod
227049001 | Fried cod
227050001 | Fried cod in batter
227051002 | Fried cod in breadcrumbs
227052009 | Fried cod goujons in breadcrumbs
227053004 | Cod in batter baked in the oven
227054005 | Dried salted cod
227055006 | Smoked poached cod
227056007 | Coley
227057003 | Conger eel
227058008 | Dab
227059000 | Haddock - dietary
227060005 | Poached haddock
227061009 | Fried haddock
227062002 | Fried haddock in batter
227063007 | Fried haddock in breadcrumbs
227064001 | Fried haddock goujons in breadcrumbs
227065000 | Smoked poached haddock
227066004 | Hake
227067008 | Halibut
227068003 | Hoki
227069006 | Fried hoki in batter
227070007 | John dory
227071006 | Kalabasu
227072004 | Lemon sole
227073009 | Poached lemon sole
227074003 | Fried lemon sole
227075002 | Fried lemon sole in breadcrumbs
227076001 | Ling
227078000 | Monkfish
227079008 | Mullet
227080006 | Gray mullet
227081005 | Red mullet
227082003 | Parrot fish - dietary
227083008 | Plaice
227084002 | Poached plaice
227085001 | Fried plaice
227086000 | Fried plaice in batter
227087009 | Fried plaice in breadcrumbs
227088004 | Fried plaice goujons in breadcrumbs
227089007 | Pollack
227091004 | Pomfret - dietary
227092006 | Black pomfret
227093001 | White pomfret
227094007 | Red snapper - dietary
227095008 | Redfish
227096009 | Rohu
227097000 | Dogfish
227098005 | Fried rock salmon in batter
227099002 | Saithe
227100005 | Sea bass - dietary
227101009 | Shark - food
227102002 | Skate
227103007 | Fried skate in batter
227104001 | Tiliapa
227105000 | Turbot
227106004 | Whiting
227107008 | Poached whiting

227108003 | Fried whiting
227109006 | Fried whiting in breadcrumbs
227110001 | Oily fish
227111002 | Bacha
227112009 | Bloater - dietary
227113004 | Carp
227114005 | Jellied eel
227115006 | Dried herring
227116007 | Fried herring in oatmeal
227117003 | Grilled herring
227118008 | Pickled herring
227119000 | Hilsa
227120006 | Jackfish
227121005 | Katla
227122003 | Kipper
227123008 | Poached kipper
227124002 | Grilled kipper
227125001 | Mackerel
227126000 | Fried mackerel
227127009 | Grilled mackerel
227128004 | Smoked mackerel
227129007 | Grilled smoked mackerel
227130002 | Orange roughy
227131003 | Pangas
227132005 | Grilled salmon
227133000 | Poached salmon
227134006 | Smoked salmon
227135007 | Salmon en croute
227136008 | Sardine
227137004 | Grilled sardine
227138009 | Fried sardine
227139001 | Sprats
227140004 | Swordfish - dietary
227141000 | Fried trout
227142007 | Baked trout
227143002 | Smoked trout
227144008 | Tuna
227145009 | Whitebait fried
227146005 | Shellfish
227147001 | Cockles
227148006 | Cockles bottled in vinegar
227149003 | Cuttlefish - dietary
227150003 | Mussels
227151004 | Prawns
227152006 | Boiled prawns
227153001 | Fried prawns
227154007 | Fried prawns in batter
227155008 | Dried prawns
227156009 | Scampi
227157000 | Fried scampi in batter
227158005 | Fried scampi in breadcrumbs
227159002 | Fried shrimp
227160007 | Dried shrimp
227161006 | Octopus - dietary
227162004 | Scallop
227163009 | Squid - dietary
227164003 | Dried squid
227165002 | Fried squid
227167005 | Fried squid in batter
227168000 | Whelks
227169008 | Winkles
227170009 | Fish products
227171008 | Crabstick
227172001 | Fish ball
227173006 | Fishcake
227174000 | Grilled fishcake
227175004 | Fried fishcake
227176003 | Fish fingers
227177007 | Fried fish fingers
227178002 | Grilled fish fingers
227179005 | Fish paste
227180008 | Fish pate

227181007 | Fish pie
227182000 | Kedgeree
227183005 | Fish roe
227184004 | Raw hard fish roe
227185003 | Fried hard fish roe
227186002 | Fried hard fish roe in batter
227187006 | Raw soft fish roe
227188001 | Fried soft fish roe
227189009 | Seafood cocktail
227190000 | Taramasalata
227191001 | Tinned fish
227193003 | Anchovies canned in oil
227194009 | Herring canned in tomato sauce
227195005 | Mackerel canned in tomato sauce
227196006 | Mackerel canned in brine
227197002 | Mussels canned in brine
227198007 | Pilchards canned in tomato sauce
227199004 | Pilchards canned in brine
227200001 | Salmon canned in brine
227201002 | Sardines canned in brine
227202009 | Sardines canned in oil
227203004 | Sardines canned in tomato sauce
227204005 | Tuna canned in brine
227205006 | Tuna canned in oil
227206007 | Crab canned in brine
227207003 | Shrimps canned in brine
227208008 | Abalone canned in brine
227209000 | Clams canned in brine
227210005 | Vegetables pulses herbs and spices
227212002 | Alfalfa sprout
227213007 | Amaranth leaf
227214001 | Arrowhead
227215000 | Artichoke
227216004 | Globe artichoke
227217008 | Jerusalem artichoke
227218003 | Asparagus
227219006 | Eggplant
227220000 | Beetroot
227221001 | Brassica vegetable
227222008 | Broccoli
227223003 | Brussels sprouts
227224009 | Cabbage - dietary
227225005 | Chinese cabbage
227226006 | Red cabbage
227227002 | Savoy cabbage
227228007 | Spring cabbage
227229004 | White cabbage
227230009 | Curly kale
227231008 | Kohl rabi
227232001 | Seakale
227233006 | Breadfruit
227234000 | Cassava - dietary
227235004 | Celeriac
227236003 | Chard
227237007 | Chicory - dietary
227238002 | Cho cho
227239005 | Zucchini
227240007 | Cole leaf
227241006 | Cucumber - dietary
227242004 | Drumstick leaf
227243009 | Endive
227244003 | Fenugreek
227245002 | Gherkin
227246001 | Gourd - dietary
227247005 | Jackfruit
227248000 | Leek - dietary
227249008 | Lotus tuber
227250008 | Marrow
227251007 | Matoki
227252000 | Mushroom - dietary
227253005 | Mustard leaf
227254004 | Okra

227255003 | Spring onion
227256002 | Pawpaw
227257006 | Parsnip
227258001 | Patra leaf
227259009 | Plantain
227260004 | Pumpkin
227261000 | Raddiccio
227262007 | Radish white - mooli
227263002 | Rape leaf
227264008 | Salsify
227265009 | Seaweed - dietary
227266005 | Swede
227267001 | Sweet potato - dietary
227268006 | Sweetcorn
227269003 | Sweetcorn kernels
227270002 | Sweetcorn on-the-cob
227271003 | Tannia
227272005 | Taro
227273000 | Taro leaves
227274006 | Tomato puree
227275007 | Turnip
227276008 | Water chestnut
227277004 | Watercress
227278009 | Yam - dietary
227279001 | Boiled potato
227280003 | Potato boiled in salted water
227281004 | Potato boiled in unsalted water
227282006 | French fries
227283001 | Thick cut French fries
227284007 | Thin cut French fries
227285008 | Crinkle cut chips
227286009 | French fries cooked in corn oil
227287000 | French fries cooked in fat drippings
227288005 | French fries cooked in blended oil
227289002 | French fries cooked in polyunsaturated oil
227290006 | Microwave French fries
227291005 | Oven French fries
227292003 | Jacket potato
227294002 | Mashed potato
227295001 | Potato mashed with butter
227296000 | Potato mashed with margarine
227297009 | Potato mashed with milk
227298004 | Potato mashed with polyunsaturated margarine
227299007 | Roast potato
227300004 | Potato roasted in vegetable oil
227301000 | Potato roasted in polyunsaturated oil
227302007 | Potato roasted in dripping
227303002 | Tinned potato
227304008 | Potato tinned in salted water
227305009 | Potato tinned in unsalted water
227306005 | Instant potato
227307001 | Instant potato made up with water
227308006 | Instant potato made up with semi-skimmed milk
227309003 | Instant potato made up with skimmed milk
227310008 | Instant potato made up with whole milk
227311007 | Potato croquette fried in blended oil
227312000 | Potato waffle
227313005 | Legumes
227314004 | Canned beans
227315003 | Baked beans
227316002 | Baked beans canned in tomato sauce
227317006 | Baked beans canned in tomato sauce reduced sugar
227318001 | Baked beans canned in tomato sauce reduced salt
227319009 | Baked beans canned in tomato sauce with burgers
227320003 | Baked beans canned in tomato sauce with pork sausages
227321004 | Balor beans canned
227322006 | Barbecue beans canned in sauce
227323001 | Canned mung beansprouts
227324007 | Canned broad beans
227325008 | Canned butter beans
227326009 | Canned chick peas
227327000 | Canned chili beans

227328005 | Canned haricot beans
227329002 | Canned lilva beans
227330007 | Canned lentils in tomato sauce
227331006 | Canned papri beans
227332004 | Canned peas
227333009 | Canned garden peas
227334003 | Canned marrowfat peas
227335002 | Canned mushy peas
227336001 | Canned pease pudding
227337005 | Canned red kidney beans
227338000 | Aduki beans
227339008 | Beansprouts
227340005 | Raw mung beansprouts
227341009 | Stir-fried mung beansprouts
227342002 | Black gram beans
227343007 | Blackeye beans
227344001 | Broad bean - dietary
227345000 | Butter beans
227346004 | Chick peas
227347008 | Cluster beans
227348003 | Haricot beans
227349006 | Hummus
227350006 | Lentils
227351005 | Green lentils
227352003 | Brown lentils
227353008 | Red split lentils
227354002 | Miso
227355001 | Mung beans
227356000 | Mung bean dahl
227357009 | Papri beans
227358004 | Pigeon peas
227359007 | Pigeon pea dahl
227360002 | Pinto beans
227361003 | Refried pinto beans
227362005 | Kidney beans
227363000 | Sword beans
227364006 | Tempeh
227365007 | Tofu
227366008 | Steamed tofu
227367004 | Steamed fried tofu
227368009 | Fu-juk tofu
227369001 | Fresh peas
227370000 | Fresh raw peas
227371001 | Fresh boiled peas
227372008 | Dried boiled peas
227373003 | Dried boiled split peas
227374009 | Herbs and spices
227375005 | Allspice
227376006 | Anise seed
227377002 | Asafoetida
227378007 | Basil - dietary
227379004 | Bay leaf
227380001 | Betel leaf
227381002 | Caraway seed
227382009 | Cardemon
227383004 | Celery seed
227384005 | Chervil, dried
227385006 | Chili powder
227386007 | Chinese five spice
227387003 | Chive - herb
227388008 | Cinnamon
227389000 | Clove
227390009 | Cilantro
227391008 | Coriander seed
227392001 | Cumin seed
227393006 | Curry leaf
227395004 | Curry powder
227396003 | Fennel
227397007 | Fennel seed
227398002 | Fenugreek seed
227399005 | Garam masala
227400003 | Ginger

227401004 | Ground ginger
227402006 | Root ginger
227403001 | Liquorice powder
227404007 | Mace
227405008 | Marjoram
227406009 | Mustard seed
227407000 | Oregano
227408005 | Paprika
227409002 | Poppy seed
227410007 | Rosemary
227411006 | Sage
227412004 | Tarragon
227413009 | Thyme
227414003 | Turmeric
227415002 | Fruit nuts and seeds
227416001 | Fresh fruit
227417005 | Bilberry
227418000 | Blackberries
227419008 | Blackcurrant
227420002 | Clementine
227421003 | Cranberries
227422005 | Damson
227423000 | Dates
227424006 | Elderberries
227425007 | Figs
227426008 | Fresh fruit salad
227427004 | Gooseberries
227428009 | Greengage
227429001 | Guava
227430006 | Lime
227431005 | Kumquat
227432003 | Loganberries
227433008 | Lychee
227434002 | Mulberries
227435001 | Nectarine
227436000 | Olives
227437009 | Ortanique
227438004 | Passion fruit
227439007 | Pawpaw
227440009 | Plum - dietary
227441008 | Pomegranate
227442001 | Pomelo
227443006 | Quince
227444000 | Raspberries
227445004 | Redcurrant
227446003 | Satsuma
227447007 | Sharon fruit
227448002 | Whitecurrant
227449005 | Dried fruit
227450005 | Dried banana
227451009 | Dried apple
227452002 | Dried apricot
227453007 | Currants
227454001 | Dried date
227455000 | Dried fig
227456004 | Dried mixed fruit
227457008 | Dried mixed peel
227458003 | Dried peach
227459006 | Dried pear
227460001 | Dried pineapple
227461002 | Glace cherry
227462009 | Prunes
227463004 | Raisin
227464005 | Sultanas
227465006 | Canned fruit
227466007 | Tinned apple
227467003 | Tinned apricot
227468008 | Tinned blackcurrants
227469000 | Tinned cherries
227470004 | Tinned figs
227471000 | Tinned fruit cocktail
227472007 | Tinned gooseberries

227473002 | Tinned grapefruit
227474008 | Tinned guava
227475009 | Tinned kumquat
227476005 | Tinned loganberries
227477001 | Tinned lychees
227478006 | Tinned mandarin oranges
227479003 | Tinned mango
227480000 | Tinned paw-paw
227481001 | Tinned peaches
227482008 | Tinned pears
227483003 | Tinned pineapple
227484009 | Tinned plums
227485005 | Tinned prunes
227486006 | Tinned raspberries
227487002 | Tinned rhubarb
227488007 | Tinned strawberries
227489004 | Tinned fruit pie filling

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.623 Value Set http://hl7.org/fhir/ValueSet/care-plan-intent

URL: https://hl7.org/fhir/R4/valueset-care-plan-intent.html
Patient Care Work Group | Maturity Level: 2 | Trial Use | Use Context: Any

This is a value set defined at http://www.hl7.org/Special/committees/patientcare/ .

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/care-plan-intent
Version: | 4.0.1
Name: | CarePlanIntent
Title: | Care Plan Intent
Definition: | Codes indicating the degree of authority/intentionality associated with a care plan.
Committee: | Patient Care Work Group
OID: | 2.16.840.1.113883.4.642.3.150 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: CarePlan.intent (code / Required)

4.4.1.623.1 Content Logical Definition

This value set includes codes from the following code systems:

http://hl7.org/fhir/request-intent

Code | Display
proposal | Proposal | The request is a suggestion made by someone/something that does not have an intention to ensure it occurs and without providing an authorization to act.
plan | Plan | The request represents an intention to ensure something occurs without providing an authorization for others to act.
order | Order | The request represents a request/demand and authorization for action by a Practitioner.
option | Option | The request represents a component or option for a RequestGroup that establishes timing, conditionality and/or other constraints among a set of requests. Refer to [[[RequestGroup]]] for additional information on how this status is used.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept

Comments | Additional notes about how to use the code

4.4.1.624 Value Set http://hl7.org/fhir/ValueSet/observation-statistics

URL: https://hl7.org/fhir/R4/valueset-observation-statistics.html
Orders and Observations Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/observation-statistics
Version: | 4.0.1
Name: | StatisticsCode
Title: | StatisticsCode
Definition: | The statistical operation parameter -"statistic" codes.
Committee: | Orders and Observations Work Group
OID: | 2.16.840.1.113883.4.642.3.405 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for StatisticsCode

4.4.1.624.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/observation-statistics

4.4.1.624.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 21 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/observation-statistics version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/observation-statistics

Code | Display | Definition
average | Average | The [mean](https://en.wikipedia.org/wiki/Arithmetic_mean) of N measurements over the stated period.
maximum | Maximum | The [maximum](https://en.wikipedia.org/wiki/Maximal_element) value of N measurements over the stated period.
minimum | Minimum | The [minimum](https://en.wikipedia.org/wiki/Minimal_element) value of N measurements over the stated period.
count | Count | The [number] of valid measurements over the stated period that contributed to the other statistical outputs.
total-count | Total Count | The total [number] of valid measurements over the stated period, including observations that were ignored because they did not contain valid result values.
median | Median | The [median](https://en.wikipedia.org/wiki/Median) of N measurements over the stated period.
std-dev | Standard Deviation | The [standard deviation](https://en.wikipedia.org/wiki/Standard_deviation) of N measurements over the stated period.
sum | Sum | The [sum](https://en.wikipedia.org/wiki/Summation) of N measurements over the stated period.
variance | Variance | The [variance](https://en.wikipedia.org/wiki/Variance) of N measurements over the stated period.
20-percent | 20th Percentile | The 20th [Percentile](https://en.wikipedia.org/wiki/Percentile) of N measurements over the stated period.
80-percent | 80th Percentile | The 80th [Percentile](https://en.wikipedia.org/wiki/Percentile) of N measurements over the stated period.
4-lower | Lower Quartile | The lower [Quartile](https://en.wikipedia.org/wiki/Quartile) Boundary of N measurements over the stated period.
4-upper | Upper Quartile | The upper [Quartile](https://en.wikipedia.org/wiki/Quartile) Boundary of N measurements over the stated period.
4-dev | Quartile Deviation | The difference between the upper and lower [Quartiles](https://en.wikipedia.org/wiki/Quartile) is called the Interquartile range. (IQR = Q3-Q1) Quartile deviation or Semi-interquartile range is one-half the difference between the first and the third quartiles.
5-1 | 1st Quintile | The lowest of four values that divide the N measurements into a frequency distribution of five classes with each containing one fifth of the total population.
5-2 | 2nd Quintile | The second of four values that divide the N measurements into a frequency distribution of five classes with each containing one fifth of the total population.
5-3 | 3rd Quintile | The third of four values that divide the N measurements into a frequency distribution of five classes with each containing one fifth of the total population.
5-4 | 4th Quintile | The fourth of four values that divide the N measurements into a frequency distribution of five classes with each containing one fifth of the total population.
skew | Skew | Skewness is a measure of the asymmetry of the probability distribution of a real-valued random variable about its mean. The skewness value can be positive or negative, or even undefined. Source: [Wikipedia](https://en.wikipedia.org/wiki/Skewness).
kurtosis | Kurtosis | Kurtosis is a measure of the "tailedness" of the probability distribution of a real-valued random variable. Source: [Wikipedia](https://en.wikipedia.org/wiki/Kurtosis).

regression | Regression | Linear regression is an approach for modeling two-dimensional sample points with one independent variable and one dependent variable (conventionally, the x and y coordinates in a Cartesian coordinate system) and finds a linear function (a non-vertical straight line) that, as accurately as possible, predicts the dependent variable values as a function of the independent variables. Source: [Wikipedia](https://en.wikipedia.org/wiki/Simple_linear_regression) This Statistic code will return both a gradient and an intercept value.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.625 Value Set http://hl7.org/fhir/ValueSet/container-material

URL: https://hl7.org/fhir/R4/valueset-container-material.html
Orders and Observations Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/container-material
Version: | 4.0.1
Name: | ContainerMaterials
Title: | Types of material for specimen containers
Definition: | This value set includes SNOMED CT codes for materials that specimen containers are made of
Committee: | Orders and Observations Work Group
OID: | 2.16.840.1.113883.4.642.3.1400 (for OID based terminology systems)
Copyright: | This resource includes content from SNOMED Clinical Terms® (SNOMED CT®) which is copyright of the International Health Terminology Standards Development Organisation (IHTSDO). Implementers of these specifications must have the appropriate SNOMED CT Affiliate license - for more information contact http://www.snomed.org/snomed-ct/get-snomed-ct or info@snomed.org
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: SpecimenDefinition.typeTested.container.material (CodeableConcept / Example)

4.4.1.625.1 Content Logical Definition

This value set includes codes from the following code systems:

http://snomed.info/sct

Code | Display
32039001 | glass
61088005 | plastic
425620007 | metal

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.626 Value Set http://hl7.org/fhir/ValueSet/participant-role

URL: https://hl7.org/fhir/R4/valueset-participant-role.html
Patient Care Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/participant-role
Version: | 4.0.1
Name: | ParticipantRoles
Title: | Participant Roles
Definition: | Roles of participants that may be included in a care team. Defined as: Is a Person, Healthcare professional (occupation) or Healthcare related organization (qualifier value).
Committee: | Patient Care Work Group
OID: | 2.16.840.1.113883.4.642.3.152 (for OID based terminology systems)
Copyright: | This resource includes content from SNOMED Clinical Terms® (SNOMED CT®) which is copyright of the International Health Terminology Standards Development Organisation (IHTSDO). Implementers of these specifications must have the appropriate SNOMED CT Affiliate license - for more information contact http://www.snomed.org/snomed-ct/get-snomed-ct or info@snomed.org
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: ServiceRequest.performerType (CodeableConcept / Example)
• Profile: ServiceRequest-Genetics: ServiceRequest.performerType (CodeableConcept / Example)
• Resource: CareTeam.participant.role (CodeableConcept / Example)
• Resource: DeviceRequest.performerType (CodeableConcept / Example)

4.4.1.626.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include codes from http://snomed.info/sct where concept is-a 125676002 (Person)
• Include codes from http://snomed.info/sct where concept is-a 223366009 (Healthcare professional)
• Include codes from http://snomed.info/sct where concept is-a 394730007 (Healthcare related organisation)

4.4.1.626.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 1006 concepts

All codes from system http://snomed.info/sct

Code | Display | Definition
270002 | Female first cousin
375005 | Sibling
444000 | Legal sibling
609005 | Adoptive father
1354005 | Orphan female
2272004 | Half-sister
2316007 | Adoptive son
2368000 | Great-great grandmother
2481008 | Working mother
2959006 | Female cousin
3425009 | Oldest daughter
3851003 | Surrogate daughter
4577005 | First cousin
6676009 | Youngest daughter
8458002 | Foster father
8674003 | Step son
9306000 | Legal parent
9947008 | Natural father
10896006 | Identical twin sibling
10960006 | Working father
11286003 | Twin sibling
11393001 | Great-great grand child
11434005 | Male second cousin
11773006 | Legal brother
11993008 | Male first cousin
12241003 | Foster son
12629003 | Single mother
13038009 | Older brother
13157002 | Older sibling
13443008 | Second cousin
13646006 | Natural parent
14469008 | Legal child
15130002 | Surrogate parent
17219007 | Male fiance
17925003 | Adoptive brother
17945006 | Natural grandmother
18205005 | Wesleyan Methodist, follower of religion

18906004 | Foster sibling
19343003 | Twin sister
19686009 | Younger sister
21093007 | Half-sibling
21464003 | Adoptive mother
21506002 | Female second cousin
22387007 | Surrogate child
22573006 | Stepdaughter
22609000 | Adoptive grandparent
22963000 | Legal sister
25211005 | Aunt
27508009 | Surrogate mother
27733009 | Sr - Sister
28010004 | Shiite muslim, follower of religion
29539002 | Younger sibling
29644004 | Fraternal twin sister
29787005 | Foster brother
30578000 | Stepfather
31656007 | Adoptive grandmother
31831004 | Foster daughter
33969000 | Great-grandparent
34581001 | Niece
34871008 | Grandfather
34972000 | Only daughter
38048003 | Uncle
38248007 | Oldest son
38265003 | Foster mother
38312007 | Grandparent
39062003 | Foster child
40683002 | Parent
41057000 | Surrogate son
41795004 | Legal son
41953004 | Adoptive parent
44181008 | Granddaughter
45929001 | Half-brother
46363003 | Stepsister
47801002 | Male cousin
48385004 | Acquaintance
48673000 | Adoptive child
50058005 | Identical twin sister
50261002 | Great-grandfather
51616000 | Sephardic Jew, follower of religion
53201003 | Older sister
54056000 | Trustee
55538000 | Cousin
58293006 | Foster sister
58626002 | Legal guardian
60614009 | Natural brother
62090008 | Legal mother
62296006 | Natural grand-father
64988008 | Sunni muslim, follower of religion
65412001 | Step-mother
65616008 | Son
65656005 | Natural mother
65853000 | Student
66089001 | Daughter
66839005 | Father
67005001 | Adoptive daughter
67147004 | Legal father
67822003 | Child
68021009 | Great-great grand-parent
70578009 | Grand son
70862002 | Contact person
70924004 | Brother
72012000 | Great grand child
72705000 | Mother
73678001 | Natural sister
74128007 | Step parent
74489007 | Younger child
75226009 | Natural child
75615008 | Surrogate father
76022008 | Adoptive sister

76087000 | Step brother
78194006 | Identical twin brother
78272007 | Natural grand-parent
78652007 | Great grand-mother
78872000 | Orphan male
79508001 | Adoptive sibling
79756005 | Boyfriend
80386000 | Great-great grand-father
80587008 | Jehovah's Witness, follower of religion
81173009 | Only son
81276006 | Twin brother
81467001 | Fraternal twin brother
81706006 | Ashkenazi Jew
82101005 | Natural sibling
83408003 | Working parent
83420006 | Natural daughter
83559000 | Nephew
84390006 | Girlfriend
85058002 | Adoptive grandfather
85411004 | Younger brother
85683001 | Single father
86372007 | Grand child
86764008 | Step child
88510002 | Step sibling
89681006 | Youngest son
90921004 | Foster parent
105428002 | Homosexual parents
105429005 | Elderly parents
105431001 | Adopted person
105436006 | Elderly person
105437002 | Elderly man
105438007 | Elderly woman
105449001 | Sick relative
105450001 | Alcoholic relative
105451002 | Demented relative
105452009 | Diabetic relative
105453004 | Schizophrenic relative
105454005 | Disabled relative
105455006 | Donor for medical or surgical procedure
105456007 | Live donor
105457003 | Cadaver donor
105458008 | Candidate donor
105459000 | Accepted donor
105460005 | Rejected donor
105461009 | Organ donor
105462002 | Donor of kidney for transplant
105463007 | Donor for liver transplant
105464001 | Eye donor for corneal transplant
105465000 | Donor for heart transplant
105466004 | Donor for lung transplant
105467008 | Donor for heart-lung transplant
105468003 | Tissue donor
105469006 | Bone tissue donor
105470007 | Blood donor
105471006 | Bone marrow donor
105472004 | Sperm donor for artificial insemination
105473009 | Husband sperm donor for artificial insemination
105474003 | Related sperm donor for artificial insemination
113157001 | Grandmother
113158006 | Fraternal twin sibling
113159003 | Legal daughter
113160008 | Natural son
113161007 | Female fiance
113163005 | Friend
116153009 | Donor of control material
116154003 | Patient
125677006 | Relative
125678001 | First degree blood relative
125679009 | Blood relative
127848009 | Spouse
127849001 | Husband
127850001 | Wife

127851002 | Fiance
133931009 | Infant
133932002 | Caregiver
133933007 | Newborn
133936004 | Adult
133937008 | Adolescent
160234004 | Salvation Army member
160498000 | School child
160505009 | Common-law husband
160506005 | Common-law wife
160540005 | Roman Catholic
160542002 | Atheist
160543007 | Jew
160544001 | Christian Science Church
160545000 | Hindu
160549006 | Christian
160557009 | Baptist
160558004 | Methodist
160560002 | Presbyterian
160562005 | Pentecostal
160563000 | Evangelist
160566008 | Christadelphian
160567004 | Agnostic
160772005 | Voluntary worker
161158003 | Immigrant
224080001 | Committed Christian, follower of religion
224526002 | Woman
224527006 | Person in the work environment
224528001 | Employee
224619008 | Migrant
224627004 | General category of person
248544006 | Rastafarian, follower of religion
255409004 | Pregnant woman
257497009 | Animal rider
257498004 | Rider of motorcycle
257499007 | Pedal cyclist
257500003 | Passenger
257501004 | Occupant of motor vehicle
257502006 | Occupant of tram
257503001 | Pillion passenger
257504007 | Person riding on outside of vehicle
257505008 | Person travelling on outside of train
257506009 | Front seat passenger
257508005 | Back seat passenger
257509002 | Occupant of animal-drawn vehicle
257510007 | Passenger on commercial watercraft
257511006 | Passenger on station
257512004 | Passenger on train
257513009 | Member of public
257514003 | Member of public involved incidentally
257516001 | Member of public on ground
257517005 | Occupant of nonmotor vehicle
257518000 | Pedestrian
257519008 | Person trespassing on railway line
257521003 | Railway employee on train
257522005 | Recreational watercraft user
257523000 | Swimmer
257524006 | User of animal-drawn vehicle
257525007 | Watercraft dweller
257526008 | Waterskier
262043009 | Companion
266943008 | Single parent
268557009 | Sportsman
271448006 | Anglican, follower of religion
276035002 | Voluntary helper
276036001 | Voluntary visitor
276037005 | Volunteer helper
276119007 | Nonconformist
303069001 | Person in the transport environment
303071001 | Person in the family
303072008 | Person in the home environment
303073003 | Person in the religious environment

303118004 | Person in the healthcare environment
303119007 | Person in the community environment
303120001 | Person in the recreational environment
303121002 | Person in the educational environment
303122009 | Crewmember
303977004 | Occupant of railway vehicle
303980003 | Driver of motor vehicle
303981004 | Driver of animal-drawn vehicle
303983001 | Passenger of animal-drawn vehicle
303984007 | Passenger on pedal cycle
303985008 | Occupant of watercraft
303986009 | Occupant of aircraft
303987000 | Horse rider
303988005 | Passenger on aircraft
303989002 | Crewmember on aircraft
303990006 | Passenger of motor vehicle
303991005 | Trespasser
303992003 | Car passenger
303993008 | Coach passenger
309456003 | Clergy
309687009 | Buddhist, follower of religion
309884000 | Muslim, follower of religion
309885004 | Protestant, follower of religion
309886003 | Quaker, follower of religion
310543008 | Graduate
313415001 | Identical twin
313416000 | Heterozygous twin
314712007 | Parachutist
314767008 | Bird fancier
339947000 | Man
360852005 | Bum
360863008 | Beggar
360984002 | Hasidic Jew, follower of religion
366740002 | Sikh, follower of religion
373864002 | Outpatient
373929005 | Orphan child
385435006 | Individual
390790000 | Asylum seeker
394566006 | Great-grandson
394567002 | Great-granddaughter
394568007 | Extended family member
394571004 | Employer
394619001 | Guardian
394738000 | Other related persons
394854006 | Immediate family member
394855007 | Non-immediate family member
394856008 | Paternal grand-father
394857004 | Maternal grand-father
394858009 | Paternal grand-mother
394859001 | Maternal grand-mother
394860006 | Distant relative
394861005 | Great-uncle
394862003 | Great-aunt
394863008 | Non-family member
394921008 | Cohabitee
407542009 | Informal carer
407543004 | Primary carer
410597007 | Person categorized by religious affiliation
410598002 | Person categorized by age
410599005 | Minor
410600008 | Pre-teen
410601007 | Child
410602000 | Preschool child
410603005 | Person in the legal environment
410604004 | Subject of record
413327003 | Student in full time education
414043009 | Domestic partner
416034003 | Primary screener
416035002 | Secondary screener
416625007 | Family reunion immigrant
416800000 | Inpatient
418119000 | Infant child

419358007 | Subject of record or other provider of history
420058008 | Provider of history other than subject
420158005 | Performer of method
427568008 | Neighbor
427729003 | Elim Pentecostalist, follower of religion
427754003 | Liberal Jew
427755002 | Lutheran, follower of religion
427874000 | Congregationalist, follower of religion
427950002 | Haredi Jew
427963008 | Advaitin Hindu, follower of religion
427981006 | Zen Buddhist
428024001 | Clinical trial participant
428315004 | Anabaptist, follower of religion
428347009 | Arya Samaj Hindu
428373004 | Coptic Orthodox, follower of religion
428376007 | Eastern Catholic
428378008 | Ethiopian Orthodox Tewahedo, follower of religion
428407001 | Theravada Buddhist
428408006 | Wiccan
428410008 | Shakti Hindu
428496003 | Tibetan Buddhist, follower of religion
428503004 | Orthodox Christian
428504005 | Christian Spiritualist, follower of religion
428506007 | Druid, follower of religion
428620006 | Smarta Hindu
428666001 | Nichiren Buddhist
428792000 | Case contact
428801007 | Ismaili Muslim
428815009 | Ahmadi, follower of religion
428816005 | Armenian Catholic
428820009 | Pure Land Buddhist, follower of religion
428821008 | Seventh Day Adventist
429158002 | Greek Orthodox
429171004 | Spiritualist
429379008 | Shiva Hindu, follower of religion
429509008 | Reform Jew, follower of religion
429511004 | Vaishnava Hindu
429527006 | New Kadampa Tradition Buddhist
429533002 | Mahayana Buddhist
429534008 | Masorti Jew, follower of religion
429539003 | Calvinist
429543004 | Taoist, follower of religion
429544005 | Russian Orthodox, follower of religion
429545006 | Sanatana Dharma follower
429547003 | Shinto
429577009 | Patient advocate
429644000 | Orthodox Jew, follower of religion
429708003 | Unitarian, follower of religion
429732005 | Baha'i
429787006 | Jain, follower of religion
429790000 | Zoroastrian, follower of religion
430098004 | Donor for intestine transplant
430101000 | Whole blood donor
430792005 | Stem cell donor
430882007 | Egg donor
431549007 | Female child
433152004 | Boy
433475001 | Human oocyte donor, age 35 and older
438347000 | Person categorized by affiliation with belief system
441463002 | Sperm donor
441465009 | Stranger is sperm donor for artificial insemination
441468006 | Unrelated known sperm donor for artificial insemination
444018008 | Person with characteristic related to subject of record
444052006 | Spouse of subject
444053001 | Husband of subject
444054007 | Wife of subject
444055008 | Uncle of subject
444145006 | Partner in relationship with subject
444146007 | Boyfriend of subject
444147003 | Girlfriend of subject
444148008 | Person in family of subject
444168002 | Homemaker

444169005 | House husband
444191003 | Aunt of subject
444192005 | Child of subject
444193000 | First degree blood relative of subject
444194006 | Daughter of subject
444199001 | Friend of subject
444210007 | Cohabitee in relationship with subject
444211006 | Common law husband of subject
444212004 | Common law wife of subject
444213009 | Domestic partner of subject
444241008 | Son of subject
444242001 | Grandparent of subject
444243006 | Maternal grandfather of subject
444244000 | Maternal grandmother of subject
444278006 | Fiance of subject
444279003 | Female fiance of subject
444280000 | Male fiance of subject
444292000 | Paternal grandfather of subject
444293005 | Paternal grandmother of subject
444294004 | Parent of subject
444295003 | Father of subject
444301002 | Mother of subject
444302009 | Sibling of subject
444303004 | Brother of subject
444304005 | Sister of subject
444464007 | Recipient of biological material from subject
444870008 | Apostolic Pentecostalist, follower of religion
446654005 | Refugee
450768005 | International student
699110007 | Second degree blood relative
699111006 | Second degree blood relative of subject
713176007 | Cadaver donor for lung transplant
713177003 | Cadaver donor for heart transplant
713178008 | Cadaver donor for liver transplant
713179000 | Cadaver donor for corneal transplant
713183000 | Cadaver donor for kidney transplant
713184006 | Blood product donor
713352007 | Skin donor
713353002 | Live related liver donor
713369009 | Live related kidney donor
715167002 | Open Episcopal Church follower of religion
471000122100 | Multi-organ donor
100217091000087103 | Maternal aunt
166557991000087101 | Maternal great grand-father
173704201000087107 | Paternal first cousin
194907391000087109 | Maternal great grand-mother
215424891000087101 | Paternal great grand-mother
286864561000087100 | Maternal cousin
396762341000087100 | Paternal cousin
398524981000087101 | Paternal great grand-father
583114651000087104 | Maternal uncle
589612571000087102 | Paternal uncle
691349461000087102 | Paternal aunt
844670641000087109 | Maternal first cousin
1421009 | Specialized surgeon
3430008 | Radiation therapist
3842006 | Chiropractor
4162009 | Dental assistant
5275007 | NA - Nursing auxiliary
6816002 | Specialized nurse
6868009 | Hospital administrator
8724009 | Plastic surgeon
11661002 | Neuropathologist
11911009 | Nephrologist
11935004 | Obstetrician
13580004 | School dental assistant
14698002 | Medical microbiologist
17561000 | Cardiologist
18803008 | Dermatologist
18850004 | Laboratory hematologist
19244007 | Gerodontist
20145008 | Removable prosthodontist

21365001 | Specialized dentist
21450003 | Neuropsychiatrist
22515006 | Medical assistant
22731001 | Orthopedic surgeon
22983004 | Thoracic surgeon
23278007 | Community health physician
24430003 | Physical medicine specialist
24590004 | Urologist
25961008 | Electroencephalography specialist
26042002 | Dental hygienist
26369006 | Public health nurse
28229004 | Optometrist
28411006 | Neonatologist
28544002 | Chemical pathologist
36682004 | PT - Physiotherapist
37154003 | Periodontist
37504001 | Orthodontist
39677007 | Internal medicine specialist
40127002 | Dietitian (general)
40204001 | Hematologist
40570005 | Interpreter
41672002 | Respiratory physician
41904004 | Medical X-ray technician
43702002 | Occupational health nurse
44652006 | Pharmaceutical assistant
45419001 | Masseur
45440000 | Rheumatologist
45544007 | Neurosurgeon
45956004 | Sanitarian
46255001 | Pharmacist
48740002 | Philologist
49203003 | Dispensing optometrist
49993003 | Maxillofacial surgeon
50149000 | Endodontist
54503009 | Faith healer
56397003 | Neurologist
56466003 | Community physician
56542007 | Medical record administrator
56545009 | Cardiovascular surgeon
57654006 | Fixed prosthodontist
59058001 | General physician
59169001 | Orthopedic technician
59944000 | Psychologist
60008001 | Community-based dietitian
61207006 | Medical pathologist
61246008 | Laboratory medicine specialist
61345009 | Otorhinolaryngologist
61894003 | Endocrinologist
62247001 | Family medicine specialist
63098009 | Clinical immunologist
66476003 | Oral pathologist
66862007 | Radiologist
68867008 | Public health dentist
68950000 | Prosthodontist
69280009 | Specialized physician
71838004 | Gastroenterologist
73265009 | Nursing aid
75271001 | MW - Midwife
76166008 | Practical aid (pharmacy)
76231001 | Osteopath
76899008 | Infectious diseases physician
78703002 | General surgeon
78729002 | Diagnostic radiologist
79898004 | Auxiliary midwife
80409005 | Translator
80546007 | OT - Occupational therapist
80584001 | Psychiatrist
80933006 | Nuclear medicine physician
81464008 | Clinical pathologist
82296001 | Pediatrician
83189004 | Other professional nurse
83273008 | Anatomic pathologist

83685006 | Gynecologist
85733003 | General pathologist
88189002 | Anesthesiologist
88475002 | Other dietitians and public health nutritionists
90201008 | Pediatric dentist
90655003 | Care of the elderly physician
106289002 | Dental surgeon
106291005 | Dietician AND/OR public health nutritionist
106292003 | Nurse
106293008 | Nursing personnel
106294002 | Midwifery personnel
106296000 | Physiotherapist AND/OR occupational therapist
106330007 | Philologist, translator AND/OR interpreter
112247003 | Medical doctor
158965000 | Medical practitioner
158966004 | Medical administrator - national
158967008 | Consultant physician
158968003 | Consultant surgeon
158969006 | Consultant gynecology and obstetrics
158970007 | Anesthetist
158971006 | Hospital registrar
158972004 | House officer
158973009 | Occupational physician
158974003 | Clinical medical officer
158975002 | Medical practitioner - teaching
158977005 | Dental administrator
158978000 | Dental consultant
158979008 | Dental general practitioner
158980006 | Dental practitioner - teaching
158983008 | Nurse administrator - national
158984002 | Nursing officer - region
158985001 | Nursing officer - district
158986000 | Nursing administrator - professional body
158987009 | Nursing officer - division
158988004 | Nurse education director
158989007 | Occupational health nursing officer
158990003 | Nursing officer
158992006 | Midwifery sister
158993001 | Nursing sister (theatre)
158994007 | Staff nurse
158995008 | Staff midwife
158996009 | State enrolled nurse
158997000 | District nurse
158998005 | Private nurse
158999002 | Community midwife
159001001 | Clinic nurse
159002008 | Practice nurse
159003003 | School nurse
159004009 | Nurse - teaching
159005005 | Student nurse
159006006 | Dental nurse
159007002 | Community pediatric nurse
159010009 | Hospital pharmacist
159011008 | Retail pharmacist
159012001 | Industrial pharmacist
159013006 | Pharmaceutical officer H.A.
159014000 | Trainee pharmacist
159016003 | Medical radiographer
159017007 | Diagnostic radiographer
159018002 | Therapeutic radiographer
159019005 | Trainee radiographer
159021000 | Ophthalmic optician
159022007 | Trainee optician
159025009 | Remedial gymnast
159026005 | Speech and language therapist
159027001 | Orthoptist
159028006 | Trainee remedial therapist
159033005 | Dietician
159034004 | Podiatrist
159035003 | Dental auxiliary
159036002 | ECG technician
159037006 | EEG technician

159038001 | Artificial limb fitter
159039009 | AT - Audiology technician
159040006 | Pharmacy technician
159041005 | Trainee medical technician
159141008 | Geneticist
159972006 | Surgical corset fitter
160008000 | Dental technician
224529009 | Clinical assistant
224530004 | Senior registrar
224531000 | Registrar
224532007 | Senior house officer
224533002 | MO - Medical officer
224534008 | Health visitor, nurse/midwife
224535009 | Registered nurse
224536005 | Midwifery tutor
224537001 | Accident and Emergency nurse
224538006 | Triage nurse
224540001 | Community nurse
224541002 | Nursing continence advisor
224542009 | Coronary care nurse
224543004 | Diabetic nurse
224544005 | Family planning nurse
224545006 | Care of the elderly nurse
224546007 | ICN - Infection control nurse
224547003 | Intensive therapy nurse
224548008 | Learning disabilities nurse
224549000 | Neonatal nurse
224550000 | Neurology nurse
224551001 | Industrial nurse
224552008 | Oncology nurse
224553003 | Macmillan nurse
224554009 | Marie Curie nurse
224555005 | Pain control nurse
224556006 | Palliative care nurse
224557002 | Chemotherapy nurse
224558007 | Radiotherapy nurse
224559004 | PACU nurse
224560009 | Stomatherapist
224561008 | Theatre nurse
224562001 | Pediatric nurse
224563006 | Psychiatric nurse
224564000 | Community mental health nurse
224565004 | Renal nurse
224566003 | Hemodialysis nurse
224567007 | Wound care nurse
224569005 | Nurse grade
224570006 | Clinical nurse specialist
224571005 | Nurse practitioner
224572003 | Nursing sister
224573008 | CN - Charge nurse
224574002 | Ward manager
224575001 | Nursing team leader
224576000 | Nursing assistant
224577009 | Healthcare assistant
224578004 | Nursery nurse
224579007 | Healthcare service manager
224580005 | Occupational health service manager
224581009 | Community nurse manager
224583007 | Behavior therapist
224584001 | Behavior therapy assistant
224585000 | Drama therapist
224586004 | Domiciliary occupational therapist
224587008 | Occupational therapy helper
224588003 | Psychotherapist
224589006 | Community-based physiotherapist
224590002 | Play therapist
224591003 | Play specialist
224592005 | Play leader
224593000 | Community-based speech/language therapist
224594006 | Speech/language assistant
224595007 | Professional counselor
224596008 | Marriage guidance counselor

224597004 | Trained nurse counselor
224598009 | Trained social worker counselor
224599001 | Trained personnel counselor
224600003 | Psychoanalyst
224601004 | Assistant psychologist
224602006 | Community-based podiatrist
224603001 | Foot care worker
224604007 | Audiometrician
224605008 | Audiometrist
224606009 | Technical healthcare occupation
224607000 | Occupational therapy technical instructor
224608005 | Administrative healthcare staff
224609002 | Complementary health worker
224610007 | Supporting services personnel
224614003 | Research associate
224615002 | Research nurse
224620002 | Human aid to communication
224621003 | Palantypist
224622005 | Note taker
224623000 | Cuer
224624006 | Lipspeaker
224625007 | Interpreter for British sign language
224626008 | Interpreter for Signs supporting English
224936003 | General practitioner locum
225726006 | Lactation consultant
225727002 | Midwife counselor
265937000 | Nursing occupation
265939002 | Medical/dental technicians
283875005 | Parkinson disease nurse
302211009 | Specialist registrar
303124005 | Member of mental health review tribunal
303129000 | Hospital manager
303133007 | Responsible medical officer
303134001 | Independent doctor
304291006 | Bereavement counselor
304292004 | Surgeon
307988006 | Medical technician
308002005 | Remedial therapist
309294001 | Accident and Emergency doctor
309295000 | Clinical oncologist
309296004 | Family planning doctor
309322005 | Associate general practitioner
309323000 | Partner of general practitioner
309324006 | Assistant GP
309326008 | Deputizing general practitioner
309327004 | General practitioner registrar
309328009 | Ambulatory pediatrician
309329001 | Community pediatrician
309330006 | Pediatric cardiologist
309331005 | Pediatric endocrinologist
309332003 | Pediatric gastroenterologist
309333008 | Pediatric nephrologist
309334002 | Pediatric neurologist
309335001 | Pediatric rheumatologist
309336000 | Pediatric oncologist
309337009 | Pain management specialist
309338004 | Intensive care specialist
309339007 | Adult intensive care specialist
309340009 | Pediatric intensive care specialist
309341008 | Blood transfusion doctor
309342001 | Histopathologist
309343006 | Physician
309345004 | Chest physician
309346003 | Thoracic physician
309347007 | Clinical hematologist
309348002 | Clinical neurophysiologist
309349005 | Clinical physiologist
309350005 | Diabetologist
309351009 | Andrologist
309352002 | Neuroendocrinologist
309353007 | Reproductive endocrinologist
309354001 | Thyroidologist

309355000 | Clinical geneticist
309356004 | Clinical cytogeneticist
309357008 | Clinical molecular geneticist
309358003 | Genitourinary medicine physician
309359006 | Palliative care physician
309360001 | Rehabilitation physician
309361002 | Child and adolescent psychiatrist
309362009 | Forensic psychiatrist
309363004 | Liaison psychiatrist
309364005 | Psychogeriatrician
309365006 | Psychiatrist for mental handicap
309366007 | Rehabilitation psychiatrist
309367003 | Obstetrician and gynecologist
309368008 | Breast surgeon
309369000 | Cardiothoracic surgeon
309371000 | Cardiac surgeon
309372007 | Ear, nose and throat surgeon
309373002 | Endocrine surgeon
309374008 | Thyroid surgeon
309375009 | Pituitary surgeon
309376005 | Gastrointestinal surgeon
309377001 | General gastrointestinal surgeon
309378006 | Upper gastrointestinal surgeon
309379003 | Colorectal surgeon
309380000 | Hand surgeon
309381001 | Hepatobiliary surgeon
309382008 | Ophthalmic surgeon
309383003 | Pediatric surgeon
309384009 | Pancreatic surgeon
309385005 | Transplant surgeon
309386006 | Trauma surgeon
309388007 | Vascular surgeon
309389004 | Medical practitioner grade
309390008 | Hospital consultant
309391007 | Visiting specialist registrar
309392000 | Research registrar
309393005 | General practitioner grade
309394004 | General practitioner principal
309395003 | Hospital specialist
309396002 | Associate specialist
309397006 | Research fellow
309398001 | Allied health professional
309399009 | Hospital dietitian
309400002 | Domiciliary physiotherapist
309401003 | General practitioner-based physiotherapist
309402005 | Hospital-based physiotherapist
309403000 | Private physiotherapist
309404006 | Physiotherapy assistant
309409001 | Hospital-based speech and language therapist
309410006 | Arts therapist
309411005 | Dance therapist
309412003 | Music therapist
309413008 | Renal dietitian
309414002 | Liver dietitian
309415001 | Oncology dietitian
309416000 | Pediatric dietitian
309417009 | Diabetes dietitian
309418004 | Audiologist
309419007 | Hearing therapist
309420001 | Audiological scientist
309421002 | Hearing aid dispenser
309422009 | Community-based occupational therapist
309423004 | Hospital occupational therapist
309427003 | Social services occupational therapist
309428008 | Orthotist
309429000 | Surgical fitter
309434001 | Hospital-based podiatrist
309435000 | Podiatry assistant
309436004 | Lymphedema nurse
309437008 | Community learning disabilities nurse
309439006 | Clinical nurse teacher
309440008 | Community practice nurse teacher

309441007 | Nurse tutor
309442000 | Nurse teacher practitioner
309443005 | Nurse lecturer practitioner
309444004 | Outreach nurse
309445003 | Anesthetic nurse
309446002 | Nurse manager
309450009 | Nurse administrator
309452001 | Midwifery grade
309453006 | Midwife
309454000 | Student midwife
309455004 | Parentcraft sister
309459005 | Healthcare professional grade
309460000 | Restorative dentist
310170009 | Pediatric audiologist
310171008 | Immunopathologist
310172001 | Audiological physician
310173006 | Clinical pharmacologist
310174000 | Private doctor
310175004 | Agency nurse
310176003 | Behavioral therapist nurse
310177007 | Cardiac rehabilitation nurse
310178002 | Genitourinary nurse
310179005 | Rheumatology nurse specialist
310180008 | Continence nurse
310181007 | Contact tracing nurse
310182000 | General nurse
310183005 | Nurse for the mentally handicapped
310184004 | Liaison nurse
310185003 | Diabetic liaison nurse
310186002 | Nurse psychotherapist
310187006 | Company nurse
310188001 | Hospital midwife
310189009 | Genetic counselor
310190000 | Mental health counselor
310191001 | Clinical psychologist
310192008 | Educational psychologist
310193003 | Coroner
310194009 | Appliance officer
310512001 | Medical oncologist
311441001 | School medical officer
312485001 | Integrated midwife
372102007 | RN First Assist
387619007 | Optician
394572006 | Medical secretary
394618009 | Hospital nurse
397824005 | Consultant anesthetist
397897005 | Paramedic
397903001 | Staff grade obstetrician
397908005 | Staff grade practitioner
398130009 | Medical student
398238009 | Acting obstetric registrar
404940000 | Physiotherapist technical instructor
405277009 | Resident physician
405278004 | Certified registered nurse anesthetist
405279007 | Attending physician
405623001 | Assigned practitioner
405684005 | Professional initiating surgical case
405685006 | Professional providing staff relief during surgical procedure
408798009 | Consultant pediatrician
408799001 | Consultant neonatologist
409974004 | Health educator
409975003 | Certified health education specialist
413854007 | Circulating nurse
415075003 | Perioperative nurse
415506007 | Scrub nurse
416160000 | Fellow of American Academy of Osteopathy
420409002 | Oculoplastic surgeon
420678001 | Retinal surgeon
421841007 | Admitting physician
422140007 | Medical ophthalmologist
422234006 | Ophthalmologist
432100008 | Health coach

442867008 | Respiratory therapist
443090005 | Podiatric surgeon
444912007 | Hypnotherapist
445313000 | Asthma nurse specialist
445451001 | Nurse case manager
446050000 | PCP - Primary care physician
446701002 | Addiction medicine specialist
449161006 | PA - physician assistant
471302004 | Government midwife
3981000175106 | Nurse complex case manager
231189271000087109 | Naturopath
236749831000087105 | Prosthetist
258508741000087105 | Hip and knee surgeon
260767431000087107 | Hepatologist
285631911000087106 | Shoulder surgeon
291705421000087106 | Interventional radiologist
341320851000087105 | Pediatric radiologist
368890881000087105 | Emergency medicine specialist
398480381000087106 | Family medicine specialist - palliative care
416186861000087101 | Surgical oncologist
450044741000087104 | Acupuncturist
465511991000087105 | Pediatric orthopedic surgeon
494782281000087101 | Pediatric hematologist
619197631000087102 | Neuroradiologist
623630151000087105 | Family medicine specialist - anesthetist
666997781000087107 | Doula
673825031000087109 | Traditional herbal medicine specialist
682131381000087105 | Occupational medicine specialist
724111801000087104 | Pediatric emergency medicine specialist
747936471000087102 | Family medicine specialist - care of the elderly
766788081000087100 | Travel medicine specialist
767205061000087108 | Spine surgeon
813758161000087106 | Maternal or fetal medicine specialist
822410621000087104 | Massage therapist
847240411000087102 | Hospitalist
853827051000087104 | Sports medicine specialist
926871431000087103 | Pediatric respirologist
954544641000087107 | Homeopath
956387501000087102 | Family medicine specialist - emergency medicine
969118571000087109 | Pediatric hematologist or oncologist
984095901000087105 | Foot and ankle surgeon
990928611000087105 | Invasive cardiologist
999480451000087102 | Case manager
999480461000087104 | Kinesthesiologist
394745000 | General practice (organisation)
394746004 | Regional office
394747008 | Health Authority
394748003 | Primary Care Group
394749006 | Primary Care Trust
394751005 | Special Health Authority
394757009 | Hospice - independent
394758004 | Independent provider
394760002 | GP practice
394762005 | GDP practice
394763000 | MoD practice
394764006 | MoD hospital
394765007 | Prison practice
398154004 | Quality Improvement Template Task Force
398255008 | Association of anesthesia clinical directors
5581000087108 | Canadian jurisdiction-based
8051000087100 | Jurisdiction driver license number
8061000087102 | Jurisdictional birth certificate registration number
8071000087106 | Interim health program certificate number
183033871000087106 | Ministry of Health Saskatchewan
189542671000087101 | Ministry of Health Quebec
249236551000087105 | Community health center
250624661000087107 | Nova Scotia Department of Health and Wellness
254693751000087105 | Nunavut Department of Health and Social Services
287697331000087102 | Provider billing number
317817411000087104 | Independent health care facility
357174231000087105 | Health regulatory body for occupational therapists
371439331000087102 | Health regulatory body for physiotherapists

392685081000087107 | Health regulatory body for dental hygienists
420314821000087103 | Health regulatory body for massage therapists
432233171000087103 | Jurisdictional health number
437208901000087109 | Provincial and/or territorial government plan - out of jurisdiction
439182891000087102 | Provider number issued by third party
441144831000087108 | Health regulatory body for physicians and surgeons
443470051000087109 | Ministry of health
460276491000087102 | New Brunswick Department of Health
539773351000087101 | Alberta Health and Wellness
548886741000087101 | Health regulatory body for denturists
549577861000087101 | Health regulatory body for chiropractors
553641311000087101 | Newfoundland and Labrador Department of Health and Community Services
567496441000087102 | Cancer Plan (Saskatchewan) and Sexually Transmitted Diseases Plan
569232871000087107 | Yukon Department of Health and Social Services
609564621000087104 | Health regulatory body for pharmacists
643545671000087106 | Health regulatory body for social workers
644191431000087105 | Ministry of Health and Long-Term Care Ontario
666298511000087107 | Health regulatory body for midwives
673433971000087102 | Provider registration number
724022191000087108 | Health regulatory body for respiratory therapists
726220521000087101 | Provincial and/or territorial government plan - resident
745424761000087106 | Workers' Compensation number
753977441000087109 | Ministry of Health Northwest Territories
762510691000087106 | Health regulatory body for audiologists and speech-language pathologists
859430851000087100 | Medical Care Plan - Hospital Technical Services (Quebec)
874107021000087108 | Health regulatory body for nurses
876612791000087101 | Health regulatory body for paramedics
882121321000087100 | Ministry of Health British Columbia
912320101000087109 | Health regulatory body for practical nurses
930012821000087105 | Health regulatory body for dietitians
998711551000087102 | Health PEI
998820271000087109 | Manitoba Health
999483971000087108 | Health regulatory body for naturopathic practitioners

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.627 Value Set http://hl7.org/fhir/ValueSet/immunization-recommendation-date-criterion

URL: https://hl7.org/fhir/R4/valueset-immunization-recommendation-date-criterion.html
Public Health and Emergency Response Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/immunization-recommendation-date-criterion
Version: | 4.0.1
Name: | ImmunizationRecommendationDateCriterionCodes
Title: | Immunization Recommendation Date Criterion Codes
Definition: | The value set to instantiate this attribute should be drawn from a terminologically robust code system that consists of or contains concepts to support the definition of dates relevant to recommendations for future doses of vaccines. This value set is provided as a suggestive example.
Committee: | Public Health and Emergency Response Work Group
OID: | 2.16.840.1.113883.4.642.3.308 (for OID based terminology systems)
Copyright: | This content from LOINC® is copyright © 1995 Regenstrief Institute, Inc. and the LOINC Committee, and available at no cost under the license at http://loinc.org/terms-of-use.
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: ImmunizationRecommendation.recommendation.dateCriterion.code (CodeableConcept / Example)

4.4.1.627.1 Content Logical Definition

This value set includes codes from the following code systems:

http://loinc.org

Code | Display
30981-5 | Earliest date to give
30980-7 | Date vaccine due
59777-3 | Latest date to give immunization
59778-1 | Date when overdue for immunization

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.628 Value Set http://hl7.org/fhir/ValueSet/choice-list-orientation

URL: https://hl7.org/fhir/R4/valueset-choice-list-orientation.html
FHIR Infrastructure Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/choice-list-orientation
Version: | 4.0.1
Name: | ChoiceListOrientation
Title: | ChoiceListOrientation
Definition: | Direction in which lists of possible answers should be displayed.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.933 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ChoiceListOrientation

4.4.1.628.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/choice-list-orientation

4.4.1.628.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 2 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/choice-list-orientation version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/choice-list-orientation

Code | Display | Definition
horizontal | Horizontal | List choices along the horizontal axis.
vertical | Vertical | List choices down the vertical axis.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.629 Value Set http://hl7.org/fhir/ValueSet/diagnostic-service-sections

URL: https://hl7.org/fhir/R4/valueset-diagnostic-service-sections.html
Orders and Observations Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/diagnostic-service-sections
Version: | 4.0.1
Name: | DiagnosticServiceSectionCodes
Title: | Diagnostic Service Section Codes
Definition: | This value set includes all the codes in HL7 V2 table 0074.
Committee: | Orders and Observations Work Group
OID: | 2.16.840.1.113883.4.642.3.234 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: DiagnosticReport.category (CodeableConcept / Example)
• Profile: LipidProfile: DiagnosticReport.category (CodeableConcept / Example)
• Profile: DiagnosticReport-Genetics: DiagnosticReport.category (CodeableConcept / Example)
• Profile: HLAResult: DiagnosticReport.category (CodeableConcept / Example)

4.4.1.629.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/v2-0074

4.4.1.629.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 41 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/v2-0074 version 2.9

All codes from system http://terminology.hl7.org/CodeSystem/v2-0074

Code | Display | Definition
AU | Audiology
BG | Blood Gases
BLB | Blood Bank
CG | Cytogenetics
CH | Chemistry
CP | Cytopathology
CT | CAT Scan
CTH | Cardiac Catheterization
CUS | Cardiac Ultrasound
EC | Electrocardiac (e.g., EKG, EEC, Holter)
EN | Electroneuro (EEG, EMG,EP,PSG)
GE | Genetics
HM | Hematology
ICU | Bedside ICU Monitoring
IMM | Immunology
LAB | Laboratory
MB | Microbiology
MCB | Mycobacteriology
MYC | Mycology
NMR | Nuclear Magnetic Resonance
NMS | Nuclear Medicine Scan
NRS | Nursing Service Measures
OSL | Outside Lab
OT | Occupational Therapy
OTH | Other
OUS | OB Ultrasound
PF | Pulmonary Function
PHR | Pharmacy
PHY | Physician (Hx. Dx, admission note, etc.)
PT | Physical Therapy
RAD | Radiology
RC | Respiratory Care (therapy)
RT | Radiation Therapy
RUS | Radiology Ultrasound

RX | Radiograph
SP | Surgical Pathology
SR | Serology
TX | Toxicology
VR | Virology
VUS | Vascular Ultrasound
XRC | Cineradiograph

Additional Language Displays

Code | Deutsch (German, de) | Nederlands (Dutch, nl)
AU | Audiologie | Audiologie
BG | Blutgase | Bloedgassen
BLB | Blutbank | Bloedbank
CG | | Cytogenetica
CH | Klinische Chemie | Chemie
CP | Zellpathologie | Cytopathologie
CT | Computertomographie | CAT-scan
CTH | Herzkatheter | Hartcatetherisatie
CUS | Kardiologische Ultraschalluntersuchung | Hart ultrasound
EC | Elektrokardiographie | Electrocardiac (bijv. EKG, EEC, Holter)
EN | Elektroneurographie, Elektroencephalographie | Electroneuro (EEG, EMG,EP,PSG)
GE | Genetik | Genetica
HM | Hämatologie | Hematologie
ICU | | IC monitoring naast het bed
IMM | Immunologie | Immunologie
LAB | Labor | Laboratorium
MB | Mikrobiologie | Microbiologie
MCB | Mykobakteriologie | Mycobacteriologie
MYC | Mykologie | Mycology
NMR | Kernspintomographie (MR) | Nucleaire magnetisch resonantie
NMS | Szintigramm | Nucleaire geneeskunde scan
NRS | Pflegemaßnahme | Verpleegkundige service maatregelen
OSL | Externes Labor | Outside Lab
OT | Beschäftigungstherapie | Ergotherapie
OTH | Andere | Anders
OUS | Geburtshilfiche Ultraschalluntersuchung | OB Ultrasound
PF | Lungenfunktion | Pulmonaire functie
PHR | Apotheke | Medicatie
PHY | Arzt (Krankengeschichte, Diagnose, Aufnahmeuntersuchung) | Arts (Hx. Dx, opnameverslag, etc.)
PT | physiklische Therapie | Lichamelijke oefening
RAD | Radiologie | Radiologie
RC | Atemtherapie | Respiratoire zorg (therapie)
RT | Strahlentherapie | Stralingstherapie
RUS | Radiologische Ultraschalluntersuchung | Radiologie ultrasound
RX | Röntgenaufnahme | Radiograph
SP | Operative Pathologie | Chirurgische pathologie
SR | Serologie | Serologie
TX | Toxikologie | Toxicologie
VR | Virologie | Virologie
VUS | Ultraschalluntersuchung der Gefäße | Vasculaire Ultrasound
XRC | Röntgenkinematographie | Cineradiograph

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.630 Value Set http://hl7.org/fhir/ValueSet/allergyintolerance-code

URL: https://hl7.org/fhir/R4/valueset-allergyintolerance-code.html
Patient Care Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/allergyintolerance-code
Version: | 4.0.1
Name: | AllergyIntoleranceSubstance/Product,ConditionAndNegationCodes
Title: | AllergyIntolerance Substance/Product, Condition and Negation Codes
Definition: | This value set includes concept codes for specific substances/pharmaceutical products, allergy or intolerance conditions, and negation/exclusion codes to specify the absence of specific types of allergies or intolerances.
Committee: | Patient Care Work Group
OID: | 2.16.840.1.113883.4.642.3.137 (for OID based terminology systems)
Copyright: | This resource includes content from SNOMED Clinical Terms® (SNOMED CT®) which is copyright of the International Health Terminology Standards Development Organisation (IHTSDO). Implementers of these specifications must have the appropriate SNOMED CT Affiliate license - for more information contact http://www.snomed.org/snomed-ct/get-snomed-ct or info@snomed.org
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: AllergyIntolerance.code (CodeableConcept / Example)

4.4.1.630.1 Content Logical Definition

This value set includes codes from the following code systems:

• Import all the codes that are contained in http://hl7.org/fhir/ValueSet/substance-code

• Include codes from http://snomed.info/sct where concept is-a 418038007 (Propensity to adverse reactions to substance)

• Include codes from http://snomed.info/sct where concept is-a 267425008 (Lactose intolerance)

• Include codes from http://snomed.info/sct where concept is-a 29736007 (Syndrome of carbohydrate intolerance)

• Include codes from http://snomed.info/sct where concept is-a 340519003 (Lysine intolerance)

• Include codes from http://snomed.info/sct where concept is-a 190753003 (Sucrose intolerance)

• Include codes from http://snomed.info/sct where concept is-a 413427002 (Acquired fructose intolerance)

• Include codes from http://snomed.info/sct where concept is-a 716186003 (No known allergy)

4.4.1.630.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 3016 concepts

Expansion based on http://hl7.org/fhir/ValueSet/substance-code version 4.0.1

All codes from system http://snomed.info/sct

Code | Display | Definition
102002 | Hemoglobin Okaloosa
120006 | Ornithine racemase
125001 | Ferrous sulfate Fe^59^
126000 | Galactosyl-N-acetylglucosaminylgalactosylglucosylceramide alpha-galactosyltransferase
130002 | Hemoglobin Hopkins-II
131003 | Dolichyl-phosphate mannosyltransferase
159002 | Ferrocyanide salt
164003 | Phosphoenolpyruvate-protein phosphotransferase
178002 | Uridyl diphosphate galactose
186002 | HLA-Cw9 antigen
187006 | Cyanocobalamin Co^57^
200001 | Berberine
217008 | Blood group antigen IH
231008 | 3-Hydroxyisobutyrate dehydrogenase
238002 | Heptachlor
261000 | Codeine phosphate
296000 | Coumachlor
322006 | Octylphenoxy P.H. ethanol
327000 | ^76^Arsenic
329002 | ^127^Antimony
336001 | Fibrinogen Tokyo II
340005 | Enzyme variant
363000 | Fibrinogen San Juan
370000 | beta>2S< Glycoprotein
371001 | Acylcarnitine hydrolase
377002 | Sparteine
392001 | ^151^Gadolinium
395004 | Immunoglobulin pentamer
412004 | Ribose-5-phosphate isomerase
424006 | Citramalyl-CoA lyase
425007 | Hemoglobin Nagoya

432003 | Carminic acid
438004 | 2-Hydroxyglutarate dehydrogenase
462009 | Urease (ATP-hydrolysing)
472007 | Vegetable textile fiber
476005 | CD1b - Cluster of differentiation antigen 1b
498001 | Nitrilase
501001 | Blood group antibody Sf^a^
505005 | Blood group antibody M'
506006 | 3-Oxosteroid delta^1^-dehydrogenase
515004 | Blood group antigen Giaigue
519005 | Free protein S
521000 | ^197^Mercury
529003 | Guanosine
538001 | 2,3-Dihydroxybenzoate 3,4-dioxygenase
566009 | Acrosin
576007 | Blood group antibody Duck
578008 | Hemoglobin Jianghua
584006 | Blood group antibody Wr^b^
585007 | SP - Substance P
591009 | 2-Oxoisovalerate dehydrogenase (acylating)
593007 | Blood group antibody Holmes
594001 | 2-Oxoglutarate synthase
597008 | ^247^Californium
604000 | Plant sapogenin glycoside
611001 | Hippurate hydrolase
620005 | Trichlorophenol
648005 | Oil of calamus
662003 | Aeromonas proteolytica aminopeptidase
668004 | ^185^Osmium
683009 | Mercuric acetate
686001 | Plastoquinol-plastocyanin reductase
693002 | Trichothecene
698006 | Erythromycin lactobionate
699003 | Coal tar extract
704006 | Blood group antigen Rx
732002 | N-valeraldehyde
735000 | Blood group antigen Jobbins
747006 | Oxamniquine
773001 | Hemoglobin M-Iwate
785009 | Dextranase
804003 | Creosotic acid
819002 | Lytic antibody
850000 | Stizolobate synthase
859004 | Peptide-N^4^-(N-acetyl-b-glucosaminyl) asparagine amidase
860009 | Immunoglobulin, aggregated
873008 | Urethan
876000 | D antigen
877009 | Carboxypeptidase A
889006 | [Acetyl-CoA carboxylase] kinase
896008 | Ice
905001 | o-Dihydroxycoumarin O^7^-glucosyltransferase
923009 | Complement component 2
925002 | Sodium iodipamide
963005 | Pyridoxine 4-dehydrogenase
974001 | Adenosylmethionine decarboxylase
979006 | Carbamate kinase
993004 | Palladium compound
1002007 | Mannotetraose 2-alpha-N-acetylglucosaminyltransferase
1010008 | N-Acetylneuraminate monooxygenase
1018001 | Nornicotine
1025008 | ^93^Molybdenum
1047008 | Guanine deaminase
1050006 | Melilotate 3-monooxygenase
1057009 | Phosphate salt
1065007 | E. coli periplasmic proteinase
1080001 | ^202^Thallium
1091008 | Coagulation factor inhibitor
1097007 | Blood group antigen M^A^
1105007 | Isochorismate synthase
1113008 | Pancreatic ribonuclease
1137008 | ^240^Uranium
1149009 | Hemoglobin Barcelona

1160000 | Antibody to antigen in ISBT system LU
1166006 | Ti - Titanium
1169004 | Hemoglobin Gower-2
1171004 | Fibrinogen Kawaguchi
1185009 | Hemoglobin Roseau-Pointe à Pitre
1189003 | Hemoglobin F-M-Osaka
1190007 | Mephenoxalone
1219001 | Diethyl xanthogen disulfide
1223009 | Blood group antigen Marks
1244009 | Fibrinogen Madrid I
1248007 | Leucostoma neutral proteinase
1269009 | Amikacin sulfate
1272002 | Pteridine oxidase
1273007 | Blood group antibody Evelyn
1313002 | Nitrate reductase (cytochrome)
1319003 | Blood group antibody K18
1320009 | Hemoglobin Manitoba
1325004 | Metocurine iodide
1331001 | Methamidophos
1334009 | Estradiol receptor
1336006 | Deoxycortone
1341003 | Hemoglobin Ta-li
1346008 | Blue shade eosin stain
1355006 | Coagulation factor IX Oxford 3 variant
1368003 | Iodine 131
1371006 | Blood group antigen Big
1373009 | ^93^Zirconium
1381005 | ^126^Iodine
1394007 | Iron pentacarbonyl
1396009 | Actinium
1405004 | Blood group antibody M^e^
1408002 | Blood group antibody 1123K
1416006 | Radium compound
1450002 | Methylpentynol
1466000 | Cyclomaltodextrinase
1471007 | Elastin
1472000 | Adenosine-phosphate deaminase
1476002 | Codeine sulfate
1477006 | Hemoglobin Yatsushiro
1496005 | Proto-oncogene
1506001 | CH1 (ISBT symbol)
1517000 | HLA - Human leukocyte antigen B21
1530004 | 6-Carboxyhexanoate-coenzyme A ligase
1535009 | Nitrogen fluoride
1536005 | Pargyline hydrochloride
1540001 | Tellurium radioisotope
1545006 | Uridine phosphorylase
1557002 | Talc dusting powder
1565004 | Blood group antibody Buckalew
1575001 | Maltose tetrapalmitate
1603001 | Cobalt isotope
1607000 | Homoserine kinase
1609002 | N-octyl isosafrole sulfoxide
1634002 | Blood group antigen Ven
1649005 | Blood group antigen Sul
1656004 | Hemoglobin Shaare Zedek
1660001 | Plant seeds
1668008 | Ceforanide
1672007 | Ligase
1673002 | Xylenol
1675009 | ^86^Rubidium
1676005 | Blood group antibody LW^ab^
1681001 | Blood group antibody BLe^b^
1696002 | 12-Hydroperoxy eicosatetraenoic acid
1701009 | ^191^Gold
1710001 | UA - Uric acid
1726000 | Diamond
1727009 | Deoxylimonate A-ring-lactonase
1740004 | Deoxy cytidine triphosphate
1764003 | Saccharopine dehydrogenase (nicotinamide adenine dinucleotide phosphate ^+^,L-glutamate-forming)
1768000 | Sucrose phosphorylase
1786002 | Leucine-transfer ribonucleic acid ligase

1793003 | Sodium trichloroacetate
1795005 | Glyodin
1798007 | Hemoglobin Hammersmith
1799004 | L-Lysine oxidase
1823002 | Hemoglobin Tochigi
1827001 | Ribonuclease T>1<
1886008 | Verdoglobin
1904005 | Galactoside 3-fucosyltransferase
1914001 | von Willebrand factor antibody
1916004 | Boroglycerin
1940007 | Immunoglobulin, GM>21< allotype
1944003 | Coagulation factor X Patient variant
1956002 | Buclizine hydrochloride
1971003 | Loxapine hydrochloride
1975007 | Blood group antibody Niemetz
1978009 | Site-specific methyltransferase (cytosine-specific)
1985008 | Vomitus
1991005 | Lignin
2000001 | Heavy nitrogen
2006007 | Inosine diphosphate
2008008 | ^67^Gallium
2009000 | Cobalt carbonyl
2017008 | Deoxyribonucleic acid topoisomerase
2027002 | Alternaria serine proteinase
2029004 | Fibrinogen Oslo II
2038002 | Blood group antibody Bg^b^
2039005 | sym-Norspermidine synthase
2050008 | Choloylglycine hydrolase
2064008 | L-Xylulokinase
2082006 | Vitronectin receptor alpha chain
2085008 | Oncogene protein TCL
2088005 | Page blue G-90 stain
2096000 | Nicotinamide adenine dinucleotide ^+^ adenosine diphosphate-ribosyltransferase
2100004 | Sulfonethylmethane
2101000 | Yeast proteinase B
2125008 | Betazole
2130007 | Cyclohexane-1,2-diol dehydrogenase
2141009 | Hydrogen
2147008 | Blood group antigen Paular
2151005 | Pyridoxamine-pyruvate aminotransferase
2154002 | Tagaturonate reductase
2159007 | Acid red 27
2163000 | Dicofol
2168009 | Bisphosphoglycerate mutase
2179004 | Malonate-semialdehyde dehydratase
2189000 | Hemoglobin F-Dammam
2194000 | ^101^Rhodium
2195004 | Tocainide hydrochloride
2197007 | Boric acid topical agent
2201007 | Bacteriopurpurin
2208001 | Phenylserine aldolase
2212007 | Fibrinogen Bethesda II
2215009 | Azuresin
2240002 | Guanidinobutyrase
2249001 | Gentamicin sulfate
2254005 | Orotate
2260005 | Human leukocyte antigen DRw18
2262002 | Cellulose polysulfatase
2264001 | Selenium isotope
2309006 | Au - Gold
2311002 | Prostacyclin synthase
2329007 | Blood group antibody Vel
2331003 | Saccharide
2338009 | Root
2343002 | Guthion
2346005 | Vascormone
2354007 | 3'-Nucleotidase
2358005 | Glass fragment, device
2369008 | Indole-3-acetate beta-glucosyltransferase
2370009 | Uridine diphosphate-N-acetylmuramate-alanine ligase
2376003 | Mercury compound
2384004 | ^230^Uranium

2404002 | Blood group antibody St^a^
2405001 | b- Propiolactone
2414006 | Prolactin receptor
2430003 | Silicon radioisotope
2431004 | Blood group antibody Friedberg
2441001 | Mercury radioisotope
2444009 | Human leukocyte antigen Dw25
2450004 | Mannosamine
2462000 | Glucose dehydrogenase (nicotinamide adenine dinucleotide phosphate ^+^)
2466002 | Chloride peroxidase
2500009 | Lymphocyte antigen CDw41b
2509005 | D-Glutamate oxidase
2516006 | Metallic sulfide compound
2522002 | Extravascular blood
2529006 | Hemoglobin Wood
2537003 | Antituberculosis agent
2568004 | Blood group antigen McAuley
2573005 | Immunoglobulin, GM>13< allotype
2575003 | Zinc alpha-2 glycoprotein
2595009 | ^119m^Tellurium
2597001 | Alpha-1 globulin
2611008 | Blood group antibody La Fave
2637006 | Indium isotope
2648004 | Bile vomitus
2649007 | Azo-dye
2660003 | Sodium dehydrocholate
2671002 | 3-Methyl-2-oxobutanoate hydroxy-methyltransferase
2674005 | ^128^Cesium
2676007 | C3(H20)
2678008 | Hemoglobin New Mexico
2680002 | Factor XIII antibody
2698003 | Natural gas
2705002 | ^72^Arsenic
2706001 | Blood group antigen Vennera
2719002 | Tartrate dehydratase
2721007 | Blood group antigen McC^f^
2728001 | Lewis system antigen
2753003 | Blood group antibody M>1<
2754009 | Hemoglobin F-Kennestone
2765004 | SC3 (ISBT symbol)
2778004 | Pleural fluid
2796008 | Methanthelinium
2799001 | Methylbenzethonium
2823004 | Hemoglobin Bristol
2832002 | Molybdenum compound
2846002 | Hemoglobin Saitama
2869004 | Ethanoic acid
2878005 | Pethidine hydrochloride
2880004 | Calcium sulfate
2883002 | Exopolygalacturonate lyase
2913009 | Immunoglobulin Epsilon chain
2916001 | ^22^Neon
2925007 | Fluoromethalone
2927004 | Rescinnamine
2938004 | Pyrazole
2942001 | Carbon^14^ D-xylose
2950005 | Hemoglobin L-Persian Gulf
2958003 | Zinc caprylate
2964005 | Dimethoxyamphetamine
2974008 | Trichophyton schoenleinii collagenase
2988007 | Human leukocyte antigen Aw
2991007 | Mecamylamine hydrochloride
2995003 | Arecoline
3027009 | ^133^Barium
3031003 | Dihydroxyaluminum sodium carbonate
3040004 | Technetium Tc^99m^ disofenin
3045009 | Nitrochlorobenzene
3052006 | Ornithine ketoacid aminotransferase
3066001 | Triiodothyroacetic acid
3070009 | Aspartate-ammonia ligase
3087006 | Oil of male fern
3107005 | Hemoglobin Shuangfeng

3108000 | Aspergillus deoxyribonuclease K>1<
3131000 | Blood group antigen Middel
3136005 | Cefoperazone sodium
3142009 | Azacyclonol
3145006 | Penicillic acid
3150000 | Sialate O-acetylesterase
3151001 | Left upper lobe mucus
3155005 | 3-Phosphoglyceroyl-phosphate-polyphosphate phosphotransferase
3161008 | 3-Methylhistidine
3167007 | Hard coal
3187008 | Blood group antigen Nielsen
3193000 | alpha-1,4-Glucan-protein synthase (uridine diphosphate-forming)
3197004 | Inosine monophosphate
3209002 | Pancuronium sodium
3212004 | Manganese sulfate
3225007 | Fibrinogen Seattle I
3232003 | o-Benzyl-parachlorophenol
3271000 | Hemoglobin Southampton
3273002 | Tyrosine-ester sulfotransferase
3300001 | Euphorbain
3318003 | Vaginal secretions
3325005 | Lipopolysaccharide
3339005 | (R)-20-Hydroxysteroid dehydrogenase
3340007 | Diastase
3342004 | Copper isotope
3346001 | Hemoglobin Brest
3378009 | Imipramine hydrochloride
3379001 | Merthiolate
3392003 | Aldehyde dehydrogenase (acceptor)
3405005 | 2-Hydroxy-3-oxoadipate synthase
3411008 | bis-(Dimethylthiocarbamyl) disulfide
3437006 | Hydroxymethylglutaryl-coenzyme A hydrolase
3440006 | Biotin carboxylase
3455002 | Discontinued pesticide
3463001 | L-Amino-acid dehydrogenase
3465008 | Deoxyribonucleic acid topoisomerase (adenosine triphosphate [ATP]-hydrolysing)
3466009 | Dimethylamine
3492002 | Galactinol-sucrose galactosyltransferase
3493007 | Smegma clitoridis
3495000 | Cystine aminopeptidase
3501003 | Isoxsuprine hydrochloride
3523004 | Hemoglobin Q-India
3532002 | Laryngeal mucus
3555004 | Blood group antigen Morrison
3579002 | ^129^Cesium
3581000 | Glucose-6-phosphatase
3587001 | Malate dehydrogenase (decarboxylating)
3588006 | Complement enzyme
3592004 | Short-acting thyroid stimulator
3597005 | Acebutolol hydrochloride
3601005 | Ether solvent
3602003 | Warm antibody
3610002 | Epoxide hydrolase
3617004 | ^79^Selenium
3648007 | Glucocorticoid receptor
3655009 | Hb - Hemoglobin Constant Spring
3672002 | Fibrinogen Caracas
3684000 | Phenylacetic acid
3689005 | Hemoglobin Mizushi
3692009 | Sodium sulfite
3693004 | Fibrinogen Dusart
3702007 | Cytidine diphosphate (CDP) glycerol glycerophosphotransferase
3710008 | Prostaglandin-endoperoxide synthase
3718001 | Cow's milk
3726009 | Valine-transfer ribonucleic acid ligase
3727000 | Hemoglobin F-Port Royal
3730007 | Blood group antigen Lanthois
3737005 | Nitrate reductase (reduced nicotinamide adenine dinucleotide)
3742002 | Extracellular crystal
3757009 | Gossypol
3771001 | Neuromelanin
3775005 | Choline dehydrogenase

3776006 | Xanthine dehydrogenase
3792001 | AA - Arachidonic acid
3793006 | Soluble barium compound
3800009 | Acetate kinase
3807007 | RH4 (ISBT symbol)
3811001 | Magnesium-protoporphyrin methyltransferase
3812008 | Beryllium isotope
3816006 | Vanadium isotope
3823007 | Prochlorperazine edisylate
3829006 | Fe - Iron
3834005 | CMP-N-acetylneuraminate-(alpha-N-acetyl-neuraminyl-2,3-beta-galactosyl-1,3)-N-acetyl-galactosaminide alpha-2,6-sialyltransferase
3836007 | Glutaminase
3844007 | Protoaphin-aglucone dehydratase (cyclizing)
3848005 | Nitrotoluene
3849002 | Carbon black
3854006 | bis-Chloro methyl ether
3874004 | Hydrocodone bitartrate
3892007 | Thymidine
3896005 | p-Hydroxybenzoate ester
3897001 | MNS30 (ISBT symbol)
3906002 | Rectified birch tar oil
3920009 | Hemoglobin Atago
3930000 | Manufactured gas
3932008 | ^64^Copper
3941003 | Metronidazole hydrochloride
3945007 | Tin isotope
3958008 | ^245^Californium
3961009 | Blood group antigen Ritherford
3976001 | Blood group antigen HEMPAS
3982003 | Oxaloacetate decarboxylase
3983008 | N,-N-dimethyltryptamine
3990003 | Alkaline phosphatase bone isoenzyme
3994007 | Hemoglobin Tampa
4014000 | Sulfisomidine
4024008 | Soft metal
4025009 | Captodiame
4043008 | Etidocaine hydrochloride
4047009 | cis-1,2-Dihydrobenzene-1,2-diol dehydrogenase
4048004 | 1,1,2,2-Tetrachloro-1,2- difluoroethane
4067000 | Chorismate mutase
4076007 | PTH - Parathyroid hormone
4077003 | Dihydrolipoamide succinyltransferase
4080002 | Hemoglobin Grady, Dakar
4091009 | Enteropeptidase
4097008 | Apo-SAA complex
4104007 | Chondroitin sulfate
4105008 | Adenylate cyclase
4115002 | Blood group antibody Norlander
4137009 | sec-Butyl acetate
4153007 | Long-chain-enoyl-coenzyme A hydratase
4167003 | PECAM-1 - Platelet endothelial cell adhesion molecule-1
4169000 | Blood group antibody Le^bH^
4177001 | Hemoglobin Long Island-Marseille
4182008 | Cytidine diphosphate (CDP) diacylglycerol-serine O-phosphatidyl-transferase
4188007 | Fibrinogen Sydney II
4200007 | Neriifolin
4201006 | 6-Aminohexanoate-dimer hydrolase
4203009 | Imipramine pamoate
4207005 | Cortisone beta-reductase
4217000 | Fluorosilicate salt
4218005 | Immunoglobulin, GM>23< allotype
4231000 | Gallium isotope
4239003 | Glycerol dehydrogenase
4255005 | Americium 241
4289006 | Keyhole-limpet hemocyanin
4290002 | Linamarin synthase
4314009 | Blood group antibody Allchurch
4334005 | Tar oil
4342006 | 2-Aminopyridine
4353000 | Di-n-butylphthalate
4355007 | Coagulation factor IX San Dimas variant

4362003 | 4-Coumarate-coenzyme A ligase
4370008 | Acetone
4393002 | Blood group antigen Fedor
4401009 | Blood group antibody H>T<
4413004 | Benzypyrinium
4422003 | Blood group antigen
4423008 | Fibrinogen New York II
4425001 | Blood group antibody Binge
4435007 | Sulfuryl fluoride
4437004 | ^127^Cesium
4471008 | ^244^Californium
4479005 | Hemoglobin Brockton
4480008 | Sulfaethidole
4509009 | Plant phenanthrene toxin
4518006 | Buthenal
4534009 | ^208^Bismuth
4540002 | Adenosine diphosphate (ADP) deaminase
4546008 | Tetradecanoic acid
4555006 | Blood group antibody Rils
4560005 | Hemoglobin Mizuho
4561009 | Arginine decarboxylase
4564001 | Blood group antibody Sisson
4567008 | Galactose-1-phosphate thymidylyltransferase
4582003 | Blood group antigen N^A^
4591004 | MNS22 (ISBT symbol)
4610008 | Senile cardiac protein
4616002 | Triclobisonium chloride
4629002 | Hypoglycin B
4635002 | Arterial blood
4643007 | Calf thymus ribonuclease H
4656000 | Alcian blue 8GX stain
4674009 | 2,3-Dihydroxybenzoate serine ligase
4681002 | Potassium permanganate
4693006 | Chromium^51^ albumin
4700006 | Bovine insulin
4706000 | Chlorine monoxide
4714006 | ^183m^Osmium
4728000 | Scopulariopsis proteinase
4731004 | Aluminum pyro powder
4732006 | Oncogene protein P55, V-MYC
4746006 | Hemoglobin Mito
4761007 | CD30 - Cluster of differentiation antigen 30
4762000 | Platelet antigen HPA-3b
4777008 | Fluroxene
4780009 | Secbutabarbital sodium
4786003 | beta-1,4-Mannosyl-glycoprotein beta-1,4-N-acetylglucosaminyltransferase
4789005 | Blood group antibody Bultar
4793004 | Azobenzene reductase
4814001 | Valethamate
4824009 | MAO - Monoamine oxidase
4825005 | Peptidyl-glycinamidase
4831008 | Arabinose-5-phosphate isomerase
4832001 | Technetium Tc^99m^ mebrofenin
4833006 | Glucan endo-1,3-alpha-glucosidase
4844003 | 3,3' Diiodothyronine
4864008 | AMP - Adenosine monophosphate
4872005 | Glucosulfone
4878009 | Human leukocyte antigen Dw3
4882006 | Ichthyoallyeinotoxin
4889002 | Xylulokinase
4901003 | Pyruvate oxidase (coenzyme A-acetylating)
4925006 | Oncogene protein V-ABC
4933007 | Leukocyte M1
4940008 | Tattoo dye
4955004 | Neoplastic structural gene
4962008 | Tree bark
4963003 | Neutral amino acid
4965005 | Glutathione reductase
4968007 | Acumentin
4986005 | Magnesium borate
5003005 | Hemoglobin Swan River
5004004 | Blood group antibody Panzar

5007006 | Papain
5024000 | Fresh water
5031001 | 3-3'Dichlorobenzidine
5040002 | Cesium
5043000 | Erythrosin Y stain
5045007 | Oncogene protein TCL4
5059000 | ^97^Technetium
5060005 | ^132^Cesium
5061009 | Protein-methionine-S-oxide reductase
5064001 | Blood group antibody D 1276
5081005 | RH31 (ISBT symbol)
5086000 | Gelsolin
5094007 | Blood group antigen Rios
5098005 | Fennel oil
5109006 | Methylated-deoxyribonucleic acid-protein-cysteine methyltransferase
5142007 | Coagulation factor II Houston variant
5160007 | Metal compound
5163009 | Scombrotoxin
5167005 | Zinc chloride fumes
5172001 | Coagulation factor Xa
5179005 | Connective tissue fiber
5200001 | trans-Epoxysuccinate hydrolase
5206007 | Cyanate compound
5220000 | Bacitracin
5226006 | Flavone O^7^-beta-glucosyltransferase
5250008 | Thymus-independent antigen
5252000 | Hafnium radioisotope
5253005 | Hemoglobin Woodville
5259009 | Blood group antigen Braden
5289002 | Scilliroside
5303002 | Hemoglobin Hoshida
5305009 | Polynucleotide
5307001 | Blood group antigen Hamet
5312000 | ^65^Zinc
5323001 | Uridine diphosphate glucuronic acid
5330007 | Actin-binding protein
5339008 | L-Glycol dehydrogenase
5340005 | Blood group antigen Swietlik
5392001 | Propylene glycol monomethyl ether
5395004 | Pyridoxamine-phosphate oxidase
5404007 | Lymphocyte antigen CD45RA
5405008 | ^60^Cobalt
5406009 | beta-L-Arabinosidase
5420002 | Accessory sinus mucus
5439007 | Blood group antibody Do^a^
5442001 | Page blue 83
5453007 | Iridium isotope
5471000 | Hemoglobin G-Coushatta
5474008 | Propionate-CoA ligase
5477001 | Ferric subsulfate
5483003 | Oxalate CoA-transferase
5504009 | Blood group antigen Fuerhart
5511008 | Inosinate nucleosidase
5513006 | Immunoglobulin, Alpha chain
5515004 | Rhodium fumes
5533005 | Blood group antibody Kp^a^
5537006 | Immunoglobulin, Delta chain
5540006 | Calcium
5547009 | ^233^Plutonium
5548004 | 2-Dehydro-3-deoxy-D-pentonate aldolase
5568005 | Hemoglobin Hijiyama
5573004 | Blood group antigen Oca
5589001 | Licodione O^2'^-methyltransferase
5590005 | Beryllium radioisotope
5628003 | Hemoglobin I-High Wycombe
5629006 | Cytidylic acid
5637003 | HLA-DQw6 antigen
5641004 | Valproate semisodium
5647000 | Griseofulvin ultramicrosize
5656008 | ^116m^Antimony
5657004 | Coal tar topical solution
5659001 | Hemoglobin J-Tongariki

5670008 | Gold isotope
5681006 | Ceftizoxime sodium
5691000 | Absorbable gelatin sponge
5692007 | Cyanocobalamin Co^58^
5699003 | Somatomedin C
5700002 | Blood group antibody Gomez
5702005 | ^106m^Silver
5704006 | Galactokinase
5705007 | 1,3-Propanediol dehydrogenase
5739006 | Stramonium
5746002 | ^118m^Antimony
5757007 | HLA-Cw8 antigen
5762008 | Heterogeneous nuclear RNA
5764009 | ^242^Plutonium
5767002 | Sulfamerazine
5774007 | White petrolatum
5800007 | tRNA (5-methylaminomethyl-2-thiouridylate)-methyltransferase
5813001 | Malate dehydrogenase
5826002 | Ethyl-4-bis-(hydroxypropyl)-1-aminobenzoate
5827006 | Crotonaldehyde
5829009 | Hemoglobin Vaasa
5830004 | Hemoglobin Bart
5840001 | Blood group antibody Wj
5858007 | ^110m^Indium
5863006 | Vitexin beta-glucosyltransferase
5896008 | Hellebrin
5899001 | Bacterial structural gene
5907009 | Quinidine polygalacturonate
5910002 | Oncogene protein PP60, V-SRC
5915007 | Blood group antigen Gladding
5927005 | Lactaldehyde dehydrogenase
5931004 | Technetium Tc^99m^ sulfur colloid
5932006 | Cysteine
5950004 | 3',5'-Cyclic-nucleotide phosphodiesterase
5955009 | Diethylene glycol
5977008 | Blood group antigen Bullock
5989005 | Immunoglobulin, GM>17< allotype
5991002 | D-Fuconate dehydratase
6021003 | ^88^Yttrium
6038004 | Oxygen radioisotope
6043006 | Bone cement
6044000 | Carbon disulfide
6054001 | Doxylamine succinate
6056004 | Blood group antibody Wk^a^
6068008 | Blood group antigen Mil
6083003 | Hydroxylysine
6085005 | Synovial fluid
6088007 | Benzfetamine hydrochloride
6089004 | Lochia alba
6091007 | Blood group antibody L Harris
6107003 | Asparagusate reductase (NADH)
6109000 | Aromatic-amino-acid aminotransferase
6115000 | Blood group antibody Anuszewska
6135004 | Blood group antigen Duck
6138002 | Blood group antigen Le Provost
6162007 | Meclocycline
6170002 | Heat labile antibody
6172005 | Fatty-acid methyltransferase
6178009 | Lymphocyte antigen CD63
6179001 | o-Methy-bufotenine
6182006 | Chloroacetone
6197009 | Blood group antigen Zd
6237004 | Bemegride
6249003 | Potassium metabisulfite
6256009 | Ribose isomerase
6257000 | Sodium chloride Na^22^
6260007 | Protokylol
6261006 | Indoklon
6263009 | Plant residue
6264003 | Diazinon
6287006 | Methidathion
6291001 | N-Acetylglucosamine-1-phosphodiester N-acetylglucosaminidase

6301006 | ^178^Tantalum
6310003 | Particulate antigen
6314007 | Phenol beta-glucosyltransferase
6333002 | Squill extract
6338006 | Imidazolonepropionase
6356006 | Chlorodiallylacetamide
6360009 | Kallidin II
6367007 | ^95m^Technetium
6386004 | N-Acetylneuraminate O^4^-acetyltransferase
6394006 | Phentermine hydrochloride
6401007 | Lichenase
6409009 | Morpholine
6411000 | Interleukin-12
6422001 | HLA-DRw14 antigen
6451002 | Chlorobenzilate
6455006 | Chloroprene
6469006 | delta^1^-Piperideine-2-carboxylate reductase
6478000 | 6-Phosphofructokinase
6495008 | Fibrinogen Montreal II
6507009 | Blood group antigen Much
6513000 | Flumethiazide
6516008 | Indium^111^-Fe(OH)>3<
6524003 | Distilled spirits
6529008 | Blood group antigen Cl^a^
6532006 | Macrophage activating factor
6590001 | Galactosylceramidase
6592009 | HLA-Dw12 antigen
6602005 | Aminoacridine
6611005 | Diethylaminoethanol
6612003 | Chloramphenicol sodium succinate
6619007 | Bilirubin Y transport protein
6642000 | Opsonin
6644004 | Homoserine dehydrogenase
6671004 | Blood group antigen Caw
6672006 | Phosphoadenylate 3'-nucleotidase
6699008 | Titanium radioisotope
6701008 | Lissamine fast red B
6702001 | Ethyl mercaptoethyl diethyl thiophosphate
6709005 | Gentamicin 2''-nucleotidyltransferase
6710000 | Nitric oxide
6713003 | ^91^Yttrium
6717002 | Nifuroxime
6725000 | Methylene blue
6730001 | ^234^Uranium
6741004 | Anti DNA antibody
6755007 | TL antigen
6786001 | Silver difluoride
6790004 | Aminopterin
6792007 | Veratrine
6808006 | Ferrous iron compound
6809003 | Phomopsin
6814004 | Discadenine synthase
6817006 | Oxidized glutathione
6826009 | Sterol hormone
6837005 | Dextropropoxyphene napsylate
6854002 | ^188^Platinum
6865007 | Theophylline calcium salicylate
6873003 | Cefapirin sodium
6879004 | 5,8,11-Eicosatrienoic acid
6881002 | Magnesium fumes
6884005 | (S)-3-Amino-2-methylpropionate aminotransferase
6890009 | 3-Deoxy-manno-octulosonate-8-phosphatase
6896003 | Thiopurine methyltransferase
6910009 | Sodium fluoride
6911008 | Deoxycytidylate methyltransferase
6916003 | Bowieine
6924008 | Exopolyphosphatase
6925009 | Leucine acetyltransferase
6927001 | ^121^Tin
6937006 | Thymidylate synthase
6945001 | Blood group antigen Le^bH^
6952004 | ^121m^Tin

6958000 | Blood group antibody Frando
6961004 | Lysolecithin acylmutase
6970001 | 4-Hydroxyproline epimerase
6973004 | Chromium^51^ chloride
6983000 | Acrylamide
6992002 | Triflupromazine hydrochloride
6993007 | Seminal fluid
6999006 | Ammonium compound
7008002 | beta-Carotene 15,15'-dioxygenase
7018007 | Malate-CoA ligase
7029006 | Blood group antigen Greenlee
7030001 | Globoside
7034005 | Diclofenac
7045008 | Lycorine
7047000 | Asphyxiant atmosphere
7049002 | Pyruvate carboxylase
7054006 | Hemoglobin Poissy
7056008 | 3-Propylmalate synthase
7059001 | N-Acylneuraminate-9-phosphatase
7061005 | Anthocyanidin O^3^-glucosyltransferase
7070008 | Convallamarin
7084003 | Fibrinogen Buenos Aires II
7110002 | ^69^Germanium
7120007 | Antigen
7132006 | ^73^Gallium
7139002 | Acid-CoA ligase (GDP-forming)
7146006 | Cyclohexene oxide
7152007 | Chlorthion
7156005 | Phosphorus isotope
7158006 | HLA-Dw19 antigen
7161007 | Complement component C2a
7179006 | Prekallikrein
7191002 | Methenyltetrahydrofolate cyclohydrolase
7208009 | Thiol oxidase
7211005 | Blood group antibody Haakestad
7237008 | Galactonate dehydratase
7243005 | Methyl isocyanate
7269004 | Th - Thorium
7271004 | Mixed dust
7280004 | dTDP4-dehydrorhamnose reductase
7281000 | Technetium Tc^99m^ lidofenin
7284008 | Mercaptan compound
7294003 | tert-Butyl acetate
7302008 | Ambuphylline
7318002 | Bacteriochlorophyll
7321000 | Pyrimidine
7325009 | Calcium hydroxide
7327001 | Sulfurous acid
7328006 | Red petrolatum
7330008 | Shellac
7337006 | Blood group antibody Tr^a^
7348004 | Factor II
7382005 | Aminoalcohol ester
7401000 | Heme-hemopexin complex
7411007 | Blood group antibody HLA-B8
7427000 | Sepiapterin reductase
7434003 | Erythrosin B
7446004 | Ruthenium
7451005 | Tobramycin ophthalmic agent
7460002 | ^127^Tellurium
7470000 | p-tert-Butyltoluene
7489000 | Homocytotropic antibody
7503004 | ^72^Gallium
7509000 | Mannitol hexanitrate
7515000 | Hepatotoxic mycotoxin
7537007 | Stizolobinate synthase
7547005 | Hemoglobin Lincoln Park
7549008 | Fibrinogen Bethesda I
7588005 | Blood group antibody Sk^a^
7608003 | Triethylene glycol
7616007 | Blood group antibody Pruitt
7648006 | HLA-Bw70 antigen

7661006 | Fish bone
7670009 | Aminobutyraldehyde dehydrogenase
7675004 | Blood group antigen Towey
7679005 | Strong oxidizing compound
7685003 | Blood group antibody Bg^c^
7696006 | Ferrovanadium dust
7716001 | Isovaleryl coenzyme A dehydrogenase
7737009 | Chlortetracycline hydrochloride
7738004 | HLA-B49 antigen
7761002 | ^111^Silver
7770004 | ^89^Strontium
7774008 | Neo-b-vitamin A>1<
7779003 | ^103^Ruthenium
7785005 | Sphingomyelin phosphodiesterase D
7790008 | 1-Monoacylglycerol
7791007 | Soy protein
7795003 | Oxalate oxidase
7801007 | Tetrahydroxypteridine cycloisomerase
7816005 | Antazoline hydrochloride
7834009 | Acetyl digitoxin
7846008 | Sphingomyelin phosphodiesterase
7848009 | 1-Phosphatidylinositol phosphodiesterase
7868003 | beta-Cyclopiazonate dehydrogenase
7879008 | ^218^Radon
7900007 | Hemoglobin Presbyterian
7904003 | Deanol
7909008 | Arginine carboxypeptidase
7924004 | Diflorasone
7938006 | D-Arabinitol dehydrogenase
7945006 | Orsellinate-depside hydrolase
7948008 | Reed-Sternberg antibody
7953003 | Thioneb
7957002 | Phosphatidate cytidylyltransferase
7961008 | Hemoglobin F-Shanghai
7970006 | Allograft
7974002 | Blood group antibody Dalman
7975001 | Amiphenazole
7979007 | 3'-Phosphoadenylylsulfate 3'-phosphatase
7983007 | Sodium rhodanide
7985000 | Sulfur isotope
7997004 | Butyl mercaptan
8000007 | Cucurbitacin delta^23^-reductase
8002004 | Blood group antibody Fleming
8025003 | Blood group antibody Gibson
8029009 | Allyl glycidyl ether
8030004 | PEG - Polyethylene glycol
8035009 | Cholestenol delta-isomerase
8048008 | Blood group antigen Th
8054009 | Orotate reductase (NADPH)
8055005 | Galactoside acetyltransferase
8105004 | Hemoglobin Leiden
8108002 | Undecaprenyl-diphosphatase
8123007 | Blood group antibody Schuppenhauer
8132009 | Magnesium acetylsalicylate
8143001 | Diosmin
8153000 | Homoproline
8156008 | Immunoglobulin, Fd fragment
8164002 | Cluster of differentiation antigen 67
8168004 | Uracil-5-carboxylate decarboxylase
8179009 | Cevadilline
8184003 | Convallamarogenin
8190004 | Diaminopimelate epimerase
8202008 | ^43^Potassium
8203003 | Human menopausal gonadotropin
8204009 | Polyester
8222007 | Coagulation factor II Padua variant
8227001 | ^106^Ruthenium
8230008 | Streptococcal cysteine proteinase
8237006 | Strobane
8252004 | Chlorothiazide sodium
8257005 | Abnormal hemoglobin
8261004 | Potassium thiosulfate

8268005 | Blood group antibody Hildebrandt
8270001 | tRNA adenylyltransferase
8275006 | Methionine-S-oxide reductase
8295000 | Uromucoid protein
8300003 | Cyclohexanol
8310007 | Hemoglobin Madrid
8313009 | RNA-directed DNA polymerase
8340009 | Procollagen-lysine,2-oxoglutarate 5-dioxygenase
8342001 | Brilliant cresyl blue
8343006 | Blood group antibody Re^a^
8354001 | Manganese ethylene bis-dithiocarbamate
8355000 | Hafnium isotope
8362009 | Anti c
8365006 | Oil of pennyroyal-European
8368008 | Xylan 1,44-beta-xylosidase
8376005 | Duffy blood group antibody
8385005 | Acid alpha-glucosidase
8397006 | Nicotine resin complex
8406008 | Nitroethane oxidase
8429000 | Brilliant orange
8450009 | Lemon grass oil
8452001 | Blood group antigen Sisson
8456003 | Methyl ethyl ketone peroxide
8460000 | Blood group antibody Vg^a^
8473001 | Homocysteine methyltransferase
8474007 | Lead oleate
8484008 | Blood group antigen Mur
8485009 | Oncogene protein P210, BCR-ABL
8486005 | HLA-DRw15 antigen
8487001 | ^48^Vanadium
8491006 | Complement inhibitor
8492004 | Allantoicase
8498000 | Short neurotoxin venom
8507001 | Cyclohexane
8514004 | Ornithine
8520003 | Hemoglobin Machida
8525008 | ^183^Osmium
8529002 | Urinary protein of low molecular weight
8534003 | ^110^Tin
8537005 | Solution
8578007 | Potassium cyanate
8591008 | Dichlorodifluoromethane
8612007 | Tumor necrosis factor
8620009 | Oncogene protein TCL6
8631001 | Potassium chloride
8653004 | Rubijervine
8660005 | Complement component C3c
8687009 | Gum arabic
8689007 | Kanamycin sulfate
8701002 | Sulfachlorpyridazine
8705006 | 4-Hydroxybenzoate decarboxylase
8731008 | Blood group antibody Austin
8740007 | C3(H20)Bb
8761000 | Adenylylsulfate kinase
8767001 | Santonin
8785008 | Chlorine dioxide
8786009 | Blood group antigen Wd^a^
8795001 | Hb F - Hemoglobin F
8817004 | LH receptor site
8818009 | Blood group antibody Tri W
8822004 | Linoleic acid
8830003 | Nitrate reductase (NAD(P)H)
8836009 | Gallocyanine
8844009 | Hydroxybutyrate-dimer hydrolase
8858006 | Strontium nitrate Sr^85^
8865003 | Graphite
8878003 | Blood group antigen Evelyn
8882001 | 3-Hydroxybenzoate 6-monooxygenase
8886003 | Flecainide acetate
8908003 | Blood group antibody I^T^
8914005 | Endolymph
8919000 | Biotin

8926000 | Azur B
8945009 | Phosphopantothenate-cysteine ligase
8953001 | 2,3-Dihydroxyindole 2,3-dioxygenase
8963009 | N-Acetylmuramoyl-L-alanine amidase
8969008 | Bulbourethral secretions
8977007 | Blood group antibody Tarplee
8982000 | Oleate hydratase
8987006 | Cycle-phase specific agent
8991001 | Ribulokinase
9010006 | Methyl blue
9013008 | Dephospho-CoA kinase
9021002 | Carbaryl
9024005 | G6PD - Glucose-6-phosphate dehydrogenase
9045003 | Radon radioisotope
9052001 | Allspice oil
9054000 | Human leukocyte antigen B15
9103003 | Retinol fatty-acyltransferase
9110009 | Mercuric compound
9125009 | Sempervirine
9159008 | Triacetate-lactonase
9172009 | Blood group antibody Alda
9174005 | Fibrinogen Poitiers
9183000 | beta-N-Acetylgalactosaminidase
9189001 | CMP-N-acetylneuraminate-lactosylceramide alpha-2,3-sialyltransferase
9195000 | Immunoglobulin gene INV allotype
9197008 | Apiose reductase
9205004 | Hemoglobin Tarrant
9220005 | Plant phenol oil
9223007 | Borneol dehydrogenase
9234005 | Chlorobutanol
9246009 | ^118^Tellurium
9253000 | HLA-DRw16 antigen
9270008 | Catecholamine receptor
9271007 | Fibrinogen Pontoise
169008 | Hypothalamic releasing factor preparation
211009 | Norethandrolone preparation
302007 | Spiramycin
439007 | Therapeutic radiopharmaceuticals
449005 | Procaine penicillin
544002 | Melphalan
669007 | Vaccinia virus vaccine
716000 | Hematological agents
796001 | Digoxin product
847003 | D-thyroxine preparation
902003 | Experimental drug
922004 | Pralidoxime
1039008 | 6MP - Mercaptopurine
1148001 | Ticarcillin
1182007 | Hypotensive agent
1206000 | Alpha 2 adrenergic blocking agent
1222004 | Metronidazole
1389007 | Beclometasone
1434005 | Calamine
1528001 | Folinic acid product
1552008 | Saluretic drug
1594006 | Azatadine
1613009 | Dexbrompheniramine maleate
1756009 | Antipruritic topical anesthetic
1758005 | Motilin preparation
1842003 | Diphemanil
1878008 | Hexachlorophane product
1887004 | Permethrin
1982006 | Bacitracin ophthalmic preparation
2016004 | Dextromethorphan
2037007 | Otic anti-inflammatory preparation
2183004 | Tetryzoline
2190009 | Benzhexol
2497003 | Hexetidine
2571007 | Busulfan
2596005 | Lincomycin
2679000 | Oxandrolone
2944000 | Depilatory agent

2949005 | Diagnostic agents
3037004 | Flumethasone preparation
3046005 | Enalapril maleate
3127006 | 5FU - Fluorouracil
3221003 | Ringer solution
3334000 | Cefotaxime
3361000 | Anti-heparin agent
3526007 | Rabies vaccine, human
3593009 | Otic anti-infective preparation
3814009 | Propylthiouracil product
3822002 | Suxamethonium
4126008 | Fluprednisolone preparation
4194004 | Mazindol
4219002 | Penicillamine
4220008 | Tolazoline
4382004 | Centrally acting antihypertensive agent
4704002 | Iothiouracil
4741001 | Sodium folate preparation
4753002 | Triamcinolone dental paste
4865009 | Prolactin releasing factor preparation
4937008 | Cefaclor
5067008 | Antithyroid drugs
5465006 | Propitocaine hydrochloride
5478006 | Trifluperidol
5487002 | Therapeutic agent
5606003 | Dexamethasone nasal preparation
5720001 | Latrodectus mactans antivenin
5737008 | Demeclocycline
5776009 | Anesthetic
5786005 | Chlorothiazide
5797005 | Clotrimazole
5924003 | Isosorbide dinitrate
5975000 | Niclosamide
6028009 | Triamcinolone preparation
6067003 | Orciprenaline
6071000 | Coal tar preparation
6102009 | Baclofen
6116004 | Oxymetholone preparation
6122008 | Class Ia antiarrhythmic drug
6232005 | Naphazoline
6247001 | Folic acid preparation
6259002 | Hydrogen peroxide solution 3%
6369005 | Penicillin -class of antibiotic-
6425004 | Antihistamine
6517004 | Butyrophenone derivative antipsychotic agent
6526001 | Nalorphine
6625006 | Zinc sulfate
6652001 | Calcium pantothenate preparation
6716006 | Abortient agent
6960003 | Polymyxin B
6985007 | Opium preparation
7092007 | Metoprolol
7140000 | Radiographic contrast media
7168001 | Magnesium carbonate
7230005 | Brucella vaccine
7235000 | Ethylenediamine derivative antihistamine
7292004 | Indocyanine green
7336002 | Trazodone
7561000 | Dexamethasone preparation
7577004 | Ciprofloxacin
7624002 | Sodium perborate product
7836006 | Expectorants
7947003 | Acetylsalicylic acid
7959004 | Teniposide
7982002 | Phenmetrazine hydrochloride
8028001 | Butacaine
8109005 | Alimemazine
8163008 | Nitroprusside
8348002 | Cyclopentolate
8372007 | Promethazine
8416000 | Dicloxacillin
8571001 | Vasoconstrictor

8658008 | Human serum albumin preparation
8661009 | Replacement preparation
8692006 | Metamfetamine
8696009 | Antispasmodic
9190005 | Tropicamide
9268004 | Butabarbitone
9307009 | Animal serum globulin preparation
9500005 | MAOI - Phenelzine
9542007 | Antihepatitis B immunoglobulin
9690006 | Nikethamide
9745007 | Sucrose product
9778000 | Cytomegalovirus immune globulin
9944001 | Chlorphenamine
10099000 | Ketoprofen
10135005 | Cinchona alkaloid
10243007 | Benzoic and salicylic acid ointment
10312003 | Prednisone preparation
10332002 | Vaccine - bacterin - toxoid
10355004 | Injectable vitamin preparation
10356003 | Undecylenic acid and undecylenate salt
10368007 | Colloidal oatmeal powder
10422008 | Nitropentaerythrol
10504007 | Doxycycline
10515002 | Lututrin preparation
10555000 | Tocainide
10632007 | Multivitamin preparation
10668002 | Skin anti-infective
10712001 | Glucagon product
10756001 | Haloperidol
10784006 | Anti-psychotic agent
11185009 | Autogenous vaccine - autogenous bacterin
11260008 | Ophthalmic anti-infective preparation
11402001 | Medicinal enzyme
11430001 | Tetracyclic antidepressant drug
11563006 | Vitamin D preparation
11719000 | Cetylpyridinium
11777007 | Ophthalmic antibiotic
11783005 | Stool softener
11796006 | Methysergide
11841005 | Doxepin
11847009 | Naproxen
11866009 | Plague vaccine
11959009 | Procainamide
12096000 | Nystatin
12236006 | Pancreatin product
12289007 | Whole blood product
12335007 | Diatrizoate
12369008 | Oxytocin product
12425002 | Human white blood cell product
12436009 | Vinblastine
12495006 | Magnesium citrate
12512008 | Triamterene
12559001 | Emetine
12657005 | Fludrocortisone acetate preparation
12839006 | Estradiol product
12968008 | Vitamin products
13132007 | Dextran product
13154009 | Chlormethine
13178004 | Rickettsial vaccine
13252002 | Salsalate
13414000 | Cefadroxil
13432000 | Nortriptyline
13512003 | Minocycline
13525006 | Acetylcholine preparation
13565005 | Bisacodyl
13592004 | Pyrazinamide
13664004 | BAL - British Anti-Lewisite
13790009 | Oral form iron preparation
13800009 | Naftifine
13813003 | Biotin preparation
13929005 | Spironolactone
13936006 | Butorphanol

13965000 | Valproic acid
14054003 | Vitamin B complex preps
14103001 | Opiate antagonist
14170004 | Capreomycin
14542005 | Caffeine and sodium benzoate injection solution vial
14601000 | Anticholinergic agent
14706000 | Pheneticillin
14728000 | Chloroquine
14745005 | Hepatitis A vaccine
14814001 | Trimethobenzamide
14816004 | Cocaine product
15117003 | Coagulant
15222008 | Enalapril
15375005 | Phenanthrene derivative
15383004 | Levodopa
15389000 | Hydantoin derivative anticonvulsant
15432003 | Ethinylestradiol
15772006 | Beta 1 blocking product
15857002 | Ethanolamine derivative antihistamine
16031005 | Dexchlorpheniramine
16037009 | Terfenadine product
16047007 | Benzodiazepine
16131008 | Animal antiserum
16403005 | Non-steroidal anti-inflammatory agent
16602005 | Hydrocortisone preparation
16787005 | Streptococcus equisimilis - suis antiserum
16791000 | Cefradine
16832004 | Conjugated estrogens
16858004 | Urea
16867004 | Sulfathiazole
16970001 | Proguanil
16977003 | Lithium carbonate
17016006 | Animal gamma globulin
17055007 | Systemic acidifier
17308007 | Dapsone
17386008 | Anti-infective agent
17502009 | Gallamine triethiodide
17554004 | Paramethasone preparation
17558001 | Corn oil
17600005 | Diagnostic radiopharmaceuticals
17805003 | Lithium citrate
17859000 | Polyvalent crotalidae antivenin
17893001 | Skeletal muscle relaxant
18002004 | Auranofin
18125000 | Fluocinonide
18335001 | Aureolic acid
18340009 | Oxychlorosene
18381001 | Prindolol
18511007 | Generic drug
18548003 | Methylphenidate
18679008 | Potassium-removing resin
18811003 | L-asparaginase preparation
18914005 | Hydroflumethiazide
18952006 | Econazole
19194001 | ddI
19225000 | Lorazepam
19232009 | Prilocaine
19261005 | Sulfinpyrazone
19315007 | Pyridoxine hydrochloride preparation
19403009 | Flurazepam
19405002 | Netilmicin
19581007 | Parasympathomimetic agent
19583005 | Diclofenamide
19630009 | Silver sulfadiazine
19768003 | Alkylating drugs
19841008 | Ceftriaxone
20091003 | Somatotropin releasing factor preparation
20201001 | Nafoxidine
20237006 | Dihydrotachysterol preparation
20249007 | Progestin preparation
20303001 | Vinblastine sulfate
20320002 | Hydrocodone

20577002 | HCG - Human chorionic gonadotrophin product
20865003 | Diflunisal
20969008 | Polyvalent snake antivenin
21069002 | Lipotropic agent
21159006 | Pargyline
21451004 | Indium-113m chloride
21691008 | Magnesium trisilicate
21701005 | Cromoglicic acid
21767006 | Iron dextran complex
21788002 | Phenazopyridine hydrochloride
21986005 | Erysipelothrix rhusiopathiae antiserum
22091006 | Hormone preparation
22168005 | Glycopyrronium
22198003 | Metolazone
22274004 | Methandriol preparation
22474002 | Aldosterone preparation
22587006 | Depolarizing muscle relaxant
22657006 | Calcitonin product
22672005 | Amfetamine group
22696000 | Hydralazine
22801004 | Sterile water solution
22826008 | Ammonia detoxicant
22969001 | Oxytetracycline
23079006 | Vincristine
23222006 | Antisera
23343005 | Human platelet product
23532003 | Phenmetrazine
23822003 | Human plasma fraction
23827009 | Sulfacetamide
23838009 | Nonspecific alpha-adrenergic blocking agent
23888001 | Cascara
24036003 | Antianemia drug
24450004 | Ethambutolol
24504000 | Methylcellulose
24866006 | Salmonella typhimurium antiserum
24891006 | Sodium amidotrizoate
25014009 | Tripelennamine
25057004 | Mercurial diuretic product
25076002 | Carisoprodol
25085002 | Cholecystokinin preparation
25142008 | Trilostane
25246002 | Allopurinol
25398003 | Ichthammol
25419009 | Barium sulfate
25465007 | Escherichia coli monoclonal antibody
25673006 | Omeprazole
25716001 | Tetanus toxoid, human
25849001 | SNP - Sodium nitroprusside
25860005 | Terconazole
25912006 | Menadiol sodium diphosphate preparation
25995007 | Triprolidine
26122009 | Dimetindene
26124005 | Glipizide product
26244009 | Antimuscarinic
26303005 | Hexestrol preparation
26370007 | Hemostatic agent
26456008 | Levothyroxine sodium preparation
26458009 | Diphenhydramine
26462003 | Cyproheptadine
26503009 | Deserpidine
26523005 | Dobutamine
26548008 | Pancreatic hormone product
26574002 | Droperidol
26580005 | Digestant
26736008 | Ferrous gluconate
26800000 | Midazolam
26842003 | Burbot liver oil preparation
27035007 | Heavy metal antagonist
27085003 | Sodium polystyrene sulphonate
27196008 | Bupivacaine
27242001 | Methylprednisolone preparation
27479000 | AZT - Azidothymidine

27518004 | Parathyroid hormone
27548001 | LA - Local anesthetic drug
27566006 | Drug vehicle preservative
27638005 | T-PA - tissue-type plasminogen activator
27658006 | Amoxicilline
27707001 | Medroxyprogesterone acetate preparation
27754002 | Pyrazinobutazone
27867009 | Antineoplastic
28028002 | Synthetic hormone preparation
28149003 | Pentostatin
28235004 | Doxapram
28240007 | Eye cosmetic
28410007 | alpha-Carboxypenicillin
28415002 | Methscopolamine
28426008 | Fluocinolone
28506006 | Flucytosine
28748001 | Chloral hydrate
28841002 | Anhydrohydroxyprogesterone
28906000 | Percoid liver oil preparation
29051009 | Loop diuretic product
29058003 | Halcinonide
29089004 | Dibromomannitol
29121001 | Mersalyl
29129004 | Oxymetazoline
29156002 | Mechlorethamine
29175007 | Rifampicin product
29237004 | Arachnida antivenin
29439004 | Captopril
29620001 | beta-Tocopherol preparation
29840005 | Amoxapine
29877002 | MAOI - Isocarboxazid
29883004 | Class IV antiarrhythmic
29896003 | Betamethasone
29987004 | Vitamins A and D preparation
30010009 | Cyanocobalamin product
30125007 | Senna product
30306003 | Vitamin B1 preparation
30317002 | Cisapride
30427009 | Erythromycin product
30466001 | Chloramiphene
30492008 | Diuretic
30666008 | Toxoid (human)
30729008 | Iron preparation
30761007 | Mannitol product
30964009 | Methyprylone
30988006 | Dienestrol preparation
31025003 | Dibenzoxazepine derivative antipsychotic agent
31087008 | Ampicillin
31181006 | Injectable testosterone preparation
31231007 | H2O2 - Hydrogen peroxide
31305008 | Streptococcus equisimilis antiserum
31306009 | Quinidine
31352002 | Normal colostral whey
31587003 | Almond oil liquid
31684002 | Buprenorphine
31690003 | Bethanechol
31692006 | Pentamidine
31716004 | Frozen plasma product, human
31865003 | Fluconazole
31872002 | Pramocaine
31970009 | Nitrate vasodilator
32249005 | Antiviral drugs
32313007 | Enflurane
32460003 | Beclometasone dipropionate nasal preparation
32462006 | Melanocyte stimulating hormone releasing factor preparation
32473004 | Spider antivenin
32474005 | Probucol
32583002 | Antiseborrheic drug
32647002 | Ergotamine
32653002 | Ergosterol preparation
32792001 | Trimethoprim
32823007 | Maprotiline

32841004 | NaCl - Sodium chloride solution
32955006 | Domperidone
32960005 | Thiosalicylate
33124007 | Butamidum
33133009 | Ophthalmic anti-inflammatory preparation
33219003 | Tricyclic antidepressant
33231001 | Pentobarbital
33234009 | Var/Vac
33236006 | Cortisone acetate preparation
33252009 | beta-adrenoceptor blocking drug
33378002 | Desipramine
33484000 | Proprietary drug
33588000 | Thioridazine
33589008 | Cardiac glycoside
33664007 | Acetazolamide
33675006 | Carbachol
33682005 | Mydriatic
33815001 | Streptococcus suis antiserum
34012005 | Sulfonylurea product
34217006 | Oxyquinoline
34364009 | Mefenamic acid
34393009 | Tolazamide
34462007 | Natamycin
34598001 | Immunobiologic agent available from the Centers for Disease Control and Prevention
34599009 | Thyroglobulin preparation
34689006 | Hepatitis B vaccine
34693000 | Zalcitabine
34696008 | Sodium ascorbate preparation
34731007 | Carbenicillin
34816007 | Cod liver oil
34833000 | Hydrocortisone ophthalmic preparation
34929006 | Benzethonium
35035001 | Orphenadrine
35063004 | Ribavirin
35097006 | Flea antigen
35282000 | Gemfibrozil
35300007 | Daunorubicin
35324004 | Paraldehyde product
35392005 | Calcium-removing resin
35476001 | Silver nitrate product
35531004 | Hydrocortamate preparation
35736007 | Cholera vaccines
35768004 | Oxybutynin
35792007 | Peritoneal dialysis solution
35967000 | Medazepam
35983000 | Cellular blood product, human
36068003 | Pyrantel
36113004 | Imipramine
36168000 | Anti-infective nasal preparations
36218003 | Thiethylperazine
36236003 | Antidepressant
36391008 | Primaquine
36537006 | Ambenonium
36594001 | Tiabendazole
36621009 | Medroxyprogesterone
36642006 | Propantheline
36763003 | Immune globulin IM
36795002 | Piperazine estrone sulfate
36893000 | Ceftazidime
36909007 | Phenindamine
36940000 | Thioxanthene derivative antipsychotic agent
36944009 | Vitamin B6 preparation
37084008 | 4-quinolones
37146000 | Typhus vaccine
37306000 | Vidarabine
37400007 | Magnesium sulfate
37628007 | Cefalotin
37643009 | Autogenous bacterin
37773009 | Diphenylbutylpiperidine derivative antipsychotic agent
37803001 | d-TC - Tubocurarine
38076006 | Sodium thyroxine
38131009 | Synthetic drug

38166006 | Tolnaftate
38181000 | Antineoplastic antibiotic
38231004 | Polysaccharide iron complex
38268001 | Propionic acid-ibuprofen
38314008 | Isotretinoin
38413003 | Otic dosage form product
38578004 | Megestrol preparation
38828006 | Sodium thiosulfate
38900001 | Proliferant agent
39064002 | Acetohexamide
39124003 | Methohexital
39128000 | Famotidine
39142008 | Phendimetrazine
39252001 | Water balance agent
39359008 | Penicillin V
39432004 | Deodorant
39487003 | Regular insulin product
39516004 | Disulfiram product
39598004 | Marijuana derivative
39608003 | Pentazocine
39707000 | Aminobenzoic acid
39741008 | Drug groups primarily affecting the gastro-intestinal system
39860005 | Fructose product
39939003 | Phenyltoloxamine
40162007 | Bulk-forming laxatives
40232005 | Ketoconazole
40339003 | Calcium lactate
40429005 | Etomidate
40430000 | Bromelains
40556005 | Phenytoin product
40562000 | Methylergonovine preparation
40589005 | Amitriptyline
40606008 | Orphan drug
40648001 | Fentanyl product
40652001 | Cromolyn sodium
40820003 | Carbamazepine product
40877002 | Streptomycin
40901001 | Desferrioxamine mesylate
40905005 | Beractant
40974005 | Phenylpiperone
40999006 | CCNU - Lomustine
41001009 | Dinoprost
41015006 | Metaraminol
41147003 | Perphenazine
41193000 | Aciclovir
41324009 | Propiomazine
41365009 | Fluphenazine
41367001 | Enterogastrone preparation
41399007 | Oxazolidinedione
41470001 | Corbadrine
41493007 | Dicycloverine
41549009 | Angiotensin-converting enzyme inhibitor
41985001 | Bitolterol
42082003 | Vancomycin
42098005 | Dexamethasone ophthalmic preparation
42271003 | Glutamic acid product
42348003 | Methyltestosterone
42374009 | Dextromethorphan hydrobromide
42444000 | Secobarbital
42514000 | Procaine product
42638008 | Methylrosanilium chloride
42714002 | Escherichia coli antiserum
42720001 | Miconazole
42863006 | Desmopressin acetate preparation
43343000 | Magaldrate
43499003 | Injectable interferon
43533002 | Chloramphenicol ophthalmic preparation
43594006 | Skin bleaching preparations
43684009 | Misoprostol product
43747001 | Drug excipient
43753001 | Dydrogesterone
43879000 | Flunisolide

43927002 | Analeptic agent
44175000 | Diperodon
44418001 | Percomorph liver oil preparation
44658005 | Promazine
44731005 | Otic form hydrocortisone
44790008 | Ethosuximide
44798001 | Dinoprostone product
44938006 | Cefoperazone
44990002 | Procyclidine
45218006 | Meclastine
45311002 | Terbutaline
45313004 | Propylpiperazine derivative of phenothiazine
45355008 | Rocky Mountain spotted fever vaccine
45518007 | Thiazide diuretic product
45680002 | Tolmetin
45844004 | Sulfasalazine
45888006 | gamma-Tocopherol preparation
45923000 | E331 - Sodium citrate
46009007 | Chlorambucil
46063005 | Psychotherapeutic agent
46123006 | Ascorbic acid product
46233009 | FLU - Influenza vaccine
46436003 | Haloprogin
46479001 | Encainide
46532003 | Brilliant green
46547007 | Labetalol
46576005 | Flecainide
46709004 | Methylphenobarbital
46741005 | SA - Salicylic acid
46913003 | Edrophonium
47065008 | Quinine
47120002 | Hexamidinum
47124006 | Aminoglutethimide
47140009 | Medrysone
47331002 | CPZ - Chlorpromazine
47527007 | Phenindione
47602007 | Nalidixic acid
47755009 | Potassium sparing diuretic
47772002 | Fenfluramine hydrochloride
47898004 | Verapamil
48028008 | Haemophilus influenzae Type b (Hib)/diphtheria/tetanus/pertussis vaccine
48174005 | Ranitidine
48256008 | Benzyl benzoate
48279009 | Emollient product
48351000 | Butadione
48546005 | Diazepam product
48603004 | Warfarin
48614001 | Clobetasol
48647005 | Pancrelipase
48698004 | Calcium-channel blocker
48836000 | Amikacin
48875009 | Dihydroergotamine product
48899009 | Hyoscyamine
49019002 | Prednisolone ophthalmic preparation
49157004 | Uricosuric product
49192007 | Antiflatulent agent
49267000 | Hydroxyphenylbutazone
49299006 | Protriptyline
49485009 | Norfloxacin
49577002 | Minoxidil
49590009 | Class I antiarrhythmic
49600008 | Normal animal plasma
49617001 | Disodium enoxolone
49663007 | Sunscreening preparations
49669006 | Escherichia coli - pasteurella multocida - salmonella typhimurium antiserum
49688004 | Hexocyclium
49694007 | Mucolytics
49705006 | Idoxuridine
49802003 | Anti-D (Rh) immunoglobulin
49860008 | Gold product
49953001 | Pheniramine
49992008 | Dexamethasone sodium phosphate

50094009 | Hetastarch
50137001 | Glycine solution
50182002 | Hemodialysis fluid
50244000 | Intravenous anesthetic
50256004 | Antitrichomonal agent
50318003 | Luteal hormone
50335004 | Levorphanol
50520001 | Neomycin B
50841004 | Otic form chloramphenicol
50868004 | Dexamfetamine
51013009 | Sulfadimethoxine
51073002 | Phenobarbital product
51103004 | Hypertonic saline
51126006 | Benzestrol preparation
51132001 | Hyaluronidase product
51326002 | BCNU - Carmustine
51334008 | Cycloserine
51361008 | Amantadine
51425004 | Scorpion antivenin
51555009 | Desiccated liver preparation
51752005 | Chologon
51758009 | Amidine
51779009 | Cephalosporin -class of antibiotic-
51908007 | Prenylamine
51992002 | Gastrin preparation
52017007 | Antiemetic
52108005 | Ferrous fumarate
52215008 | Desonide
52388000 | Prednisolone
52412009 | Methylergonovine maleate preparation
52423008 | Tar preparation
52451004 | Dihydroindolone derivative antipsychotic agent
52489009 | Vasopressin tannate preparation
52628005 | Typhoid-paratyphoid vaccine
52761007 | Anticholinesterase parasympathomimetic
52883001 | para-Hydroxyamphetamine
52896000 | Chinoform
53009005 | Analgesic
53480001 | Phentermine
53584007 | Methacholine
53640004 | Fluoxetine
53641000 | Flavoxate
53691001 | Calcium gluconate
53793005 | Escherichia coli antibody
53800008 | Dithranol
53848009 | Metyrapone
53855006 | Neostigmine bromide
54142000 | Domiphen
54344006 | Flurbiprofen
54391004 | Levamisole
54406003 | Methoxamine
54541002 | Ergometrine preparation
54544005 | Pethidine
54577009 | Ceftizoxime
54659006 | Thiamine hydrochloride preparation
54705000 | Temazepam
54765002 | Phenylephrine
54824008 | Isometheptene
54882005 | Amfepramone
54887004 | Cefalexin
54972005 | Tretinoin
54982006 | Promethestrol preparation
55015008 | E325 - Sodium lactate
55217007 | Calcium carbonate
55432002 | Azlocillin
55556000 | Tetracaine
55673009 | Iothalamate sodium I^125^
55745002 | Propranolol
55782006 | Amide-type anesthetic
55784007 | Natural hormone preparation
55830003 | Menotrophin
55867006 | Aminophylline

55889005 | Praziquantel
56011005 | Stinging insects immunotherapy extract
56014002 | Hydroxyprogesterone
56032002 | Androstanolone
56059005 | Mebendazole product
56123002 | Hexamine
56230001 | Bretylium
56234005 | Growth hormone product
56240003 | Paramethasone acetate preparation
56281006 | Anesthetic ether
56480005 | Parabromdylamine
56549003 | Metoclopramide
56602009 | Hydroxycarbamide
56615000 | Polyestradiol phosphate preparation
56656002 | Magnesium-based antacid
56844000 | Yellow fever vaccine product
56928005 | Etoposide product
56934003 | PVP - Polyvinylpyrrolidone
57002000 | Chlorprothixene
57066004 | Cisplatin product
57123008 | Urinary acidifier
57191001 | Chloramphenicol product
57263002 | Oxiconazole
57376006 | Monosodium carbonate
57538001 | Chlortetracycline
57616006 | STD - Sodium tetradecyl sulfate
57670008 | Cefoxitin
57752001 | Gentamicin product
57811004 | Dihydrocodeine
57819002 | Somatotropin inhibiting factor preparation
57845006 | Isoprenaline
57853003 | Clidinium
57893000 | Chlortalidone
57952007 | Lipid lowering drug
58050004 | Antiparkinsonian agent
58098008 | Tetanus toxoid adsorbed
58360000 | Phenazocine
58467001 | Papaverine
58502006 | Propylamine derivative antihistamine
58760003 | Antimetabolites
58805000 | Pituitary hormone preparation
58883005 | Clindamycin
58892008 | F3T - trifluorothymidine
58905004 | Diazoxide
58944007 | Vasodilator
59057006 | Antihemophilic factor preparation
59187003 | Dopamine product
59240002 | MMC-Mitomycin-C
59245007 | Hypophysiostatic factor preparation
59255006 | Sulfonamide
59261009 | Oral contraceptive preparation
59270007 | Loxapine
59456005 | Astemizole
59589008 | Pyrimethamine product
59594008 | Nondepolarizing agent
59751001 | Cough suppressant
59799001 | Osmotic diuretic product
59941008 | Diltiazem
59953007 | Pyridostigmine
59999009 | Tetanus and diphtheria toxoid adsorbed for adult use
60090003 | Soft drug
60093001 | Phenylpropanolamine hydrochloride
60149003 | Indometacin
60169008 | Antacid
60468008 | Magnesium hydroxide
60520004 | Thiamine mononitrate preparation
60533005 | Astringent drug
60541005 | Lanatoside C
60682004 | Echothiophate
60731009 | Diethylcarbamazine
60767002 | Normal animal serum
60881009 | Heroin

60972002 | Irrigating solution
60978003 | Barbiturate
61020000 | Thyroid preparation
61093008 | Prolactin inhibiting factor preparation
61123001 | Amobarbital sodium
61132004 | Gas gangrene antitoxin
61153008 | Measles, mumps and rubella vaccine
61181002 | Meprednisone preparation
61408004 | Molindone
61457001 | Adrenal hormone preparation
61602008 | Pertussis vaccine
61621000 | Laxative
61623002 | Buclizine
61650007 | Niacinamide preparation
61651006 | Cefamandole
61862008 | Meticillin
61939007 | Diphenylmethane laxative
61946003 | Estrogenic preparation
62051009 | Dichlorisone preparation
62063004 | Nasal anti-inflammatory preparation
62191005 | Injectable liver preparation
62288001 | Anthelmintic
62294009 | Varicella-zoster immune globulin
62353003 | Clomifene citrate
62354009 | Bronchitis vaccine
62529008 | Tiotixene
62560008 | Fluorometholone ophthalmic preparation
62782004 | Clonidine
63094006 | Anticonvulsant
63136007 | Phytonadione preparation
63318000 | Benzoic acid
63338004 | Drug flavoring
63470003 | Fluoxymesterone preparation
63639004 | Niacin preparation
63682003 | Halothane
63758001 | Norethindrone preparation
63822004 | Vitamin E preparation
64083002 | Genitourinary smooth muscle relaxant
64115004 | Amodiaquin
64127001 | Dactinomycin
64240003 | Methandrostenolone preparation
64296005 | Staphylococcus toxoid
64349004 | Griseofulvin
64462001 | Terpin
64558005 | Metixene
64851009 | Diiodohydroxyquin
64878006 | Methylthiouracil
65020006 | Benzocaine
65026000 | Ephedrine
65041000 | Biperiden
65092008 | Chloropyrilene
65265006 | Thromboplastin
65461004 | Adrenal cortex extract
65484006 | Prostacycline PGI>2< preparation
65502005 | Epinephrine preparation
65505007 | Gastrointestinal smooth muscle relaxant
65627005 | 4-Amino-2-methyl-1-naphthol hydrate preparation
65628000 | Skin agent
65638005 | Corticotropin zinc hydroxide preparation
65695006 | General inhalation anesthetic
65823007 | Protective agent
65884003 | Antrapurol
65965000 | Micrurus fulvius antivenin
66065008 | Saline laxative
66094001 | Probenecid
66125007 | Flunisolide nasal preparation
66243004 | Norethindrone acetate preparation
66261008 | Tetracycline
66349002 | Androgenic preparation
66441000 | Pantothenic acid preparation
66492008 | Isoflurane
66493003 | Theophylline

66602007 | Stanozolol preparation
66672008 | Ester type local anesthetic
66742008 | Pigmenting agent
66854004 | Betamethasone valerate
66859009 | Dipyridamole
66860004 | Aluminum chloride
66919007 | Methyclothiazide
66971004 | Colestipol
67213005 | Lymphocyte immune globulin
67423003 | Acylaminopenicillin
67440007 | Alpha-adrenergic blocking agent
67477004 | Valproate
67507000 | Antiarrhythmic drug
67585005 | Topical anesthetic
67735003 | Paclitaxel
67891001 | Second generation cephalosporin
67939000 | Apomorphine
68088000 | Acebutolol
68113000 | Hib oligosaccharide conjugate (diphtheria CRM>197< protein conjugate) vaccine
68206008 | Calcitriol preparation
68208009 | Diphtheria and tetanus toxoid adsorbed
68395000 | Calcium chloride
68398003 | Somatomedin preparation
68402007 | Carbonic anhydrase inhibitor
68407001 | Hydrogen peroxide solution 30%
68422006 | Cloxacillin
68424007 | Isoflurophate
68444001 | Doxorubicin
68461003 | Surgical jelly
68490009 | Sodium propionate
68622003 | Secretin preparation
68702006 | Sodium aurothiomalate
68774008 | Isoxsuprine
68864001 | Polyvalent reptile antivenin
68887009 | Methotrexate
68892006 | Penicillinase-resistant penicillin
69107004 | Dantrolene
69204002 | Guanadrel
69227005 | Class II antiarrhythmic drug
69236009 | Amiodarone
69242008 | Miotic
69331001 | Aminocyclohexylpenicillin
69431002 | Immunosuppressant
69509008 | Biological agent
69576000 | Menadione preparation
69708003 | Clonazepam
69801001 | Sodium estrone sulfate preparation
69879000 | Altretamine
69918003 | Aztreonam
69967001 | Sucralfate
70047000 | Sulfamethoxazole
70216000 | Undecylenate salt
70254000 | Sulfamethizole
70343008 | Piperazine derivative antihistamine
70379000 | Sodium chloride
70447008 | Polyvalent pneumococcal vaccine
70460002 | Fish liver oil preparation
70702006 | Deferoxamine
70776005 | Pemoline
70841003 | Chymotrypsin preparation
70864001 | Meprobamate
70934008 | Demecarium
71181003 | Vaccine
71289008 | Snake antivenin
71338005 | Vasopressin preparation
71451001 | Kanamycin
71453003 | Mupirocin
71455005 | Flurandrenolone preparation
71462001 | Podophyllum resin
71516007 | Ergocalciferol preparation
71584004 | Sodium phosphate
71634000 | Chlormezanone

71699007 | Trifluoperazine
71724000 | Ferrous sulfate
71731001 | Medrysone ophthalmic preparation
71759000 | Nitroglycerin
71770007 | Monoamine oxidase inhibitor
71798005 | Fenoprofen
71837009 | Cytotoxic agent
72312007 | Cyclandelate
72398005 | Ophthalmic sulfonamide preparation
72416006 | Metacycline
72623000 | 2-Amino-6-mercaptopurine
72824008 | Cholestyramine
72870001 | Scopolamine
72924009 | Clofazimine
72968006 | Sodium salicylate
72996000 | Drug builder
73074003 | Colistin
73093001 | Neomycin
73133000 | Colchicine
73170009 | Menthol
73212002 | Adipiodone
73228007 | Non-steroidal estrogen preparation
73274006 | Human plasma cryoprecipitate
73277004 | Mecamylamine
73454001 | Desmopressin preparation
73572009 | Morphine
73585007 | Phenytoin sodium
73627008 | Caloric agent
73639000 | Prescription drug
73647000 | Dipivefrine
73756003 | Amobarbital
73763003 | Extended spectrum penicillin
73805002 | Thyrotropin releasing hormone preparation
73949004 | Atropine
73986003 | Cefuroxime
74022005 | Mepenzolate
74065006 | Prazepam
74074008 | Atracurium
74147005 | Edetate disodium
74213004 | Indapamide
74226000 | Vitamin K preparation
74470007 | Cyclophosphamide
74480006 | Potassium supplement
74575000 | Piperacillin
74583006 | Hydroquinone product
74626007 | Drug diluent
74632002 | Succinimide
74674007 | Propofol
74771007 | Phenoxybenzamine
74780007 | Anthraquinone laxatives
74782004 | Naturally occurring alkaloid
74798006 | Pipenzolate
74819009 | Acetohydroxamic acid
75029008 | Desoxycorticosterone preparation
75201000 | Bacterin - toxoid
75203002 | Mometasone
75292001 | Vincristine sulfate
75366006 | Sodium calciumedetate
75429004 | Dexbrompheniramine
75501004 | Bromazine
75661008 | delta-Tocopherol preparation
75748004 | Theophylline sodium glycinate
75770001 | Urea topical preparation
75927008 | Floxuridine
75959001 | Tamoxifen
75969007 | Gonadotropin releasing factor preparation
76058001 | Prazosin
76155001 | Iopanoic acid
76286000 | Gallamine
76289007 | Xylometazoline
76385003 | Alpha 1 adrenergic blocking agent
76390000 | Practolol

76591000 | Bleomycin
76696004 | Noscapine
76759004 | Disopyramide product
76962009 | Iproniazid
77035009 | Clofibrate
77048008 | Dip/ser
77237006 | Emetic
77331005 | Normal animal globulin concentrate
77390008 | Benzatropine
77398001 | Antidiarrhea agent
77549006 | Terpene
77657003 | Illegal drug
77731008 | N-Acetylcysteine
77750008 | Dacarbazine
77856005 | Esmolol
77885004 | Mestranol product
77898008 | Simethicone
78025001 | Ganciclovir product
78057000 | Carbol-fuchsin topical solution
78174002 | Mezlocillin
78379001 | Reserpine
78411008 | Thiopental sodium
78439009 | Citrate salt
78449007 | Nitrazepam
78507004 | Benzylpenicillin
78542000 | Potassium citrate product
15911003 | Cow's milk protein sensitivity
25868003 | Soy protein sensitivity
59037007 | Drug intolerance
61712006 | Transient gluten sensitivity
72354005 | Oral contraceptive intolerance
91929009 | Anti-infectives allergy
91930004 | Allergy to eggs
91931000 | Erythromycin allergy
91932007 | Allergy to fruit
91933002 | Macrolide allergy
91934008 | Nut allergy
91935009 | Allergy to peanuts
91936005 | Penicillin allergy
91937001 | Seafood allergy
91938006 | Allergy to strawberries
91939003 | Sulfonamide allergy
91940001 | Allergy to walnut
188336009 | Laboratory animal dander allergy
197493001 | Malabsorption due to intolerance to soy protein
197494007 | Intestinal malabsorption of fat
200848004 | Allergic contact dermatitis due to plants, except food
213020009 | Egg protein allergy
232346004 | Allergy to cat dander
232347008 | Allergy to animal dander
232348003 | Feather allergy
232349006 | House dust allergy
232350006 | House dust mite allergy
235719002 | Food intolerance
235720008 | Malabsorption caused by intolerance to carbohydrate
292545003 | Adverse reaction to oxitropium
293580007 | Allergy to over-the-counter drug
293582004 | Analgesic allergy
293583009 | Non-opioid analgesic allergy
293584003 | Acetaminophen allergy
293585002 | Salicylate allergy
293586001 | Aspirin allergy
293587005 | Opioid analgesic allergy
293588000 | Pentazocine allergy
293589008 | Phenazocine allergy
293590004 | Methadone analog allergy
293591000 | Dextromoramide allergy
293592007 | Dextropropoxyphene allergy
293593002 | Dipipanone allergy
293594008 | Methadone allergy
293595009 | Morphinan opioid allergy
293596005 | Buprenorphine allergy

293597001 | Codeine allergy
293598006 | Diamorphine allergy
293599003 | Dihydrocodeine allergy
293600000 | Nalbuphine allergy
293601001 | Morphine allergy
293602008 | Opium alkaloid allergy
293603003 | Pethidine analog allergy
293604009 | Alfentanil allergy
293605005 | Fentanyl allergy
293606006 | Pethidine allergy
293607002 | Phenoperidine allergy
293608007 | Meptazinol allergy
293609004 | Levorphanol allergy
293610009 | Non-steroidal anti-inflammatory drug allergy
293611008 | Acemetacin allergy
293612001 | Azapropazone allergy
293613006 | Diclofenac allergy
293614000 | Etodolac allergy
293615004 | Felbinac allergy
293616003 | Fenbufen allergy
293617007 | Fenoprofen allergy
293618002 | Flurbiprofen allergy
293619005 | Ibuprofen allergy
293620004 | Indometacin allergy
293621000 | Ketoprofen allergy
293622007 | Ketorolac allergy
293623002 | Mefenamic acid allergy
293624008 | Nabumetone allergy
293625009 | Naproxen allergy
293626005 | Nefopam allergy
293627001 | Oxyphenbutazone allergy
293628006 | Phenylbutazone allergy
293629003 | Piroxicam allergy
293630008 | Sulindac allergy
293631007 | Tenoxicam allergy
293632000 | Tiaprofenic acid allergy
293633005 | Tolmetin allergy
293634004 | Diagnostic agent allergy
293635003 | Tuberculin allergy
293636002 | Radiopharmaceutical allergy
293637006 | Contrast media allergy
293638001 | X-ray contrast media allergy
293639009 | Magnetic resonance imaging contrast media allergy
293640006 | Allergen extract vaccine allergy
293641005 | Diagnostic dye allergy
293643008 | Drug groups primarily affecting the gastrointestinal system allergy
293644002 | Ulcer healing drug allergy
293645001 | Bismuth chelate allergy
293646000 | Sucralfate allergy
293647009 | Liquorice allergy
293648004 | Misoprostol allergy
293649007 | H2 receptor antagonist allergy
293650007 | Cimetidine allergy
293651006 | Famotidine allergy
293652004 | Nizatidine allergy
293653009 | Ranitidine allergy
293654003 | Proton pump inhibitor allergy
293655002 | Omeprazole allergy
293656001 | Lansoprazole allergy
293657005 | Carbenoxolone allergy
293658000 | Pirenzepine allergy
293659008 | Pancreatin allergy
293660003 | Mesalazine allergy
293662006 | Olsalazine allergy
293663001 | Sulfasalazine allergy
293664007 | Antacid allergy
293665008 | Magnesium trisilicate allergy
293666009 | Aluminum hydroxide allergy
293667000 | Antidiarrheal drug allergy
293668005 | Loperamide allergy
293669002 | Kaolin allergy
293670001 | Motility stimulant allergy

293671002 | Cisapride allergy
293672009 | Antiemetic allergy
293673004 | Nabilone allergy
293674005 | Domperidone allergy
293675006 | Metoclopramide allergy
293676007 | 5-Hydroxytryptamine-3-receptor antagonist allergy
293677003 | Laxative allergy
293678008 | Bisacodyl allergy
293679000 | Dantron allergy
293680002 | Sodium picosulfate allergy
293681003 | Lactulose allergy
293682005 | Magnesium sulfate allergy
293683000 | Bulk-forming laxative allergy
293684006 | Anthraquinone laxative allergy
293685007 | Cascara allergy
293686008 | Senna allergy
293687004 | Docusate allergy
293688009 | Antispasmodic allergy
293689001 | Hyoscine butylbromide allergy
293690005 | Peppermint oil allergy
293691009 | Alverine allergy
293692002 | Mebeverine allergy
293693007 | Dicycloverine allergy
293694001 | Mepenzolate allergy
293695000 | Pipenzolate allergy
293696004 | Poldine allergy
293697008 | Propantheline allergy
293698003 | Bile agent allergy
293699006 | Chenodeoxycholic acid allergy
293700007 | Dehydrocholic acid allergy
293701006 | Ursodeoxycholic acid allergy
293702004 | Chenodeoxycholic and ursodeoxycholic acid allergy
293703009 | Anesthetics and medical gases allergy
293704003 | General anesthetic drug allergy
293705002 | Intravenous anesthetics allergy
293706001 | Etomidate allergy
293707005 | Ketamine allergy
293708000 | Propofol allergy
293709008 | Thiopental allergy
293710003 | Methohexitone allergy
293711004 | Inhalational anesthetics allergy
293712006 | Enflurane allergy
293713001 | Ether, anesthetic allergy
293714007 | Halothane allergy
293715008 | Isoflurane allergy
293716009 | Trichloroethylene allergy
293717000 | Desflurane allergy
293718005 | Local anesthetic drug allergy
293719002 | Bupivacaine allergy
293720008 | Cinchocaine allergy
293721007 | Prilocaine allergy
293722000 | Lidocaine allergy
293723005 | Cocaine allergy
293724004 | Benzocaine allergy
293725003 | Tetracaine allergy
293726002 | Oxybuprocaine allergy
293727006 | Procaine allergy
293728001 | Proxymetacaine allergy
293730004 | Immunostimulant allergy
293732007 | Amifostine allergy
293733002 | Aldesleukin allergy
293734008 | Colony stimulating factors allergy
293735009 | Molgramostim allergy
293736005 | Lenograstim allergy
293737001 | Filgrastim allergy
293738006 | Levamisole allergy
293739003 | Antineoplastic allergy
293740001 | Alkylating drug allergy
293741002 | Mitobronitol allergy
293742009 | Busulfan allergy
293743004 | Treosulfan allergy
293745006 | Thiotepa allergy

293746007 | Nitrogen mustard derivative allergy
293747003 | Chlorambucil allergy
293748008 | Cyclophosphamide allergy
293749000 | Ethoglucid allergy
293750000 | Ifosfamide allergy
293751001 | Melphalan allergy
293752008 | Estramustine allergy
293753003 | Chlormethine allergy
293754009 | Nitrosurea allergy
293755005 | Carmustine allergy
293756006 | Lomustine allergy
293757002 | Triazene antineoplastic allergy
293758007 | Dacarbazine allergy
293759004 | Cytotoxic antibiotic allergy
293760009 | Dactinomycin allergy
293761008 | Bleomycin allergy
293762001 | Mitomycin allergy
293763006 | Plicamycin allergy
293764000 | Aclarubicin allergy
293765004 | Mitoxantrone allergy
293766003 | Doxorubicin allergy
293767007 | Epirubicin allergy
293768002 | Idarubicin allergy
293769005 | Antimetabolite allergy
293770006 | Mercuric oxide allergy
293771005 | Methotrexate allergy
293772003 | Mercaptopurine allergy
293773008 | Tioguanine allergy
293774002 | Pentostatin allergy
293775001 | Cytarabine allergy
293776000 | Fluorouracil allergy
293777009 | Etoposide allergy
293778004 | Amsacrine allergy
293779007 | Carboplatin allergy
293780005 | Cisplatin allergy
293781009 | Hydroxycarbamide allergy
293782002 | Procarbazine allergy
293783007 | Razoxane allergy
293784001 | Crisantaspase allergy
293785000 | Paclitaxel allergy
293786004 | Fludarabine allergy
293787008 | Aminoglutethimide allergy
293788003 | Estrogen antagonist allergy
293789006 | Trilostane allergy
293790002 | Tamoxifen allergy
293791003 | Formestane allergy
293792005 | Vinca alkaloid allergy
293793000 | Vinblastine allergy
293794006 | Vincristine allergy
293795007 | Vindesine allergy
293796008 | Dimethyl sulfoxide allergy
293797004 | Immunosuppressant allergy
293798009 | Ciclosporin allergy
293799001 | Azathioprine allergy
293800002 | Drug groups primarily affecting the central nervous system allergy
293801003 | Centrally acting appetite suppressant allergy
293802005 | Mazindol allergy
293803000 | Phentermine allergy
293804006 | Dexfenfluramine allergy
293805007 | Diethylpropion allergy
293806008 | Fenfluramine allergy
293807004 | Dopaminergic drug used in parkinsonism allergy
293808009 | Levodopa allergy
293809001 | Benserazide + levodopa allergy
293810006 | Carbidopa + levodopa allergy
293811005 | Amantadine allergy
293812003 | Apomorphine allergy
293813008 | Lisuride allergy
293814002 | Pergolide allergy
293815001 | Bromocriptine allergy
293816000 | Antidepressant drug allergy
293817009 | Lithium allergy

293818004 | Lithium carbonate allergy
293819007 | Lithium citrate allergy
293821002 | Tricyclic antidepressant drug allergy
293822009 | Butriptyline allergy
293823004 | Doxepin allergy
293824005 | Iprindole allergy
293825006 | Lofepramine allergy
293826007 | Nortriptyline allergy
293827003 | Trimipramine allergy
293828008 | Amoxapine allergy
293829000 | Amitriptyline allergy
293830005 | Clomipramine allergy
293831009 | Desipramine allergy
293832002 | Dosulepin allergy
293833007 | Imipramine allergy
293834001 | Protriptyline allergy
293835000 | Monoamine oxidase inhibitor allergy
293836004 | Phenelzine allergy
293837008 | Iproniazid allergy
293838003 | Isocarboxazid allergy
293839006 | Tranylcypromine allergy
293840008 | Moclobemide allergy
293841007 | Compound antidepressants allergy
293842000 | Tryptophan allergy
293843005 | Venlafaxine allergy
293844004 | Selective serotonin re-uptake inhibitor allergy
293845003 | Sertraline allergy
293847006 | Paroxetine allergy
293848001 | Nefazodone allergy
293849009 | Citalopram allergy
293850009 | Fluoxetine allergy
293851008 | Fluvoxamine allergy
293852001 | Tetracyclic antidepressant drug allergy
293853006 | Maprotiline allergy
293854000 | Mianserin allergy
293855004 | Trazodone allergy
293856003 | Viloxazine allergy
293857007 | Antiepileptic allergy
293858002 | Beclamide allergy
293859005 | Lamotrigine allergy
293860000 | Piracetam allergy
293861001 | Gabapentin allergy
293862008 | Sodium valproate allergy
293863003 | Barbiturate antiepileptic allergy
293864009 | Methylphenobarbital allergy
293865005 | Phenobarbital allergy
293866006 | Primidone allergy
293867002 | Carbamazepine allergy
293868007 | Vigabatrin allergy
293869004 | Phenytoin allergy
293870003 | Ethosuximide allergy
293871004 | Clonazepam allergy
293873001 | Sedative allergy
293874007 | Zopiclone allergy
293875008 | Zolpidem allergy
293876009 | Chlormezanone allergy
293877000 | Methyprylone allergy
293878005 | Paraldehyde allergy
293879002 | Barbiturate sedative allergy
293880004 | Amobarbital allergy
293881000 | Butobarbital allergy
293882007 | Cyclobarbitone allergy
293883002 | Amobarbital sodium allergy
293884008 | Secobarbital allergy
293885009 | Benzodiazepine sedative allergy
293886005 | Flunitrazepam allergy
293887001 | Flurazepam allergy
293888006 | Loprazolam allergy
293889003 | Lormetazepam allergy
293890007 | Nitrazepam allergy
293891006 | Triazolam allergy
293892004 | Alprazolam allergy

293893009 | Bromazepam allergy
293894003 | Chlordiazepoxide allergy
293895002 | Clobazam allergy
293896001 | Dipotassium clorazepate allergy
293897005 | Ketazolam allergy
293898000 | Medazepam allergy
293899008 | Oxazepam allergy
293900003 | Prazepam allergy
293901004 | Midazolam allergy
293902006 | Diazepam allergy
293903001 | Lorazepam allergy
293904007 | Temazepam allergy
293905008 | Carbamate sedative allergy
293906009 | Meprobamate allergy
293907000 | Chloral sedative allergy
293908005 | Chloral hydrate allergy
293909002 | Dichloralphenazone allergy
293910007 | Triclofos sodium allergy
293911006 | Buspirone allergy
293912004 | Clomethiazole allergy
293913009 | Neuroleptic allergy
293914003 | Sulpiride allergy
293915002 | Loxapine allergy
293916001 | Clozapine allergy
293917005 | Risperidone allergy
293918000 | Tetrabenazine allergy
293919008 | Butyrophenone allergy
293920002 | Benperidol allergy
293921003 | Trifluperidol allergy
293922005 | Haloperidol decanoate allergy
293923000 | Droperidol allergy
293924006 | Haloperidol allergy
293925007 | Diphenylbutylpiperidine allergy
293926008 | Pimozide allergy
293927004 | Fluspirilene allergy
293928009 | Phenothiazine allergy
293929001 | Levomepromazine allergy
293930006 | Pericyazine allergy
293931005 | Fluphenazine decanoate allergy
293932003 | Fluphenazine enanthate allergy
293933008 | Thiethylperazine allergy
293934002 | Fluphenazine allergy
293935001 | Chlorpromazine allergy
293936000 | Pipotiazine allergy
293937009 | Promazine allergy
293938004 | Thioridazine allergy
293939007 | Perphenazine allergy
293940009 | Prochlorperazine allergy
293941008 | Trifluoperazine allergy
293942001 | Thioxanthene allergy
293943006 | Chlorprothixene allergy
293944000 | Flupentixol decanoate allergy
293945004 | Zuclopenthixol decanoate allergy
293946003 | Zuclopenthixol allergy
293948002 | Flupentixol allergy
293949005 | Oxypertine allergy
293950005 | Remoxipride allergy
293951009 | Decarboxylase inhibitor allergy
293952002 | Selegiline allergy
293953007 | Central stimulant allergy
293954001 | Pemoline allergy
293955000 | Methylphenidate allergy
293956004 | Prolintane allergy
293957008 | Amfetamine group allergy
293958003 | Dexamfetamine allergy
293959006 | Alcohol metabolism modifier allergy
293960001 | Disulfiram allergy
293961002 | Drug groups primarily affecting the autonomic nervous system allergy
293962009 | Beta-adrenoceptor blocking drug allergy
293963004 | Cardioselective beta-blocker allergy
293964005 | Acebutolol allergy
293965006 | Atenolol allergy

293966007 | Betaxolol allergy
293967003 | Bisoprolol allergy
293968008 | Celiprolol allergy
293969000 | Esmolol allergy
293970004 | Metoprolol allergy
293971000 | Non-cardioselective beta-blocker allergy
293972007 | Nadolol allergy
293973002 | Pindolol allergy
293974008 | Carvedilol allergy
293975009 | Metipranolol allergy
293976005 | Carteolol allergy
293977001 | Labetalol allergy
293978006 | Levobunolol allergy
293979003 | Oxprenolol allergy
293980000 | Penbutolol allergy
293981001 | Practolol allergy
293982008 | Propranolol allergy
293983003 | Sotalol allergy
293984009 | Timolol allergy
293985005 | Alpha-adrenoceptor blocking drug allergy
293986006 | Alfuzosin allergy
293987002 | Doxazosin allergy
293988007 | Indoramin allergy
293989004 | Phenoxybenzamine allergy
293990008 | Phentolamine allergy
293991007 | Prazosin allergy
293992000 | Terazosin allergy
293993005 | Nicotine allergy
293994004 | Calcium-channel blocker allergy
293995003 | Lidoflazine allergy
293996002 | Nifedipine allergy
293997006 | Prenylamine allergy
293998001 | Isradipine allergy
293999009 | Felodipine allergy
294000006 | Lacidipine allergy
294001005 | Nimodipine allergy
294002003 | Amlodipine allergy
294003008 | Diltiazem allergy
294004002 | Nicardipine allergy
294005001 | Verapamil allergy
294006000 | Parasympathomimetic allergy
294007009 | Pilocarpine allergy
294008004 | Acetylcholine chloride allergy
294009007 | Methacholine allergy
294010002 | Anticholinesterase parasympathomimetic allergy
294011003 | Physostigmine allergy
294012005 | Demecarium allergy
294013000 | Distigmine allergy
294014006 | Ecothiopate allergy
294015007 | Edrophonium allergy
294016008 | Pyridostigmine allergy
294017004 | Neostigmine allergy
294018009 | Bethanechol allergy
294019001 | Carbachol allergy
294020007 | Sympathomimetic allergy
294021006 | Isoetharine hydrochloride allergy
294022004 | Phenylpropanolamine hydrochloride allergy
294023009 | Pseudoephedrine allergy
294024003 | Alpha-adrenoceptor agonist allergy
294025002 | Oxedrine tartrate allergy
294026001 | Metaraminol allergy
294027005 | Methoxamine allergy
294028000 | Naphazoline allergy
294029008 | Norepinephrine allergy
294030003 | Phenylephrine allergy
294031004 | Xylometazoline allergy
294032006 | Beta-adrenoceptor agonist allergy
294033001 | Selective beta-2 adrenoceptor stimulants allergy
294035008 | Pirbuterol allergy
294036009 | Salmeterol allergy
294037000 | Salbutamol allergy
294038005 | Bambuterol allergy

294039002 | Fenoterol allergy
294040000 | Orciprenaline allergy
294041001 | Reproterol allergy
294042008 | Rimiterol allergy
294043003 | Ritodrine allergy
294044009 | Terbutaline allergy
294045005 | Tulobuterol allergy
294046006 | Selective beta-1 adrenoceptor stimulants allergy
294047002 | Dobutamine allergy
294048007 | Dopexamine allergy
294050004 | Isoprenaline allergy
294051000 | Oral isoprenaline allergy
294052007 | Inhaled isoprenaline allergy
294053002 | Parenteral isoprenaline allergy
294054008 | Central alpha-adrenoceptor agonist allergy
294055009 | Methyldopa allergy
294056005 | Methyldopa and diuretic allergy
294057001 | Apraclonidine allergy
294058006 | Clonidine allergy
294059003 | Lofexidine allergy
294060008 | Dipivefrine allergy
294061007 | Dopamine allergy
294062000 | Ephedrine allergy
294063005 | Oxymetazoline allergy
294064004 | Xamoterol allergy
294065003 | Anticholinergic allergy
294066002 | Antimuscarinic allergy
294067006 | Belladonna alkaloids allergy
294068001 | Piperidolate hydrochloride allergy
294069009 | Biperiden allergy
294070005 | Emepronium bromide allergy
294071009 | Terodiline hydrochloride allergy
294072002 | Lachesine chloride allergy
294073007 | Tropicamide allergy
294074001 | Hyoscine allergy
294075000 | Hyoscine hydrobromide allergy
294076004 | Atropine allergy
294077008 | Trihexyphenidyl allergy
294078003 | Benzatropine allergy
294079006 | Cyclopentolate allergy
294080009 | Glycopyrronium allergy
294081008 | Homatropine allergy
294082001 | Ipratropium allergy
294083006 | Methixene allergy
294084000 | Orphenadrine allergy
294085004 | Orphenadrine hydrochloride allergy
294086003 | Orphenadrine citrate allergy
294087007 | Oxitropium allergy
294088002 | Oxybutynin allergy
294089005 | Procyclidine allergy
294090001 | Drug groups primarily affecting the respiratory system allergy
294091002 | Dornase alfa allergy
294092009 | Mucolytics allergy
294093004 | Tyloxapol allergy
294094005 | Bromhexine hydrochloride allergy
294095006 | Carbocisteine allergy
294096007 | Methylcysteine allergy
294097003 | Acetylcysteine allergy
294098008 | Respiratory stimulant allergy
294099000 | Nikethamide allergy
294100008 | Ethamivan allergy
294101007 | Doxapram allergy
294102000 | Respiratory surfactant allergy
294103005 | Beractant allergy
294104004 | Phospholipid fraction allergy
294105003 | Pumactant allergy
294106002 | Colfosceril allergy
294107006 | Antiallergenic drug allergy
294108001 | Topical antihistamine allergy
294109009 | H1 antihistamine allergy
294110004 | Non-sedating antihistamine allergy
294111000 | Astemizole allergy

294112007 | Terfenadine allergy
294113002 | Acrivastine allergy
294114008 | Loratadine allergy
294115009 | Azelastine allergy
294116005 | Cetirizine allergy
294117001 | Sedating antihistamine allergy
294118006 | Clemastine allergy
294119003 | Mebhydrolin allergy
294120009 | Mequitazine allergy
294121008 | Oxatomide allergy
294122001 | Cyclizine allergy
294123006 | Dimenhydrinate allergy
294124000 | Meclozine hydrochloride allergy
294125004 | Antazoline allergy
294126003 | Promethazine allergy
294127007 | Azatadine allergy
294128002 | Brompheniramine allergy
294129005 | Chlorphenamine allergy
294130000 | Cinnarizine allergy
294131001 | Cyproheptadine allergy
294132008 | Dimetindene allergy
294133003 | Diphenhydramine allergy
294134009 | Diphenylpyraline allergy
294135005 | Hydroxyzine allergy
294136006 | Mepyramine allergy
294137002 | Phenindamine allergy
294138007 | Pheniramine allergy
294139004 | Triprolidine allergy
294140002 | Alimemazine allergy
294141003 | Cromoglycate and related antiallergenic allergy
294142005 | Nedocromil allergy
294143000 | Sodium cromoglicate allergy
294144006 | Ketotifen allergy
294145007 | Lodoxamide allergy
294146008 | Cough/decongestant preparation allergy
294147004 | Cough suppressant allergy
294148009 | Isoaminile allergy
294149001 | Morphinan cough suppressant allergy
294151002 | Dextromethorphan hydrobromide allergy
294152009 | Noscapine allergy
294153004 | Pholcodine allergy
294154005 | Compound cough suppressant allergy
294155006 | Proprietary decongestant allergy
294156007 | Expectorant allergy
294157003 | Xanthine allergy
294158008 | Aminophylline allergy
294159000 | Choline theophyllinate allergy
294160005 | Theophylline allergy
294161009 | Bronchodilator preparations allergy
294162002 | Allergy to inhaled corticosteroids
294163007 | Drug groups and agents primarily acting on the skin allergy
294164001 | Topical dermatological preparation allergy
294165000 | Urea creams allergy
294166004 | Emollient bath additives allergy
294167008 | Dusting powders allergy
294168003 | Calamine allergy
294169006 | Coal tar allergy
294170007 | Coal tar scalp preparations allergy
294171006 | Coal tar bath preparations allergy
294172004 | Bufexamac allergy
294173009 | Dithranol allergy
294174003 | Dithranol ointment allergy
294175002 | Dithranol cream allergy
294176001 | Dithranol paste allergy
294177005 | Ichthammol allergy
294178000 | Calcipotriol allergy
294179008 | Sulfur allergy
294180006 | Azelaic acid allergy
294181005 | Bromine complexes allergy
294182003 | Podophyllum resin allergy
294183008 | Podophyllotoxin allergy
294184002 | Sunscreening preparations allergy

294185001 | Camouflaging preparations allergy
294187009 | Desloughing agents allergy
294188004 | Aserbine allergy
294189007 | Surgical tissue adhesive allergy
294190003 | Enbucrilate allergy
294191004 | Collodion allergy
294192006 | Counter irritants allergy
294193001 | Emollient allergy
294194007 | Poultices allergy
294195008 | Anionic surfactant allergy
294196009 | Alkali metal soap allergy
294197000 | Astringent allergy
294198005 | Aluminum astringent allergy
294199002 | Topical selenium allergy
294200004 | Crotamiton allergy
294201000 | Topical salicylic acid allergy
294202007 | Topical abrasive agent allergy
294203002 | Benzoyl peroxide allergy
294204008 | Silver nitrate allergy
294206005 | Gamolenic acid allergy
294207001 | Retinoid allergy
294208006 | Etretinate allergy
294209003 | Acitretin allergy
294210008 | Tretinoin allergy
294211007 | Isotretinoin allergy
294212000 | Drug groups primarily affecting the musculoskeletal system allergy
294213005 | Drug for the treatment of gout allergy
294214004 | Colchicum alkaloid allergy
294215003 | Colchicine allergy
294216002 | Uricosuric agent allergy
294217006 | Probenecid allergy
294218001 | Sulfinpyrazone allergy
294219009 | Xanthine oxidase inhibitor allergy
294220003 | Allopurinol allergy
294221004 | Skeletal muscle relaxant allergy
294222006 | Neuromuscular transmission drug allergy
294223001 | Depolarizing muscle relaxant allergy
294224007 | Suxamethonium allergy
294225008 | Non-depolarizing muscle relaxant allergy
294226009 | Mivacurium allergy
294227000 | Alcuronium allergy
294228005 | Atracurium allergy
294229002 | Gallamine allergy
294230007 | Pancuronium allergy
294231006 | Tubocurarine allergy
294232004 | Vecuronium allergy
294233009 | Rocuronium allergy
294234003 | Baclofen allergy
294235002 | Carisoprodol allergy
294236001 | Methocarbamol allergy
294237005 | Dantrolene allergy
294238000 | Gold allergy
294239008 | Sodium aurothiomalate allergy
294240005 | Auranofin allergy
294241009 | Smooth muscle relaxant allergy
294242002 | Papaverine allergy
294243007 | Flavoxate allergy
294244001 | Drug groups primarily used in obstetrics, gynecological and urinary tract disorders allergy
294245000 | Mifepristone allergy
294246004 | Non-ionic surfactant allergy
294247008 | Nonoxinol allergy
294248003 | Octoxinol allergy
294249006 | Prostaglandin allergy
294250006 | A series prostaglandin allergy
294252003 | E series prostaglandin allergy
294253008 | Dinoprostone allergy
294254002 | Gemeprost allergy
294255001 | Alprostadil allergy
294256000 | F series prostaglandin allergy
294257009 | Dinoprost allergy
294258004 | Carboprost allergy
294259007 | I series prostaglandin allergy

294260002 | Epoprostenol allergy
294261003 | Terpenes allergy
294262005 | Chelating agents and antidotes allergy
294263000 | Antidote allergy
294264006 | Ipecacuanha allergy
294265007 | Charcoal-activated allergy
294266008 | Sodium nitrite allergy
294268009 | Digoxin-specific-antibody allergy
294269001 | Mesna allergy
294270000 | Benzodiazepine antagonist allergy
294271001 | Flumazenil allergy
294272008 | Cholinesterase reactivator allergy
294273003 | Pralidoxime allergy
294274009 | Hydrofluoric acid burn antidote allergy
294275005 | Opioid antagonist allergy
294276006 | Naltrexone allergy
294277002 | Naloxone allergy
294278007 | Protamine allergy
294279004 | Antidotes for pesticides allergy
294280001 | Fullers earth powder allergy
294281002 | Bentonite powder allergy
294282009 | Chelating agent allergy
294283004 | Dimercaprol allergy
294284005 | Desferrioxamine allergy
294285006 | Edetate allergy
294286007 | Dicobalt edetate allergy
294287003 | Sodium calcium edetate allergy
294288008 | Trisodium edetate allergy
294289000 | Disodium edetate allergy
294290009 | Trientine allergy
294291008 | Penicillamine allergy
294293006 | Homeopathic medicine allergy
294295004 | Herbal medicine allergy
294297007 | Sodium hyaluronate allergy
294298002 | Glycine allergy
294299005 | Dialysis fluid allergy
294300002 | Peritoneal dialysis solution allergy
294301003 | Hemodialysis fluid allergy
294302005 | Hemofiltration solution allergy
294304006 | Cardioplegia solution allergy
294306008 | Dimethyl-ether propane allergy
294307004 | Citrate allergy
294308009 | Hemorrhoid preparation allergy
294309001 | Ear wax removal preparation allergy
294310006 | Topical potassium allergy
294311005 | Topical zinc allergy
294312003 | Oropharyngeal preparations allergy
294313008 | Artificial saliva allergy
294315001 | Fixed oil allergy
294316000 | Olive oil allergy
294317009 | Arachis oil allergy
294318004 | Castor oil allergy
294319007 | Glycerol/glycol/macrogol allergy
294320001 | Glycerol allergy
294321002 | Lubricant allergy
294322009 | Base allergy
294324005 | Paraffin allergy
294325006 | Paraffin-yellow soft allergy
294326007 | Paraffin-white soft allergy
294327003 | Liquid paraffin allergy
294328008 | Silicone allergy
294329000 | Dimethicone allergy
294330005 | Wool alcohol allergy
294331009 | Viscosity modifier allergy
294332002 | Polyvinyl alcohol allergy
294333007 | Carbomer-940 allergy
294334001 | Cellulose-derived viscosity modifier allergy
294335000 | Hypromellose allergy
294336004 | Hypromellose eye drops allergy
294337008 | Hydroxyethylcellulose allergy
294339006 | Carmellose allergy
294340008 | Antifungal drug allergy

294341007 | Flucytosine allergy
294342000 | Terbinafine allergy
294343005 | Nitrophenol allergy
294344004 | Tolnaftate allergy
294346002 | Amorolfine allergy
294348001 | Griseofulvin allergy
294349009 | Amphotericin allergy
294350009 | Natamycin allergy
294351008 | Nystatin allergy
294352001 | Azole antifungal allergy
294354000 | Undecenoates allergy
294355004 | Imidazole antifungal allergy
294356003 | Clotrimazole allergy
294357007 | Fenticonazole allergy
294358002 | Tioconazole allergy
294359005 | Econazole allergy
294360000 | Isoconazole allergy
294361001 | Sulconazole allergy
294362008 | Ketoconazole allergy
294363003 | Miconazole allergy
294364009 | Triazole antifungals allergy
294365005 | Fluconazole allergy
294366006 | Itraconazole allergy
294367002 | Antiviral drug allergy
294368007 | Inosine pranobex allergy
294369004 | Zidovudine allergy
294370003 | Ganciclovir allergy
294371004 | Famciclovir allergy
294372006 | Didanosine allergy
294373001 | Zalcitabine allergy
294374007 | Valaciclovir allergy
294375008 | Interferons allergy
294376009 | Human interferon gamma-1b allergy
294377000 | Interferon-A-2a allergy
294378005 | Interferon-A-2b allergy
294379002 | Interferon-A-N1 allergy
294380004 | Ribavirin allergy
294381000 | Trifluorothymidine allergy
294382007 | Foscarnet allergy
294383002 | Vidarabine allergy
294384008 | Aciclovir allergy
294385009 | Idoxuridine allergy
294386005 | Idoxuridine in dimethylsulfoxide allergy
294387001 | Antimalarial drug allergy
294388006 | Pyrimethamine allergy
294389003 | Aminoquinoline antimalarial allergy
294390007 | Amodiaquine allergy
294391006 | Primaquine allergy
294392004 | Mefloquine allergy
294393009 | Hydroxychloroquine allergy
294394003 | Chloroquine allergy
294395002 | Biguanide antimalarial allergy
294396001 | Proguanil allergy
294397005 | Cinchona antimalarial allergy
294398000 | Quinine allergy
294399008 | Halofantrine allergy
294400001 | Mepacrine allergy
294401002 | Lactobacill acidophil vaccine allergy
294402009 | Anti-infective nasal preparations allergy
294403004 | Disinfectants and cleansers allergy
294404005 | Acetic acid allergy
294405006 | Hydrargaphen allergy
294406007 | Polynoxylin allergy
294407003 | Hexetidine allergy
294408008 | Sodium perborate allergy
294409000 | Chlorinated solutions allergy
294410005 | Potassium permanganate allergy
294411009 | Bismuth subnitrate and iodoform paste impregnated gauze allergy
294412002 | Phenolics allergy
294413007 | Thymol allergy
294414001 | Chlorinated phenol disinfectant allergy
294415000 | Chloroxylenol allergy

294416004 | Hexachlorophene allergy
294417008 | Triclosan allergy
294418003 | Phenol allergy
294419006 | Alcoholic disinfectant allergy
294420000 | Alcohol products allergy
294421001 | Industrial methylated spirit allergy
294422008 | Aldehyde disinfectant allergy
294423003 | Glutaraldehyde allergy
294425005 | Noxythiolin allergy
294426006 | Formaldehyde allergy
294427002 | Amidine disinfectant allergy
294428007 | Propamidine isethionate allergy
294429004 | Dibrompropamidine isethionate allergy
294430009 | Biguanide disinfectant allergy
294431008 | Chlorhexidine allergy
294432001 | Chlorhexidine hydrochloride and neomycin sulfate allergy
294433006 | Borate allergy
294434000 | Boric acid allergy
294435004 | Cationic surfactant allergy
294436003 | Quaternary ammonium surfactant allergy
294437007 | Cetrimide allergy
294438002 | Benzalkonium allergy
294439005 | Domiphen allergy
294440007 | Quaternary pyridinium surfactant allergy
294441006 | Cetylpyridinium allergy
294442004 | Quaternary quinolinium surfactant allergy
294443009 | Dequalinium allergy
294444003 | Disinfectant dye allergy
294445002 | Acridine azo disinfectant dye allergy
294446001 | Triphenylmethane azo disinfectant dye allergy
294447005 | Crystal violet allergy
294448000 | Brilliant green allergy
294449008 | Hydrogen peroxide allergy
294450008 | Anthelmintics allergy
294451007 | Piperazine allergy
294452000 | Pyrantel allergy
294453005 | Niclosamide allergy
294455003 | Bephenium allergy
294456002 | Diethylcarbamazine allergy
294457006 | Benzimidazole anthelmintic allergy
294458001 | Mebendazole allergy
294459009 | Albendazole allergy
294460004 | Tiabendazole allergy
294461000 | Antibacterial drug allergy
294462007 | Aminoglycosides allergy
294463002 | Amikacin allergy
294464008 | Kanamycin allergy
294465009 | Netilmicin allergy
294466005 | Streptomycin allergy
294467001 | Framycetin allergy
294468006 | Neomycin allergy
294469003 | Gentamicin allergy
294470002 | Tobramycin allergy
294471003 | Clarithromycin allergy
294472005 | Azithromycin allergy
294473000 | Sodium fusidate allergy
294474006 | Spectinomycin allergy
294475007 | Vancomycin allergy
294476008 | Teicoplanin allergy
294477004 | Trimethoprim allergy
294478009 | Nitrofurantoin allergy
294479001 | Methenamine hippurate allergy
294480003 | Mupirocin allergy
294481004 | Nitrofurazone allergy
294482006 | Fusidic acid allergy
294483001 | 4-quinolones allergy
294484007 | Acrosoxacin allergy
294485008 | Cinoxacin allergy
294486009 | Nalidixic acid allergy
294487000 | Ciprofloxacin allergy
294488005 | Enoxacin allergy
294489002 | Ofloxacin allergy

294490006 | Norfloxacin allergy
294491005 | Temafloxacin allergy
294492003 | Penicillinase-sensitive penicillins allergy
294494002 | Benethamine penicillin allergy
294495001 | Benzathine penicillin allergy
294496000 | Phenethicillin allergy
294497009 | Phenoxymethylpenicillin allergy
294498004 | Procaine benzylpenicillin allergy
294499007 | Benzylpenicillin allergy
294500003 | Penicillinase-resistant penicillins allergy
294501004 | Cloxacillin allergy
294502006 | Flucloxacillin allergy
294503001 | Methicillin allergy
294504007 | Broad spectrum penicillins allergy
294505008 | Amoxicillin allergy
294506009 | Ampicillin allergy
294507000 | Ciclacillin allergy
294508005 | Mezlocillin allergy
294509002 | Pivampicillin allergy
294510007 | Carbenicillin allergy
294511006 | Bacampicillin allergy
294512004 | Talampicillin allergy
294513009 | Antipseudomonal penicillins allergy
294514003 | Temocillin allergy
294515002 | Piperacillin allergy
294516001 | Azlocillin allergy
294517005 | Ticarcillin allergy
294518000 | Carfecillin allergy
294519008 | Mecillinam allergy
294520002 | Pivmecillinam allergy
294521003 | Combined penicillin preparation allergy
294522005 | Ampicillin and cloxacillin allergy
294523000 | Amoxicillin + clavulanate potassium allergy
294524006 | Ampicillin + floxacillin allergy
294525007 | Piperacillin and tazobactam allergy
294526008 | Pivampicillin and pivmecillinam allergy
294527004 | Ticarcillin and clavulanic acid allergy
294528009 | Polymyxins allergy
294529001 | Colistin allergy
294530006 | Polymyxin B allergy
294531005 | Carbapenem allergy
294532003 | Cephalosporin allergy
294533008 | First generation cephalosporin allergy
294534002 | Cefadroxil allergy
294535001 | Cefalexin allergy
294536000 | Cephalothin allergy
294537009 | Cefazolin allergy
294538004 | Cefradine allergy
294539007 | Latamoxef allergy
294540009 | Second generation cephalosporin allergy
294541008 | Cefaclor allergy
294542001 | Cefuroxime allergy
294543006 | Cephamandole allergy
294544000 | Third generation cephalosporin allergy
294545004 | Cefotaxime allergy
294546003 | Ceftazidime allergy
294547007 | Ceftizoxime allergy
294548002 | Cefixime allergy
294549005 | Cefodizime allergy
294550005 | Cefpodoxime allergy
294551009 | Ceftriaxone allergy
294552002 | Ceftibuten allergy
294554001 | Cefsulodin allergy
294555000 | Fourth generation cephalosporin allergy
294556004 | Cefpirome allergy
294557008 | Cephamycin allergy
294558003 | Cefoxitin allergy
294559006 | Fosfomycin allergy
294560001 | Lincomycin and derivatives allergy
294561002 | Clindamycin allergy
294562009 | Lincomycin allergy
294563004 | Mandelic acid allergy

294564005 | Monobactam allergy
294565006 | Aztreonam allergy
294566007 | Nitroimidazole allergy
294567003 | Metronidazole allergy
294568008 | Tinidazole allergy
294569000 | Nimorazole allergy
294570004 | Calcium sulfaloxate allergy
294571000 | Phthalylsulfathiazole allergy
294572007 | Sulfametopyrazine allergy
294573002 | Sulfadiazine allergy
294574008 | Sulfadimethoxine allergy
294575009 | Sulfadimidine allergy
294576005 | Sulfafurazole allergy
294577001 | Sulfaguanidine allergy
294578006 | Sulfaurea allergy
294579003 | Mafenide allergy
294580000 | Silver sulfadiazine allergy
294581001 | Combined sulfonamides allergy
5388008 | Congenital lactase deficiency
25744000 | Hereditary gastrogenic lactose intolerance
29512005 | Lactase deficiency in diseases other than of the small intestine
38032004 | Ontogenic late onset lactase deficiency
54250004 | Lactose intolerance in children without lactase deficiency
60414003 | Acquired lactase deficiency
190751001 | Primary lactose intolerance
52070001 | Acquired monosaccharide malabsorption
237978005 | Glycerol intolerance
700094005 | Milk intolerance
409137002 | No known history of drug allergy
428197003 | No known insect allergy
428607008 | No known environmental allergy
429625007 | No known food allergy
716184000 | No known latex allergy
716220001 | No known animal allergy

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.631 Value Set http://hl7.org/fhir/ValueSet/procedure-progress-status-codes

URL: https://hl7.org/fhir/R4/valueset-procedure-progress-status-codes.html
Patient Care Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/procedure-progress-status-codes
Version: | 4.0.1
Name: | ProcedureProgressStatusCodes
Title: | Procedure Progress Status Codes
Definition: | This value set is provided as an example. The value set to instantiate this attribute should be drawn from a robust terminology code system that consists of or contains concepts to support the procedure performance process.
Committee: | Patient Care Work Group
OID: | 2.16.840.1.113883.4.642.3.946 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ProcedureProgressStatusCodes

• Extension: http://hl7.org/fhir/StructureDefinition/procedure-progressStatus: progressStatus (CodeableConcept / Example)

4.4.1.631.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/procedure-progress-status-code

4.4.1.631.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 6 concepts

Expansion based on http://hl7.org/fhir/procedure-progress-status-code version 4.0.1

All codes from system http://hl7.org/fhir/procedure-progress-status-code

Code | Display | Definition
in-operating-room | In Operating Room | A patient is in the Operating Room.
prepared | Prepared | The patient is prepared for a procedure.
anesthesia-induced | Anesthesia Induced | The patient is under anesthesia.
open-incision | Open Incision | The patient has open incision(s).
closed-incision | Closed Incision | The patient has incision(s) closed.
in-recovery-room | In Recovery Room | The patient is in the recovery room.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.632 Value Set http://hl7.org/fhir/ValueSet/measure-report-status

URL: https://hl7.org/fhir/R4/valueset-measure-report-status.html
Clinical Quality Information Work Group | Maturity Level: 2 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/measure-report-status
Version: | 4.0.1
Name: | MeasureReportStatus
Title: | MeasureReportStatus
Definition: | The status of the measure report.
Committee: | Clinical Quality Information Work Group
OID: | 2.16.840.1.113883.4.642.3.776 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for MeasureReportStatus
• ConceptMap: Translation source in Canonical Mapping for "MeasureReportStatus"
• Resource: MeasureReport.status (code / Required)

4.4.1.632.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/measure-report-status

4.4.1.632.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 3 concepts

Expansion based on http://hl7.org/fhir/measure-report-status version 4.0.1

All codes from system http://hl7.org/fhir/measure-report-status

Code | Display | Definition | Canonical Status
complete | Complete | The report is complete and ready for use. | ~complete
pending | Pending | The report is currently being generated. | ~active
error | Error | An error occurred attempting to generate the report. | ~failed

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.633 Value Set http://hl7.org/fhir/ValueSet/entformula-type

URL: https://hl7.org/fhir/R4/valueset-entformula-type.html
Orders and Observations Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/entformula-type
Version: | 4.0.1
Name: | EnteralFormulaTypeCodes
Title: | Enteral Formula Type Codes
Definition: | EnteralFormulaType : Codes for type of enteral formula to be administered to patient. This value set is composed of SNOMED CT (US Extension) Concepts from SCTID 470581016 (Enteral+supplement feeds hierarchy (product)) and is provided as a suggestive example.
Committee: | Orders and Observations Work Group
OID: | 2.16.840.1.113883.4.642.3.391 (for OID based terminology systems)
Copyright: | This resource includes content from SNOMED Clinical Terms® (SNOMED CT®) which is copyright of the International Health Terminology Standards Development Organisation (IHTSDO). Implementers of these specifications must have the appropriate SNOMED CT Affiliate license - for more information contact http://www.snomed.org/snomed-ct/get-snomed-ct or info@snomed.org
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: NutritionOrder.enteralFormula.baseFormulaType (CodeableConcept / Example)

4.4.1.633.1 Content Logical Definition

This value set includes codes from the following code systems:

http://snomed.info/sct

Code	Display
443031000124106	Adult critical care formula
443051000124104	Adult diabetes specialty formula
442911000124109	Adult elemental formula
443021000124108	Adult hepatic specialty formula
442971000124100	Adult high energy formula
442981000124102	Adult hydrolyzed protein formula
442991000124104	Adult high protein formula
443011000124100	Adult high protein high fiber formula
442961000124107	Adult low carbohydrate formula
442951000124105	Adult pulmonary specialty formula
442941000124108	Adult renal specialty formula
442921000124101	Adult standard formula
442931000124103	Adult soy protein isolate formula
443361000124100	Pediatric Formula
443401000124105	Pediatric elemental formula
443491000124103	Pediatric high energy formula
443501000124106	Pediatric high energy formula with increased fiber
443421000124100	Pediatric hydrolyzed protein formula
443471000124104	Pediatric increased fiber formula
444431000124104	Pediatric reduced energy formula
443451000124109	Pediatric standard formula
441561000124106	Standard enteral formula with fiber
443461000124106	Standard Formula
441531000124102	Standard Enteral Formula
443561000124107	Soy based formula
443481000124101	Renal Formula
441571000124104	High energy enteral formula with fiber
441591000124103	Diabetic enteral formula with fiber

441601000124106 | Diabetic high calorie high protein enteral formula with fiber
443351000124102 | Increased fiber formula
443771000124106 | Hydrolyzed protein formula
441671000124100 | Hydrolyzed peptide-based high protein enteral formula
443111000124101 | High protein formula
443431000124102 | High Energy Formula
443411000124108 | Elemental Formula
442651000124102 | Adult formula

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.634 Value Set http://hl7.org/fhir/ValueSet/medicationrequest-category

URL: https://hl7.org/fhir/R4/valueset-medicationrequest-category.html
Pharmacy Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/medicationrequest-category
Version: | 4.0.1
Name: | medicationRequest Category Codes
Title: | Medication request category codes
Definition: | MedicationRequest Category Codes
Committee: | Pharmacy Work Group
OID: | 2.16.840.1.113883.4.642.3.1322 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for medicationRequest Category Codes

• Resource: MedicationRequest.category (CodeableConcept / Example)

4.4.1.634.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/medicationrequest-category

4.4.1.634.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 4 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/medicationrequest-category version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/medicationrequest-category

Code | Display | Definition
inpatient | Inpatient | Includes requests for medications to be administered or consumed in an inpatient or acute care setting
outpatient | Outpatient | Includes requests for medications to be administered or consumed in an outpatient setting (for example, Emergency Department, Outpatient Clinic, Outpatient Surgery, Doctor's office)
community | Community | Includes requests for medications to be administered or consumed by the patient in their home (this would include long term care or nursing homes, hospices, etc.)
discharge | Discharge | Includes requests for medications created when the patient is being released from a facility

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application

Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.635 Value Set http://hl7.org/fhir/ValueSet/adjudication

URL: https://hl7.org/fhir/R4/valueset-adjudication.html
Financial Management Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/adjudication
Version: | 4.0.1
Name: | AdjudicationValueCodes
Title: | Adjudication Value Codes
Definition: | This value set includes a smattering of Adjudication Value codes which includes codes to indicate the amounts eligible under the plan, the amount of benefit, copays etc.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.596 (for OID based terminology systems)
Copyright: | This is an example set.
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for AdjudicationValueCodes
• Resource: ExplanationOfBenefit.item.adjudication.category (CodeableConcept / Example)
• Resource: ExplanationOfBenefit.total.category (CodeableConcept / Example)
• Resource: ClaimResponse.item.adjudication.category (CodeableConcept / Example)
• Resource: ClaimResponse.total.category (CodeableConcept / Example)

4.4.1.635.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/adjudication

4.4.1.635.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 8 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/adjudication version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/adjudication

Code | Display | Definition
submitted | Submitted Amount | The total submitted amount for the claim or group or line item.
copay | CoPay | Patient Co-Payment
eligible | Eligible Amount | Amount of the change which is considered for adjudication.
deductible | Deductible | Amount deducted from the eligible amount prior to adjudication.
unallocdeduct | Unallocated Deductible | The amount of deductible which could not allocated to other line items.
eligpercent | Eligible % | Eligible Percentage.
tax | Tax | The amount of tax.
benefit | Benefit Amount | Amount payable under the coverage

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.636 Value Set http://hl7.org/fhir/ValueSet/contract-action

URL: https://hl7.org/fhir/R4/valueset-contract-action.html
Financial Management Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/contract-action
Version: | 4.0.1
Name: | ContractActionCodes
Title: | Contract Action Codes
Definition: | This value set includes sample Contract Action codes.
Committee: | Financial Management Work Group
OID: | 2.16.840.1.113883.4.642.3.739 (for OID based terminology systems)
Copyright: | This is an example set.
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ContractActionCodes

• Resource: Contract.term.action.type (CodeableConcept / Example)

4.4.1.636.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/contractaction

4.4.1.636.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 2 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/contractaction version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/contractaction

Code | Display | Definition
action-a | Action A | Definition for Action A
action-b | Action B | Definition for Action B

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.637 Value Set http://hl7.org/fhir/ValueSet/ucum-bodylength

URL: https://hl7.org/fhir/R4/valueset-ucum-bodylength.html
Orders and Observations Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/ucum-bodylength
Version: | 4.0.1
Name: | BodyLengthUnits
Title: | Body Length Units
Definition: | UCUM units for recording body length measures such as height and head circumference
Committee: | Orders and Observations Work Group
OID: | 2.16.840.1.113883.4.642.3.958 (for OID based terminology systems)
Copyright: | UCUM is Copyright © 1999-2013 Regenstrief Institute, Inc. and The UCUM Organization, Indianapolis, IN. All rights reserved. See http://unitsofmeasure.org/trac//wiki/TermsOfUse for details
Source Resource | XML / JSON

This value set is not currently used

4.4.1.637.1 Content Logical Definition

This value set includes codes from the following code systems:

http://unitsofmeasure.org

Code | Display
cm | cm
[in_i] | in_i

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.638 Value Set http://hl7.org/fhir/ValueSet/specimen-processing-procedure

URL: https://hl7.org/fhir/R4/valueset-specimen-processing-procedure.html
Orders and Observations Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/specimen-processing-procedure
Version: | 4.0.1
Name: | SpecimenProcessingProcedure
Title: | Specimen processing procedure
Definition: | The technique that is used to perform the process or preserve the specimen.
Committee: | Orders and Observations Work Group
OID: | 2.16.840.1.113883.4.642.3.469 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: Specimen.processing.procedure (CodeableConcept / Example)

4.4.1.638.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/v2-0373

4.4.1.638.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 8 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/v2-0373 version 2.9

All codes from system http://terminology.hl7.org/CodeSystem/v2-0373

Code | Display | Definition
ACID | Acidification
ALK | Alkalization
DEFB | Defibrination
FILT | Filtration
LDLP | LDL Precipitation
NEUT | Neutralization
RECA | Recalification
UFIL | Ultrafiltration

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.639 Value Set http://hl7.org/fhir/ValueSet/copy-number-event

URL: https://hl7.org/fhir/R4/valueset-copy-number-event.html
Clinical Genomics Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/copy-number-event
Version: | 4.0.1
Name: | CopyNumberEvent
Title: | CopyNumberEvent
Definition: | Copy Number Event.
Committee: | Clinical Genomics Work Group
OID: | 2.16.840.1.113883.4.642.3.226 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for CopyNumberEvent

4.4.1.639.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/copy-number-event

4.4.1.639.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 3 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/copy-number-event version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/copy-number-event

Code | Display | Definition
amp | amplification | amplification.
del | deletion | deletion.
lof | loss of function | loss of function.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.640 Value Set http://hl7.org/fhir/ValueSet/goal-priority

URL: https://hl7.org/fhir/R4/valueset-goal-priority.html
Patient Care Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined at http://www.hl7.org/Special/committees/patientcare/ .

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/goal-priority
Version: | 4.0.1
Name: | GoalPriority
Title: | Goal priority
Definition: | Indicates the level of importance associated with reaching or sustaining a goal.
Committee: | Patient Care Work Group
OID: | 2.16.840.1.113883.4.642.3.273 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for GoalPriority
• Resource: Goal.priority (CodeableConcept / Preferred)
• Resource: PlanDefinition.goal.priority (CodeableConcept / Preferred)
• Profile: ShareablePlanDefinition: PlanDefinition.goal.priority (CodeableConcept / Preferred)
• Profile: ComputablePlanDefinition: PlanDefinition.goal.priority (CodeableConcept / Preferred)
• Profile: CDSHooksServicePlanDefinition: PlanDefinition.goal.priority (CodeableConcept / Preferred)
• Extension: http://hl7.org/fhir/StructureDefinition/goal-acceptance: acceptance (CodeableConcept / Example)

4.4.1.640.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/goal-priority

4.4.1.640.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 3 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/goal-priority version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/goal-priority

Code | Display | Definition
high-priority | High Priority | Indicates that the goal is of considerable importance and should be a primary focus of care delivery.
medium-priority | Medium Priority | Indicates that the goal has a reasonable degree of importance and that concrete action should be taken towards the goal. Attainment is not as critical as high-priority goals.
low-priority | Low Priority | The goal is desirable but is not sufficiently important to devote significant resources to. Achievement of the goal may be sought when incidental to achieving other goals.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.641 Value Set http://hl7.org/fhir/ValueSet/endpoint-status

URL: https://hl7.org/fhir/R4/valueset-endpoint-status.html
Patient Administration Work Group | Maturity Level: 2 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/endpoint-status
Version: | 4.0.1
Name: | EndpointStatus
Title: | EndpointStatus
Definition: | The status of the endpoint.
Committee: | Patient Administration Work Group
OID: | 2.16.840.1.113883.4.642.3.494 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for EndpointStatus

• ConceptMap: Translation source in Canonical Mapping for "EndpointStatus"

• Resource: Endpoint.status (code / Required)

4.4.1.641.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/endpoint-status

4.4.1.641.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 6 concepts

Expansion based on http://hl7.org/fhir/endpoint-status version 4.0.1

All codes from system http://hl7.org/fhir/endpoint-status

Code | Display | Definition | Canonical Status
active | Active | This endpoint is expected to be active and can be used. | ~active
suspended | Suspended | This endpoint is temporarily unavailable. | ~suspended
error | Error | This endpoint has exceeded connectivity thresholds and is considered in an error state and should no longer be attempted to connect to until corrective action is taken. | ~failed
off | Off | This endpoint is no longer to be used. | ~inactive
entered-in-error | Entered in error | This instance should not have been part of this patient's medical record. | ~error
test | Test | This endpoint is not intended for production usage. | ~draft

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.642 Value Set http://hl7.org/fhir/ValueSet/variants

URL: https://hl7.org/fhir/R4/valueset-variants.html

This value set could not be expanded by the publication tooling: Error from server: Unable to provide support for code system http://varnomen.hgvs.org/

Orders and Observations Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/variants
Version: | 4.0.1
Name: | HGVS
Title: | H g v s
Definition: | HGVS : Human Genome Variation Society
Committee: | Orders and Observations Work Group
OID: | 2.16.840.1.113883.6.282 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is not currently used

4.4.1.642.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://varnomen.hgvs.org/

4.4.1.642.2 Expansion

This expansion generated 01 Nov 2019

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.643 Value Set http://hl7.org/fhir/ValueSet/prepare-patient-prior-specimen-collection

URL: https://hl7.org/fhir/R4/valueset-prepare-patient-prior-specimen-collection.html
Orders and Observations Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/prepare-patient-prior-specimen-collection
Version: | 4.0.1
Name: | PreparePatient
Title: | Patient preparation prior specimen collection
Definition: | Checks on the patient prior specimen collection. All SNOMED CT concepts descendants of 703763000 |Precondition value (qualifier value)|
Committee: | Orders and Observations Work Group
OID: | 2.16.840.1.113883.4.642.3.1399 (for OID based terminology systems)
Copyright: | This resource includes content from SNOMED Clinical Terms® (SNOMED CT®) which is copyright of the International Health Terminology Standards Development Organisation (IHTSDO). Implementers of these specifications must have the

appropriate SNOMED CT Affiliate license - for more information contact http://www.snomed.org/snomed-ct/get-snomed-ct or info@snomed.org
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: SpecimenDefinition.patientPreparation (CodeableConcept / Example)

4.4.1.643.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include codes from http://snomed.info/sct where concept is-a 703763000 (Precondition value)

4.4.1.643.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 45 concepts

All codes from system http://snomed.info/sct

Code	Display	Definition
703760002	1H post 50 g glucose PO	
703761003	2H post 75 g glucose PO	
703762005	1H post 100 g glucose PO	
703764006	At time of peak level	
703765007	At time of trough level	
703766008	1H post 75 g glucose PO	
703767004	One hour after a meal	
703768009	Two hours after a meal	
703769001	1H post dose glucose	
703770000	Two hours after glucose dose	
703771001	pre 100 g glucose PO	
703772008	2H post 100 g glucose PO	
703773003	Three hours after glucose dose	
703774009	3H post 100 g glucose PO	
703775005	30M post 75 g glucose PO	
707707000	Thirty minutes after resting	
707708005	One hour after resting	
707709002	Ninety minutes after resting	
707710007	Twenty minutes after incubation	
707712004	Thirty minutes after incubation	
707714003	One hour after incubation	
707715002	Two hours after incubation	
707716001	Three hours after incubation	
707717005	Four hours after incubation	
707718000	One day after incubation	
707719008	Seven days after incubation	
707720002	Two hours after peritoneal dialysis	
707721003	Four hours after peritoneal dialysis	
707722005	Twelve hours after peritoneal dialysis	
707723000	Twenty four hours after peritoneal dialysis	
707757008	30M post dose corticotropin	
707758003	1H post dose corticotropin	
707759006	15M pre dose ornithine	
707760001	After urea dose	
707761002	Pre contrast	
707762009	Pre washing	
707763004	After precipitation	
707764005	Post transfusion reaction	
707765006	On ventilator	
707815009	Immediately after addition of normal plasma	
707816005	Immediately after addition of protamine sulfate	
707817001	On menstrual cycle day 10	
707818006	On menstrual cycle day 11	
707819003	On menstrual cycle day 21	
707820009	On menstrual cycle day 3	

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')

Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.644 Value Set http://hl7.org/fhir/ValueSet/address-use

URL: https://hl7.org/fhir/R4/valueset-address-use.html
FHIR Infrastructure Work Group | Maturity Level: N | Normative (from v4.0.0) | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/address-use
Version: | 4.0.1
Name: | AddressUse
Title: | AddressUse
Definition: | The use of an address.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.67 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for AddressUse
• ConceptMap: Translation source in v3 map for AddressUse
• ConceptMap: Translation source in v2 map for AddressUse
• Resource: Address.use (code / Required)

4.4.1.644.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/address-use

4.4.1.644.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 5 concepts

Expansion based on http://hl7.org/fhir/address-use version 4.0.1

All codes from system http://hl7.org/fhir/address-use

Code | Display | Definition
home | Home | A communication address at a home.
work | Work | An office address. First choice for business related contacts during business hours.
temp | Temporary | A temporary address. The period can provide more detailed information.
old | Old / Incorrect | This address is no longer in use (or was never correct but retained for records).
billing | Billing | An address to be used to send bills, invoices, receipts etc.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.645 Value Set http://hl7.org/fhir/ValueSet/genenames

URL: https://hl7.org/fhir/R4/valueset-genenames.html

This value set could not be expanded by the publication tooling: Error from server: Unable to provide support for code system http://www.genenames.org

Orders and Observations Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/genenames
Version: | 4.0.1
Name: | HGNC

Title: | H g n c
Definition: | HGNC: Human Gene Nomenclature Committee
Committee: | Orders and Observations Work Group
OID: | 2.16.840.1.113883.6.281 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• Extension: http://hl7.org/fhir/StructureDefinition/observation-geneticsGene: Gene (CodeableConcept / Preferred)

4.4.1.645.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://www.genenames.org

4.4.1.645.2 Expansion

This expansion generated 01 Nov 2019

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.646 Value Set http://hl7.org/fhir/ValueSet/effect-estimate-type

URL: https://hl7.org/fhir/R4/valueset-effect-estimate-type.html
Clinical Decision Support Work Group | Maturity Level: 0 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/effect-estimate-type
Version: | 4.0.1
Name: | EffectEstimateType
Title: | EffectEstimateType
Definition: | Whether the effect estimate is an absolute effect estimate (absolute difference) or a relative effect estimate (relative difference), and the specific type of effect estimate (eg relative risk or median difference).
Committee: | Clinical Decision Support Work Group
OID: | 2.16.840.1.113883.4.642.3.1355 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for EffectEstimateType

• Resource: EffectEvidenceSynthesis.effectEstimate.type (CodeableConcept / Extensible)

4.4.1.646.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/effect-estimate-type

4.4.1.646.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 7 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/effect-estimate-type version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/effect-estimate-type

Code | Display | Definition
relative-RR | relative risk | relative risk (a type of relative effect estimate).
relative-OR | odds ratio | odds ratio (a type of relative effect estimate).
relative-HR | hazard ratio | hazard ratio (a type of relative effect estimate).
absolute-ARD | absolute risk difference | absolute risk difference (a type of absolute effect estimate).
absolute-MeanDiff | mean difference | mean difference (a type of absolute effect estimate).
absolute-SMD | standardized mean difference | standardized mean difference (a type of absolute effect estimate).

absolute-MedianDiff | median difference | median difference (a type of absolute effect estimate).

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.647 Value Set http://hl7.org/fhir/ValueSet/service-type

URL: https://hl7.org/fhir/R4/valueset-service-type.html
Patient Administration Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/service-type
Version: | 4.0.1
Name: | ServiceType
Title: | Service type
Definition: | This value set defines an example set of codes of service-types.
Committee: | Patient Administration Work Group
OID: | 2.16.840.1.113883.4.642.3.518 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for ServiceType
• Resource: Appointment.serviceType (CodeableConcept / Example)
• Resource: Slot.serviceType (CodeableConcept / Example)
• Resource: Encounter.serviceType (CodeableConcept / Example)
• Resource: HealthcareService.type (CodeableConcept / Example)
• Resource: Schedule.serviceType (CodeableConcept / Example)

4.4.1.647.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/service-type

4.4.1.647.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 596 concepts

All codes from system http://terminology.hl7.org/CodeSystem/service-type

Code | Display | Definition
1 | Adoption/Permanent Care Info/Support | Adoption & permanent care information/support
2 | Aged Care Assessment | Aged care assessment
3 | Aged Care Information/Referral | Aged Care information/referral
4 | Aged Residential Care | Aged Residential Care
5 | Case Management for Older Persons | Case management for older persons
6 | Delivered Meals (Meals On Wheels) | Delivered meals (meals on wheels)
7 | Friendly Visiting | Friendly visiting
8 | Home Care/Housekeeping Assistance | Home care/housekeeping assistance
9 | Home Maintenance and Repair | Home maintenance and repair
10 | Personal Alarms/Alerts | Personal alarms/alerts
11 | Personal Care for Older Persons | Personal care for older persons
12 | Planned Activity Groups | Planned activity groups
13 | Acupuncture | Acupuncture
14 | Alexander Technique Therapy | Alexander technique therapy
15 | Aromatherapy | Aromatherapy
16 | Biorhythm Services | Biorhythm services
17 | Bowen Therapy | Bowen therapy
18 | Chinese Herbal Medicine | Chinese herbal medicine
19 | Feldenkrais | Feldenkrais
20 | Homoeopathy | Homoeopathy
21 | Hydrotherapy | Hydrotherapy

22 | Hypnotherapy | Hypnotherapy
23 | Kinesiology | Kinesiology
24 | Magnetic Therapy | Magnetic therapy
25 | Massage Therapy | Massage therapy
26 | Meditation | Meditation
27 | Myotherapy | Myotherapy
28 | Naturopathy | Naturopathy
29 | Reflexology | Reflexology
30 | Reiki | Reiki
31 | Relaxation Therapy | Relaxation therapy
32 | Shiatsu | Shiatsu
33 | Western Herbal Medicine | Western herbal medicine
34 | Family Day care | Family day care
35 | Holiday Programs | Holiday programs
36 | Kindergarten Inclusion Support | Kindergarten inclusion support for children with a disability
37 | Kindergarten/Preschool | Kindergarten/preschool
38 | Long Day Child Care | Long day child care
39 | Occasional Child Care | Occasional child care
40 | Outside School Hours Care | Outside school hours care
41 | Children's Play Programs | Children's play programs
42 | Parenting/Family Support/Education | Parenting & family management support/education
43 | Playgroup | Playgroup
44 | School Nursing | School nursing
45 | Toy Library | Toy library
46 | Child Protection/Child Abuse Report | Child protection/child abuse report
47 | Foster Care | Foster care
48 | Residential/Out-of-Home Care | Residential/ out of home care
49 | Support - Young People Leaving Care | Support for young people leaving care
50 | Audiology | Audiology
51 | Blood Donation | Blood donation
52 | Chiropractic | Chiropractic
53 | Dietetics | Dietetics
54 | Family Planning | Family planning
55 | Health Advocacy/Liaison Service | Health advocacy/Liaison service
56 | Health Information/Referral | Health information/referral
57 | Immunization | Immunization
58 | Maternal & Child Health | Maternal & child health
59 | Nursing | Nursing
60 | Nutrition | Nutrition
61 | Occupational Therapy | Occupational therapy
62 | Optometry | Optometry
63 | Osteopathy | Osteopathy
64 | Pharmacy | Pharmacy
65 | Physiotherapy | Physiotherapy
66 | Podiatry | Podiatry
67 | Sexual Health | Sexual health
68 | Speech Pathology/Therapy | Speech pathology/therapy
69 | Bereavement Counselling | Bereavement counselling
70 | Crisis Counselling | Crisis counselling
71 | Family Counselling/Therapy | Family counselling and/or family therapy
72 | Family Violence Counselling | Family violence counselling
73 | Financial Counselling | Financial counselling
74 | Generalist Counselling | Generalist counselling
75 | Genetic Counselling | Genetic counselling
76 | Health Counselling | Health counselling
77 | Mediation | Mediation
78 | Problem Gambling Counselling | Problem gambling counselling
79 | Relationship Counselling | Relationship counselling
80 | Sexual Assault Counselling | Sexual assault counselling
81 | Trauma Counselling | Trauma counselling
82 | Victims of Crime Counselling | Victims of crime counselling
83 | Cemetery Operation | Cemetery operation
84 | Cremation | Cremation
85 | Death Service Information | Death service information
86 | Funeral Services | Funeral services
87 | Endodontic | Endodontic
88 | General Dental | General dental
89 | Oral Medicine | Oral medicine
90 | Oral Surgery | Oral surgery
91 | Orthodontic | Orthodontic
92 | Paediatric Dentistry | Paediatric Dentistry
93 | Periodontic | Periodontic

94 | Prosthodontic | Prosthodontic
95 | Acquired Brain Injury Info/Referral | Acquired brain injury information/referral
96 | Disability Advocacy | Disability advocacy
97 | Disability Aids & Equipment | Disability aids & equipment
98 | Disability Case Management | Disability case management
99 | Disability Day Programs/Activities | Disability day programs & activities
100 | Disability Information/Referral | Disability information/referral
101 | Disability Support Packages | Disability support packages
102 | Disability Supported Accommodation | Disability supported accommodation
103 | Early Childhood Intervention | Early childhood intervention
104 | Hearing Aids & Equipment | Hearing aids & equipment
105 | Drug and/or Alcohol Counselling | Drug and/or alcohol counselling
106 | Drug/Alcohol Information/Referral | Drug and/or alcohol information/referral
107 | Needle & Syringe Exchange | Needle & Syringe exchange
108 | Non-resid. Alcohol/Drug Treatment | Non-residential alcohol and/or drug dependence treatment
109 | Pharmacotherapy | Pharmacotherapy (eg. methadone) program
110 | Quit Program | Quit program
111 | Residential Alcohol/Drug Treatment | Residential alcohol and/or drug dependence treatment
112 | Adult/Community Education | Adult/community education
113 | Higher Education | Higher education
114 | Primary Education | Primary education
115 | Secondary Education | Secondary education
116 | Training & Vocational Education | Training & vocational education
117 | Emergency Medical | Emergency medical
118 | Employment Placement and/or Support | Employment placement and/or support
119 | Vocational Rehabilitation | Vocational Rehabilitation
120 | Work Safety/Accident Prevention | Workplace safety and/or accident prevention
121 | Financial Assistance | Financial assistance
122 | Financial Information/Advice | Financial information/advice
123 | Material Aid | Material aid
124 | General Practice | General Practice/GP (doctor)
125 | Accommodation Placement/Support | Accommodation placement and/or support
126 | Crisis/Emergency Accommodation | Crisis/emergency accommodation
127 | Homelessness Support | Homelessness support
128 | Housing Information/Referral | Housing information/referral
129 | Public Rental Housing | Public rental housing
130 | Interpreting/Multilingual Service | Interpreting/Multilingual/Language service
131 | Juvenile Justice | Juvenile Justice
132 | Legal Advocacy | Legal advocacy
133 | Legal Information/Advice/Referral | Legal information/advice/referral
134 | Mental Health Advocacy | Mental health advocacy
135 | Mental Health Assess/Triage/Crisis Response | Mental health assessment/triage/crisis response
136 | Mental Health Case Management | Mental health case management/continuing care
137 | Mental Health Information/Referral | Mental health information/referral
138 | Mental Health Inpatient Services | Mental health inpatient services (hospital psychiatric unit) - requires referral
139 | Mental Health Non-residential Rehab | Mental health non-residential rehabilitation
140 | Mental Health Residential Rehab/CCU | Mental health residential rehabilitation/community care unit
141 | Psychiatry (Requires Referral) | Psychiatry (requires referral)
142 | Psychology | Psychology
143 | Martial Arts | Martial arts
144 | Personal Fitness Training | Personal fitness training
145 | Physical Activity Group | Physical activity group
146 | Physical Activity Programs | Physical activity programs
147 | Physical Fitness Testing | Physical fitness testing
148 | Pilates | Pilates
149 | Self-Defence | Self defence
150 | Sporting Club | Sporting club
151 | Yoga | Yoga
152 | Food Safety | Food safety
153 | Health Regulatory /Inspection /Cert. | Health regulatory, inspection and/or certification
154 | Work Health/Safety Inspection/Cert. | Workplace health and/or safety inspection and/or certification
155 | Carer Support | Carer support
156 | Respite Care | Respite care
157 | Anatomical Pathology | Anatomical Pathology (including Cytopathology & Forensic Pathology)
158 | Pathology - Clinical Chemistry | Pathology - Clinical Chemistry
159 | Pathology - General | Pathology - General
160 | Pathology - Genetics | Pathology - Genetics
161 | Pathology - Haematology | Pathology - Haematology
162 | Pathology - Immunology | Pathology - Immunology
163 | Pathology - Microbiology | Pathology - Microbiology
164 | Anaesthesiology - Pain Medicine | Anaesthesiology - Pain Medicine
165 | Cardiology | Cardiology

166 | Clinical Genetics | Clinical Genetics
167 | Clinical Pharmacology | Clinical Pharmacology
168 | Dermatology | Dermatology
169 | Endocrinology | Endocrinology
170 | Gastroenterology & Hepatology | Gastroenterology & Hepatology
171 | Geriatric Medicine | Geriatric medicine
172 | Immunology & Allergy | Immunology & Allergy
173 | Infectious Diseases | Infectious diseases
174 | Intensive Care Medicine | Intensive care medicine
175 | Medical Oncology | Medical Oncology
176 | Nephrology | Nephrology
177 | Neurology | Neurology
178 | Occupational Medicine | Occupational Medicine
179 | Palliative Medicine | Palliative Medicine
180 | Public Health Medicine | Public Health Medicine
181 | Rehabilitation Medicine | Rehabilitation Medicine
182 | Rheumatology | Rheumatology
183 | Sleep Medicine | Sleep Medicine
184 | Thoracic Medicine | Thoracic medicine
185 | Gynaecological Oncology | Gynaecological Oncology
186 | Obstetrics & Gynaecology | Obstetrics & Gynaecology
187 | Reproductive Endocrinology/Infertility | Reproductive Endocrinology & Infertility
188 | Urogynaecology | Urogynaecology
189 | Neonatology & Perinatology | Neonatology & Perinatology
190 | Paediatric Cardiology | Paediatric Cardiology
191 | Paediatric Clinical Genetics | Paediatric Clinical Genetics
192 | Paediatric Clinical Pharmacology | Paediatric Clinical Pharmacology
193 | Paediatric Endocrinology | Paediatric Endocrinology
194 | Paed. Gastroenterology/Hepatology | Paediatric Gastroenterology & Hepatology
195 | Paediatric Haematology | Paediatric Haematology
196 | Paediatric Immunology & Allergy | Paediatric Immunology & Allergy
197 | Paediatric Infectious Diseases | Paediatric Infectious diseases
198 | Paediatric Intensive Care Medicine | Paediatric intensive care medicine
199 | Paediatric Medical Oncology | Paediatric Medical Oncology
200 | Paediatric Medicine | Paediatric Medicine
201 | Paediatric Nephrology | Paediatric Nephrology
202 | Paediatric Neurology | Paediatric Neurology
203 | Paediatric Nuclear Medicine | Paediatric Nuclear Medicine
204 | Paediatric Rehabilitation Medicine | Paediatric Rehabilitation Medicine
205 | Paediatric Rheumatology | Paediatric Rheumatology
206 | Paediatric Sleep Medicine | Paediatric Sleep Medicine
207 | Paediatric Surgery | Paediatric Surgery
208 | Paediatric Thoracic Medicine | Paediatric Thoracic Medicine
209 | Diag. Radiology /Xray /CT /Fluoroscopy | Diagnostic Radiology/Xray/CT/Fluoroscopy
210 | Diagnostic Ultrasound | Diagnostic Ultrasound
211 | Magnetic Resonance Imaging (MRI) | Magnetic Resonance Imaging (MRI)
212 | Nuclear Medicine | Nuclear Medicine
213 | Obstetric/Gynaecological Ultrasound | Obstetric & Gynaecological Ultrasound
214 | Radiation Oncology | Radiation oncology
215 | Cardiothoracic Surgery | Cardiothoracic surgery
216 | Neurosurgery | Neurosurgery
217 | Ophthalmology | Ophthalmology
218 | Orthopaedic Surgery | Orthopaedic surgery
219 | Otolaryngology/Head & Neck Surgery | Otolaryngology - Head & Neck Surgery
220 | Plastic & Reconstructive Surgery | Plastic & Reconstructive Surgery
221 | Surgery - General | Surgery - General
222 | Urology | Urology
223 | Vascular Surgery | Vascular surgery
224 | Support Groups | Support groups
225 | Air ambulance | Air ambulance
226 | Ambulance | Ambulance
227 | Blood Transport | Blood transport
228 | Community Bus | Community bus
229 | Flying Doctor Service | Flying doctor service
230 | Patient Transport | Patient transport
231 | A&E | A&E
232 | A&EP | A&EP
233 | Abuse | Abuse
234 | ACAS | ACAS
235 | Access | Access
236 | Accident | Accident
237 | Acute Inpatient Serv | Acute Inpatient Service's

238 | Adult Day Programs | Adult Day Programs
239 | Adult Mental Health Services | Adult Mental Health Services
240 | Advice | Advice
241 | Advocacy | Advocacy
242 | Aged Persons Mental | Aged Persons Mental Health Residential Units
243 | Aged Persons Mental | Aged Persons Mental Health Services
244 | Aged Persons Mental | Aged Persons Mental Health Teams
245 | Aids | Aids
246 | Al-Anon | Al-Anon
247 | Alcohol | Alcohol
248 | Al-Teen | Al-Teen
249 | Antenatal | Antenatal
250 | Anxiety | Anxiety
251 | Arthritis | Arthritis
252 | Assessment | Assessment
253 | Assistance | Assistance
254 | Asthma | Asthma
255 | ATSS | ATSS
256 | Attendant Care | Attendant Care
257 | Babies | Babies
258 | Bathroom Modificatio | Bathroom Modification
259 | Behavior | Behavior
260 | Behavior Interventi | Behavior Intervention
261 | Bereavement | Bereavement
262 | Bipolar | Bipolar
263 | Birth | Birth
264 | Birth Control | Birth Control
265 | Birthing Options | Birthing Options
266 | BIST | BIST
267 | Blood | Blood
268 | Bone | Bone
269 | Bowel | Bowel
270 | Brain | Brain
271 | Breast Feeding | Breast Feeding
272 | Breast Screen | Breast Screen
273 | Brokerage | Brokerage
274 | Cancer | Cancer
275 | Cancer Support | Cancer Support
276 | Cardiovascular Disea | Cardiovascular Disease
277 | Care Packages | Care Packages
278 | Carer | Carer
279 | Case Management | Case Management
280 | Casualty | Casualty
281 | Centrelink | Centrelink
282 | Chemists | Chemists
283 | Child And Adolescent | Child And Adolescent Mental Health Services
284 | Child Care | Child Care
285 | Child Services | Child Services
286 | Children | Children
287 | Children's Services | Children's Services
288 | Cholesterol | Cholesterol
289 | Clothing | Clothing
290 | Community Based Acco | Community Based Accommodation
291 | Community Care Unit | Community Care Unit
292 | Community Child And | Community Child And Adolescent Mental Health Services
293 | Community Health | Community Health
294 | Community Residentia | Community Residential Unit
295 | Community Transport | Community Transport
296 | Companion Visiting | Companion Visiting
297 | Companionship | Companionship
298 | Consumer Advice | Consumer Advice
299 | Consumer Issues | Consumer Issues
300 | Continuing Care Serv | Continuing Care Services
301 | Contraception Inform | Contraception Information
302 | Coordinating Bodies | Coordinating Bodies
303 | Correctional Service | Correctional Services
304 | Council Environmenta | Council Environmental Health
305 | Counselling | Counselling
306 | Criminal | Criminal
307 | Crises | Crises
308 | Crisis Assessment An | Crisis Assessment And Treatment Services (Cats)
309 | Crisis Assistance | Crisis Assistance

310 | Crisis Refuge | Crisis Refuge
311 | Day Program | Day Program
312 | Deaf | Deaf
313 | Dental Hygiene | Dental Hygiene
314 | Dentistry | Dentistry
315 | Dentures | Dentures
316 | Depression | Depression
317 | Detoxification | Detoxification
318 | Diabetes | Diabetes
319 | Diaphragm Fitting | Diaphragm Fitting
320 | Dieticians | Dieticians
321 | Disabled Parking | Disabled Parking
322 | District Nursing | District Nursing
323 | Divorce | Divorce
324 | Doctors | Doctors
325 | Drink-Drive | Drink-Drive
326 | Dual Diagnosis Servi | Dual Diagnosis Services
327 | Early Choice | Early Choice
328 | Eating Disorder | Eating Disorder
330 | Emergency Relief | Emergency Relief
331 | Employment And Train | Employment And Training
332 | Environment | Environment
333 | Equipment | Equipment
334 | Exercise | Exercise
335 | Facility | Facility
336 | Family Choice | Family Choice
337 | Family Law | Family Law
338 | Family Options | Family Options
339 | Family Services | Family Services
340 | FFYA | FFYA
341 | Financial Aid | Financial Aid
342 | Fitness | Fitness
343 | Flexible Care Packag | Flexible Care Packages
344 | Food | Food
345 | Food Vouchers | Food Vouchers
346 | Forensic Mental Heal | Forensic Mental Health Services
347 | Futures | Futures
348 | Futures For Young Ad | Futures For Young Adults
349 | General Practitioner | General Practitioners
350 | Grants | Grants
351 | Grief | Grief
352 | Grief Counselling | Grief Counselling
353 | HACC | HACC
354 | Heart Disease | Heart Disease
355 | Help | Help
356 | High Blood Pressure | High Blood Pressure
357 | Home Help | Home Help
358 | Home Nursing | Home Nursing
359 | Homefirst | Homefirst
360 | Hospice Care | Hospice Care
361 | Hospital Services | Hospital Services
362 | Hospital To Home | Hospital To Home
364 | Hostel | Hostel
365 | Hostel Accommodation | Hostel Accommodation
366 | Household Items | Household Items
367 | Hypertension | Hypertension
368 | Illness | Illness
369 | Independent Living | Independent Living
370 | Information | Information
371 | Injury | Injury
372 | Intake | Intake
373 | Intensive Mobile You | Intensive Mobile Youth Outreach Services (Imyos)
374 | Intervention | Intervention
375 | Job Searching | Job Searching
376 | Justice | Justice
377 | Leisure | Leisure
378 | Loans | Loans
379 | Low Income Earners | Low Income Earners
380 | Lung | Lung
381 | Making A Difference | Making A Difference
382 | Medical Services | Medical Services
383 | Medical Specialists | Medical Specialists

384 | Medication Administr | Medication Administration
385 | Menstrual Informatio | Menstrual Information
386 | Methadone | Methadone
387 | Mobile Support And T | Mobile Support And Treatment Services (MSTS)
388 | Motor Neurone | Motor Neurone
389 | Multiple Sclerosis | Multiple Sclerosis
390 | Neighbourhood House | Neighbourhood House
391 | Nursing Home | Nursing Home
392 | Nursing Mothers | Nursing Mothers
393 | Obesity | Obesity
394 | Occupational Health | Occupational Health & Safety
395 | Optometrist | Optometrist
396 | Oral Hygiene | Oral Hygiene
397 | Outpatients | Outpatients
398 | Outreach Service | Outreach Service
399 | PADP | PADP
400 | Pain | Pain
401 | Pap Smear | Pap Smear
402 | Parenting | Parenting
403 | Peak Organizations | Peak Organizations
404 | Personal Care | Personal Care
405 | Pharmacies | Pharmacies
406 | Phobias | Phobias
407 | Physical | Physical
408 | Physical Activity | Physical Activity
409 | Postnatal | Postnatal
410 | Pregnancy | Pregnancy
411 | Pregnancy Tests | Pregnancy Tests
412 | Preschool | Preschool
413 | Prescriptions | Prescriptions
414 | Primary Mental Healt | Primary Mental Health And Early Intervention Teams
415 | Property Maintenance | Property Maintenance
416 | Prostate | Prostate
417 | Psychiatric | Psychiatric
418 | Psychiatric Disabili | Psychiatric Disability Support Services - Home-Based Outreach
419 | Psychiatric Disabili | Psychiatric Disability Support Services - Planned Respite
420 | Psychiatric Disabili | Psychiatric Disability Support Services - Residential Rehabilitation
421 | Psychiatric Disabili | Psychiatric Disability Support Services Home-Based Outreach
422 | Psychiatric Disabili | Psychiatric Disability Support Services Mutual Support And Self Help
423 | Psychiatric Support | Psychiatric Support
424 | Recreation | Recreation
425 | Referral | Referral
426 | Refuge | Refuge
427 | Rent Assistance | Rent Assistance
428 | Residential Faciliti | Residential Facilities
429 | Residential Respite | Residential Respite
430 | Respiratory | Respiratory
431 | Response | Response
432 | Rooming Houses | Rooming Houses
433 | Safe Sex | Safe Sex
434 | Secure Extended Care | Secure Extended Care Inpatient Services
435 | Self Help | Self Help
436 | Separation | Separation
437 | Services | Services
438 | Sex Education | Sex Education
439 | Sexual Abuse | Sexual Abuse
440 | Sexual Issues | Sexual Issues
441 | Sexually Transmitted | Sexually Transmitted Diseases
442 | SIDS | SIDS
443 | Social Support | Social Support
444 | Socialisation | Socialisation
445 | Special Needs | Special Needs
446 | Speech Therapist | Speech Therapist
447 | Splinting | Splinting
448 | Sport | Sport
449 | Statewide And Specia | Statewide And Specialist Services
450 | STD | STD
451 | STI | STI
452 | Stillbirth | Stillbirth
453 | Stomal Care | Stomal Care
454 | Stroke | Stroke
455 | Substance Abuse | Substance Abuse

456 | Support | Support
457 | Syringes | Syringes
458 | Teeth | Teeth
459 | Tenancy Advice | Tenancy Advice
460 | Terminal Illness | Terminal Illness
461 | Therapy | Therapy
462 | Transcription | Transcription
463 | Translating Services | Translating Services
464 | Translator | Translator
465 | Transport | Transport
466 | Vertebrae | Vertebrae
467 | Violence | Violence
468 | Vocational Guidance | Vocational Guidance
469 | Weight | Weight
470 | Welfare Assistance | Welfare Assistance
471 | Welfare Counselling | Welfare Counselling
472 | Wheelchairs | Wheelchairs
473 | Wound Management | Wound Management
474 | Young People At Risk | Young People At Risk
475 | Further Desc. - Community Health Care | Further Description - Community Health Care
476 | Library | Library
477 | Community Hours | Community Hours
478 | Further Desc. - Specialist Medical | Further Description - Specialist Medical
479 | Hepatology | Hepatology
480 | Gastroenterology | Gastroenterology
481 | Gynaecology | Gynaecology
482 | Obstetrics | Obstetrics
483 | Further Desc. - Specialist Surgical | Further Description - Specialist Surgical
484 | Placement Protection | Placement Protection
485 | Family Violence | Family Violence
486 | Integrated Family Services | Integrated Family Services
488 | Diabetes Educator | Diabetes Educator
489 | Kinship Care | Kinship Care
490 | General Mental Health Services | General Mental Health Services
491 | Exercise Physiology | Exercise Physiology
492 | Medical Research | Medical Research
493 | Youth | Youth
494 | Youth Services | Youth Services
495 | Youth Health | Youth Health
496 | Child and Family Ser | Child and Family Services
497 | Home Visits | Home Visits
498 | Mobile Services | Mobile Services
500 | Before and/or After | Before and/or After School Care
501 | Cancer Services | Cancer Services
502 | Integrated Cancer Se | Integrated Cancer Services
503 | Multidisciplinary Se | Multidisciplinary Services
504 | Multidisciplinary Ca | Multidisciplinary Cancer Services
505 | Meetings | Meetings
506 | Blood pressure monit | Blood pressure monitoring
507 | Dose administration | Dose administration aid
508 | Medical Equipment Hi | Medical Equipment Hire
509 | Parenting/Family Support/Education | Parenting & family support/education
510 | Deputising Service | Deputising Service
513 | Cancer Support Groups | Cancer Support Groups
514 | Community Cancer Services | Community Cancer Services
530 | Disability Care Transport | Disability Care Transport
531 | Aged Care Transport | Aged Care Transport
532 | Diabetes Education s | Diabetes Education service
533 | Cardiac Rehabilitati | Cardiac Rehabilitation Service
534 | Young Adult Diabetes | Young Adult Diabetes services (YADS)
535 | Pulmonary Rehabilita | Pulmonary Rehabilitation Service
536 | Art therapy | Art therapy
537 | Medication Reviews | Medication Reviews
538 | Telephone Counselling | Telephone Counselling
539 | Telephone Help Line | Telephone Help Line
540 | Online Service | Online Service
541 | Crisis - Mental Health | Crisis - Mental Health
542 | Youth Crisis | Youth Crisis
543 | Sexual Assault | Sexual Assault
544 | GPAH Other | GPAH Other
545 | Paediatric Dermatology | Paediatric Dermatology
546 | Veterans Services | Veterans Services

547 | Veterans | Veterans
548 | Food Relief/Food/Meals | Food Relief/food/meals
550 | Dementia Care | Dementia Care
551 | Alzheimer | Alzheimer
552 | Drug and/or Alcohol Support Groups | Drug and/or alcohol support groups
553 | 1-on-1 Support /Mentoring /Coaching | One on One Support/Mentoring/Coaching
554 | Chronic Disease Management | Chronic Disease Management
555 | Liaison Services | Liaison Services
556 | Walk-in Centre /Non-Emergency | Walk in Centre / non emergency
557 | Inpatients | Inpatients
558 | Spiritual Counselling | Spiritual Counselling
559 | Women's Health | Women's Health
560 | Men's Health | Men's Health
561 | Health Education/Awareness Program | Health education/Health awareness program
562 | Test Message | Test Message
563 | Remedial Massage | Remedial Massage
564 | Adolescent Mental Health Services | Adolescent Mental Health Services
565 | Youth Drop In/Assistance/Support | Youth drop in/assistance/support
566 | Aboriginal Health Worker | Aboriginal Health Worker
567 | Women's Health Clinic | Women's Health Clinic
568 | Men's Health Clinic | Men's Health Clinic
569 | Migrant Health Clinic | Migrant Health Clinic
570 | Refugee Health Clinic | Refugee Health Clinic
571 | Aboriginal Health Clinic | Aboriginal Health Clinic
572 | Nurse Practitioner Lead Clinic/s | Nurse Practitioner lead Clinic/s
573 | Nurse Lead Clinic/s | Nurse lead Clinic/s
574 | Culturally Tailored Support Groups | Culturally tailored support groups
575 | Culturally Tailored Health Promotion | Culturally tailored health promotion
576 | Rehabilitation | Rehabilitation
577 | Education Information/Referral | Education information/referral
580 | Social Work | Social Work
581 | Haematology | Haematology
582 | Maternity Shared Car | Maternity Shared Care
583 | Rehabilitation Servi | Rehabilitation Service
584 | Cranio-sacral Therapy | Cranio-Sacral Therapy
585 | Prosthetics & Orthotics | Prosthetics & Orthotics
589 | Home Medicine Review | Home Medicine Review
590 | GPAH - Medical | GPAH - Medical
591 | Music Therapy | Music Therapy
593 | Falls Prevention | Falls Prevention
599 | Accommodation/Tenancy | Accommodation/Tenancy
600 | Assess-Skill, Ability, Needs | Assess-Skill, Ability, Needs
601 | Assist Access/Maintain Employ | Assist Access/Maintain Employ
602 | Assist Prod-Pers Care/Safety | Assist Prod-Pers Care/Safety
603 | Assist-Integrate School/Ed | Assist-Integrate School/Ed
604 | Assist-Life Stage, Transition | Assist-Life Stage, Transition
605 | Assist-Personal Activities | Assist-Personal Activities
606 | Assist-Travel/Transport | Assist-Travel/Transport
607 | Assistive Equip-General Tasks | Assistive Equip-General Tasks
608 | Assistive Equip-Recreation | Assistive Equip-Recreation
609 | Assistive Prod-Household Task | Assistive Prod-Household Task
610 | Behavior Support | Behavior Support
611 | Comms & Info Equipment | Comms & Info Equipment
612 | Community Nursing Care | Community Nursing Care
613 | Daily Tasks/Shared Living | Daily Tasks/Shared Living
614 | Development-Life Skills | Development-Life Skills
615 | Early Childhood Supports | Early Childhood Supports
616 | Equipment Special Assess Setup | Equipment Special Assess Setup
617 | Hearing Equipment | Hearing Equipment
618 | Home Modification | Home Modification
619 | Household Tasks | Household Tasks
620 | Interpret/Translate | Interpret/Translate
621 | Other Innovative Supports | Other Innovative Supports
622 | Participate Community | Participate Community
623 | Personal Mobility Equipment | Personal Mobility Equipment
624 | Physical Wellbeing | Physical Wellbeing
625 | Plan Management | Plan Management
626 | Therapeutic Supports | Therapeutic Supports
627 | Training-Travel Independence | Training-Travel Independence
628 | Vehicle modifications | Vehicle modifications
629 | Vision Equipment | Vision Equipment

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.648 Value Set http://hl7.org/fhir/ValueSet/map-target-list-mode

URL: https://hl7.org/fhir/R4/valueset-map-target-list-mode.html
FHIR Infrastructure Work Group | Maturity Level: 2 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/map-target-list-mode
Version: | 4.0.1
Name: | StructureMapTargetListMode
Title: | StructureMapTargetListMode
Definition: | If field is a list, how to manage the production.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.685 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for StructureMapTargetListMode

• Resource: StructureMap.group.rule.target.listMode (code / Required)

4.4.1.648.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/map-target-list-mode

4.4.1.648.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 4 concepts

Expansion based on http://hl7.org/fhir/map-target-list-mode version 4.0.1

All codes from system http://hl7.org/fhir/map-target-list-mode

Code | Display | Definition
first | First | when the target list is being assembled, the items for this rule go first. If more than one rule defines a first item (for a given instance of mapping) then this is an error.
share | Share | the target instance is shared with the target instances generated by another rule (up to the first common n items, then create new ones).
last | Last | when the target list is being assembled, the items for this rule go last. If more than one rule defines a last item (for a given instance of mapping) then this is an error.
collate | Collate | re-use the first item in the list, and keep adding content to it.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.649 Value Set http://hl7.org/fhir/ValueSet/verificationresult-validation-process

URL: https://hl7.org/fhir/R4/valueset-verificationresult-validation-process.html

Patient Administration Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/verificationresult-validation-process
Version: | 4.0.1
Name: | validation-process
Title: | Validation-process
Definition: | The primary process by which the target is validated
Committee: | Patient Administration Work Group
OID: | 2.16.840.1.113883.4.642.3.888 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for validation-process

• Resource: VerificationResult.validationProcess (CodeableConcept / Example)

4.4.1.649.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/validation-process

4.4.1.649.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 6 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/validation-process version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/validation-process

Code | Display | Definition
edit-check | edit check
valueset | value set
primary | primary source
multi | multiple sources
standalone | standalone
in-context | in context

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.650 Value Set http://hl7.org/fhir/ValueSet/detectedissue-category

URL: https://hl7.org/fhir/R4/valueset-detectedissue-category.html
Clinical Decision Support Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/detectedissue-category
Version: | 4.0.1
Name: | DetectedIssueCategory
Title: | Detected Issue Category
Definition: | Kinds of issues or contraindications, such as 'drug-drug interaction', 'duplicate therapy', etc.
Committee: | Clinical Decision Support Work Group
OID: | 2.16.840.1.113883.4.642.3.204 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: DetectedIssue.code (CodeableConcept / Preferred)

4.4.1.650.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include codes from http://terminology.hl7.org/CodeSystem/v3-ActCode where concept is-a _AdministrationDetectedIssueCode

• Include codes from http://terminology.hl7.org/CodeSystem/v3-ActCode where concept is-a _ActSuppliedItemDetectedIssueCode

• Include codes from http://terminology.hl7.org/CodeSystem/v3-ActCode where concept is-a _SupplyDetectedIssueCode

4.4.1.650.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 27 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/v3-ActCode version 2018-08-12

All codes from system http://terminology.hl7.org/CodeSystem/v3-ActCode

Code | Display | Definition
FOOD | Food Interaction Alert | Proposed therapy may interact with certain foods
TPROD | Therapeutic Product Alert | Proposed therapy may interact with an existing or recent therapeutic product
DRG | Drug Interaction Alert | Proposed therapy may interact with an existing or recent drug therapy
NHP | Natural Health Product Alert | Proposed therapy may interact with existing or recent natural health product therapy
NONRX | Non-Prescription Interaction Alert | Proposed therapy may interact with a non-prescription drug (e.g. alcohol, tobacco, Aspirin)
PREVINEF | previously ineffective | Definition:The same or similar treatment has previously been attempted with the patient without achieving a positive effect.
DACT | drug action detected issue | Description:Proposed therapy may be contraindicated or ineffective based on an existing or recent drug therapy.
TIME | timing detected issue | Description:Proposed therapy may be inappropriate or ineffective based on the proposed start or end time.
ALRTENDLATE | end too late alert | Definition:Proposed therapy may be inappropriate or ineffective because the end of administration is too close to another planned therapy.
ALRTSTRTLATE | start too late alert | Definition:Proposed therapy may be inappropriate or ineffective because the start of administration is too late after the onset of the condition.
ENDLATE | End Too Late Alert | Proposed therapy may be inappropriate or ineffective because the end of administration is too close to another planned therapy
STRTLATE | Start Too Late Alert | Proposed therapy may be inappropriate or ineffective because the start of administration is too late after the onset of the condition
ALLDONE | already performed | Definition:The requested action has already been performed and so this request has no effect
FULFIL | fulfillment alert | Definition:The therapy being performed is in some way out of alignment with the requested therapy.
NOTACTN | no longer actionable | Definition:The status of the request being fulfilled has changed such that it is no longer actionable. This may be because the request has expired, has already been completely fulfilled or has been otherwise stopped or disabled. (Not used for 'suspended' orders.)
NOTEQUIV | not equivalent alert | Definition:The therapy being performed is not sufficiently equivalent to the therapy which was requested.
NOTEQUIVGEN | not generically equivalent alert | Definition:The therapy being performed is not generically equivalent (having the identical biological action) to the therapy which was requested.
NOTEQUIVTHER | not therapeutically equivalent alert | Definition:The therapy being performed is not therapeutically equivalent (having the same overall patient effect) to the therapy which was requested.
TIMING | event timing incorrect alert | Definition:The therapy is being performed at a time which diverges from the time the therapy was requested
INTERVAL | outside requested time | Definition:The therapy action is being performed outside the bounds of the time period requested
MINFREQ | too soon within frequency based on the usage | Definition:The therapy action is being performed too soon after the previous occurrence based on the requested frequency
HELD | held/suspended alert | Definition:There should be no actions taken in fulfillment of a request that has been held or suspended.
TOOLATE | Refill Too Late Alert | The patient is receiving a subsequent fill significantly later than would be expected based on the amount previously supplied and the therapy dosage instructions
TOOSOON | Refill Too Soon Alert | The patient is receiving a subsequent fill significantly earlier than would be expected based on the amount previously supplied and the therapy dosage instructions
HISTORIC | record recorded as historical | Description: While the record was accepted in the repository, there is a more recent version of a record of this type.
PATPREF | violates stated preferences | Definition:The proposed therapy goes against preferences or consent constraints recorded in the patient's record.
PATPREFALT | violates stated preferences, alternate available | Definition:The proposed therapy goes against preferences or consent constraints recorded in the patient's record. An alternate therapy meeting those constraints is available.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')

Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.651 Value Set http://hl7.org/fhir/ValueSet/report-result-codes

URL: https://hl7.org/fhir/R4/valueset-report-result-codes.html
FHIR Infrastructure Work Group | Maturity Level: 0 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/report-result-codes
Version: | 4.0.1
Name: | TestReportResult
Title: | TestReportResult
Definition: | The reported execution result.
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.718 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for TestReportResult
• Resource: TestReport.result (code / Required)

4.4.1.651.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://hl7.org/fhir/report-result-codes

4.4.1.651.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 3 concepts

Expansion based on http://hl7.org/fhir/report-result-codes version 4.0.1

All codes from system http://hl7.org/fhir/report-result-codes

Code | Display | Definition
pass | Pass | All test operations successfully passed all asserts.
fail | Fail | One or more test operations failed one or more asserts.
pending | Pending | One or more test operations is pending execution completion.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.652 Value Set http://hl7.org/fhir/ValueSet/mimetypes

URL: https://hl7.org/fhir/R4/valueset-mimetypes.html
FHIR Infrastructure Work Group | Maturity Level: N | Normative (from v4.0.0) | Use Context: Any

| This page has been approved as part of an ANSI standard. See the Infrastructure Package for further details.

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/mimetypes
Version: | 4.0.1
Name: | Mime Types
Title: | MimeType
Definition: | This value set includes all possible codes from BCP-13 (http://tools.ietf.org/html/bcp13)
Committee: | FHIR Infrastructure Work Group
OID: | 2.16.840.1.113883.4.642.3.1024 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: Endpoint.payloadMimeType (code / Required)
• Resource: CapabilityStatement.format (code / Required)
• Resource: CapabilityStatement.patchFormat (code / Required)
• Resource: Subscription.channel.payload (code / Required)
• Resource: Binary.contentType (code / Required)
• Resource: TestScript.setup.action.operation.accept (code / Required)
• Resource: TestScript.setup.action.operation.contentType (code / Required)
• Resource: TestScript.setup.action.assert.contentType (code / Required)
• Resource: Attachment.contentType (code / Required)
• Resource: Signature.targetFormat (code / Required)
• Resource: Signature.sigFormat (code / Required)
• Resource: ElementDefinition.mapping.language (code / Required)

4.4.1.652.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in urn:ietf:bcp:13

4.4.1.652.2 Expansion

This expansion generated 01 Nov 2019

This value set cannot be expanded because of the way it is defined - it has an infinite number of members

Code | System | Display

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.653 Value Set http://hl7.org/fhir/ValueSet/communication-topic

URL: https://hl7.org/fhir/R4/valueset-communication-topic.html
Patient Care Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/communication-topic
Version: | 4.0.1
Name: | CommunicationTopic
Title: | CommunicationTopic
Definition: | Codes describing the purpose or content of the communication.
Committee: | Patient Care Work Group
OID: | 2.16.840.1.113883.4.642.3.983 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for CommunicationTopic
• Resource: Communication.topic (CodeableConcept / Example)

4.4.1.653.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/communication-topic

4.4.1.653.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 6 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/communication-topic version 4.0.1

All codes from system http://terminology.hl7.org/CodeSystem/communication-topic

Code | Display | Definition
prescription-refill-request | Prescription Refill Request | The purpose or content of the communication is a prescription refill request.
progress-update | Progress Update | The purpose or content of the communication is a progress update.
report-labs | Report Labs | The purpose or content of the communication is to report labs.
appointment-reminder | Appointment Reminder | The purpose or content of the communication is an appointment reminder.
phone-consult | Phone Consult | The purpose or content of the communication is a phone consult.
summary-report | Summary Report | The purpose or content of the communication is a summary report.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.654 Value Set http://hl7.org/fhir/ValueSet/insuranceplan-type

URL: https://hl7.org/fhir/R4/valueset-insuranceplan-type.html
Patient Administration Work Group | Maturity Level: 1 | Draft | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/insuranceplan-type
Version: | 4.0.1
Name: | InsurancePlanType
Title: | Insurance plan type
Definition: | This example value set defines a set of codes that can be used to indicate a type of insurance plan.
Committee: | Patient Administration Work Group
OID: | 2.16.840.1.113883.4.642.3.870 (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• CodeSystem: This value set is the designated 'entire code system' value set for InsurancePlanType

• Resource: InsurancePlan.type (CodeableConcept / Example)

4.4.1.654.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/insurance-plan-type

4.4.1.654.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 10 concepts

All codes from system http://terminology.hl7.org/CodeSystem/insurance-plan-type

Code | Display | Definition
medical | Medical
dental | Dental
mental | Mental Health
subst-ab | Substance Abuse
vision | Vision
Drug | Drug
short-term | Short Term
long-term | Long Term Care
hospice | Hospice
home | Home Health

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.655 Value Set http://hl7.org/fhir/ValueSet/all-time-units

URL: https://hl7.org/fhir/R4/valueset-all-time-units.html
Vocabulary Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/all-time-units
Version: | 4.0.1
Name: | AllUCUMExpressionForTime
Title: | All UCUM Expression for Time
Definition: | Unified Code for Units of Measure (UCUM). This value set includes all UCUM codes for units of time
Committee: | ??
OID: | (for OID based terminology systems)
Copyright: | The UCUM codes, UCUM table (regardless of format), and UCUM Specification are copyright © 1999-2009, Regenstrief Institute, Inc. and the Unified Codes for Units of Measures (UCUM) Organization. All rights reserved.
Source Resource | XML / JSON

This value set is not currently used

4.4.1.655.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include codes from http://unitsofmeasure.org where canonical = a

4.4.1.655.2 Expansion

This expansion generated 01 Nov 2019

This value set cannot be expanded because of the way it is defined - it has an infinite number of members

Code | System | Display

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.656 Value Set http://hl7.org/fhir/ValueSet/all-distance-units

URL: https://hl7.org/fhir/R4/valueset-all-distance-units.html
Vocabulary Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/all-distance-units
Version: | 4.0.1
Name: | AllUCUMExpressionForDistance
Title: | All UCUM Expression for Distance
Definition: | Unified Code for Units of Measure (UCUM). This value set includes all UCUM codes for units of length
Committee: | ??
OID: | (for OID based terminology systems)
Copyright: | The UCUM codes, UCUM table (regardless of format), and UCUM Specification are copyright © 1999-2009, Regenstrief Institute, Inc. and the Unified Codes for Units of Measures (UCUM) Organization. All rights reserved.
Source Resource | XML / JSON

This value set is not currently used

4.4.1.656.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include codes from http://unitsofmeasure.org where canonical = m

4.4.1.656.2 Expansion

This expansion generated 01 Nov 2019

This value set has >1000 codes in it. In order to keep the publication size manageable, only a selection (1000 codes) of the whole set of codes is shown

All codes from system http://unitsofmeasure.org

Code	Display
[Ch]	French (catheter gauge)
[ft_i]	Feet
[fth_i]	fathom
[in_i]	inch (international)
[mi_i]	statute mile
[mi_us]	mile
[nmi_i]	nautical mile
[yd_i]	yard
cm	centimeter
dm	decimeter
fm	femtometer
km	kilometer
m	meter
mL/({h'b}.m2)	milliliter / h'b * (meter ^ 2)
mL/[sin_i]	milliliter per square inch (international)
mL/m2	milliliter per square meter
mm	millimeter
nm	nanometer
pm	picometer
um	micrometer
[ch_br]	Gunter's chain
[ch_us]	Gunter's chain Surveyor's chain
[cicero]	cicero Didot's pica
[didot]	didot Didot's point
[ft_br]	foot
[ft_us]	foot
[fth_br]	fathom
[fth_us]	fathom
[fur_us]	furlong
[hd_i]	hand
[in_br]	inch
[in_us]	inch
[ligne]	ligne French line
[lk_br]	link for Gunter's chain
[lk_us]	link for Gunter's chain
[lne]	line
[ly]	light-year
[mi_br]	mile
[mil_i]	mil
[mil_us]	mil
[nmi_br]	nautical mile
[pc_br]	pace
[pca]	pica
[pca_pr]	Printer's pica
[pied]	pied French foot
[pnt]	point
[pnt_pr]	Printer's point
[pouce]	pouce French inch
[rch_us]	Ramden's chain Engineer's chain
[rd_br]	rod
[rd_us]	rod
[rlk_us]	link for Ramden's chain
[smoot]	Smoot
[yd_br]	yard
[yd_us]	yard
Ao	Ångström
AU	astronomic unit

pc | parsec

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.657 Value Set http://hl7.org/fhir/ValueSet/iso3166-1-N

URL: https://hl7.org/fhir/R4/valueset-iso3166-1-N.html
Vocabulary Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/iso3166-1-N
Version: | 4.0.1
Name: | Iso3166-1-N
Title: | Iso 3166 Part 1: Numeric Codes
Definition: | This value set defines the ISO 3166 Part 1 Numeric codes
Committee: | ??
OID: | (for OID based terminology systems)
Source Resource | XML / JSON

This value set is not currently used

4.4.1.657.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include codes from urn:iso:std:iso:3166 where code matches (by regex) [0-9]{3}

4.4.1.657.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 249 concepts

All codes from system urn:iso:std:iso:3166

Code | Display
004 | Afghanistan
008 | Albania
010 | Antarctica
012 | Algeria
016 | American Samoa
020 | Andorra
024 | Angola
028 | Antigua and Barbuda
031 | Azerbaijan
032 | Argentina
036 | Australia
040 | Austria
044 | Bahamas
048 | Bahrain
050 | Bangladesh
051 | Armenia
052 | Barbados
056 | Belgium
060 | Bermuda
064 | Bhutan
068 | Bolivia, Plurinational State of
070 | Bosnia and Herzegovina
072 | Botswana
074 | Bouvet Island
076 | Brazil
084 | Belize
086 | British Indian Ocean Territory
090 | Solomon Islands
092 | Virgin Islands, British

096 | Brunei Darussalam
100 | Bulgaria
104 | Myanmar
108 | Burundi
112 | Belarus
116 | Cambodia
120 | Cameroon
124 | Canada
132 | Cabo Verde
136 | Cayman Islands
140 | Central African Republic
144 | Sri Lanka
148 | Chad
152 | Chile
156 | China
158 | Taiwan, Province of China
162 | Christmas Island
166 | Cocos (Keeling) Islands
170 | Colombia
174 | Comoros
175 | Mayotte
178 | Congo
180 | Congo, the Democratic Republic of the
184 | Cook Islands
188 | Costa Rica
191 | Croatia
192 | Cuba
196 | Cyprus
203 | Czechia
204 | Benin
208 | Denmark
212 | Dominica
214 | Dominican Republic
218 | Ecuador
222 | El Salvador
226 | Equatorial Guinea
231 | Ethiopia
232 | Eritrea
233 | Estonia
234 | Faroe Islands
238 | Falkland Islands (Malvinas)
239 | South Georgia and the South Sandwich Islands
242 | Fiji
246 | Finland
248 | Åland Islands
250 | France
254 | French Guiana
258 | French Polynesia
260 | French Southern Territories
262 | Djibouti
266 | Gabon
268 | Georgia
270 | Gambia
275 | Palestine, State of
276 | Germany
288 | Ghana
292 | Gibraltar
296 | Kiribati
300 | Greece
304 | Greenland
308 | Grenada
312 | Guadeloupe
316 | Guam
320 | Guatemala
324 | Guinea
328 | Guyana
332 | Haiti
334 | Heard Island and McDonald Islands
336 | Holy See
340 | Honduras
344 | Hong Kong
348 | Hungary

352 | Iceland
356 | India
360 | Indonesia
364 | Iran, Islamic Republic of
368 | Iraq
372 | Ireland
376 | Israel
380 | Italy
384 | Côte d'Ivoire
388 | Jamaica
392 | Japan
398 | Kazakhstan
400 | Jordan
404 | Kenya
408 | Korea, Democratic People's Republic of
410 | Korea, Republic of
414 | Kuwait
417 | Kyrgyzstan
418 | Lao People's Democratic Republic
422 | Lebanon
426 | Lesotho
428 | Latvia
430 | Liberia
434 | Libya
438 | Liechtenstein
440 | Lithuania
442 | Luxembourg
446 | Macao
450 | Madagascar
454 | Malawi
458 | Malaysia
462 | Maldives
466 | Mali
470 | Malta
474 | Martinique
478 | Mauritania
480 | Mauritius
484 | Mexico
492 | Monaco
496 | Mongolia
498 | Moldova, Republic of
499 | Montenegro
500 | Montserrat
504 | Morocco
508 | Mozambique
512 | Oman
516 | Namibia
520 | Nauru
524 | Nepal
528 | Netherlands
531 | Curaçao
533 | Aruba
534 | Sint Maarten (Dutch part)
535 | Bonaire, Sint Eustatius and Saba
540 | New Caledonia
548 | Vanuatu
554 | New Zealand
558 | Nicaragua
562 | Niger
566 | Nigeria
570 | Niue
574 | Norfolk Island
578 | Norway
580 | Northern Mariana Islands
581 | United States Minor Outlying Islands
583 | Micronesia, Federated States of
584 | Marshall Islands
585 | Palau
586 | Pakistan
591 | Panama
598 | Papua New Guinea
600 | Paraguay

604 | Peru
608 | Philippines
612 | Pitcairn
616 | Poland
620 | Portugal
624 | Guinea-Bissau
626 | Timor-Leste
630 | Puerto Rico
634 | Qatar
638 | Réunion
642 | Romania
643 | Russian Federation
646 | Rwanda
652 | Saint Barthélemy
654 | Saint Helena, Ascension and Tristan da Cunha
659 | Saint Kitts and Nevis
660 | Anguilla
662 | Saint Lucia
663 | Saint Martin (French part)
666 | Saint Pierre and Miquelon
670 | Saint Vincent and the Grenadines
674 | San Marino
678 | Sao Tome and Principe
682 | Saudi Arabia
686 | Senegal
688 | Serbia
690 | Seychelles
694 | Sierra Leone
702 | Singapore
703 | Slovakia
704 | Viet Nam
705 | Slovenia
706 | Somalia
710 | South Africa
716 | Zimbabwe
724 | Spain
728 | South Sudan
729 | Sudan
732 | Western Sahara
740 | Suriname
744 | Svalbard and Jan Mayen
748 | Swaziland
752 | Sweden
756 | Switzerland
760 | Syrian Arab Republic
762 | Tajikistan
764 | Thailand
768 | Togo
772 | Tokelau
776 | Tonga
780 | Trinidad and Tobago
784 | United Arab Emirates
788 | Tunisia
792 | Turkey
795 | Turkmenistan
796 | Turks and Caicos Islands
798 | Tuvalu
800 | Uganda
804 | Ukraine
807 | Macedonia, the former Yugoslav Republic of
818 | Egypt
826 | United Kingdom
831 | Guernsey
832 | Jersey
833 | Isle of Man
834 | Tanzania, United Republic of
840 | United States of America
850 | Virgin Islands, U.S.
854 | Burkina Faso
858 | Uruguay
860 | Uzbekistan
862 | Venezuela, Bolivarian Republic of

876 | Wallis and Futuna
882 | Samoa
887 | Yemen
894 | Zambia

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.658 Value Set http://hl7.org/fhir/ValueSet/performer-function

URL: https://hl7.org/fhir/R4/valueset-performer-function.html
Vocabulary Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined at http://www.hl7.org/Special/committees/orders/index.cfm .

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/performer-function
Version: | 4.0.1
Name: | Performer Function Codes
Title:
Definition: | The types of involvement of the performer in the Event.
Committee: | ??
OID: | (for OID based terminology systems)
Source Resource | XML / JSON

This value set is used in the following places:

• Extension: http://hl7.org/fhir/StructureDefinition/event-performerFunction: performerFunction (CodeableConcept / Extensible)

4.4.1.658.1 Content Logical Definition

This value set includes codes from the following code systems:

http://terminology.hl7.org/CodeSystem/v3-ParticipationType

Code | Display
TRANS | Transcriber | An entity entering the data into the originating system. The data entry entity is collected optionally for internal quality control purposes. This includes the transcriptionist for dictated text transcribed into electronic form.
PART | Participation | Indicates that the target of the participation is involved in some manner in the act, but does not qualify how.
ATND | attender | The practitioner that has responsibility for overseeing a patient's care during a patient encounter.
CON | consultant | An advisor participating in the service by performing evaluations and making recommendations.
AUT | author (originator) | Definition: A party that originates the Act and therefore has responsibility for the information given in the Act and ownership of this Act. Example: the report writer, the person writing the act definition, the guideline author, the placer of an order, the EKG cart (device) creating a report etc. Every Act should have an author. Authorship is regardless of mood always actual authorship. Examples of such policies might include: The author and anyone they explicitly delegate may update the report; All administrators within the same clinic may cancel and reschedule appointments created by other administrators within that clinic; A party that is neither an author nor a party who is extended authorship maintenance rights by policy, may only amend, reverse, override, replace, or follow up in other ways on this Act, whereby the Act remains intact and is linked to another Act authored by that other party.
INF | informant | A source of reported information (e.g., a next of kin who answers questions about the patient's history). For history questions, the patient is logically an informant, yet the informant of history questions is implicitly the subject.
ENT | data entry person | A person entering the data into the originating system. The data entry person is collected optionally for internal quality control purposes. This includes the transcriptionist for dictated text.
WIT | witness | Only with service events. A person witnessing the action happening without doing anything. A witness is not necessarily aware, much less approves of anything stated in the service event. Example for a witness is students watching an operation or an advanced directive witness.
PPRF | primary performer | The principal or primary performer of the act.
SPRF | secondary performer | A person assisting in an act through his substantial presence and involvement This includes: assistants, technicians, associates, or whatever the job titles may be.
RESP | responsible party | The person or organization that has primary responsibility for the act. The responsible party is not necessarily present in an action, but is accountable for the action through the power to delegate, and the duty to review actions with the performing actor after the fact. This responsibility may be ethical, legal, contractual, fiscal, or fiduciary in nature. Example: A person who is the head of a biochemical laboratory; a sponsor for a policy or government program.
VRF | verifier | A person who verifies the correctness and appropriateness of the service (plan, order, event, etc.) and hence takes on accountability.

AUTHEN | authenticator | A verifier who attests to the accuracy of an act, but who does not have privileges to legally authenticate the act. An example would be a resident physician who sees a patient and dictates a note, then later signs it. Their signature constitutes an authentication.
LA | legal authenticator | A verifier who legally authenticates the accuracy of an act. An example would be a staff physician who sees a patient and dictates a note, then later signs it. Their signature constitutes a legal authentication.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.659 Value Set http://hl7.org/fhir/ValueSet/written-language

URL: https://hl7.org/fhir/R4/valueset-written-language.html
Vocabulary Work Group | Maturity Level: 0 | Informative | Use Context: Not Intended for Production use

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/written-language
Version: | 4.0.1
Name: | Common Languages (Written)
Title:
Definition: | This value set includes common codes from BCP-47 (http://tools.ietf.org/html/bcp47) for the purpose of writing; this value set (unlike the common languages value set) doesn't include dialects except where they are relevant for written languages
Committee: | ??
OID: | (for OID based terminology systems)
Source Resource | XML / JSON

This value set is not currently used

4.4.1.659.1 Content Logical Definition

This value set includes codes from the following code systems:

urn:ietf:bcp:47

Code | Display
ar | Arabic
bn | Bengali
cs | Czech
da | Danish
de | German
el | Greek
en | English
es | Spanish
fi | Finnish
fr | French
fy | Frysian
hi | Hindi
hr | Croatian
it | Italian
ja | Japanese
ko | Korean
nl | Dutch
no | Norwegian
pa | Punjabi
pl | Polish
pt | Portuguese
ru | Russian
sr | Serbian
sv | Swedish
te | Telegu
zh | Chinese

Additional Language Displays

Code | Dansk (Danish, da) | Deutsch (German, de) | English (English, en) | Nederlands (Dutch, nl) | Polskie (Polish, pl) | Русский (Russian, ru) | □□ (Chinese, zh)
ar | Arabisk | | Arabic | Arabisch
bn | Bengalsk | Bengalisch | Bengali | Bengaals | | Бенгальский | □□□□
cs | Tjekkisk | Tschechisch | Czech | Tsjechisch | | Чешский | □□□
da | Dansk | Dänisch | Danish | Deens | | Датский | □□□
de | Tysk | Deutsch | German | Duits | | Немецкий | □□
el | Græsk | Griechisch | Greek | Grieks | | Греческий | □□□
en | Engelsk | Englisch | English | Engels | | Английский | □□
es | Spansk | Spanisch | Spanish | Spaans | | Испанский | □□□□
fi | Finsk | Finnisch | Finnish | Fins | | Финский | □□□
fr | Fransk | Französisch | French | Frans | | Французский | □□
fy | Frisisk | Friesisch | Frysian | Fries | | | □□□□□
hi | Hindi | | Hindi | Hindi
hr | Kroatisk | Kroatisch | Croatian | Kroatisch | | Хорватский | □□□□□
it | Italiensk | Italienisch | Italian | Italiaans | | Итальянский | □□□□
ja | Japansk | Japanisch | Japanese | Japans | | Японский | □□
ko | Koreansk | Koreanisch | Korean | Koreaans | | Корейский | □□□
nl | Hollandsk | Niederländisch | Dutch | Nederlands | | Нидерландский | □□□
no | Norsk | Norwegisch | Norwegian | Noors | | Норвежский | □□□
pa | Punjabi | | Punjabi | Punjabi
pl | | | | | Polskie
pt | Portugisisk | Portugiesisch | Portuguese | Portugees | | Португальский | □□□□
ru | Russisk | Russisch | Russian | Russisch | | Русский | □□
sr | Serbisk | Serbisch | Serbian | Servisch | | Сербский | □□□□□
sv | Svensk | Schwedisch | Swedish | Zweeds | | Шведский | □□□
te | Telugu | Telugu | Telegu | Teloegoe | | Телугу | □□□□
zh | Kinesisk | Chinesisch | Chinese | Chinees | | Китайский | □□

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.660 Value Set http://hl7.org/fhir/ValueSet/iso3166-1-2

URL: https://hl7.org/fhir/R4/valueset-iso3166-1-2.html
Vocabulary Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/iso3166-1-2
Version: | 4.0.1
Name: | Iso3166-1-2
Title: | Iso 3166 Part 1: 2 Letter Codes
Definition: | This value set defines the ISO 3166 Part 1 2-letter codes
Committee: | ??
OID: | (for OID based terminology systems)
Source Resource | XML / JSON

This value set is not currently used

4.4.1.660.1 Content Logical Definition

This value set includes codes from the following code systems:

urn:iso:std:iso:3166

Code | Display
AD | Andorra
AE | United Arab Emirates
AF | Afghanistan
AG | Antigua and Barbuda
AI | Anguilla
AL | Albania
AM | Armenia
AO | Angola

AQ | Antarctica
AR | Argentina
AS | American Samoa
AT | Austria
AU | Australia
AW | Aruba
AX | Åland Islands
AZ | Azerbaijan
BA | Bosnia and Herzegovina
BB | Barbados
BD | Bangladesh
BE | Belgium
BF | Burkina Faso
BG | Bulgaria
BH | Bahrain
BI | Burundi
BJ | Benin
BL | Saint Barthélemy
BM | Bermuda
BN | Brunei Darussalam
BO | Bolivia, Plurinational State of
BQ | Bonaire, Sint Eustatius and Saba
BR | Brazil
BS | Bahamas
BT | Bhutan
BV | Bouvet Island
BW | Botswana
BY | Belarus
BZ | Belize
CA | Canada
CC | Cocos (Keeling) Islands
CD | Congo, the Democratic Republic of the
CF | Central African Republic
CG | Congo
CH | Switzerland
CI | Côte d''Ivoire
CK | Cook Islands
CL | Chile
CM | Cameroon
CN | China
CO | Colombia
CR | Costa Rica
CU | Cuba
CV | Cabo Verde
CW | Curaçao
CX | Christmas Island
CY | Cyprus
CZ | Czechia
DE | Germany
DJ | Djibouti
DK | Denmark
DM | Dominica
DO | Dominican Republic
DZ | Algeria
EC | Ecuador
EE | Estonia
EG | Egypt
EH | Western Sahara
ER | Eritrea
ES | Spain
ET | Ethiopia
FI | Finland
FJ | Fiji
FK | Falkland Islands (Malvinas)
FM | Micronesia, Federated States of
FO | Faroe Islands
FR | France
GA | Gabon
GB | United Kingdom of Great Britain and Northern Ireland
GD | Grenada
GE | Georgia
GF | French Guiana

GG | Guernsey
GH | Ghana
GI | Gibraltar
GL | Greenland
GM | Gambia
GN | Guinea
GP | Guadeloupe
GQ | Equatorial Guinea
GR | Greece
GS | South Georgia and the South Sandwich Islands
GT | Guatemala
GU | Guam
GW | Guinea-Bissau
GY | Guyana
HK | Hong Kong
HM | Heard Island and McDonald Islands
HN | Honduras
HR | Croatia
HT | Haiti
HU | Hungary
ID | Indonesia
IE | Ireland
IL | Israel
IM | Isle of Man
IN | India
IO | British Indian Ocean Territory
IQ | Iraq
IR | Iran, Islamic Republic of
IS | Iceland
IT | Italy
JE | Jersey
JM | Jamaica
JO | Jordan
JP | Japan
KE | Kenya
KG | Kyrgyzstan
KH | Cambodia
KI | Kiribati
KM | Comoros
KN | Saint Kitts and Nevis
KP | Korea, Democratic People''s Republic of
KR | Korea, Republic of
KW | Kuwait
KY | Cayman Islands
KZ | Kazakhstan
LA | Lao People''s Democratic Republic
LB | Lebanon
LC | Saint Lucia
LI | Liechtenstein
LK | Sri Lanka
LR | Liberia
LS | Lesotho
LT | Lithuania
LU | Luxembourg
LV | Latvia
LY | Libya
MA | Morocco
MC | Monaco
MD | Moldova, Republic of
ME | Montenegro
MF | Saint Martin (French part)
MG | Madagascar
MH | Marshall Islands
MK | Macedonia, the former Yugoslav Republic of
ML | Mali
MM | Myanmar
MN | Mongolia
MO | Macao
MP | Northern Mariana Islands
MQ | Martinique
MR | Mauritania
MS | Montserrat

MT | Malta
MU | Mauritius
MV | Maldives
MW | Malawi
MX | Mexico
MY | Malaysia
MZ | Mozambique
NA | Namibia
NC | New Caledonia
NE | Niger
NF | Norfolk Island
NG | Nigeria
NI | Nicaragua
NL | Netherlands
NO | Norway
NP | Nepal
NR | Nauru
NU | Niue
NZ | New Zealand
OM | Oman
PA | Panama
PE | Peru
PF | French Polynesia
PG | Papua New Guinea
PH | Philippines
PK | Pakistan
PL | Poland
PM | Saint Pierre and Miquelon
PN | Pitcairn
PR | Puerto Rico
PS | Palestine, State of
PT | Portugal
PW | Palau
PY | Paraguay
QA | Qatar
RE | Réunion
RO | Romania
RS | Serbia
RU | Russian Federation
RW | Rwanda
SA | Saudi Arabia
SB | Solomon Islands
SC | Seychelles
SD | Sudan
SE | Sweden
SG | Singapore
SH | Saint Helena, Ascension and Tristan da Cunha
SI | Slovenia
SJ | Svalbard and Jan Mayen
SK | Slovakia
SL | Sierra Leone
SM | San Marino
SN | Senegal
SO | Somalia
SR | Suriname
SS | South Sudan
ST | Sao Tome and Principe
SV | El Salvador
SX | Sint Maarten (Dutch part)
SY | Syrian Arab Republic
SZ | Swaziland
TC | Turks and Caicos Islands
TD | Chad
TF | French Southern Territories
TG | Togo
TH | Thailand
TJ | Tajikistan
TK | Tokelau
TL | Timor-Leste
TM | Turkmenistan
TN | Tunisia
TO | Tonga

TR | Turkey
TT | Trinidad and Tobago
TV | Tuvalu
TW | Taiwan, Province of China
TZ | Tanzania, United Republic of
UA | Ukraine
UG | Uganda
UM | United States Minor Outlying Islands
US | United States of America
UY | Uruguay
UZ | Uzbekistan
VA | Holy See
VC | Saint Vincent and the Grenadines
VE | Venezuela, Bolivarian Republic of
VG | Virgin Islands, British
VI | Virgin Islands,
VN | Viet Nam
VU | Vanuatu
WF | Wallis and Futuna
WS | Samoa
YE | Yemen
YT | Mayotte
ZA | South Africa
ZM | Zambia
ZW | Zimbabwe

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.661 Value Set http://hl7.org/fhir/ValueSet/iso3166-1-3

URL: https://hl7.org/fhir/R4/valueset-iso3166-1-3.html
Vocabulary Work Group | Maturity Level: 0 | Informative | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/iso3166-1-3
Version: | 4.0.1
Name: | Iso3166-1-3
Title: | Iso 3166 Part 1: 3 Letter Codes
Definition: | This value set defines the ISO 3166 Part 1 3-letter codes
Committee: | ??
OID: | (for OID based terminology systems)
Source Resource | XML / JSON

This value set is not currently used

4.4.1.661.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include codes from urn:iso:std:iso:3166 where code matches (by regex) [A-Z]{3}

4.4.1.661.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 249 concepts

All codes from system urn:iso:std:iso:3166

Code | Display
ABW | Aruba
AFG | Afghanistan
AGO | Angola
AIA | Anguilla
ALA | Åland Islands

ALB | Albania
AND | Andorra
ARE | United Arab Emirates
ARG | Argentina
ARM | Armenia
ASM | American Samoa
ATA | Antarctica
ATF | French Southern Territories
ATG | Antigua and Barbuda
AUS | Australia
AUT | Austria
AZE | Azerbaijan
BDI | Burundi
BEL | Belgium
BEN | Benin
BES | Bonaire, Sint Eustatius and Saba
BFA | Burkina Faso
BGD | Bangladesh
BGR | Bulgaria
BHR | Bahrain
BHS | Bahamas
BIH | Bosnia and Herzegovina
BLM | Saint Barthélemy
BLR | Belarus
BLZ | Belize
BMU | Bermuda
BOL | Bolivia, Plurinational State of
BRA | Brazil
BRB | Barbados
BRN | Brunei Darussalam
BTN | Bhutan
BVT | Bouvet Island
BWA | Botswana
CAF | Central African Republic
CAN | Canada
CCK | Cocos (Keeling) Islands
CHE | Switzerland
CHL | Chile
CHN | China
CIV | Côte d'Ivoire
CMR | Cameroon
COD | Congo, the Democratic Republic of the
COG | Congo
COK | Cook Islands
COL | Colombia
COM | Comoros
CPV | Cabo Verde
CRI | Costa Rica
CUB | Cuba
CUW | Curaçao
CXR | Christmas Island
CYM | Cayman Islands
CYP | Cyprus
CZE | Czechia
DEU | Germany
DJI | Djibouti
DMA | Dominica
DNK | Denmark
DOM | Dominican Republic
DZA | Algeria
ECU | Ecuador
EGY | Egypt
ERI | Eritrea
ESH | Western Sahara
ESP | Spain
EST | Estonia
ETH | Ethiopia
FIN | Finland
FJI | Fiji
FLK | Falkland Islands (Malvinas)
FRA | France
FRO | Faroe Islands

FSM | Micronesia, Federated States of
GAB | Gabon
GBR | United Kingdom
GEO | Georgia
GGY | Guernsey
GHA | Ghana
GIB | Gibraltar
GIN | Guinea
GLP | Guadeloupe
GMB | Gambia
GNB | Guinea-Bissau
GNQ | Equatorial Guinea
GRC | Greece
GRD | Grenada
GRL | Greenland
GTM | Guatemala
GUF | French Guiana
GUM | Guam
GUY | Guyana
HKG | Hong Kong
HMD | Heard Island and McDonald Islands
HND | Honduras
HRV | Croatia
HTI | Haiti
HUN | Hungary
IDN | Indonesia
IMN | Isle of Man
IND | India
IOT | British Indian Ocean Territory
IRL | Ireland
IRN | Iran, Islamic Republic of
IRQ | Iraq
ISL | Iceland
ISR | Israel
ITA | Italy
JAM | Jamaica
JEY | Jersey
JOR | Jordan
JPN | Japan
KAZ | Kazakhstan
KEN | Kenya
KGZ | Kyrgyzstan
KHM | Cambodia
KIR | Kiribati
KNA | Saint Kitts and Nevis
KOR | Korea, Republic of
KWT | Kuwait
LAO | Lao People's Democratic Republic
LBN | Lebanon
LBR | Liberia
LBY | Libya
LCA | Saint Lucia
LIE | Liechtenstein
LKA | Sri Lanka
LSO | Lesotho
LTU | Lithuania
LUX | Luxembourg
LVA | Latvia
MAC | Macao
MAF | Saint Martin (French part)
MAR | Morocco
MCO | Monaco
MDA | Moldova, Republic of
MDG | Madagascar
MDV | Maldives
MEX | Mexico
MHL | Marshall Islands
MKD | Macedonia, the former Yugoslav Republic of
MLI | Mali
MLT | Malta
MMR | Myanmar
MNE | Montenegro

MNG | Mongolia
MNP | Northern Mariana Islands
MOZ | Mozambique
MRT | Mauritania
MSR | Montserrat
MTQ | Martinique
MUS | Mauritius
MWI | Malawi
MYS | Malaysia
MYT | Mayotte
NAM | Namibia
NCL | New Caledonia
NER | Niger
NFK | Norfolk Island
NGA | Nigeria
NIC | Nicaragua
NIU | Niue
NLD | Netherlands
NOR | Norway
NPL | Nepal
NRU | Nauru
NZL | New Zealand
OMN | Oman
PAK | Pakistan
PAN | Panama
PCN | Pitcairn
PER | Peru
PHL | Philippines
PLW | Palau
PNG | Papua New Guinea
POL | Poland
PRI | Puerto Rico
PRK | Korea, Democratic People's Republic of
PRT | Portugal
PRY | Paraguay
PSE | Palestine, State of
PYF | French Polynesia
QAT | Qatar
REU | Réunion
ROU | Romania
RUS | Russian Federation
RWA | Rwanda
SAU | Saudi Arabia
SDN | Sudan
SEN | Senegal
SGP | Singapore
SGS | South Georgia and the South Sandwich Islands
SHN | Saint Helena, Ascension and Tristan da Cunha
SJM | Svalbard and Jan Mayen
SLB | Solomon Islands
SLE | Sierra Leone
SLV | El Salvador
SMR | San Marino
SOM | Somalia
SPM | Saint Pierre and Miquelon
SRB | Serbia
SSD | South Sudan
STP | Sao Tome and Principe
SUR | Suriname
SVK | Slovakia
SVN | Slovenia
SWE | Sweden
SWZ | Swaziland
SXM | Sint Maarten (Dutch part)
SYC | Seychelles
SYR | Syrian Arab Republic
TCA | Turks and Caicos Islands
TCD | Chad
TGO | Togo
THA | Thailand
TJK | Tajikistan
TKL | Tokelau

TKM | Turkmenistan
TLS | Timor-Leste
TON | Tonga
TTO | Trinidad and Tobago
TUN | Tunisia
TUR | Turkey
TUV | Tuvalu
TWN | Taiwan, Province of China
TZA | Tanzania, United Republic of
UGA | Uganda
UKR | Ukraine
UMI | United States Minor Outlying Islands
URY | Uruguay
USA | United States of America
UZB | Uzbekistan
VAT | Holy See
VCT | Saint Vincent and the Grenadines
VEN | Venezuela, Bolivarian Republic of
VGB | Virgin Islands, British
VIR | Virgin Islands, U.S.
VNM | Viet Nam
VUT | Vanuatu
WLF | Wallis and Futuna
WSM | Samoa
YEM | Yemen
ZAF | South Africa
ZMB | Zambia
ZWE | Zimbabwe

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.1.662 Value Set http://hl7.org/fhir/ValueSet/object-lifecycle-events

URL: https://hl7.org/fhir/R4/valueset-object-lifecycle-events.html
Security Work Group | Maturity Level: 3 | Trial Use | Use Context: Any

This is a value set defined by the FHIR project.

Summary

Defining URL: | http://hl7.org/fhir/ValueSet/object-lifecycle-events
Version: | 4.0.1
Name: | ObjectLifecycleEvents
Title:
Definition: | This example FHIR value set is comprised of lifecycle event codes. The FHIR Actor value set is based on DICOM Audit Message, ParticipantObjectDataLifeCycle; ISO Standard, TS 21089-2017;
Committee: | Security Work Group
OID: | (for OID based terminology systems)
Copyright: | This is an example FHIR value set based on ASTM Standard, E1762-95 (2013) HL7 RoleClass OID 2.16.840.1.113883.5.110, HL7 Role Code 2.16.840.1.113883.5.111, HL7 ParticipationType OID: 2.16.840.1.113883.5.90, HL7 ParticipationFunction codes at OID: 2.16.840.1.113883.5.88, and HL7 Security and Privacy Domain Analysis Model roles classes. These codes are excerpted from Digital Imaging and Communications in Medicine (DICOM) Standard, Part 16: Content Mapping Resource, Copyright © 2011 by the National Electrical Manufacturers Association.
Source Resource | XML / JSON

This value set is used in the following places:

• Resource: AuditEvent.entity.lifecycle (Coding / Extensible)

4.4.1.662.1 Content Logical Definition

This value set includes codes from the following code systems:

• Include all codes defined in http://terminology.hl7.org/CodeSystem/dicom-audit-lifecycle
• Include all codes defined in http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle

4.4.1.662.2 Expansion

This expansion generated 01 Nov 2019

This value set contains 42 concepts

Expansion based on http://terminology.hl7.org/CodeSystem/dicom-audit-lifecycle version 20100826

Code	System	Display	Definition
1	http://terminology.hl7.org/CodeSystem/dicom-audit-lifecycle	Origination / Creation	
2	http://terminology.hl7.org/CodeSystem/dicom-audit-lifecycle	Import / Copy	
3	http://terminology.hl7.org/CodeSystem/dicom-audit-lifecycle	Amendment	
4	http://terminology.hl7.org/CodeSystem/dicom-audit-lifecycle	Verification	
5	http://terminology.hl7.org/CodeSystem/dicom-audit-lifecycle	Translation	
6	http://terminology.hl7.org/CodeSystem/dicom-audit-lifecycle	Access / Use	
7	http://terminology.hl7.org/CodeSystem/dicom-audit-lifecycle	De-identification	
8	http://terminology.hl7.org/CodeSystem/dicom-audit-lifecycle	Aggregation / summarization / derivation	
9	http://terminology.hl7.org/CodeSystem/dicom-audit-lifecycle	Report	
10	http://terminology.hl7.org/CodeSystem/dicom-audit-lifecycle	Export	
11	http://terminology.hl7.org/CodeSystem/dicom-audit-lifecycle	Disclosure	
12	http://terminology.hl7.org/CodeSystem/dicom-audit-lifecycle	Receipt of disclosure	
13	http://terminology.hl7.org/CodeSystem/dicom-audit-lifecycle	Archiving	
14	http://terminology.hl7.org/CodeSystem/dicom-audit-lifecycle	Logical deletion	
15	http://terminology.hl7.org/CodeSystem/dicom-audit-lifecycle	Permanent erasure / Physical destruction	
access	http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle	Access/View Record Lifecycle Event	Occurs when an agent causes the system to obtain and open a record entry for inspection or review.
hold	http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle	Add Legal Hold Record Lifecycle Event	Occurs when an agent causes the system to tag or otherwise indicate special access management and suspension of record entry deletion/destruction, if deemed relevant to a lawsuit or which are reasonably anticipated to be relevant or to fulfill organizational policy under the legal doctrine of “duty to preserve”.
amend	http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle	Amend (Update) Record Lifecycle Event	Occurs when an agent makes any change to record entry content currently residing in storage considered permanent (persistent).
archive	http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle	Archive Record Lifecycle Event	Occurs when an agent causes the system to create and move archive artifacts containing record entry content, typically to long-term offline storage.
attest	http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle	Attest Record Lifecycle Event	Occurs when an agent causes the system to capture the agent’s digital signature (or equivalent indication) during formal validation of record entry content.
decrypt	http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle	Decrypt Record Lifecycle Event	Occurs when an agent causes the system to decode record entry content from a cipher.
deidentify	http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle	De-Identify (Anononymize) Record Lifecycle Event	Occurs when an agent causes the system to scrub record entry content to reduce the association between a set of identifying data and the data subject in a way that might or might not be reversible.
deprecate	http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle	Deprecate Record Lifecycle Event	Occurs when an agent causes the system to tag record entry(ies) as obsolete, erroneous or untrustworthy, to warn against its future use.
destroy	http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle	Destroy/Delete Record Lifecycle Event	Occurs when an agent causes the system to permanently erase record entry content from the system.
disclose	http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle	Disclose Record Lifecycle Event	Occurs when an agent causes the system to release, transfer, provision access to, or otherwise divulge record entry content.
encrypt	http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle	Encrypt Record Lifecycle Event	Occurs when an agent causes the system to encode record entry content in a cipher.
extract	http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle	Extract Record Lifecycle Event	Occurs when an agent causes the system to selectively pull out a subset of record entry content, based on explicit criteria.
link	http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle	Link Record Lifecycle Event	Occurs when an agent causes the system to connect related record entries.
merge	http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle	Merge Record Lifecycle Event	Occurs when an agent causes the system to combine or join content from two or more record entries, resulting in a single logical record entry.
originate	http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle	Originate/Retain Record Lifecycle Event	Occurs when an agent causes the system to: a) initiate capture of potential record content, and b) incorporate that content into the storage considered a permanent part of the health record.
pseudonymize	http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle	Pseudonymize Record Lifecycle Event	Occurs when an agent causes the system to remove record entry content to reduce the association between a set of identifying data and the data subject in a way that may be reversible.
reactivate	http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle	Re-activate Record Lifecycle Event	Occurs when an agent causes the system to recreate or restore full status to record entries previously deleted or deprecated.
receive	http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle	Receive/Retain Record Lifecycle Event	Occurs when an agent causes the system to a) initiate capture of data content from elsewhere, and b) incorporate that content into the storage considered a permanent part of the health record.
reidentify	http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle	Re-identify Record Lifecycle Event	Occurs when an agent causes the system to restore information to data that allows identification of information source and/or information subject.
unhold	http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle	Remove Legal Hold Record Lifecycle Event	Occurs when an agent causes the system to remove a tag or other cues for special access management had required to fulfill organizational policy under the legal doctrine of “duty to preserve”.
report	http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle	Report (Output) Record Lifecycle Event	Occurs when an agent causes the system to produce and deliver record entry content in a particular form and manner.
restore	http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle	Restore Record Lifecycle Event	Occurs when an agent causes the system to recreate record entries and their content from a previous created archive artefact.
transform	http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle	Transform/Translate Record Lifecycle Event	Occurs when an agent causes the system to change the form, language or code system used to represent record entry content.

transmit | http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle | Transmit Record Lifecycle Event | Occurs when an agent causes the system to send record entry content from one (EHR/PHR/other) system to another.
unlink | http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle | Unlink Record Lifecycle Event | Occurs when an agent causes the system to disconnect two or more record entries previously connected, rendering them separate (disconnected) again.
unmerge | http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle | Unmerge Record Lifecycle Event | Occurs when an agent causes the system to reverse a previous record entry merge operation, rendering them separate again.
verify | http://terminology.hl7.org/CodeSystem/iso-21089-lifecycle | Verify Record Lifecycle Event | Occurs when an agent causes the system to confirm compliance of data or data objects with regulations, requirements, specifications, or other imposed conditions based on organizational policy.

See the full registry of value sets defined as part of FHIR.

Explanation of the columns that may appear on this page:

Lvl | A few code lists that FHIR defines are hierarchical - each code is assigned a level. For value sets, levels are mostly used to organize codes for user convenience, but may follow code system hierarchy - see Code System for further information
Source | The source of the definition of the code (when the value set draws in codes defined elsewhere)
Code | The code (used as the code in the resource instance). If the code is in italics, this indicates that the code is not selectable ('Abstract')
Display | The display (used in the display element of a Coding). If there is no display, implementers should not simply display the code, but map the concept into their application
Definition | An explanation of the meaning of the concept
Comments | Additional notes about how to use the code

4.4.2.1 HL7 v3 Code System AdministrativeGender

URL: https://hl7.org/fhir/R4/v3/AdministrativeGender/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-AdministrativeGender) is defined as part of HL7 v3. Related FHIR content: AdministrativeGender.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-AdministrativeGender
Version: | 2018-08-12
Name: | v3.AdministrativeGender
Title: | v3 Code System AdministrativeGender
Definition: | The gender of a person used for adminstrative purposes (as opposed to clinical gender)
OID: | 2.16.840.1.113883.5.1 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System AdministrativeGender (The gender of a person used for adminstrative purposes (as opposed to clinical gender))

4.4.2.1.1 Code System Content

• All Languages
• English
• Nederlands (Dutch)

This code system http://terminology.hl7.org/CodeSystem/v3-AdministrativeGender defines the following codes:

Code | Display | Definition
F | en: Female nl: Vrouw | en: Female nl: Vrouwelijk
M | en: Male nl: Man | en: Male nl: Man
UN | en: Undifferentiated nl: Ongedifferentieerd | en: The gender of a person could not be uniquely defined as male or female, such as hermaphrodite. nl: Het geslacht van een persoon kan niet uniek worden gedefinieerd als man of vrouw, zoals een hermafrodiet.

This code system http://terminology.hl7.org/CodeSystem/v3-AdministrativeGender defines the following codes:

Code | Display | Definition
F | Female | Female
M | Male | Male
UN | Undifferentiated | The gender of a person could not be uniquely defined as male or female, such as hermaphrodite.

Definition: Het geslacht van een persoon bedoeld voor administratieve doeleinden (in tegenstelling tot klinisch geslacht)

Dit code systeem http://terminology.hl7.org/CodeSystem/v3-AdministrativeGender definieert de volgende codes:

Code | Display | Definition
F | Vrouw | Vrouwelijk
M | Man | Man
UN | Ongedifferentieerd | Het geslacht van een persoon kan niet uniek worden gedefinieerd als man of vrouw, zoals een hermafrodiet.

4.4.2.2 HL7 v3 Code System MaritalStatus

URL: https://hl7.org/fhir/R4/v3/MaritalStatus/cs.html
Vocabulary Work Group | Maturity Level: N | Normative (from v4.0.0) | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-MaritalStatus) is defined as part of HL7 v3. Related FHIR content: MaritalStatus.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-MaritalStatus
Version: | 2018-08-12
Name: | v3.MaritalStatus
Title: | v3 Code System MaritalStatus
Definition: | No description supplied * * * Open Issue: The specific meanings of these codes can vary somewhat by jurisdiction and implementation so caution should be used when determining equivalency.
OID: | 2.16.840.1.113883.5.2 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System MaritalStatus (* * * No description supplied * * * Open Issue: The specific meanings of these codes can vary somewhat by jurisdiction and implementation so caution should be used when determining equivalency.)

• ValueSet: MaritalStatus (This value set defines the set of codes that can be used to indicate the marital status of a person.)

• ValueSet: MaritalStatus (This value set defines the set of codes that can be used to indicate the marital status of a person.)

4.4.2.2.1 Code System Content

Release Date: 2018-08-12

Level	Code	Display	Definition
1	A	Annulled	Marriage contract has been declared null and to not have existed
1	D	Divorced	Marriage contract has been declared dissolved and inactive
1	I	Interlocutory	Subject to an Interlocutory Decree.
1	L	Legally Separated	
1	M	Married	A current marriage contract is active
1	P	Polygamous	More than 1 current spouse
1	S	Never Married	No marriage contract has ever been entered
1	T	Domestic partner	Person declares that a domestic partner relationship exists.
1	U	unmarried	Currently not in a marriage contract.
1	W	Widowed	The spouse has died

4.4.2.4 HL7 v3 Code System ActCode

URL: https://hl7.org/fhir/R4/v3/ActCode/cs.html
Vocabulary Work Group | Maturity Level: 1 | Draft | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-ActCode) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-ActCode
Version: | 2018-08-12
Name: | v3.ActCode
Title: | v3 Code System ActCode
Definition: | A code specifying the particular kind of Act that the Act-instance represents within its class. Constraints: The kind of Act (e.g. physical examination, serum potassium, inpatient encounter, charge financial transaction, etc.) is specified with a code from one of several, typically external, coding systems. The coding system will depend on the class of Act, such as LOINC for observations, etc. Conceptually, the Act.code must be a specialization of the Act.classCode. This is why the structure of ActClass domain should be reflected in the superstructure of the ActCode domain and then individual codes or externally referenced vocabularies subordinated under these domains that reflect the ActClass structure. Act.classCode and Act.code are not modifiers of each other but the Act.code concept should really imply the Act.classCode concept. For a negative example, it is not appropriate to use an Act.code "potassium" together with and Act.classCode for "laboratory observation" to somehow mean "potassium laboratory observation" and then use the same Act.code for "potassium" together with Act.classCode for "medication" to mean "substitution of potassium". This mutually modifying use of Act.code and Act.classCode is not permitted.
OID: | 2.16.840.1.113883.5.4 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: V3 Value SetRefrainPolicy (Conveys prohibited actions which an information custodian, receiver, or user is not permitted to perform unless otherwise authorized or permitted under specified circumstances. Examples: prohibit redisclosure without consent directive)

• ValueSet: V3 Value SetCompartment (A named tag set for metadata used to populate a security category label field that "segments" an IT resource per policy by indicating that access and use is restricted to members of a defined community or project. (HL7 Healthcare Privacy and Security Classification System) Usage Note: This is the healthcare analog to the US Intelligence Community's concept of a Special Access Program. Compartment codes may be used in as a field value in an initiator's clearance to indicate permission to access and use an IT Resource with a security label having the same compartment value in security category label field. Map: Aligns with ISO 2382-8 definition of Compartment - "A division of data into isolated blocks with separate security controls for the purpose of reducing risk.")

• ValueSet: v3 Code System ActCode (A code specifying the particular kind of Act that the Act-instance represents within its class. Constraints: The kind of Act (e.g. physical examination, serum potassium, inpatient encounter, charge financial transaction, etc.) is specified with a code from one of several, typically external, coding systems. The coding system will depend on the class of Act, such as LOINC for observations, etc. Conceptually, the Act.code must be a specialization of the Act.classCode. This is why the structure of ActClass domain should be reflected in the superstructure of the ActCode domain and then individual codes or externally referenced vocabularies subordinated under these domains that reflect the ActClass structure. Act.classCode and Act.code are not modifiers of each other but the Act.code concept should really imply the Act.classCode concept. For a negative example, it is not appropriate to use an Act.code "potassium" together with and Act.classCode for "laboratory observation" to somehow mean "potassium laboratory observation" and then use the same Act.code for "potassium" together with Act.classCode for "medication" to mean "substitution of potassium". This mutually modifying use of Act.code and Act.classCode is not permitted.)

• ValueSet: V3 Value SetActInvoiceGroupCode (Type of invoice element that is used to assist in describing an Invoice that is either submitted for adjudication or for which is returned on adjudication results. Invoice elements of this type signify a grouping of one or more children (detail) invoice elements. They do not have intrinsic costing associated with them, but merely reflect the sum of all costing for it's immediate children invoice elements.)

• ValueSet: V3 Value SetObligationPolicy (Conveys the mandated workflow action that an information custodian, receiver, or user must perform. Examples: encrypt Usage Note: Per OASIS XACML, an obligation is an operation specified in a policy or policy that is performed in conjunction with the enforcement of an access control decision.)

• ValueSet: V3 Value SetActIncidentCode (Set of codes indicating the type of incident or accident.)

• ValueSet: V3 Value SetActEncounterCode (Domain provides codes that qualify the ActEncounterClass (ENC))

• ValueSet: Detected Issue Mitigation Action (Kinds of mitigating actions and observations that can be associated with a detected issue or contraindication, such as 'added concurrent therapy', 'prior therapy documented', etc.)

• ValueSet: Provenance activity type (This value set contains representative Activity Type codes, which includes codes from the HL7 DocumentCompletion, ActStatus, and DataOperations code system, W3C PROV-DM and PROV-N concepts and display names, several HL7 Lifecycle Event codes for which there are agreed upon definitions, and non-duplicated codes from the HL7 Security and Privacy Ontology Operations codes.)

• ValueSet: V3 Value SetInformationSensitivityPolicy (Sensitivity codes are not useful for interoperability outside of a policy domain because sensitivity policies are typically localized and vary drastically across policy domains even for the same information category because of differing organizational business rules, security policies, and jurisdictional requirements. For example, an "employee" sensitivity code would make little sense for use outside of a policy domain. "Taboo" would rarely be useful outside of a policy domain unless there are jurisdictional requirements requiring that a provider disclose sensitive information to a patient directly. Sensitivity codes may be more appropriate in a legacy system's Master Files in order to notify those who access a patient's orders and observations about the sensitivity policies that apply. Newer systems may have a security engine that uses a sensitivity policy's criteria directly. The specializable Sensitivity Act.code may be useful in some scenarious if used in combination with a sensitivity identifier and/or Act.title.)

• ValueSet: V3 Value SetActPharmacySupplyType (Identifies types of dispensing events)

• ValueSet: Provenance activity type (This value set contains representative Activity Type codes, which includes codes from the HL7 DocumentCompletion, ActStatus, and DataOperations code system, W3C PROV-DM and PROV-N concepts and display names, several HL7 Lifecycle Event codes for which there are agreed upon definitions, and non-duplicated codes from the HL7 Security and Privacy Ontology Operations codes.)

• ValueSet: V3 Value SetActTaskCode (Description: A task or action that a user may perform in a clinical information system.)

• ValueSet: Account Types (This examples value set defines the set of codes that can be used to represent the type of an account.)

• ValueSet: V3 Value SetObservationType (Identifies the kinds of observations that can be performed)

• ValueSet: Detected Issue Category (Kinds of issues or contraindications, such as 'drug-drug interaction', 'duplicate therapy', etc.)

• ValueSet: Account Types (This examples value set defines the set of codes that can be used to represent the type of an account.)

• ValueSet: V3 Value SetActCoverageTypeCode (Definition: Set of codes indicating the type of insurance policy or program that pays for the cost of benefits provided to covered parties.)

• ValueSet: V3 Value SetActConsentType (Definition: The type of consent directive, e.g., to consent or dissent to collect, access, or use in specific ways within an EHRS or for health information exchange; or to disclose health information for purposes such as research.)

• ValueSet: Detected Issue Mitigation Action (Kinds of mitigating actions and observations that can be associated with a detected issue or contraindication, such as 'added concurrent therapy', 'prior therapy documented', etc.)

• ValueSet: V3 Value SetActConsentDirective (ActConsentDirective codes are used to specify the type of Consent Directive to which a Consent Directive Act conforms.)

• ValueSet: V3 Value SetSecurityPolicy (Types of security policies that further specify the ActClassPolicy value set. Examples: encrypt prohibit redisclosure without consent directive)

• ValueSet: Detected Issue Category (Kinds of issues or contraindications, such as 'drug-drug interaction', 'duplicate therapy', etc.)

4.4.2.4.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition

1 | (_ActAccountCode) Abstract | | An account represents a grouping of financial transactions that are tracked and reported together with a single balance. Examples of account codes (types) are Patient billing accounts (collection of charges), Cost centers; Cash.

2 | ACCTRECEIVABLE | account receivable | An account for collecting charges, reversals, adjustments and payments, including deductibles, copayments, coinsurance (financial transactions) credited or debited to the account receivable account for a patient's encounter.

2 | CASH | Cash

2 | CC | credit card | Description: Types of advance payment to be made on a plastic card usually issued by a financial institution used of purchasing services and/or products.

3 | AE | American Express

3 | DN | Diner's Club

3 | DV | Discover Card

3 | MC | Master Card

3 | V | Visa

2 | PBILLACCT | patient billing account | An account representing charges and credits (financial transactions) for a patient's encounter.

1 | (_ActAdjudicationCode) Abstract | | Includes coded responses that will occur as a result of the adjudication of an electronic invoice at a summary level and provides guidance on interpretation of the referenced adjudication results.

2 | (_ActAdjudicationGroupCode) Abstract | | Catagorization of grouping criteria for the associated transactions and/or summary (totals, subtotals).

3 | CONT | contract | Transaction counts and value totals by Contract Identifier.

3 | DAY | day | Transaction counts and value totals for each calendar day within the date range specified.

3 | LOC | location | Transaction counts and value totals by service location (e.g clinic).

3 | MONTH | month | Transaction counts and value totals for each calendar month within the date range specified.

3 | PERIOD | period | Transaction counts and value totals for the date range specified.

3 | PROV | provider | Transaction counts and value totals by Provider Identifier.

3 | WEEK | week | Transaction counts and value totals for each calendar week within the date range specified.

3 | YEAR | year | Transaction counts and value totals for each calendar year within the date range specified.

2 | AA | adjudicated with adjustments | The invoice element has been accepted for payment but one or more adjustment(s) have been made to one or more invoice element line items (component charges). Also includes the concept 'Adjudicate as zero' and items not covered under a particular Policy. Invoice element can be reversed (nullified). Recommend that the invoice element is saved for DUR (Drug Utilization Reporting).

3 | ANF | adjudicated with adjustments and no financial impact | The invoice element has been accepted for payment but one or more adjustment(s) have been made to one or more invoice element line items (component charges) without changing the amount. Invoice element can be reversed (nullified). Recommend that the invoice element is saved for DUR (Drug Utilization Reporting).

2 | AR | adjudicated as refused | The invoice element has passed through the adjudication process but payment is refused due to one or more reasons. Includes items such as patient not covered, or invoice element is not constructed according to payer rules (e.g. 'invoice submitted too late'). If one invoice element line item in the invoice element structure is rejected, the remaining line items may not be adjudicated and the complete group is treated as rejected. A refused invoice element can be forwarded to the next payer (for Coordination of Benefits) or modified and resubmitted to refusing payer. Invoice element cannot be reversed (nullified) as there is nothing to reverse. Recommend that the invoice element is not saved for DUR (Drug Utilization Reporting).

2 | AS | adjudicated as submitted | The invoice element was/will be paid exactly as submitted, without financial adjustment(s). If the dollar amount stays the same, but the billing codes have been amended or financial adjustments have been applied through the adjudication process, the invoice element is treated as "Adjudicated with Adjustment". If information items are included in the adjudication results that do not affect the monetary amounts paid, then this is still Adjudicated as Submitted (e.g. 'reached Plan Maximum on this Claim'). Invoice element can be reversed (nullified). Recommend that the invoice element is saved for DUR (Drug Utilization Reporting).

1 | (_ActAdjudicationResultActionCode) Abstract | | Actions to be carried out by the recipient of the Adjudication Result information.
2 | DISPLAY | Display | The adjudication result associated is to be displayed to the receiver of the adjudication result.
2 | FORM | Print on Form | The adjudication result associated is to be printed on the specified form, which is then provided to the covered party.
1 | (_ActBillableModifierCode) Abstract | | Definition:An identifying modifier code for healthcare interventions or procedures.
2 | CPTM | CPT modifier codes | Description:CPT modifier codes are found in Appendix A of CPT 2000 Standard Edition.
2 | HCPCSA | HCPCS Level II and Carrier-assigned | Description:HCPCS Level II (HCFA-assigned) and Carrier-assigned (Level III) modifiers are reported in Appendix A of CPT 2000 Standard Edition and in the Medicare Bulletin.
1 | (_ActBillingArrangementCode) Abstract | | The type of provision(s) made for reimbursing for the deliver of healthcare services and/or goods provided by a Provider, over a specified period.
2 | BLK | block funding | A billing arrangement where a Provider charges a lump sum to provide a prescribed group (volume) of services to a single patient which occur over a period of time. Services included in the block may vary. This billing arrangement is also known as Program of Care for some specific Payors and Program Fees for other Payors.
2 | CAP | capitation funding | A billing arrangement where the payment made to a Provider is determined by analyzing one or more demographic attributes about the persons/patients who are enrolled with the Provider (in their practice).
2 | CONTF | contract funding | A billing arrangement where a Provider charges a lump sum to provide a particular volume of one or more interventions/procedures or groups of interventions/procedures.
2 | FINBILL | financial | A billing arrangement where a Provider charges for non-clinical items. This includes interest in arrears, mileage, etc. Clinical content is not included in Invoices submitted with this type of billing arrangement.
2 | ROST | roster funding | A billing arrangement where funding is based on a list of individuals registered as patients of the Provider.
2 | SESS | sessional funding | A billing arrangement where a Provider charges a sum to provide a group (volume) of interventions/procedures to one or more patients within a defined period of time, typically on the same date. Interventions/procedures included in the session may vary.
2 | FFS | fee for service | A billing arrangement where a Provider charges a separate fee for each intervention/procedure/event or product. Fee for Service is used when an individual intervention/procedure/event is used for billing purposes. In other words, fees are associated with the intervention/procedure/event. For example, a specific CCI (Canadian Classification of Interventions) code has an associated fee and is used for billing purposes.
3 | FFPS | first fill, part fill, partial strength | A first fill where the quantity supplied is less than one full repetition of the ordered amount. (e.g. If the order was 90 tablets, 3 refills, a partial fill might be for only 30 tablets.) and also where the strength supplied is less than the ordered strength (e.g. 10mg for an order of 50mg where a subsequent fill will dispense 40mg tablets)
3 | FFCS | first fill complete, partial strength | A first fill where the quantity supplied is equal to one full repetition of the ordered amount. (e.g. If the order was 90 tablets, 3 refills, a complete fill would be for the full 90 tablets) and also where the strength supplied is less than the ordered strength (e.g. 10mg for an order of 50mg where a subsequent fill will dispense 40mg tablets).
3 | TFS | trial fill partial strength | A fill where a small portion is provided to allow for determination of the therapy effectiveness and patient tolerance and also where the strength supplied is less than the ordered strength (e.g. 10mg for an order of 50mg where a subsequent fill will dispense 40mg tablets).
1 | (_ActBoundedROICode) Abstract | | Type of bounded ROI.
2 | ROIFS | fully specified ROI | A fully specified bounded Region of Interest (ROI) delineates a ROI in which only those dimensions participate that are specified by boundary criteria, whereas all other dimensions are excluded. For example a ROI to mark an episode of "ST elevation" in a subset of the EKG leads V2, V3, and V4 would include 4 boundaries, one each for time, V2, V3, and V4.
2 | ROIPS | partially specified ROI | A partially specified bounded Region of Interest (ROI) specifies a ROI in which at least all values in the dimensions specified by the boundary criteria participate. For example, if an episode of ventricular fibrillations (VFib) is observed, it usually doesn't make sense to exclude any EKG leads from the observation and the partially specified ROI would contain only one boundary for time indicating the time interval where VFib was observed.
1 | (_ActCareProvisionCode) Abstract | | Description:The type and scope of responsibility taken-on by the performer of the Act for a specific subject of care.
2 | (_ActCredentialedCareCode) Abstract | | Description:The type and scope of legal and/or professional responsibility taken-on by the performer of the Act for a specific subject of care as described by a credentialing agency, i.e. government or non-government agency. Failure in executing this Act may result in loss of credential to the person or organization who participates as performer of the Act. Excludes employment agreements. Example:Hospital license; physician license; clinic accreditation.
3 | (_ActCredentialedCareProvisionPersonCode) Abstract | | Description:The type and scope of legal and/or professional responsibility taken-on by the performer of the Act for a specific subject of care as described by an agency for credentialing individuals.
4 | CACC | certified anatomic pathology and clinical pathology care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CAIC | certified allergy and immunology care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CAMC | certified aerospace medicine care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CANC | certified anesthesiology care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CAPC | certified anatomic pathology care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CBGC | certified clinical biochemical genetics care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CCCC | certified clinical cytogenetics care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CCGC | certified clinical genetics (M.D.) care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CCPC | certified clinical pathology care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CCSC | certified colon and rectal surgery care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.

4 | CDEC | certified dermatology care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CDRC | certified diagnostic radiology care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CEMC | certified emergency medicine care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CFPC | certified family practice care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CIMC | certified internal medicine care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CMGC | certified clinical molecular genetics care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CNEC | certified neurology care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board
4 | CNMC | certified nuclear medicine care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CNQC | certified neurology with special qualifications in child neurology care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CNSC | certified neurological surgery care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | COGC | certified obstetrics and gynecology care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | COMC | certified occupational medicine care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | COPC | certified ophthalmology care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | COSC | certified orthopaedic surgery care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | COTC | certified otolaryngology care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CPEC | certified pediatrics care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CPGC | certified Ph.D. medical genetics care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CPHC | certified public health and general preventive medicine care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CPRC | certified physical medicine and rehabilitation care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CPSC | certified plastic surgery care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CPYC | certified psychiatry care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CROC | certified radiation oncology care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CRPC | certified radiological physics care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CSUC | certified surgery care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CTSC | certified thoracic surgery care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CURC | certified urology care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | CVSC | certified vascular surgery care | Description:Scope of responsibility taken on for specialty care as defined by the respective Specialty Board.
4 | LGPC | licensed general physician care | Description:Scope of responsibility taken-on for physician care of a patient as defined by a governmental licensing agency.
3 | (_ActCredentialedCareProvisionProgramCode) Abstract | | Description:The type and scope of legal and/or professional responsibility taken-on by the performer of the Act for a specific subject of care as described by an agency for credentialing programs within organizations.
4 | AALC | accredited assisted living care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the respective accreditation agency.
4 | AAMC | accredited ambulatory care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the respective accreditation agency.
4 | ABHC | accredited behavioral health care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the respective accreditation agency.
4 | ACAC | accredited critical access hospital care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the respective accreditation agency.
4 | ACHC | accredited hospital care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the respective accreditation agency.
4 | AHOC | accredited home care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the respective accreditation agency.
4 | ALTC | accredited long term care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the respective accreditation agency.
4 | AOSC | accredited office-based surgery care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the respective accreditation agency.

4 | CACS | certified acute coronary syndrome care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CAMI | certified acute myocardial infarction care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CAST | certified asthma care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CBAR | certified bariatric surgery care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CCAD | certified coronary artery disease care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CCAR | certified cardiac care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CDEP | certified depression care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CDGD | certified digestive/gastrointestinal disorders care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CDIA | certified diabetes care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CEPI | certified epilepsy care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CFEL | certified frail elderly care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CHFC | certified heart failure care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CHRO | certified high risk obstetrics care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CHYP | certified hyperlipidemia care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CMIH | certified migraine headache care | Description:.
4 | CMSC | certified multiple sclerosis care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | COJR | certified orthopedic joint replacement care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CONC | certified oncology care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | COPD | certified chronic obstructive pulmonary disease care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CORT | certified organ transplant care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CPAD | certified parkinsons disease care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CPND | certified pneumonia disease care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CPST | certified primary stroke center care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CSDM | certified stroke disease management care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CSIC | certified sickle cell care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CSLD | certified sleep disorders care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CSPT | certified spine treatment care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CTBU | certified trauma/burn center care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CVDC | certified vascular diseases care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CWMA | certified wound management care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
4 | CWOH | certified women's health care | Description:Scope of responsibility taken on by an organization for care of a patient as defined by the disease management certification agency.
2 | (_ActEncounterCode) Abstract | | Domain provides codes that qualify the ActEncounterClass (ENC)
3 | AMB | ambulatory | A comprehensive term for health care provided in a healthcare facility (e.g. a practitioneraTMs office, clinic setting, or hospital) on a nonresident basis. The term ambulatory usually implies that the patient has come to the location and is not assigned to a bed. Sometimes referred to as an outpatient encounter.
3 | EMER | emergency | A patient encounter that takes place at a dedicated healthcare service delivery location where the patient receives immediate evaluation and treatment, provided until the patient can be discharged or responsibility for the patient's care is transferred elsewhere (for example, the patient could be admitted as an inpatient or transferred to another facility.)
3 | FLD | field | A patient encounter that takes place both outside a dedicated service delivery location and outside a patient's residence. Example locations might include an accident site and at a supermarket.
3 | HH | home health | Healthcare encounter that takes place in the residence of the patient or a designee
3 | IMP | inpatient encounter | A patient encounter where a patient is admitted by a hospital or equivalent facility, assigned to a location where patients generally stay at least overnight and provided with room, board, and continuous nursing service.
4 | ACUTE | inpatient acute | An acute inpatient encounter.
4 | NONAC | inpatient non-acute | Any category of inpatient encounter except 'acute'

3 | OBSENC | observation encounter | An encounter where the patient usually will start in different encounter, such as one in the emergency department (EMER) but then transition to this type of encounter because they require a significant period of treatment and monitoring to determine whether or not their condition warrants an inpatient admission or discharge. In the majority of cases the decision about admission or discharge will occur within a time period determined by local, regional or national regulation, often between 24 and 48 hours.
3 | PRENC | pre-admission | A patient encounter where patient is scheduled or planned to receive service delivery in the future, and the patient is given a pre-admission account number. When the patient comes back for subsequent service, the pre-admission encounter is selected and is encapsulated into the service registration, and a new account number is generated. Usage Note: This is intended to be used in advance of encounter types such as ambulatory, inpatient encounter, virtual, etc.
3 | SS | short stay | An encounter where the patient is admitted to a health care facility for a predetermined length of time, usually less than 24 hours.
3 | VR | virtual | A patient encounter where the patient and the practitioner(s) are not in the same physical location. Examples include telephone conference, email exchange, robotic surgery, and televideo conference.
2 | (_ActMedicalServiceCode) Abstract | | General category of medical service provided to the patient during their encounter.
3 | ALC | Alternative Level of Care | Provision of Alternate Level of Care to a patient in an acute bed. Patient is waiting for placement in a long-term care facility and is unable to return home.
3 | CARD | Cardiology | Provision of diagnosis and treatment of diseases and disorders affecting the heart
3 | CHR | Chronic | Provision of recurring care for chronic illness.
3 | DNTL | Dental | Provision of treatment for oral health and/or dental surgery.
3 | DRGRHB | Drug Rehab | Provision of treatment for drug abuse.
3 | GENRL | General | General care performed by a general practitioner or family doctor as a responsible provider for a patient.
3 | MED | Medical | Provision of diagnostic and/or therapeutic treatment.
3 | OBS | Obstetrics | Provision of care of women during pregnancy, childbirth and immediate postpartum period. Also known as Maternity.
3 | ONC | Oncology | Provision of treatment and/or diagnosis related to tumors and/or cancer.
3 | PALL | Palliative | Provision of care for patients who are living or dying from an advanced illness.
3 | PED | Pediatrics | Provision of diagnosis and treatment of diseases and disorders affecting children.
3 | PHAR | Pharmaceutical | Pharmaceutical care performed by a pharmacist.
3 | PHYRHB | Physical Rehab | Provision of treatment for physical injury.
3 | PSYCH | Psychiatric | Provision of treatment of psychiatric disorder relating to mental illness.
3 | SURG | Surgical | Provision of surgical treatment.
1 | (_ActClaimAttachmentCategoryCode) Abstract | | Description: Coded types of attachments included to support a healthcare claim.
2 | AUTOATTCH | auto attachment | Description: Automobile Information Attachment
2 | DOCUMENT | document | Description: Document Attachment
2 | HEALTHREC | health record | Description: Health Record Attachment
2 | IMG | image attachment | Description: Image Attachment
2 | LABRESULTS | lab results | Description: Lab Results Attachment
2 | MODEL | model | Description: Digital Model Attachment
2 | WIATTCH | work injury report attachment | Description: Work Injury related additional Information Attachment
2 | XRAY | x-ray | Description: Digital X-Ray Attachment
1 | (_ActConsentType) Abstract | | Definition: The type of consent directive, e.g., to consent or dissent to collect, access, or use in specific ways within an EHRS or for health information exchange; or to disclose health information for purposes such as research.
2 | ICOL | information collection | Definition: Consent to have healthcare information collected in an electronic health record. This entails that the information may be used in analysis, modified, updated.
2 | IDSCL | information disclosure | Definition: Consent to have collected healthcare information disclosed.
2 | INFA | information access | Definition: Consent to access healthcare information.
3 | INFAO | access only | Definition: Consent to access or "read" only, which entails that the information is not to be copied, screen printed, saved, emailed, stored, re-disclosed or altered in any way. This level ensures that data which is masked or to which access is restricted will not be. Example: Opened and then emailed or screen printed for use outside of the consent directive purpose.
3 | INFASO | access and save only | Definition: Consent to access and save only, which entails that access to the saved copy will remain locked.
2 | IRDSCL | information redisclosure | Definition: Information re-disclosed without the patient's consent.
2 | RESEARCH | research information access | Definition: Consent to have healthcare information in an electronic health record accessed for research purposes.
3 | RSDID | de-identified information access | Definition: Consent to have de-identified healthcare information in an electronic health record that is accessed for research purposes, but without consent to re-identify the information under any circumstance.
3 | RSREID | re-identifiable information access | Definition: Consent to have de-identified healthcare information in an electronic health record that is accessed for research purposes re-identified under specific circumstances outlined in the consent. Example:: Where there is a need to inform the subject of potential health issues.
1 | (_ActContainerRegistrationCode) Abstract | | Constrains the ActCode to the domain of Container Registration
2 | ID | Identified | Used by one system to inform another that it has received a container.
2 | IP | In Position | Used by one system to inform another that the container is in position for specimen transfer (e.g., container removal from track, pipetting, etc.).
2 | L | Left Equipment | Used by one system to inform another that the container has been released from that system.
2 | M | Missing | Used by one system to inform another that the container did not arrive at its next expected location.
2 | O | In Process | Used by one system to inform another that the specific container is being processed by the equipment. It is useful as a response to a query about Container Status, when the specific step of the process is not relevant.
2 | R | Process Completed | Status is used by one system to inform another that the processing has been completed, but the container has not been released from that system.
2 | X | Container Unavailable | Used by one system to inform another that the container is no longer available within the scope of the system (e.g., tube broken or discarded).
1 | (_ActControlVariable) Abstract | | An observation form that determines parameters or attributes of an Act. Examples are the settings of a ventilator machine as parameters of a ventilator treatment act; the controls on dillution factors of a chemical

analyzer as a parameter of a laboratory observation act; the settings of a physiologic measurement assembly (e.g., time skew) or the position of the body while measuring blood pressure. Control variables are forms of observations because just as with clinical observations, the Observation.code determines the parameter and the Observation.value assigns the value. While control variables sometimes can be observed (by noting the control settings or an actually measured feedback loop) they are not primary observations, in the sense that a control variable without a primary act is of no use (e.g., it makes no sense to record a blood pressure position without recording a blood pressure, whereas it does make sense to record a systolic blood pressure without a diastolic blood pressure).

2 | AUTO | auto-repeat permission | Specifies whether or not automatic repeat testing is to be initiated on specimens.

2 | ENDC | endogenous content | A baseline value for the measured test that is inherently contained in the diluent. In the calculation of the actual result for the measured test, this baseline value is normally considered.

2 | REFLEX | reflex permission | Specifies whether or not further testing may be automatically or manually initiated on specimens.

1 | (_ActCoverageConfirmationCode) Abstract | | Response to an insurance coverage eligibility query or authorization request.

2 | (_ActCoverageAuthorizationConfirmationCode) Abstract | | Indication of authorization for healthcare service(s) and/or product(s). If authorization is approved, funds are set aside.

3 | AUTH | Authorized | Authorization approved and funds have been set aside to pay for specified healthcare service(s) and/or product(s) within defined criteria for the authorization.

3 | NAUTH | Not Authorized | Authorization for specified healthcare service(s) and/or product(s) denied.

2 | (_ActCoverageEligibilityConfirmationCode) Abstract | | Indication of eligibility coverage for healthcare service(s) and/or product(s).

3 | ELG | Eligible | Insurance coverage is in effect for healthcare service(s) and/or product(s).

3 | NELG | Not Eligible | Insurance coverage is not in effect for healthcare service(s) and/or product(s). May optionally include reasons for the ineligibility.

1 | (_ActCoverageLimitCode) Abstract | | Criteria that are applicable to the authorized coverage.

2 | (_ActCoverageQuantityLimitCode) Abstract | | Maximum amount paid or maximum number of services/products covered; or maximum amount or number covered during a specified time period under the policy or program.

3 | COVPRD | coverage period | Codes representing the time period during which coverage is available; or financial participation requirements are in effect.

3 | LFEMX | life time maximum | Definition: Maximum amount paid by payer or covered party; or maximum number of services or products covered under the policy or program during a covered party's lifetime.

3 | NETAMT | Net Amount | Maximum net amount that will be covered for the product or service specified.

3 | PRDMX | period maximum | Definition: Maximum amount paid by payer or covered party; or maximum number of services/products covered under the policy or program by time period specified by the effective time on the act.

3 | UNITPRICE | Unit Price | Maximum unit price that will be covered for the authorized product or service.

3 | UNITQTY | Unit Quantity | Maximum number of items that will be covered of the product or service specified.

2 | COVMX | coverage maximum | Definition: Codes representing the maximum coverate or financial participation requirements.

3 | LFEMX

3 | PRDMX

2 | (_ActCoveredPartyLimitCode) Abstract | | Codes representing the types of covered parties that may receive covered benefits under a policy or program.

1 | (_ActCoverageTypeCode) Abstract | | Definition: Set of codes indicating the type of insurance policy or program that pays for the cost of benefits provided to covered parties.

2 | (_ActInsurancePolicyCode) Abstract | | Set of codes indicating the type of insurance policy or other source of funds to cover healthcare costs.

3 | EHCPOL | extended healthcare | Private insurance policy that provides coverage in addition to other policies (e.g. in addition to a Public Healthcare insurance policy).

3 | HSAPOL | health spending account | Insurance policy that provides for an allotment of funds replenished on a periodic (e.g. annual) basis. The use of the funds under this policy is at the discretion of the covered party.

3 | AUTOPOL | automobile | Insurance policy for injuries sustained in an automobile accident. Will also typically covered non-named parties to the policy, such as pedestrians and passengers.

4 | COL | collision coverage policy | Definition: An automobile insurance policy under which the insurance company will cover the cost of damages to an automobile owned by the named insured that are caused by accident or intentionally by another party.

4 | UNINSMOT | uninsured motorist policy | Definition: An automobile insurance policy under which the insurance company will indemnify a loss for which another motorist is liable if that motorist is unable to pay because he or she is uninsured. Coverage under the policy applies to bodily injury damages only. Injuries to the covered party caused by a hit-and-run driver are also covered.

3 | PUBLICPOL | public healthcare | Insurance policy funded by a public health system such as a provincial or national health plan. Examples include BC MSP (British Columbia Medical Services Plan) OHIP (Ontario Health Insurance Plan), NHS (National Health Service).

4 | DENTPRG | dental program | Definition: A public or government health program that administers and funds coverage for dental care to assist program eligible who meet financial and health status criteria.

4 | DISEASEPRG | public health program | Definition: A public or government health program that administers and funds coverage for health and social services to assist program eligible who meet financial and health status criteria related to a particular disease. Example: Reproductive health, sexually transmitted disease, and end renal disease programs.

5 | CANPRG | women's cancer detection program | Definition: A program that provides low-income, uninsured, and underserved women access to timely, high-quality screening and diagnostic services, to detect breast and cervical cancer at the earliest stages. Example: To improve women's access to screening for breast and cervical cancers, Congress passed the Breast and Cervical Cancer Mortality Prevention Act of 1990, which guided CDC in creating the National Breast and Cervical Cancer Early Detection Program (NBCCEDP), which provides access to critical breast and cervical cancer screening services for underserved women in the United States. An estimated 7 to 10% of U.S. women of screening age are eligible to receive NBCCEDP services. Federal guidelines establish an eligibility baseline to direct services to uninsured and underinsured women at or below 250% of federal poverty level; ages 18 to 64 for cervical screening; ages 40 to 64 for breast screening.

5 | ENDRENAL | end renal program | Definition: A public or government program that administers publicly funded coverage of kidney dialysis and kidney transplant services. Example: In the U.S., the Medicare End-stage Renal Disease program (ESRD), the National Kidney Foundation (NKF) American Kidney Fund (AKF) The Organ Transplant Fund.

5 | HIVAIDS | HIV-AIDS program | Definition: Government administered and funded HIV-AIDS program for beneficiaries meeting financial and health status criteria. Administration, funding levels, eligibility criteria, covered benefits, provider types, and financial

participation are typically set by a regulatory process. Payer responsibilities for administering the program may be delegated to contractors. Example: In the U.S., the Ryan White program, which is administered by the Health Resources and Services Administration.

4 | MANDPOL | mandatory health program

4 | MENTPRG | mental health program | Definition: Government administered and funded mental health program for beneficiaries meeting financial and mental health status criteria. Administration, funding levels, eligibility criteria, covered benefits, provider types, and financial participation are typically set by a regulatory process. Payer responsibilities for administering the program may be delegated to contractors. Example: In the U.S., states receive funding for substance use programs from the Substance Abuse Mental Health Administration (SAMHSA).

4 | SAFNET | safety net clinic program | Definition: Government administered and funded program to support provision of care to underserved populations through safety net clinics. Example: In the U.S., safety net providers such as federally qualified health centers (FQHC) receive funding under PHSA Section 330 grants administered by the Health Resources and Services Administration.

4 | SUBPRG | substance use program | Definition: Government administered and funded substance use program for beneficiaries meeting financial, substance use behavior, and health status criteria. Beneficiaries may be required to enroll as a result of legal proceedings. Administration, funding levels, eligibility criteria, covered benefits, provider types, and financial participation are typically set by a regulatory process. Payer responsibilities for administering the program may be delegated to contractors. Example: In the U.S., states receive funding for substance use programs from the Substance Abuse Mental Health Administration (SAMHSA).

4 | SUBSIDIZ | subsidized health program | Definition: A government health program that provides coverage for health services to persons meeting eligibility criteria such as income, location of residence, access to other coverages, health condition, and age, the cost of which is to some extent subsidized by public funds.

5 | SUBSIDMC | subsidized managed care program | Definition: A government health program that provides coverage through managed care contracts for health services to persons meeting eligibility criteria such as income, location of residence, access to other coverages, health condition, and age, the cost of which is to some extent subsidized by public funds. Discussion: The structure and business processes for underwriting and administering a subsidized managed care program is further specified by the Underwriter and Payer Role.class and Role.code.

5 | SUBSUPP | subsidized supplemental health program | Definition: A government health program that provides coverage for health services to persons meeting eligibility criteria for a supplemental health policy or program such as income, location of residence, access to other coverages, health condition, and age, the cost of which is to some extent subsidized by public funds. Example: Supplemental health coverage program may cover the cost of a health program or policy financial participations, such as the copays and the premiums, and may provide coverage for services in addition to those covered under the supplemented health program or policy. In the U.S., Medicaid programs may pay the premium for a covered party who is also covered under the Medicare program or a private health policy. Discussion: The structure and business processes for underwriting and administering a subsidized supplemental retiree health program is further specified by the Underwriter and Payer Role.class and Role.code.

3 | WCBPOL | worker's compensation | Insurance policy for injuries sustained in the work place or in the course of employment.

2 | (_ActInsuranceTypeCode) Abstract | | Definition: Set of codes indicating the type of insurance policy. Insurance, in law and economics, is a form of risk management primarily used to hedge against the risk of potential financial loss. Insurance is defined as the equitable transfer of the risk of a potential loss, from one entity to another, in exchange for a premium and duty of care. A policy holder is an individual or an organization enters into a contract with an underwriter which stipulates that, in exchange for payment of a sum of money (a premium), one or more covered parties (insureds) is guaranteed compensation for losses resulting from certain perils under specified conditions. The underwriter analyzes the risk of loss, makes a decision as to whether the risk is insurable, and prices the premium accordingly. A policy provides benefits that indemnify or cover the cost of a loss incurred by a covered party, and may include coverage for services required to remediate a loss. An insurance policy contains pertinent facts about the policy holder, the insurance coverage, the covered parties, and the insurer. A policy may include exemptions and provisions specifying the extent to which the indemnification clause cannot be enforced for intentional tortious conduct of a covered party, e.g., whether the covered parties are jointly or severably insured. Discussion: In contrast to programs, an insurance policy has one or more policy holders, who own the policy. The policy holder may be the covered party, a relative of the covered party, a partnership, or a corporation, e.g., an employer. A subscriber of a self-insured health insurance policy is a policy holder. A subscriber of an employer sponsored health insurance policy is holds a certificate of coverage, but is not a policy holder; the policy holder is the employer. See CoveredRoleType.

3 | AUTOPOL

3 | (_ActHealthInsuranceTypeCode) Abstract | | Definition: Set of codes indicating the type of health insurance policy that covers health services provided to covered parties. A health insurance policy is a written contract for insurance between the insurance company and the policyholder, and contains pertinent facts about the policy owner (the policy holder), the health insurance coverage, the insured subscribers and dependents, and the insurer. Health insurance is typically administered in accordance with a plan, which specifies (1) the type of health services and health conditions that will be covered under what circumstances (e.g., exclusion of a pre-existing condition, service must be deemed medically necessary; service must not be experimental; service must provided in accordance with a protocol; drug must be on a formulary; service must be prior authorized; or be a referral from a primary care provider); (2) the type and affiliation of providers (e.g., only allopathic physicians, only in network, only providers employed by an HMO); (3) financial participations required of covered parties (e.g., co-pays, coinsurance, deductibles, out-of-pocket); and (4) the manner in which services will be paid (e.g., under indemnity or fee-for-service health plans, the covered party typically pays out-of-pocket and then file a claim for reimbursement, while health plans that have contractual relationships with providers, i.e., network providers, typically do not allow the providers to bill the covered party for the cost of the service until after filing a claim with the payer and receiving reimbursement).

4 | DENTAL | dental care policy | Definition: A health insurance policy that that covers benefits for dental services.

4 | DISEASE | disease specific policy | Definition: A health insurance policy that covers benefits for healthcare services provided for named conditions under the policy, e.g., cancer, diabetes, or HIV-AIDS.

4 | DRUGPOL | drug policy | Definition: A health insurance policy that covers benefits for prescription drugs, pharmaceuticals, and supplies.

4 | EHCPOL

4 | HIP | health insurance plan policy | Definition: A health insurance policy that covers healthcare benefits by protecting covered parties from medical expenses arising from health conditions, sickness, or accidental injury as well as preventive care. Health insurance policies explicitly exclude coverage for losses insured under a disability policy, workers' compensation program, liability insurance (including automobile insurance); or for medical expenses, coverage for on-site medical clinics or for limited dental or vision benefits when these are provided under a separate policy. Discussion: Health insurance policies are offered by health insurance plans that typically reimburse providers for covered services on a fee-for-service basis, that is, a fee that is the allowable amount that a provider may charge. This is in contrast to managed care plans, which typically prepay providers a

per-member/per-month amount or capitation as reimbursement for all covered services rendered. Health insurance plans include indemnity and healthcare services plans.

4 | HSAPOL

4 | LTC | long term care policy | Definition: An insurance policy that covers benefits for long-term care services people need when they no longer can care for themselves. This may be due to an accident, disability, prolonged illness or the simple process of aging. Long-term care services assist with activities of daily living including: Help at home with day-to-day activities, such as cooking, cleaning, bathing and dressing Care in the community, such as in an adult day care facility Supervised care provided in an assisted living facility Skilled care provided in a nursing home

4 | MCPOL | managed care policy | Definition: Government mandated program providing coverage, disability income, and vocational rehabilitation for injuries sustained in the work place or in the course of employment. Employers may either self-fund the program, purchase commercial coverage, or pay a premium to a government entity that administers the program. Employees may be required to pay premiums toward the cost of coverage as well. Managed care policies specifically exclude coverage for losses insured under a disability policy, workers' compensation program, liability insurance (including automobile insurance); or for medical expenses, coverage for on-site medical clinics or for limited dental or vision benefits when these are provided under a separate policy. Discussion: Managed care policies are offered by managed care plans that contract with selected providers or health care organizations to provide comprehensive health care at a discount to covered parties and coordinate the financing and delivery of health care. Managed care uses medical protocols and procedures agreed on by the medical profession to be cost effective, also known as medical practice guidelines. Providers are typically reimbursed for covered services by a capitated amount on a per member per month basis that may reflect difference in the health status and level of services anticipated to be needed by the member.

5 | POS | point of service policy | Definition: A policy for a health plan that has features of both an HMO and a FFS plan. Like an HMO, a POS plan encourages the use its HMO network to maintain discounted fees with participating providers, but recognizes that sometimes covered parties want to choose their own provider. The POS plan allows a covered party to use providers who are not part of the HMO network (non-participating providers). However, there is a greater cost associated with choosing these non-network providers. A covered party will usually pay deductibles and coinsurances that are substantially higher than the payments when he or she uses a plan provider. Use of non-participating providers often requires the covered party to pay the provider directly and then to file a claim for reimbursement, like in an FFS plan.

5 | HMO | health maintenance organization policy | Definition: A policy for a health plan that provides coverage for health care only through contracted or employed physicians and hospitals located in particular geographic or service areas. HMOs emphasize prevention and early detection of illness. Eligibility to enroll in an HMO is determined by where a covered party lives or works.

5 | PPO | preferred provider organization policy | Definition: A network-based, managed care plan that allows a covered party to choose any health care provider. However, if care is received from a "preferred" (participating in-network) provider, there are generally higher benefit coverage and lower deductibles.

4 | MENTPOL | mental health policy | Definition: A health insurance policy that covers benefits for mental health services and prescriptions.

4 | POS

4 | SUBPOL | substance use policy | Definition: A health insurance policy that covers benefits for substance use services.

4 | VISPOL | vision care policy | Definition: Set of codes for a policy that provides coverage for health care expenses arising from vision services. A health insurance policy that covers benefits for vision care services, prescriptions, and products.

3 | DIS | disability insurance policy | Definition: An insurance policy that provides a regular payment to compensate for income lost due to the covered party's inability to work because of illness or injury.

3 | EWB | employee welfare benefit plan policy | Definition: An insurance policy under a benefit plan run by an employer or employee organization for the purpose of providing benefits other than pension-related to employees and their families. Typically provides health-related benefits, benefits for disability, disease or unemployment, or day care and scholarship benefits, among others. An employer sponsored health policy includes coverage of health care expenses arising from sickness or accidental injury, coverage for on-site medical clinics or for dental or vision benefits, which are typically provided under a separate policy. Coverage excludes health care expenses covered by accident or disability, workers' compensation, liability or automobile insurance.

3 | FLEXP | flexible benefit plan policy | Definition: An insurance policy that covers qualified benefits under a Flexible Benefit plan such as group medical insurance, long and short term disability income insurance, group term life insurance for employees only up to $50,000 face amount, specified disease coverage such as a cancer policy, dental and/or vision insurance, hospital indemnity insurance, accidental death and dismemberment insurance, a medical expense reimbursement plan and a dependent care reimbursement plan. Discussion: See UnderwriterRoleTypeCode flexible benefit plan which is defined as a benefit plan that allows employees to choose from several life, health, disability, dental, and other insurance plans according to their individual needs. Also known as cafeteria plans. Authorized under Section 125 of the Revenue Act of 1978.

3 | LIFE | life insurance policy | Definition: A policy under which the insurer agrees to pay a sum of money upon the occurrence of the covered partys death. In return, the policyholder agrees to pay a stipulated amount called a premium at regular intervals. Life insurance indemnifies the beneficiary for the loss of the insurable interest that a beneficiary has in the life of a covered party. For persons related by blood, a substantial interest established through love and affection, and for all other persons, a lawful and substantial economic interest in having the life of the insured continue. An insurable interest is required when purchasing life insurance on another person. Specific exclusions are often written into the contract to limit the liability of the insurer; for example claims resulting from suicide or relating to war, riot and civil commotion. Discussion:A life insurance policy may be used by the covered party as a source of health care coverage in the case of a viatical settlement, which is the sale of a life insurance policy by the policy owner, before the policy matures. Such a sale, at a price discounted from the face amount of the policy but usually in excess of the premiums paid or current cash surrender value, provides the seller an immediate cash settlement. Generally, viatical settlements involve insured individuals with a life expectancy of less than two years. In countries without state-subsidized healthcare and high healthcare costs (e.g. United States), this is a practical way to pay extremely high health insurance premiums that severely ill people face. Some people are also familiar with life settlements, which are similar transactions but involve insureds with longer life expectancies (two to fifteen years).

4 | ANNU | annuity policy | Definition: A policy that, after an initial premium or premiums, pays out a sum at pre-determined intervals. For example, a policy holder may pay $10,000, and in return receive $150 each month until he dies; or $1,000 for each of 14 years or death benefits if he dies before the full term of the annuity has elapsed.

4 | TLIFE | term life insurance policy | Definition: Life insurance under which the benefit is payable only if the insured dies during a specified period. If an insured dies during that period, the beneficiary receives the death payments. If the insured survives, the policy ends and the beneficiary receives nothing.

4 | ULIFE | universal life insurance policy | Definition: Life insurance under which the benefit is payable upon the insuredaTMs death or diagnosis of a terminal illness. If an insured dies during that period, the beneficiary receives the death payments. If the insured survives, the policy ends and the beneficiary receives nothing

3 | PNC | property and casualty insurance policy | Definition: A type of insurance that covers damage to or loss of the policyholderaTMs property by providing payments for damages to property damage or the injury or death of living subjects. The terms "casualty" and "liability" insurance are often used interchangeably. Both cover the policyholder's legal liability for damages caused to other persons and/or their property.

3 | REI | reinsurance policy | Definition: An agreement between two or more insurance companies by which the risk of loss is proportioned. Thus the risk of loss is spread and a disproportionately large loss under a single policy does not fall on one insurance company. Acceptance by an insurer, called a reinsurer, of all or part of the risk of loss of another insurance company. Discussion: Reinsurance is a means by which an insurance company can protect itself against the risk of losses with other insurance companies. Individuals and corporations obtain insurance policies to provide protection for various risks (hurricanes, earthquakes, lawsuits, collisions, sickness and death, etc.). Reinsurers, in turn, provide insurance to insurance companies. For example, an HMO may purchase a reinsurance policy to protect itself from losing too much money from one insured's particularly expensive health care costs. An insurance company issuing an automobile liability policy, with a limit of $100,000 per accident may reinsure its liability in excess of $10,000. A fire insurance company which issues a large policy generally reinsures a portion of the risk with one or several other companies. Also called risk control insurance or stop-loss insurance.

3 | SURPL | surplus line insurance policy | Definition: A risk or part of a risk for which there is no normal insurance market available. Insurance written by unauthorized insurance companies. Surplus lines insurance is insurance placed with unauthorized insurance companies through licensed surplus lines agents or brokers.

3 | UMBRL | umbrella liability insurance policy | Definition: A form of insurance protection that provides additional liability coverage after the limits of your underlying policy are reached. An umbrella liability policy also protects you (the insured) in many situations not covered by the usual liability policies.

2 | (_ActProgramTypeCode) Abstract | | Definition: A set of codes used to indicate coverage under a program. A program is an organized structure for administering and funding coverage of a benefit package for covered parties meeting eligibility criteria, typically related to employment, health, financial, and demographic status. Programs are typically established or permitted by legislation with provisions for ongoing government oversight. Regulations may mandate the structure of the program, the manner in which it is funded and administered, covered benefits, provider types, eligibility criteria and financial participation. A government agency may be charged with implementing the program in accordance to the regulation. Risk of loss under a program in most cases would not meet what an underwriter would consider an insurable risk, i.e., the risk is not random in nature, not financially measurable, and likely requires subsidization with government funds. Discussion: Programs do not have policy holders or subscribers. Program eligibles are enrolled based on health status, statutory eligibility, financial status, or age. Program eligibles who are covered parties under the program may be referred to as members, beneficiaries, eligibles, or recipients. Programs risk are underwritten by not for profit organizations such as governmental entities, and the beneficiaries typically do not pay for any or some portion of the cost of coverage. See CoveredPartyRoleType.

3 | PUBLICPOL

3 | WCBPOL

3 | CHAR | charity program | Definition: A program that covers the cost of services provided directly to a beneficiary who typically has no other source of coverage without charge.

3 | CRIME | crime victim program | Definition: A program that covers the cost of services provided to crime victims for injuries or losses related to the occurrence of a crime.

3 | EAP | employee assistance program | Definition: An employee assistance program is run by an employer or employee organization for the purpose of providing benefits and covering all or part of the cost for employees to receive counseling, referrals, and advice in dealing with stressful issues in their lives. These may include substance abuse, bereavement, marital problems, weight issues, or general wellness issues. The services are usually provided by a third-party, rather than the company itself, and the company receives only summary statistical data from the service provider. Employee's names and services received are kept confidential.

3 | GOVEMP | government employee health program | Definition: A set of codes used to indicate a government program that is an organized structure for administering and funding coverage of a benefit package for covered parties meeting eligibility criteria, typically related to employment, health and financial status. Government programs are established or permitted by legislation with provisions for ongoing government oversight. Regulation mandates the structure of the program, the manner in which it is funded and administered, covered benefits, provider types, eligibility criteria and financial participation. A government agency is charged with implementing the program in accordance to the regulation Example: Federal employee health benefit program in the U.S.

3 | HIRISK | high risk pool program | Definition: A government program that provides health coverage to individuals who are considered medically uninsurable or high risk, and who have been denied health insurance due to a serious health condition. In certain cases, it also applies to those who have been quoted very high premiums a" again, due to a serious health condition. The pool charges premiums for coverage. Because the pool covers high-risk people, it incurs a higher level of claims than premiums can cover. The insurance industry pays into the pool to make up the difference and help it remain viable.

3 | IND | indigenous peoples health program | Definition: Services provided directly and through contracted and operated indigenous peoples health programs. Example: Indian Health Service in the U.S.

3 | MILITARY | military health program | Definition: A government program that provides coverage for health services to military personnel, retirees, and dependents. A covered party who is a subscriber can choose from among Fee-for-Service (FFS) plans, and their Preferred Provider Organizations (PPO), or Plans offering a Point of Service (POS) Product, or Health Maintenance Organizations. Example: In the U.S., TRICARE, CHAMPUS.

3 | RETIRE | retiree health program | Definition: A government mandated program with specific eligibility requirements based on premium contributions made during employment, length of employment, age, and employment status, e.g., being retired, disabled, or a dependent of a covered party under this program. Benefits typically include ambulatory, inpatient, and long-term care, such as hospice care, home health care and respite care.

3 | SOCIAL | social service program | Definition: A social service program funded by a public or governmental entity. Example: Programs providing habilitation, food, lodging, medicine, transportation, equipment, devices, products, education, training, counseling, alteration of living or work space, and other resources to persons meeting eligibility criteria.

3 | VET | veteran health program | Definition: Services provided directly and through contracted and operated veteran health programs.

1 | (_ActDetectedIssueManagementCode) Abstract | | Codes dealing with the management of Detected Issue observations

2 | (_ActAdministrativeDetectedIssueManagementCode) Abstract | | Codes dealing with the management of Detected Issue observations for the administrative and patient administrative acts domains.

3 | (_AuthorizationIssueManagementCode) Abstract

4 | EMAUTH | emergency authorization override | Used to temporarily override normal authorization rules to gain access to data in a case of emergency. Use of this override code will typically be monitored, and a procedure to verify its proper use may be triggered when used.

5 | 21 | authorization confirmed | Description: Indicates that the permissions have been externally verified and the request should be processed.
2 | 1 | Therapy Appropriate | Confirmed drug therapy appropriate
3 | 19 | Consulted Supplier | Consulted other supplier/pharmacy, therapy confirmed
3 | 2 | Assessed Patient | Assessed patient, therapy is appropriate
3 | 22 | appropriate indication or diagnosis | Description: The patient has the appropriate indication or diagnosis for the action to be taken.
3 | 23 | prior therapy documented | Description: It has been confirmed that the appropriate pre-requisite therapy has been tried.
3 | 3 | Patient Explanation | Patient gave adequate explanation
3 | 4 | Consulted Other Source | Consulted other supply source, therapy still appropriate
3 | 5 | Consulted Prescriber | Consulted prescriber, therapy confirmed
4 | 6 | Prescriber Declined Change | Consulted prescriber and recommended change, prescriber declined
3 | 7 | Interacting Therapy No Longer Active/Planned | Concurrent therapy triggering alert is no longer on-going or planned
2 | 14 | Supply Appropriate | Confirmed supply action appropriate
3 | 15 | Replacement | Patient's existing supply was lost/wasted
3 | 16 | Vacation Supply | Supply date is due to patient vacation
3 | 17 | Weekend Supply | Supply date is intended to carry patient over weekend
3 | 18 | Leave of Absence | Supply is intended for use during a leave of absence from an institution.
3 | 20 | additional quantity on separate dispense | Description: Supply is different than expected as an additional quantity has been supplied in a separate dispense.
2 | 8 | Other Action Taken | Order is performed as issued, but other action taken to mitigate potential adverse effects
3 | 10 | Provided Patient Education | Provided education or training to the patient on appropriate therapy use
3 | 11 | Added Concurrent Therapy | Instituted an additional therapy to mitigate potential negative effects
3 | 12 | Temporarily Suspended Concurrent Therapy | Suspended existing therapy that triggered interaction for the duration of this therapy
3 | 13 | Stopped Concurrent Therapy | Aborted existing therapy that triggered interaction.
3 | 9 | Instituted Ongoing Monitoring Program | Arranged to monitor patient for adverse effects
1 | (_ActExposureCode) Abstract | | Concepts that identify the type or nature of exposure interaction. Examples include "household", "care giver", "intimate partner", "common space", "common substance", etc. to further describe the nature of interaction.
2 | CHLDCARE | Day care - Child care Interaction | Description: Exposure participants' interaction occurred in a child care setting
2 | CONVEYNC | Common Conveyance Interaction | Description: An interaction where the exposure participants traveled in/on the same vehicle (not necessarily concurrently, e.g. both are passengers of the same plane, but on different flights of that plane).
2 | HLTHCARE | Health Care Interaction - Not Patient Care | Description: Exposure participants' interaction occurred during the course of health care delivery or in a health care delivery setting, but did not involve the direct provision of care (e.g. a janitor cleaning a patient's hospital room).
2 | HOMECARE | Care Giver Interaction | Description: Exposure interaction occurred in context of one providing care for the other, i.e. a babysitter providing care for a child, a home-care aide providing assistance to a paraplegic.
2 | HOSPPTNT | Hospital Patient Interaction | Description: Exposure participants' interaction occurred when both were patients being treated in the same (acute) health care delivery facility.
2 | HOSPVSTR | Hospital Visitor Interaction | Description: Exposure participants' interaction occurred when one visited the other who was a patient being treated in a health care delivery facility.
2 | HOUSEHLD | Household Interaction | Description: Exposure interaction occurred in context of domestic interaction, i.e. both participants reside in the same household.
2 | INMATE | Inmate Interaction | Description: Exposure participants' interaction occurred in the course of one or both participants being incarcerated at a correctional facility
2 | INTIMATE | Intimate Interaction | Description: Exposure interaction was intimate, i.e. participants are intimate companions (e.g. spouses, domestic partners).
2 | LTRMCARE | Long Term Care Facility Interaction | Description: Exposure participants' interaction occurred in the course of one or both participants being resident at a long term care facility (second participant may be a visitor, worker, resident or a physical place or object within the facility).
2 | PLACE | Common Space Interaction | Description: An interaction where the exposure participants were both present in the same location/place/space.
2 | PTNTCARE | Health Care Interaction - Patient Care | Description: Exposure participants' interaction occurred during the course of health care delivery by a provider (e.g. a physician treating a patient in her office).
2 | SCHOOL2 | School Interaction | Description: Exposure participants' interaction occurred in an academic setting (e.g., participants are fellow students, or student and teacher).
2 | SOCIAL2 | Social/Extended Family Interaction | Description: An interaction where the exposure participants are social associates or members of the same extended family
2 | SUBSTNCE | Common Substance Interaction | Description: An interaction where the exposure participants shared or co-used a common substance (e.g. drugs, needles, or common food item).
2 | TRAVINT | Common Travel Interaction | Description: An interaction where the exposure participants traveled together in/on the same vehicle/trip (e.g. concurrent co-passengers).
2 | WORK2 | Work Interaction | Description: Exposure interaction occurred in a work setting, i.e. participants are co-workers.
1 | (_ActFinancialTransactionCode) Abstract
2 | CHRG | Standard Charge | A type of transaction that represents a charge for a service or product. Expressed in monetary terms.
2 | REV | Standard Charge Reversal | A type of transaction that represents a reversal of a previous charge for a service or product. Expressed in monetary terms. It has the opposite effect of a standard charge.
1 | (_ActIncidentCode) Abstract | | Set of codes indicating the type of incident or accident.
2 | MVA | Motor vehicle accident | Incident or accident as the result of a motor vehicle accident
2 | SCHOOL | School Accident | Incident or accident is the result of a school place accident.
2 | SPT | Sporting Accident | Incident or accident is the result of a sporting accident.
2 | WPA | Workplace accident | Incident or accident is the result of a work place accident
1 | (_ActInformationAccessCode) Abstract | | Description: The type of health information to which the subject of the information or the subject's delegate consents or dissents.

2 | ACADR | adverse drug reaction access | Description: Provide consent to collect, use, disclose, or access adverse drug reaction information for a patient.
2 | ACALL | all access | Description: Provide consent to collect, use, disclose, or access all information for a patient.
2 | ACALLG | allergy access | Description: Provide consent to collect, use, disclose, or access allergy information for a patient.
2 | ACCONS | informational consent access | Description: Provide consent to collect, use, disclose, or access informational consent information for a patient.
2 | ACDEMO | demographics access | Description: Provide consent to collect, use, disclose, or access demographics information for a patient.
2 | ACDI | diagnostic imaging access | Description: Provide consent to collect, use, disclose, or access diagnostic imaging information for a patient.
2 | ACIMMUN | immunization access | Description: Provide consent to collect, use, disclose, or access immunization information for a patient.
2 | ACLAB | lab test result access | Description: Provide consent to collect, use, disclose, or access lab test result information for a patient.
2 | ACMED | medication access | Description: Provide consent to collect, use, disclose, or access medical condition information for a patient.
2 | ACMEDC | medical condition access | Definition: Provide consent to view or access medical condition information for a patient.
2 | ACMEN | mental health access | Description:Provide consent to collect, use, disclose, or access mental health information for a patient.
2 | ACOBS | common observations access | Description: Provide consent to collect, use, disclose, or access common observation information for a patient.
2 | ACPOLPRG | policy or program information access | Description: Provide consent to collect, use, disclose, or access coverage policy or program for a patient.
2 | ACPROV | provider information access | Description: Provide consent to collect, use, disclose, or access provider information for a patient.
2 | ACPSERV | professional service access | Description: Provide consent to collect, use, disclose, or access professional service information for a patient.
2 | ACSUBSTAB | substance abuse access | Description:Provide consent to collect, use, disclose, or access substance abuse information for a patient.
1 | (_ActInformationAccessContextCode) Abstract | | Concepts conveying the context in which authorization given under jurisdictional law, by organizational policy, or by a patient consent directive permits the collection, access, use or disclosure of specified patient health information.
2 | INFAUT | authorized information transfer | Authorization to collect, access, use, or disclose specified patient health information in accordance with jurisdictional law, organizational policy, or a patient's consent directive, which may be implied, deemed, opt-in, opt-out, or explicit.
3 | INFCON | after explicit consent | Authorization to collect, access, use, or disclose specified patient health information as explicitly consented to by the subject of the information or the subject's representative.
2 | INFCRT | only on court order | Authorization to collect, access, use, or disclose specified patient health information in accordance with judicial system protocol, such as in the case of a subpoena or court order.
2 | INFDNG | only if danger to others | Authorization to collect, access, use, or disclose specified patient health information where deemed necessary to avert potential danger to other persons in accordance with jurisdictional law, organizational policy, or standards of practice. For example, disclosure about a person threatening violence.
2 | INFEMER | only in an emergency | Authorization to collect, access, use, or disclose specified patient health information in accordance with emergency information transfer protocol dictated by jurisdictional law, organization policy, or standards of practice. For example, sharing of health information during disaster response.
2 | INFPWR | only if public welfare risk | Authorization to collect, access, use, or disclose specified patient health information necessary to avert potential public welfare risk in accordance with jurisdictional law, organizational policy, or standards of practice. For example, reporting that a person is a victim of abuse or demonstrating suicidal tendencies.
2 | INFREG | regulatory information transfer | Authorization to collect, access, use, or disclose specified patient health information for public health, welfare, and safety purposes in accordance with jurisdictional law, organizational policy, or standards of practice. For example, public health reporting of notifiable conditions.
1 | (_ActInformationCategoryCode) Abstract | | Definition:Indicates the set of information types which may be manipulated or referenced, such as for recommending access restrictions.
2 | ALLCAT | all categories | Description: All patient information.
2 | ALLGCAT | allergy category | Definition:All information pertaining to a patient's allergy and intolerance records.
2 | ARCAT | adverse drug reaction category | Description: All information pertaining to a patient's adverse drug reactions.
2 | COBSCAT | common observation category | Definition:All information pertaining to a patient's common observation records (height, weight, blood pressure, temperature, etc.).
2 | DEMOCAT | demographics category | Definition:All information pertaining to a patient's demographics (such as name, date of birth, gender, address, etc).
2 | DICAT | diagnostic image category | Definition:All information pertaining to a patient's diagnostic image records (orders & results).
2 | IMMUCAT | immunization category | Definition:All information pertaining to a patient's vaccination records.
2 | LABCAT | lab test category | Description: All information pertaining to a patient's lab test records (orders & results)
2 | MEDCCAT | medical condition category | Definition:All information pertaining to a patient's medical condition records.
2 | MENCAT | mental health category | Description: All information pertaining to a patient's mental health records.
2 | PSVCCAT | professional service category | Definition:All information pertaining to a patient's professional service records (such as smoking cessation, counseling, medication review, mental health).
2 | RXCAT | medication category | Definition:All information pertaining to a patient's medication records (orders, dispenses and other active medications).
1 | (_ActInvoiceElementCode) Abstract | | Type of invoice element that is used to assist in describing an Invoice that is either submitted for adjudication or for which is returned on adjudication results.
2 | (_ActInvoiceAdjudicationPaymentCode) Abstract | | Codes representing a grouping of invoice elements (totals, sub-totals), reported through a Payment Advice or a Statement of Financial Activity (SOFA). The code can represent summaries by day, location, payee and other cost elements such as bonus, retroactive adjustment and transaction fees.

3 | (_ActInvoiceAdjudicationPaymentGroupCode) Abstract | | Codes representing adjustments to a Payment Advice such as retroactive, clawback, garnishee, etc.
4 | ALEC | alternate electronic | Payment initiated by the payor as the result of adjudicating a submitted invoice that arrived to the payor from an electronic source that did not provide a conformant set of HL7 messages (e.g. web claim submission).
4 | BONUS | bonus | Bonus payments based on performance, volume, etc. as agreed to by the payor.
4 | CFWD | carry forward adjusment | An amount still owing to the payor but the payment is 0$ and this cannot be settled until a future payment is made.
4 | EDU | education fees | Fees deducted on behalf of a payee for tuition and continuing education.
4 | EPYMT | early payment fee | Fees deducted on behalf of a payee for charges based on a shorter payment frequency (i.e. next day versus biweekly payments.
4 | GARN | garnishee | Fees deducted on behalf of a payee for charges based on a per-transaction or time-period (e.g. monthly) fee.
4 | INVOICE | submitted invoice | Payment is based on a payment intent for a previously submitted Invoice, based on formal adjudication results..
4 | PINV | paper invoice | Payment initiated by the payor as the result of adjudicating a paper (original, may have been faxed) invoice.
4 | PPRD | prior period adjustment | An amount that was owed to the payor as indicated, by a carry forward adjusment, in a previous payment advice
4 | PROA | professional association deduction | Professional association fee that is collected by the payor from the practitioner/provider on behalf of the association
4 | RECOV | recovery | Retroactive adjustment such as fee rate adjustment due to contract negotiations.
4 | RETRO | retro adjustment | Bonus payments based on performance, volume, etc. as agreed to by the payor.
4 | TRAN | transaction fee | Fees deducted on behalf of a payee for charges based on a per-transaction or time-period (e.g. monthly) fee.
3 | (_ActInvoiceAdjudicationPaymentSummaryCode) Abstract | | Codes representing a grouping of invoice elements (totals, sub-totals), reported through a Payment Advice or a Statement of Financial Activity (SOFA). The code can represent summaries by day, location, payee, etc.
4 | CONT
4 | DAY
4 | LOC
4 | MONTH
4 | PERIOD
4 | PROV
4 | WEEK
4 | YEAR
4 | INVTYPE | invoice type | Transaction counts and value totals by invoice type (e.g. RXDINV - Pharmacy Dispense)
4 | PAYEE | payee | Transaction counts and value totals by each instance of an invoice payee.
4 | PAYOR | payor | Transaction counts and value totals by each instance of an invoice payor.
4 | SENDAPP | sending application | Transaction counts and value totals by each instance of a messaging application on a single processor. It is a registered identifier known to the receivers.
2 | (_ActInvoiceDetailCode) Abstract | | Codes representing a service or product that is being invoiced (billed). The code can represent such concepts as "office visit", "drug X", "wheelchair" and other billable items such as taxes, service charges and discounts.
3 | (_ActInvoiceDetailClinicalProductCode) Abstract | | An identifying data string for healthcare products.
4 | UNSPSC | United Nations Standard Products and Services Classification | Description:United Nations Standard Products and Services Classification, managed by Uniform Code Council (UCC): www.unspsc.org
3 | (_ActInvoiceDetailDrugProductCode) Abstract | | An identifying data string for A substance used as a medication or in the preparation of medication.
4 | GTIN | Global Trade Item Number | Description:Global Trade Item Number is an identifier for trade items developed by GS1 (comprising the former EAN International and Uniform Code Council).
4 | UPC | Universal Product Code | Description:Universal Product Code is one of a wide variety of bar code languages widely used in the United States and Canada for items in stores.
3 | (_ActInvoiceDetailGenericCode) Abstract | | The detail item codes to identify charges or changes to the total billing of a claim due to insurance rules and payments.
4 | (_ActInvoiceDetailGenericAdjudicatorCode) Abstract | | The billable item codes to identify adjudicator specified components to the total billing of a claim.
5 | COIN | coinsurance | That portion of the eligible charges which a covered party must pay for each service and/or product. It is a percentage of the eligible amount for the service/product that is typically charged after the covered party has met the policy deductible. This amount represents the covered party's coinsurance that is applied to a particular adjudication result. It is expressed as a negative dollar amount in adjudication results.
5 | COPAYMENT | patient co-pay | That portion of the eligible charges which a covered party must pay for each service and/or product. It is a defined amount per service/product of the eligible amount for the service/product. This amount represents the covered party's copayment that is applied to a particular adjudication result. It is expressed as a negative dollar amount in adjudication results.
5 | DEDUCTIBLE | deductible | That portion of the eligible charges which a covered party must pay in a particular period (e.g. annual) before the benefits are payable by the adjudicator. This amount represents the covered party's deductible that is applied to a particular adjudication result. It is expressed as a negative dollar amount in adjudication results.
5 | PAY | payment | The guarantor, who may be the patient, pays the entire charge for a service. Reasons for such action may include: there is no insurance coverage for the service (e.g. cosmetic surgery); the patient wishes to self-pay for the service; or the insurer denies payment for the service due to contractual provisions such as the need for prior authorization.
5 | SPEND | spend down | That total amount of the eligible charges which a covered party must periodically pay for services and/or products prior to the Medicaid program providing any coverage. This amount represents the covered party's spend down that is applied to a particular adjudication result. It is expressed as a negative dollar amount in adjudication results
5 | COINS | co-insurance | The covered party pays a percentage of the cost of covered services.

4 | (_ActInvoiceDetailGenericModifierCode) Abstract | | The billable item codes to identify modifications to a billable item charge. As for example after hours increase in the office visit fee.
5 | AFTHRS | non-normal hours | Premium paid on service fees in compensation for practicing outside of normal working hours.
5 | ISOL | isolation allowance | Premium paid on service fees in compensation for practicing in a remote location.
5 | OOO | out of office | Premium paid on service fees in compensation for practicing at a location other than normal working location.
4 | (_ActInvoiceDetailGenericProviderCode) Abstract | | The billable item codes to identify provider supplied charges or changes to the total billing of a claim.
5 | CANCAPT | cancelled appointment | A charge to compensate the provider when a patient cancels an appointment with insufficient time for the provider to make another appointment with another patient.
5 | DSC | discount | A reduction in the amount charged as a percentage of the amount. For example a 5% discount for volume purchase.
5 | ESA | extraordinary service assessment | A premium on a service fee is requested because, due to extenuating circumstances, the service took an extraordinary amount of time or supplies.
5 | FFSTOP | fee for service top off | Under agreement between the parties (payor and provider), a guaranteed level of income is established for the provider over a specific, pre-determined period of time. The normal course of business for the provider is submission of fee-for-service claims. Should the fee-for-service income during the specified period of time be less than the agreed to amount, a top-up amount is paid to the provider equal to the difference between the fee-for-service total and the guaranteed income amount for that period of time. The details of the agreement may specify (or not) a requirement for repayment to the payor in the event that the fee-for-service income exceeds the guaranteed amount.
5 | FNLFEE | final fee | Anticipated or actual final fee associated with treating a patient.
5 | FRSTFEE | first fee | Anticipated or actual initial fee associated with treating a patient.
5 | MARKUP | markup or up-charge | An increase in the amount charged as a percentage of the amount. For example, 12% markup on product cost.
5 | MISSAPT | missed appointment | A charge to compensate the provider when a patient does not show for an appointment.
5 | PERFEE | periodic fee | Anticipated or actual periodic fee associated with treating a patient. For example, expected billing cycle such as monthly, quarterly. The actual period (e.g. monthly, quarterly) is specified in the unit quantity of the Invoice Element.
5 | PERMBNS | performance bonus | The amount for a performance bonus that is being requested from a payor for the performance of certain services (childhood immunizations, influenza immunizations, mammograms, pap smears) on a sliding scale. That is, for 90% of childhood immunizations to a maximum of $2200/yr. An invoice is created at the end of the service period (one year) and a code is submitted indicating the percentage achieved and the dollar amount claimed.
5 | RESTOCK | restocking fee | A charge is requested because the patient failed to pick up the item and it took an amount of time to return it to stock for future use.
5 | TRAVEL | travel | A charge to cover the cost of travel time and/or cost in conjuction with providing a service or product. It may be charged per kilometer or per hour based on the effective agreement.
5 | URGENT | urgent | Premium paid on service fees in compensation for providing an expedited response to an urgent situation.
4 | (_ActInvoiceDetailTaxCode) Abstract | | The billable item codes to identify modifications to a billable item charge by a tax factor applied to the amount. As for example 7% provincial sales tax.
5 | FST | federal sales tax | Federal tax on transactions such as the Goods and Services Tax (GST)
5 | HST | harmonized sales Tax | Joint Federal/Provincial Sales Tax
5 | PST | provincial/state sales tax | Tax levied by the provincial or state jurisdiction such as Provincial Sales Tax
3 | (_ActInvoiceDetailPreferredAccommodationCode) Abstract | | An identifying data string for medical facility accommodations.
4 | (_ActEncounterAccommodationCode) Abstract | | Accommodation type. In Intent mood, represents the accommodation type requested. In Event mood, represents accommodation assigned/used. In Definition mood, represents the available accommodation type.
5 | (_HL7AccommodationCode) Abstract | | Description:Accommodation type. In Intent mood, represents the accommodation type requested. In Event mood, represents accommodation assigned/used. In Definition mood, represents the available accommodation type.
6 | I | Isolation | Accommodations used in the care of diseases that are transmitted through casual contact or respiratory transmission.
6 | P | Private | Accommodations in which there is only 1 bed.
6 | S | Suite | Uniquely designed and elegantly decorated accommodations with many amenities available for an additional charge.
6 | SP | Semi-private | Accommodations in which there are 2 beds.
6 | W | Ward | Accommodations in which there are 3 or more beds.
3 | (_ActInvoiceDetailClinicalServiceCode) Abstract | | An identifying data string for healthcare procedures.
2 | (_ActInvoiceGroupCode) Abstract | | Type of invoice element that is used to assist in describing an Invoice that is either submitted for adjudication or for which is returned on adjudication results. Invoice elements of this type signify a grouping of one or more children (detail) invoice elements. They do not have intrinsic costing associated with them, but merely reflect the sum of all costing for it's immediate children invoice elements.
3 | (_ActInvoiceInterGroupCode) Abstract | | Type of invoice element that is used to assist in describing an Invoice that is either submitted for adjudication or for which is returned on adjudication results. Invoice elements of this type signify a grouping of one or more children (detail) invoice elements. They do not have intrinsic costing associated with them, but merely reflect the sum of all costing for it's immediate children invoice elements. The domain is only specified for an intermediate invoice element group (non-root or non-top level) for an Invoice.
4 | CPNDDRGING | compound drug invoice group | A grouping of invoice element groups and details including the ones specifying the compound ingredients being invoiced. It may also contain generic detail items such as markup.
4 | CPNDINDING | compound ingredient invoice group | A grouping of invoice element details including the one specifying an ingredient drug being invoiced. It may also contain generic detail items such as tax or markup.
4 | CPNDSUPING | compound supply invoice group | A grouping of invoice element groups and details including the ones specifying the compound supplies being invoiced. It may also contain generic detail items such as markup.
4 | DRUGING | drug invoice group | A grouping of invoice element details including the one specifying the drug being invoiced. It may also contain generic detail items such as markup.
4 | FRAMEING | frame invoice group | A grouping of invoice element details including the ones specifying the frame fee and the frame dispensing cost that are being invoiced.
4 | LENSING | lens invoice group | A grouping of invoice element details including the ones specifying the lens fee and the lens dispensing cost that are being invoiced.

4 | PRDING | product invoice group | A grouping of invoice element details including the one specifying the product (good or supply) being invoiced. It may also contain generic detail items such as tax or discount.
3 | (_ActInvoiceRootGroupCode) Abstract | | Type of invoice element that is used to assist in describing an Invoice that is either submitted for adjudication or for which is returned on adjudication results. Invoice elements of this type signify a grouping of one or more children (detail) invoice elements. They do not have intrinsic costing associated with them, but merely reflect the sum of all costing for it's immediate children invoice elements. Codes from this domain reflect the type of Invoice such as Pharmacy Dispense, Clinical Service and Clinical Product. The domain is only specified for the root (top level) invoice element group for an Invoice.
4 | CPINV | clinical product invoice | Clinical product invoice where the Invoice Grouping contains one or more billable item and is supported by clinical product(s). For example, a crutch or a wheelchair.
4 | CSINV | clinical service invoice | Clinical Services Invoice which can be used to describe a single service, multiple services or repeated services. [1] Single Clinical services invoice where the Invoice Grouping contains one billable item and is supported by one clinical service. For example, a single service for an office visit or simple clinical procedure (e.g. knee mobilization). [2] Multiple Clinical services invoice where the Invoice Grouping contains more than one billable item, supported by one or more clinical services. The services can be distinct and over multiple dates, but for the same patient. This type of invoice includes a series of treatments which must be adjudicated together. For example, an adjustment and ultrasound for a chiropractic session where fees are associated for each of the services and adjudicated (invoiced) together. [3] Repeated Clinical services invoice where the Invoice Grouping contains one or more billable item, supported by the same clinical service repeated over a period of time. For example, the same Chiropractic adjustment (service or treatment) delivered on 3 separate occasions over a period of time at the discretion of the provider (e.g. month).
4 | CSPINV | clinical service and product | A clinical Invoice Grouping consisting of one or more services and one or more product. Billing for these service(s) and product(s) are supported by multiple clinical billable events (acts). All items in the Invoice Grouping must be adjudicated together to be acceptable to the Adjudicator. For example , a brace (product) invoiced together with the fitting (service).
4 | FININV | financial invoice | Invoice Grouping without clinical justification. These will not require identification of participants and associations from a clinical context such as patient and provider. Examples are interest charges and mileage.
4 | OHSINV | oral health service | A clinical Invoice Grouping consisting of one or more oral health services. Billing for these service(s) are supported by multiple clinical billable events (acts). All items in the Invoice Grouping must be adjudicated together to be acceptable to the Adjudicator.
4 | PAINV | preferred accommodation invoice | HealthCare facility preferred accommodation invoice.
4 | RXCINV | Rx compound invoice | Pharmacy dispense invoice for a compound.
4 | RXDINV | Rx dispense invoice | Pharmacy dispense invoice not involving a compound
4 | SBFINV | sessional or block fee invoice | Clinical services invoice where the Invoice Group contains one billable item for multiple clinical services in one or more sessions.
4 | VRXINV | vision dispense invoice | Vision dispense invoice for up to 2 lens (left and right), frame and optional discount. Eye exams are invoiced as a clinical service invoice.
1 | (_ActInvoiceElementSummaryCode) Abstract | | Identifies the different types of summary information that can be reported by queries dealing with Statement of Financial Activity (SOFA). The summary information is generally used to help resolve balance discrepancies between providers and payors.
2 | (_InvoiceElementAdjudicated) Abstract | | Total counts and total net amounts adjudicated for all Invoice Groupings that were adjudicated within a time period based on the adjudication date of the Invoice Grouping.
3 | ADNFPPELAT | adjud. nullified prior-period electronic amount | Identifies the total net amount of all Invoice Groupings that were adjudicated as payable prior to the specified time period (based on adjudication date), subsequently cancelled in the specified period and submitted electronically.
3 | ADNFPPELCT | adjud. nullified prior-period electronic count | Identifies the total number of all Invoice Groupings that were adjudicated as payable prior to the specified time period (based on adjudication date), subsequently cancelled in the specified period and submitted electronically.
3 | ADNFPPMNAT | adjud. nullified prior-period manual amount | Identifies the total net amount of all Invoice Groupings that were adjudicated as payable prior to the specified time period (based on adjudication date), subsequently cancelled in the specified period and submitted manually.
3 | ADNFPPMNCT | adjud. nullified prior-period manual count | Identifies the total number of all Invoice Groupings that were adjudicated as payable prior to the specified time period (based on adjudication date), subsequently cancelled in the specified period and submitted manually.
3 | ADNFSPELAT | adjud. nullified same-period electronic amount | Identifies the total net amount of all Invoice Groupings that were adjudicated as payable during the specified time period (based on adjudication date), subsequently nullified in the specified period and submitted electronically.
3 | ADNFSPELCT | adjud. nullified same-period electronic count | Identifies the total number of all Invoice Groupings that were adjudicated as payable during the specified time period (based on adjudication date), subsequently nullified in the specified period and submitted electronically.
3 | ADNFSPMNAT | adjud. nullified same-period manual amount | Identifies the total net amount of all Invoice Groupings that were adjudicated as payable during the specified time period (based on adjudication date), subsequently cancelled in the specified period and submitted manually.
3 | ADNFSPMNCT | adjud. nullified same-period manual count | Identifies the total number of all Invoice Groupings that were adjudicated as payable during the specified time period (based on adjudication date), subsequently cancelled in the specified period and submitted manually.
3 | ADNPPPELAT | adjud. non-payee payable prior-period electronic amount | Identifies the total net amount of all Invoice Groupings that were adjudicated as payable prior to the specified time period (based on adjudication date) that do not match a specified payee (e.g. pay patient) and submitted electronically.
3 | ADNPPPELCT | adjud. non-payee payable prior-period electronic count | Identifies the total number of all Invoice Groupings that were adjudicated as payable prior to the specified time period (based on adjudication date) that do not match a specified payee (e.g. pay patient) and submitted electronically.
3 | ADNPPPMNAT | adjud. non-payee payable prior-period manual amount | Identifies the total net amount of all Invoice Groupings that were adjudicated as payable prior to the specified time period (based on adjudication date) that do not match a specified payee (e.g. pay patient) and submitted manually.
3 | ADNPPPMNCT | adjud. non-payee payable prior-period manual count | Identifies the total number of all Invoice Groupings that were adjudicated as payable prior to the specified time period (based on adjudication date) that do not match a specified payee (e.g. pay patient) and submitted manually.

3 | ADNPSPELAT | adjud. non-payee payable same-period electronic amount | Identifies the total net amount of all Invoice Groupings that were adjudicated as payable during the specified time period (based on adjudication date) that do not match a specified payee (e.g. pay patient) and submitted electronically.
3 | ADNPSPELCT | adjud. non-payee payable same-period electronic count | Identifies the total number of all Invoice Groupings that were adjudicated as payable during the specified time period (based on adjudication date) that do not match a specified payee (e.g. pay patient) and submitted electronically.
3 | ADNPSPMNAT | adjud. non-payee payable same-period manual amount | Identifies the total net amount of all Invoice Groupings that were adjudicated as payable during the specified time period (based on adjudication date) that do not match a specified payee (e.g. pay patient) and submitted manually.
3 | ADNPSPMNCT | adjud. non-payee payable same-period manual count | Identifies the total number of all Invoice Groupings that were adjudicated as payable during the specified time period (based on adjudication date) that do not match a specified payee (e.g. pay patient) and submitted manually.
3 | ADPPPPELAT | adjud. payee payable prior-period electronic amount | Identifies the total net amount of all Invoice Groupings that were adjudicated as payable prior to the specified time period (based on adjudication date) that match a specified payee (e.g. pay provider) and submitted electronically.
3 | ADPPPPELCT | adjud. payee payable prior-period electronic count | Identifies the total number of all Invoice Groupings that were adjudicated as payable prior to the specified time period (based on adjudication date) that match a specified payee (e.g. pay provider) and submitted electronically.
3 | ADPPPPMNAT | adjud. payee payable prior-period manual amout | Identifies the total net amount of all Invoice Groupings that were adjudicated as payable prior to the specified time period (based on adjudication date) that match a specified payee (e.g. pay provider) and submitted manually.
3 | ADPPPPMNCT | adjud. payee payable prior-period manual count | Identifies the total number of all Invoice Groupings that were adjudicated as payable prior to the specified time period (based on adjudication date) that match a specified payee (e.g. pay provider) and submitted manually.
3 | ADPPSPELAT | adjud. payee payable same-period electronic amount | Identifies the total net amount of all Invoice Groupings that were adjudicated as payable during the specified time period (based on adjudication date) that match a specified payee (e.g. pay provider) and submitted electronically.
3 | ADPPSPELCT | adjud. payee payable same-period electronic count | Identifies the total number of all Invoice Groupings that were adjudicated as payable during the specified time period (based on adjudication date) that match a specified payee (e.g. pay provider) and submitted electronically.
3 | ADPPSPMNAT | adjud. payee payable same-period manual amount | Identifies the total net amount of all Invoice Groupings that were adjudicated as payable during the specified time period (based on adjudication date) that match a specified payee (e.g. pay provider) and submitted manually.
3 | ADPPSPMNCT | adjud. payee payable same-period manual count | Identifies the total number of all Invoice Groupings that were adjudicated as payable during the specified time period (based on adjudication date) that match a specified payee (e.g. pay provider) and submitted manually.
3 | ADRFPPELAT | adjud. refused prior-period electronic amount | Identifies the total net amount of all Invoice Groupings that were adjudicated as refused prior to the specified time period (based on adjudication date) and submitted electronically.
3 | ADRFPPELCT | adjud. refused prior-period electronic count | Identifies the total number of all Invoice Groupings that were adjudicated as refused prior to the specified time period (based on adjudication date) and submitted electronically.
3 | ADRFPPMNAT | adjud. refused prior-period manual amount | Identifies the total net amount of all Invoice Groupings that were adjudicated as refused prior to the specified time period (based on adjudication date) and submitted manually.
3 | ADRFPPMNCT | adjud. refused prior-period manual count | Identifies the total number of all Invoice Groupings that were adjudicated as refused prior to the specified time period (based on adjudication date) and submitted manually.
3 | ADRFSPELAT | adjud. refused same-period electronic amount | Identifies the total net amount of all Invoice Groupings that were adjudicated as refused during the specified time period (based on adjudication date) and submitted electronically.
3 | ADRFSPELCT | adjud. refused same-period electronic count | Identifies the total number of all Invoice Groupings that were adjudicated as refused during the specified time period (based on adjudication date) and submitted electronically.
3 | ADRFSPMNAT | adjud. refused same-period manual amount | Identifies the total net amount of all Invoice Groupings that were adjudicated as refused during the specified time period (based on adjudication date) and submitted manually.
3 | ADRFSPMNCT | adjud. refused same-period manual count | Identifies the total number of all Invoice Groupings that were adjudicated as refused during the specified time period (based on adjudication date) and submitted manually.
2 | (_InvoiceElementPaid) Abstract | | Total counts and total net amounts paid for all Invoice Groupings that were paid within a time period based on the payment date.
3 | PDNFPPELAT | paid nullified prior-period electronic amount | Identifies the total net amount of all Invoice Groupings that were paid prior to the specified time period (based on payment date), subsequently nullified in the specified period and submitted electronically.
3 | PDNFPPELCT | paid nullified prior-period electronic count | Identifies the total number of all Invoice Groupings that were paid prior to the specified time period (based on payment date), subsequently nullified in the specified period and submitted electronically.
3 | PDNFPPMNAT | paid nullified prior-period manual amount | Identifies the total net amount of all Invoice Groupings that were paid prior to the specified time period (based on payment date), subsequently nullified in the specified period and submitted manually.
3 | PDNFPPMNCT | paid nullified prior-period manual count | Identifies the total number of all Invoice Groupings that were paid prior to the specified time period (based on payment date), subsequently nullified in the specified period and submitted manually.
3 | PDNFSPELAT | paid nullified same-period electronic amount | Identifies the total net amount of all Invoice Groupings that were paid during the specified time period (based on payment date), subsequently nullified in the specified period and submitted electronically.
3 | PDNFSPELCT | paid nullified same-period electronic count | Identifies the total number of all Invoice Groupings that were paid during the specified time period (based on payment date), subsequently cancelled in the specified period and submitted electronically.
3 | PDNFSPMNAT | paid nullified same-period manual amount | Identifies the total net amount of all Invoice Groupings that were paid during the specified time period (based on payment date), subsequently nullified in the specified period and submitted manually.
3 | PDNFSPMNCT | paid nullified same-period manual count | Identifies the total number of all Invoice Groupings that were paid during the specified time period (based on payment date), subsequently nullified in the specified period and submitted manually.

3 | PDNPPPELAT | paid non-payee payable prior-period electronic amount | Identifies the total net amount of all Invoice Groupings that were paid prior to the specified time period (based on payment date) that do not match a specified payee (e.g. pay patient) and submitted electronically.
3 | PDNPPPELCT | paid non-payee payable prior-period electronic count | Identifies the total number of all Invoice Groupings that were paid prior to the specified time period (based on payment date) that do not match a specified payee (e.g. pay patient) and submitted electronically.
3 | PDNPPPMNAT | paid non-payee payable prior-period manual amount | Identifies the total net amount of all Invoice Groupings that were paid prior to the specified time period (based on payment date) that do not match a specified payee (e.g. pay patient) and submitted manually.
3 | PDNPPPMNCT | paid non-payee payable prior-period manual count | Identifies the total number of all Invoice Groupings that were paid prior to the specified time period (based on payment date) that do not match a specified payee (e.g. pay patient) and submitted manually.
3 | PDNPSPELAT | paid non-payee payable same-period electronic amount | Identifies the total net amount of all Invoice Groupings that were paid during the specified time period (based on payment date) that do not match a specified payee (e.g. pay patient) and submitted electronically.
3 | PDNPSPELCT | paid non-payee payable same-period electronic count | Identifies the total number of all Invoice Groupings that were paid during the specified time period (based on payment date) that do not match a specified payee (e.g. pay patient) and submitted electronically.
3 | PDNPSPMNAT | paid non-payee payable same-period manual amount | Identifies the total net amount of all Invoice Groupings that were paid during the specified time period (based on payment date) that do not match a specified payee (e.g. pay patient) and submitted manually.
3 | PDNPSPMNCT | paid non-payee payable same-period manual count | Identifies the total number of all Invoice Groupings that were paid during the specified time period (based on payment date) that do not match a specified payee (e.g. pay patient) and submitted manually.
3 | PDPPPPELAT | paid payee payable prior-period electronic amount | Identifies the total net amount of all Invoice Groupings that were paid prior to the specified time period (based on payment date) that match a specified payee (e.g. pay provider) and submitted electronically.
3 | PDPPPPELCT | paid payee payable prior-period electronic count | Identifies the total number of all Invoice Groupings that were paid prior to the specified time period (based on payment date) that match a specified payee (e.g. pay provider) and submitted electronically.
3 | PDPPPPMNAT | paid payee payable prior-period manual amount | Identifies the total net amount of all Invoice Groupings that were paid prior to the specified time period (based on payment date) that match a specified payee (e.g. pay provider) and submitted manually.
3 | PDPPPPMNCT | paid payee payable prior-period manual count | Identifies the total number of all Invoice Groupings that were paid prior to the specified time period (based on payment date) that match a specified payee (e.g. pay provider) and submitted manually.
3 | PDPPSPELAT | paid payee payable same-period electronic amount | Identifies the total net amount of all Invoice Groupings that were paid during the specified time period (based on payment date) that match a specified payee (e.g. pay provider) and submitted electronically.
3 | PDPPSPELCT | paid payee payable same-period electronic count | Identifies the total number of all Invoice Groupings that were paid during the specified time period (based on payment date) that match a specified payee (e.g. pay provider) and submitted electronically.
3 | PDPPSPMNAT | paid payee payable same-period manual amount | Identifies the total net amount of all Invoice Groupings that were paid during the specified time period (based on payment date) that match a specified payee (e.g. pay provider) and submitted manually.
3 | PDPPSPMNCT | paid payee payable same-period manual count | Identifies the total number of all Invoice Groupings that were paid during the specified time period (based on payment date) that match a specified payee (e.g. pay provider) and submitted manually.
2 | (_InvoiceElementSubmitted) Abstract | | Total counts and total net amounts billed for all Invoice Groupings that were submitted within a time period. Adjudicated invoice elements are included.
3 | SBBLELAT | submitted billed electronic amount | Identifies the total net amount billed for all submitted Invoice Groupings within a time period and submitted electronically. Adjudicated invoice elements are included.
3 | SBBLELCT | submitted billed electronic count | Identifies the total number of submitted Invoice Groupings within a time period and submitted electronically. Adjudicated invoice elements are included.
3 | SBNFELAT | submitted nullified electronic amount | Identifies the total net amount billed for all submitted Invoice Groupings that were nullified within a time period and submitted electronically. Adjudicated invoice elements are included.
3 | SBNFELCT | submitted cancelled electronic count | Identifies the total number of submitted Invoice Groupings that were nullified within a time period and submitted electronically. Adjudicated invoice elements are included.
3 | SBPDELAT | submitted pending electronic amount | Identifies the total net amount billed for all submitted Invoice Groupings that are pended or held by the payor, within a time period and submitted electronically. Adjudicated invoice elements are not included.
3 | SBPDELCT | submitted pending electronic count | Identifies the total number of submitted Invoice Groupings that are pended or held by the payor, within a time period and submitted electronically. Adjudicated invoice elements are not included.
1 | (_ActInvoiceOverrideCode) Abstract | | Includes coded responses that will occur as a result of the adjudication of an electronic invoice at a summary level and provides guidance on interpretation of the referenced adjudication results.
2 | COVGE | coverage problem | Insurance coverage problems have been encountered. Additional explanation information to be supplied.
2 | EFORM | electronic form to follow | Electronic form with supporting or additional information to follow.
2 | FAX | fax to follow | Fax with supporting or additional information to follow.
2 | GFTH | good faith indicator | The medical service was provided to a patient in good faith that they had medical coverage, although no evidence of coverage was available before service was rendered.
2 | LATE | late invoice | Knowingly over the payor's published time limit for this invoice possibly due to a previous payor's delays in processing. Additional reason information will be supplied.
2 | MANUAL | manual review | Manual review of the invoice is requested. Additional information to be supplied. This may be used in the case of an appeal.
2 | OOJ | out of jurisdiction | The medical service and/or product was provided to a patient that has coverage in another jurisdiction.

2 | ORTHO | orthodontic service | The service provided is required for orthodontic purposes. If the covered party has orthodontic coverage, then the service may be paid.
2 | PAPER | paper documentation to follow | Paper documentation (or other physical format) with supporting or additional information to follow.
2 | PIE | public insurance exhausted | Public Insurance has been exhausted. Invoice has not been sent to Public Insuror and therefore no Explanation Of Benefits (EOB) is provided with this Invoice submission.
2 | PYRDELAY | delayed by a previous payor | Allows provider to explain lateness of invoice to a subsequent payor.
2 | REFNR | referral not required | Rules of practice do not require a physician's referral for the provider to perform a billable service.
2 | REPSERV | repeated service | The same service was delivered within a time period that would usually indicate a duplicate billing. However, the repeated service is a medical necessity and therefore not a duplicate.
2 | UNRELAT | unrelated service | The service provided is not related to another billed service. For example, 2 unrelated services provided on the same day to the same patient which may normally result in a refused payment for one of the items.
2 | VERBAUTH | verbal authorization | The provider has received a verbal permission from an authoritative source to perform the service or supply the item being invoiced.
1 | (_ActListCode) Abstract | | Provides codes associated with ActClass value of LIST (working list)
2 | (_ActObservationList) Abstract
3 | CARELIST | care plan | List of acts representing a care plan. The acts can be in a varierty of moods including event (EVN) to record acts that have been carried out as part of the care plan.
3 | CONDLIST | condition list | List of condition observations.
4 | INTOLIST | intolerance list | List of intolerance observations.
4 | PROBLIST | problem list | List of problem observations.
4 | RISKLIST | risk factors | List of risk factor observations.
3 | GOALLIST | goal list | List of observations in goal mood.
2 | (_ActTherapyDurationWorkingListCode) Abstract | | Codes used to identify different types of 'duration-based' working lists. Examples include "Continuous/Chronic", "Short-Term" and "As-Needed".
3 | (_ActMedicationTherapyDurationWorkingListCode) Abstract | | Definition:A collection of concepts that identifies different types of 'duration-based' mediation working lists. Examples:"Continuous/Chronic" "Short-Term" and "As Needed"
4 | ACU | short term/acute | Definition:A list of medications which the patient is only expected to consume for the duration of the current order or limited set of orders and which is not expected to be renewed.
4 | CHRON | continuous/chronic | Definition:A list of medications which are expected to be continued beyond the present order and which the patient should be assumed to be taking unless explicitly stopped.
4 | ONET | one time | Definition:A list of medications which the patient is intended to be administered only once.
4 | PRN | as needed | Definition:A list of medications which the patient will consume intermittently based on the behavior of the condition for which the medication is indicated.
2 | MEDLIST | medication list | List of medications.
3 | CURMEDLIST | current medication list | List of current medications.
3 | DISCMEDLIST | discharge medication list | List of discharge medications.
3 | HISTMEDLIST | medication history | Historical list of medications.
1 | (_ActMonitoringProtocolCode) Abstract | | Identifies types of monitoring programs
2 | CTLSUB | Controlled Substance | A monitoring program that focuses on narcotics and/or commonly abused substances that are subject to legal restriction.
2 | INV | investigational | Definition:A monitoring program that focuses on a drug which is under investigation and has not received regulatory approval for the condition being investigated
2 | LU | limited use | Description:A drug that can be prescribed (and reimbursed) only if it meets certain criteria.
2 | OTC | non prescription medicine | Medicines designated in this way may be supplied for patient use without a prescription. The exact form of categorisation will vary in different realms.
2 | RX | prescription only medicine | Some form of prescription is required before the related medicine can be supplied for a patient. The exact form of regulation will vary in different realms.
2 | SA | special authorization | Definition:A drug that requires prior approval (to be reimbursed) before being dispensed
2 | SAC | special access | Description:A drug that requires special access permission to be prescribed and dispensed.
1 | (_ActNonObservationIndicationCode) Abstract | | Description:Concepts representing indications (reasons for clinical action) other than diagnosis and symptoms.
2 | IND01 | imaging study requiring contrast | Description:Contrast agent required for imaging study.
2 | IND02 | colonoscopy prep | Description:Provision of prescription or direction to consume a product for purposes of bowel clearance in preparation for a colonoscopy.
2 | IND03 | prophylaxis | Description:Provision of medication as a preventative measure during a treatment or other period of increased risk.
2 | IND04 | surgical prophylaxis | Description:Provision of medication during pre-operative phase; e.g., antibiotics before dental surgery or bowel prep before colon surgery.
2 | IND05 | pregnancy prophylaxis | Description:Provision of medication for pregnancy --e.g., vitamins, antibiotic treatments for vaginal tract colonization, etc.
1 | (_ActObservationVerificationType) Abstract | | Identifies the type of verification investigation being undertaken with respect to the subject of the verification activity. Examples: Verification of eligibility for coverage under a policy or program - aka enrolled/covered by a policy or program Verification of record - e.g., person has record in an immunization registry Verification of enumeration - e.g. NPI Verification of Board Certification - provider specific Verification of Certification - e.g. JAHCO, NCQA, URAC Verification of Conformance - e.g. entity use with HIPAA, conformant to the CCHIT EHR system criteria Verification of Provider Credentials Verification of no adverse findings - e.g. on National Provider Data Bank, Health Integrity Protection Data Base (HIPDB)
2 | VFPAPER | verify paper | Definition:Indicates that the paper version of the record has, should be or is being verified against the electronic version.
1 | (_ActPaymentCode) Abstract | | Code identifying the method or the movement of payment instructions. Codes are drawn from X12 data element 591 (PaymentMethodCode)
2 | ACH | Automated Clearing House | Automated Clearing House (ACH).
2 | CHK | Cheque | A written order to a bank to pay the amount specified from funds on deposit.
2 | DDP | Direct Deposit | Electronic Funds Transfer (EFT) deposit into the payee's bank account

2 | NON | Non-Payment Data | Non-Payment Data.
1 | (_ActPharmacySupplyType) Abstract | | Identifies types of dispensing events
2 | DF | Daily Fill | A fill providing sufficient supply for one day
2 | EM | Emergency Supply | A supply action where there is no 'valid' order for the supplied medication. E.g. Emergency vacation supply, weekend supply (when prescriber is unavailable to provide a renewal prescription)
3 | SO | Script Owing | An emergency supply where the expectation is that a formal order authorizing the supply will be provided at a later date.
2 | FF | First Fill | The initial fill against an order. (This includes initial fills against refill orders.)
3 | FFS
3 | FFC | First Fill - Complete | A first fill where the quantity supplied is equal to one full repetition of the ordered amount. (e.g. If the order was 90 tablets, 3 refills, a complete fill would be for the full 90 tablets).
4 | FFCS
3 | FFP | First Fill - Part Fill | A first fill where the quantity supplied is less than one full repetition of the ordered amount. (e.g. If the order was 90 tablets, 3 refills, a partial fill might be for only 30 tablets.)
3 | FFPS
3 | FFSS | first fill, partial strength | A first fill where the strength supplied is less than the ordered strength. (e.g. 10mg for an order of 50mg where a subsequent fill will dispense 40mg tablets).
4 | FFPS
4 | FFCS
4 | TFS
3 | TF | Trial Fill | A fill where a small portion is provided to allow for determination of the therapy effectiveness and patient tolerance.
4 | TFS
2 | FS | Floor stock | A supply action to restock a smaller more local dispensary.
2 | MS | Manufacturer Sample | A supply of a manufacturer sample
2 | RF | Refill | A fill against an order that has already been filled (or partially filled) at least once.
3 | DF
3 | UD | Unit Dose | A supply action that provides sufficient material for a single dose.
3 | RFC | Refill - Complete | A refill where the quantity supplied is equal to one full repetition of the ordered amount. (e.g. If the order was 90 tablets, 3 refills, a complete fill would be for the full 90 tablets.)
4 | RFCS | refill complete partial strength | A refill where the quantity supplied is equal to one full repetition of the ordered amount. (e.g. If the order was 90 tablets, 3 refills, a complete fill would be for the full 90 tablets.) and where the strength supplied is less than the ordered strength (e.g. 10mg for an order of 50mg where a subsequent fill will dispense 40mg tablets).
3 | RFF | Refill (First fill this facility) | The first fill against an order that has already been filled at least once at another facility.
4 | RFFS | refill partial strength (first fill this facility) | The first fill against an order that has already been filled at least once at another facility and where the strength supplied is less than the ordered strength (e.g. 10mg for an order of 50mg where a subsequent fill will dispense 40mg tablets).
3 | RFP | Refill - Part Fill | A refill where the quantity supplied is less than one full repetition of the ordered amount. (e.g. If the order was 90 tablets, 3 refills, a partial fill might be for only 30 tablets.)
4 | RFPS | refill part fill partial strength | A refill where the quantity supplied is less than one full repetition of the ordered amount. (e.g. If the order was 90 tablets, 3 refills, a partial fill might be for only 30 tablets.) and where the strength supplied is less than the ordered strength (e.g. 10mg for an order of 50mg where a subsequent fill will dispense 40mg tablets).
3 | RFS | refill partial strength | A fill against an order that has already been filled (or partially filled) at least once and where the strength supplied is less than the ordered strength (e.g. 10mg for an order of 50mg where a subsequent fill will dispense 40mg tablets).
3 | TB | Trial Balance | A fill where the remainder of a 'complete' fill is provided after a trial fill has been provided.
4 | TBS | trial balance partial strength | A fill where the remainder of a 'complete' fill is provided after a trial fill has been provided and where the strength supplied is less than the ordered strength (e.g. 10mg for an order of 50mg where a subsequent fill will dispense 40mg tablets).
2 | UD
2 | UDE | unit dose equivalent | A supply action that provides sufficient material for a single dose via multiple products. E.g. 2 50mg tablets for a 100mg unit dose.
1 | (_ActPolicyType) Abstract | | Description:Types of policies that further specify the ActClassPolicy value set.
2 | (_ActPrivacyPolicy) Abstract | | A policy deeming certain information to be private to an individual or organization. Definition: A mandate, obligation, requirement, rule, or expectation relating to privacy. Discussion: ActPrivacyPolicyType codes support the designation of the 1..* policies that are applicable to an Act such as a Consent Directive, a Role such as a VIP Patient, or an Entity such as a patient who is a minor. 1..* ActPrivacyPolicyType values may be associated with an Act or Role to indicate the policies that govern the assignment of an Act or Role confidentialityCode. Use of multiple ActPrivacyPolicyType values enables fine grain specification of applicable policies, but must be carefully assigned to ensure cogency and avoid creation of conflicting policy mandates. Usage Note: Statutory title may be named in the ActClassPolicy Act Act.title to specify which privacy policy is being referenced.
3 | (_ActConsentDirective) Abstract | | Specifies the type of agreement between one or more grantor and grantee in which rights and obligations related to one or more shared items of interest are allocated. Usage Note: Such agreements may be considered "consent directives" or "contracts" depending on the context, and are considered closely related or synonymous from a legal perspective. Examples: Healthcare Privacy Consent Directive permitting or restricting in whole or part the collection, access, use, and disclosure of health information, and any associated handling caveats. Healthcare Medical Consent Directive to receive medical procedures after being informed of risks and benefits, thereby reducing the grantee's liability. Research Informed Consent for participation in clinical trials and disclosure of health information after being informed of risks and benefits, thereby reducing the grantee's liability. Substitute decision maker delegation in which the grantee assumes responsibility to act on behalf of the grantor. Contracts in which the agreement requires assent/dissent by the grantor of terms offered by a grantee, a consumer opts out of an "award" system for use of a retailer's marketing or credit card vendor's point collection cards in exchange for allowing purchase tracking and profiling. A mobile device or App privacy policy and terms of service to which a user must agree in whole or in part in order to utilize the service. Agreements between a client and an authorization server or between an authorization server and a resource operator and/or resource owner permitting or restricting e.g., collection, access, use, and disclosure of information, and any associated handling caveats.

4 | EMRGONLY | emergency only | This general consent directive specifically limits disclosure of health information for purpose of emergency treatment. Additional parameters may further limit the disclosure to specific users, roles, duration, types of information, and impose uses obligations. Definition: Opt-in to disclosure of health information for emergency only consent directive.

4 | GRANTORCHOICE | grantor choice | A grantor's terms of agreement to which a grantee may assent or dissent, and which may include an opportunity for a grantee to request restrictions or extensions. Comment: A grantor typically is able to stipulate preferred terms of agreement when the grantor has control over the topic of the agreement, which a grantee must accept in full or may be offered an opportunity to extend or restrict certain terms. Usage Note: If the grantor's term of agreement must be accepted in full, then this is considered "basic consent". If a grantee is offered an opportunity to extend or restrict certain terms, then the agreement is considered "granular consent". Examples: Healthcare: A PHR account holder [grantor] may require any PHR user [grantee] to accept the terms of agreement in full, or may permit a PHR user to extend or restrict terms selected by the account holder or requested by the PHR user. Non-healthcare: The owner of a resource server [grantor] may require any authorization server [grantee] to meet authorization requirements stipulated in the grantor's terms of agreement.

4 | IMPLIED | implied consent | A grantor's presumed assent to the grantee's terms of agreement is based on the grantor's behavior, which may result from not expressly assenting to the consent directive offered, or from having no right to assent or dissent offered by the grantee. Comment: Implied or "implicit" consent occurs when the behavior of the grantor is understood by a reasonable person to signal agreement to the grantee's terms. Usage Note: Implied consent with no opportunity to assent or dissent to certain terms is considered "basic consent". Examples: Healthcare: A patient schedules an appointment with a provider, and either does not take the opportunity to expressly assent or dissent to the provider's consent directive, does not have an opportunity to do so, as in the case where emergency care is required, or simply behaves as though the patient [grantor] agrees to the rights granted to the provider [grantee] in an implicit consent directive. An injured and unconscious patient is deemed to have assented to emergency treatment by those permitted to do so under jurisdictional laws, e.g., Good Samaritan laws. Non-healthcare: Upon receiving a driver's license, the driver is deemed to have assented without explicitly consenting to undergoing field sobriety tests. A corporation that does business in a foreign nation is deemed to have deemed to have assented without explicitly consenting to abide by that nation's laws.

4 | IMPLIEDD | implied consent with opportunity to dissent | A grantor's presumed assent to the grantee's terms of agreement, which is based on the grantor's behavior, and includes a right to dissent to certain terms. Comment: A grantor assenting to the grantee's terms of agreement may or may not exercise a right to dissent to grantor selected terms or to grantee's selected terms to which a grantor may dissent. Usage Note: Implied or "implicit" consent with an "opportunity to dissent" occurs when the grantor's behavior is understood by a reasonable person to signal assent to the grantee's terms of agreement whether the grantor requests or the grantee approves further restrictions, is considered "granular consent". Examples: Healthcare Examples: A healthcare provider deems a patient's assent to disclosure of health information to family members and friends, but offers an opportunity or permits the patient to dissent to such disclosures. A health information exchanges deems a patient to have assented to disclosure of health information for treatment purposes, but offers the patient an opportunity to dissents to disclosure to particular provider organizations. Non-healthcare Examples: A bank deems a banking customer's assent to specified collection, access, use, or disclosure of financial information as a requirement of holding a bank account, but provides the user an opportunity to limit third-party collection, access, use or disclosure of that information for marketing purposes.

4 | NOCONSENT | no consent | No notification or opportunity is provided for a grantor to assent or dissent to a grantee's terms of agreement. Comment: A "No Consent" policy scheme provides no opportunity for accommodation of an individual's preferences, and may not comply with Fair Information Practice Principles [FIPP] by enabling the data subject to object, access collected information, correct errors, or have accounting of disclosures. Usage Note: The grantee's terms of agreement, may be available to the grantor by reviewing the grantee's privacy policies, but there is no notice by which a grantor is apprised of the policy directly or able to acknowledge. Examples: Healthcare: Without notification or an opportunity to assent or dissent, a patient's health information is automatically included in and available (often according to certain rules) through a health information exchange. Note that this differs from implied consent, where the patient is assumed to have consented. Without notification or an opportunity to assent or dissent, a patient's health information is collected, accessed, used, or disclosed for research, public health, security, fraud prevention, court order, or law enforcement. Non-healthcare: Without notification or an opportunity to assent or dissent, a consumer's healthcare or non-healthcare internet searches are aggregated for secondary uses such as behavioral tracking and profiling. Without notification or an opportunity to assent or dissent, a consumer's location and activities in a shopping mall are tracked by RFID tags on purchased items.

4 | NOPP | notice of privacy practices | Acknowledgement of custodian notice of privacy practices. Usage Notes: This type of consent directive acknowledges a custodian's notice of privacy practices including its permitted collection, access, use and disclosure of health information to users and for purposes of use specified.

4 | OPTIN | opt-in | A grantor's assent to the terms of an agreement offered by a grantee without an opportunity for to dissent to any terms. Comment: Acceptance of a grantee's terms pertaining, for example, to permissible activities, purposes of use, handling caveats, expiry date, and revocation policies. Usage Note: Opt-in with no opportunity for a grantor to restrict certain permissions sought by the grantee is considered "basic consent". Examples: Healthcare: A patient [grantor] signs a provider's [grantee's] consent directive form, which lists permissible collection, access, use, or disclosure activities, purposes of use, handling caveats, and revocation policies. Non-healthcare: An employee [grantor] signs an employer's [grantee's] non-disclosure and non-compete agreement.

4 | OPTINR | opt-in with restrictions | A grantor's assent to the grantee's terms of an agreement with an opportunity for to dissent to certain grantor or grantee selected terms. Comment: A grantor dissenting to the grantee's terms of agreement may or may not exercise a right to assent to grantor's pre-approved restrictions or to grantee's selected terms to which a grantor may dissent. Usage Note: Opt-in with restrictions is considered "granular consent" because the grantor has an opportunity to narrow the permissions sought by the grantee. Examples: Healthcare: A patient assent to grantee's consent directive terms for collection, access, use, or disclosure of health information, and dissents to disclosure to certain recipients as allowed by the provider's pre-approved restriction list. Non-Healthcare: A cell phone user assents to the cell phone's privacy practices and terms of use, but dissents from location tracking by turning off the cell phone's tracking capability.

4 | OPTOUT | op-out | A grantor's dissent to the terms of agreement offered by a grantee without an opportunity for to assent to any terms. Comment: Rejection of a grantee's terms of agreement pertaining, for example, to permissible activities, purposes of use, handling caveats, expiry date, and revocation policies. Usage Note: Opt-out with no opportunity for a grantor to permit certain permissions sought by the grantee is considered "basic consent". Examples: Healthcare: A patient [grantor] declines to sign a provider's [grantee's] consent directive form, which lists permissible collection, access, use, or disclosure activities, purposes of use, handling caveats, revocation policies, and consequences of not assenting. Non-healthcare: An employee [grantor] refuses to sign an employer's [grantee's] agreement not to join unions or participate in a strike where state law protects employee's collective bargaining rights. A citizen [grantor] refuses to enroll in mandatory government [grantee] health insurance based on religious beliefs, which is an exemption.

4 | OPTOUTE | opt-out with exceptions | A grantor's dissent to the grantee's terms of agreement except for certain grantor or grantee selected terms. Comment: A rejection of a grantee's terms of agreement while assenting to certain permissions sought by

the grantee or requesting approval of additional grantor terms. Usage Note: Opt-out with exceptions is considered a "granular consent" because the grantor has an opportunity to accept certain permissions sought by the grantee or request additional grantor terms, while rejecting other grantee terms. Examples: Healthcare: A patient [grantor] dissents to a health information exchange consent directive with the exception of disclosure based on a limited "time to live" shared secret [e.g., a token or password], which the patient can give to a provider when seeking care. Non-healthcare: A social media user [grantor] dissents from public access to their account, but assents to access to a circle of friends.

3 | (_ActPrivacyLaw) Abstract | | A jurisdictional mandate, regulation, obligation, requirement, rule, or expectation deeming certain information to be private to an individual or organization, which is imposed on: The activity of a governed party The behavior of a governed party The manner in which an act is executed by a governed party

4 | _ActUSPrivacyLaw | _ActUSPrivacyLaw | Definition: A jurisdictional mandate in the U.S. relating to privacy. Usage Note: ActPrivacyLaw codes may be associated with an Act or a Role to indicate the legal provision to which the assignment of an Act.confidentialityCode or Role.confidentialtyCode complies. May be used to further specify rationale for assignment of other ActPrivacyPolicy codes in the US realm, e.g., ETH and 42CFRPart2 can be differentiated from ETH and Title38Part1.

5 | 42CFRPart2 | 42 CFR Part2 | 42 CFR Part 2 stipulates the right of an individual who has applied for or been given diagnosis or treatment for alcohol or drug abuse at a federally assisted program. Definition: Non-disclosure of health information relating to health care paid for by a federally assisted substance abuse program without patient consent. Usage Note: May be associated with an Act or a Role to indicate the legal provision to which the assignment of an Act.confidentialityCode or Role.confidentialityCode complies.

5 | CommonRule | Common Rule | U.S. Federal regulations governing the protection of human subjects in research (codified at Subpart A of 45 CFR part 46) that has been adopted by 15 U.S. Federal departments and agencies in an effort to promote uniformity, understanding, and compliance with human subject protections. Existing regulations governing the protection of human subjects in Food and Drug Administration (FDA)-regulated research (21 CFR parts 50, 56, 312, and 812) are separate from the Common Rule but include similar requirements. Definition: U.S. federal laws governing research-related privacy policies. Usage Note: May be associated with an Act or a Role to indicate the legal provision to which the assignment of an Act.confidentialityCode or Role.confidentialtyCode complies.

5 | HIPAANOPP | HIPAA notice of privacy practices | The U.S. Public Law 104-191 Health Insurance Portability and Accountability Act (HIPAA) Privacy Rule (45 CFR Part 164 Subpart E) permits access, use and disclosure of certain personal health information (PHI as defined under the law) for purposes of Treatment, Payment, and Operations, and requires that the provider ask that patients acknowledge the Provider's Notice of Privacy Practices as permitted conduct under the law. Definition: Notification of HIPAA Privacy Practices. Usage Note: May be associated with an Act or a Role to indicate the legal provision to which the assignment of an Act.confidentialityCode or Role.confidentialtyCode complies.

5 | HIPAAPsyNotes | HIPAA psychotherapy notes | The U.S. Public Law 104-191 Health Insurance Portability and Accountability Act (HIPAA) Privacy Rule (45 CFR Part 164 Section 164.508) requires authorization for certain uses and disclosure of psychotherapy notes. Definition: Authorization that must be obtained for disclosure of psychotherapy notes. Usage Note: May be associated with an Act or a Role to indicate the legal provision to which the assignment of an Act.confidentialityCode or Role.confidentialityCode complies.

5 | HIPAASelfPay | HIPAA self-pay | Section 13405(a) of the Health Information Technology for Economic and Clinical Health Act (HITECH) stipulates the right of an individual to have disclosures regarding certain health care items or services for which the individual pays out of pocket in full restricted from a health plan. Definition: Non-disclosure of health information to a health plan relating to health care items or services for which an individual pays out of pocket in full. Usage Note: May be associated with an Act or a Role to indicate the legal provision to which the assignment of an Act.confidentialityCode or Role.confidentialityCode complies.

5 | Title38Section7332 | Title 38 Section 7332 | Title 38 Part 1-protected information may only be disclosed to a third party with the special written consent of the patient except where expressly authorized by 38 USC 7332. VA may disclose this information for specific purposes to: VA employees on a need to know basis - more restrictive than Privacy Act need to know; contractors who need the information in order to perform or fulfil the duties of the contract; and researchers who provide assurances that the information will not be identified in any report. This information may also be disclosed without consent where patient lacks decision-making capacity; in a medical emergency for the purpose of treating a condition which poses an immediate threat to the health of any individual and which requires immediate medical intervention; for eye, tissue, or organ donation purposes; and disclosure of HIV information for public health purposes. Definition: Title 38 Part 1 - Section 1.462 Confidentiality restrictions. (a) General. The patient records to which Sections 1.460 through 1.499 of this part apply may be disclosed or used only as permitted by these regulations and may not otherwise be disclosed or used in any civil, criminal, administrative, or legislative proceedings conducted by any Federal, State, or local authority. Any disclosure made under these regulations must be limited to that information which is necessary to carry out the purpose of the disclosure. SUBCHAPTER III--PROTECTION OF PATIENT RIGHTS Sec. 7332. Confidentiality of certain medical records (a)(1) Records of the identity, diagnosis, prognosis, or treatment of any patient or subject which are maintained in connection with the performance of any program or activity (including education, training, treatment, rehabilitation, or research) relating to drug abuse, alcoholism or alcohol abuse, infection with the human immunodeficiency virus, or sickle cell anemia which is carried out by or for the Department under this title shall, except as provided in subsections (e) and (f), be confidential, and (section 5701 of this title to the contrary notwithstanding) such records may be disclosed only for the purposes and under the circumstances expressly authorized under subsection (b). Usage Note: May be associated with an Act or a Role to indicate the legal provision to which the assignment of an Act.confidentialityCode or Role.confidentialityCode complies.

3 | (_InformationSensitivityPolicy) Abstract | | A mandate, obligation, requirement, rule, or expectation characterizing the value or importance of a resource and may include its vulnerability. (Based on ISO7498-2:1989. Note: The vulnerability of personally identifiable sensitive information may be based on concerns that the unauthorized disclosure may result in social stigmatization or discrimination.) Description: Types of Sensitivity policy that apply to Acts or Roles. A sensitivity policy is adopted by an enterprise or group of enterprises (a 'policy domain') through a formal data use agreement that stipulates the value, importance, and vulnerability of information. A sensitivity code representing a sensitivity policy may be associated with criteria such as categories of information or sets of information identifiers (e.g., a value set of clinical codes or branch in a code system hierarchy). These criteria may in turn be used for the Policy Decision Point in a Security Engine. A sensitivity code may be used to set the confidentiality code used on information about Acts and Roles to trigger the security mechanisms required to control how security principals (i.e., a person, a machine, a software application) may act on the information (e.g., collection, access, use, or disclosure). Sensitivity codes are never assigned to the transport or business envelope containing patient specific information being exchanged outside of a policy domain as this would disclose the information intended to be protected by the policy. When sensitive information is exchanged with others outside of a policy domain, the confidentiality code on the transport or business envelope conveys the receiver's responsibilities and indicates the how the information is to be safeguarded without unauthorized disclosure of the sensitive information. This ensures that sensitive information is treated by receivers as the sender intends, accomplishing interoperability without point to point negotiations. Usage Note: Sensitivity codes are not useful for interoperability outside of a policy domain because sensitivity policies are typically localized and vary drastically across policy domains even for

the same information category because of differing organizational business rules, security policies, and jurisdictional requirements. For example, an employee's sensitivity code would make little sense for use outside of a policy domain. 'Taboo' would rarely be useful outside of a policy domain unless there are jurisdictional requirements requiring that a provider disclose sensitive information to a patient directly. Sensitivity codes may be more appropriate in a legacy system's Master Files in order to notify those who access a patient's orders and observations about the sensitivity policies that apply. Newer systems may have a security engine that uses a sensitivity policy's criteria directly. The specializable InformationSensitivityPolicy Act.code may be useful in some scenarios if used in combination with a sensitivity identifier and/or Act.title.

4 | (_ActInformationSensitivityPolicy) Abstract | | Types of sensitivity policies that apply to Acts. Act.confidentialityCode is defined in the RIM as "constraints around appropriate disclosure of information about this Act, regardless of mood." Usage Note: ActSensitivity codes are used to bind information to an Act.confidentialityCode according to local sensitivity policy so that those confidentiality codes can then govern its handling across enterprises. Internally to a policy domain, however, local policies guide the access control system on how end users in that policy domain are able to use information tagged with these sensitivity values.

5 | ETH | substance abuse information sensitivity | Policy for handling alcohol or drug-abuse information, which will be afforded heightened confidentiality. Information handling protocols based on organizational policies related to alcohol or drug-abuse information that is deemed sensitive. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

5 | GDIS | genetic disease information sensitivity | Policy for handling genetic disease information, which will be afforded heightened confidentiality. Information handling protocols based on organizational policies related to genetic disease information that is deemed sensitive. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

5 | HIV | HIV/AIDS information sensitivity | Policy for handling HIV or AIDS information, which will be afforded heightened confidentiality. Information handling protocols based on organizational policies related to HIV or AIDS information that is deemed sensitive. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

5 | MST | military sexual trauma information sensitivity | Policy for handling information related to sexual assault or repeated, threatening sexual harassment that occurred while the patient was in the military, which is afforded heightened confidentiality. Access control concerns for military sexual trauma is based on the patient being subject to control by a higher ranking military perpetrator and/or censure by others within the military unit. Due to the relatively unfettered access to healthcare information by higher ranking military personnel and those who have command over the patient, there is a need to sequester this information outside of the typical controls on access to military health records. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law in addition to this more generic code.

5 | SCA | sickle cell anemia information sensitivity | Policy for handling sickle cell disease information, which is afforded heightened confidentiality. Information handling protocols are based on organizational policies related to sickle cell disease information, which is deemed sensitive. Usage Note: If there is a jurisdictional mandate, then the Act valued with this ActCode should be associated with an Act valued with any applicable laws from the ActPrivacyLaw code system.

5 | SDV | sexual assault, abuse, or domestic violence information sensitivity | Policy for handling sexual assault, abuse, or domestic violence information, which will be afforded heightened confidentiality. Information handling protocols based on organizational policies related to sexual assault, abuse, or domestic violence information that is deemed sensitive. SDV code covers violence perpetrated by related and non-related persons. This code should be specific to physical and mental trauma caused by a related person only. The access control concerns are keeping the patient safe from the perpetrator who may have an abusive psychological control over the patient, may be stalking the patient, or may try to manipulate care givers into allowing the perpetrator to make contact with the patient. The definition needs to be clarified. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

5 | SEX | sexuality and reproductive health information sensitivity | Policy for handling sexuality and reproductive health information, which will be afforded heightened confidentiality. Information handling protocols based on organizational policies related to sexuality and reproductive health information that is deemed sensitive. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

5 | SPI | specially protected information sensitivity | Policy for handling information deemed specially protected by law or policy including substance abuse, substance use, psychiatric, mental health, behavioral health, and cognitive disorders, which is afforded heightened confidentiality. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law in addition to this more generic code.

6 | BH | behavioral health information sensitivity | Policy for handling information related to behavioral and emotional disturbances affecting social adjustment and physical health, which is afforded heightened confidentiality. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law in addition to this more generic code.

7 | COGN | cognitive disability information sensitivity | Policy for handling information related to cognitive disability disorders and conditions caused by these disorders, which are afforded heightened confidentiality. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law in addition to this more generic code. Examples may include dementia, traumatic brain injury, attention deficit, hearing and visual disability such as dyslexia and other disorders and related conditions which impair learning and self-sufficiency. However, the cognitive disabilities to which this term may apply versus other behavioral health categories varies by jurisdiction and organizational policy in part due to overlap with other behavioral health conditions. Implementers should constrain to those diagnoses applicable in the domain in which this code is used.

7 | DVD | developmental disability information sensitivity | Policy for handling information related to developmental disability disorders and conditions caused by these disorders, which is afforded heightened confidentiality. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law in addition to this more generic code. A diverse group of chronic conditions that are due to mental or physical impairments impacting activities of daily living, self-care, language acuity, learning, mobility, independent living and economic self-sufficiency. Examples may include Down syndrome and Autism spectrum. However, the developmental disabilities to which this term applies versus other behavioral health categories varies by jurisdiction and organizational policy in part due to overlap with other behavioral health conditions. Implementers should constrain to those diagnoses applicable in the domain in which this code is used.

7 | EMOTDIS | emotional disturbance information sensitivity | Policy for handling information related to emotional disturbance disorders and conditions caused by these disorders, which is afforded heightened confidentiality. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law in addition to this more generic code. Typical used to characterize behavioral and mental health issues of adolescents where the disorder may be temporarily diagnosed in order to avoid the potential and unnecessary stigmatizing diagnoses of disorder long term.

6 | MH | mental health information sensitivity | Policy for handling information related to psychological disorders, which is afforded heightened confidentiality. Mental health information may be deemed specifically sensitive and distinct from physical health, substance use disorders, and behavioral disabilities and disorders in some jurisdictions. Usage Note: If there is a jurisdictional

mandate, then use the applicable ActPrivacyLaw code system, and specify the law in addition to this more generic code.
6 | PSY | psychiatry disorder information sensitivity | Policy for handling psychiatry psychiatric disorder information, which is afforded heightened confidentiality. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.
6 | PSYTHPN | psychotherapy note information sensitivity | Policy for handling psychotherapy note information, which is afforded heightened confidentiality. Usage Note: In some jurisdiction, disclosure of psychotherapy notes requires patient consent. If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.
6 | SUD | substance use disorder information sensitivity | Policy for handling information related to alcohol or drug use disorders and conditions caused by these disorders, which is afforded heightened confidentiality. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law in addition to this more generic code.
7 | ETHUD | alcohol use disorder information sensitivity | Policy for handling information related to alcohol use disorders and conditions caused by these disorders, which is afforded heightened confidentiality. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law in addition to this more generic code.
7 | OPIOIDUD | opioid use disorder information sensitivity | Policy for handling information related to opioid use disorders and conditions caused by these disorders, which is afforded heightened confidentiality. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law in addition to this more generic code.
5 | STD | sexually transmitted disease information sensitivity | Policy for handling sexually transmitted disease information, which will be afforded heightened confidentiality. Information handling protocols based on organizational policies related to sexually transmitted disease information that is deemed sensitive. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.
5 | TBOO | taboo | Policy for handling information not to be initially disclosed or discussed with patient except by a physician assigned to patient in this case. Information handling protocols based on organizational policies related to sensitive patient information that must be initially discussed with the patient by an attending physician before being disclosed to the patient. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code. Open Issue: This definition conflates a rule and a characteristic, and there may be a similar issue with ts sibling codes.
5 | VIO | violence information sensitivity | Policy for handling information related to harm by violence, which is afforded heightened confidentiality. Harm by violence is perpetrated by an unrelated person. Access control concerns for information about mental or physical harm resulting from violence caused by an unrelated person may include manipulation of care givers or access to records that enable the perpetrator contact or locate the patient, but the perpetrator will likely not have established abusive psychological control over the patient. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law in addition to this more generic code.
5 | SICKLE | sickle cell | Types of sensitivity policies that apply to Acts. Act.confidentialityCode is defined in the RIM as "constraints around appropriate disclosure of information about this Act, regardless of mood." Usage Note: ActSensitivity codes are used to bind information to an Act.confidentialityCode according to local sensitivity policy so that those confidentiality codes can then govern its handling across enterprises. Internally to a policy domain, however, local policies guide the access control system on how end users in that policy domain are able to use information tagged with these sensitivity values.
4 | (_EntitySensitivityPolicyType) Abstract | | Types of sensitivity policies that may apply to a sensitive attribute on an Entity. Usage Note: EntitySensitivity codes are used to convey a policy that is applicable to sensitive information conveyed by an entity attribute. May be used to bind a Role.confidentialityCode associated with an Entity per organizational policy. Role.confidentialityCode is defined in the RIM as "an indication of the appropriate disclosure of information about this Role with respect to the playing Entity."
5 | DEMO | all demographic information sensitivity | Policy for handling all demographic information about an information subject, which will be afforded heightened confidentiality. Policies may govern sensitivity of information related to all demographic about an information subject, the disclosure of which could impact the privacy, well-being, or safety of that subject. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.
5 | DOB | date of birth information sensitivity | Policy for handling information related to an information subject's date of birth, which will be afforded heightened confidentiality.Policies may govern sensitivity of information related to an information subject's date of birth, the disclosure of which could impact the privacy, well-being, or safety of that subject. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.
5 | GENDER | gender and sexual orientation information sensitivity | Policy for handling information related to an information subject's gender and sexual orientation, which will be afforded heightened confidentiality. Policies may govern sensitivity of information related to an information subject's gender and sexual orientation, the disclosure of which could impact the privacy, well-being, or safety of that subject. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.
5 | LIVARG | living arrangement information sensitivity | Policy for handling information related to an information subject's living arrangement, which will be afforded heightened confidentiality. Policies may govern sensitivity of information related to an information subject's living arrangement, the disclosure of which could impact the privacy, well-being, or safety of that subject. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.
5 | MARST | marital status information sensitivity | Policy for handling information related to an information subject's marital status, which will be afforded heightened confidentiality. Policies may govern sensitivity of information related to an information subject's marital status, the disclosure of which could impact the privacy, well-being, or safety of that subject. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.
5 | RACE | race information sensitivity | Policy for handling information related to an information subject's race, which will be afforded heightened confidentiality. Policies may govern sensitivity of information related to an information subject's race, the disclosure of which could impact the privacy, well-being, or safety of that subject. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.
5 | REL | religion information sensitivity | Policy for handling information related to an information subject's religious affiliation, which will be afforded heightened confidentiality. Policies may govern sensitivity of information related to an information subject's religion, the disclosure of which could impact the privacy, well-being, or safety of that subject. Usage Notes: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

4 | (_RoleInformationSensitivityPolicy) Abstract | | Types of sensitivity policies that apply to Roles. Usage Notes: RoleSensitivity codes are used to bind information to a Role.confidentialityCode per organizational policy. Role.confidentialityCode is defined in the RIM as "an indication of the appropriate disclosure of information about this Role with respect to the playing Entity."
5 | B | business information sensitivity | Policy for handling trade secrets such as financial information or intellectual property, which will be afforded heightened confidentiality. Description: Since the service class can represent knowledge structures that may be considered a trade or business secret, there is sometimes (though rarely) the need to flag those items as of business level confidentiality. Usage Notes: No patient related information may ever be of this confidentiality level. If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.
5 | EMPL | employer information sensitivity | Policy for handling information related to an employer which is deemed classified to protect an employee who is the information subject, and which will be afforded heightened confidentiality. Description: Policies may govern sensitivity of information related to an employer, such as law enforcement or national security, the identity of which could impact the privacy, well-being, or safety of an information subject who is an employee. Usage Notes: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.
5 | LOCIS | location information sensitivity | Policy for handling information related to the location of the information subject, which will be afforded heightened confidentiality. Description: Policies may govern sensitivity of information related to the location of the information subject, the disclosure of which could impact the privacy, well-being, or safety of that subject. Usage Notes: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.
5 | SSP | sensitive service provider information sensitivity | Policy for handling information related to a provider of sensitive services, which will be afforded heightened confidentiality. Description: Policies may govern sensitivity of information related to providers who deliver sensitive healthcare services in order to protect the privacy, well-being, and safety of the provider and of patients receiving sensitive services. Usage Notes: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.
4 | ADOL | adolescent information sensitivity | Policy for handling information related to an adolescent, which will be afforded heightened confidentiality per applicable organizational or jurisdictional policy. An enterprise may have a policy that requires that adolescent patient information be provided heightened confidentiality. Information deemed sensitive typically includes health information and patient role information including patient status, demographics, next of kin, and location. Usage Note: For use within an enterprise in which an adolescent is the information subject. If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.
4 | CEL | celebrity information sensitivity | Policy for handling information related to a celebrity (people of public interest (VIP), which will be afforded heightened confidentiality. Celebrities are people of public interest (VIP) about whose information an enterprise may have a policy that requires heightened confidentiality. Information deemed sensitive may include health information and patient role information including patient status, demographics, next of kin, and location. Usage Note: For use within an enterprise in which the information subject is deemed a celebrity or very important person. If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.
4 | DIA | diagnosis information sensitivity | Policy for handling information related to a diagnosis, health condition or health problem, which will be afforded heightened confidentiality. Diagnostic, health condition or health problem related information may be deemed sensitive by organizational policy, and require heightened confidentiality. Usage Note: For use within an enterprise that provides heightened confidentiality to diagnostic, health condition or health problem related information deemed sensitive. If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.
4 | DRGIS | drug information sensitivity | Policy for handling information related to a drug, which will be afforded heightened confidentiality. Drug information may be deemed sensitive by organizational policy, and require heightened confidentiality. Usage Note: For use within an enterprise that provides heightened confidentiality to drug information deemed sensitive. If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.
4 | EMP | employee information sensitivity | Policy for handling information related to an employee, which will be afforded heightened confidentiality. When a patient is an employee, an enterprise may have a policy that requires heightened confidentiality. Information deemed sensitive typically includes health information and patient role information including patient status, demographics, next of kin, and location. Usage Note: Policy for handling information related to an employee, which will be afforded heightened confidentiality. Description: When a patient is an employee, an enterprise may have a policy that requires heightened confidentiality. Information deemed sensitive typically includes health information and patient role information including patient status, demographics, next of kin, and location.
4 | PDS | patient default information sensitivity | Policy for specially protecting information reported by or about a patient, which is deemed sensitive within the enterprise (i.e., by default regardless of whether the patient requested that the information be deemed sensitive for another reason.) For example information reported by the patient about another person, e.g., a family member, may be deemed sensitive by default. Organizational policy may allow the sensitivity tag to be cleared on patient's request. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law in addition to this more generic code. For example, VA deems employee information sensitive by default. Information about a patient who is being stalked or a victim of abuse or violence may be deemed sensitive by default per a provider organization's policies.
4 | PHY | physician requested information sensitivity | Policy for handling information about a patient, which a physician or other licensed healthcare provider deems sensitive. Once tagged by the provider, this may trigger alerts for follow up actions according to organizational policy or jurisdictional law. Usage Note: For use within an enterprise that provides heightened confidentiality to certain types of information designated by a physician as sensitive. If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code. Use cases in which this code could be used are, e.g., in systems that lack the ability to automatically detect sensitive information and must rely on manual tagging; a system that lacks an applicable sensitivity tag, or for ad hoc situations where criticality of the situation requires that the tagging be done immediately by the provider before coding or transcription of consult notes can be completed, e.g., upon detection of a patient with suicidal tendencies or potential for violence.
4 | PRS | patient requested information sensitivity | Policy for specially protecting information reported by or about a patient, which the patient deems sensitive, and the patient requests that collection, access, use, or disclosure of that information be restricted. For example, a minor patient may request that information about reproductive health not be disclosed to the patient's family or to particular providers and payers. Usage Note: If there is a jurisdictional mandate, then use the applicable ActPrivacyLaw code system, and specify the law rather than or in addition to this more generic code.

3 | COMPT | compartment | This is the healthcare analog to the US Intelligence Community's concept of a Special Access Program. Compartment codes may be used in as a field value in an initiator's clearance to indicate permission to access and use an IT Resource with a security label having the same compartment value in security category label field. Map: Aligns with ISO 2382-8 definition of Compartment - "A division of data into isolated blocks with separate security controls for the purpose of reducing risk."

4 | ACOCOMPT | accountable care organization compartment | A group of health care entities, which may include health care providers, care givers, hospitals, facilities, health plans, and other health care constituents who coordinate care for reimbursement based on quality metrics for improving outcomes and lowering costs, and may be authorized to access the consumer's health information because of membership in that group. Security Compartment Labels assigned to a consumer's information use in accountable care workflows should be met or exceeded by the Security Compartment attribute claimed by a participant in a an accountable care workflow who is requesting access to that information

4 | CTCOMPT | care team compartment | Care coordination across participants in a care plan requires sharing of a healthcare consumer's information specific to that workflow. A care team member should only have access to that information while participating in that workflow or for other authorized uses. Security Compartment Labels assigned to a consumer's information use in care coordination workflows should be met or exceeded by the Security Compartment attribute claimed by a participant in a care team member workflow who is requesting access to that information

4 | FMCOMPT | financial management compartment | Financial management department members who have access to healthcare consumer information as part of a patient account, billing and claims workflows. Security Compartment Labels assigned to consumer information used in these workflows should be met or exceeded by the Security Compartment attribute claimed by a participant in a financial management workflow who is requesting access to that information.

4 | HRCOMPT | human resource compartment | A security category label field value, which indicates that access and use of an IT resource is restricted to members of human resources department or workflow.

4 | LRCOMPT | legitimate relationship compartment | Providers and care givers who have an established relationship per criteria determined by policy are considered to have an established care provision relations with a healthcare consumer, and may be authorized to access the consumer's health information because of that relationship. Providers and care givers should only have access to that information while participating in legitimate relationship workflows or for other authorized uses. Security Compartment Labels assigned to a consumer's information use in legitimate relationship workflows should be met or exceeded by the Security Compartment attribute claimed by a participant in a legitimate relationship workflow who is requesting access to that information.

4 | PACOMPT | patient administration compartment | Patient administration members who have access to healthcare consumer information as part of a patient administration workflows. Security Compartment Labels assigned to consumer information used in these workflows should be met or exceeded by the Security Compartment attribute claimed by a participant in a patient administration workflow who is requesting access to that information.

4 | RESCOMPT | research project compartment | A security category label field value, which indicates that access and use of an IT resource is restricted to members of a research project.

4 | RMGTCOMPT | records management compartment | A security category label field value, which indicates that access and use of an IT resource is restricted to members of records management department or workflow.

2 | (ActTrustPolicyType) Abstract | | A mandate, obligation, requirement, rule, or expectation conveyed as security metadata between senders and receivers required to establish the reliability, authenticity, and trustworthiness of their transactions. Trust security metadata are observation made about aspects of trust applicable to an IT resource (data, information object, service, or system capability). Trust applicable to IT resources is established and maintained in and among security domains, and may be comprised of observations about the domain's trust authority, trust framework, trust policy, trust interaction rules, means for assessing and monitoring adherence to trust policies, mechanisms that enforce trust, and quality and reliability measures of assurance in those mechanisms. [Based on ISO IEC 10181-1 and NIST SP 800-63-2] For example, identity proofing , level of assurance, and Trust Framework.

3 | TRSTACCRD | trust accreditation | Type of security metadata about the formal declaration by an authority or neutral third party that validates the technical, security, trust, and business practice conformance of Trust Agents to facilitate security, interoperability, and trust among participants within a security domain or trust framework.

3 | TRSTAGRE | trust agreement | Type of security metadata about privacy and security requirements with which a security domain must comply. [ISO IEC 10181-1]

3 | TRSTASSUR | trust assurance | Type of security metadata about the digital quality or reliability of a trust assertion, activity, capability, information exchange, mechanism, process, or protocol.

3 | TRSTCERT | trust certificate | Type of security metadata about a set of security-relevant data issued by a security authority or trusted third party, together with security information which is used to provide the integrity and data origin authentication services for an IT resource (data, information object, service, or system capability). [Based on ISO IEC 10181-1]

3 | TRSTFWK | trust framework | Type of security metadata about a complete set of contracts, regulations, or commitments that enable participating actors to rely on certain assertions by other actors to fulfill their information security requirements. [Kantara Initiative]

3 | TRSTMEC | trust mechanism | Type of security metadata about a security architecture system component that supports enforcement of security policies.

2 | COVPOL | benefit policy | Description:A mandate, obligation, requirement, rule, or expectation unilaterally imposed on benefit coverage under a policy or program by a sponsor, underwriter or payor on: The activity of another party The behavior of another party The manner in which an act is executed Examples:A clinical protocol imposed by a payer to which a provider must adhere in order to be paid for providing the service. A formulary from which a provider must select prescribed drugs in order for the patient to incur a lower copay.

2 | SecurityPolicy | security policy | Types of security policies that further specify the ActClassPolicy value set. Examples: obligation to encrypt refrain from redisclosure without consent

3 | AUTHPOL | authorization policy | Authorisation policies are essentially security policies related to access-control and specify what activities a subject is permitted or forbidden to do, to a set of target objects. They are designed to protect target objects so are interpreted by access control agents or the run-time systems at the target system. A positive authorisation policy defines the actions that a subject is permitted to perform on a target. A negative authorisation policy specifies the actions that a subject is forbidden to perform on a target. Positive authorisation policies may also include filters to transform the parameters associated with their actions. (Based on PONDERS)

4 | ACCESSCONSCHEME | access control scheme | An access control policy specific to the type of access control scheme, which is used to enforce one or more authorization policies. Usage Note: Access control schemes are the type of access control policy, which is comprised of access control policy rules concerning the provision of the access control service. There are two categories of access control policies, rule-based and identity-based, which are identified in CCITT Rec. X.800 aka ISO 7498-2. Rule-based access control policies are intended to apply to all access requests by any initiator on any target in a security domain.

Identity-based access control policies are based on rules specific to an individual initiator, a group of initiators, entities acting on behalf of initiators, or originators acting in a specific role. Context can modify rule-based or identity-based access control policies. Context rules may define the entire policy in effect. Real systems will usually employ a combination of these policy types; if a rule-based policy is used, then an identity-based policy is usually in effect also. An access control scheme may be based on access control lists, capabilities, labels, and context or a combination of these. An access control scheme is a component of an access control mechanism or "service") along with the supporting mechanisms required by that scheme to provide access control decision information (ADI) supplied by the scheme to the access decision facility (ADF also known as a PDP). (Based on ISO/IEC 10181-3:1996) Examples: Attribute Based Access Control (ABAC) Discretionary Access Control (DAC) History Based Access Control (HBAC) Identity Based Access Control (IBAC) Mandatory Access Control (MAC) Organization Based Access Control (OrBAC) Relationship Based Access Control (RelBac) Responsibility Based Access Control (RespBAC) Risk Adaptable Access Control (RAdAC) >

3 | DELEPOL | delegation policy | Delegation policies specify which actions subjects are allowed to delegate to others. A delegation policy thus specifies an authorisation to delegate. Subjects must already possess the access rights to be delegated. Delegation policies are aimed at subjects delegating rights to servers or third parties to perform actions on their behalf and are not meant to be the means by which security administrators would assign rights to subjects. A negative delegation policy identifies what delegations are forbidden. A Delegation policy specifies the authorisation policy from which delegated rights are derived, the grantors, which are the entities which can delegate these access rights, and the grantees, which are the entities to which the access rights can be delegated. There are two types of delegation policy, positive and negative. (Based on PONDERS)

3 | ObligationPolicy | obligation policy | Conveys the mandated workflow action that an information custodian, receiver, or user must perform. Usage Notes: Per ISO 22600-2, ObligationPolicy instances 'are event-triggered and define actions to be performed by manager agent'. Per HL7 Composite Security and Privacy Domain Analysis Model: This value set refers to the action required to receive the permission specified in the privacy rule. Per OASIS XACML, an obligation is an operation specified in a policy or policy that is performed in conjunction with the enforcement of an access control decision.

4 | ANONY | anonymize | Custodian system must remove any information that could result in identifying the information subject.

4 | AOD | accounting of disclosure | Custodian system must make available to an information subject upon request an accounting of certain disclosures of the individualâ€™s protected health information over a period of time. Policy may dictate that the accounting include information about the information disclosed, the date of disclosure, the identification of the receiver, the purpose of the disclosure, the time in which the disclosing entity must provide a response and the time period for which accountings of disclosure can be requested.

4 | AUDIT | audit | Custodian system must monitor systems to ensure that all users are authorized to operate on information objects.

4 | AUDTR | audit trail | Custodian system must monitor and maintain retrievable log for each user and operation on information.

4 | CPLYCC | comply with confidentiality code | Custodian security system must retrieve, evaluate, and comply with the information handling directions of the Confidentiality Code associated with an information target.

4 | CPLYCD | comply with consent directive | Custodian security system must retrieve, evaluate, and comply with applicable information subject consent directives.

4 | CPLYJPP | comply with jurisdictional privacy policy | Custodian security system must retrieve, evaluate, and comply with applicable jurisdictional privacy policies associated with the target information.

4 | CPLYOPP | comply with organizational privacy policy | Custodian security system must retrieve, evaluate, and comply with applicable organizational privacy policies associated with the target information.

4 | CPLYOSP | comply with organizational security policy | Custodian security system must retrieve, evaluate, and comply with the organizational security policies associated with the target information.

4 | CPLYPOL | comply with policy | Custodian security system must retrieve, evaluate, and comply with applicable policies associated with the target information.

4 | DECLASSIFYLABEL | declassify security label | Custodian security system must declassify information assigned security labels by instantiating a new version of the classified information so as to break the binding of the classifying security label when assigning a new security label that marks the information as unclassified in accordance with applicable jurisdictional privacy policies associated with the target information. The system must retain an immutable record of the previous assignment and binding.

4 | DEID | deidentify | Custodian system must strip information of data that would allow the identification of the source of the information or the information subject.

4 | DELAU | delete after use | Custodian system must remove target information from access after use.

4 | DOWNGRDLABEL | downgrade security label | Custodian security system must downgrade information assigned security labels by instantiating a new version of the classified information so as to break the binding of the classifying security label when assigning a new security label that marks the information as classified at a less protected level in accordance with applicable jurisdictional privacy policies associated with the target information. The system must retain an immutable record of the previous assignment and binding.

4 | DRIVLABEL | derive security label | Custodian security system must assign and bind security labels derived from compilations of information by aggregation or disaggregation in order to classify information compiled in the information systems under its control for collection, access, use and disclosure in accordance with applicable jurisdictional privacy policies associated with the target information. The system must retain an immutable record of the previous assignment and binding.

4 | ENCRYPT | encrypt | Custodian system must render information unreadable by algorithmically transforming plaintext into ciphertext. Usage Notes: A mathematical transposition of a file or data stream so that it cannot be deciphered at the receiving end without the proper key. Encryption is a security feature that assures that only the parties who are supposed to be participating in a videoconference or data transfer are able to do so. It can include a password, public and private keys, or a complex combination of all. (Per Infoway.)

5 | ENCRYPTR | encrypt at rest | Custodian system must render information unreadable and unusable by algorithmically transforming plaintext into ciphertext when "at rest" or in storage.

5 | ENCRYPTT | encrypt in transit | Custodian system must render information unreadable and unusable by algorithmically transforming plaintext into ciphertext while "in transit" or being transported by any means.

5 | ENCRYPTU | encrypt in use | Custodian system must render information unreadable and unusable by algorithmically transforming plaintext into ciphertext while in use such that operations permitted on the target information are limited by the license granted to the end user.

4 | HUAPRV | human approval | Custodian system must require human review and approval for permission requested.

4 | LABEL | assign security label | Custodian security system must assign and bind security labels in order to classify information created in the information systems under its control for collection, access, use and disclosure in accordance with applicable jurisdictional privacy policies associated with the target information. The system must retain an immutable record of the

assignment and binding. Usage Note: In security systems, security policy label assignments do not change, they may supersede prior assignments, and such reassignments are always tracked for auditing and other purposes.
4 | MASK | mask | Custodian system must render information unreadable and unusable by algorithmically transforming plaintext into ciphertext. User may be provided a key to decrypt per license or "shared secret".
4 | MINEC | minimum necessary | Custodian must limit access and disclosure to the minimum information required to support an authorized user's purpose of use. Usage Note: Limiting the information available for access and disclosure to that an authorized user or receiver "needs to know" in order to perform permitted workflow or purpose of use.
4 | PERSISTLABEL | persist security label | Custodian security system must persist the binding of security labels to classify information received or imported by information systems under its control for collection, access, use and disclosure in accordance with applicable jurisdictional privacy policies associated with the target information. The system must retain an immutable record of the assignment and binding.
4 | PRIVMARK | privacy mark | Custodian must create and/or maintain human readable security label tags as required by policy. Map: Aligns with ISO 22600-3 Section A.3.4.3 description of privacy mark: "If present, the privacy-mark is not used for access control. The content of the privacy-mark may be defined by the security policy in force (identified by the security-policy-identifier) which may define a list of values to be used. Alternately, the value may be determined by the originator of the security-label."
4 | PSEUD | pseudonymize | Custodian system must strip information of data that would allow the identification of the source of the information or the information subject. Custodian may retain a key to relink data necessary to reidentify the information subject.
4 | REDACT | redact | Custodian system must remove information, which is not authorized to be access, used, or disclosed from records made available to otherwise authorized users.
4 | UPGRDLABEL | upgrade security label | Custodian security system must declassify information assigned security labels by instantiating a new version of the classified information so as to break the binding of the classifying security label when assigning a new security label that marks the information as classified at a more protected level in accordance with applicable jurisdictional privacy policies associated with the target information. The system must retain an immutable record of the previous assignment and binding.
3 | RefrainPolicy | refrain policy | Conveys prohibited actions which an information custodian, receiver, or user is not permitted to perform unless otherwise authorized or permitted under specified circumstances. Usage Notes: ISO 22600-2 species that a Refrain Policy "defines actions the subjects must refrain from performing". Per HL7 Composite Security and Privacy Domain Analysis Model: May be used to indicate that a specific action is prohibited based on specific access control attributes e.g., purpose of use, information type, user role, etc.
4 | NOAUTH | no disclosure without subject authorization | Prohibition on disclosure without information subject's authorization.
4 | NOCOLLECT | no collection | Prohibition on collection or storage of the information.
4 | NODSCLCD | no disclosure without consent directive | Prohibition on disclosure without organizational approved patient restriction.
4 | NODSCLCDS | no disclosure without information subject's consent directive | Prohibition on disclosure without a consent directive from the information subject.
4 | NOINTEGRATE | no integration | Prohibition on Integration into other records.
4 | NOLIST | no unlisted entity disclosure | Prohibition on disclosure except to entities on specific access list.
4 | NOMOU | no disclosure without MOU | Prohibition on disclosure without an interagency service agreement or memorandum of understanding (MOU).
4 | NOORGPOL | no disclosure without organizational authorization | Prohibition on disclosure without organizational authorization.
4 | NOPAT | no disclosure to patient, family or caregivers without attending provider's authorization | Prohibition on disclosing information to patient, family or caregivers without attending provider's authorization. Usage Note: The information may be labeled with the ActInformationSensitivity TBOO code, triggering application of this RefrainPolicy code as a handling caveat controlling access. Maps to FHIR NOPAT: Typically, this is used on an Alert resource, when the alert records information on patient abuse or non-compliance. FHIR print name is "keep information from patient". Maps to the French realm - code: INVISIBLE_PATIENT. displayName: Document non visible par le patient codingScheme: 1.2.250.1.213.1.1.4.13 French use case: A label for documents that the author chose to hide from the patient until the content can be disclose to the patient in a face to face meeting between a healthcare professional and the patient (in French law some results like cancer diagnosis or AIDS diagnosis must be announced to the patient by a healthcare professional and should not be find out by the patient alone).
4 | NOPERSISTP | no collection beyond purpose of use | Prohibition on collection of the information beyond time necessary to accomplish authorized purpose of use is prohibited.
4 | NORDSCLCD | no redisclosure without consent directive | Prohibition on redisclosure without patient consent directive.
4 | NORDSCLCDS | no redisclosure without information subject's consent directive | Prohibition on redisclosure without a consent directive from the information subject.
4 | NORDSCLW | no disclosure without jurisdictional authorization | Prohibition on disclosure without authorization under jurisdictional law.
4 | NORELINK | no relinking | Prohibition on associating de-identified or pseudonymized information with other information in a manner that could or does result in disclosing information intended to be masked.
4 | NOREUSE | no reuse beyond purpose of use | Prohibition on use of the information beyond the purpose of use initially authorized.
4 | NOVIP | no unauthorized VIP disclosure | Prohibition on disclosure except to principals with access permission to specific VIP information.
4 | ORCON | no disclosure without originator authorization | Prohibition on disclosure except as permitted by the information originator.
1 | (_ActProductAcquisitionCode) Abstract | | The method that a product is obtained for use by the subject of the supply act (e.g. patient). Product examples are consumable or durable goods.
2 | LOAN | Loan | Temporary supply of a product without transfer of ownership for the product.
3 | RENT | Rent | Temporary supply of a product with financial compensation, without transfer of ownership for the product.
2 | TRANSFER | Transfer | Transfer of ownership for a product.
3 | SALE | Sale | Transfer of ownership for a product for financial compensation.
1 | (_ActSpecimenTransportCode) Abstract | | Transportation of a specimen.
2 | SREC | specimen received | Description:Specimen has been received by the participating organization/department.
2 | SSTOR | specimen in storage | Description:Specimen has been placed into storage at a participating location.
2 | STRAN | specimen in transit | Description:Specimen has been put in transit to a participating receiver.
1 | (_ActSpecimenTreatmentCode) Abstract | | Set of codes related to specimen treatments

2 | ACID | Acidification | The lowering of specimen pH through the addition of an acid
2 | ALK | Alkalization | The act rendering alkaline by impregnating with an alkali; a conferring of alkaline qualities.
2 | DEFB | Defibrination | The removal of fibrin from whole blood or plasma through physical or chemical means
2 | FILT | Filtration | The passage of a liquid through a filter, accomplished by gravity, pressure or vacuum (suction).
2 | LDLP | LDL Precipitation
2 | NEUT | Neutralization | The act or process by which an acid and a base are combined in such proportions that the resulting compound is neutral.
2 | RECA | Recalcification | The addition of calcium back to a specimen after it was removed by chelating agents
2 | UFIL | Ultrafiltration | The filtration of a colloidal substance through a semipermeable medium that allows only the passage of small molecules.
1 | (_ActSubstanceAdministrationCode) Abstract | | Description: Describes the type of substance administration being performed. This should not be used to carry codes for identification of products. Use an associated role or entity to carry such information.
2 | DRUG | Drug therapy | The introduction of a drug into a subject with the intention of altering its biologic state with the intent of improving its health status.
2 | FD | food | Description: The introduction of material into a subject with the intent of providing nutrition or other dietary supplements (e.g. minerals or vitamins).
2 | IMMUNIZ | Immunization | The introduction of an immunogen with the intent of stimulating an immune response, aimed at preventing subsequent infections by more viable agents.
3 | BOOSTER | Booster Immunization | An additional immunization administration within a series intended to bolster or enhance immunity.
3 | INITIMMUNIZ | Initial Immunization | The first immunization administration in a series intended to produce immunity
1 | (_ActTaskCode) Abstract | | Description: A task or action that a user may perform in a clinical information system (e.g., medication order entry, laboratory test results review, problem list entry).
2 | OE | order entry task | A clinician creates a request for a service to be performed for a given patient.
3 | LABOE | laboratory test order entry task | A clinician creates a request for a laboratory test to be done for a given patient.
3 | MEDOE | medication order entry task | A clinician creates a request for the administration of one or more medications to a given patient.
2 | PATDOC | patient documentation task | A person enters documentation about a given patient.
3 | ALLERLREV | allergy list review | Description: A person reviews a list of known allergies of a given patient.
3 | CLINNOTEE | clinical note entry task | A clinician enters a clinical note about a given patient
4 | DIAGLISTE | diagnosis list entry task | A clinician enters a diagnosis for a given patient.
4 | DISCHINSTE | discharge instruction entry | A person provides a discharge instruction to a patient.
4 | DISCHSUME | discharge summary entry task | A clinician enters a discharge summary for a given patient.
4 | PATEDUE | patient education entry | A person provides a patient-specific education handout to a patient.
4 | PATREPE | pathology report entry task | A pathologist enters a report for a given patient.
4 | PROBLISTE | problem list entry task | A clinician enters a problem for a given patient.
4 | RADREPE | radiology report entry task | A radiologist enters a report for a given patient.
3 | IMMLREV | immunization list review | Description: A person reviews a list of immunizations due or received for a given patient.
3 | REMLREV | reminder list review | Description: A person reviews a list of health care reminders for a given patient.
4 | WELLREMLREV | wellness reminder list review | Description: A person reviews a list of wellness or preventive care reminders for a given patient.
2 | PATINFO | patient information review task | A person (e.g., clinician, the patient herself) reviews patient information in the electronic medical record.
3 | ALLERLE | allergy list entry | Description: A person enters a known allergy for a given patient.
3 | CDSREV | clinical decision support intervention review | A person reviews a recommendation/assessment provided automatically by a clinical decision support application for a given patient.
3 | CLINNOTEREV | clinical note review task | A person reviews a clinical note of a given patient.
4 | DISCHSUMREV | discharge summary review task | A person reviews a discharge summary of a given patient.
3 | DIAGLISTREV | diagnosis list review task | A person reviews a list of diagnoses of a given patient.
3 | IMMLE | immunization list entry | Description: A person enters an immunization due or received for a given patient.
3 | LABRREV | laboratory results review task | A person reviews a list of laboratory results of a given patient.
3 | MICRORREV | microbiology results review task | A person reviews a list of microbiology results of a given patient.
4 | MICROORGRREV | microbiology organisms results review task | A person reviews organisms of microbiology results of a given patient.
4 | MICROSENSRREV | microbiology sensitivity test results review task | A person reviews the sensitivity test of microbiology results of a given patient.
3 | MLREV | medication list review task | A person reviews a list of medication orders submitted to a given patient
4 | MARWLREV | medication administration record work list review task | A clinician reviews a work list of medications to be administered to a given patient.
3 | OREV | orders review task | A person reviews a list of orders submitted to a given patient.
3 | PATREPREV | pathology report review task | A person reviews a pathology report of a given patient.
3 | PROBLISTREV | problem list review task | A person reviews a list of problems of a given patient.
3 | RADREPREV | radiology report review task | A person reviews a radiology report of a given patient.
3 | REMLE | reminder list entry | Description: A person enters a health care reminder for a given patient.
4 | WELLREMLE | wellness reminder list entry | Description: A person enters a wellness or preventive care reminder for a given patient.
3 | RISKASSESS | risk assessment instrument task | A person reviews a Risk Assessment Instrument report of a given patient.
4 | FALLRISK | falls risk assessment instrument task | A person reviews a Falls Risk Assessment Instrument report of a given patient.
1 | (_ActTransportationModeCode) Abstract | | Characterizes how a transportation act was or will be carried out. Examples: Via private transport, via public transit, via courier.
2 | (_ActPatientTransportationModeCode) Abstract | | Definition: Characterizes how a patient was or will be transported to the site of a patient encounter. Examples: Via ambulance, via public transit, on foot.
3 | AFOOT | pedestrian transport

3 | AMBT | ambulance transport
4 | AMBAIR | fixed-wing ambulance transport
4 | AMBGRND | ground ambulance transport
4 | AMBHELO | helicopter ambulance transport
3 | LAWENF | law enforcement transport
3 | PRVTRN | private transport
3 | PUBTRN | public transport
1 | (_ObservationType) Abstract | | Identifies the kinds of observations that can be performed
2 | (_ActSpecObsCode) Abstract | | Identifies the type of observation that is made about a specimen that may affect its processing, analysis or further result interpretation
3 | ARTBLD | ActSpecObsArtBldCode | Describes the artificial blood identifier that is associated with the specimen.
3 | DILUTION | ActSpecObsDilutionCode | An observation that reports the dilution of a sample.
4 | AUTO-HIGH | Auto-High Dilution | The dilution of a sample performed by automated equipment. The value is specified by the equipment
4 | AUTO-LOW | Auto-Low Dilution | The dilution of a sample performed by automated equipment. The value is specified by the equipment
4 | PRE | Pre-Dilution | The dilution of the specimen made prior to being loaded onto analytical equipment
4 | RERUN | Rerun Dilution | The value of the dilution of a sample after it had been analyzed at a prior dilution value
3 | EVNFCTS | ActSpecObsEvntfctsCode | Domain provides codes that qualify the ActLabObsEnvfctsCode domain. (Environmental Factors)
3 | INTFR | ActSpecObsInterferenceCode | An observation that relates to factors that may potentially cause interference with the observation
4 | FIBRIN | Fibrin | The Fibrin Index of the specimen. In the case of only differentiating between Absent and Present, recommend using 0 and 1
4 | HEMOLYSIS | Hemolysis | An observation of the hemolysis index of the specimen in g/L
4 | ICTERUS | Icterus | An observation that describes the icterus index of the specimen. It is recommended to use mMol/L of bilirubin
4 | LIPEMIA | Lipemia | An observation used to describe the Lipemia Index of the specimen. It is recommended to use the optical turbidity at 600 nm (in absorbance units).
3 | VOLUME | ActSpecObsVolumeCode | An observation that reports the volume of a sample.
4 | AVAILABLE | Available Volume | The available quantity of specimen. This is the current quantity minus any planned consumption (e.g., tests that are planned)
4 | CONSUMPTION | Consumption Volume | The quantity of specimen that is used each time the equipment uses this substance
4 | CURRENT | Current Volume | The current quantity of the specimen, i.e., initial quantity minus what has been actually used.
4 | INITIAL | Initial Volume | The initial quantity of the specimen in inventory
2 | (_AnnotationType) Abstract
3 | _ActPatientAnnotationType | ActPatientAnnotationType | Description:Provides a categorization for annotations recorded directly against the patient .
4 | ANNDI | diagnostic image note | Description:A note that is specific to a patient's diagnostic images, either historical, current or planned.
4 | ANNGEN | general note | Description:A general or uncategorized note.
4 | ANNIMM | immunization note | A note that is specific to a patient's immunizations, either historical, current or planned.
4 | ANNLAB | laboratory note | Description:A note that is specific to a patient's laboratory results, either historical, current or planned.
4 | ANNMED | medication note | Description:A note that is specific to a patient's medications, either historical, current or planned.
2 | (_GeneticObservationType) Abstract | | Description: None provided
3 | GENE | gene | Description: A DNA segment that contributes to phenotype/function. In the absence of demonstrated function a gene may be characterized by sequence, transcription or homology
2 | _ImmunizationObservationType | ImmunizationObservationType | Description: Observation codes which describe characteristics of the immunization material.
3 | OBSANTC | antigen count | Description: Indicates the valid antigen count.
3 | OBSANTV | antigen validity | Description: Indicates whether an antigen is valid or invalid.
2 | (_IndividualCaseSafetyReportType) Abstract | | A code that is used to indicate the type of case safety report received from sender. The current code example reference is from the International Conference on Harmonisation (ICH) Expert Workgroup guideline on Clinical Safety Data Management: Data Elements for Transmission of Individual Case Safety Reports. The unknown/unavailable option allows the transmission of information from a secondary sender where the initial sender did not specify the type of report. Example concepts include: Spontaneous, Report from study, Other.
3 | PAT_ADV_EVNT | patient adverse event | Indicates that the ICSR is describing problems that a patient experienced after receiving a vaccine product.
3 | VAC_PROBLEM | vaccine product problem | Indicates that the ICSR is describing a problem with the actual vaccine product such as physical defects (cloudy, particulate matter) or inability to confer immunity.
2 | (_LOINCObservationActContextAgeType) Abstract | | Definition:The set of LOINC codes for the act of determining the period of time that has elapsed since an entity was born or created.
3 | 21611-9 | age patient qn est | Definition:Estimated age.
3 | 21612-7 | age patient qn reported | Definition:Reported age.
3 | 29553-5 | age patient qn calc | Definition:Calculated age.
3 | 30525-0 | age patient qn definition | Definition:General specification of age with no implied method of determination.
3 | 30972-4 | age at onset of adverse event | Definition:Age at onset of associated adverse event; no implied method of determination.
2 | (_MedicationObservationType) Abstract
3 | REP_HALF_LIFE | representative half-life | Description:This observation represents an 'average' or 'expected' half-life typical of the product.
3 | SPLCOATING | coating | Definition: A characteristic of an oral solid dosage form of a medicinal product, indicating whether it has one or more coatings such as sugar coating, film coating, or enteric coating. Only coatings to the external surface or the dosage form should be considered (for example, coatings to individual pellets or granules inside a capsule or tablet are excluded from

consideration). Constraints: The Observation.value must be a Boolean (BL) with true for the presence or false for the absence of one or more coatings on a solid dosage form.

3 | SPLCOLOR | color | Definition: A characteristic of an oral solid dosage form of a medicinal product, specifying the color or colors that most predominantly define the appearance of the dose form. SPLCOLOR is not an FDA specification for the actual color of solid dosage forms or the names of colors that can appear in labeling. Constraints: The Observation.value must be a single coded value or a list of multiple coded values, specifying one or more distinct colors that approximate of the color(s) of distinct areas of the solid dosage form, such as the different sides of a tablet or one-part capsule, or the different halves of a two-part capsule. Bands on banded capsules, regardless of the color, are not considered when assigning an SPLCOLOR. Imprints on the dosage form, regardless of their color are not considered when assigning an SPLCOLOR. If more than one color exists on a particular side or half, then the most predominant color on that side or half is recorded. If the gelatin capsule shell is colorless and transparent, use the predominant color of the contents that appears through the colorless and transparent capsule shell. Colors can include: Black;Gray;White;Red;Pink;Purple;Green;Yellow;Orange;Brown;Blue;Turquoise.

3 | SPLIMAGE | image | Description: A characteristic representing a single file reference that contains two or more views of the same dosage form of the product; in most cases this should represent front and back views of the dosage form, but occasionally additional views might be needed in order to capture all of the important physical characteristics of the dosage form. Any imprint and/or symbol should be clearly identifiable, and the viewer should not normally need to rotate the image in order to read it. Images that are submitted with SPL should be included in the same directory as the SPL file.

3 | SPLIMPRINT | imprint | Definition: A characteristic of an oral solid dosage form of a medicinal product, specifying the alphanumeric text that appears on the solid dosage form, including text that is embossed, debossed, engraved or printed with ink. The presence of other non-textual distinguishing marks or symbols is recorded by SPLSYMBOL. Examples: Included in SPLIMPRINT are alphanumeric text that appears on the bands of banded capsules and logos and other symbols that can be interpreted as letters or numbers. Constraints: The Observation.value must be of type Character String (ST). Excluded from SPLIMPRINT are internal and external cut-outs in the form of alphanumeric text and the letter 'R' with a circle around it (when referring to a registered trademark) and the letters 'TM' (when referring to a 'trade mark'). To record text, begin on either side or part of the dosage form. Start at the top left and progress as one would normally read a book. Enter a semicolon to show separation between words or line divisions.

3 | SPLSCORING | scoring | Definition: A characteristic of an oral solid dosage form of a medicinal product, specifying the number of equal pieces that the solid dosage form can be divided into using score line(s). Example: One score line creating two equal pieces is given a value of 2, two parallel score lines creating three equal pieces is given a value of 3. Constraints: Whether three parallel score lines create four equal pieces or two intersecting score lines create two equal pieces using one score line and four equal pieces using both score lines, both have the scoring value of 4. Solid dosage forms that are not scored are given a value of 1. Solid dosage forms that can only be divided into unequal pieces are given a null-value with nullFlavor other (OTH).

3 | SPLSHAPE | shape | Description: A characteristic of an oral solid dosage form of a medicinal product, specifying the two dimensional representation of the solid dose form, in terms of the outside perimeter of a solid dosage form when the dosage form, resting on a flat surface, is viewed from directly above, including slight rounding of corners. SPLSHAPE does not include embossing, scoring, debossing, or internal cut-outs. SPLSHAPE is independent of the orientation of the imprint and logo. Shapes can include: Triangle (3 sided); Square; Round; Semicircle; Pentagon (5 sided); Diamond; Double circle; Bullet; Hexagon (6 sided); Rectangle; Gear; Capsule; Heptagon (7 sided); Trapezoid; Oval; Clover; Octagon (8 sided); Tear; Freeform.

3 | SPLSIZE | size | Definition: A characteristic of an oral solid dosage form of a medicinal product, specifying the longest single dimension of the solid dosage form as a physical quantity in the dimension of length (e.g., 3 mm). The length is should be specified in millimeters and should be rounded to the nearest whole millimeter. Example: SPLSIZE for a rectangular shaped tablet is the length and SPLSIZE for a round shaped tablet is the diameter.

3 | SPLSYMBOL | symbol | Definition: A characteristic of an oral solid dosage form of a medicinal product, to describe whether or not the medicinal product has a mark or symbol appearing on it for easy and definite recognition. Score lines, letters, numbers, and internal and external cut-outs are not considered marks or symbols. See SPLSCORING and SPLIMPRINT for these characteristics. Constraints: The Observation.value must be a Boolean (BL) with <u>true</u> indicating the presence and <u>false</u> for the absence of marks or symbols. Example:

2 | (_ObservationIssueTriggerCodedObservationType) Abstract | | Distinguishes the kinds of coded observations that could be the trigger for clinical issue detection. These are observations that are not measurable, but instead can be defined with codes. Coded observation types include: Allergy, Intolerance, Medical Condition, Pregnancy status, etc.

3 | (_CaseTransmissionMode) Abstract | | Code for the mechanism by which disease was acquired by the living subject involved in the public health case. Includes sexually transmitted, airborne, bloodborne, vectorborne, foodborne, zoonotic, nosocomial, mechanical, dermal, congenital, environmental exposure, indeterminate.

4 | AIRTRNS | airborne transmission | Communication of an agent from a living subject or environmental source to a living subject through indirect contact via oral or nasal inhalation.

4 | ANANTRNS | animal to animal transmission | Communication of an agent from one animal to another proximate animal.

4 | ANHUMTRNS | animal to human transmission | Communication of an agent from an animal to a proximate person.

4 | BDYFLDTRNS | body fluid contact transmission | Communication of an agent from one living subject to another living subject through direct contact with any body fluid.

4 | BLDTRNS | blood borne transmission | Communication of an agent to a living subject through direct contact with blood or blood products whether the contact with blood is part of a therapeutic procedure or not.

4 | DERMTRNS | transdermal transmission | Communication of an agent from a living subject or environmental source to a living subject via agent migration through intact skin.

4 | ENVTRNS | environmental exposure transmission | Communication of an agent from an environmental surface or source to a living subject by direct contact.

4 | FECTRNS | fecal-oral transmission | Communication of an agent from a living subject or environmental source to a living subject through oral contact with material contaminated by person or animal fecal material.

4 | FOMTRNS | fomite transmission | Communication of an agent from an non-living material to a living subject through direct contact.

4 | FOODTRNS | food-borne transmission | Communication of an agent from a food source to a living subject via oral consumption.

4 | HUMHUMTRNS | human to human transmission | Communication of an agent from a person to a proximate person.

4 | INDTRNS | indeterminate disease transmission mode | Communication of an agent to a living subject via an undetermined route.

4 | LACTTRNS | lactation transmission | Communication of an agent from one living subject to another living subject through direct contact with mammalian milk or colostrum.

4 | NOSTRNS | nosocomial transmission | Communication of an agent from any entity to a living subject while the living subject is in the patient role in a healthcare facility.

4 | PARTRNS | parenteral transmission | Communication of an agent from a living subject or environmental source to a living subject where the acquisition of the agent is not via the alimentary canal.
4 | PLACTRNS | transplacental transmission | Communication of an agent from a living subject to the progeny of that living subject via agent migration across the maternal-fetal placental membranes while in utero.
4 | SEXTRNS | sexual transmission | Communication of an agent from one living subject to another living subject through direct contact with genital or oral tissues as part of a sexual act.
4 | TRNSFTRNS | transfusion transmission | Communication of an agent from one living subject to another living subject through direct contact with blood or blood products where the contact with blood is part of a therapeutic procedure.
4 | VECTRNS | vector-borne transmission | Communication of an agent from a living subject acting as a required intermediary in the agent transmission process to a recipient living subject via direct contact.
4 | WATTRNS | water-borne transmission | Communication of an agent from a contaminated water source to a living subject whether the water is ingested as a food or not. The route of entry of the water may be through any bodily orifice.
2 | _ObservationQualityMeasureAttribute | ObservationQualityMeasureAttribute | Codes used to define various metadata aspects of a health quality measure.
3 | AGGREGATE | aggregate measure observation | Indicates that the observation is carrying out an aggregation calculation, contained in the value element.
3 | CMPMSRMTH | composite measure method | Indicates what method is used in a quality measure to combine the component measure results included in an composite measure.
3 | CMPMSRSCRWGHT | component measure scoring weight | An attribute of a quality measure describing the weight this component measure score is to carry in determining the overall composite measure final score. The value is real value greater than 0 and less than 1.0. Each component measure score will be multiplied by its CMPMSRSCRWGHT and then summed with the other component measures to determine the final overall composite measure score. The sum across all CMPMSRSCRWGHT values within a single composite measure SHALL be 1.0. The value assigned is scoped to the composite measure referencing this component measure only.
3 | COPY | copyright | Identifies the organization(s) who own the intellectual property represented by the eMeasure.
3 | CRS | clinical recommendation statement | Summary of relevant clinical guidelines or other clinical recommendations supporting this eMeasure.
3 | DEF | definition | Description of individual terms, provided as needed.
3 | DISC | disclaimer | Disclaimer information for the eMeasure.
3 | FINALDT | finalized date/time | The timestamp when the eMeasure was last packaged in the Measure Authoring Tool.
3 | GUIDE | guidance | Used to allow measure developers to provide additional guidance for implementers to understand greater specificity than could be provided in the logic for data criteria.
3 | IDUR | improvement notation | Information on whether an increase or decrease in score is the preferred result (e.g., a higher score indicates better quality OR a lower score indicates better quality OR quality is within a range).
3 | ITMCNT | items counted | Describes the items counted by the measure (e.g., patients, encounters, procedures, etc.)
3 | KEY | keyword | A significant word that aids in discoverability.
3 | MEDT | measurement end date | The end date of the measurement period.
3 | MSD | measurement start date | The start date of the measurement period.
3 | MSRADJ | risk adjustment | The method of adjusting for clinical severity and conditions present at the start of care that can influence patient outcomes for making valid comparisons of outcome measures across providers. Indicates whether an eMeasure is subject to the statistical process for reducing, removing, or clarifying the influences of confounding factors to allow more useful comparisons.
3 | MSRAGG | rate aggregation | Describes how to combine information calculated based on logic in each of several populations into one summarized result. It can also be used to describe how to risk adjust the data based on supplemental data elements described in the eMeasure. (e.g., pneumonia hospital measures antibiotic selection in the ICU versus non-ICU and then the roll-up of the two). Open Issue: The description does NOT align well with the definition used in the HQMF specfication; correct the MSGAGG definition, and the possible distinction of MSRAGG as a child of AGGREGATE.
3 | MSRIMPROV | health quality measure improvement notation | Information on whether an increase or decrease in score is the preferred result. This should reflect information on which way is better, an increase or decrease in score.
3 | MSRJUR | jurisdiction | The list of jurisdiction(s) for which the measure applies.
3 | MSRRPTR | reporter type | Type of person or organization that is expected to report the issue.
3 | MSRRPTTIME | timeframe for reporting | The maximum time that may elapse following completion of the measure until the measure report must be sent to the receiver.
3 | MSRSCORE | measure scoring | Indicates how the calculation is performed for the eMeasure (e.g., proportion, continuous variable, ratio)
3 | MSRSET | health quality measure care setting | Location(s) in which care being measured is rendered Usage Note: MSRSET is used rather than RoleCode because the setting applies to what is being measured, as opposed to participating directly in the health quality measure documantion itself).
3 | MSRTOPIC | health quality measure topic type
3 | MSRTP | measurement period | The time period for which the eMeasure applies.
3 | MSRTYPE | measure type | Indicates whether the eMeasure is used to examine a process or an outcome over time (e.g., Structure, Process, Outcome).
3 | RAT | rationale | Succinct statement of the need for the measure. Usually includes statements pertaining to Importance criterion: impact, gap in care and evidence.
3 | REF | reference | Identifies bibliographic citations or references to clinical practice guidelines, sources of evidence, or other relevant materials supporting the intent and rationale of the eMeasure.
3 | SDE | supplemental data elements | Comparison of results across strata can be used to show where disparities exist or where there is a need to expose differences in results. For example, Centers for Medicare & Medicaid Services (CMS) in the U.S. defines four required Supplemental Data Elements (payer, ethnicity, race, and gender), which are variables used to aggregate data into various subgroups. Additional supplemental data elements required for risk adjustment or other purposes of data aggregation can be included in the Supplemental Data Element section.
3 | STRAT | stratification | Describes the strata for which the measure is to be evaluated. There are three examples of reasons for stratification based on existing work. These include: (1) evaluate the measure based on different age groupings within the population described in the measure (e.g., evaluate the whole [age 14-25] and each sub-stratum [14-19] and [20-25]); (2) evaluate the eMeasure based on either a specific condition, a specific discharge location, or both; (3) evaluate the eMeasure based on different locations within a facility (e.g., evaluate the overall rate for all intensive care units and also some strata include

additional findings [specific birth weights for neonatal intensive care units]).
3 | TRANF | transmission format | Can be a URL or hyperlinks that link to the transmission formats that are specified for a particular reporting program.
3 | USE | notice of use | Usage notes.
2 | (_ObservationSequenceType) Abstract
3 | TIME_ABSOLUTE | absolute time sequence | A sequence of values in the "absolute" time domain. This is the same time domain that all HL7 timestamps use. It is time as measured by the Gregorian calendar
3 | TIME_RELATIVE | relative time sequence | A sequence of values in a "relative" time domain. The time is measured relative to the earliest effective time in the Observation Series containing this sequence.
2 | (_ObservationSeriesType) Abstract
3 | (_ECGObservationSeriesType) Abstract
4 | REPRESENTATIVE_BEAT | ECG representative beat waveforms | This Observation Series type contains waveforms of a "representative beat" (a.k.a. "median beat" or "average beat"). The waveform samples are measured in relative time, relative to the beginning of the beat as defined by the Observation Series effective time. The waveforms are not directly acquired from the subject, but rather algorithmically derived from the "rhythm" waveforms.
4 | RHYTHM | ECG rhythm waveforms | This Observation type contains ECG "rhythm" waveforms. The waveform samples are measured in absolute time (a.k.a. "subject time" or "effective time"). These waveforms are usually "raw" with some minimal amount of noise reduction and baseline filtering applied.
2 | _PatientImmunizationRelatedObservationType | PatientImmunizationRelatedObservationType | Description: Reporting codes that are related to an immunization event.
3 | CLSSRM | classroom | Description: The class room associated with the patient during the immunization event.
3 | GRADE | grade | Description: The school grade or level the patient was in when immunized.
3 | SCHL | school | Description: The school the patient attended when immunized.
3 | SCHLDIV | school division | Description: The school division or district associated with the patient during the immunization event.
3 | TEACHER | teacher | Description: The patient's teacher when immunized.
2 | (_PopulationInclusionObservationType) Abstract | | Observation types for specifying criteria used to assert that a subject is included in a particular population.
3 | DENEX | denominator exclusions | Criteria which specify subjects who should be removed from the eMeasure population and denominator before determining if numerator criteria are met. Denominator exclusions are used in proportion and ratio measures to help narrow the denominator.
3 | DENEXCEP | denominator exceptions | Criteria which specify the removal of a subject, procedure or unit of measurement from the denominator, only if the numerator criteria are not met. Denominator exceptions allow for adjustment of the calculated score for those providers with higher risk populations. Denominator exceptions are used only in proportion eMeasures. They are not appropriate for ratio or continuous variable eMeasures. Denominator exceptions allow for the exercise of clinical judgment and should be specifically defined where capturing the information in a structured manner fits the clinical workflow. Generic denominator exception reasons used in proportion eMeasures fall into three general categories: Medical reasons Patient (or subject) reasons System reasons
3 | DENOM | denominator | Criteria for specifying the entities to be evaluated by a specific quality measure, based on a shared common set of characteristics (within a specific measurement set to which a given measure belongs). The denominator can be the same as the initial population, or it may be a subset of the initial population to further constrain it for the purpose of the eMeasure. Different measures within an eMeasure set may have different denominators. Continuous Variable eMeasures do not have a denominator, but instead define a measure population.
3 | IPOP | initial population | Criteria for specifying the entities to be evaluated by a specific quality measure, based on a shared common set of characteristics (within a specific measurement set to which a given measure belongs).
4 | IPPOP | initial patient population | Criteria for specifying the patients to be evaluated by a specific quality measure, based on a shared common set of characteristics (within a specific measurement set to which a given measure belongs). Details often include information based upon specific age groups, diagnoses, diagnostic and procedure codes, and enrollment periods.
3 | MSROBS | measure observation | Defines the observation to be performed for each patient or event in the measure population. Measure observations for each case in the population are aggregated to determine the overall measure score for the population. Examples: the median time from arrival in the Emergency Room to departure the median time from decision to admit to a hospital to the actual admission for Emergency Room patients
3 | MSRPOPL | measure population | Criteria for specifying the measure population as a narrative description (e.g., all patients seen in the Emergency Department during the measurement period). This is used only in continuous variable eMeasures.
3 | MSRPOPLEX | measure population exclusions | Criteria for specifying subjects who should be removed from the eMeasure's Initial Population and Measure Population. Measure Population Exclusions are used in Continuous Variable measures to help narrow the Measure Population before determining the value(s) of the continuous variable(s).
3 | NUMER | numerator | Criteria for specifying the processes or outcomes expected for each patient, procedure, or other unit of measurement defined in the denominator for proportion measures, or related to (but not directly derived from) the denominator for ratio measures (e.g., a numerator listing the number of central line blood stream infections and a denominator indicating the days per thousand of central line usage in a specific time period).
3 | NUMEX | numerator exclusions | Criteria for specifying instances that should not be included in the numerator data. (e.g., if the number of central line blood stream infections per 1000 catheter days were to exclude infections with a specific bacterium, that bacterium would be listed as a numerator exclusion). Numerator Exclusions are used only in ratio eMeasures.
2 | (_PreferenceObservationType) Abstract | | Types of observations that can be made about Preferences.
3 | PREFSTRENGTH | preference strength | An observation about how important a preference is to the target of the preference.
2 | ADVERSE_REACTION | Adverse Reaction | Indicates that the observation is of an unexpected negative occurrence in the subject suspected to result from the subject's exposure to one or more agents. Observation values would be the symptom resulting from the reaction.
2 | ASSERTION | Assertion | Description:Refines classCode OBS to indicate an observation in which observation.value contains a finding or other nominalized statement, where the encoded information in Observation.value is not altered by Observation.code. For instance, observation.code="ASSERTION" and observation.value="fracture of femur present" is an assertion of a clinical finding of femur fracture.
2 | CASESER | case seriousness criteria | Definition:An observation that provides a characterization of the level of harm to an investigation subject as a result of a reaction or event.

2 | CDIO | case disease imported observation | An observation that states whether the disease was likely acquired outside the jurisdiction of observation, and if so, the nature of the inter-jurisdictional relationship. OpenIssue: This code could be moved to LOINC if it can be done before there are significant implemenations using it.
2 | CRIT | criticality | A clinical judgment as to the worst case result of a future exposure (including substance administration). When the worst case result is assessed to have a life-threatening or organ system threatening potential, it is considered to be of high criticality.
2 | CTMO | case transmission mode observation | An observation that states the mechanism by which disease was acquired by the living subject involved in the public health case. OpenIssue: This code could be moved to LOINC if it can be done before there are significant implemenations using it.
2 | DX | ObservationDiagnosisTypes | Includes all codes defining types of indications such as diagnosis, symptom and other indications such as contrast agents for lab tests.
3 | ADMDX | admitting diagnosis | Admitting diagnosis are the diagnoses documented for administrative purposes as the basis for a hospital admission.
3 | DISDX | discharge diagnosis | Discharge diagnosis are the diagnoses documented for administrative purposes as the time of hospital discharge.
3 | INTDX | intermediate diagnosis | Intermediate diagnoses are those diagnoses documented for administrative purposes during the course of a hospital stay.
3 | NOI | nature of injury | The type of injury that the injury coding specifies.
2 | GISTIER | GIS tier | Description: Accuracy determined as per the GIS tier code system.
2 | HHOBS | household situation observation | Indicates that the observation is of a person’s living situation in a household including the household composition and circumstances.
2 | ISSUE | detected issue | There is a clinical issue for the therapy that makes continuation of the therapy inappropriate. Open Issue: The definition of this code does not correctly represent the concept space of its specializations (children)
3 | (_ActAdministrativeDetectedIssueCode) Abstract | | Identifies types of detectyed issues for Act class "ALRT" for the administrative and patient administrative acts domains.
4 | (_ActAdministrativeAuthorizationDetectedIssueCode) Abstract
5 | NAT | Insufficient authorization | The requesting party has insufficient authorization to invoke the interaction.
5 | SUPPRESSED | record suppressed | Description: One or more records in the query response have been suppressed due to consent or privacy restrictions.
5 | VALIDAT | validation issue | Description:The specified element did not pass business-rule validation.
6 | KEY204 | Unknown key identifier | The ID of the patient, order, etc., was not found. Used for transactions other than additions, e.g. transfer of a non-existent patient.
6 | KEY205 | Duplicate key identifier | The ID of the patient, order, etc., already exists. Used in response to addition transactions (Admit, New Order, etc.).
6 | COMPLY | Compliance Alert | There may be an issue with the patient complying with the intentions of the proposed therapy
7 | DUPTHPY | Duplicate Therapy Alert | The proposed therapy appears to duplicate an existing therapy
8 | DUPTHPCLS | duplicate therapeutic alass alert | Description:The proposed therapy appears to have the same intended therapeutic benefit as an existing therapy, though the specific mechanisms of action vary.
8 | DUPTHPGEN | duplicate generic alert | Description:The proposed therapy appears to have the same intended therapeutic benefit as an existing therapy and uses the same mechanisms of action as the existing therapy.
7 | ABUSE | commonly abused/misused alert | Description:The proposed therapy is frequently misused or abused and therefore should be used with caution and/or monitoring.
7 | FRAUD | potential fraud | Description:The request is suspected to have a fraudulent basis.
7 | PLYDOC | Poly-orderer Alert | A similar or identical therapy was recently ordered by a different practitioner.
7 | PLYPHRM | Poly-supplier Alert | This patient was recently supplied a similar or identical therapy from a different pharmacy or supplier.
6 | DOSE | Dosage problem | Proposed dosage instructions for therapy differ from standard practice.
7 | DOSECOND | dosage-condition alert | Description:Proposed dosage is inappropriate due to patient's medical condition.
7 | DOSEDUR | Dose-Duration Alert | Proposed length of therapy differs from standard practice.
8 | DOSEDURH | Dose-Duration High Alert | Proposed length of therapy is longer than standard practice
9 | DOSEDURHIND | Dose-Duration High for Indication Alert | Proposed length of therapy is longer than standard practice for the identified indication or diagnosis
8 | DOSEDURL | Dose-Duration Low Alert | Proposed length of therapy is shorter than that necessary for therapeutic effect
9 | DOSEDURLIND | Dose-Duration Low for Indication Alert | Proposed length of therapy is shorter than standard practice for the identified indication or diagnosis
7 | DOSEH | High Dose Alert | Proposed dosage exceeds standard practice
8 | DOSEHINDA | High Dose for Age Alert | Proposed dosage exceeds standard practice for the patient's age
8 | DOSEHIND | High Dose for Indication Alert
8 | DOSEHINDSA | High Dose for Height/Surface Area Alert | Proposed dosage exceeds standard practice for the patient's height or body surface area
8 | DOSEHINDW | High Dose for Weight Alert | Proposed dosage exceeds standard practice for the patient's weight
7 | DOSEIVL | Dose-Interval Alert | Proposed dosage interval/timing differs from standard practice
8 | DOSEIVLIND | Dose-Interval for Indication Alert | Proposed dosage interval/timing differs from standard practice for the identified indication or diagnosis
7 | DOSEL | Low Dose Alert | Proposed dosage is below suggested therapeutic levels
8 | DOSELINDA | Low Dose for Age Alert | Proposed dosage is below suggested therapeutic levels for the patient's age
8 | DOSELIND | Low Dose for Indication Alert
8 | DOSELINDSA | Low Dose for Height/Surface Area Alert | Proposed dosage is below suggested therapeutic levels for the patient's height or body surface area
8 | DOSELINDW | Low Dose for Weight Alert | Proposed dosage is below suggested therapeutic levels for the patient's weight
7 | MDOSE | maximum dosage reached | Description:The maximum quantity of this drug allowed to be administered within a particular time-range (month, year, lifetime) has been reached or exceeded.
6 | OBSA | Observation Alert | Proposed therapy may be inappropriate or contraindicated due to conditions or characteristics of the patient
7 | AGE | Age Alert | Proposed therapy may be inappropriate or contraindicated due to patient age

8 | ADALRT | adult alert | Proposed therapy is outside of the standard practice for an adult patient.
8 | DOSEHINDA
8 | DOSELINDA
8 | GEALRT | geriatric alert | Proposed therapy is outside of standard practice for a geriatric patient.
8 | PEALRT | pediatric alert | Proposed therapy is outside of the standard practice for a pediatric patient.
7 | COND | Condition Alert | Proposed therapy may be inappropriate or contraindicated due to an existing/recent patient condition or diagnosis
8 | HGHT
8 | LACT | Lactation Alert | Proposed therapy may be inappropriate or contraindicated when breast-feeding
8 | PREG | Pregnancy Alert | Proposed therapy may be inappropriate or contraindicated during pregnancy
8 | WGHT
7 | CREACT | common reaction alert | Description:Proposed therapy may be inappropriate or contraindicated because of a common but non-patient specific reaction to the product. Example:There is no record of a specific sensitivity for the patient, but the presence of the sensitivity is common and therefore caution is warranted.
7 | GEN | Genetic Alert | Proposed therapy may be inappropriate or contraindicated due to patient genetic indicators.
7 | GEND | Gender Alert | Proposed therapy may be inappropriate or contraindicated due to patient gender.
7 | LAB | Lab Alert | Proposed therapy may be inappropriate or contraindicated due to recent lab test results
7 | REACT | Reaction Alert | Proposed therapy may be inappropriate or contraindicated based on the potential for a patient reaction to the proposed product
8 | ALGY | Allergy Alert | Proposed therapy may be inappropriate or contraindicated because of a recorded patient allergy to the proposed product. (Allergies are immune based reactions.)
8 | INT | Intolerance Alert | Proposed therapy may be inappropriate or contraindicated because of a recorded patient intolerance to the proposed product. (Intolerances are non-immune based sensitivities.)
7 | RREACT | Related Reaction Alert | Proposed therapy may be inappropriate or contraindicated because of a potential patient reaction to a cross-sensitivity related product.
8 | RALG | Related Allergy Alert | Proposed therapy may be inappropriate or contraindicated because of a recorded patient allergy to a cross-sensitivity related product. (Allergies are immune based reactions.)
8 | RAR | Related Prior Reaction Alert | Proposed therapy may be inappropriate or contraindicated because of a recorded prior adverse reaction to a cross-sensitivity related product.
8 | RINT | Related Intolerance Alert | Proposed therapy may be inappropriate or contraindicated because of a recorded patient intolerance to a cross-sensitivity related product. (Intolerances are non-immune based sensitivities.)
6 | BUS | business constraint violation | Description:A local business rule relating multiple elements has been violated.
6 | CODE_INVAL | code is not valid | Description:The specified code is not valid against the list of codes allowed for the element.
7 | CODE_DEPREC | code has been deprecated | Description:The specified code has been deprecated and should no longer be used. Select another code from the code system.
6 | FORMAT | invalid format | Description:The element does not follow the formatting or type rules defined for the field.
6 | ILLEGAL | illegal | Description:The request is missing elements or contains elements which cause it to not meet the legal standards for actioning.
6 | LEN_RANGE | length out of range | Description:The length of the data specified falls out of the range defined for the element.
7 | LEN_LONG | length is too long | Description:The length of the data specified is greater than the maximum length defined for the element.
7 | LEN_SHORT | length is too short | Description:The length of the data specified is less than the minimum length defined for the element.
6 | MISSCOND | conditional element missing | Description:The specified element must be specified with a non-null value under certain conditions. In this case, the conditions are true but the element is still missing or null.
6 | MISSMAND | mandatory element missing | Description:The specified element is mandatory and was not included in the instance.
6 | NODUPS | duplicate values are not permitted | Description:More than one element with the same value exists in the set. Duplicates not permission in this set in a set.
6 | NOPERSIST | element will not be persisted | Description: Element in submitted message will not persist in data storage based on detected issue.
6 | REP_RANGE | repetitions out of range | Description:The number of repeating elements falls outside the range of the allowed number of repetitions.
7 | MAXOCCURS | repetitions above maximum | Description:The number of repeating elements is above the maximum number of repetitions allowed.
7 | MINOCCURS | repetitions below minimum | Description:The number of repeating elements is below the minimum number of repetitions allowed.
4 | (_ActAdministrativeRuleDetectedIssueCode) Abstract
5 | KEY204
5 | KEY205
5 | KEY206 | non-matching identification | Description: Metadata associated with the identification (e.g. name or gender) does not match the identification being verified.
5 | OBSOLETE | obsolete record returned | Description: One or more records in the query response have a status of 'obsolete'.
3 | (_ActSuppliedItemDetectedIssueCode) Abstract | | Identifies types of detected issues regarding the administration or supply of an item to a patient.
4 | (_AdministrationDetectedIssueCode) Abstract | | Administration of the proposed therapy may be inappropriate or contraindicated as proposed
5 | (_AppropriatenessDetectedIssueCode) Abstract
6 | (_InteractionDetectedIssueCode) Abstract
7 | FOOD | Food Interaction Alert | Proposed therapy may interact with certain foods
7 | TPROD | Therapeutic Product Alert | Proposed therapy may interact with an existing or recent therapeutic product
8 | DRG | Drug Interaction Alert | Proposed therapy may interact with an existing or recent drug therapy
8 | NHP | Natural Health Product Alert | Proposed therapy may interact with existing or recent natural health product therapy
8 | NONRX | Non-Prescription Interaction Alert | Proposed therapy may interact with a non-prescription drug (e.g. alcohol, tobacco, Aspirin)

6 | OBSA
6 | PREVINEF | previously ineffective | Definition:The same or similar treatment has previously been attempted with the patient without achieving a positive effect.
5 | COMPLY
5 | DACT | drug action detected issue | Description:Proposed therapy may be contraindicated or ineffective based on an existing or recent drug therapy.
5 | DOSE
5 | DUPTHPY
5 | TIME | timing detected issue | Description:Proposed therapy may be inappropriate or ineffective based on the proposed start or end time.
6 | ALRTENDLATE | end too late alert | Definition:Proposed therapy may be inappropriate or ineffective because the end of administration is too close to another planned therapy.
6 | ALRTSTRTLATE | start too late alert | Definition:Proposed therapy may be inappropriate or ineffective because the start of administration is too late after the onset of the condition.
5 | (_TimingDetectedIssueCode) Abstract | | Proposed therapy may be inappropriate or ineffective based on the proposed start or end time.
6 | ENDLATE | End Too Late Alert | Proposed therapy may be inappropriate or ineffective because the end of administration is too close to another planned therapy
6 | STRTLATE | Start Too Late Alert | Proposed therapy may be inappropriate or ineffective because the start of administration is too late after the onset of the condition
4 | (_SupplyDetectedIssueCode) Abstract | | Supplying the product at this time may be inappropriate or indicate compliance issues with the associated therapy
5 | ALLDONE | already performed | Definition:The requested action has already been performed and so this request has no effect
5 | FULFIL | fulfillment alert | Definition:The therapy being performed is in some way out of alignment with the requested therapy.
6 | NOTACTN | no longer actionable | Definition:The status of the request being fulfilled has changed such that it is no longer actionable. This may be because the request has expired, has already been completely fulfilled or has been otherwise stopped or disabled. (Not used for 'suspended' orders.)
6 | NOTEQUIV | not equivalent alert | Definition:The therapy being performed is not sufficiently equivalent to the therapy which was requested.
7 | NOTEQUIVGEN | not generically equivalent alert | Definition:The therapy being performed is not generically equivalent (having the identical biological action) to the therapy which was requested.
7 | NOTEQUIVTHER | not therapeutically equivalent alert | Definition:The therapy being performed is not therapeutically equivalent (having the same overall patient effect) to the therapy which was requested.
6 | TIMING | event timing incorrect alert | Definition:The therapy is being performed at a time which diverges from the time the therapy was requested
7 | INTERVAL | outside requested time | Definition:The therapy action is being performed outside the bounds of the time period requested
7 | MINFREQ | too soon within frequency based on the usage | Definition:The therapy action is being performed too soon after the previous occurrence based on the requested frequency
5 | HELD | held/suspended alert | Definition:There should be no actions taken in fulfillment of a request that has been held or suspended.
5 | TOOLATE | Refill Too Late Alert | The patient is receiving a subsequent fill significantly later than would be expected based on the amount previously supplied and the therapy dosage instructions
5 | TOOSOON | Refill Too Soon Alert | The patient is receiving a subsequent fill significantly earlier than would be expected based on the amount previously supplied and the therapy dosage instructions
4 | HISTORIC | record recorded as historical | Description: While the record was accepted in the repository, there is a more recent version of a record of this type.
4 | PATPREF | violates stated preferences | Definition:The proposed therapy goes against preferences or consent constraints recorded in the patient's record.
5 | PATPREFALT | violates stated preferences, alternate available | Definition:The proposed therapy goes against preferences or consent constraints recorded in the patient's record. An alternate therapy meeting those constraints is available.
2 | KSUBJ | knowledge subject | Categorization of types of observation that capture the main clinical knowledge subject which may be a medication, a laboratory test, a disease.
2 | KSUBT | knowledge subtopic | Categorization of types of observation that capture a knowledge subtopic which might be treatment, etiology, or prognosis.
2 | OINT | intolerance | Hypersensitivity resulting in an adverse reaction upon exposure to an agent.
3 | ALG | Allergy | Hypersensitivity to an agent caused by an immunologic response to an initial exposure
4 | DALG | Drug Allergy | An allergy to a pharmaceutical product.
4 | EALG | Environmental Allergy | An allergy to a substance other than a drug or a food. E.g. Latex, pollen, etc.
4 | FALG | Food Allergy | An allergy to a substance generally consumed for nutritional purposes.
3 | DINT | Drug Intolerance | Hypersensitivity resulting in an adverse reaction upon exposure to a drug.
4 | DALG
4 | DNAINT | Drug Non-Allergy Intolerance | Hypersensitivity to an agent caused by a mechanism other than an immunologic response to an initial exposure
3 | EINT | Environmental Intolerance | Hypersensitivity resulting in an adverse reaction upon exposure to environmental conditions.
4 | EALG
4 | ENAINT | Environmental Non-Allergy Intolerance | Hypersensitivity to an agent caused by a mechanism other than an immunologic response to an initial exposure
3 | FINT | Food Intolerance | Hypersensitivity resulting in an adverse reaction upon exposure to food.
4 | FALG
4 | FNAINT | Food Non-Allergy Intolerance | Hypersensitivity to an agent caused by a mechanism other than an immunologic response to an initial exposure
3 | NAINT | Non-Allergy Intolerance | Hypersensitivity to an agent caused by a mechanism other than an immunologic response to an initial exposure

4 | DNAINT
4 | ENAINT
4 | FNAINT
2 | SEV | Severity Observation | A subjective evaluation of the seriousness or intensity associated with another observation.
2 | (_FDALabelData) Abstract | | FDA label data
3 | FDACOATING | coating | FDA label coating
3 | FDACOLOR | color | FDA label color
3 | FDAIMPRINTCD | imprint code | FDA label imprint code
3 | FDALOGO | logo | FDA label logo
3 | FDASCORING | scoring | FDA label scoring
3 | FDASHAPE | shape | FDA label shape
3 | FDASIZE | size | FDA label size
1 | (_ROIOverlayShape) Abstract | | Shape of the region on the object being referenced
2 | CIRCLE | circle | A circle defined by two (column,row) pairs. The first point is the center of the circle and the second point is a point on the perimeter of the circle.
2 | ELLIPSE | ellipse | An ellipse defined by four (column,row) pairs, the first two points specifying the endpoints of the major axis and the second two points specifying the endpoints of the minor axis.
2 | POINT | point | A single point denoted by a single (column,row) pair, or multiple points each denoted by a (column,row) pair.
2 | POLY | polyline | A series of connected line segments with ordered vertices denoted by (column,row) pairs; if the first and last vertices are the same, it is a closed polygon.
1 | C | corrected | Description:Indicates that result data has been corrected.
1 | DIET | Diet | Code set to define specialized/allowed diets
2 | BR | breikost (GE) | A diet exclusively composed of oatmeal, semolina, or rice, to be extremely easy to eat and digest.
2 | DM | diabetes mellitus diet | A diet that uses carbohydrates sparingly. Typically with a restriction in daily energy content (e.g. 1600-2000 kcal).
2 | FAST | fasting | No enteral intake of foot or liquids whatsoever, no smoking. Typically 6 to 8 hours before anesthesia.
2 | FORMULA | formula diet | A diet consisting of a formula feeding, either for an infant or an adult, to provide nutrition either orally or through the gastrointestinal tract via tube, catheter or stoma.
2 | GF | gluten free | Gluten free diet for celiac disease.
2 | LF | low fat | A diet low in fat, particularly to patients with hepatic diseases.
2 | LP | low protein | A low protein diet for patients with renal failure.
2 | LQ | liquid | A strictly liquid diet, that can be fully absorbed in the intestine, and therefore may not contain fiber. Used before enteral surgeries.
2 | LS | low sodium | A diet low in sodium for patients with congestive heart failure and/or renal failure.
2 | N | normal diet | A normal diet, i.e. no special preparations or restrictions for medical reasons. This is notwithstanding any preferences the patient might have regarding special foods, such as vegetarian, kosher, etc.
2 | NF | no fat | A no fat diet for acute hepatic diseases.
2 | PAF | phenylalanine free | Phenylketonuria diet.
2 | PAR | parenteral | Patient is supplied with parenteral nutrition, typically described in terms of i.v. medications.
2 | RD | reduction diet | A diet that seeks to reduce body fat, typically low energy content (800-1600 kcal).
2 | SCH | schonkost (GE) | A diet that avoids ingredients that might cause digestion problems, e.g., avoid excessive fat, avoid too much fiber (cabbage, peas, beans).
2 | SUPPLEMENT | nutritional supplement | A diet that is not intended to be complete but is added to other diets.
2 | T | tea only | This is not really a diet, since it contains little nutritional value, but is essentially just water. Used before coloscopy examinations.
2 | VLI | low valin, leucin, isoleucin | Diet with low content of the amino-acids valin, leucin, and isoleucin, for "maple syrup disease."
1 | DRUGPRG | drug program | Definition: A public or government health program that administers and funds coverage for prescription drugs to assist program eligible who meet financial and health status criteria.
1 | F | final | Description:Indicates that a result is complete. No further results are to come. This maps to the 'complete' state in the observation result status code.
1 | PRLMN | preliminary | Description:Indicates that a result is incomplete. There are further results to come. This maps to the 'active' state in the observation result status code.
1 | (SECOBS) Abstract | | An observation identifying security metadata about an IT resource (data, information object, service, or system capability), which may be used to make access control decisions. Security metadata are used to name security labels. Rationale: According to ISO/TS 22600-3:2009(E) A.9.1.7 SECURITY LABEL MATCHING, Security label matching compares the initiator's clearance to the target's security label. All of the following must be true for authorization to be granted: The security policy identifiers shall be identical The classification level of the initiator shall be greater than or equal to that of the target (that is, there shall be at least one value in the classification list of the clearance greater than or equal to the classification of the target), and For each security category in the target label, there shall be a security category of the same type in the initiator's clearance and the initiator's classification level shall dominate that of the target. Examples: SecurityObservationType security label fields include: Confidentiality classification Compartment category Sensitivity category Security mechanisms used to ensure data integrity or to perform authorized data transformation Indicators of an IT resource completeness, veracity, reliability, trustworthiness, or provenance. Usage Note: SecurityObservationType codes designate security label field types, which are valued with an applicable SecurityObservationValue code as the "security label tag".
2 | SECCATOBS | security category observation | Type of security metadata observation made about the category of an IT resource (data, information object, service, or system capability), which may be used to make access control decisions. Security category metadata is defined by ISO/IEC 2382-8:1998(E/F)/ T-REC-X.812-1995 as: "A nonhierarchical grouping of sensitive information used to control access to data more finely than with hierarchical security classification alone." Rationale: A security category observation supports requirement to specify the type of IT resource to facilitate application of appropriate levels of information security according to a range of levels of impact or consequences that might result from the unauthorized disclosure, modification, or use of the information or information system. A resource is assigned to a specific category of information (e.g., privacy, medical, proprietary, financial, investigative, contractor sensitive, security management) defined by an organization or in some instances, by a specific law, Executive Order, directive, policy, or regulation. [FIPS 199] Examples: Types of security categories include: Compartment: A division of data into isolated blocks with separate security controls for the purpose of reducing risk. (ISO 2382-8). A security label tag that "segments" an IT resource by indicating that access and use is restricted to members of a

defined community or project. (HL7 Healthcare Classification System) Sensitivity: The characteristic of an IT resource which implies its value or importance and may include its vulnerability. (ISO 7492-2) Privacy metadata for information perceived as undesirable to share. (HL7 Healthcare Classification System)

2 | SECCLASSOBS | security classification observation | Type of security metadata observation made about the classification of an IT resource (data, information object, service, or system capability), which may be used to make access control decisions. Security classification is defined by ISO/IEC 2382-8:1998(E/F)/ T-REC-X.812-1995 as: "The determination of which specific degree of protection against access the data or information requires, together with a designation of that degree of protection." Security classification metadata is based on an analysis of applicable policies and the risk of financial, reputational, or other harm that could result from unauthorized disclosure. Rationale: A security classification observation may indicate that the confidentiality level indicated by an Act or Role confidentiality attribute has been overridden by the entity responsible for ascribing the SecurityClassificationObservationValue. This supports the business requirement for increasing or decreasing the level of confidentiality (classification or declassification) based on parameters beyond the original assignment of an Act or Role confidentiality. Examples: Types of security classification include: HL7 Confidentiality Codes such as very restricted, unrestricted, and normal. Intelligence community examples include top secret, secret, and confidential. Usage Note: Security classification observation type codes designate security label field types, which are valued with an applicable SecurityClassificationObservationValue code as the "security label tag".

2 | SECCONOBS | security control observation | Type of security metadata observation made about the control of an IT resource (data, information object, service, or system capability), which may be used to make access control decisions. Security control metadata convey instructions to users and receivers for secure distribution, transmission, and storage; dictate obligations or mandated actions; specify any action prohibited by refrain policy such as dissemination controls; and stipulate the permissible purpose of use of an IT resource. Rationale: A security control observation supports requirement to specify applicable management, operational, and technical controls (i.e., safeguards or countermeasures) prescribed for an information system to protect the confidentiality, integrity, and availability of the system and its information. [FIPS 199] Examples: Types of security control metadata include: handling caveats dissemination controls obligations refrain policies purpose of use constraints

2 | SECINTOBS | security integrity observation | Type of security metadata observation made about the integrity of an IT resource (data, information object, service, or system capability), which may be used to make access control decisions. Rationale: A security integrity observation supports the requirement to guard against improper information modification or destruction, and includes ensuring information non-repudiation and authenticity. (44 U.S.C., SEC. 3542) Examples: Types of security integrity metadata include: Integrity status, which indicates the completeness or workflow status of an IT resource (data, information object, service, or system capability) Integrity confidence, which indicates the reliability and trustworthiness of an IT resource Integrity control, which indicates pertinent handling caveats, obligations, refrain policies, and purpose of use for the resource Data integrity, which indicate the security mechanisms used to ensure that the accuracy and consistency are preserved regardless of changes made (ISO/IEC DIS 2382-8) Alteration integrity, which indicate the security mechanisms used for authorized transformations of the resource Integrity provenance, which indicates the entity responsible for a report or assertion relayed "second-hand" about an IT resource

3 | SECALTINTOBS | security alteration integrity observation | Type of security metadata observation made about the alteration integrity of an IT resource (data, information object, service, or system capability), which indicates the mechanism used for authorized transformations of the resource. Examples: Types of security alteration integrity observation metadata, which may value the observation with a code used to indicate the mechanism used for authorized transformation of an IT resource, including: translation syntactic transformation semantic mapping redaction masking pseudonymization anonymization

3 | SECDATINTOBS | security data integrity observation | Type of security metadata observation made about the data integrity of an IT resource (data, information object, service, or system capability), which indicates the security mechanism used to preserve resource accuracy and consistency. Data integrity is defined by ISO 22600-23.3.21 as: "The property that data has not been altered or destroyed in an unauthorized manner", and by ISO/IEC 2382-8: The property of data whose accuracy and consistency are preserved regardless of changes made." Examples: Types of security data integrity observation metadata, which may value the observation, include cryptographic hash function and digital signature.

3 | SECINTCONOBS | security integrity confidence observation | Type of security metadata observation made about the integrity confidence of an IT resource (data, information object, service, or system capability), which may be used to make access control decisions. Examples: Types of security integrity confidence observation metadata, which may value the observation, include highly reliable, uncertain reliability, and not reliable. Usage Note: A security integrity confidence observation on an Act may indicate that a valued Act.uncertaintycode attribute has been overridden by the entity responsible for ascribing the SecurityIntegrityConfidenceObservationValue. This supports the business requirements for increasing or decreasing the assessment of the reliability or trustworthiness of an IT resource based on parameters beyond the original assignment of an Act statement level of uncertainty.

3 | (SECINTPRVOBS) Abstract | | Type of security metadata observation made about the provenance integrity of an IT resource (data, information object, service, or system capability), which indicates the lifecycle completeness of an IT resource in terms of workflow status such as its creation, modification, suspension, and deletion; locations in which the resource has been collected or archived, from which it may be retrieved, and the history of its distribution and disclosure. Integrity provenance metadata about an IT resource may be used to assess its veracity, reliability, and trustworthiness. Examples: Types of security integrity provenance observation metadata, which may value the observation about an IT resource, include: completeness or workflow status, such as authentication the entity responsible for original authoring or informing about an IT resource the entity responsible for a report or assertion about an IT resource relayed â€œsecond-handâ€? the entity responsible for excerpting, transforming, or compiling an IT resource

4 | SECINTPRVABOBS | security integrity provenance asserted by observation | Type of security metadata observation made about the integrity provenance of an IT resource (data, information object, service, or system capability), which indicates the entity that made assertions about the resource. The asserting entity may not be the original informant about the resource. Examples: Types of security integrity provenance asserted by observation metadata, which may value the observation, including: assertions about an IT resource by a patient assertions about an IT resource by a clinician assertions about an IT resource by a device

4 | SECINTPRVRBOBS | security integrity provenance reported by observation | Type of security metadata observation made about the integrity provenance of an IT resource (data, information object, service, or system capability), which indicates the entity that reported the existence of the resource. The reporting entity may not be the original author of the resource. Examples: Types of security integrity provenance reported by observation metadata, which may value the observation, include: reports about an IT resource by a patient reports about an IT resource by a clinician reports about an IT resource by a device

3 | SECINTSTOBS | security integrity status observation | Type of security metadata observation made about the integrity status of an IT resource (data, information object, service, or system capability), which may be used to make access control decisions. Indicates the completeness of an IT resource in terms of workflow status, which may impact users that are authorized to access and use the resource. Examples: Types of security integrity status observation metadata, which may value the observation, include codes from the HL7 DocumentCompletion code system such as legally authenticated, in progress, and incomplete.

2 | (SECTRSTOBS) Abstract | | An observation identifying trust metadata about an IT resource (data, information object, service, or system capability), which may be used as a trust attribute to populate a computable trust policy, trust credential, trust assertion, or trust label field in a security label or trust policy, which are principally used for authentication, authorization, and access control decisions.

3 | TRSTACCRDOBS | trust accreditation observation | Type of security metadata observation made about the formal declaration by an authority or neutral third party that validates the technical, security, trust, and business practice conformance of Trust Agents to facilitate security, interoperability, and trust among participants within a security domain or trust framework.

3 | TRSTAGREOBS | trust agreement observation | Type of security metadata observation made about privacy and security requirements with which a security domain must comply. [ISO IEC 10181-1]

3 | TRSTCERTOBS | trust certificate observation | Type of security metadata observation made about a set of security-relevant data issued by a security authority or trusted third party, together with security information which is used to provide the integrity and data origin authentication services for an IT resource (data, information object, service, or system capability). [Based on ISO IEC 10181-1] For example, A Certificate Policy (CP), which is a named set of rules that indicates the applicability of a certificate to a particular community and/or class of application with common security requirements. For example, a particular Certificate Policy might indicate the applicability of a type of certificate to the authentication of electronic data interchange transactions for the trading of goods within a given price range. [Trust Service Principles and Criteria for Certification Authorities Version 2.0 March 2011 Copyright 2011 by Canadian Institute of Chartered Accountants. A Certificate Practice Statement (CSP), which is a statement of the practices which an Authority employs in issuing and managing certificates. [Trust Service Principles and Criteria for Certification Authorities Version 2.0 March 2011 Copyright 2011 by Canadian Institute of Chartered Accountants.]

3 | TRSTFWKOBS | trust framework observation | Type of security metadata observation made about a complete set of contracts, regulations or commitments that enable participating actors to rely on certain assertions by other actors to fulfill their information security requirements. [Kantara Initiative]

3 | TRSTLOAOBS | trust assurance observation | Type of security metadata observation made about the digital quality or reliability of a trust assertion, activity, capability, information exchange, mechanism, process, or protocol.

3 | TRSTMECOBS | trust mechanism observation | Type of security metadata observation made about a security architecture system component that supports enforcement of security policies.

1 | SUBSIDFFS | subsidized fee for service program | Definition: A government health program that provides coverage on a fee for service basis for health services to persons meeting eligibility criteria such as income, location of residence, access to other coverages, health condition, and age, the cost of which is to some extent subsidized by public funds. Discussion: The structure and business processes for underwriting and administering a subsidized fee for service program is further specified by the Underwriter and Payer Role.class and Role.code.

1 | WRKCOMP | (workers compensation program | Definition: Government mandated program providing coverage, disability income, and vocational rehabilitation for injuries sustained in the work place or in the course of employment. Employers may either self-fund the program, purchase commercial coverage, or pay a premium to a government entity that administers the program. Employees may be required to pay premiums toward the cost of coverage as well.

1 | (_ActProcedureCode) Abstract | | An identifying code for healthcare interventions/procedures.

2 | (_ActBillableServiceCode) Abstract | | Definition: An identifying code for billable services, as opposed to codes for similar services used to identify them for functional purposes.

1 | (_HL7DefinedActCodes) Abstract | | Domain provides the root for HL7-defined detailed or rich codes for the Act classes.

1 | COPAY

1 | DEDUCT

1 | DOSEIND

1 | PRA

1 | STORE Deprecated | Storage | The act of putting something away for safe keeping. The "something" may be physical object such as a specimen, or information, such as observations regarding a specimen.

4.4.2.6 HL7 v3 Code System ActClass

URL: https://hl7.org/fhir/R4/v3/ActClass/cs.html

Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-ActClass) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-ActClass

Version: | 2018-08-12

Name: | v3.ActClass

Title: | v3 Code System ActClass

Definition: | **** MISSING DEFINITIONS ****

OID: | 2.16.840.1.113883.5.6 (for OID based terminology systems)

Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: V3 Value SetActClassObservation (Description: An act that is intended to result in new information about a subject. The main difference between Observations and other Acts is that Observations have a value attribute. Thecode attribute of Observation and thevalue attribute of Observation must be considered in combination to determine the semantics of the observation. Discussion: Structurally, many observations are name-value-pairs, where the Observation.code (inherited from Act) is the name and the Observation.value is the value of the property. Such a construct is also known as a variable (a named feature that can assume a value) hence, the Observation class is always used to hold generic name-value-pairs or variables, even though the variable valuation may not be the result of an elaborate observation method. It may be a simple answer to a question or it may be an assertion or setting of a parameter. As with all Act statements, Observation statements describe what was done, and in the case of Observations, this includes a description of what was actually observed (results or answers); and those results or answers are part of the

observation and not split off into other objects. The method of action is asserted by the Observation classCode or its subclasses at the least granular level, by the Observation.code attribute value at the medium level of granularity, and by the attribute value of observation.methodCode when a finer level of granularity is required. The method in whole or in part may also appear in the attribute value of Observation.value when using coded data types to express the value of the attribute. Relevant aspects of methodology may also be restated in value when the results themselves imply or state a methodology. An observation may consist of component observations each having their own Observation.code and Observation.value. In this case, the composite observation may not have an Observation.value for itself. For instance, a white blood cell count consists of the sub-observations for the counts of the various granulocytes, lymphocytes and other normal or abnormal blood cells (e.g., blasts). The overall white blood cell count Observation itself may therefore not have a value by itself (even though it could have one, e.g., the sum total of white blood cells). Thus, as long as an Act is essentially an Act of recognizing and noting information about a subject, it is an Observation, regardless of whether it has a simple value by itself or whether it has sub-observations. Even though observations are professional acts (see Act) and as such are intentional actions, this does not require that every possible outcome of an observation be pondered in advance of it being actually made. For instance, differential white blood cell counts (WBC) rarely show blasts, but if they do, this is part of the WBC observation even though blasts might not be predefined in the structure of a normal WBC. Clinical documents commonly have Subjective and Objective findings, both of which are kinds of Observations. In addition, clinical documents commonly contain Assessments, which are also kinds of Observations. Thus, the establishment of a diagnosis is an Observation. Examples: Recording the results of a Family History Assessment Laboratory test and associated result Physical exam test and associated result Device temperature Soil lead level)

• ValueSet: V3 Value SetActClassDocument (Specialization of Act to add the characteristics unique to document management services.)

• ValueSet: v3 Code System ActClass (**** MISSING DEFINITIONS ****)

• ValueSet: V3 Value SetActClassInvestigation (An formalized inquiry into the circumstances surrounding a particular unplanned event or potential event for the purposes of identifying possible causes and contributing factors for the event. This investigation could be conducted at a local institutional level or at the level of a local or national government.)

• ValueSet: V3 Value SetActClassClinicalDocument (A clinical document is a documentation of clinical observations and services, with the following characteristics: (1) Persistence - A clinical document continues to exist in an unaltered state, for a time period defined by local and regulatory requirements; (2) Stewardship - A clinical document is maintained by a person or organization entrusted with its care; (3) Potential for authentication - A clinical document is an assemblage of information that is intended to be legally authenticated; (4) Wholeness - Authentication of a clinical document applies to the whole and does not apply to portions of the document without the full context of the document; (5) Human readability - A clinical document is human readable.")

• ValueSet: V3 Value SetActClassROI (Regions of Interest (ROI) within a subject Act. Primarily used for making secondary observations on a subset of a subject observation. The relationship between a ROI and its referenced Act is specified through an ActRelationship of type "subject" (SUBJ), which must always be present.)

• ValueSet: V3 Value SetActClassProcedure (An Act whose immediate and primary outcome (post-condition) is the alteration of the physical condition of the subject. Examples: : Procedures may involve the disruption of some body surface (e.g. an incision in a surgical procedure), but they also include conservative procedures such as reduction of a luxated join, chiropractic treatment, massage, balneotherapy, acupuncture, shiatsu, etc. Outside of clinical medicine, procedures may be such things as alteration of environments (e.g. straightening rivers, draining swamps, building dams) or the repair or change of machinery etc.)

• ValueSet: V3 Value SetActClassSupply (Supply orders and deliveries are simple Acts that focus on the delivered product. The product is associated with the Supply Act via Participation.typeCode="product". With general Supply Acts, the precise identification of the Material (manufacturer, serial numbers, etc.) is important. Most of the detailed information about the Supply should be represented using the Material class. If delivery needs to be scheduled, tracked, and billed separately, one can associate a Transportation Act with the Supply Act. Pharmacy dispense services are represented as Supply Acts, associated with a SubstanceAdministration Act. The SubstanceAdministration class represents the administration of medication, while dispensing is supply.)

4.4.2.6.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition

1 | ACT | act | A record of something that is being done, has been done, can be done, or is intended or requested to be done. Examples:The kinds of acts that are common in health care are (1) a clinical observation, (2) an assessment of health condition (such as problems and diagnoses), (3) healthcare goals, (4) treatment services (such as medication, surgery, physical and psychological therapy), (5) assisting, monitoring or attending, (6) training and education services to patients and their next of kin, (7) and notary services (such as advanced directives or living will), (8) editing and maintaining documents, and many others. Discussion and Rationale: Acts are the pivot of the RIM; all domain information and processes are represented primarily in Acts. Any profession or business, including healthcare, is primarily constituted of intentional and occasionally non-intentional actions,

performed and recorded by responsible actors. An Act-instance is a record of such an action. Acts connect to Entities in their Roles through Participations and connect to other Acts through ActRelationships. Participations are the authors, performers and other responsible parties as well as subjects and beneficiaries (which includes tools and material used in the performance of the act, which are also subjects). The moodCode distinguishes between Acts that are meant as factual records, vs. records of intended or ordered services, and the other modalities in which act can appear. One of the Participations that all acts have (at least implicitly) is a primary author, who is responsible of the Act and who "owns" the act. Responsibility for the act means responsibility for what is being stated in the Act and as what it is stated. Ownership of the act is assumed in the sense of who may operationally modify the same act. Ownership and responsibility of the Act is not the same as ownership or responsibility of what the Act-object refers to in the real world. The same real world activity can be described by two people, each being the author of their Act, describing the same real world activity. Yet one can be a witness while the other can be a principal performer. The performer has responsibilities for the physical actions; the witness only has responsibility for making a true statement to the best of his or her ability. The two Act-instances may even disagree, but because each is properly attributed to its author, such disagreements can exist side by side and left to arbitration by a recipient of these Act-instances. In this sense, an Act-instance represents a "statement" according to Rector and Nowlan (1991) [Foundations for an electronic medical record. Methods Inf Med. 30.] Rector and Nowlan have emphasized the importance of understanding the medical record not as a collection of facts, but "a faithful record of what clinicians have heard, seen, thought, and done." Rector and Nowlan go on saying that "the other requirements for a medical record, e.g., that it be attributable and permanent, follow naturally from this view." Indeed the Act class is this attributable statement, and the rules of updating acts (discussed in the state-transition model, see Act.statusCode) versus generating new Act-instances are designed according to this principle of permanent attributable statements. Rector and Nolan focus on the electronic medical record as a collection of statements, while attributed statements, these are still mostly factual statements. However, the Act class goes beyond this limitation to attributed factual statements, representing what is known as "speech-acts" in linguistics and philosophy. The notion of speech-act includes that there is pragmatic meaning in language utterances, aside from just factual statements; and that these utterances interact with the real world to change the state of affairs, even directly cause physical activities to happen. For example, an order is a speech act that (provided it is issued adequately) will cause the ordered action to be physically performed. The speech act theory has culminated in the seminal work by Austin (1962) [How to do things with words. Oxford University Press]. An activity in the real world may progress from defined, through planned and ordered to executed, which is represented as the mood of the Act. Even though one might think of a single activity as progressing from planned to executed, this progression is reflected by multiple Act-instances, each having one and only one mood that will not change along the Act-instance life cycle. This is because the attribution and content of speech acts along this progression of an activity may be different, and it is often critical that a permanent and faithful record be maintained of this progression. The specification of orders or promises or plans must not be overwritten by the specification of what was actually done, so as to allow comparing actions with their earlier specifications. Act-instances that describe this progression of the same real world activity are linked through the ActRelationships (of the relationship category "sequel"). Act as statements or speech-acts are the only representation of real world facts or processes in the HL7 RIM. The truth about the real world is constructed through a combination (and arbitration) of such attributed statements only, and there is no class in the RIM whose objects represent "objective state of affairs" or "real processes" independent from attributed statements. As such, there is no distinction between an activity and its documentation. Every Act includes both to varying degrees. For example, a factual statement made about recent (but past) activities, authored (and signed) by the performer of such activities, is commonly known as a procedure report or original documentation (e.g., surgical procedure report, clinic note etc.). Conversely, a status update on an activity that is presently in progress, authored by the performer (or a close observer) is considered to capture that activity (and is later superceded by a full procedure report). However, both status update and procedure report are acts of the same kind, only distinguished by mood and state (see statusCode) and completeness of the information.

2 | (_ActClassRecordOrganizer) Abstract | | Used to group a set of acts sharing a common context. Organizer structures can nest within other context structures - such as where a document is contained within a folder, or a folder is contained within an EHR extract.

3 | COMPOSITION | composition | A context representing a grouped commitment of information to the EHR. It is considered the unit of modification of the record, the unit of transmission in record extracts, and the unit of attestation by authorizing clinicians. A composition represents part of a patient record originating from a single interaction between an authenticator and the record. Unless otherwise stated all statements within a composition have the same authenticator, apply to the same patient and were recorded in a single session of use of a single application. A composition contains organizers and entries.

4 | DOC | document | The notion of a document comes particularly from the paper world, where it corresponds to the contents recorded on discrete pieces of paper. In the electronic world, a document is a kind of composition that bears resemblance to their paper world counter-parts. Documents typically are meant to be human-readable. HL7's notion of document differs from that described in the W3C XML Recommendation, in which a document refers specifically to the contents that fall between the root element's start-tag and end-tag. Not all XML documents are HL7 documents.

5 | DOCCLIN | clinical document | A clinical document is a documentation of clinical observations and services, with the following characteristics: Persistence - A clinical document continues to exist in an unaltered state, for a time period defined by local and regulatory requirements; Stewardship - A clinical document is maintained by a person or organization entrusted with its care; Potential for authentication - A clinical document is an assemblage of information that is intended to be legally authenticated; Wholeness - Authentication of a clinical document applies to the whole and does not apply to portions of the document without the full context of the document; Human readability - A clinical document is human readable.

6 | CDALVLONE | CDA Level One clinical document | A clinical document that conforms to Level One of the HL7 Clinical Document Architecture (CDA)

3 | CONTAINER | record container | Description: Container of clinical statements. Navigational. No semantic content. Knowledge of the section code is not required to interpret contained observations. Represents a heading in a heading structure, or "container tree". The record entries relating to a single clinical session are usually grouped under headings that represent phases of the encounter, or assist with layout and navigation. Clinical headings usually reflect the clinical workflow during a care session, and might also reflect the main author's reasoning processes. Much research has demonstrated that headings are used differently by different professional groups and specialties, and that headings are not used consistently enough to support safe automatic processing of the E H R.

4 | CATEGORY | category | A group of entries within a composition or topic that have a common characteristic - for example, Examination, Diagnosis, Management OR Subjective, Objective, Analysis, Plan. The distinction from Topic relates to value sets. For Category there is a bounded list of things like "Examination", "Diagnosis" or SOAP categories. For Topic the list is wide open to any clinical condition or reason for a part of an encounter. A CATEGORY MAY CONTAIN ENTRIES.

4 | DOCBODY | document body | A context that distinguishes the body of a document from the document header. This is seen, for instance, in HTML documents, which have discrete <head> and <body> elements.

4 | DOCSECT | document section | A context that subdivides the body of a document. Document sections are typically used for human navigation, to give a reader a clue as to the expected content. Document sections are used to organize and provide consistency to the contents of a document body. Document sections can contain document sections and can contain entries.

4 | TOPIC | topic | A group of entries within a composition that are related to a common clinical theme - such as a specific disorder or problem, prevention, screening and provision of contraceptive services. A topic may contain categories and entries.
3 | EXTRACT | extract | This context represents the part of a patient record conveyed in a single communication. It is drawn from a providing system for the purposes of communication to a requesting process (which might be another repository, a client application or a middleware service such as an electronic guideline engine), and supporting the faithful inclusion of the communicated data in the receiving system. An extract may be the entirety of the patient record as held by the sender or it may be a part of that record (e.g. changes since a specified date). An extract contains folders or compositions. An extract cannot contain another extract.
4 | EHR | electronic health record | A context that comprises all compositions. The EHR is an extract that includes the entire chart. NOTE: In an exchange scenario, an EHR is a specialization of an extract.
3 | FOLDER | folder | A context representing the high-level organization of an extract e.g. to group parts of the record by episode, care team, clinical specialty, clinical condition, or source application. Internationally, this kind of organizing structure is used variably: in some centers and systems the folder is treated as an informal compartmentalization of the overall health record; in others it might represent a significant legal portion of the EHR relating to the originating enterprise or team. A folder contains compositions. Folders may be nested within folders.
3 | GROUPER | grouper | Definition: An ACT that organizes a set of component acts into a semantic grouping that share a particular context such as timeframe, patient, etc. UsageNotes: The focus in a GROUPER act is the grouping of the contained acts. For example "a request to group" (RQO), "a type of grouping that is allowed to occur" (DEF), etc. Unlike WorkingList, which represents a dynamic, shared, continuously updated collection to provide a "view" of a set of objects, GROUPER collections tend to be static and simply indicate a shared set of semantics. Note that sharing of semantics can be achieved using ACT as well. However, with GROUPER, the sole semantic is of grouping.
4 | CLUSTER | Cluster | Description:An ACT that organizes a set of component acts into a semantic grouping that have a shared subject. The subject may be either a subject participation (SBJ), subject act relationship (SUBJ), or child participation/act relationship types. Discussion: The focus in a CLUSTER act is the grouping of the contained acts. For example "a request to cluster" (RQO), "a type of cluster that is allowed to occur" (DEF), etc. Examples: Radiologic investigations that might include administration of a dye, followed by radiographic observations; "Isolate cluster" which includes all testing and specimen processing performed on a specific isolate; a set of actions to perform at a particular stage in a clinical trial.
2 | ACCM | accommodation | An accommodation is a service provided for a Person or other LivingSubject in which a place is provided for the subject to reside for a period of time. Commonly used to track the provision of ward, private and semi-private accommodations for a patient.
2 | ACCT | account | A financial account established to track the net result of financial acts.
2 | ACSN | accession | A unit of work, a grouper of work items as defined by the system performing that work. Typically some laboratory order fulfillers communicate references to accessions in their communications regarding laboratory orders. Often one or more specimens are related to an accession such that in some environments the accession number is taken as an identifier for a specimen (group).
2 | ADJUD | financial adjudication | A transformation process where a requested invoice is transformed into an agreed invoice. Represents the adjudication processing of an invoice (claim). Adjudication results can be adjudicated as submitted, with adjustments or refused. Adjudication results comprise 2 components: the adjudication processing results and a restated (or adjudicated) invoice or claim
2 | CACT | control act | An act representing a system action such as the change of state of another act or the initiation of a query. All control acts represent trigger events in the HL7 context. ControlActs may occur in different moods.
3 | ACTN | action | Sender asks addressee to do something depending on the focal Act of the payload. An example is "fulfill this order". Addressee has responsibilities to either reject the message or to act on it in an appropriate way (specified by the specific receiver responsibilities for the interaction).
3 | INFO | information | Sender sends payload to addressee as information. Addressee does not have responsibilities beyond serving addressee's own interest (i.e., read and memorize if you see fit). This is equivalent to an FYI on a memo.
3 | STC | state transition control | Description: Sender transmits a status change pertaining to the focal act of the payload. This status of the focal act is the final state of the state transition. This can be either a request or an event, according to the mood of the control act.
2 | CNTRCT | contract | An agreement of obligation between two or more parties that is subject to contractual law and enforcement.
3 | FCNTRCT | financial contract | A contract whose value is measured in monetary terms.
4 | COV | coverage | When used in the EVN mood, this concept means with respect to a covered party: A health care insurance policy or plan that is contractually binding between two or more parties; or A health care program, usually administered by government entities, that provides coverage to persons determined eligible under the terms of the program. When used in the definition (DEF) mood, COV means potential coverage for a patient who may or may not be a covered party. The concept's meaning is fully specified by the choice of ActCoverageTypeCode (abstract) ActProgramCode or ActInsurancePolicyCode.
2 | CONC | concern | Definition: A worry that tends to persist over time and has as its subject a state or process. The subject of the worry has the potential to require intervention or management. Examples: an observation result, procedure, substance administration, equipment repair status, device recall status, a health risk, a financial risk, public health risk, pregnancy, health maintenance, allergy, and acute or chronic illness.
3 | HCASE | public health case | A public health case is a Concern about an observation or event that has a specific significance for public health. The creation of a PublicHealthCase initiates the tracking of the object of concern. The decision to track is related to but somewhat independent of the underlying event or observation. UsageNotes: Typically a Public Health Case involves an instance or instances of a reportable infectious disease or other condition. The public health case can include a health-related event concerning a single individual or it may refer to multiple health-related events that are occurrences of the same disease or condition of interest to public health. A public health case definition (Act.moodCode = "definition") includes the description of the clinical, laboratory, and epidemiologic indicators associated with a disease or condition of interest to public health. There are case definitions for conditions that are reportable, as well as for those that are not. A public health case definition is a construct used by public health for the purpose of counting cases, and should not be used as clinical indications for treatment. Examples include AIDS, toxic-shock syndrome, and salmonellosis and their associated indicators that are used to define a case.
3 | OUTBR | outbreak | An Outbreak is a concern resulting from a series of public health cases. UsageNotes: The date on which an outbreak starts is the earliest date of onset among the cases assigned to the outbreak and its ending date is the last date of onset among the cases assigned to the outbreak. The effectiveTime attribute is used to convey the relevant dates for the case. An outbreak definition (Act.moodCode = "definition" includes the criteria for the number, types and occurrence pattern of cases necessary to declare an outbreak and to judge the severity of an outbreak.
2 | CONS | consent | The Consent class represents informed consents and all similar medico-legal transactions between the patient (or his legal guardian) and the provider. Examples are informed consent for surgical procedures, informed consent for clinical

trials, advanced beneficiary notice, against medical advice decline from service, release of information agreement, etc. The details of consents vary. Often an institution has a number of different consent forms for various purposes, including reminding the physician about the topics to mention. Such forms also include patient education material. In electronic medical record communication, consents thus are information-generating acts on their own and need to be managed similar to medical activities. Thus, Consent is modeled as a special class of Act. The "signatures" to the consent document are represented electronically through Participation instances to the consent object. Typically an informed consent has Participation.typeCode of "performer", the healthcare provider informing the patient, and "consenter", the patient or legal guardian. Some consent may associate a witness or a notary public (e.g., living wills, advanced directives). In consents where a healthcare provider is not required (e.g. living will), the performer may be the patient himself or a notary public. Some consent has a minimum required delay between the consent and the service, so as to allow the patient to rethink his decisions. This minimum delay can be expressed in the act definition by the ActRelationship.pauseQuantity attribute that delays the service until the pause time has elapsed after the consent has been completed.

2 | CONTREG | container registration | An Act where a container is registered either via an automated sensor, such as a barcode reader, or by manual receipt

2 | CTTEVENT | clinical trial timepoint event | An identified point during a clinical trial at which one or more actions are scheduled to be performed (definition mood), or are actually performed (event mood). The actions may or may not involve an encounter between the subject and a healthcare professional.

2 | DISPACT | disciplinary action | An action taken with respect to a subject Entity by a regulatory or authoritative body with supervisory capacity over that entity. The action is taken in response to behavior by the subject Entity that body finds to be undesirable. Suspension, license restrictions, monetary fine, letter of reprimand, mandated training, mandated supervision, etc.Examples:

2 | EXPOS | exposure | An interaction between entities that provides opportunity for transmission of a physical, chemical, or biological agent from an exposure source entity to an exposure target entity. Examples: The following examples are provided to indicate what interactions are considered exposures rather than other types of Acts: A patient accidentally receives three times the recommended dose of their medication due to a dosing error. This is a substance administration. Public health and/or safety authorities may also be interested in documenting this with an associated exposure. A patient accidentally is dispensed an incorrect medicine (e.g., clomiphene instead of clomipramine). They have taken several doses before the mistake is detected. They are therefore "exposed" to a medicine that there was no therapeutic indication for them to receive. There are several substance administrations in this example. Public health and/or safety authorities may also be interested in documenting this with associated exposures. In a busy medical ward, a patient is receiving chemotherapy for a lymphoma. Unfortunately, the IV infusion bag containing the medicine splits, spraying cytotoxic medication over the patient being treated and the patient in the adjacent bed. There are three substance administrations in this example. The first is the intended one (IV infusion) with its associated (implicit) exposure. There is an incident with an associated substance administration to the same patient involving the medication sprayed over the patient as well as an associated exposure. Additionally, the incident includes a substance administration involving the spraying of medication on the adjacent patient, also with an associated exposure. A patient who is a refugee from a war-torn African nation arrives in a busy inner city A&E department suffering from a cough with bloody sputum. Not understanding the registration and triage process, they sit in the waiting room for several hours before it is noticed that they have not booked in. As soon as they are being processed, it is suspected that they are suffering from TB. Vulnerable (immunosuppressed) patients who were sharing the waiting room with this patient may have been exposed to the tubercule bacillus, and must be traced for investigation. This is an exposure (or possibly multiple exposures) in the waiting room involving the refugee and everyone else in the waiting room during the period. There might also be a number of known or presumed substance administrations (coughing) via several possible routes. The substance administrations are only hypotheses until confirmed by further testing. A patient who has received an elective total hip replacement procedure suffers a prolonged stay in hospital, due to contracting an MRSA infection in the surgical wound site after the surgery. This is an exposure to MRSA. Although there was some sort of substance administration, it's possible the exact mechanism for introduction of the MRSA into the wound will not be identified. Routine maintenance of the X-ray machines at a local hospital reveals a serious breach of the shielding on one of the machines. Patients who have undergone investigations using that machine in the last month are likely to have been exposed to significantly higher doses of X-rays than was intended, and must be tracked for possible adverse effects. There has been an exposure of each patient who used the machine in the past 30 days. Some patients may have had substance administrations. A new member of staff is employed in the laundry processing room of a small cottage hospital, and a misreading of the instructions for adding detergents results in fifty times the usual concentration of cleaning materials being added to a batch of hospital bedding. As a result, several patients have been exposed to very high levels of detergents still present in the "clean" bedding, and have experienced dermatological reactions to this. There has been an incident with multiple exposures to several patients. Although there are substance administrations involving the application of the detergent to the skin of the patients, it is expected that the substance administrations would not be directly documented. Seven patients who are residents in a health care facility for the elderly mentally ill have developed respiratory problems. After several months of various tests having been performed and various medications prescribed to these patients, the problem is traced to their being "sensitive" to a new fungicide used in the wall plaster of the ward where these patients reside. The patients have been continuously exposed to the fungicide. Although there have been continuous substance administrations (via breathing) this would not normally be documented as a substance administration. A patient with osteoarthritis of the knees is treated symptomatically using analgesia, paracetamol (acetaminophen) 1g up to four times a day for pain relief. His GP does not realize that the patient has, 20 years previously (while at college) had severe alcohol addiction problems, and now, although this is completely under control, his liver has suffered significantly, leaving him more sensitive to hepatic toxicity from paracetamol use. Later that year, the patient returns with a noticeable level of jaundice. Paracetamol is immediately withdrawn and alternative solutions for the knee pain are sought. The jaundice gradually subsides with conservative management, but referral to the gastroenterologist is required for advice and monitoring. There is a substance administration with an associated exposure. The exposure component is based on the relative toxic level of the substance to a patient with a compromised liver function. A patient goes to their GP complaining of abdominal pain, having been discharged from the local hospital ten days' previously after an emergency appendectomy. The GP can find nothing particularly amiss, and presumes it is post operative surgical pain that will resolve. The patient returns a fortnight later, when the GP prescribes further analgesia, but does decide to request an outpatient surgical follow-up appointment. At this post-surgical outpatient review, the registrar decides to order an ultrasound, which, when performed three weeks later, shows a small faint inexplicable mass. A laparoscopy is then performed, as a day case procedure, and a piece of a surgical swab is removed from the patient's abdominal cavity. Thankfully, a full recovery then takes place. This is a procedural sequelae. There may be an Incident recorded for this also. A patient is slightly late for a regular pacemaker battery check in the Cardiology department of the local hospital. They are hurrying down the second floor corridor. A sudden summer squall has recently passed over the area, and rain has come in through an open corridor window leaving a small puddle on the corridor floor. In their haste, the patient slips in the puddle and falls so badly that they have to be taken to the A&E department, where it is discovered on investigation they have slightly torn the cruciate ligament in their left knee. This is not an exposure. There has been an incident. Usage Notes: This class deals only with opportunity and not the outcome of the exposure; i.e. not all exposed parties will

necessarily experience actual harm or benefit. Exposure differs from Substance Administration by the absence of the participation of a performer in the act. The following participations SHOULD be used with the following participations to distinguish the specific entities: The exposed entity participates via the "exposure target" (EXPTRGT) participation. An entity that has carried the agent transmitted in the exposure participates via the "exposure source" (EXSRC) participation. For example: a person or animal who carried an infectious disease and interacts (EXSRC) with another person or animal (EXPTRGT) transmitting the disease agent; a place or other environment (EXSRC) and a person or animal (EXPTRGT) who is exposed in the presence of this environment. When it is unknown whether a participating entity is the source of the agent (EXSRC) or the target of the transmission (EXPTRGT), the "exposure participant" (EXPART) is used. The physical (including energy), chemical or biological substance which is participating in the exposure uses the "exposure agent" (EXPAGNT) participation. There are at least three scenarios: the player of the Role that participates as EXPAGNT is the chemical or biological substance mixed or carried by the scoper-entity of the Role (e.g., ingredient role); or the player of the Role that participates as EXPAGNT is a mixture known to contain the chemical, radiological or biological substance of interest; or the player of the Role that participates as a EXPAGNT is known to carry the agent (i.e., the player is a fomite, vector, etc.). The Exposure.statusCode attribute should be interpreted as the state of the Exposure business object (e.g., active, aborted, completed) and not the clinical status of the exposure (e.g., probable, confirmed). The clinical status of the exposure should be associated with the exposure via a subject observation. Design Comment: The usage notes require a clear criterion for determining whether an act is an exposure or substance administration-deleterious potential, uncertainty of actual transmission, or otherwise. SBADM states that the criterion is the presence of a performer-but there are examples above that call this criterion into question (e.g., the first one, concerning a dosing error).

3 | AEXPOS | acquisition exposure | Description: An acquisition exposure act describes the proximity (location and time) through which the participating entity was potentially exposed to a physical (including energy), chemical or biological agent from another entity. The acquisition exposure act is used in conjunction with transmission exposure acts as part of an analysis technique for contact tracing. Although an exposure can be decomposed into transmission and acquisition exposures, there is no requirement that all exposures be treated in this fashion. Constraints: The Acquisition Exposure inherits the participation constraints that apply to Exposure with the following exception. The EXPSRC (exposure source) participation must never be associated with the Transmission Exposure either directly or via context conduction.

3 | TEXPOS | transmission exposure | Description: A transmission exposure act describes the proximity (time and location) over which the participating source entity was capable of transmitting a physical (including energy), chemical or biological substance agent to another entity. The transmission exposure act is used in conjunction with acquisition exposure acts as part of an analysis technique for contact tracing. Although an exposure can be decomposed into transmission and acquisition exposures, there is no requirement that all exposures be treated in this fashion. Constraints: The Transmission Exposure inherits the participation constraints that apply to Exposure with the following exception. The EXPTRGT (exposure target) participation must never be associated with the Transmission Exposure either directly or via context conduction.

2 | INC | incident | An event that occurred outside of the control of one or more of the parties involved. Includes the concept of an accident.

2 | INFRM | inform | The act of transmitting information and understanding about a topic to a subject where the participation association must be SBJ. Discussion: This act may be used to request that a patient or provider be informed about an Act, or to indicate that a person was informed about a particular act.

2 | INVE | invoice element | Represents concepts related to invoice processing in health care

2 | LIST | working list | Working list collects a dynamic list of individual instances of Act via ActRelationship which reflects the need of an individual worker, team of workers, or an organization to manage lists of acts for many different clinical and administrative reasons. Examples of working lists include problem lists, goal lists, allergy lists, and to-do lists.

2 | MPROT | monitoring program | An officially or unofficially instituted program to track acts of a particular type or categorization.

2 | OBS | observation | Description:An act that is intended to result in new information about a subject. The main difference between Observations and other Acts is that Observations have a value attribute. The code attribute of Observation and the value attribute of Observation must be considered in combination to determine the semantics of the observation. Discussion: Structurally, many observations are name-value-pairs, where the Observation.code (inherited from Act) is the name and the Observation.value is the value of the property. Such a construct is also known as a variable (a named feature that can assume a value) hence, the Observation class is always used to hold generic name-value-pairs or variables, even though the variable valuation may not be the result of an elaborate observation method. It may be a simple answer to a question or it may be an assertion or setting of a parameter. As with all Act statements, Observation statements describe what was done, and in the case of Observations, this includes a description of what was actually observed (results or answers); and those results or answers are part of the observation and not split off into other objects. The method of action is asserted by the Observation classCode or its subclasses at the least granular level, by the Observation.code attribute value at the medium level of granularity, and by the attribute value of observation.methodCode when a finer level of granularity is required. The method in whole or in part may also appear in the attribute value of Observation.value when using coded data types to express the value of the attribute. Relevant aspects of methodology may also be restated in value when the results themselves imply or state a methodology. An observation may consist of component observations each having their own Observation.code and Observation.value. In this case, the composite observation may not have an Observation.value for itself. For instance, a white blood cell count consists of the sub-observations for the counts of the various granulocytes, lymphocytes and other normal or abnormal blood cells (e.g., blasts). The overall white blood cell count Observation itself may therefore not have a value by itself (even though it could have one, e.g., the sum total of white blood cells). Thus, as long as an Act is essentially an Act of recognizing and noting information about a subject, it is an Observation, regardless of whether it has a simple value by itself or whether it has sub-observations. Even though observations are professional acts (see Act) and as such are intentional actions, this does not require that every possible outcome of an observation be pondered in advance of it being actually made. For instance, differential white blood cell counts (WBC) rarely show blasts, but if they do, this is part of the WBC observation even though blasts might not be predefined in the structure of a normal WBC. Clinical documents commonly have Subjective and Objective findings, both of which are kinds of Observations. In addition, clinical documents commonly contain Assessments, which are also kinds of Observations. Thus, the establishment of a diagnosis is an Observation. Examples: Recording the results of a Family History Assessment Laboratory test and associated result Physical exam test and associated result Device temperature Soil lead level

3 | (_ActClassROI) Abstract | | Regions of Interest (ROI) within a subject Act. Primarily used for making secondary observations on a subset of a subject observation. The relationship between a ROI and its referenced Act is specified through an ActRelationship of type "subject" (SUBJ), which must always be present.

4 | ROIBND | bounded ROI | A Region of Interest (ROI) specified for a multidimensional observation, such as an Observation Series (OBSSER). The ROI is specified using a set of observation criteria, each delineating the boundary of the region in one of the dimensions in the multidimensional observation. The relationship between a ROI and its referenced Act is specified through an ActRelationship of type subject (SUBJ), which must always be present. Each of the boundary criteria observations is connected with the ROI using ActRelationships of type "has component" (COMP). In each boundary criterion, the Act.code names the dimension and the Observation.value specifies the range of values inside the region. Typically the bounded dimension is continuous, and so the Observation.value will be an interval (IVL) data type. The Observation.value need not be specified if the

respective dimension is only named but not constrained. For example, an ROI for the QT interval of a certain beat in ECG Lead II would contain 2 boundary criteria, one naming the interval in time (constrained), and the other naming the interval in ECG Lead II (only named, but not constrained).

4 | ROIOVL | overlay ROI | A Region of Interest (ROI) specified for an image using an overlay shape. Typically used to make reference to specific regions in images, e.g., to specify the location of a radiologic finding in an image or to specify the site of a physical finding by "circling" a region in a schematic picture of a human body. The units of the coordinate values are in pixels. The origin is in the upper left hand corner, with positive X values going to the right and positive Y values going down. The relationship between a ROI and its referenced Act is specified through an ActRelationship of type "subject" (SUBJ), which must always be present.

3 | (_SubjectPhysicalPosition) Abstract | | The spatial relationship of a subject whether human, other animal, or plant, to a frame of reference such as gravity or a collection device.

4 | (_SubjectBodyPosition) Abstract Deprecated | | Contains codes for defining the observed, physical position of a subject, such as during an observation, assessment, collection of a specimen, etc. ECG waveforms and vital signs, such as blood pressure, are two examples where a general, observed position typically needs to be noted. Deprecation Comment: This concept has been deprecated because it does not describe a type of Act (as it should in the ActClass code system), but rather encodes the result or value of an observation. The same code has been added to the ObservationValue code system.

5 | LLD Deprecated | left lateral decubitus | Lying on the left side. Deprecation Comment: This concept has been deprecated because it does not describe a type of Act (as it should in the ActClass code system), but rather encodes the result or value of an observation. The same code has been added to the ObservationValue code system.

5 | PRN Deprecated | prone | Lying with the front or ventral surface downward; lying face down. Deprecation Comment: This concept has been deprecated because it does not describe a type of Act (as it should in the ActClass code system), but rather encodes the result or value of an observation. The same code has been added to the ObservationValue code system.

5 | RLD Deprecated | right lateral decubitus | Lying on the right side. Deprecation Comment: This concept has been deprecated because it does not describe a type of Act (as it should in the ActClass code system), but rather encodes the result or value of an observation. The same code has been added to the ObservationValue code system.

5 | SFWL Deprecated | Semi-Fowler's | A semi-sitting position in bed with the head of the bed elevated approximately 45 degrees. Deprecation Comment: This concept has been deprecated because it does not describe a type of Act (as it should in the ActClass code system), but rather encodes the result or value of an observation. The same code has been added to the ObservationValue code system.

5 | SIT Deprecated | sitting | Resting the body on the buttocks, typically with upper torso erect or semi erect. Deprecation Comment: This concept has been deprecated because it does not describe a type of Act (as it should in the ActClass code system), but rather encodes the result or value of an observation. The same code has been added to the ObservationValue code system.

5 | STN Deprecated | standing | To be stationary, upright, vertical, on one's legs. Deprecation Comment: This concept has been deprecated because it does not describe a type of Act (as it should in the ActClass code system), but rather encodes the result or value of an observation. The same code has been added to the ObservationValue code system.

5 | SUP Deprecated | supine | Deprecation Comment: This concept has been deprecated because it does not describe a type of Act (as it should in the ActClass code system), but rather encodes the result or value of an observation. The same code has been added to the ObservationValue code system.

6 | RTRD Deprecated | reverse trendelenburg | Lying on the back, on an inclined plane, typically about 30-45 degrees with head raised and feet lowered. Deprecation Comment: This concept has been deprecated because it does not describe a type of Act (as it should in the ActClass code system), but rather encodes the result or value of an observation. The same code has been added to the ObservationValue code system.

6 | TRD Deprecated | trendelenburg | Lying on the back, on an inclined plane, typically about 30-45 degrees, with head lowered and feet raised. Deprecation Comment: This concept has been deprecated because it does not describe a type of Act (as it should in the ActClass code system), but rather encodes the result or value of an observation. The same code has been added to the ObservationValue code system.

3 | ALRT | detected issue | An observation identifying a potential adverse outcome as a result of an Act or combination of Acts. Examples: Detection of a drug-drug interaction; Identification of a late-submission for an invoice; Requesting discharge for a patient who does not meet hospital-defined discharge criteria. Discussion: This class is commonly used for identifying 'business rule' or 'process' problems that may result in a refusal to carry out a particular request. In some circumstances it may be possible to 'bypass' a problem by modifying the request to acknowledge the issue and/or by providing some form of mitigation. Constraints: the Act or Acts that may cause the the adverse outcome are the target of a subject ActRelationship. The subbtypes of this concept indicate the type of problem being detected (e.g. drug-drug interaction) while the Observation.value is used to repesent a specific problem code (e.g. specific drug-drug interaction id).

3 | BATTERY | battery | Definition: An observation that is composed of a set of observations. These observations typically have a logical or practical grouping for generally accepted clinical or functional purposes, such as observations that are run together because of automation. A battery can define required and optional component observations and, in some cases, will define complex rules that determine whether or not a particular observation is made. BATTERY is a constraint on the Observation class in that it is understood to always be composed of component observations. UsageNotes: The focus in a BATTERY is that it is composed of individual observations. In request (RQO) mood, a battery is a request to perform the component observations. In event (EVN) mood a battery is a reporting of associated set of observation events. In definition mood a battery is the definition of the associated set of observations. Examples: Vital signs, Full blood count, Chemistry panel.

3 | CLNTRL | clinical trial | The set of actions that define an experiment to assess the effectiveness and/or safety of a biopharmaceutical product (food, drug, device, etc.). In definition mood, this set of actions is often embodied in a clinical trial protocol; in event mood, this designates the aggregate act of applying the actions to one or more subjects.

3 | CNOD Deprecated | Condition Node | An instance of Observation of a Condition at a point in time that includes any Observations or Procedures associated with that Condition as well as links to previous instances of Condition Node for the same Condition Deprecation Comment: This concept has been deprecated because an alternative structure for tracking the evolution of a problem has been presented and adopted by the Care Provision Work Group.

3 | COND Deprecated | Condition | An observable finding or state that persists over time and tends to require intervention or management, and, therefore, distinguished from an Observation made at a point in time; may exist before an Observation of the Condition is made or after interventions to manage the Condition are undertaken. Examples: equipment repair status, device recall status, a health risk, a financial risk, public health risk, pregnancy, health maintenance, chronic illness

4 | CASE Deprecated | public health case | A public health case is an Observation representing a condition or event that has a specific significance for public health. Typically it involves an instance or instances of a reportable infectious disease or other condition. The public health case can include a health-related event concerning a single individual or it may refer to multiple health-related events that are occurrences of the same disease or condition of interest to public health. An outbreak involving multiple individuals may be considered as a type of public health case. A public health case definition (Act.moodCode =

"definition") includes the description of the clinical, laboratory, and epidemiologic indicators associated with a disease or condition of interest to public health. There are case definitions for conditions that are reportable, as well as for those that are not. There are also case definitions for outbreaks. A public health case definition is a construct used by public health for the purpose of counting cases, and should not be used as clinical indications for treatment. Examples include AIDS, toxic-shock syndrome, and salmonellosis and their associated indicators that are used to define a case.
5 | OUTB Deprecated | outbreak | An outbreak represents a series of public health cases. The date on which an outbreak starts is the earliest date of onset among the cases assigned to the outbreak, and its ending date is the last date of onset among the cases assigned to the outbreak.
3 | DGIMG | diagnostic image | Class for holding attributes unique to diagnostic images.
3 | GEN | genomic observation | Description:An observation of genomic phenomena.
4 | DETPOL | determinant peptide | Description:A determinant peptide in a polypeptide as described by polypeptide.
4 | EXP | expression level | Description:An expression level of genes/proteins or other expressed genomic entities.
4 | LOC | locus | Description:The position of a gene (or other significant sequence) on the genome.
4 | PHN | phenotype | Description:A genomic phenomenon that is expressed externally in the organism.
4 | POL | polypeptide | Description:A polypeptide resulting from the translation of a gene.
4 | SEQ | bio sequence | Description:A sequence of biomolecule like the DNA, RNA, protein and the like.
4 | SEQVAR | bio sequence variation | Description:A variation in a sequence as described by BioSequence.
3 | INVSTG | investigation | An formalized inquiry into the circumstances surrounding a particular unplanned event or potential event for the purposes of identifying possible causes and contributing factors for the event. This investigation could be conducted at a local institutional level or at the level of a local or national government.
3 | OBSSER | observation series | Container for Correlated Observation Sequences sharing a common frame of reference. All Observations of the same cd must be comparable and relative to the common frame of reference. For example, a 3-channel ECG device records a 12-lead ECG in 4 steps (3 leads at a time). Each of the separate 3-channel recordings would be in their own "OBSCOR". And, all 4 OBSCOR would be contained in one OBSSER because all the times are relative to the same origin (beginning of the recording) and all the ECG signals were from a fixed set of electrodes.
4 | OBSCOR | correlated observation sequences | Container for Observation Sequences (Observations whose values are contained in LIST<>'s) having values correlated with each other. Each contained Observation Sequence LIST<> must be the same length. Values in the LIST<>'s are correlated based on index. E.g. the values in position 2 in all the LIST<>'s are correlated. This is analogous to a table where each column is an Observation Sequence with a LIST<> of values, and each row in the table is a correlation between the columns. For example, a 12-lead ECG would contain 13 sequences: one sequence for time, and a sequence for each of the 12 leads.
3 | POS | position | An observation denoting the physical location of a person or thing based on a reference coordinate system.
4 | POSACC | position accuracy | Description:An observation representing the degree to which the assignment of the spatial coordinates, based on a matching algorithm by a geocoding engine against a reference spatial database, matches true or accepted values.
4 | POSCOORD | position coordinate | Description:An observation representing one of a set of numerical values used to determine the position of a place. The name of the coordinate value is determined by the reference coordinate system.
3 | SPCOBS | specimen observation | An observation on a specimen in a laboratory environment that may affect processing, analysis or result interpretation
3 | VERIF | Verification | An act which describes the process whereby a 'verifying party' validates either the existence of the Role attested to by some Credential or the actual Vetting act and its details.
2 | PCPR | care provision | An Act that of taking on whole or partial responsibility for, or attention to, safety and well-being of a subject of care. Discussion: A care provision event may exist without any other care actions taking place. For example, when a patient is assigned to the care of a particular health professional. In request (RQO) mood care provision communicates a referral, which is a request: from one party (linked as a participant of type author (AUT)), to another party (linked as a participant of type performer (PRF), to take responsibility for a scope specified by the code attribute, for an entity (linked as a participant of type subject (SBJ)). The scope of the care for which responsibility is taken is identified by code attribute. In event (EVN) mood care provision indicates the effective time interval of a specified scope of responsibility by a performer (PRF) or set of performers (PRF) for a subject (SBJ). Examples: Referral from GP to a specialist. Assignment of a patient or group of patients to the case list of a health professional. Assignment of inpatients to the care of particular nurses for a working shift.
3 | ENC | encounter | An interaction between a patient and healthcare participant(s) for the purpose of providing patient service(s) or assessing the health status of a patient. For example, outpatient visit to multiple departments, home health support (including physical therapy), inpatient hospital stay, emergency room visit, field visit (e.g., traffic accident), office visit, occupational therapy, telephone call.
2 | POLICY | policy | Description:A mandate, regulation, obligation, requirement, rule, or expectation unilaterally imposed by one party on: The activity of another party The behavior of another party The manner in which an act is executed
3 | JURISPOL | jurisdictional policy | Description:A mandate, regulation, obligation, requirement, rule, or expectation unilaterally imposed by a jurisdiction on: The activity of another party The behavior of another party The manner in which an act is executed Examples:A jurisdictional mandate regarding the prescribing and dispensing of a particular medication. A jurisdictional privacy or security regulation dictating the manner in which personal health information is disclosed. A jurisdictional requirement that certain services or health conditions are reported to a monitoring program, e.g., immunizations, methadone treatment, or cancer registries.
3 | ORGPOL | organizational policy | Description:A mandate, obligation, requirement, rule, or expectation unilaterally imposed by an organization on: The activity of another party The behavior of another party The manner in which an act is executed Examples:A clinical or research protocols imposed by a payer, a malpractice insurer, or an institution to which a provider must adhere. A mandate imposed by a denominational institution for a provider to provide or withhold certain information from the patient about treatment options.
3 | SCOPOL | scope of practice policy | Description:An ethical or clinical obligation, requirement, rule, or expectation imposed or strongly encouraged by organizations that oversee particular clinical domains or provider certification which define the boundaries within which a provider may practice and which may have legal basis or ramifications on: The activity of another party The behavior of another party The manner in which an act is executed Examples:An ethical obligation for a provider to fully inform a patient about all treatment options. An ethical obligation for a provider not to disclose personal health information that meets certain criteria, e.g., where disclosure might result in harm to the patient or another person. The set of health care services which a provider is credentialed or privileged to provide.
3 | STDPOL | standard of practice policy | Description:A requirement, rule, or expectation typically documented as guidelines, protocols, or formularies imposed or strongly encouraged by an organization that oversees or has authority over the practices within a domain, and which may have legal basis or ramifications on: The activity of another party The behavior of another party

The manner in which an act is executed Examples:A payer may require a prescribing provider to adhere to formulary guidelines. An institution may adopt clinical guidelines and protocols and implement these within its electronic health record and decision support systems.

2 | PROC | procedure | An Act whose immediate and primary outcome (post-condition) is the alteration of the physical condition of the subject. Examples: : Procedures may involve the disruption of some body surface (e.g. an incision in a surgical procedure), but they also include conservative procedures such as reduction of a luxated join, chiropractic treatment, massage, balneotherapy, acupuncture, shiatsu, etc. Outside of clinical medicine, procedures may be such things as alteration of environments (e.g. straightening rivers, draining swamps, building dams) or the repair or change of machinery etc.

3 | SBADM | substance administration | The act of introducing or otherwise applying a substance to the subject. Discussion: The effect of the substance is typically established on a biochemical basis, however, that is not a requirement. For example, radiotherapy can largely be described in the same way, especially if it is a systemic therapy such as radio-iodine. This class also includes the application of chemical treatments to an area. Examples: Chemotherapy protocol; Drug prescription; Vaccination record

3 | SBEXT | Substance Extraction | Description: The act of removing a substance from the subject.

4 | SPECCOLLECT | Specimen Collection | A procedure for obtaining a specimen from a source entity.

2 | REG | registration | Represents the act of maintaining information about the registration of its associated registered subject. The subject can be either an Act or a Role, and includes subjects such as lab exam definitions, drug protocol definitions, prescriptions, persons, patients, practitioners, and equipment. The registration may have a unique identifier - separate from the unique identification of the subject - as well as a core set of related participations and act relationships that characterize the registration event and aid in the disposition of the subject information by a receiving system.

2 | REV | review | The act of examining and evaluating the subject, usually another act. For example, "This prescription needs to be reviewed in 2 months."

2 | SPCTRT | specimen treatment | A procedure or treatment performed on a specimen to prepare it for analysis

2 | SPLY | supply | Supply orders and deliveries are simple Acts that focus on the delivered product. The product is associated with the Supply Act via Participation.typeCode="product". With general Supply Acts, the precise identification of the Material (manufacturer, serial numbers, etc.) is important. Most of the detailed information about the Supply should be represented using the Material class. If delivery needs to be scheduled, tracked, and billed separately, one can associate a Transportation Act with the Supply Act. Pharmacy dispense services are represented as Supply Acts, associated with a SubstanceAdministration Act. The SubstanceAdministration class represents the administration of medication, while dispensing is supply.

3 | DIET Deprecated | diet | Diet services are supply services, with some aspects resembling Medication services: the detail of the diet is given as a description of the Material associated via Participation.typeCode="product". Medically relevant diet types may be communicated in the Diet.code attribute using domain ActDietCode, however, the detail of the food supplied and the various combinations of dishes should be communicated as Material instances. Deprecation Note Class: Use either the Supply class (if dealing with what should be given to the patient) or SubstanceAdministration class (if dealing with what the patient should consume) energyQuantity: This quantity can be conveyed by using a Content relationship with a quantity attribute expressing the calories carbohydrateQuantity:This quantity can be conveyed using a Content relationship to an Entity with a code of carbohydrate and a quantity attribute on the content relationship.

2 | STORE | storage | The act of putting something away for safe keeping. The "something" may be physical object such as a specimen, or information, such as observations regarding a specimen.

2 | SUBST | Substitution | Definition: Indicates that the subject Act has undergone or should undergo substitution of a type indicated by Act.code. Rationale: Used to specify "allowed" substitution when creating orders, "actual" susbstitution when sending events, as well as the reason for the substitution and who was responsible for it.

2 | TRFR | transfer | Definition: The act of transferring information without the intent of imparting understanding about a topic to the subject that is the recipient or holder of the transferred information where the participation association must be RCV or HLD.

2 | TRNS | transportation | Transportation is the moving of a payload (people or material) from a location of origin to a destination location. Thus, any transport service has the three target instances of type payload, origin, and destination, besides the targets that are generally used for any service (i.e., performer, device, etc.)

2 | XACT | financial transaction | A sub-class of Act representing any transaction between two accounts whose value is measured in monetary terms. In the "intent" mood, communicates a request for a transaction to be initiated, or communicates a transfer of value between two accounts. In the "event" mood, communicates the posting of a transaction to an account.

2 | (_ActClassContainer) Abstract

3 | ENTRY | entry | This context represents the information acquired and recorded for an observation, a clinical statement such as a portion of the patient's history or an inference or assertion, or an action that might be intended or has actually been performed. This class may represent both the actual data describing the observation, inference, or action, and optionally the details supporting the clinical reasoning process such as a reference to an electronic guideline, decision support system, or other knowledge reference.

3 | ORGANIZER | organizer | Organizer of entries. Navigational. No semantic content. Knowledge of the section code is not required to interpret contained observations. Represents a heading in a heading structure, or "organizer tree". The record entries relating to a single clinical session are usually grouped under headings that represent phases of the encounter, or assist with layout and navigation. Clinical headings usually reflect the clinical workflow during a care session, and might also reflect the main author's reasoning processes. Much research has demonstrated that headings are used differently by different professional groups and specialties, and that headings are not used consistently enough to support safe automatic processing of the E H R.

1 | DOCCNTNT

1 | DOCLIST

1 | DOCLSTITM

1 | DOCPARA

1 | DOCTBL

1 | LINKHTML

1 | LOCALATTR

1 | LOCALMRKP

1 | ordered

1 | REFR

1 | TBLCOL

1 | TBLCOLGP

1 | TBLDATA

1 | TBLHDR

1 | TBLROW
1 | tbody
1 | tfoot
1 | thead
1 | unordered

4.4.2.7 HL7 v3 Code System ActPriority

URL: https://hl7.org/fhir/R4/v3/ActPriority/cs.html
Vocabulary Work Group | Maturity Level: 1 | Draft | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-ActPriority) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-ActPriority
Version: | 2018-08-12
Name: | v3.ActPriority
Title: | v3 Code System ActPriority
Definition: | A set of codes (e.g., for routine, emergency), specifying the urgency under which the Act happened, can happen, is happening, is intended to happen, or is requested/demanded to happen.
OID: | 2.16.840.1.113883.5.7 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System ActPriority (A set of codes (e.g., for routine, emergency), specifying the urgency under which the Act happened, can happen, is happening, is intended to happen, or is requested/demanded to happen.)

4.4.2.7.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | A | ASAP | As soon as possible, next highest priority after stat.
1 | CR | callback results | Filler should contact the placer as soon as results are available, even for preliminary results. (Was "C" in HL7 version 2.3's reporting priority.)
1 | CS | callback for scheduling | Filler should contact the placer (or target) to schedule the service. (Was "C" in HL7 version 2.3's TQ-priority component.)
2 | CSP | callback placer for scheduling | Filler should contact the placer to schedule the service. (Was "C" in HL7 version 2.3's TQ-priority component.)
2 | CSR | contact recipient for scheduling | Filler should contact the service recipient (target) to schedule the service. (Was "C" in HL7 version 2.3's TQ-priority component.)
1 | EL | elective | Beneficial to the patient but not essential for survival.
1 | EM | emergency | An unforeseen combination of circumstances or the resulting state that calls for immediate action.
1 | P | preop | Used to indicate that a service is to be performed prior to a scheduled surgery. When ordering a service and using the pre-op priority, a check is done to see the amount of time that must be allowed for performance of the service. When the order is placed, a message can be generated indicating the time needed for the service so that it is not ordered in conflict with a scheduled operation.
1 | PRN | as needed | An "as needed" order should be accompanied by a description of what constitutes a need. This description is represented by an observation service predicate as a precondition.
1 | R | routine | Routine service, do at usual work hours.
1 | RR | rush reporting | A report should be prepared and sent as quickly as possible.
1 | S | stat | With highest priority (e.g., emergency).
1 | T | timing critical | It is critical to come as close as possible to the requested time (e.g., for a through antimicrobial level).
1 | UD | use as directed | Drug is to be used as directed by the prescriber.
1 | UR | urgent | Calls for prompt action.

4.4.2.8 HL7 v3 Code System ActReason

URL: https://hl7.org/fhir/R4/v3/ActReason/cs.html
Vocabulary Work Group | Maturity Level: 1 | Draft | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-ActReason) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-ActReason
Version: | 2018-08-12
Name: | v3.ActReason
Title: | v3 Code System ActReason
Definition: | A set of codes specifying the motivation, cause, or rationale of an Act, when such rationale is not reasonably represented as an ActRelationship of type "has reason" linking to another Act. Examples: Example reasons that might qualify for being coded in this field might be: "routine requirement", "infectious disease reporting requirement", "on patient request", "required by law".
OID: | 2.16.840.1.113883.5.8 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: V3 Value SetSubstanceAdminSubstitutionReason (No Description Provided)

• ValueSet: Immunization Status Reason Codes (The value set to instantiate this attribute should be drawn from a terminologically robust code system that consists of or contains concepts to support describing the reason why a dose of vaccine was not administered. This value set is provided as a suggestive example.)

• ValueSet: V3 Value SetGeneralPurposeOfUse (Supports communication of purpose of use at a general level.)

• ValueSet: Immunization Status Reason Codes (The value set to instantiate this attribute should be drawn from a terminologically robust code system that consists of or contains concepts to support describing the reason why a dose of vaccine was not administered. This value set is provided as a suggestive example.)

• ValueSet: v3 Code System ActReason (A set of codes specifying the motivation, cause, or rationale of an Act, when such rationale is not reasonably represented as an ActRelationship of type "has reason" linking to another Act. Examples: Example reasons that might qualify for being coded in this field might be: "routine requirement", "infectious disease reporting requirement", "on patient request", "required by law".)

• ValueSet: V3 Value SetPurposeOfUse (Supports communication of purpose of use at a general level.)

4.4.2.8.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition

1 | (_ActAccommodationReason) Abstract | | Identifies the reason the patient is assigned to this accommodation type

2 | ACCREQNA | Accommodation Requested Not Available | Accommodation requested is not available.

2 | FLRCNV | Floor Convenience | Accommodation is assigned for floor convenience.

2 | MEDNEC | Medical Necessity | Required for medical reasons(s).

2 | PAT | Patient request | The Patient requested the action

1 | (_ActCoverageReason) Abstract | | Description:Codes used to specify reasons or criteria relating to coverage provided under a policy or program. May be used to convey reasons pertaining to coverage contractual provisions, including criteria for eligibility, coverage limitations, coverage maximums, or financial participation required of covered parties.

2 | (_EligibilityActReasonCode) Abstract | | Identifies the reason or rational for why a person is eligibile for benefits under an insurance policy or progam. Examples: A person is a claimant under an automobile insurance policy are client deceased & adopted client has been given a new policy identifier. A new employee is eligible for health insurance as an employment benefit. A person meets a government program eligibility criteria for financial, age or health status.

3 | (_ActIneligibilityReason) Abstract | | Identifies the reason or rational for why a person is not eligibile for benefits under an insurance policy. Examples are client deceased & adopted client has been given a new policy identifier.

4 | COVSUS | coverage suspended | When a client has no contact with the health system for an extended period, coverage is suspended. Client will be reinstated to original start date upon proof of identification, residency etc. Example: Coverage may be suspended during a strike situation, when employer benefits for employees are not covered (i.e. not in effect).

4 | DECSD | deceased | Client deceased.

4 | REGERR | registered in error | Client was registered in error.

3 | (_CoverageEligibilityReason) Abstract | | Definition: Identifies the reason or rational for why a person is eligibile for benefits under an insurance policy or progam. Examples: A person is a claimant under an automobile insurance policy are client deceased & adopted client has been given a new policy identifier. A new employee is eligible for health insurance as an employment benefit. A person meets a government program eligibility criteria for financial, age or health status.

4 | AGE | age eligibility | A person becomes eligible for a program based on age. Example: In the U.S., a person who is 65 years of age or older is eligible for Medicare.

4 | CRIME | crime victim | A person becomes eligible for insurance or a program because of crime related health condition or injury. Example: A person is a claimant under the U.S. Crime Victims Compensation program.

4 | DIS | disability | A person becomes a claimant under a disability income insurance policy or a disability rehabilitation program because of a health condition or injury which limits the person's ability to earn an income or function without institutionalization.

4 | EMPLOY | employment benefit | A person becomes eligible for insurance provided as an employment benefit based on employment status.

4 | FINAN | financial eligibility | A person becomes eligible for a program based on financial criteria. Example: A person whose family income is below a financial threshold for eligibility for Medicaid or SCHIP.

4 | HEALTH | health status | A person becomes eligible for a program because of a qualifying health condition or injury. Examples: A person is determined to have a qualifying health conditions include pregnancy, HIV/AIDs, tuberculosis, end stage renal disease, breast or cervical cancer, or other condition requiring specialized health services, hospice, institutional or community based care provided under a program

4 | MULTI | multiple criteria eligibility | A person becomes eligible for a program based on more than one criterion. Examples: In the U.S., a child whose familiy income meets Medicaid financial thresholds and whose age is less than 18 is eligible for the Early and Periodic Screening, Diagnostic, and Treatment program (EPSDT). A person whose family income meets Medicaid financial thresholds and whose age is 65 years or older is eligible for Medicaid and Medicare, and are referred to as dual eligibles.

4 | PNC | property and casualty condition | A person becomes a claimant under a property and casualty insurance policy because of a related health condition or injury resulting from a circumstance covered under the terms of the policy. Example: A person is a claimant under a homeowners insurance policy because of an injury sustained on the policyholderaTMs premises.

4 | STATUTORY | statutory eligibility | A person becomes eligible for a program based on statutory criteria. Examples: A person is a member of an indigenous group, a veteran of military service, or in the U.S., a recipient of adoption assistance and foster care under Title IV-E of the Social Security.

4 | VEHIC | motor vehicle accident victim | A person becomes a claimant under a motor vehicle accident insurance because of a motor vehicle accident related health condition or injury.

4 | WORK | work related | A person becomes eligible for insurance or a program because of a work related health condition or injury. Example: A person is a claimant under the U.S. Black Lung Program.

1 | (_ActInformationManagementReason) Abstract | | Description:The rationale or purpose for an act relating to information management, such as archiving information for the purpose of complying with an enterprise data retention policy.

2 | (_ActHealthInformationManagementReason) Abstract | | Description:The rationale or purpose for an act relating to health information management, such as archiving information for the purpose of complying with an organization policy or jurisdictional law relating to data retention.

3 | (_ActConsentInformationAccessOverrideReason) Abstract | | To perform one or more operations on information to which the patient has not consented as deemed necessary by authorized entities for providing care in the best interest of the patient; providing immediately needed health care for an emergent condition; or for protecting public or third party safety. Usage Notes: Used to convey the reason that a provider or other entity may or has accessed personal healthcare information. Typically, this involves overriding the subject's consent directives.

4 | OVRER | emergency treatment override | To perform one or more operations on information to which the patient has not consented by authorized entities for treating a condition which poses an immediate threat to the patient's health and which requires immediate medical intervention. Usage Notes: The patient is unable to provide consent, but the provider determines they have an urgent healthcare related reason to access the record.

4 | OVRINCOMP | incompetency override | To perform one or more operations on information to which the patient has not consented because deemed incompetent to provide consent. Usage Note: Maps to v2 CON-16 Subject Competence Indicator (ID) 01791 Definition: Identifies whether the subject was deemed competent to provide consent. Refer to table HL7 Table 0136 - Yes/No Indicator and CON-23 Non-Subject Consenter Reason User-defined Table 0502 - Non-Subject Consenter Reason code NC "Subject is not competent to consent".

4 | OVRPJ | professional judgment override | To perform one or more operations on information to which the patient declined to consent for providing health care. Usage Notes: The patient, while able to give consent, has not. However the provider believes it is in the patient's interest to access the record without patient consent.

4 | OVRPS | public safety override | To perform one or more operations on information to which the patient has not consented for public safety reasons. Usage Notes: The patient, while able to give consent, has not. However, the provider believes that access to masked patient information is justified because of concerns related to public safety.

4 | OVRTPS | third party safety override | To perform one or more operations on information to which the patient has not consented for third party safety. Usage Notes: The patient, while able to give consent, has not. However, the provider believes that access to masked patient information is justified because of concerns related to the health and safety of one or more third parties.

3 | PurposeOfUse | purpose of use | Reason for performing one or more operations on information, which may be permitted by source system's security policy in accordance with one or more privacy policies and consent directives. Usage Notes: The rationale or purpose for an act relating to the management of personal health information, such as collecting personal health information for research or public health purposes.

4 | HMARKT | healthcare marketing | To perform one or more operations on information for marketing services and products related to health care.

4 | HOPERAT | healthcare operations | To perform one or more operations on information used for conducting administrative and contractual activities related to the provision of health care.

5 | CAREMGT | care management | To perform analytics, evaluation and other secondary uses of treatment and healthcare related information to manage the quality, efficacy, patient safety, population health, and cost effectiveness of healthcare delivery. Explicitly excludes the use of information to organize the delivery of health care for care coordination and case management, or to provide healthcare treatment. Usage Note: The concept of care management is narrower than the list of activities related to more general organizational objectives such as provider profiling, education of healthcare and non-healthcare professionals; insurance underwriting, premium rating, reinsurance; organizational legal, medical review, auditing, compliance and fraud and abuse detection; business planning, development, and restructuring; fund-raising; and customer service. Map: Maps to ISO 14265 Classification Term "Health service management and quality assurance" described as "To inform persons or processes responsible for determining the availability, quality, safety, equity and cost-effectiveness of health care services." There is a semantic gap in concepts. This classification term is described as activities, i.e., "to inform persons" or "to inform processes" rather than the rationale for performing actions/operations on information related to the activity.

5 | DONAT | donation | To perform one or more operations on information used for cadaveric organ, eye or tissue donation.

5 | FRAUD | fraud | To perform one or more operations on information used for fraud detection and prevention processes.

5 | GOV | government | To perform one or more operations on information used within government processes.

5 | HACCRED | health accreditation | To perform one or more operations on information for conducting activities related to meeting accreditation criteria.

5 | HCOMPL | health compliance | To perform one or more operations on information used for conducting activities required to meet a mandate.

5 | HDECD | decedent | To perform one or more operations on information used for handling deceased patient matters.

5 | HDIRECT | directory | To perform one or more operation operations on information used to manage a patient directory. Examples: facility enterprise payer health information exchange patient directory

5 | HDM | healthcare delivery management | To perform one or more actions on information used for conducting administrative and contractual activities by or on behalf of organizational entities responsible for delivery of an individual's benefits in a healthcare program, health plan or insurance. Explicitly excludes the use of information to organize the delivery of health care for care coordination and case management, or to provide healthcare treatment. Usage Note: Examples of activities conducted under this purpose of use: provider profiling, risk adjustment, underwriting, fraud and abuse, quality improvement population health and care management. Aligns with HIPAA Operation POU minus coordination of care or other treatment related activities. Similar to the description in SAMHSA Confidentiality of Substance Use Disorder Patient Records Supplemental notice of proposed rulemaking. Map: Maps to ISO 14265 Classification Term "Administration of care for an individual subject of care" described as "To inform persons or processes responsible for enabling the availability of resources or funding or permissions for providing health care services to the subject of care." However, this classification term is described as activities, i.e., "to inform persons" or "to inform processes" rather than the rationale for performing actions/operations on information related to the activity.

5 | HLEGAL | legal | To perform one or more operations on information for conducting activities required by legal proceeding.

5 | HOUTCOMS | health outcome measure | To perform one or more operations on information used for assessing results and comparative effectiveness achieved by health care practices and interventions.

5 | HPRGRP | health program reporting | To perform one or more operations on information used for conducting activities to meet program accounting requirements.

5 | HQUALIMP | health quality improvement | To perform one or more operations on information used for conducting administrative activities to improve health care quality.

5 | HSYSADMIN | health system administration | To perform one or more operations on information to administer the electronic systems used for the delivery of health care.
6 | LABELING | labeling | To perform one or more operations on information to assign, persist, and manage labels to healthcare data to characterize various aspects, such as its security classification, sensitivity, compartment, integrity, and provenance; applicable privacy, consent, security, provenance, and trust policies; and handling caveats such as purpose of use, obligations, and refrain policies. Label management includes classification of target data by constructing and binding of a label set per applicable policies, security policy information file semantics, and classification guides. Label management also includes process and procedures for subsequent revision of a label for, e.g., reclassification, downgrading classification, and declassification. Label revisions may be triggered by, e.g., expiry of classification period; changes in applicable policy, e.g., revocation of a consent directive; or changes in the governing policy domain in which the data is relocated or a copy of the data is sent. If a label is revised, an audit log should be kept and the provenance of the label changes should be tracked.
6 | METAMGT | metadata management | To perform one or more operations on information to assign, persist, and manage metadata to healthcare data to characterize various aspects used for its indexing, discovery, retrieval, and processing by systems, applications, and end users. For example, master index identifier, media type, and location.
5 | MEMADMIN | member administration | To perform one or more operations on information to administer health care coverage to an enrollee under a policy or program.
5 | MILCDM | military command | To perform one or more operations on information for conducting activities required by military processes, procedures, policies, or law.
5 | PATADMIN | patient administration | To perform one or more operations on information used for operational activities conducted to administer the delivery of health care to a patient.
5 | PATSFTY | patient safety | To perform one or more operations on information in processes related to ensuring the safety of health care.
5 | PERFMSR | performance measure | To perform one or more operations on information used for monitoring performance of recommended health care practices and interventions.
5 | RECORDMGT | records management | To perform one or more operations on information used within the health records management process.
5 | SYSDEV | system development | To perform one or more operations on information to design, develop, implement, test, or deploy a healthcare system or application.
6 | HTEST | test health data | To perform one or more operations on information that is simulated or synthetic health data used for testing system capabilities outside of a production or operational system environment. Usage Note: Data marked with a HTEST security label enables an access control system to permit interfacing systems or end users provisioned with a clearance, which includes a HTEST purpose of use attribute, to test, verify, or validate that a system or application will operate in production as intended based on design specifications.
5 | TRAIN | training | To perform one or more operations on information used in training and education.
4 | HPAYMT | healthcare payment | To perform one or more operations on information for conducting financial or contractual activities related to payment for provision of health care.
5 | CLMATTCH | claim attachment | To perform one or more operations on information for provision of additional clinical evidence in support of a request for coverage or payment for health services.
5 | COVAUTH | coverage authorization | To perform one or more operations on information for conducting prior authorization or predetermination of coverage for services.
5 | COVERAGE | coverage under policy or program | To perform one or more operations on information for conducting activities related to coverage under a program or policy.
6 | ELIGDTRM | eligibility determination | To perform one or more operations on information used for conducting eligibility determination for coverage in a program or policy. May entail review of financial status or disability assessment.
6 | ELIGVER | eligibility verification | To perform one or more operations on information used for conducting eligibility verification of coverage in a program or policy. May entail provider contacting coverage source (e.g., government health program such as workers compensation or health plan) for confirmation of enrollment, eligibility for specific services, and any applicable copays.
6 | ENROLLM | enrollment | To perform one or more operations on information used for enrolling a covered party in a program or policy. May entail recording of covered party's and any dependent's demographic information and benefit choices.
6 | MILDCRG | military discharge | To perform one or more operations on information for the process of releasing military personnel from their service obligations, which may include determining service merit, discharge benefits, and disability assessment.
5 | REMITADV | remittance advice | To perform one or more operations on information about the amount remitted for a health care claim.
4 | HRESCH | healthcare research | To perform one or more operations on information for conducting scientific investigations to obtain health care knowledge. Use of the data iincludes basic and applied research such as biomedical, population origin or ancestry, translational research, and disease, discipline, specialty specific healthcare research and clinical trial research.
5 | BIORCH | biomedical research | To perform one or more operations on information for conducting scientific investigations to obtain health care knowledge. Use of the data must be related to specified biomedical basic or applied research. For example, research on rare plants to determine whether biologic properties may be useful for pharmaceutical development. May be used in combination with clinical trial and other healthcare research purposes of use.
5 | CLINTRCH | clinical trial research | To perform one or more operations on information for conducting scientific investigations in accordance with clinical trial protocols to obtain health care knowledge.
6 | CLINTRCHNPC | clinical trial research without patient care | To perform one or more operations on information for conducting scientific investigations in accordance with clinical trial protocols to obtain health care knowledge without provision of patient care. May be post-coordinated or used with other purposes of use such as disease, discipline, specialty, population origins or ancestry, translational healthcare research. For example, a clinical trial conducted on laboratory specimens collected from a specified patient population.
6 | CLINTRCHPC | clinical trial research with patient care | To perform one or more operations on information for conducting scientific investigations with patient care in accordance with clinical trial protocols to obtain health care knowledge. May be post-coordinated or used with other purposes of use such as disease, discipline, specialty, population origins or ancestry, translational healthcare research. For example, an "off-label" drug used for cancer therapy administer to a specified patient population.
6 | PRECLINTRCH | preclinical trial research | To perform one or more operations on information in preparation for conducting scientific investigation to obtain health care knowledge, such as research on animals or review of patient health records, to determine the feasibility of a clinical trial study; assist with protocol design; or in preparation for institutional review board or ethics committee approval process. May be post-coordinated or used with other purposes of use such as disease, discipline,

specialty, population origins or ancestry, translational healthcare research.

5 | DSRCH | disease specific healthcare research | To perform one or more operations on information for conducting scientific investigations to obtain health care knowledge. Use of the data must be related to specified conditions, diagnosis, or disease healthcare research. For example, conducting cancer research by testing reaction of tumor cells to certain biologics. May be used in combination with clinical trial and other healthcare research purposes of use.

5 | POARCH | population origins or ancestry healthcare research | To perform one or more operations on information, including genealogical pedigrees, historical records, surveys, family health data, health records, and genetic information, for conducting scientific investigations to obtain health care knowledge. Use of the data must be related to population origins and/or ancestry healthcare research. For example, gathering genetic specimens from a specific population in order to determine the ancestry and population origins of that group. May be used in combination with clinical trial and other healthcare research purposes of use.

5 | TRANSRCH | translational healthcare research | To perform one or more operations on information for conducting scientific investigations to obtain health care knowledge related to evidence based medicine during the course of providing healthcare treatment. Sometimes referred to as "bench to bedside", which is the iterative feedback loop between healthcare research and clinical trials with input from information collected in the course of routine provision of healthcare. For example, by extending a patient encounter to conduct a survey related to a research topic such as attitudes about use of a wellness device that a patient agreed to use. May be used in combination with clinical trial and other healthcare research purposes of use.

4 | PATRQT | patient requested | To perform one or more operations on information in response to a patient's request.

5 | FAMRQT | family requested | To perform one or more operations on information in response to a request by a family member authorized by the patient.

5 | PWATRNY | power of attorney | To perform one or more operations on information in response to a request by a person appointed as the patient's legal representative.

5 | SUPNWK | support network | To perform one or more operations on information in response to a request by a person authorized by the patient.

4 | PUBHLTH | public health | To perform one or more operations on information for conducting public health activities, such as the reporting of notifiable conditions.

5 | DISASTER | disaster | To perform one or more operations on information used for provision of immediately needed health care to a population of living subjects located in a disaster zone.

5 | THREAT | threat | To perform one or more operations on information used to prevent injury or disease to living subjects who may be the target of violence.

4 | TREAT | treatment | To perform one or more operations on information for provision of health care.

5 | CLINTRL | clinical trial | To perform health care as part of the clinical trial protocol.

5 | COC | coordination of care | To perform one or more actions on information in order to organize the provision and case management of an individualâ€™s healthcare, including: Monitoring a person's goals, needs, and preferences; acting as the communication link between two or more participants concerned with a person's health and wellness; organizing and facilitating care activities and promoting self-management by advocating for, empowering, and educating a person; and ensuring safe, appropriate, non-duplicative, and effective integrated care. Usage Note: Use when describing these functions: 1. Monitoring a personâ€™s goals, needs, and preferences. 2. Acting as the communication link between two or more participants concerned with a person's health and wellness. 3. Organizing and facilitating care activities and promoting self-management by advocating for, empowering, and educating a person. 4. Ensuring safe, appropriate, non-duplicative, and effective integrated care. The goal is to clearly differentiate this type of coordination of care from HIPAA Operations by specifying that these actions on information are undertaken in the provision of healthcare treatment. For similar uses of this concept, see SAMHSA Confidentiality of Substance Use Disorder Patient Records Supplemental notice of proposed rulemaking, which differentiates concepts of care coordination and case management for the provision of treatment as specifically distinct from activities related to health care delivery management and the operations of organizational entities involved in the delivery of healthcare. Map: Maps to ISO 14265 Classification Terms: "Support of care activities within the provider organisation for an individual subject of care" described as "To inform persons or processes enabling others to provide health care services to the subject of care." "Subject of Care Uses" described as "To inform the subject of care in support of his or her own interests."

5 | ETREAT | Emergency Treatment | To perform one or more operations on information for provision of immediately needed health care for an emergent condition.

6 | BTG | break the glass | To perform policy override operations on information for provision of immediately needed health care for an emergent condition affecting potential harm, death or patient safety by end users who are not provisioned for this purpose of use. Includes override of organizational provisioning policies and may include override of subject of care consent directive restricting access. Map: Partially Maps to ISO 14265 Classification Term "Emergency care provision to an individual subject of care" described as "To inform persons needing to provide health care services to the subject of care urgently, possibly needing to over-ride the policies and consents pertaining to Purpose 1 above." Purpose 1 is equivalent to HL7 treatment purpose of use: "Clinical care provision to an individual subject of care" described as "To inform persons or processes responsible for providing health care services to the subject of care." The ISO description conflates both of the proposed specializations of HL7 ETREAT: break the glass and the typically broader access to health information normally available to providers who are provisioned for emergency workflows on a regular basis, e.g., Emergency Room providers. Examples of greater access than is normally accessible by providers based on the need to know are access to sensitive information for which access typically requires a patient's consent. This is not an override of a patient's dissent to disclose sensitive information in cases where the applicable policy waives the need for that consent to access this information. In US, Title 38 Section 7332 and 42 CFR Part 2 both permit emergency access without the need to override a patient's consent directive; rather, this access is a limitation to the patient's right to dissent from disclosure.

6 | ERTREAT | emergency room treatment | To perform one or more operations on information for provision of immediately needed health care for an emergent condition in an emergency room or similar emergent care context by end users provisioned for this purpose, which does not constitute as policy override such as in a "Break the Glass" purpose of use. Map:Partially Maps to ISO 14265 Classification Term "Emergency care provision to an individual subject of care" described as "To inform persons needing to provide health care services to the subject of care urgently, possibly needing to over-ride the policies and consents pertaining to Purpose 1 above." Purpose 1 is equivalent to HL7 treatment purpose of use: "Clinical care provision to an individual subject of care" described as "To inform persons or processes responsible for providing health care services to the subject of care." The ISO description conflates both of the proposed specializations of HL7 ETREAT: break the glass and the typically broader access to health information normally available to providers who are provisioned for emergency workflows on a regular basis, e.g., Emergency Room providers. Examples of greater access than is normally accessible by providers based on the need to know are access to sensitive information for which access typically requires a patient's consent. This is not an override of a patient's dissent to disclose sensitive information in cases where the applicable policy waives the need for that consent to access this information. In US, Title 38 Section 7332 and 42 CFR Part 2 both permit emergency access without the need to override a patient's consent directive; rather, this access is a limitation to the patient's right to dissent from disclosure. There is a semantic gap in concepts.

This classification term is described as activities â€œto inform personsâ€? rather than the rationale for performing actions/operations on information related to the activity.
5 | POPHLTH | population health | To perform one or more operations on information for provision of health care to a population of living subjects, e.g., needle exchange program.
2 | (_ActInformationPrivacyReason) Abstract | | Description:The rationale or purpose for an act relating to the management of personal information, such as disclosing personal tax information for the purpose of complying with a court order.
3 | MARKT | marketing | Description:
3 | OPERAT | operations | Description:Administrative and contractual processes required to support an activity, product, or service
4 | LEGAL | subpoena | Definition:To provide information as a result of a subpoena.
4 | ACCRED | accreditation | Description:Operational activities conducted for the purposes of meeting of criteria defined by an accrediting entity for an activity, product, or service
4 | COMPL | compliance | Description:Operational activities required to meet a mandate related to an activity, product, or service
4 | ENADMIN | entity administration | Description:Operational activities conducted to administer information relating to entities involves with an activity, product, or service
4 | OUTCOMS | outcome measure | Description:Operational activities conducted for the purposes of assessing the results of an activity, product, or service
4 | PRGRPT | program reporting | Description:Operational activities conducted to meet program accounting requirements related to an activity, product, or service
4 | QUALIMP | quality improvement | Description:Operational activities conducted for the purposes of improving the quality of an activity, product, or service
4 | SYSADMN | system administration | Description:Operational activities conducted to administer the electronic systems used for an activity, product, or service
3 | PAYMT | payment | Description:Administrative, financial, and contractual processes related to payment for an activity, product, or service
3 | RESCH | research | Description:Investigative activities conducted for the purposes of obtaining knowledge
3 | SRVC | service | Description:Provision of a service, product, or capability to an individual or organization
1 | _ActInvalidReason | ActInvalidReason | Description: Types of reasons why a substance is invalid for use.
2 | ADVSTORAGE | adverse storage condition | Description: Storage conditions caused the substance to be ineffective.
3 | COLDCHNBRK | cold chain break | Description: Cold chain was not maintained for the substance.
2 | EXPLOT | expired lot | Description: The lot from which the substance was drawn was expired.
2 | OUTSIDESCHED | administered outside recommended schedule or practice | The substance was administered outside of the recommended schedule or practice.
2 | PRODRECALL | product recall | Description: The substance was recalled by the manufacturer.
1 | (_ActInvoiceCancelReason) Abstract | | Domain specifies the codes used to describe reasons why a Provider is cancelling an Invoice or Invoice Grouping.
2 | INCCOVPTY | incorrect covered party as patient | The covered party (patient) specified with the Invoice is not correct.
2 | INCINVOICE | incorrect billing | The billing information, specified in the Invoice Elements, is not correct. This could include incorrect costing for items included in the Invoice.
2 | INCPOLICY | incorrect policy | The policy specified with the Invoice is not correct. For example, it may belong to another Adjudicator or Covered Party.
2 | INCPROV | incorrect provider | The provider specified with the Invoice is not correct.
1 | (_ActNoImmunizationReason) Abstract | | A coded description of the reason for why a patient did not receive a scheduled immunization. (important for public health strategy
2 | IMMUNE | immunity | Definition:Testing has shown that the patient already has immunity to the agent targeted by the immunization.
2 | MEDPREC | medical precaution | Definition:The patient currently has a medical condition for which the vaccine is contraindicated or for which precaution is warranted.
2 | OSTOCK | product out of stock | Definition:There was no supply of the product on hand to perform the service.
2 | PATOBJ | patient objection | Definition:The patient or their guardian objects to receiving the vaccine.
2 | PHILISOP | philosophical objection | Definition:The patient or their guardian objects to receiving the vaccine because of philosophical beliefs.
2 | RELIG | religious objection | Definition:The patient or their guardian objects to receiving the vaccine on religious grounds.
2 | VACEFF | vaccine efficacy concerns | Definition:The intended vaccine has expired or is otherwise believed to no longer be effective. Example:Due to temperature exposure.
2 | VACSAF | vaccine safety concerns | Definition:The patient or their guardian objects to receiving the vaccine because of concerns over its safety.
1 | (_ActSupplyFulfillmentRefusalReason) Abstract | | Indicates why a fulfiller refused to fulfill a supply order, and considered it important to notify other providers of their decision. E.g. "Suspect fraud", "Possible abuse", "Contraindicated". (used when capturing 'refusal to fill' annotations)
2 | FRR01 | order stopped | Definition:The order has been stopped by the prescriber but this fact has not necessarily captured electronically. Example:A verbal stop, a fax, etc.
2 | FRR02 | stale-dated order | Definition:Order has not been fulfilled within a reasonable amount of time, and may not be current.
2 | FRR03 | incomplete data | Definition:Data needed to safely act on the order which was expected to become available independent of the order is not yet available Example:Lab results, diagnostic imaging, etc.
2 | FRR04 | product unavailable | Definition:Product not available or manufactured. Cannot supply.
2 | FRR05 | ethical/religious | Definition:The dispenser has ethical, religious or moral objections to fulfilling the order/dispensing the product.
2 | FRR06 | unable to provide care | Definition:Fulfiller not able to provide appropriate care associated with fulfilling the order. Example:Therapy requires ongoing monitoring by fulfiller and fulfiller will be ending practice, leaving town, unable to schedule necessary time, etc.
1 | (_ClinicalResearchEventReason) Abstract | | Definition:Specifies the reason that an event occurred in a clinical research study.
2 | RET | retest | Definition:The event occurred so that a test or observation performed at a prior event could be performed again due to conditions set forth in the protocol.
2 | SCH | scheduled | Definition:The event occurred due to it being scheduled in the research protocol.
2 | TRM | termination | Definition:The event occurred in order to terminate the subject's participation in the study.

2 | UNS | unscheduled | Definition:The event that occurred was initiated by a study participant (e.g. the subject or the investigator), and did not occur for protocol reasons.
1 | (_ClinicalResearchObservationReason) Abstract | | Definition:SSpecifies the reason that a test was performed or observation collected in a clinical research study. Note:This set of codes are not strictly reasons, but are used in the currently Normative standard. Future revisions of the specification will model these as ActRelationships and thes codes may subsequently be retired. Thus, these codes should not be used for new specifications.
2 | NPT | non-protocol | Definition:The observation or test was neither defined or scheduled in the study protocol.
2 | PPT | per protocol | Definition:The observation or test occurred due to it being defined in the research protocol, and during an activity or event that was scheduled in the protocol.
2 | UPT | per definition | :The observation or test occurred as defined in the research protocol, but at a point in time not specified in the study protocol.
1 | (_CombinedPharmacyOrderSuspendReasonCode) Abstract | | Description:Indicates why the prescription should be suspended.
2 | ALTCHOICE | try another treatment first | Description:This therapy has been ordered as a backup to a preferred therapy. This order will be released when and if the preferred therapy is unsuccessful.
2 | CLARIF | prescription requires clarification | Description:Clarification is required before the order can be acted upon.
2 | DRUGHIGH | drug level too high | Description:The current level of the medication in the patient's system is too high. The medication is suspended to allow the level to subside to a safer level.
2 | HOSPADM | admission to hospital | Description:The patient has been admitted to a care facility and their community medications are suspended until hospital discharge.
2 | LABINT | lab interference issues | Description:The therapy would interfere with a planned lab test and the therapy is being withdrawn until the test is completed.
2 | NON-AVAIL | patient not-available | Description:Patient not available for a period of time due to a scheduled therapy, leave of absence or other reason.
2 | PREG | parent is pregnant/breast feeding | Description:The patient is pregnant or breast feeding. The therapy will be resumed when the pregnancy is complete and the patient is no longer breastfeeding.
2 | SALG | allergy | Description:The patient is believed to be allergic to a substance that is part of the therapy and the therapy is being temporarily withdrawn to confirm.
2 | SDDI | drug interacts with another drug | Description:The drug interacts with a short-term treatment that is more urgently required. This order will be resumed when the short-term treatment is complete.
2 | SDUPTHER | duplicate therapy | Description:Another short-term co-occurring therapy fulfills the same purpose as this therapy. This therapy will be resumed when the co-occuring therapy is complete.
2 | SINTOL | suspected intolerance | Description:The patient is believed to have an intolerance to a substance that is part of the therapy and the therapy is being temporarily withdrawn to confirm.
2 | SURG | patient scheduled for surgery | Description:The drug is contraindicated for patients receiving surgery and the patient is scheduled to be admitted for surgery in the near future. The drug will be resumed when the patient has sufficiently recovered from the surgery.
2 | WASHOUT | waiting for old drug to wash out | Description:The patient was previously receiving a medication contraindicated with the current medication. The current medication will remain on hold until the prior medication has been cleansed from their system.
1 | _ControlActNullificationReasonCode | ControlActNullificationReasonCode | Description:Identifies reasons for nullifying (retracting) a particular control act.
2 | ALTD | altered decision | Description:The decision on which the recorded information was based was changed before the decision had an effect. Example:Aborted prescription before patient left office, released prescription before suspend took effect.
2 | EIE | entered in error | Description:The information was recorded incorrectly or was recorded in the wrong record.
2 | NORECMTCH | no record match | Description: There is no match for the record in the database.
1 | (_ControlActNullificationRefusalReasonType) Abstract | | Description: Reasons to refuse a transaction to be undone.
2 | INRQSTATE | in requested state | The record is already in the requested state.
2 | NOMATCH | no match | Description: There is no match.
3 | NORECMTCH
3 | NOPRODMTCH | no product match | Description: There is no match for the product in the master file repository.
3 | NOSERMTCH | no service match | Description: There is no match for the service in the master file repository.
3 | NOVERMTCH | no version match | Description: There is no match for the record and version.
2 | NOPERM | no permission | Description: There is no permission.
3 | NOUSERPERM | no user permission | Definition:The user does not have permission
3 | NOAGNTPERM | no agent permission | Description: The agent does not have permission.
3 | NOUSRPERM | no user permission | Description: The user does not have permission.
2 | WRNGVER | wrong version | Description: The record and version requested to update is not the current version.
1 | (_ControlActReason) Abstract | | Identifies why a specific query, request, or other trigger event occurred.
2 | (_MedicationOrderAbortReasonCode) Abstract | | Description:Indicates the reason the medication order should be aborted.
3 | DISCONT | product discontinued | Description:The medication is no longer being manufactured or is otherwise no longer available.
3 | INEFFECT | ineffective | Description:The therapy has been found to not have the desired therapeutic benefit on the patient.
3 | MONIT | response to monitoring | Description:Monitoring the patient while taking the medication, the decision has been made that the therapy is no longer appropriate.
3 | NOREQ | no longer required for treatment | Description:The underlying condition has been resolved or has evolved such that a different treatment is no longer needed.
3 | NOTCOVER | not covered | Description:The product does not have (or no longer has) coverage under the patientaTMs insurance policy.
3 | PREFUS | patient refuse | Description:The patient refused to take the product.
3 | RECALL | product recalled | Description:The manufacturer or other agency has requested that stocks of a medication be removed from circulation.
3 | REPLACE | change in order | Description:Item in current order is no longer in use as requested and a new one has/will be created to replace it.
4 | DOSECHG | change in medication/dose | Description:The medication is being re-prescribed at a different dosage.

3 | REPLACEFIX | error in order | Description:Current order was issued with incorrect data and a new order has/will be created to replace it.
3 | UNABLE | unable to use | Description:<The patient is not (or is no longer) able to use the medication in a manner prescribed. Example:CanaTMt swallow.
2 | (_MedicationOrderReleaseReasonCode) Abstract | | Definition:A collection of concepts that indicate why the prescription should be released from suspended state.
3 | HOLDDONE | suspend reason no longer applies | Definition:The original reason for suspending the medication has ended.
3 | HOLDINAP | suspend reason inappropriate | Definition:
2 | (_ModifyPrescriptionReasonType) Abstract | | Types of reason why a prescription is being changed.
3 | ADMINERROR | administrative error in order | Order was created with incorrect data and is changed to reflect the intended accuracy of the order.
3 | CLINMOD | clinical modification | Order is changed based on a clinical reason.
2 | _PharmacySupplyEventAbortReason | PharmacySupplyEventAbortReason | Definition:Identifies why the dispense event was not completed.
3 | CONTRA | contraindication | Definition:Contraindication identified
3 | FOABORT | order aborted | Definition:Order to be fulfilled was aborted
3 | FOSUSP | order suspended | Definition:Order to be fulfilled was suspended
3 | NOPICK | not picked up | Definition:Patient did not come to get medication
3 | PATDEC | patient changed mind | Definition:Patient changed their mind regarding obtaining medication
3 | QUANTCHG | change supply quantity | Definition:Patient requested a revised quantity of medication
2 | (_PharmacySupplyEventStockReasonCode) Abstract | | Definition:A collection of concepts that indicates the reason for a "bulk supply" of medication.
3 | FLRSTCK | floor stock | Definition:The bulk supply is issued to replenish a ward for local dispensing. (Includes both mobile and fixed-location ward stocks.)
3 | LTC | long term care use | Definition:The bulk supply will be administered within a long term care facility.
3 | OFFICE | office use | Definition:The bulk supply is intended for general clinician office use.
3 | PHARM | pharmacy transfer | Definition:The bulk supply is being transferred to another dispensing facility to. Example:Alleviate a temporary shortage.
3 | PROG | program use | Definition:The bulk supply is intended for dispensing according to a specific program. Example:Mass immunization.
2 | (_PharmacySupplyRequestRenewalRefusalReasonCode) Abstract | | Definition:A collection of concepts that identifies why a renewal prescription has been refused.
3 | DISCONT
3 | ALREADYRX | new prescription exists | Definition:Patient has already been given a new (renewal) prescription.
3 | FAMPHYS | family physician must authorize further fills | Definition:Request for further authorization must be done through patient's family physician.
3 | MODIFY | modified prescription exists | Definition:Therapy has been changed and new prescription issued
3 | NEEDAPMT | patient must make appointment | Definition:Patient must see prescriber prior to further fills.
3 | NOTAVAIL | prescriber not available | Definition:Original prescriber is no longer available to prescribe and no other prescriber has taken responsibility for the patient.
3 | NOTPAT | patient no longer in this practice | Definition:Patient no longer or has never been under this prescribers care.
3 | ONHOLD | medication on hold | Definition:This medication is on hold.
3 | PRNA | product not available | Description:This product is not available or manufactured.
3 | STOPMED | prescriber stopped medication for patient | Renewing or original prescriber informed patient to stop using the medication.
3 | TOOEARLY | too early | Definition:The patient should have medication remaining.
2 | (_SupplyOrderAbortReasonCode) Abstract | | Definition:A collection of concepts that indicates why the prescription should no longer be allowed to be dispensed (but can still administer what is already being dispensed).
3 | IMPROV | condition improved | Definition:The patient's medical condition has nearly abated.
3 | INTOL | intolerance | Description:The patient has an intolerance to the medication.
3 | NEWSTR | new strength | Definition:The current medication will be replaced by a new strength of the same medication.
3 | NEWTHER | new therapy | Definition:A new therapy will be commenced when current supply exhausted.
1 | (_GenericUpdateReasonCode) Abstract | | Description:Identifies why a change is being made to a record.
2 | CHGDATA | information change | Description:Information has changed since the record was created.
2 | FIXDATA | error correction | Description:Previously recorded information was erroneous and is being corrected.
2 | MDATA | merge data | Information is combined into the record.
2 | NEWDATA | new information | Description:New information has become available to supplement the record.
2 | UMDATA | unmerge data | Information is separated from the record.
1 | (_PatientProfileQueryReasonCode) Abstract | | Definition:A collection of concepts identifying why the patient's profile is being queried.
2 | ADMREV | administrative review | Definition: To evaluate for service authorization, payment, reporting, or performance/outcome measures.
2 | LEGAL
2 | PATCAR | patient care | Definition:To obtain records as part of patient care.
2 | PATREQ | patient request query | Definition:Patient requests information from their profile.
2 | PRCREV | practice review | Definition:To evaluate the provider's current practice for professional-improvement reasons.
2 | REGUL | regulatory review | Description:Review for the purpose of regulatory compliance.
2 | RSRCH | research | Definition:To provide research data, as authorized by the patient.
2 | VALIDATION | validation review | Description:To validate the patient's record. Example:Merging or unmerging records.
1 | _PharmacySupplyRequestFulfillerRevisionRefusalReasonCode | PharmacySupplyRequestFulfillerRevisionRefusalReasonCode | Definition:Indicates why the request to transfer a prescription from one dispensing facility to another has been refused.
2 | LOCKED | locked | Definition:The prescription may not be reassigned from the original pharmacy.
2 | NOUSERPERM
2 | UNKWNTARGET | unknown target | Definition:The target facility does not recognize the dispensing facility.

1 | (_RefusalReasonCode) Abstract | | Description: Identifies why a request to add (or activate) a record is being refused. Examples include the receiving system not able to match the identifier and find that record in the receiving system, having no permission, or a detected issue exists which precludes the requested action.
2 | NOMATCH
2 | NOPERM
1 | (_SchedulingActReason) Abstract | | Reasons for cancelling or rescheduling an Appointment
2 | PAT
2 | BLK | Unexpected Block (of Schedule) | The time slots previously allocated are now blocked and no longer available for booking Appointments
2 | DEC | Patient Deceased | The Patient is deceased
2 | FIN | No Financial Backing | Patient unable to pay and not covered by insurance
2 | MED | Medical Status Altered | The medical condition of the Patient has changed
2 | MTG | In an outside meeting | The Physician is in a meeting. For example, he/she may request administrative time to talk to family after appointment
2 | PHY | Physician request | The Physician requested the action
1 | _StatusRevisionRefusalReasonCode | StatusRevisionRefusalReasonCode | Indicates why the act revision (status update) is being refused.
2 | NORECMTCH
2 | INRQSTATE
2 | NOUSERPERM
2 | FILLED | fully filled | Ordered quantity has already been completely fulfilled.
1 | _SubstanceAdministrationPermissionRefusalReasonCode | SubstanceAdministrationPermissionRefusalReasonCode | Definition:Indicates why the requested authorization to prescribe or dispense a medication has been refused.
2 | NOUSERPERM
2 | PATINELIG | patient not eligible | Definition:Patient not eligible for drug
2 | PROTUNMET | protocol not met | Definition:Patient does not meet required protocol
2 | PROVUNAUTH | provider not authorized | Definition:Provider is not authorized to prescribe or dispense
1 | (_SubstanceAdminSubstitutionNotAllowedReason) Abstract | | Reasons why substitution of a substance administration request is not permitted.
2 | PAT
2 | ALGINT | allergy intolerance | Definition: Patient has had a prior allergic intolerance response to alternate product or one of its components.
2 | COMPCON | compliance concern | Definition: Patient has compliance issues with medication such as differing appearance, flavor, size, shape or consistency.
2 | THERCHAR | therapeutic characteristics | The prescribed product has specific clinical release or other therapeutic characteristics not shared by other substitutable medications.
2 | TRIAL | clinical trial drug | Definition: The specific manufactured drug is part of a clinical trial.
1 | (_SubstanceAdminSubstitutionReason) Abstract
2 | CT | continuing therapy | Indicates that the decision to substitute or to not substitute was driven by a desire to maintain consistency with a pre-existing therapy. I.e. The performer provided the same item/service as had been previously provided rather than providing exactly what was ordered, or rather than substituting with a lower-cost equivalent.
2 | FP | formulary policy | Indicates that the decision to substitute or to not substitute was driven by a policy expressed within the formulary.
2 | OS | out of stock | In the case of 'substitution', indicates that the substitution occurred because the ordered item was not in stock. In the case of 'no substitution', indicates that a cheaper equivalent was not substituted because it was not in stock.
2 | RR | regulatory requirement | Indicates that the decision to substitute or to not substitute was driven by a jurisdictional regulatory requirement mandating or prohibiting substitution.
1 | (_TransferActReason) Abstract | | The explanation for why a patient is moved from one location to another within the organization
2 | ER | Error | Moved to an error in placing the patient in the original location.
2 | RQ | Request | Moved at the request of the patient.
1 | (_ActBillableServiceReason) Abstract | | Definition: This domain is used to document reasons for providing a billable service; the billable services may include both clinical services and social services.
2 | (_ActBillableClinicalServiceReason) Abstract | | Reason for Clinical Service being performed. This domain excludes reasons specified by diagnosed conditions. Examples of values from this domain include duplicate therapy and fraudulent prescription.
1 | BONUS
1 | CHD | Children only | Description:The level of coverage under the policy or program is available only to children
1 | DEP | Dependents only | Description:The level of coverage under the policy or program is available only to a subscriber's dependents.
1 | ECH | Employee and children | Description:The level of coverage under the policy or program is available to an employee and his or her children.
1 | EDU
1 | EMP | Employee only | Description:The level of coverage under the policy or program is available only to an employee.
1 | ESP | Employee and spouse | Description:The level of coverage under the policy or program is available to an employee and his or her spouse.
1 | FAM | Family | Description:The level of coverage under the policy or program is available to a subscriber's family.
1 | IND | Individual | Description:The level of coverage under the policy or program is available to an individual.
1 | INVOICE
1 | PROA
1 | RECOV
1 | RETRO
1 | SPC | Spouse and children | Description:The level of coverage under the policy or program is available to a subscriber's spouse and children
1 | SPO | Spouse only | Description:The level of coverage under the policy or program is available only to a subscribers spouse

1 | TRAN

4.4.2.9 HL7 v3 Code System CalendarCycle

URL: https://hl7.org/fhir/R4/v3/CalendarCycle/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-CalendarCycle) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-CalendarCycle
Version: | 2018-08-12
Name: | v3.CalendarCycle
Title: | v3 Code System CalendarCycle
Definition: | Calendar cycle identifiers
OID: | 2.16.840.1.113883.5.9 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System CalendarCycle (Calendar cycle identifiers)

4.4.2.9.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | (_CalendarCycleOneLetter) Abstract Deprecated
2 | CW | week (continuous)
2 | CY | year
2 | D | day of the month
2 | DW | day of the week (begins with Monday)
2 | H | hour of the day
2 | M | month of the year
2 | N | minute of the hour
2 | S | second of the minute
1 | (_CalendarCycleTwoLetter) Abstract Deprecated
2 | CW
2 | CY
2 | D
2 | DW
2 | H
2 | M
2 | N
2 | S
2 | CD | day (continuous)
2 | CH | hour (continuous)
2 | CM | month (continuous)
2 | CN | minute (continuous)
2 | CS | second (continuous)
2 | DY | day of the year
2 | WY | week of the year
1 | WM | week of the month | The week with the month's first Thursday in it (analagous to the ISO 8601 definition for week of the year).

4.4.2.10 HL7 v3 Code System ActRelationshipCheckpoint

URL: https://hl7.org/fhir/R4/v3/ActRelationshipCheckpoint/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-ActRelationshipCheckpoint) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-ActRelationshipCheckpoint
Version: | 2018-08-12
Name: | v3.ActRelationshipCheckpoint
Title: | v3 Code System ActRelationshipCheckpoint
Definition: | **** MISSING DEFINITIONS ****
OID: | 2.16.840.1.113883.5.10 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System ActRelationshipCheckpoint (**** MISSING DEFINITIONS ****)

4.4.2.10.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | B | beginning | Condition is tested every time before execution of the service (WHILE condition DO service).
1 | E | end | Condition is tested at the end of a repeated service execution. The service is repeated only if the condition is true (DO service WHILE condition).
1 | S | entry | Condition is tested once before the service is executed (IF condition THEN service).
1 | T | through | Condition must be true throughout the execution and the service is interrupted (asynchronously) as soon as the condition turns false (asynchronous WHILE loop). The service must be interruptible.
1 | X | exit | Condition is a loop checkpoint, i.e. it is a step of an activity plan and, if negative causes the containing loop to exit.

4.4.2.12 HL7 v3 Code System ActRelationshipJoin

URL: https://hl7.org/fhir/R4/v3/ActRelationshipJoin/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-ActRelationshipJoin) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-ActRelationshipJoin
Version: | 2018-08-12
Name: | v3.ActRelationshipJoin
Title: | v3 Code System ActRelationshipJoin
Definition: | **** MISSING DEFINITIONS ****
OID: | 2.16.840.1.113883.5.12 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System ActRelationshipJoin (**** MISSING DEFINITIONS ****)

4.4.2.12.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | D | detached | Detach this branch from the other branches so it will not be resynchronized with the other branches.
1 | K | kill | When all other concurrent branches are terminated, interrupt and discontinue this branch.
1 | W | wait | Wait for this branch to terminate.
1 | X | exclusive wait | Wait for any one of the branches in the set of exclusive wait branches to terminate, then discontinue all the other exclusive wait branches.

4.4.2.13 HL7 v3 Code System ActRelationshipSplit

URL: https://hl7.org/fhir/R4/v3/ActRelationshipSplit/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-ActRelationshipSplit) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-ActRelationshipSplit
Version: | 2018-08-12
Name: | v3.ActRelationshipSplit
Title: | v3 Code System ActRelationshipSplit
Definition: | **** MISSING DEFINITIONS ****
OID: | 2.16.840.1.113883.5.13 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System ActRelationshipSplit (**** MISSING DEFINITIONS ****)

4.4.2.13.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | E1 | exclusive try once | The pre-condition associated with the branch is evaluated once and if true the branch may be entered. All other exclusive branches compete with each other and only one will be selected. This implements a COND, IF and CASE conditionals, or "XOR-split." The order in which the branches are considered may be specified in the priorityNumber attribute.
1 | EW | exclusive wait | A branch is selected as soon as the pre-condition associated with the branch evaluates to true. If the condition is false, the branch may be entered later, when the condition turns true. All other exclusive branches compete with each other and only one will be selected. Each waiting branch executes in parallel with the default join code wait (see below). The order in which the branches are considered may be specified in the Service_relationship.priority_nmb.
1 | I1 | inclusive try once | A branch is executed if its associated preconditions permit. If associated preconditions do not permit, the branch is dropped. Inclusive branches are not suppressed and do not suppress other branches.
1 | IW | inclusive wait | A branch is executed as soon as its associated conditions permit. If the condition is false, the branch may be entered later, when the condition turns true. Inclusive branches are not suppressed and do not suppress other branches. Each waiting branch executes in parallel with the default join code wait (see below).

4.4.2.14 HL7 v3 Code System ActStatus

URL: https://hl7.org/fhir/R4/v3/ActStatus/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-ActStatus) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-ActStatus
Version: | 2018-08-12
Name: | v3.ActStatus
Title: | v3 Code System ActStatus
Definition: | Codes representing the defined possible states of an Act, as defined by the Act class state machine.
OID: | 2.16.840.1.113883.5.14 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System ActStatus (Codes representing the defined possible states of an Act, as defined by the Act class state machine.)

• ValueSet: V3 Value SetProvenanceEventCurrentState-AS (Specifies the state change of a target Act, using ActStatus codes, from its previous state as a predecessor Act. For example, if the target Act is the result of a predecessor Act being "obsoleted" and replaced with the target Act, the source ProvenanceEventCurrentState Act code would be "obsoleted".)

4.4.2.14.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | normal | normal | Encompasses the expected states of an Act, but excludes "nullified" and "obsolete" which represent unusual terminal states for the life-cycle.
2 | aborted | aborted | The Act has been terminated prior to the originally intended completion.
2 | active | active | The Act can be performed or is being performed
2 | cancelled | cancelled | The Act has been abandoned before activation.
2 | completed | completed | An Act that has terminated normally after all of its constituents have been performed.
2 | held | held | An Act that is still in the preparatory stages has been put aside. No action can occur until the Act is released.
2 | new | new | An Act that is in the preparatory stages and may not yet be acted upon
2 | suspended | suspended | An Act that has been activated (actions could or have been performed against it), but has been temporarily disabled. No further action should be taken against it until it is released
1 | nullified | nullified | This Act instance was created in error and has been 'removed' and is treated as though it never existed. A record is retained for audit purposes only.
1 | obsolete | obsolete | This Act instance has been replaced by a new instance.

4.4.2.16 HL7 v3 Code System AddressPartType

URL: https://hl7.org/fhir/R4/v3/AddressPartType/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-AddressPartType) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-AddressPartType
Version: | 2018-08-12
Name: | v3.AddressPartType
Title: | v3 Code System AddressPartType
Definition: | Description: Code that specifies whether an address part names the street, city, country, postal code, post box, etc. Discussion: The hierarchical nature of these concepts shows composition. E.g. "Street Name" is part of "Street Address Line"
OID: | 2.16.840.1.113883.5.16 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System AddressPartType (Description: Code that specifies whether an address part names the street, city, country, postal code, post box, etc. Discussion: The hierarchical nature of these concepts shows composition. E.g. "Street Name" is part of "Street Address Line")

4.4.2.16.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition | PartOf
1 | ADL | additional locator | This can be a unit designator, such as apartment number, suite number, or floor. There may be several unit designators in an address (e.g., "3rd floor, Appt. 342"). This can also be a designator pointing away from the location, rather than specifying a smaller location within some larger one (e.g., Dutch "t.o." means "opposite to" for house boats located across the street facing houses). | AL
1 | AL | address line | Description: An address line is for either an additional locator, a delivery address or a street address.

2 | DAL | delivery address line | A delivery address line is frequently used instead of breaking out delivery mode, delivery installation, etc. An address generally has only a delivery address line or a street address line, but not both.
2 | SAL | street address line | Description: A street address line is frequently used instead of breaking out build number, street name, street type, etc. An address generally has only a delivery address line or a street address line, but not both.
1 | BNN | building number numeric | The numeric portion of a building number | BNR
1 | BNR | building number | The number of a building, house or lot alongside the street. Also known as "primary street number". This does not number the street but rather the building. | SAL
1 | BNS | building number suffix | Any alphabetic character, fraction or other text that may appear after the numeric portion of a building number | BNR
1 | CAR | care of | The name of the party who will take receipt at the specified address, and will take on responsibility for ensuring delivery to the target recipient
1 | CEN | census tract | A geographic sub-unit delineated for demographic purposes.
1 | CNT | country | Country
1 | CPA | county or parish | A sub-unit of a state or province. (49 of the United States of America use the term "county;" Louisiana uses the term "parish".)
1 | CTY | municipality | The name of the city, town, village, or other community or delivery center
1 | DEL | delimiter | Delimiters are printed without framing white space. If no value component is provided, the delimiter appears as a line break.
1 | DINST | delivery installation type | Indicates the type of delivery installation (the facility to which the mail will be delivered prior to final shipping via the delivery mode.) Example: post office, letter carrier depot, community mail center, station, etc. | DAL
1 | DINSTA | delivery installation area | The location of the delivery installation, usually a town or city, and is only required if the area is different from the municipality. Area to which mail delivery service is provided from any postal facility or service such as an individual letter carrier, rural route, or postal route. | DAL
1 | DINSTQ | delivery installation qualifier | A number, letter or name identifying a delivery installation. E.g., for Station A, the delivery installation qualifier would be 'A'. | DAL
1 | DIR | direction | Direction (e.g., N, S, W, E) | SAL
1 | DMOD | delivery mode | Indicates the type of service offered, method of delivery. For example: post office box, rural route, general delivery, etc. | DAL
1 | DMODID | delivery mode identifier | Represents the routing information such as a letter carrier route number. It is the identifying number of the designator (the box number or rural route number). | DAL
1 | DPID | delivery point identifier | A value that uniquely identifies the postal address.
1 | INT | intersection | Description:An intersection denotes that the actual address is located AT or CLOSE TO the intersection OF two or more streets. | AL
1 | POB | post box | A numbered box located in a post station. | DMODID
1 | PRE | precinct | A subsection of a municipality
1 | STA | state or province | A sub-unit of a country with limited sovereignty in a federally organized country.
1 | STB | street name base | The base name of a roadway or artery recognized by a municipality (excluding street type and direction) | STR
1 | STR | street name | | SAL
1 | STTYP | street type | The designation given to the street. (e.g. Street, Avenue, Crescent, etc.) | STR
1 | UNID | unit identifier | The number or name of a specific unit contained within a building or complex, as assigned by that building or complex. | ADL
1 | UNIT | unit designator | Indicates the type of specific unit contained within a building or complex. E.g. Appartment, Floor | ADL
1 | ZIP | postal code | A postal code designating a region defined by the postal service.

4.4.2.18 HL7 v3 Code System AcknowledgementType

URL: https://hl7.org/fhir/R4/v3/AcknowledgementType/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-AcknowledgementType) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-AcknowledgementType
Version: | 2018-08-12
Name: | v3.AcknowledgementType
Title: | v3 Code System AcknowledgementType
Definition: | This attribute contains an acknowledgement code as described in the HL7 message processing rules. OpenIssue: Description was copied from attribute and needs to be improved to be appropriate for a code system.
OID: | 2.16.840.1.113883.5.18 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System AcknowledgementType (This attribute contains an acknowledgement code as described in the HL7 message processing rules. OpenIssue: Description was copied from attribute and needs to be improved to be appropriate for a code system.)

4.4.2.18.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | AA | Application Acknowledgement Accept | Receiving application successfully processed message.

1 | AE | Application Acknowledgement Error | Receiving application found error in processing message. Sending error response with additional error detail information.
1 | AR | Application Acknowledgement Reject | Receiving application failed to process message for reason unrelated to content or format. Original message sender must decide on whether to automatically send message again.
1 | CA | Accept Acknowledgement Commit Accept | Receiving message handling service accepts responsibility for passing message onto receiving application.
1 | CE | Accept Acknowledgement Commit Error | Receiving message handling service cannot accept message for any other reason (e.g. message sequence number, etc.).
1 | CR | Accept Acknowledgement Commit Reject | Receiving message handling service rejects message if interaction identifier, version or processing mode is incompatible with known receiving application role information.

4.4.2.21 HL7 v3 Code System Charset

URL: https://hl7.org/fhir/R4/v3/Charset/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-Charset) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-Charset
Version: | 2018-08-12
Name: | v3.Charset
Title: | v3 Code System Charset
Definition: | Internet Assigned Numbers Authority (IANA) Charset Types
OID: | 2.16.840.1.113883.5.21 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System Charset (Internet Assigned Numbers Authority (IANA) Charset Types)

4.4.2.21.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | EBCDIC | EBCDIC | HL7 is indifferent to the use of this Charset.
1 | ISO-10646-UCS-2 | ISO-10646-UCS-2 | Deprecated for HL7 use.
1 | ISO-10646-UCS-4 | ISO-10646-UCS-4 | Deprecated for HL7 use.
1 | ISO-8859-1 | ISO-8859-1 | HL7 is indifferent to the use of this Charset.
1 | ISO-8859-2 | ISO-8859-2 | HL7 is indifferent to the use of this Charset.
1 | ISO-8859-5 | ISO-8859-5 | HL7 is indifferent to the use of this Charset.
1 | JIS-2022-JP | JIS-2022-JP | HL7 is indifferent to the use of this Charset.
1 | US-ASCII | US-ASCII | Required for HL7 use.
1 | UTF-7 | UTF-7 | HL7 is indifferent to the use of this Charset.
1 | UTF-8 | UTF-8 | Required for Unicode support.

4.4.2.25 HL7 v3 Code System Confidentiality

URL: https://hl7.org/fhir/R4/v3/Confidentiality/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-Confidentiality) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-Confidentiality
Version: | 2018-08-12
Name: | v3.Confidentiality
Title: | v3 Code System Confidentiality
Definition: | A set of codes specifying the security classification of acts and roles in accordance with the definition for concept domain "Confidentiality".
OID: | 2.16.840.1.113883.5.25 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: V3 Value Setx_BasicConfidentialityKind (Description: Used to enumerate the typical confidentiality constraints placed upon a clinical document. Usage Note: x_BasicConfidentialityKind is a subset of Confidentiality codes that are used as metadata indicating the receiver responsibility to comply with normally applicable jurisdictional privacy law or disclosure authorization; that the receiver may not disclose this information except as directed by the information custodian, who may be the information subject; or that the receiver may not disclose this information except as directed by the information custodian, who may be the information subject.)

• ValueSet: v3 Code System Confidentiality (A set of codes specifying the security classification of acts and roles in accordance with the definition for concept domain "Confidentiality".)

• ValueSet: V3 Value SetConfidentialityClassification (Set of codes used to value Act.Confidentiality and Role.Confidentiality attribute in accordance with the definition for concept domain "Confidentiality".)

4.4.2.25.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition

1 | _Confidentiality | Confidentiality | A specializable code and its leaf codes used in Confidentiality value sets to value the Act.Confidentiality and Role.Confidentiality attribute in accordance with the definition for concept domain "Confidentiality".

2 | L | low | Definition: Privacy metadata indicating that the information has been de-identified, and there are mitigating circumstances that prevent re-identification, which minimize risk of harm from unauthorized disclosure. The information requires protection to maintain low sensitivity. Examples: Includes anonymized, pseudonymized, or non-personally identifiable information such as HIPAA limited data sets. Map: No clear map to ISO 13606-4 Sensitivity Level (1) Care Management: RECORD_COMPONENTs that might need to be accessed by a wide range of administrative staff to manage the subject of care's access to health services. Usage Note: This metadata indicates the receiver may have an obligation to comply with a data use agreement.

2 | M | moderate | Definition: Privacy metadata indicating moderately sensitive information, which presents moderate risk of harm if disclosed without authorization. Examples: Includes allergies of non-sensitive nature used inform food service; health information a patient authorizes to be used for marketing, released to a bank for a health credit card or savings account; or information in personal health record systems that are not governed under health privacy laws. Map: Partial Map to ISO 13606-4 Sensitivity Level (2) Clinical Management: Less sensitive RECORD_COMPONENTs that might need to be accessed by a wider range of personnel not all of whom are actively caring for the patient (e.g. radiology staff). Usage Note: This metadata indicates that the receiver may be obligated to comply with the receiver's terms of use or privacy policies.

2 | N | normal | Definition: Privacy metadata indicating that the information is typical, non-stigmatizing health information, which presents typical risk of harm if disclosed without authorization. Examples: In the US, this includes what HIPAA identifies as the minimum necessary protected health information (PHI) given a covered purpose of use (treatment, payment, or operations). Includes typical, non-stigmatizing health information disclosed in an application for health, workers compensation, disability, or life insurance. Map: Partial Map to ISO 13606-4 Sensitivity Level (3) Clinical Care: Default for normal clinical care access (i.e. most clinical staff directly caring for the patient should be able to access nearly all of the EHR). Maps to normal confidentiality for treatment information but not to ancillary care, payment and operations. Usage Note: This metadata indicates that the receiver may be obligated to comply with applicable jurisdictional privacy law or disclosure authorization.

2 | R | restricted | Privacy metadata indicating highly sensitive, potentially stigmatizing information, which presents a high risk to the information subject if disclosed without authorization. May be pre-empted by jurisdictional law, e.g., for public health reporting or emergency treatment. Examples: Includes information that is additionally protected such as sensitive conditions mental health, HIV, substance abuse, domestic violence, child abuse, genetic disease, and reproductive health; or sensitive demographic information such as a patient's standing as an employee or a celebrity. May be used to indicate proprietary or classified information that is not related to an individual, e.g., secret ingredients in a therapeutic substance; or the name of a manufacturer. Map: Partial Map to ISO 13606-4 Sensitivity Level (3) Clinical Care: Default for normal clinical care access (i.e. most clinical staff directly caring for the patient should be able to access nearly all of the EHR). Maps to normal confidentiality for treatment information but not to ancillary care, payment and operations.. Usage Note: This metadata indicates that the receiver may be obligated to comply with applicable, prevailing (default) jurisdictional privacy law or disclosure authorization..

2 | U | unrestricted | Definition: Privacy metadata indicating that the information is not classified as sensitive. Examples: Includes publicly available information, e.g., business name, phone, email or physical address. Usage Note: This metadata indicates that the receiver has no obligation to consider additional policies when making access control decisions. Note that in some jurisdictions, personally identifiable information must be protected as confidential, so it would not be appropriate to assign a confidentiality code of "unrestricted" to that information even if it is publicly available.

2 | V | very restricted | . Privacy metadata indicating that the information is extremely sensitive and likely stigmatizing health information that presents a very high risk if disclosed without authorization. This information must be kept in the highest confidence. Examples: Includes information about a victim of abuse, patient requested information sensitivity, and taboo subjects relating to health status that must be discussed with the patient by an attending provider before sharing with the patient. May also include information held under â€œlegal lockâ€? or attorney-client privilege Map: This metadata indicates that the receiver may not disclose this information except as directed by the information custodian, who may be the information subject. Usage Note: This metadata indicates that the receiver may not disclose this information except as directed by the information custodian, who may be the information subject.

1 | (_ConfidentialityByAccessKind) Abstract Deprecated | | Description: By accessing subject / role and relationship based rights (These concepts are mutually exclusive, one and only one is required for a valid confidentiality coding.) Deprecation Comment:Deprecated due to updated confidentiality codes under ActCode

2 | B Deprecated | business | Description: Since the service class can represent knowledge structures that may be considered a trade or business secret, there is sometimes (though rarely) the need to flag those items as of business level confidentiality. However, no patient related information may ever be of this confidentiality level. Deprecation Comment: Replced by ActCode.B

2 | D Deprecated | clinician | Description: Only clinicians may see this item, billing and administration persons can not access this item without special permission. Deprecation Comment:Deprecated due to updated confidentiality codes under ActCode

2 | I Deprecated | individual | Description: Access only to individual persons who are mentioned explicitly as actors of this service and whose actor type warrants that access (cf. to actor type code). Deprecation Comment:Deprecated due to updated confidentiality codes under ActCode

1 | (_ConfidentialityByInfoType) Abstract Deprecated | | Description: By information type, only for service catalog entries (multiples allowed). Not to be used with actual patient data! Deprecation Comment:Deprecated due to updated confidentiality codes under ActCode

2 | ETH Deprecated | substance abuse related | Description: Alcohol/drug-abuse related item Deprecation Comment:Replced by ActCode.ETH

2 | HIV Deprecated | HIV related | Description: HIV and AIDS related item Deprecation Comment:Replced by ActCode.HIV

2 | PSY Deprecated | psychiatry relate | Description: Psychiatry related item Deprecation Comment:Replced by ActCode.PSY

2 | SDV Deprecated | sexual and domestic violence related | Description: Sexual assault / domestic violence related item Deprecation Comment:Replced by ActCode.SDV

1 | (_ConfidentialityModifiers) Abstract Deprecated | | Description: Modifiers of role based access rights (multiple allowed) Deprecation Comment:Deprecated due to updated confidentiality codes under ActCode

2 | C Deprecated | celebrity | Description: Celebrities are people of public interest (VIP) including employees, whose information require special protection. Deprecation Comment:Replced by ActCode.CEL
2 | S Deprecated | sensitive | Description: Information for which the patient seeks heightened confidentiality. Sensitive information is not to be shared with family members. Information reported by the patient about family members is sensitive by default. Flag can be set or cleared on patient's request. Deprecation Comment:Deprecated due to updated confidentiality codes under ActCode
2 | T Deprecated | taboo | Description: Information not to be disclosed or discussed with patient except through physician assigned to patient in this case. This is usually a temporary constraint only, example use is a new fatal diagnosis or finding, such as malignancy or HIV. Deprecation Note:Replced by ActCode.TBOO

4.4.2.26 HL7 v3 Code System ContainerCap

URL: https://hl7.org/fhir/R4/v3/ContainerCap/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-ContainerCap) is defined as part of HL7 v3. Related FHIR content: ContainerCap.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-ContainerCap
Version: | 2018-08-12
Name: | v3.ContainerCap
Title: | v3 Code System ContainerCap
Definition: | The type of cap associated with a container
OID: | 2.16.840.1.113883.5.26 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System ContainerCap (The type of cap associated with a container)

4.4.2.26.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | (_MedicationCap) Abstract | | Cap types for medication containers
2 | CHILD | ChildProof | A cap designed to be difficult to open for children. Generally requires multiple simultaneous actions (e.g. squeeze and twist) to open. Used for products that may be dangerous if ingested or overdosed by children.
2 | EASY | EasyOpen | A cap designed to be easily removed. For products intended to be opened by persons with limited strength or dexterity.
1 | FILM | Film | A non-reactive plastic film covering over the opening of a container.
1 | FOIL | Foil | A foil covering (type of foil not specified) over the opening of a container
1 | PUSH | Push Cap | A non-threaded cap that forms a tight-fitting closure on a container by pushing the fitted end into the conatiner opening
1 | SCR | Screw Cap | A threaded cap that is screwed onto the opening of a container

4.4.2.27 HL7 v3 Code System ContainerSeparator

URL: https://hl7.org/fhir/R4/v3/ContainerSeparator/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-ContainerSeparator) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-ContainerSeparator
Version: | 2018-08-12
Name: | v3.ContainerSeparator
Title: | v3 Code System ContainerSeparator
Definition: | A material in a blood collection container that facilites the separation of of blood cells from serum or plasma
OID: | 2.16.840.1.113883.5.27 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System ContainerSeparator (A material in a blood collection container that facilites the separation of of blood cells from serum or plasma)

4.4.2.27.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | GEL | Gel | A gelatinous type of separator material.
1 | NONE | None | No separator material is present in the container.

4.4.2.30 HL7 v3 Code System EntityDeterminer

URL: https://hl7.org/fhir/R4/v3/EntityDeterminer/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-EntityDeterminer) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-EntityDeterminer
Version: | 2018-08-12
Name: | v3.EntityDeterminer
Title: | v3 Code System EntityDeterminer
Definition: | EntityDeterminer in natural language grammar is the class of words that comprises articles, demonstrative pronouns, and quantifiers. In the RIM, determiner is a structural code in the Entity class to distinguish whether any given Entity object stands for some, any one, or a specific thing.
OID: | 2.16.840.1.113883.5.30 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System EntityDeterminer (EntityDeterminer in natural language grammar is the class of words that comprises articles, demonstrative pronouns, and quantifiers. In the RIM, determiner is a structural code in the Entity class to distinguish whether any given Entity object stands for some, any one, or a specific thing.)

• ValueSet: V3 Value SetEntityDeterminerDetermined (The described determiner is used to indicate that the given Entity is taken as a general description of a kind of thing that can be taken in whole, in part, or in multiples.)

4.4.2.30.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | INSTANCE | specific | Description:A determiner that specifies that the Entity object represents a particular physical thing (as opposed to a universal, kind, or class of physical thing). Discussion: It does not matter whether an INSTANCE still exists as a whole at the point in time (or process) when we mention it, for example, a drug product lot is an INSTANCE even though it has been portioned out for retail purpose.
2 | GROUP | specific group | A determiner that specifies that the Entity object represents a particular collection of physical things (as opposed to a universal, kind, or class of physical thing). While the collection may resolve to having only a single individual (or even no individuals), the potential should exist for it to cover multiple individuals.
1 | KIND | described | Description:A determiner that specifies that the Entity object represents a universal, kind or class of physical thing (as opposed to a particular thing).
2 | GROUPKIND | described group | A determiner that specifies that the Entity object represents a universal, kind or class of collections physical things. While the collection may resolve to having only a single individual (or even no individuals), the potential should exist for it to cover multiple individuals.
2 | QUANTIFIED_KIND Deprecated | described quantified | The described quantified determiner indicates that the given Entity is taken as a general description of a specific amount of a thing. For example, QUANTIFIED_KIND of syringe (quantity = 3,) stands for exactly three syringes.

4.4.2.31 HL7 v3 Code System DeviceAlertLevel

URL: https://hl7.org/fhir/R4/v3/DeviceAlertLevel/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-DeviceAlertLevel) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-DeviceAlertLevel
Version: | 2018-08-12
Name: | v3.DeviceAlertLevel
Title: | v3 Code System DeviceAlertLevel
Definition: | Domain values for the Device.Alert_levelCode
OID: | 2.16.840.1.113883.5.31 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System DeviceAlertLevel (Domain values for the Device.Alert_levelCode)

4.4.2.31.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | C | Critical | Shut Down, Fix Problem and Re-initialize
1 | N | Normal | No Corrective Action Needed
1 | S | Serious | Corrective Action Required
1 | W | Warning | Corrective Action Anticipated

4.4.2.33 HL7 v3 Code System DocumentCompletion

URL: https://hl7.org/fhir/R4/v3/DocumentCompletion/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-DocumentCompletion) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-DocumentCompletion
Version: | 2018-08-12
Name: | v3.DocumentCompletion
Title: | v3 Code System DocumentCompletion
Definition: | Identifies the current completion state of a clinical document.
OID: | 2.16.840.1.113883.5.33 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: Provenance activity type (This value set contains representative Activity Type codes, which includes codes from the HL7 DocumentCompletion, ActStatus, and DataOperations code system, W3C PROV-DM and PROV-N concepts and display names, several HL7 Lifecycle Event codes for which there are agreed upon definitions, and non-duplicated codes from the HL7 Security and Privacy Ontology Operations codes.)

• ValueSet: v3 Code System DocumentCompletion (Identifies the current completion state of a clinical document.)

• ValueSet: Provenance activity type (This value set contains representative Activity Type codes, which includes codes from the HL7 DocumentCompletion, ActStatus, and DataOperations code system, W3C PROV-DM and PROV-N concepts and display names, several HL7 Lifecycle Event codes for which there are agreed upon definitions, and non-duplicated codes from the HL7 Security and Privacy Ontology Operations codes.)

• ValueSet: V3 Value SetProvenanceEventCurrentState-DC (Specifies the state change of a target Act using DocuymentCompletion codes, from its previous state as a predecessor Act. For example, if the target Act is the result of a predecessor Act being "obsoleted" and replaced with the target Act, the source ProvenanceEventCurrentState Act code would be "obsoleted".)

4.4.2.33.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | AU | authenticated | A completion status in which a document has been signed manually or electronically by one or more individuals who attest to its accuracy. No explicit determination is made that the assigned individual has performed the authentication. While the standard allows multiple instances of authentication, it would be typical to have a single instance of authentication, usually by the assigned individual.
1 | DI | dictated | A completion status in which information has been orally recorded but not yet transcribed.
1 | DO | documented | A completion status in which document content, other than dictation, has been received but has not been translated into the final electronic format. Examples include paper documents, whether hand-written or typewritten, and intermediate electronic forms, such as voice to text.
1 | IN | incomplete | A completion status in which information is known to be missing from a transcribed document.
1 | IP | in progress | A workflow status where the material has been assigned to personnel to perform the task of transcription. The document remains in this state until the document is transcribed.
1 | LA | legally authenticated | A completion status in which a document has been signed manually or electronically by the individual who is legally responsible for that document. This is the most mature state in the workflow progression.
1 | NU | nullified document | A completion status in which a document was created in error or was placed in the wrong chart. The document is no longer available.
1 | PA | pre-authenticated | A completion status in which a document is transcribed but not authenticated.
1 | UC | unsigned completed document | A completion status where the document is complete and there is no expectation that the document will be signed.

4.4.2.34 HL7 v3 Code System DocumentStorage

URL: https://hl7.org/fhir/R4/v3/DocumentStorage/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-DocumentStorage) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-DocumentStorage
Version: | 2018-08-12
Name: | v3.DocumentStorage
Title: | v3 Code System DocumentStorage
Definition: | Identifies the storage status of a document.
OID: | 2.16.840.1.113883.5.34 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System DocumentStorage (Identifies the storage status of a document.)

4.4.2.34.1 Code System Content

Release Date: 2018-08-12

Level	Code	Display	Definition
1	AC	active	A storage status in which a document is available on-line.
2	AA	active and archived	A storage status in which a document is available on-line and is also stored off-line for long-term access.
1	AR	archived (not active)	A storage status in which a document has been stored off-line for long-term access.
1	PU	purged	A storage status in which a document is no longer available in this system.

4.4.2.37 HL7 v3 Code System EncounterAdmissionSource

URL: https://hl7.org/fhir/R4/v3/EncounterAdmissionSource/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-EncounterAdmissionSource) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-EncounterAdmissionSource
Version: | 2018-08-12
Name: | v3.EncounterAdmissionSource
Title: | v3 Code System EncounterAdmissionSource
Definition: | **** MISSING DEFINITIONS ****
OID: | 2.16.840.1.113883.5.37 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System EncounterAdmissionSource (**** MISSING DEFINITIONS ****)

4.4.2.37.1 Code System Content

Release Date: 2018-08-12

Level	Code	Display	Definition
1	E	emergency	
1	LD	labor and delivery	
1	NB	newborn	

4.4.2.40 HL7 v3 Code System EncounterSpecialCourtesy

URL: https://hl7.org/fhir/R4/v3/EncounterSpecialCourtesy/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-EncounterSpecialCourtesy) is defined as part of HL7 v3. Related FHIR content: Special courtesy.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-EncounterSpecialCourtesy
Version: | 2018-08-12
Name: | v3.EncounterSpecialCourtesy
Title: | v3 Code System EncounterSpecialCourtesy
Definition: | **** MISSING DEFINITIONS ****
OID: | 2.16.840.1.113883.5.40 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: Special courtesy (This value set defines a set of codes that can be used to indicate special courtesies provided to the patient.)

• ValueSet: v3 Code System EncounterSpecialCourtesy (**** MISSING DEFINITIONS ****)

• ValueSet: Special courtesy (This value set defines a set of codes that can be used to indicate special courtesies provided to the patient.)

4.4.2.40.1 Code System Content

Release Date: 2018-08-12

Level	Code	Display	Definition
1	EXT	extended courtesy	
1	NRM	normal courtesy	
1	PRF	professional courtesy	
1	STF	staff	Courtesies extended to the staff of the entity providing service.
1	VIP	very important person	

4.4.2.41 HL7 v3 Code System EntityClass

URL: https://hl7.org/fhir/R4/v3/EntityClass/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-EntityClass) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-EntityClass
Version: | 2018-08-12
Name: | v3.EntityClass
Title: | v3 Code System EntityClass
Definition: | Classifies the Entity class and all of its subclasses. The terminology is hierarchical. At the top is this HL7-defined domain of high-level categories (such as represented by the Entity subclasses). Each of these terms must be harmonized and is specializable. The value sets beneath are drawn from multiple, frequently external, domains that reflect much more fine-grained typing.
OID: | 2.16.840.1.113883.5.41 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: V3 Value SetEntityClassDevice (A subtype of ManufacturedMaterial used in an activity, without being substantially changed through that activity. The kind of device is identified by the code attribute inherited from Entity. Usage: This includes durable (reusable) medical equipment as well as disposable equipment.)

• ValueSet: V3 Value SetEntityClassManufacturedMaterial (Corresponds to the ManufacturedMaterial class)

• ValueSet: v3 Code System EntityClass (Classifies the Entity class and all of its subclasses. The terminology is hierarchical. At the top is this HL7-defined domain of high-level categories (such as represented by the Entity subclasses). Each of these terms must be harmonized and is specializable. The value sets beneath are drawn from multiple, frequently external, domains that reflect much more fine-grained typing.)

• ValueSet: V3 Value SetEntityClassLivingSubject (Anything that essentially has the property of life, independent of current state (a dead human corpse is still essentially a living subject.))

• ValueSet: V3 Value SetEntityClassOrganization (A social or legal structure formed by human beings.)

• ValueSet: V3 Value SetEntityClassPlace (A physicial place or site with its containing structure. May be natural or man-made. The geographic position of a place may or may not be constant.)

• ValueSet: V3 Value SetEntityClassRoot (Corresponds to the Entity class)

4.4.2.41.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | ENT | entity | Corresponds to the Entity class
2 | HCE | health chart entity | A health chart included to serve as a document receiving entity in the management of medical records.
2 | LIV | living subject | Anything that essentially has the property of life, independent of current state (a dead human corpse is still essentially a living subject).
3 | NLIV | non-person living subject | A subtype of living subject that includes all living things except the species Homo Sapiens.
4 | ANM | animal | A living subject from the animal kingdom.
4 | MIC | microorganism | All single celled living organisms including protozoa, bacteria, yeast, viruses, etc.
4 | PLNT | plant | A living subject from the order of plants.
3 | PSN | person | A living subject of the species homo sapiens.
2 | MAT | material | Any thing that has extension in space and mass, may be of living or non-living origin.
3 | CHEM | chemical substance | A substance that is fully defined by an organic or inorganic chemical formula, includes mixtures of other chemical substances. Refine using, e.g., IUPAC codes.
3 | FOOD | food | Naturally occurring, processed or manufactured entities that are primarily used as food for humans and animals.
3 | MMAT | manufactured material | Corresponds to the ManufacturedMaterial class
4 | CONT | container | A container of other entities.
5 | HOLD | holder | A type of container that can hold other containers or other holders.
4 | DEV | device | A subtype of ManufacturedMaterial used in an activity, without being substantially changed through that activity. The kind of device is identified by the code attribute inherited from Entity. Usage: This includes durable (reusable) medical equipment as well as disposable equipment.
5 | CER | certificate representation | A physical artifact that stores information about the granting of authorization.
5 | MODDV | imaging modality | Class to contain unique attributes of diagnostic imaging equipment.
2 | ORG | organization | A social or legal structure formed by human beings.
3 | PUB | public institution | An agency of the people of a state often assuming some authority over a certain matter. Includes government, governmental agencies, associations.
3 | STATE | state | A politically organized body of people bonded by territory, culture, or ethnicity, having sovereignty (to a certain extent) granted by other states (enclosing or neighboring states). This includes countries (nations), provinces (e.g., one of the United States of America or a French departement), counties or municipalities. Refine using, e.g., ISO country codes, FIPS-PUB state codes, etc.
4 | NAT | Nation | A politically organized body of people bonded by territory and known as a nation.

2 | PLC | place | A physical place or site with its containing structure. May be natural or man-made. The geographic position of a place may or may not be constant.
3 | CITY | city or town | The territory of a city, town or other municipality.
3 | COUNTRY | country | The territory of a sovereign nation.
3 | COUNTY | county or parish | The territory of a county, parish or other division of a state or province.
3 | PROVINCE | state or province | The territory of a state, province, department or other division of a sovereign country.
2 | RGRP | group | A grouping of resources (personnel, material, or places) to be used for scheduling purposes. May be a pool of like-type resources, a team, or combination of personnel, material and places.

4.4.2.42 HL7 v3 Code System EntityHandling

URL: https://hl7.org/fhir/R4/v3/EntityHandling/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-EntityHandling) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-EntityHandling
Version: | 2018-08-12
Name: | v3.EntityHandling
Title: | v3 Code System EntityHandling
Definition: | **** MISSING DEFINITIONS ****
OID: | 2.16.840.1.113883.5.42 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System EntityHandling (**** MISSING DEFINITIONS ****)

4.4.2.42.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | AMB | Ambient Temperature | Keep at ambient temperature, 22 +/- 2C
1 | C37 | Body Temperature | Critical to keep at body temperature 36-38C
1 | CAMB | Critical Ambient temperature | Critical ambient - must not be refrigerated or frozen.
1 | CATM | Protect from Air | Critical. Do not expose to atmosphere. Do not uncap.
1 | CFRZ | Critical frozen | Critical frozen. Specimen must not be allowed to thaw until immediately prior to testing.
1 | CREF | Critical refrigerated temperature | Critical refrigerated - must not be allowed to freeze or warm until imediately prior to testing.
1 | DFRZ | Deep Frozen | Deep Frozen -16 to -20C.
1 | DRY | dry | Keep in a dry environment
1 | FRZ | frozen | Keep frozen below 0 ?C
1 | MTLF | Metal Free | Container is free of heavy metals, including lead.
1 | NTR | nitrogen | Keep in liquid nitrogen
1 | PRTL | Protect from Light | Protect from light (eg. Wrap in aluminum foil).
1 | PSA | do not shake | Do not shake
1 | PSO | no shock | Protect against shock
1 | REF | Refrigerated temperature | Keep at refrigerated temperature:4-8C Accidental warming or freezing is of little consequence.
1 | SBU | Shake before use | Shake thoroughly before using
1 | UFRZ | Ultra frozen | Ultra cold frozen -75 to -85C. Ultra cold freezer is typically at temperature of dry ice.
1 | UPR | upright | Keep upright, do not turn upside down

4.4.2.43 HL7 v3 Code System EntityNamePartQualifier

URL: https://hl7.org/fhir/R4/v3/EntityNamePartQualifier/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-EntityNamePartQualifier) is defined as part of HL7 v3. Related FHIR content: EntityNamePartQualifier.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-EntityNamePartQualifier
Version: | 2018-08-12
Name: | v3.EntityNamePartQualifier
Title: | v3 Code System EntityNamePartQualifier
Definition: | OpenIssue: Needs description
OID: | 2.16.840.1.113883.5.43 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: EntityNamePartQualifier (A set of codes each of which specifies a certain subcategory of the name part in addition to the main name part type.)

• ValueSet: EntityNamePartQualifier (A set of codes each of which specifies a certain subcategory of the name part in addition to the main name part type.)

• ValueSet: v3 Code System EntityNamePartQualifier (OpenIssue: Needs description)

4.4.2.43.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition

1 | (_OrganizationNamePartQualifier) Abstract

2 | AC | academic | Indicates that a prefix like "Dr." or a suffix like "M.D." or "Ph.D." is an academic title.

2 | AD | adopted | The name the person was given at the time of adoption.

2 | BR | birth | A name that a person had shortly after being born. Usually for family names but may be used to mark given names at birth that may have changed later.

2 | CL | callme | A callme name is (usually a given name) that is preferred when a person is directly addressed.

2 | IN | initial | Indicates that a name part is just an initial. Initials do not imply a trailing period since this would not work with non-Latin scripts. Initials may consist of more than one letter, e.g., "Ph." could stand for "Philippe" or "Th." for "Thomas".

2 | LS | Legal status | For organizations a suffix indicating the legal status, e.g., "Inc.", "Co.", "AG", "GmbH", "B.V." "S.A.", "Ltd." etc.

2 | NB | nobility | In Europe and Asia, there are still people with nobility titles (aristocrats). German "von" is generally a nobility title, not a mere voorvoegsel. Others are "Earl of" or "His Majesty King of..." etc. Rarely used nowadays, but some systems do keep track of this.

2 | PR | professional | Primarily in the British Imperial culture people tend to have an abbreviation of their professional organization as part of their credential suffices.

2 | SP | spouse | The name assumed from the partner in a marital relationship (hence the "SP"). Usually the spouse's family name. Note that no inference about gender can be made from the existence of spouse names.

2 | TITLE | title | Indicates that a prefix or a suffix is a title that applies to the whole name, not just the adjacent name part.

2 | VV | voorvoegsel | A Dutch "voorvoegsel" is something like "van" or "de" that might have indicated nobility in the past but no longer so. Similar prefixes exist in other languages such as Spanish, French or Portugese.

1 | (PharmaceuticalEntityNamePartQualifiers) Abstract | | Description: Medication Name Parts are a means of specifying a range of acceptable "official" forms of the name of a product. They are used as patterns against which input name strings may be matched for automatic identification of products from input text reports. While they cover the concepts held under "doseForm" or "route" or "strength" the name parts are not the same and do not fit into a controlled vocabulary in the same way. By specifying up to 8 name parts a much larger range of possible names can be generated.

2 | CON | container name | Description: This refers to the container if present in the medicinal product name. EXAMPLES: For Optaflu suspension for injection in pre-filled syringe Influenza vaccine (surface antigen, inactivated, prepared in cell culture) (2007/2008 season): pre-filled syringe

2 | DEV | device name | Description: This refers to the qualifiers in the name for devices and is at the moment mainly applicable to insulins and inhalation products. EXAMPLES: For the medicinal product Actrapid FlexPen 100 IU/ml Solution for injection Subcutaneous use: FlexPen.

2 | FLAV | FlavorName | Description: This refers to a flavor of the medicinal product if present in the medicinal product name. Examples: For 'CoughCure Linctus Orange Flavor', the flavor part is "Orange" For 'Wonderdrug Syrup Cherry Flavor', the flavor part is "Cherry"

2 | FORMUL | FormulationPartName | Description: This refers to the formulation of the medicinal product if present in the medicinal product name. Examples: For 'SpecialMed Sugar Free Cough Syrup', the formulation name part is "Sugar Free" For 'QuickCure Gluten-free Bulk Fibre', the formulation name part is "gluten-free"

2 | FRM | form name | Description: This refers to the pharmaceutical form/ if present in the medicinal product name. EXAMPLES: For Agenerase 50 mg soft capsules: Soft Capsules For Ludiomil 25mg-Filmtabletten: Filmtabletten For Optaflu suspension for injection in pre-filled syringe Influenza vaccine (surface antigen, inactivated, prepared in cell culture) (2007/2008 season): suspension for injection

2 | INV | invented name | Description: This refers to the product name without the trademark or the name of the marketing authorization holder or any other descriptor reflected in the product name and, if appropriate, whether it is intended e.g. for babies, children or adults. EXAMPLES: Agenerase Optaflu Ludiomil

2 | POPUL | TargetPopulationName | Description: This refers to the target population for the medicinal product if present in the medicinal product name Examples: For 'Broncho-Drug 3.5 mg-capsules for children', the target population part is "children" For 'Adult Chesty Cough Syrup', the target population part is "adult"

2 | SCI | scientific name | Description: This refers to the product common or scientific name without the trademark or the name of the marketing authorization holder or any other descriptor reflected in the product name. EXAMPLES: For Agenerase: N/A For Optaflu: Influenza vaccine (surface antigen, inactivated, prepared in cell culture) (2007/2008 season) For Ludiomil: N/A

2 | STR | strength name | Description: This refers to the strength if present in the medicinal product name. The use of decimal points should be accommodated if required. EXAMPLES: For Agenerase 50 mg soft capsules: 50mg For Ludiomil 25mg-Filmtabletten: 25 mg For Optaflu suspension for injection in pre-filled syringe Influenza vaccine (surface antigen, inactivated, prepared in cell culture) (2007/2008 season): N/A

2 | TIME | TimeOrPeriodName | Description: This refers to a time or time period that may be specified in the text of the medicinal product name Example: For an influenza vaccine 'Drug-FLU season 2008/2009', the time/period part is "2008/2009 season"

2 | TMK | trademark name | Description: This refers to trademark/company element if present in the medicinal product name. EXAMPLES: for Insulin Human Winthrop Comb 15: Winthrop

2 | USE | intended use name | Description: This refers to the intended use if present in the medicinal product name without the trademark or the name of the marketing authorization holder or any other descriptor reflected in the product name. Examples: For 'Drug-BI Caplets - Heartburn Relief', the intended use part is: "Heartburn Relief" For 'Medicine Honey Syrup for Soothing Coughs' the intended use part is "Soothing Coughs"

1 | (_PersonNamePartQualifier) Abstract

2 | IN

2 | TITLE

2 | (_PersonNamePartAffixTypes) Abstract

3 | AC

3 | NB

3 | PR
3 | VV
2 | (_PersonNamePartChangeQualifier) Abstract
3 | AD
3 | BR
3 | SP
2 | (_PersonNamePartMiscQualifier) Abstract
3 | CL

4.4.2.44 HL7 v3 Code System EntityNamePartType

URL: https://hl7.org/fhir/R4/v3/EntityNamePartType/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-EntityNamePartType) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-EntityNamePartType
Version: | 2018-08-12
Name: | v3.EntityNamePartType
Title: | v3 Code System EntityNamePartType
Definition: | **** MISSING DEFINITIONS ****
OID: | 2.16.840.1.113883.5.44 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System EntityNamePartType (**** MISSING DEFINITIONS ****)

4.4.2.44.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | DEL | delimiter | A delimiter has no meaning other than being literally printed in this name representation. A delimiter has no implicit leading and trailing white space.
1 | FAM | family | Family name, this is the name that links to the genealogy. In some cultures (e.g. Eritrea) the family name of a son is the first name of his father.
1 | GIV | given | Given name (don't call it "first name" since this given names do not always come first)
1 | PFX | prefix | A prefix has a strong association to the immediately following name part. A prefix has no implicit trailing white space (it has implicit leading white space though). Note that prefixes can be inverted.
1 | SFX | suffix | Description:A suffix has a strong association to the immediately preceding name part. A suffix has no implicit leading white space (it has implicit trailing white space though). Suffices cannot be inverted.

4.4.2.45 HL7 v3 Code System EntityNameUse

URL: https://hl7.org/fhir/R4/v3/EntityNameUse/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-EntityNameUse) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-EntityNameUse
Version: | 2018-08-12
Name: | v3.EntityNameUse
Title: | v3 Code System EntityNameUse
Definition: | **** MISSING DEFINITIONS ****
OID: | 2.16.840.1.113883.5.45 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System EntityNameUse (**** MISSING DEFINITIONS ****)
• ValueSet: NameRepresentationUse (A set of codes for each different representation of a name.)
• ValueSet: NameRepresentationUse (A set of codes for each different representation of a name.)

4.4.2.45.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | (_NameRepresentationUse) Abstract | | Identifies the different representations of a name. The representation may affect how the name is used. (E.g. use of Ideographic for formal communications.)
2 | ABC | Alphabetic | Alphabetic transcription of name (Japanese: romaji)
2 | IDE | Ideographic | Ideographic representation of name (e.g., Japanese kanji, Chinese characters)
2 | SYL | Syllabic | Syllabic transcription of name (e.g., Japanese kana, Korean hangul)

1 | ASGN | assigned | A name assigned to a person. Reasons some organizations assign alternate names may include not knowing the person's name, or to maintain anonymity. Some, but not necessarily all, of the name types that people call "alias" may fit into this category.
1 | C | License | As recorded on a license, record, certificate, etc. (only if different from legal name)
1 | I | Indigenous/Tribal | e.g. Chief Red Cloud
1 | L | Legal | Known as/conventional/the one you use
2 | OR | official registry | Definition:The formal name as registered in an official (government) registry, but which name might not be commonly used. Particularly used in countries with a law system based on Napoleonic law.
1 | P | pseudonym | A self asserted name that the person is using or has used.
2 | A | Artist/Stage | Includes writer's pseudonym, stage name, etc
1 | R | Religious | e.g. Sister Mary Francis, Brother John
1 | SRCH | search | A name intended for use in searching or matching.
2 | PHON | phonetic | A name spelled phonetically. There are a variety of phonetic spelling algorithms. This code value does not distinguish between these.Discussion:
2 | SNDX | Soundex | A name spelled according to the SoundEx algorithm.

4.4.2.46 HL7 v3 Code System EntityRisk

URL: https://hl7.org/fhir/R4/v3/EntityRisk/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-EntityRisk) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-EntityRisk
Version: | 2018-08-12
Name: | v3.EntityRisk
Title: | v3 Code System EntityRisk
Definition: | Kinds of risks associated with the handling of the material..
OID: | 2.16.840.1.113883.5.46 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System EntityRisk (Kinds of risks associated with the handling of the material..)

4.4.2.46.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | AGG | aggressive | A danger that can be associated with certain living subjects, including humans.
1 | BIO | Biological | The dangers associated with normal biological materials. I.e. potential risk of unknown infections. Routine biological materials from living subjects.
1 | COR | Corrosive | Material is corrosive and may cause severe injury to skin, mucous membranes and eyes. Avoid any unprotected contact.
1 | ESC | Escape Risk | The entity is at risk for escaping from containment or control.
1 | IFL | inflammable | Material is highly inflammable and in certain mixtures (with air) may lead to explosions. Keep away from fire, sparks and excessive heat.
2 | EXP | explosive | Material is an explosive mixture. Keep away from fire, sparks, and heat.
1 | INF | infectious | Material known to be infectious with human pathogenic microorganisms. Those who handle this material must take precautions for their protection.
2 | BHZ | biohazard | Material contains microorganisms that is an environmental hazard. Must be handled with special care.
1 | INJ | injury hazard | Material is solid and sharp (e.g., cannulas). Dispose in hard container.
1 | POI | poison | Material is poisonous to humans and/or animals. Special care must be taken to avoid incorporation, even of small amounts.
1 | RAD | radioactive | Material is a source for ionizing radiation and must be handled with special care to avoid injury of those who handle it and to avoid environmental hazards.

4.4.2.49 HL7 v3 Code System EquipmentAlertLevel

URL: https://hl7.org/fhir/R4/v3/EquipmentAlertLevel/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-EquipmentAlertLevel) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-EquipmentAlertLevel
Version: | 2018-08-12
Name: | v3.EquipmentAlertLevel
Title: | v3 Code System EquipmentAlertLevel
Definition: | **** MISSING DEFINITIONS ****
OID: | 2.16.840.1.113883.5.49 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System EquipmentAlertLevel (**** MISSING DEFINITIONS ****)

4.4.2.49.1 Code System Content

Release Date: 2018-08-12

Level	Code	Display	Definition
1	C	Critical	Shut Down, Fix Problem and Re-init
1	N	Normal	No Corrective Action Needed
1	S	Serious	Corrective Action Required
1	W	Warning	Corrective Action Anticipated

4.4.2.50 HL7 v3 Code System Ethnicity

URL: https://hl7.org/fhir/R4/v3/Ethnicity/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-Ethnicity) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-Ethnicity
Version: | 2018-08-12
Name: | v3.Ethnicity
Title: | v3 Code System Ethnicity
Definition: | In the United States, federal standards for classifying data on ethnicity determine the categories used by federal agencies and exert a strong influence on categorization by state and local agencies and private sector organizations. The federal standards do not conceptually define ethnicity, and they recognize the absence of an anthropological or scientific basis for ethnicity classification. Instead, the federal standards acknowledge that ethnicity is a social-political construct in which an individual's own identification with a particular ethnicity is preferred to observer identification. The standards specify two minimum ethnicity categories: Hispanic or Latino, and Not Hispanic or Latino. The standards define a Hispanic or Latino as a person of "Mexican, Puerto Rican, Cuban, South or Central America, or other Spanish culture or origin, regardless of race." The standards stipulate that ethnicity data need not be limited to the two minimum categories, but any expansion must be collapsible to those categories. In addition, the standards stipulate that an individual can be Hispanic or Latino or can be Not Hispanic or Latino, but cannot be both.
OID: | 2.16.840.1.113883.5.50 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System Ethnicity (In the United States, federal standards for classifying data on ethnicity determine the categories used by federal agencies and exert a strong influence on categorization by state and local agencies and private sector organizations. The federal standards do not conceptually define ethnicity, and they recognize the absence of an anthropological or scientific basis for ethnicity classification. Instead, the federal standards acknowledge that ethnicity is a social-political construct in which an individual's own identification with a particular ethnicity is preferred to observer identification. The standards specify two minimum ethnicity categories: Hispanic or Latino, and Not Hispanic or Latino. The standards define a Hispanic or Latino as a person of "Mexican, Puerto Rican, Cuban, South or Central America, or other Spanish culture or origin, regardless of race." The standards stipulate that ethnicity data need not be limited to the two minimum categories, but any expansion must be collapsible to those categories. In addition, the standards stipulate that an individual can be Hispanic or Latino or can be Not Hispanic or Latino, but cannot be both.)

4.4.2.50.1 Code System Content

Release Date: 2018-08-12

Level	Code	Display	Definition
1	2135-2	Hispanic or Latino	
2	2137-8	Spaniard	
3	2138-6	Andalusian	
3	2139-4	Asturian	
3	2140-2	Castillian	
3	2141-0	Catalonian	
3	2142-8	Belearic Islander	
3	2143-6	Gallego	
3	2144-4	Valencian	
3	2145-1	Canarian	
3	2146-9	Spanish Basque	
2	2148-5	Mexican	
3	2149-3	Mexican American	
3	2150-1	Mexicano	
3	2151-9	Chicano	
3	2152-7	La Raza	
3	2153-5	Mexican American Indian	
2	2155-0	Central American	
3	2156-8	Costa Rican	
3	2157-6	Guatemalan	

3 | 2158-4 | Honduran
3 | 2159-2 | Nicaraguan
3 | 2160-0 | Panamanian
3 | 2161-8 | Salvadoran
3 | 2162-6 | Central American Indian
3 | 2163-4 | Canal Zone
2 | 2165-9 | South American
3 | 2166-7 | Argentinean
3 | 2167-5 | Bolivian
3 | 2168-3 | Chilean
3 | 2169-1 | Colombian
3 | 2170-9 | Ecuadorian
3 | 2171-7 | Paraguayan
3 | 2172-5 | Peruvian
3 | 2173-3 | Uruguayan
3 | 2174-1 | Venezuelan
3 | 2175-8 | South American Indian
3 | 2176-6 | Criollo
2 | 2178-2 | Latin American
2 | 2180-8 | Puerto Rican
2 | 2182-4 | Cuban
2 | 2184-0 | Dominican
1 | 2186-5 | Not Hispanic or Latino | Note that this term remains in the table for completeness, even though within HL7, the notion of "not otherwise coded" term is deprecated.

4.4.2.51 HL7 v3 Code System GenderStatus

URL: https://hl7.org/fhir/R4/v3/GenderStatus/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-GenderStatus) is defined as part of HL7 v3. Related FHIR content: Gender status.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-GenderStatus
Version: | 2018-08-12
Name: | v3.GenderStatus
Title: | v3 Code System GenderStatus
Definition: | **** MISSING DEFINITIONS ****
OID: | 2.16.840.1.113883.5.51 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System GenderStatus (**** MISSING DEFINITIONS ****)

4.4.2.51.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | I | Intact | Reproductively intact
1 | N | Neutered | Reproductively neutered

4.4.2.57 HL7 v3 Code System HL7UpdateMode

URL: https://hl7.org/fhir/R4/v3/HL7UpdateMode/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-HL7UpdateMode) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-HL7UpdateMode
Version: | 2018-08-12
Name: | v3.HL7UpdateMode
Title: | v3 Code System HL7UpdateMode
Definition: | The possible modes of updating that occur when an attribute is received by a system that already contains values for that attribute.
OID: | 2.16.840.1.113883.5.57 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System HL7UpdateMode (The possible modes of updating that occur when an attribute is received by a system that already contains values for that attribute.)

4.4.2.57.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | A | Add | Description:The item was (or is to be) added, having not been present immediately before. (If it is already present, this may be treated as an error condition.)
1 | AR | Add or Replace | Description:The item was (or is to be) either added or replaced.
1 | D | Remove | Description:The item was (or is to be) removed (sometimes referred to as deleted). If the item is part of a collection, delete any matching items.
1 | K | Key | Description:This item is part of the identifying information for this object.
1 | N | No Change | Description:There was (or is to be) no change to the item. This is primarily used when this element has not changed, but other attributes in the instance have changed.
1 | R | Replace | Description:The item existed previously and has (or is to be) revised. (If an item does not already exist, this may be treated as an error condition.)
1 | REF | Reference | Description:This item provides enough information to allow a processing system to locate the full applicable record by identifying the object.
1 | U | Unknown | Description:Description:It is not specified whether or what kind of change has occurred to the item, or whether the item is present as a reference or identifying property.
1 | (_SetUpdateMode) Abstract | | These concepts apply when the element and/or message is updating a set of items.
2 | ESA | Set Add | Add the message element to the collection of items on the receiving system that correspond to the message element.
2 | ESAC | Set Add or Change | Change the item on the receiving system that corresponds to this message element; if a matching element does not exist, add a new one created with the values in the message.
2 | ESC | Set Change | Change the item on the receiving system that corresponds to this message element; do not process if a matching element does not exist.
2 | ESD | Set Delete | Delete the item on the receiving system that corresponds to this message element.
1 | AU | Add or Update | Description: AU: If this item exists, update it with these values. If it does not exist, create it with these values. If the item is part of the collection, update each item that matches this item, and if no items match, add a new item to the collection.
1 | I | Ignore | Ignore this role, it is not relevant to the update.
1 | V | Verify | Verify - this message element must match a value already in the receiving systems database in order to process the message.

4.4.2.58 HL7 v3 Code System HtmlLinkType

URL: https://hl7.org/fhir/R4/v3/HtmlLinkType/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-HtmlLinkType) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-HtmlLinkType
Version: | 2018-08-12
Name: | v3.HtmlLinkType
Title: | v3 Code System HtmlLinkType
Definition: | HtmlLinkType values are drawn from HTML 4.0 and describe the relationship between the current document and the anchor that is the target of the link
OID: | 2.16.840.1.113883.5.58 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System HtmlLinkType (HtmlLinkType values are drawn from HTML 4.0 and describe the relationship between the current document and the anchor that is the target of the link)

4.4.2.58.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | alternate | alternate | Designates substitute versions for the document in which the link occurs. When used together with the lang attribute, it implies a translated version of the document. When used together with the media attribute, it implies a version designed for a different medium (or media).
1 | appendix | appendix | Refers to a document serving as an appendix in a collection of documents.
1 | bookmark | bookmark | Refers to a bookmark. A bookmark is a link to a key entry point within an extended document. The title attribute may be used, for example, to label the bookmark. Note that several bookmarks may be defined in each document.
1 | chapter | chapter | Refers to a document serving as a chapter in a collection of documents.
1 | contents | contents | Refers to a document serving as a table of contents. Some user agents also support the synonym ToC (from "Table of Contents").
1 | copyright | copyright | Refers to a copyright statement for the current document.
1 | glossary | glossary | Refers to a document providing a glossary of terms that pertain to the current document.
1 | help | help | Refers to a document offering help (more information, links to other sources of information, etc.).
1 | index | index | Refers to a document providing an index for the current document.
1 | next | next | Refers to the next document in a linear sequence of documents. User agents may choose to preload the "next" document, to reduce the perceived load time.
1 | prev | prev | Refers to the previous document in an ordered series of documents. Some user agents also support the synonym "Previous".
1 | section | section | Refers to a document serving as a section in a collection of documents.

1 | start | start | Refers to the first document in a collection of documents. This link type tells search engines which document is considered by the author to be the starting point of the collection.
1 | stylesheet | stylesheet | Refers to an external style sheet. See the section on external style sheets for details. This is used together with the link type "Alternate" for user-selectable alternate style sheets.
1 | subsection | subsection | Refers to a document serving as a subsection in a collection of documents.

4.4.2.60 HL7 v3 Code System LanguageAbilityMode

URL: https://hl7.org/fhir/R4/v3/LanguageAbilityMode/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-LanguageAbilityMode) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-LanguageAbilityMode
Version: | 2018-08-12
Name: | v3.LanguageAbilityMode
Title: | v3 Code System LanguageAbilityMode
Definition: | A value representing the method of expression of the language. Example: Expressed spoken, expressed written, expressed signed, received spoken, received written, received signed. OpenIssue: Description copied from Concept Domain of same name. Must be verified.
OID: | 2.16.840.1.113883.5.60 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System LanguageAbilityMode (A value representing the method of expression of the language. Example: Expressed spoken, expressed written, expressed signed, received spoken, received written, received signed. OpenIssue: Description copied from Concept Domain of same name. Must be verified.)

4.4.2.60.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | ESGN | Expressed signed
1 | ESP | Expressed spoken
1 | EWR | Expressed written
1 | RSGN | Received signed
1 | RSP | Received spoken
1 | RWR | Received written

4.4.2.61 HL7 v3 Code System LanguageAbilityProficiency

URL: https://hl7.org/fhir/R4/v3/LanguageAbilityProficiency/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-LanguageAbilityProficiency) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-LanguageAbilityProficiency
Version: | 2018-08-12
Name: | v3.LanguageAbilityProficiency
Title: | v3 Code System LanguageAbilityProficiency
Definition: | A value representing the level of proficiency in a language. Example: Excellent, good, fair, poor. OpenIssue: Description copied from Concept Domain of same name. Must be verified.
OID: | 2.16.840.1.113883.5.61 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System LanguageAbilityProficiency (A value representing the level of proficiency in a language. Example: Excellent, good, fair, poor. OpenIssue: Description copied from Concept Domain of same name. Must be verified.)

4.4.2.61.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | E | Excellent
1 | F | Fair
1 | G | Good
1 | P | Poor

4.4.2.63 HL7 v3 Code System LivingArrangement

URL: https://hl7.org/fhir/R4/v3/LivingArrangement/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-LivingArrangement) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-LivingArrangement
Version: | 2018-08-12
Name: | v3.LivingArrangement
Title: | v3 Code System LivingArrangement
Definition: | A code depicting the living arrangements of a person
OID: | 2.16.840.1.113883.5.63 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System LivingArrangement (A code depicting the living arrangements of a person)

4.4.2.63.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | HL | homeless | Definition: Living arrangements lacking a permanent residence.
2 | M | Nomadic | Nomadic
2 | T | Transient | Transient
1 | I | Institution | Institution
2 | CS | community shelter | Definition: A group living arrangement specifically for the care of those in need of temporary and crisis housing assistance. Examples include domestic violence shelters, shelters for displaced or homeless individuals, Salvation Army, Jesus House, etc. Community based services may be provided.
2 | G | Group Home | Group Home
2 | N | Nursing Home | Nursing Home
2 | X | Extended care facility | Extended care facility
1 | PR | private residence | Definition: A living arrangement within a private residence for single family.
2 | H | Independent Household | Independent Household
2 | R | Retirement Community | Retirement Community
2 | SL | supported living | Definition: Assisted living in a single family residence for persons with physical, behavioral, or functional health, or socio-economic challenges. There may or may not be on-site supervision but the housing is designed to assist the client with developing independent living skills. Community based services may be provided.

4.4.2.65 HL7 v3 Code System LocalMarkupIgnore

URL: https://hl7.org/fhir/R4/v3/LocalMarkupIgnore/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-LocalMarkupIgnore) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-LocalMarkupIgnore
Version: | 2018-08-12
Name: | v3.LocalMarkupIgnore
Title: | v3 Code System LocalMarkupIgnore
Definition: | Tells a receiver to ignore just the local markup tags (local_markup, local_header, local_attr) when value="markup", or to ignore the local markup tags and all contained content when value="all"
OID: | 2.16.840.1.113883.5.65 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System LocalMarkupIgnore (Tells a receiver to ignore just the local markup tags (local_markup, local_header, local_attr) when value="markup", or to ignore the local markup tags and all contained content when value="all")

4.4.2.65.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | all | all
1 | markup | markup

4.4.2.66 HL7 v3 Code System LocalRemoteControlState

URL: https://hl7.org/fhir/R4/v3/LocalRemoteControlState/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-LocalRemoteControlState) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-LocalRemoteControlState
Version: | 2018-08-12
Name: | v3.LocalRemoteControlState
Title: | v3 Code System LocalRemoteControlState
Definition: | **** MISSING DEFINITIONS ****
OID: | 2.16.840.1.113883.5.66 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System LocalRemoteControlState (**** MISSING DEFINITIONS ****)

4.4.2.66.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | L | Local | An equipment can either work autonomously ('Local' control state).
1 | R | Remote | An equipment can be controlled by another system, e.g., LAS computer ('Remote' control state).

4.4.2.67 HL7 v3 Code System MapRelationship

URL: https://hl7.org/fhir/R4/v3/MapRelationship/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-MapRelationship) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-MapRelationship
Version: | 2018-08-12
Name: | v3.MapRelationship
Title: | v3 Code System MapRelationship
Definition: | The closeness or quality of the mapping between the HL7 concept (as represented by the HL7 concept identifier) and the source coding system. The values are patterned after the similar relationships used in the UMLS Metathesaurus. Because the HL7 coding sy
OID: | 2.16.840.1.113883.5.67 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System MapRelationship (The closeness or quality of the mapping between the HL7 concept (as represented by the HL7 concept identifier) and the source coding system. The values are patterned after the similar relationships used in the UMLS Metathesaurus. Because the HL7 coding sy)

4.4.2.67.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | BT | Broader Than | The first concept is at a more abstract level than the second concept. For example, Hepatitis is broader than Hepatitis A, and endocrine disease is broader than Diabetes Mellitus. Broader than is the opposite of the narrower than relationship.
1 | E | Exact | The two concepts have identical meaning.
1 | NT | Narrower Than | The first concept is at a more detailed level than the second concept. For example, Pennicillin G is narrower than Pennicillin, and vellus hair is narrower than hair. Narrower than is the opposite of broader than.

4.4.2.81 HL7 v3 Code System ModifyIndicator

URL: https://hl7.org/fhir/R4/v3/ModifyIndicator/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-ModifyIndicator) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-ModifyIndicator
Version: | 2018-08-12
Name: | v3.ModifyIndicator
Title: | v3 Code System ModifyIndicator
Definition: | **** MISSING DEFINITIONS ****
OID: | 2.16.840.1.113883.5.81 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System ModifyIndicator (**** MISSING DEFINITIONS ****)

4.4.2.81.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | M | Modified subscription | Modified subscription to a query server.
1 | N | New subscription | New subscription to a query server.

4.4.2.83 HL7 v3 Code System ObservationInterpretation

URL: https://hl7.org/fhir/R4/v3/ObservationInterpretation/cs.html
Vocabulary Work Group | Maturity Level: N | Normative (from v4.0.0) | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-ObservationInterpretation) is defined as part of HL7 v3. Related FHIR content: Observation Interpretation Codes.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-ObservationInterpretation
Version: | 2018-08-12
Name: | v3.ObservationInterpretation
Title: | v3 Code System ObservationInterpretation
Definition: | One or more codes providing a rough qualitative interpretation of the observation, such as "normal" / "abnormal", "low" / "high", "better" / "worse", "resistant" / "susceptible", "expected" / "not expected". The value set is intended to be for ANY use where coded representation of an interpretation is needed.
OID: | 2.16.840.1.113883.5.83 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System ObservationInterpretation (One or more codes providing a rough qualitative interpretation of the observation, such as "normal" / "abnormal", "low" / "high", "better" / "worse", "resistant" / "susceptible", "expected" / "not expected". The value set is intended to be for ANY use where coded representation of an interpretation is needed.)

• ValueSet: Observation Interpretation Codes (A categorical assessment, providing a rough qualitative interpretation of the observation value, such as “normal”/ “abnormal”,”low” / “high”, “better” / “worse”, “susceptible” / “resistant”, “expected”/ “not expected”. The value set is intended to be for ANY use where coded representation of an interpretation is needed. Notes: This is being communicated in v2.x in OBX-8 (Observation Interpretation), in v3 in ObservationInterpretation (CWE) in R1 (Representative Realm) and in FHIR in Observation.interpretation. Historically these values come from the laboratory domain, and these codes are extensively used. The value set binding is extensible, so codes outside the value set that are needed for interpretation concepts (i.e. particular meanings) that are not included in the value set can be used, and these new codes may also be added to the value set and published in a future version.)

• ValueSet: Observation Interpretation Codes (A categorical assessment, providing a rough qualitative interpretation of the observation value, such as “normal”/ “abnormal”,”low” / “high”, “better” / “worse”, “susceptible” / “resistant”, “expected”/ “not expected”. The value set is intended to be for ANY use where coded representation of an interpretation is needed. Notes: This is being communicated in v2.x in OBX-8 (Observation Interpretation), in v3 in ObservationInterpretation (CWE) in R1 (Representative Realm) and in FHIR in Observation.interpretation. Historically these values come from the laboratory domain, and these codes are extensively used. The value set binding is extensible, so codes outside the value set that are needed for interpretation concepts (i.e. particular meanings) that are not included in the value set can be used, and these new codes may also be added to the value set and published in a future version.)

4.4.2.83.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | (_GeneticObservationInterpretation) Abstract | | Codes that specify interpretation of genetic analysis, such as "positive", "negative", "carrier", "responsive", etc.
2 | CAR | Carrier | The patient is considered as carrier based on the testing results. A carrier is an individual who carries an altered form of a gene which can lead to having a child or offspring in future generations with a genetic disorder.
2 | Carrier Deprecated | Carrier | The patient is considered as carrier based on the testing results. A carrier is an individual who carries an altered form of a gene which can lead to having a child or offspring in future generations with a genetic disorder. Deprecation Comment: This code is currently the same string as the print name for this concept and is inconsistent with the conventions being used for the other codes in the coding system, as it is a full word with initial capitalization, rather than an all upper case mnemonic. The recommendation from OO is to deprecate the code "Carrier" and to add "CAR" as the new active code representation for this concept.
1 | (_ObservationInterpretationChange) Abstract | | Interpretations of change of quantity and/or severity. At most one of B or W and one of U or D allowed.
2 | B | Better | The current result or observation value has improved compared to the previous result or observation value (the change is significant as defined in the respective test procedure). [Note: This can be applied to quantitative or qualitative observations.]
2 | D | Significant change down | The current result has decreased from the previous result for a quantitative observation (the change is significant as defined in the respective test procedure).

2 | U | Significant change up | The current result has increased from the previous result for a quantitative observation (the change is significant as defined in the respective test procedure).
2 | W | Worse | The current result or observation value has degraded compared to the previous result or observation value (the change is significant as defined in the respective test procedure). [Note: This can be applied to quantitative or qualitative observations.]
1 | (_ObservationInterpretationExceptions) Abstract | | Technical exceptions resulting in the inability to provide an interpretation. At most one allowed. Does not imply normality or severity.
2 | < | Off scale low | The result is below the minimum detection limit (the test procedure or equipment is the limiting factor). Synonyms: Below analytical limit, low off scale.
2 | > | Off scale high | The result is above the maximum quantifiable limit (the test procedure or equipment is the limiting factor). Synonyms: Above analytical limit, high off scale.
2 | AC Deprecated | Anti-complementary substances present | A valid result cannot be obtained for the specified component / analyte due to the presence of anti-complementary substances in the sample. Deprecation Comment: This code is being deprecated to match the status in V2 Table 0078 "Interpretation Codes.
2 | IE | Insufficient evidence | There is insufficient evidence that the species in question is a good target for therapy with the drug. A categorical interpretation is not possible. [Note: A MIC with "IE" and/or a comment may be reported (without an accompanying S, I or R-categorization).]
2 | QCF Deprecated | Quality control failure | A result cannot be considered valid for the specified component / analyte or organism due to failure in the quality control testing component. Deprecation Comment: This code is being deprecated to match the status in V2 Table 0078 "Interpretation Codes.
2 | TOX Deprecated | Cytotoxic substance present | A valid result cannot be obtained for the specified organism or cell line due to the presence of cytotoxic substances in the sample or culture. Deprecation Comment: This code is being deprecated to match the status in V2 Table 0078 "Interpretation Codes.
1 | (_ObservationInterpretationNormality) Abstract | | Interpretation of normality or degree of abnormality (including critical or "alert" level). Concepts in this category are mutually exclusive, i.e., at most one is allowed.
2 | A | Abnormal | The result or observation value is outside the reference range or expected norm (as defined for the respective test procedure). [Note: Typically applies to non-numeric results.]
3 | AA | Critical abnormal | The result or observation value is outside a reference range or expected norm at a level at which immediate action should be considered for patient safety (as defined for the respective test procedure). [Note: Typically applies to non-numeric results. Analogous to critical/panic limits for numeric results.]
4 | HH | Critical high | The result for a quantitative observation is above a reference level at which immediate action should be considered for patient safety (as defined for the respective test procedure). Synonym: Above upper panic limits.
4 | LL | Critical low | The result for a quantitative observation is below a reference level at which immediate action should be considered for patient safety (as defined for the respective test procedure). Synonym: Below lower panic limits.
3 | H | High | The result for a quantitative observation is above the upper limit of the reference range (as defined for the respective test procedure). Synonym: Above high normal
4 | H> Deprecated | Significantly high | A test result that is significantly higher than the reference (normal) or therapeutic interval, but has not reached the critically high value and might need special attention, as defined by the laboratory or the clinician.[Note: This level is situated between 'H' and 'HH'.] Deprecation Comment: The code 'H>' is being deprecated in order to align with the use of the code 'HU' for "Very high" in V2 Table 0078 "Interpretation Codes". [Note: The use of code 'H>' is non-preferred, as this code is deprecated and on track to be retired; use code 'HU' instead.
5 | HH
4 | HU | Significantly high | A test result that is significantly higher than the reference (normal) or therapeutic interval, but has not reached the critically high value and might need special attention, as defined by the laboratory or the clinician.
5 | HH
3 | L | Low | The result for a quantitative observation is below the lower limit of the reference range (as defined for the respective test procedure). Synonym: Below low normal
4 | L< Deprecated | Significantly low | A test result that is significantly lower than the reference (normal) or therapeutic interval, but has not reached the critically low value and might need special attention, as defined by the laboratory or the clinician.[Note: This level is situated between 'L' and 'LL'.] Deprecation Comment: The code 'L<' is being deprecated in order to align with the use of the code 'LU' for "Very low" in V2 Table 0078 "Interpretation Codes". [Note: The use of code 'L<' is non-preferred, as this code is deprecated and on track to be retired; use code 'LU' instead.
5 | LL
4 | LU | Significantly low | A test result that is significantly lower than the reference (normal) or therapeutic interval, but has not reached the critically low value and might need special attention, as defined by the laboratory or the clinician.
5 | LL
2 | N | Normal | The result or observation value is within the reference range or expected norm (as defined for the respective test procedure). [Note: Applies to numeric or non-numeric results.]
1 | (_ObservationInterpretationSusceptibility) Abstract | | Interpretations of anti-microbial susceptibility testing results (microbiology). At most one allowed.
2 | IE
2 | I | Intermediate | Bacterial strain inhibited in vitro by a concentration of an antimicrobial agent that is associated with uncertain therapeutic effect. Reference: CLSI (http://www.clsi.org/Content/NavigationMenu/Resources/HarmonizedTerminologyDatabase/Harmonized_Terminolo.htm) Projects: ISO 20776-1, ISO 20776-2 [Note 1: Bacterial strains are categorized as intermediate by applying the appropriate breakpoints in a defined phenotypic test system.] [Note 2: This class of susceptibility implies that an infection due to the isolate can be appropriately treated in body sites where the drugs are physiologically concentrated or when a high dosage of drug can be used.] [Note 3: This class also indicates a "buffer zone," to prevent small, uncontrolled, technical factors from causing major discrepancies in interpretations.] [Note 4: These breakpoints can be altered due to changes in circumstances (e.g., changes in commonly used drug dosages, emergence of new resistance mechanisms).]
2 | MS Deprecated | moderately susceptible | The patient is considered as carrier based on the testing results. A carrier is an individual who carries an altered form of a gene which can lead to having a child or offspring in future generations with a genetic disorder. Deprecation Comment: This antimicrobial susceptibility test interpretation concept is recommended by OO to be deprecated as it is no longer recommended for use in susceptibility testing by CLSI (reference CLSI document M100-S22; Vol. 32 No.3; CLSI Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Second Informational Supplement. Jan 2012).
2 | NCL | No CLSI defined breakpoint | Use when not enough clinical trial data published by the Clinical and Laboratory Standards Institutes (CLSI) is available to establish the breakpoints for susceptible / intermediate and resistant.

2 | NS | Non-susceptible | A category used for isolates for which only a susceptible interpretive criterion has been designated because of the absence or rare occurrence of resistant strains. Isolates that have MICs above or zone diameters below the value indicated for the susceptible breakpoint should be reported as non-susceptible. NOTE 1: An isolate that is interpreted as non-susceptible does not necessarily mean that the isolate has a resistance mechanism. It is possible that isolates with MICs above the susceptible breakpoint that lack resistance mechanisms may be encountered within the wild-type distribution subsequent to the time the susceptible-only breakpoint is set. NOTE 2: For strains yielding results in the "nonsusceptible" category, organism identification and antimicrobial susceptibility test results should be confirmed. Synonym: decreased susceptibility.

2 | R | Resistant | Bacterial strain inhibited in vitro by a concentration of an antimicrobial agent that is associated with a high likelihood of therapeutic failure. Reference: CLSI (http://www.clsi.org/Content/NavigationMenu/Resources/HarmonizedTerminologyDatabase/Harmonized_Terminolo.htm) Projects: ISO 20776-1, ISO 20776-2 [Note 1: Bacterial strains are categorized as resistant by applying the appropriate breakpoints in a defined phenotypic test system.] [Note 2: This breakpoint can be altered due to changes in circumstances (e.g., changes in commonly used drug dosages, emergence of new resistance mechanisms).]

3 | SYN-R | Synergy - resistant | A category for isolates where the bacteria (e.g. enterococci) are not susceptible in vitro to a combination therapy (e.g., high-level aminoglycoside and cell wall active agent). This is predictive that this combination therapy will not be effective. Usage Note: Since the use of penicillin or ampicillin alone often results in treatment failure of serious enterococcal or other bacterial infections, combination therapy is usually indicated to enhance bactericidal activity. The synergy between a cell wall active agent (such as penicillin, ampicillin, or vancomycin) and an aminoglycoside (such as gentamicin, kanamycin or streptomycin) is best predicted by screening for high-level bacterial resistance to the aminoglycoside. Open Issue: The print name of the code is very general and the description is very specific to a pair of classes of agents, which may lead to confusion of these concepts in the future should other synergies be found.

2 | S | Susceptible | Bacterial strain inhibited by in vitro concentration of an antimicrobial agent that is associated with a high likelihood of therapeutic success. Reference: CLSI (http://www.clsi.org/Content/NavigationMenu/Resources/HarmonizedTerminologyDatabase/Harmonized_Terminolo.htm) Synonym (earlier term): Sensitive Projects: ISO 20776-1, ISO 20776-2 [Note 1: Bacterial strains are categorized as susceptible by applying the appropriate breakpoints in a defined phenotypic system.] [Note 2: This breakpoint can be altered due to changes in circumstances (e.g., changes in commonly used drug dosages, emergence of new resistance mechanisms).]

3 | SDD | Susceptible-dose dependent | A category that includes isolates with antimicrobial agent minimum inhibitory concentrations (MICs) that approach usually attainable blood and tissue levels and for which response rates may be lower than for susceptible isolates. Reference: CLSI document M44-A2 2009 "Method for antifungal disk diffusion susceptibility testing of yeasts; approved guideline - second edition" - page 2.

3 | SYN-S | Synergy - susceptible | A category for isolates where the bacteria (e.g. enterococci) are susceptible in vitro to a combination therapy (e.g., high-level aminoglycoside and cell wall active agent). This is predictive that this combination therapy will be effective. Usage Note: Since the use of penicillin or ampicillin alone often results in treatment failure of serious enterococcal or other bacterial infections, combination therapy is usually indicated to enhance bactericidal activity. The synergy between a cell wall active agent (such as penicillin, ampicillin, or vancomycin) and an aminoglycoside (such as gentamicin, kanamycin or streptomycin) is best predicted by screening for high-level bacterial resistance to the aminoglycoside. Open Issue: The print name of the code is very general and the description is very specific to a pair of classes of agents, which may lead to confusion of these concepts in the future should other synergies be found.

2 | VS Deprecated | very susceptible | The patient is considered as carrier based on the testing results. A carrier is an individual who carries an altered form of a gene which can lead to having a child or offspring in future generations with a genetic disorder. Deprecation Comment: This antimicrobial susceptibility test interpretation concept is recommended by OO to be deprecated as it is no longer recommended for use in susceptibility testing by CLSI (reference CLSI document M100-S22; Vol. 32 No.3; CLSI Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Second Informational Supplement. Jan 2012).

1 | EX | outside threshold | The observation/test result is interpreted as being outside the inclusion range for a particular protocol within which the result is being reported. Example: A positive result on a Hepatitis screening test. Open Issue: EX, HX, LX: These three concepts do not seem to meet a clear need in the vocabulary, and their use in observation interpretation appears likely to be covered by other existing concepts (e.g., A, H, L). The only apparent significant difference is their reference to use in protocols for exclusion of study subjects. These concepts/codes were proposed by RCRIM for use in the CTLaboratory message. They were submitted and approved in the November 2005 Harmonization cycle in proposal "030103C_VOCAB_RCRIM_l_quade_RCRIM Obs Interp_20051028154455". However, this proposal was not fully implemented in the vocabulary. The proposal recommended creation of the x_ClinicalResearchExclusion domain in ObservationInterpretation with a value set including those three concepts/codes, but there is no subdomain of that name or equivalent with a binding to either of the value sets that contain these concepts/codes. Members of the OO WG have recently attempted to contact members of RCRIM regarding these concepts, both by email and at the recent WGM in Atlanta, without response. It is felt by OO that the best course of action to take at this time is to add this comprehensive Open Issue rather than deprecate these three concepts at this time, until further discussion is held.

2 | HX | above high threshold | The observation/test result is interpreted as being outside the inclusion range for a particular protocol within which the result is being reported. Example: A positive result on a Hepatitis screening test. Open Issue: EX, HX, LX: These three concepts do not seem to meet a clear need in the vocabulary, and their use in observation interpretation appears likely to be covered by other existing concepts (e.g., A, H, L). The only apparent significant difference is their reference to use in protocols for exclusion of study subjects. These concepts/codes were proposed by RCRIM for use in the CTLaboratory message. They were submitted and approved in the November 2005 Harmonization cycle in proposal "030103C_VOCAB_RCRIM_l_quade_RCRIM Obs Interp_20051028154455". However, this proposal was not fully implemented in the vocabulary. The proposal recommended creation of the x_ClinicalResearchExclusion domain in ObservationInterpretation with a value set including those three concepts/codes, but there is no subdomain of that name or equivalent with a binding to either of the value sets that contain these concepts/codes. Members of the OO WG have recently attempted to contact members of RCRIM regarding these concepts, both by email and at the recent WGM in Atlanta, without response. It is felt by OO that the best course of action to take at this time is to add this comprehensive Open Issue rather than deprecate these three concepts at this time, until further discussion is held.

2 | LX | below low threshold | The numeric observation/test result is interpreted as being below the low threshold value for a particular protocol within which the result is being reported. Example: A Total White Blood Cell Count falling below a protocol-defined threshold value of 3000/mm^3 Open Issue: EX, HX, LX: These three concepts do not seem to meet a clear need in the vocabulary, and their use in observation interpretation appears likely to be covered by other existing concepts (e.g., A, H, L). The only apparent significant difference is their reference to use in protocols for exclusion of study subjects. These concepts/codes were proposed by RCRIM for use in the CTLaboratory message. They were submitted and approved in the November 2005 Harmonization cycle in proposal "030103C_VOCAB_RCRIM_l_quade_RCRIM Obs Interp_20051028154455". However, this proposal was not fully implemented in the vocabulary. The proposal recommended creation of the x_ClinicalResearchExclusion domain in ObservationInterpretation with a value set including those three concepts/codes, but there

is no subdomain of that name or equivalent with a binding to either of the value sets that contain these concepts/codes. Members of the OO WG have recently attempted to contact members of RCRIM regarding these concepts, both by email and at the recent WGM in Atlanta, without response. It is felt by OO that the best course of action to take at this time is to add this comprehensive Open Issue rather than deprecate these three concepts at this time, until further discussion is held.

1 | HM Deprecated | Hold for Medical Review | Hold for Medical Review Usage Note: This code is not intended for use in V3 artifacts. It is included in the code system to maintain alignment with the V2 Table 0078 "Interpretation Codes."

1 | (ObservationInterpretationDetection) Abstract | | Interpretations of the presence or absence of a component / analyte or organism in a test or of a sign in a clinical observation. In keeping with laboratory data processing practice, these concepts provide a categorical interpretation of the "meaning" of the quantitative value for the same observation.

2 | IND | Indeterminate | The specified component / analyte, organism or clinical sign could neither be declared positive / negative nor detected / not detected by the performed test or procedure. Usage Note: For example, if the specimen was degraded, poorly processed, or was missing the required anatomic structures, then "indeterminate" (i.e. "cannot be determined") is the appropriate response, not "equivocal".

3 | E | Equivocal | The test or procedure was successfully performed, but the results are borderline and can neither be declared positive / negative nor detected / not detected according to the current established criteria.

2 | NEG | Negative | An absence finding of the specified component / analyte, organism or clinical sign based on the established threshold of the performed test or procedure. [Note: Negative does not necessarily imply the complete absence of the specified item.]

3 | ND | Not detected | The presence of the specified component / analyte, organism or clinical sign could not be determined within the limit of detection of the performed test or procedure.

2 | POS | Positive | A presence finding of the specified component / analyte, organism or clinical sign based on the established threshold of the performed test or procedure.

3 | DET | Detected | The measurement of the specified component / analyte, organism or clinical sign above the limit of detection of the performed test or procedure.

1 | (ObservationInterpretationExpectation) Abstract | | Interpretation of the observed result taking into account additional information (contraindicators) about the patient's situation. Concepts in this category are mutually exclusive, i.e., at most one is allowed.

2 | EXP | Expected | This result has been evaluated in light of known contraindicators. Once those contraindicators have been taken into account the result is determined to be "Expected" (e.g., presence of drugs in a patient that is taking prescription medication for pain management).

2 | UNE | Unexpected | This result has been evaluated in light of known contraindicators. Once those contraindicators have been taken into account the result is determined to be "Unexpected" (e.g., presence of non-prescribed drugs in a patient that is taking prescription medication for pain management).

1 | OBX Deprecated | Interpretation qualifiers in separate OBX segments | Interpretation qualifiers in separate OBX segments Usage Note: This code is not intended for use in V3 artifacts. It is included in the code system to maintain alignment with the V2 Table 0078 "Interpretation Codes."

1 | (ReactivityObservationInterpretation) Abstract | | Interpretations of the presence and level of reactivity of the specified component / analyte with the reagent in the performed laboratory test.

2 | NR | Non-reactive | An absence finding used to indicate that the specified component / analyte did not react measurably with the reagent.

2 | RR | Reactive | A presence finding used to indicate that the specified component / analyte reacted with the reagent above the reliably measurable limit of the performed test.

3 | WR | Weakly reactive | A weighted presence finding used to indicate that the specified component / analyte reacted with the reagent, but below the reliably measurable limit of the performed test.

4.4.2.84 HL7 v3 Code System ObservationMethod

URL: https://hl7.org/fhir/R4/v3/ObservationMethod/cs.html

Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-ObservationMethod) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-ObservationMethod

Version: | 2018-08-12

Name: | v3.ObservationMethod

Title: | v3 Code System ObservationMethod

Definition: | A code that provides additional detail about the means or technique used to ascertain the observation. Examples: Blood pressure measurement method: arterial puncture vs. sphygmomanometer (Riva-Rocci), sitting vs. supine position, etc.

OpenIssue: Description copied from Concept Domain of same name. Must be verified. Note that the Domain has a full discussion about use of the attribute and constraining that is not appropriate for the code system description. Needs to be improved.

OID: | 2.16.840.1.113883.5.84 (for OID based terminology systems)

Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: V3 Value SetVerificationMethod (No Description Provided)

• ValueSet: v3 Code System ObservationMethod (A code that provides additional detail about the means or technique used to ascertain the observation. Examples: Blood pressure measurement method: arterial puncture vs. sphygmomanometer (Riva-Rocci), sitting vs. supine position, etc. OpenIssue: Description copied from Concept Domain of same name. Must be verified. Note that the Domain has a full discussion about use of the attribute and constraining that is not appropriate for the code system description. Needs to be improved.)

4.4.2.84.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | (_DecisionObservationMethod) Abstract | | Provides codes for decision methods, initially for assessing the causality of events.
2 | ALGM | algorithm | Reaching a decision through the application of an algorithm designed to weigh the different factors involved.
3 | BYCL | bayesian calculation | Reaching a decision through the use of Bayesian statistical analysis.
2 | GINT | global introspection | Reaching a decision by consideration of the totality of factors involved in order to reach a judgement.
1 | (_GeneticObservationMethod) Abstract | | A code that provides additional detail about the means or technique used to ascertain the genetic analysis. Example, PCR, Micro Array
2 | PCR | PCR | Description: Polymerase Chain Reaction
1 | (_ObservationMethodAggregate) Abstract | | Provides additional detail about the aggregation methods used to compute the aggregated values for an observation. This is an abstract code.
2 | AVERAGE | average | Average of non-null values in the referenced set of values
2 | COUNT | count | Count of non-null values in the referenced set of values
2 | MAX | maxima | Largest of all non-null values in the referenced set of values.
2 | MEDIAN | median | The median of all non-null values in the referenced set of values.
2 | MIN | minima | Smallest of all non-null values in the referenced set of values.
2 | MODE | mode | The most common value of all non-null values in the referenced set of values.
2 | STDEV.P | population standard deviation | Standard Deviation of the values in the referenced set of values, computed over the population.
2 | STDEV.S | sample standard deviation | Standard Deviation of the values in the referenced set of values, computed over a sample of the population.
2 | SUM | sum | Sum of non-null values in the referenced set of values
2 | VARIANCE.P | population variance | Variance of the values in the referenced set of values, computed over the population.
2 | VARIANCE.S | sample variance | Variance of the values in the referenced set of values, computed over a sample of the population.
1 | (_VerificationMethod) Abstract
2 | VDOC | document verification | Verification by means of document. Example: Fax, letter, attachment to e-mail.
2 | VREG | registry verification | verification by means of a response to an electronic query Example: query message to a Covered Party registry application or Coverage Administrator.
2 | VTOKEN | electronic token verification | Verification by means of electronic token. Example: smartcard, magnetic swipe card, RFID device.
2 | VVOICE | voice-based verification | Verification by means of voice. Example: By speaking with or calling the Coverage Administrator or Covered Party
1 | 0001 | Complement fixation | Complement fixation
1 | 0002 | Computed axial tomography | Computed axial tomography
1 | 0003 | HLAR agar test | Susceptibility, High Level Aminoglycoside Resistance agar test
1 | 0004 | Macroscopic observation | Visual, Macroscopic observation
1 | 0005 | Magnetic resonance | Computed, Magnetic resonance
1 | 0006 | Morphometry | Computed, Morphometry
1 | 0007 | Positron emission tomography | Computed, Positron emission tomography
1 | 0008 | SAMHSA confirmation | SAMHSA drug assay confirmation
1 | 0009 | SAMHSA screening | SAMHSA drug assay screening
1 | 0010 | Serum Neutralization | Serum Neutralization
1 | 0011 | Titration | Titration
1 | 0012 | Ultrasound | Ultrasound
1 | 0013 | X-ray crystallography | X-ray crystallography
1 | 0014 | Agglutination | Agglutination
1 | 0015 | Buffered acidified plate agglutination | Agglutination, Buffered acidified plate
1 | 0016 | Card agglutination | Agglutination, Card
1 | 0017 | Hemagglutination | Agglutination, Hemagglutination
1 | 0018 | Hemagglutination inhibition | Agglutination, Hemagglutination inhibition
1 | 0019 | Latex agglutination | Agglutination, Latex
1 | 0020 | Plate agglutination | Agglutination, Plate
1 | 0021 | Rapid agglutination | Agglutination, Rapid Plate
1 | 0022 | RBC agglutination | Agglutination, RBC
1 | 0023 | Rivanol agglutination | Agglutination, Rivanol
1 | 0024 | Tube agglutination | Agglutination, Tube
1 | 0025 | Bioassay | Bioassay
1 | 0026 | Animal Inoculation | Bioassay, Animal Inoculation
1 | 0027 | Cytotoxicity | Bioassay, Cytotoxicity
1 | 0028 | Embryo infective dose 50 | Bioassay, Embryo Infective Dose 50
1 | 0029 | Embryo lethal dose 50 | Bioassay, Embryo Lethal Dose 50
1 | 0030 | Mouse intercerebral inoculation | Bioassay, Mouse intercerebral inoculation
1 | 0031 | Bioassay, qualitative | Bioassay, qualitative
1 | 0032 | Bioassay, quantitative | Bioassay, quantitative
1 | 0033 | Chemical method | Chemical
1 | 0034 | Differential light absorption chemical test | Chemical, Differential light absorption
1 | 0035 | Dipstick | Chemical, Dipstick
1 | 0036 | Dipstick colorimetric laboratory test | Chemical, Dipstick colorimetric laboratory test
1 | 0037 | Test strip | Chemical, Test strip
1 | 0038 | Chromatography | Chromatography

1 | 0039 | Affinity chromatography | Chromatography, Affinity
1 | 0040 | Gas liquid chromatography | Chromatography, Gas liquid
1 | 0041 | High performance liquid chromatography | Chromatography, High performance liquid
1 | 0042 | Liquid Chromatography | Chromatography, Liquid
1 | 0043 | Protein A affinity chromatography | Chromatography, Protein A affinity
1 | 0044 | Coagulation | Coagulation
1 | 0045 | Tilt tube coagulation time | Coagulation, Tilt tube
1 | 0046 | Tilt tube reptilase induced coagulation | Coagulation, Tilt tube reptilase induced
1 | 0047 | Automated count | Count, Automated
1 | 0048 | Manual cell count | Count, Manual
1 | 0049 | Platelet count, Rees-Ecker | Count, Platelet, Rees-Ecker
1 | 0050 | Aerobic Culture | Culture, Aerobic
1 | 0051 | Anaerobic Culture | Culture, Anaerobic
1 | 0052 | Chicken embryo culture | Culture, Chicken Embryo
1 | 0053 | Delayed secondary enrichment | Culture, Delayed secondary enrichment
1 | 0054 | Microaerophilic Culture | Culture, Microaerophilic
1 | 0055 | Quantitative microbial culture, cup | Culture, Quantitative microbial, cup
1 | 0056 | Quantitative microbial culture, droplet | Culture, Quantitative microbial, droplet
1 | 0057 | Quantitative microbial culture, filter paper | Culture, Quantitative microbial, filter paper
1 | 0058 | Quantitative microbial culture, pad culture | Culture, Quantitative microbial, pad
1 | 0059 | Quantitative microbial culture, pour plate | Culture, Quantitative microbial, pour plate
1 | 0060 | Quantitative microbial culture, surface streak | Culture, Quantitative microbial, surface streak
1 | 0061 | Somatic Cell culture | Culture, Somatic Cell
1 | 0062 | Agar diffusion | Diffusion, Agar
1 | 0063 | Agar Gel Immunodiffusion | Diffusion, Agar Gel Immunodiffusion
1 | 0064 | Electrophoresis | Electrophoresis
1 | 0065 | Agaorse gel electrophoresis | Electrophoresis, Agaorse gel
1 | 0066 | Electrophoresis, citrate agar | Electrophoresis, citrate agar
1 | 0067 | Immunoelectrophoresis | Electrophoresis, Immuno
1 | 0068 | Polyacrylamide gel electrophoresis | Electrophoresis, Polyacrylamide gel
1 | 0069 | Starch gel electrophoresis | Electrophoresis, Starch gel
1 | 0070 | ELISA | ELISA
1 | 0071 | ELISA, antigen capture | ELISA, antigen capture
1 | 0072 | ELISA, avidin biotin peroxidase complex | ELISA, avidin biotin peroxidase complex
1 | 0073 | Kinetic ELISA | ELISA, Kinetic
1 | 0074 | ELISA, peroxidase-antiperoxidase | ELISA, peroxidase-antiperoxidase
1 | 0075 | API 20 Strep | Identification, API 20 Strep
1 | 0076 | API 20A | Identification, API 20A
1 | 0077 | API 20C AUX | Identification, API 20C AUX
1 | 0078 | API 20E | Identification, API 20E
1 | 0079 | API 20NE | Identification, API 20NE
1 | 0080 | API 50 CH | Identification, API 50 CH
1 | 0081 | API An-IDENT | Identification, API An-IDENT
1 | 0082 | API Coryne | Identification, API Coryne
1 | 0083 | API Rapid 20E | Identification, API Rapid 20E
1 | 0084 | API Staph | Identification, API Staph
1 | 0085 | API ZYM | Identification, API ZYM
1 | 0086 | Bacterial identification | Identification, Bacterial
1 | 0087 | mini VIDAS | Identification, mini VIDAS
1 | 0088 | Phage susceptibility typing | Identification, Phage susceptibility typing
1 | 0089 | Quad-FERM+ | Identification, Quad-FERM+
1 | 0090 | RAPIDEC Staph | Identification, RAPIDEC Staph
1 | 0091 | Staphaurex | Identification, Staphaurex
1 | 0092 | VIDAS | Identification, VIDAS
1 | 0093 | Vitek | Identification, Vitek
1 | 0094 | VITEK 2 | Identification, VITEK 2
1 | 0095 | Immune stain | Immune stain
1 | 0096 | Immunofluorescent antibody, direct | Immune stain, Immunofluorescent antibody, direct
1 | 0097 | Immunofluorescent antibody, indirect | Immune stain, Immunofluorescent antibody, indirect
1 | 0098 | Immunoperoxidase, Avidin-Biotin Complex | Immune stain, Immunoperoxidase, Avidin-Biotin Complex
1 | 0099 | Immunoperoxidase, Peroxidase anti-peroxidase complex | Immune stain, Immunoperoxidase, Peroxidase anti-peroxidase complex
1 | 0100 | Immunoperoxidase, Protein A-peroxidase complex | Immune stain, Immunoperoxidase, Protein A-peroxidase complex
1 | 0101 | Immunoassay | Immunoassay
1 | 0102 | Immunoassay, qualitative, multiple step | Immunoassay, qualitative, multiple step
1 | 0103 | Immunoassay, qualitative, single step | Immunoassay, qualitative, single step
1 | 0104 | Radioimmunoassay | Immunoassay, Radioimmunoassay
1 | 0105 | Immunoassay, semi-quantitative, multiple step | Immunoassay, semi-quantitative, multiple step
1 | 0106 | Immunoassay, semi-quantitative, single step | Immunoassay, semi-quantitative, single step
1 | 0107 | Microscopy | Microscopy
1 | 0108 | Darkfield microscopy | Microscopy, Darkfield
1 | 0109 | Electron microscopy | Microscopy, Electron

1 | 0110 | Electron microscopy tomography | Microscopy, Electron microscopy tomography
1 | 0111 | Electron microscopy, negative stain | Microscopy, Electron, negative stain
1 | 0112 | Electron microscopy, thick section | Microscopy, Electron, thick section transmission
1 | 0113 | Electron microscopy, thin section | Microscopy, Electron, thin section transmission
1 | 0114 | Microscopy, Light | Microscopy, Light
1 | 0115 | Polarizing light microscopy | Microscopy, Polarized light
1 | 0116 | Scanning electron microscopy | Microscopy, Scanning electron
1 | 0117 | Transmission electron microscopy | Microscopy, Transmission electron
1 | 0118 | Transparent tape direct examination | Microscopy, Transparent tape direct examination
1 | 0119 | 3 Self-Sustaining Sequence Replication | Molecular, 3 Self-Sustaining Sequence Replication
1 | 0120 | Branched Chain DNA | Molecular, Branched Chain DNA
1 | 0121 | Hybridization Protection Assay | Molecular, Hybridization Protection Assay
1 | 0122 | Immune blot | Molecular, Immune blot
1 | 0123 | In-situ hybridization | Molecular, In-situ hybridization
1 | 0124 | Ligase Chain Reaction | Molecular, Ligase Chain Reaction
1 | 0125 | Ligation Activated Transcription | Molecular, Ligation Activated Transcription
1 | 0126 | Nucleic Acid Probe | Molecular, Nucleic Acid Probe
1 | 0128 | Nucleic acid probe with amplification | Molecular, Nucleic acid probe with amplification Rationale: Duplicate of code 0126. Use code 0126 instead.
1 | 0129 | Nucleic acid probe with target amplification | Molecular, Nucleic acid probe with target amplification
1 | 0130 | Nucleic acid reverse transcription | Molecular, Nucleic acid reverse transcription
1 | 0131 | Nucleic Acid Sequence Based Analysis | Molecular, Nucleic Acid Sequence Based Analysis
1 | 0132 | Polymerase chain reaction | Molecular, Polymerase chain reaction
1 | 0133 | Q-Beta Replicase or probe amplification category method | Molecular, Q-Beta Replicase or probe amplification category method
1 | 0134 | Restriction Fragment Length Polymorphism | Molecular, Restriction Fragment Length Polymorphism
1 | 0135 | Southern Blot | Molecular, Southern Blot
1 | 0136 | Strand Displacement Amplification | Molecular, Strand Displacement Amplification
1 | 0137 | Transcription Mediated Amplification | Molecular, Transcription Mediated Amplification
1 | 0138 | Western Blot | Molecular, Western Blot
1 | 0139 | Flocculation | Precipitation, Flocculation
1 | 0140 | Immune precipitation | Precipitation, Immune precipitation
1 | 0141 | Milk ring test | Precipitation, Milk ring test
1 | 0142 | Precipitin | Precipitation, Precipitin
1 | 0143 | Acid fast stain | Stain, Acid fast
1 | 0144 | Acid fast stain, fluorochrome | Stain, Acid fast, fluorochrome
1 | 0145 | Acid fast stain, Kinyoun's cold carbolfuchsin | Stain, Acid fast, Kinyoun's cold carbolfuchsin
1 | 0146 | Acid fast stain, Ziehl-Neelsen | Stain, Acid fast, Ziehl-Neelsen
1 | 0147 | Acid phosphatase stain | Stain, Acid phosphatase
1 | 0148 | Acridine orange stain | Stain, Acridine orange
1 | 0149 | Active brilliant orange KH stain | Stain, Active brilliant orange KH
1 | 0150 | Alazarin red S stain | Stain, Alazarin red S
1 | 0151 | Alcian blue stain | Stain, Alcian blue
1 | 0152 | Alcian blue with Periodic acid Schiff stain | Stain, Alcian blue with Periodic acid Schiff
1 | 0153 | Argentaffin stain | Stain, Argentaffin
1 | 0154 | Argentaffin silver stain | Stain, Argentaffin silver
1 | 0155 | Azure-eosin stain | Stain, Azure-eosin
1 | 0156 | Basic Fuschin stain | Stain, Basic Fuschin
1 | 0157 | Bennhold stain | Stain, Bennhold
1 | 0158 | Bennhold's Congo red stain | Stain, Bennhold's Congo red
1 | 0159 | Bielschowsky stain | Stain, Bielschowsky
1 | 0160 | Bielschowsky's silver stain | Stain, Bielschowsky's silver
1 | 0161 | Bleach stain | Stain, Bleach
1 | 0162 | Bodian stain | Stain, Bodian
1 | 0163 | Brown-Brenn stain | Stain, Brown-Brenn
1 | 0164 | Butyrate-esterase stain | Stain, Butyrate-esterase
1 | 0165 | Calcofluor white fluorescent stain | Stain, Calcofluor white fluorescent
1 | 0166 | Carbol-fuchsin stain | Stain, Carbol-fuchsin
1 | 0167 | Carmine stain | Stain, Carmine
1 | 0168 | Churukian-Schenk stain | Stain, Churukian-Schenk
1 | 0169 | Congo red stain | Stain, Congo red
1 | 0170 | Cresyl echt violet stain | Stain, Cresyl echt violet
1 | 0171 | Crystal violet stain | Stain, Crystal violet
1 | 0172 | De Galantha stain | Stain, De Galantha
1 | 0173 | Dieterle silver impregnation stain | Stain, Dieterle silver impregnation
1 | 0174 | Fite-Farco stain | Stain, Fite-Farco
1 | 0175 | Fontana-Masson silver stain | Stain, Fontana-Masson silver
1 | 0176 | Fouchet stain | Stain, Fouchet
1 | 0177 | Gomori stain | Stain, Gomori
1 | 0178 | Gomori methenamine silver stain | Stain, Gomori methenamine silver
1 | 0179 | Gomori-Wheatly trichrome stain | Stain, Gomori-Wheatly trichrome
1 | 0180 | Gridley stain | Stain, Gridley

1 | 0181 | Grimelius silver stain | Stain, Grimelius silver
1 | 0182 | Grocott stain | Stain, Grocott
1 | 0183 | Grocott methenamine silver stain | Stain, Grocott methenamine silver
1 | 0184 | Hale's colloidal ferric oxide stain | Stain, Hale's colloidal ferric oxide
1 | 0185 | Hale's colloidal iron stain | Stain, Hale's colloidal iron
1 | 0186 | Hansel stain | Stain, Hansel
1 | 0187 | Harris regressive hematoxylin and eosin stain | Stain, Harris regressive hematoxylin and eosin
1 | 0188 | Hematoxylin and eosin stain | Stain, Hematoxylin and eosin
1 | 0189 | Highman stain | Stain, Highman
1 | 0190 | Holzer stain | Stain, Holzer
1 | 0191 | Iron hematoxylin stain | Stain, Iron hematoxylin
1 | 0192 | Jones stain | Stain, Jones
1 | 0193 | Jones methenamine silver stain | Stain, Jones methenamine silver
1 | 0194 | Kossa stain | Stain, Kossa
1 | 0195 | Lawson-Van Gieson stain | Stain, Lawson-Van Gieson
1 | 0196 | Loeffler methylene blue stain | Stain, Loeffler methylene blue
1 | 0197 | Luxol fast blue with cresyl violet stain | Stain, Luxol fast blue with cresyl violet
1 | 0198 | Luxol fast blue with Periodic acid-Schiff stain | Stain, Luxol fast blue with Periodic acid-Schiff
1 | 0199 | MacNeal's tetrachrome blood stain | Stain, MacNeal's tetrachrome blood
1 | 0200 | Mallory-Heidenhain stain | Stain, Mallory-Heidenhain
1 | 0201 | Masson trichrome stain | Stain, Masson trichrome
1 | 0202 | Mayer mucicarmine stain | Stain, Mayer mucicarmine
1 | 0203 | Mayers progressive hematoxylin and eosin stain | Stain, Mayers progressive hematoxylin and eosin
1 | 0204 | May-Grunwald Giemsa stain | Stain, May-Grunwald Giemsa
1 | 0205 | Methyl green stain | Stain, Methyl green
1 | 0206 | Methyl green pyronin stain | Stain, Methyl green pyronin
1 | 0207 | Modified Gomori-Wheatly trichrome stain | Stain, Modified Gomori-Wheatly trichrome
1 | 0208 | Modified Masson trichrome stain | Stain, Modified Masson trichrome
1 | 0209 | Modified trichrome stain | Stain, Modified trichrome
1 | 0210 | Movat pentachrome stain | Stain, Movat pentachrome
1 | 0211 | Mucicarmine stain | Stain, Mucicarmine
1 | 0212 | Neutral red stain | Stain, Neutral red
1 | 0213 | Night blue stain | Stain, Night blue
1 | 0214 | Non-specific esterase stain | Stain, Non-specific esterase
1 | 0215 | Oil red-O stain | Stain, Oil red-O
1 | 0216 | Orcein stain | Stain, Orcein
1 | 0217 | Perls' stain | Stain, Perls'
1 | 0218 | Phosphotungstic acid-hematoxylin stain | Stain, Phosphotungstic acid-hematoxylin
1 | 0219 | Potassium ferrocyanide stain | Stain, Potassium ferrocyanide
1 | 0220 | Prussian blue stain | Stain, Prussian blue
1 | 0221 | Putchler modified Bennhold stain | Stain, Putchler modified Bennhold
1 | 0222 | Quinacrine fluorescent stain | Stain, Quinacrine fluorescent
1 | 0223 | Reticulin stain | Stain, Reticulin
1 | 0224 | Rhodamine stain | Stain, Rhodamine
1 | 0225 | Safranin stain | Stain, Safranin
1 | 0226 | Schmorl stain | Stain, Schmorl
1 | 0227 | Seiver-Munger stain | Stain, Seiver-Munger
1 | 0228 | Silver stain | Stain, Silver
1 | 0229 | Specific esterase stain | Stain, Specific esterase
1 | 0230 | Steiner silver stain | Stain, Steiner silver
1 | 0231 | Sudan III stain | Stain, Sudan III
1 | 0232 | Sudan IVI stain | Stain, Sudan IVI
1 | 0233 | Sulfated alcian blue stain | Stain, Sulfated alcian blue
1 | 0234 | Supravital stain | Stain, Supravital
1 | 0235 | Thioflavine-S stain | Stain, Thioflavine-S
1 | 0236 | Three micron Giemsa stain | Stain, Three micron Giemsa
1 | 0237 | Vassar-Culling stain | Stain, Vassar-Culling
1 | 0238 | Vital Stain | Stain, Vital
1 | 0239 | von Kossa stain | Stain, von Kossa
1 | 0243 | Minimum bactericidal concentration test, macrodilution | Susceptibility, Minimum bactericidal concentration, macrodilution
1 | 0244 | Minimum bactericidal concentration test, microdilution | Susceptibility, Minimum bactericidal concentration, microdilution
1 | 0247 | Turbidometric | Turbidometric
1 | 0248 | Refractometric | Turbidometric, Refractometric
1 | 0249 | Thin layer chromatography (TLC) | Chromatography, Thin Layer
1 | 0250 | EMIT | Immunoassay, enzyme-multiplied technique (EMIT)
1 | 0251 | Flow cytometry (FC) | Flow Cytometry
1 | 0252 | Radial immunodiffusion (RID) | Radial Immunodiffusion
1 | 0253 | Fluorescence polarization immunoassay (FPIA) | Immunoassay, Fluorescence Polarization
1 | 0254 | Immunofixation electrophoresis (IFE) | Electrophoresis, Immunofixation
1 | 0255 | Equilibrium dialysis | Dialysis, Direct Equilibrium

1 | 0256 | Kleihauer-Betke acid elution | Acid Elution, Kleihauer-Betke Method
1 | 0257 | Anti-complement immunofluorescence (ACIF) | Immunofluorescence, Anti-Complement
1 | 0258 | GC/MS | Gas Chromatography/Mass Spectroscopy
1 | 0259 | Nephelometry | Light Scatter, Nephelometry
1 | 0260 | IgE immunoassay antibody | Immunoassay, IgE Antibody Test
1 | 0261 | Lymphocyte Microcytotoxicity Assay | Lymphocyte Microcytotoxicity Assay
1 | 0262 | Spectrophotometry | Spectrophotometry
1 | 0263 | Atomic absorption spectrophotometry (AAS) | Spectrophotometry, Atomic Absorption
1 | 0264 | Ion selective electrode (ISE) | Electrochemical, Ion Selective Electrode
1 | 0265 | Gas chromatography (GC) | Chromatography, Gas
1 | 0266 | Isoelectric focusing (IEF) | Isoelectric Focusing
1 | 0267 | Immunochemiluminescence | Immunoassay, Chemiluminescent
1 | 0268 | Microparticle enzyme immunoassay (MEIA) | Immunoassay, Microparticle Enzyme
1 | 0269 | ICP/MS | Inductively-Coupled Plasma/Mass Spectrometry
1 | 0270 | Immunoradiometric assay (IRMA) | Immunoassay, Immunoradiometric Assay
1 | 0271 | Photo optical clot detection | Coagulation, Photo Optical Clot Detection
1 | 0280 | Susceptibility Testing | Test methods designed to determine a microorganismaTMs susceptibility to being killed by an antibiotic.
2 | 0240 | Antibiotic sensitivity, disk | Susceptibility, Antibiotic sensitivity, disk
2 | 0241 | BACTEC susceptibility test | Susceptibility, BACTEC susceptibility test
2 | 0242 | Disk dilution | Susceptibility, Disk dilution
2 | 0272 | Minimum Inhibitory Concentration | Testing to measure the minimum concentration of the antibacterial agent in a given culture medium below which bacterial growth is not inhibited.
3 | 0245 | Minimum Inhibitory Concentration, macrodilution | Susceptibility, Minimum Inhibitory concentration, macrodilution
3 | 0246 | Minimum Inhibitory Concentration, microdilution | Susceptibility, Minimum Inhibitory concentration, microdilution
2 | 0273 | Viral Genotype Susceptibility
2 | 0274 | Viral Phenotype Susceptibility
2 | 0275 | Gradient Strip
2 | 0275a | Minimum Lethal Concentration (MLC)
2 | 0276 | Slow Mycobacteria Susceptibility | Testing to measure the minimum concentration of the antibacterial agent in a given culture medium below which bacterial growth is not inhibited.
2 | 0277 | Serum bactericidal titer
2 | 0278 | Agar screen
2 | 0279 | Disk induction
1 | 0127 | Nucleic acid probe | Molecular, Nucleic acid probe

4.4.2.85 HL7 v3 Code System orderableDrugForm

URL: https://hl7.org/fhir/R4/v3/orderableDrugForm/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-orderableDrugForm) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-orderableDrugForm
Version: | 2018-08-12
Name: | v3.orderableDrugForm
Title: | v3 Code System orderableDrugForm
Definition: | OpenIssue: Missing description.
OID: | 2.16.840.1.113883.5.85 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System orderableDrugForm (OpenIssue: Missing description.)

4.4.2.85.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | (_AdministrableDrugForm) Abstract
2 | APPFUL | Applicatorful
2 | DROP | Drops
3 | NDROP | Nasal Drops
3 | OPDROP | Ophthalmic Drops
3 | ORDROP | Oral Drops
3 | OTDROP | Otic Drops
2 | PUFF | Puff
2 | SCOOP | Scoops
2 | SPRY | Sprays
1 | (_DispensableDrugForm) Abstract
2 | (_GasDrugForm) Abstract | | Any elastic aeriform fluid in which the molecules are separated from one another and have free paths.

3 | GASINHL | Gas for Inhalation
2 | (_GasLiquidMixture) Abstract
3 | AER | Aerosol
4 | BAINHL | Breath Activated Inhaler
4 | INHLSOL | Inhalant Solution
4 | MDINHL | Metered Dose Inhaler
4 | NASSPRY | Nasal Spray
3 | DERMSPRY | Dermal Spray
3 | FOAM | Foam
4 | FOAMAPL | Foam with Applicator
4 | RECFORM | Rectal foam
4 | VAGFOAM | Vaginal foam
5 | VAGFOAMAPL | Vaginal foam with applicator
3 | RECSPRY | Rectal Spray
3 | VAGSPRY | Vaginal Spray
2 | (_GasSolidSpray) Abstract
3 | INHL | Inhalant
4 | BAINHLPWD | Breath Activated Powder Inhaler
4 | INHLPWD | Inhalant Powder
4 | MDINHLPWD | Metered Dose Powder Inhaler
4 | NASINHL | Nasal Inhalant
4 | ORINHL | Oral Inhalant
3 | PWDSPRY | Powder Spray
3 | SPRYADAPT | Spray with Adaptor
2 | (_Liquid) Abstract | | A state of substance that is an intermediate one entered into as matter goes from solid to gas; liquids are also intermediate in that they have neither the orderliness of a crystal nor the randomness of a gas. (Note: This term should not be used to describe solutions, only pure chemicals in their liquid state.)
3 | LIQCLN | Liquid Cleanser
4 | LIQSOAP | Medicated Liquid Soap
4 | SHMP | Shampoo | A liquid soap or detergent used to clean the hair and scalp and is often used as a vehicle for dermatologic agents.
3 | OIL | Oil | An unctuous, combustible substance which is liquid, or easily liquefiable, on warming, and is soluble in ether but insoluble in water. Such substances, depending on their origin, are classified as animal, mineral, or vegetable oils.
4 | TOPOIL | Topical Oil
3 | SOL | Solution | A liquid preparation that contains one or more chemical substances dissolved, i.e., molecularly dispersed, in a suitable solvent or mixture of mutually miscible solvents.
4 | DROP
4 | IPSOL | Intraperitoneal Solution
4 | IRSOL | Irrigation Solution | A sterile solution intended to bathe or flush open wounds or body cavities; they're used topically, never parenterally.
5 | DOUCHE | Douche | A liquid preparation, intended for the irrigative cleansing of the vagina, that is prepared from powders, liquid solutions, or liquid concentrates and contains one or more chemical substances dissolved in a suitable solvent or mutually miscible solvents.
5 | ENEMA | Enema | A rectal preparation for therapeutic, diagnostic, or nutritive purposes.
5 | OPIRSOL | Ophthalmic Irrigation Solution
4 | IVSOL | Intravenous Solution
4 | ORALSOL | Oral Solution
5 | ORDROP
5 | ELIXIR | Elixir | A clear, pleasantly flavored, sweetened hydroalcoholic liquid containing dissolved medicinal agents; it is intended for oral use.
5 | RINSE | Mouthwash/Rinse | An aqueous solution which is most often used for its deodorant, refreshing, or antiseptic effect.
5 | SYRUP | Syrup | An oral solution containing high concentrations of sucrose or other sugars; the term has also been used to include any other liquid dosage form prepared in a sweet and viscid vehicle, including oral suspensions.
4 | RECSOL | Rectal Solution
4 | TOPSOL | Topical Solution
5 | LIN | Liniment | A solution or mixture of various substances in oil, alcoholic solutions of soap, or emulsions intended for external application.
5 | MUCTOPSOL | Mucous Membrane Topical Solution
5 | TINC | Tincture
2 | (_LiquidLiquidEmulsion) Abstract | | A two-phase system in which one liquid is dispersed throughout another liquid in the form of small droplets.
3 | CRM | Cream | A semisolid dosage form containing one or more drug substances dissolved or dispersed in a suitable base; more recently, the term has been restricted to products consisting of oil-in-water emulsions or aqueous microcrystalline dispersions of long chain fatty acids or alcohols that are water washable and more cosmetically and aesthetically acceptable.
4 | NASCRM | Nasal Cream
4 | OPCRM | Ophthalmic Cream
4 | ORCRM | Oral Cream
4 | OTCRM | Otic Cream
4 | RECCRM | Rectal Cream
4 | TOPCRM | Topical Cream
4 | VAGCRM | Vaginal Cream
5 | VAGCRMAPL | Vaginal Cream with Applicator
3 | LTN | Lotion | The term "lotion" has been used to categorize many topical suspensions, solutions and emulsions intended for application to the skin.

4 | TOPLTN | Topical Lotion
3 | OINT | Ointment | A semisolid preparation intended for external application to the skin or mucous membranes.
4 | NASOINT | Nasal Ointment
4 | OINTAPL | Ointment with Applicator
4 | OPOINT | Ophthalmic Ointment
4 | OTOINT | Otic Ointment
4 | RECOINT | Rectal Ointment
4 | TOPOINT | Topical Ointment
4 | VAGOINT | Vaginal Ointment
5 | VAGOINTAPL | Vaginal Ointment with Applicator
2 | (_LiquidSolidSuspension) Abstract | | A liquid preparation which consists of solid particles dispersed throughout a liquid phase in which the particles are not soluble.
3 | GEL | Gel | A semisolid system consisting of either suspensions made up of small inorganic particles or large organic molecules interpenetrated by a liquid.
4 | GELAPL | Gel with Applicator
4 | NASGEL | Nasal Gel
4 | OPGEL | Ophthalmic Gel
4 | OTGEL | Otic Gel
4 | TOPGEL | Topical Gel
4 | URETHGEL | Urethral Gel
4 | VAGGEL | Vaginal Gel
5 | VGELAPL | Vaginal Gel with Applicator
3 | PASTE | Paste | A semisolid dosage form that contains one or more drug substances intended for topical application.
4 | PUD | Pudding
4 | TPASTE | Toothpaste | A paste formulation intended to clean and/or polish the teeth, and which may contain certain additional agents.
3 | SUSP | Suspension
4 | ITSUSP | Intrathecal Suspension
4 | OPSUSP | Ophthalmic Suspension
4 | ORSUSP | Oral Suspension
5 | ERSUSP | Extended-Release Suspension
6 | ERSUSP12 | 12 Hour Extended-Release Suspension
6 | ERSUSP24 | 24 Hour Extended Release Suspension
4 | OTSUSP | Otic Suspension
4 | RECSUSP | Rectal Suspension
2 | (_SolidDrugForm) Abstract
3 | BAR | Bar
4 | BARSOAP | Bar Soap
5 | MEDBAR | Medicated Bar Soap
4 | CHEWBAR | Chewable Bar | A solid dosage form usually in the form of a rectangle that is meant to be chewed.
3 | BEAD | Beads | A solid dosage form in the shape of a small ball.
3 | CAKE | Cake
3 | CEMENT | Cement | A substance that serves to produce solid union between two surfaces.
3 | CRYS | Crystals | A naturally produced angular solid of definite form in which the ultimate units from which it is built up are systematically arranged; they are usually evenly spaced on a regular space lattice.
3 | DISK | Disk | A circular plate-like organ or structure.
3 | FLAKE | Flakes
3 | GRAN | Granules | A small particle or grain.
3 | GUM | ChewingGum | A sweetened and flavored insoluble plastic material of various shapes which when chewed, releases a drug substance into the oral cavity.
3 | PAD | Pad
4 | MEDPAD | Medicated Pad
3 | PATCH | Patch | A drug delivery system that contains an adhesived backing and that permits its ingredients to diffuse from some portion of it (e.g., the backing itself, a reservoir, the adhesive, or some other component) into the body from the external site where it is applied.
4 | TPATCH | Transdermal Patch
5 | TPATH16 | 16 Hour Transdermal Patch
5 | TPATH24 | 24 Hour Transdermal Patch
5 | TPATH2WK | Biweekly Transdermal Patch
5 | TPATH72 | 72 Hour Transdermal Patch
5 | TPATHWK | Weekly Transdermal Patch
3 | PELLET | Pellet | A small sterile solid mass consisting of a highly purified drug (with or without excipients) made by the formation of granules, or by compression and molding.
3 | PILL | Pill | A small, round solid dosage form containing a medicinal agent intended for oral administration.
4 | CAP | Capsule | A solid dosage form in which the drug is enclosed within either a hard or soft soluble container or "shell" made from a suitable form of gelatin.
5 | ORCAP | Oral Capsule
6 | ENTCAP | Enteric Coated Capsule
7 | ERENTCAP | Extended Release Enteric Coated Capsule
6 | ERCAP | Extended Release Capsule | A solid dosage form in which the drug is enclosed within either a hard or soft soluble container made from a suitable form of gelatin, and which releases a drug (or drugs) in such a manner to allow a reduction in dosing frequency as compared to that drug (or drugs) presented as a conventional dosage form.
7 | ERENTCAP

7 | ERCAP12 | 12 Hour Extended Release Capsule
7 | ERCAP24 | 24 Hour Extended Release Capsule
7 | ERECCAP | Extended Release Enteric Coated Capsule | Rationale: Duplicate of code ERENTCAP. Use code ERENTCAP instead.
4 | TAB | Tablet | A solid dosage form containing medicinal substances with or without suitable diluents.
5 | ORTAB | Oral Tablet
6 | BUCTAB | Buccal Tablet
7 | SRBUCTAB | Sustained Release Buccal Tablet
6 | CAPLET | Caplet
6 | CHEWTAB | Chewable Tablet | A solid dosage form containing medicinal substances with or without suitable diluents that is intended to be chewed, producing a pleasant tasting residue in the oral cavity that is easily swallowed and does not leave a bitter or unpleasant after-taste.
6 | CPTAB | Coated Particles Tablet
6 | DISINTAB | Disintegrating Tablet | A solid dosage form containing medicinal substances which disintegrates rapidly, usually within a matter of seconds, when placed upon the tongue.
6 | DRTAB | Delayed Release Tablet
6 | ECTAB | Enteric Coated Tablet
7 | ERECTAB | Extended Release Enteric Coated Tablet
6 | ERTAB | Extended Release Tablet | A solid dosage form containing a drug which allows at least a reduction in dosing frequency as compared to that drug presented in conventional dosage form.
7 | ERECTAB
7 | ERTAB12 | 12 Hour Extended Release Tablet
7 | ERTAB24 | 24 Hour Extended Release Tablet
6 | ORTROCHE | Lozenge/Oral Troche | A solid preparation containing one or more medicaments, usually in a flavored, sweetened base which is intended to dissolve or disintegrate slowly in the mouth.
6 | SLTAB | Sublingual Tablet
5 | VAGTAB | Vaginal Tablet
3 | POWD | Powder | An intimate mixture of dry, finely divided drugs and/or chemicals that may be intended for internal or external use.
4 | TOPPWD | Topical Powder
5 | RECPWD | Rectal Powder
5 | VAGPWD | Vaginal Powder
3 | SUPP | Suppository | A solid body of various weights and shapes, adapted for introduction into the rectal, vaginal, or urethral orifice of the human body; they usually melt, soften, or dissolve at body temperature.
4 | RECSUPP | Rectal Suppository
4 | URETHSUPP | Urethral suppository
4 | VAGSUPP | Vaginal Suppository
3 | SWAB | Swab | A wad of absorbent material usually wound around one end of a small stick and used for applying medication or for removing material from an area.
4 | MEDSWAB | Medicated swab
3 | WAFER | Wafer | A thin slice of material containing a medicinal agent.

4.4.2.88 HL7 v3 Code System ParticipationFunction

URL: https://hl7.org/fhir/R4/v3/ParticipationFunction/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-ParticipationFunction) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-ParticipationFunction
Version: | 2018-08-12
Name: | v3.ParticipationFunction
Title: | v3 Code System ParticipationFunction
Definition: | This code is used to specify the exact function an actor had in a service in all necessary detail. This domain may include local extensions (CWE).
OID: | 2.16.840.1.113883.5.88 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: ParticipationRoleType (This FHIR value set is comprised of Actor participation Type codes, which can be used to value FHIR agents, actors, and other role elements. The FHIR Actor participation type value set is based on DICOM Audit Message, C402; ASTM Standard, E1762-95 [2013]; selected codes and derived actor roles from HL7 RoleClass OID 2.16.840.1.113883.5.110; HL7 Role Code 2.16.840.1.113883.5.111, including AgentRoleType; HL7 ParticipationType OID: 2.16.840.1.113883.5.90; and HL7 ParticipationFunction codes OID: 2.16.840.1.113883.5.88. This value set includes, by reference, role codes from external code systems: NUCC Health Care Provider Taxonomy OID: 2.16.840.1.113883.6.101; North American Industry Classification System [NAICS]OID: 2.16.840.1.113883.6.85; IndustryClassificationSystem 2.16.840.1.113883.1.11.16039; and US Census Occupation Code OID: 2.16.840.1.113883.6.243 for relevant recipient or custodian codes not included in this value set. If no source is indicated in the definition comments, then these are example FHIR codes.)

• ValueSet: ParticipationRoleType (This FHIR value set is comprised of Actor participation Type codes, which can be used to value FHIR agents, actors, and other role elements. The FHIR Actor participation type value set is based on DICOM Audit Message, C402; ASTM Standard, E1762-95 [2013]; selected codes and derived actor roles from HL7 RoleClass OID 2.16.840.1.113883.5.110; HL7 Role Code 2.16.840.1.113883.5.111, including AgentRoleType; HL7 ParticipationType OID: 2.16.840.1.113883.5.90; and HL7 ParticipationFunction codes OID: 2.16.840.1.113883.5.88. This value set includes, by reference, role codes from external code systems: NUCC Health Care Provider Taxonomy OID: 2.16.840.1.113883.6.101; North American Industry Classification System [NAICS]OID: 2.16.840.1.113883.6.85; IndustryClassificationSystem 2.16.840.1.113883.1.11.16039; and US Census Occupation Code OID: 2.16.840.1.113883.6.243 for relevant recipient or custodian codes not included in this value set. If no source is indicated in the definition comments, then these are example FHIR codes.)

• ValueSet: SecurityRoleType (This example FHIR value set is comprised of example Actor Type codes, which can be used to value FHIR agents, actors, and other role elements such as those specified in financial transactions. The FHIR Actor value set is based on DICOM Audit Message, C402; ASTM Standard, E1762-95 [2013]; selected codes and derived actor roles from HL7 RoleClass OID 2.16.840.1.113883.5.110; HL7 Role Code 2.16.840.1.113883.5.111, including AgentRoleType; HL7 ParticipationType OID: 2.16.840.1.113883.5.90; and HL7 ParticipationFunction codes OID: 2.16.840.1.113883.5.88. This value set includes, by reference, role codes from external code systems: NUCC Health Care Provider Taxonomy OID: 2.16.840.1.113883.6.101; North American Industry Classification System [NAICS]OID: 2.16.840.1.113883.6.85; IndustryClassificationSystem 2.16.840.1.113883.1.11.16039; and US Census Occupation Code OID: 2.16.840.1.113883.6.243 for relevant recipient or custodian codes not included in this value set. If no source is indicated in the definition comments, then these are example FHIR codes. It can be extended with appropriate roles described by SNOMED as well as those described in the HL7 Role Based Access Control Catalog and the HL7 Healthcare (Security and Privacy) Access Control Catalog. In Role-Based Access Control (RBAC), permissions are operations on an object that a user wishes to access. Permissions are grouped into roles. A role characterizes the functions a user is allowed to perform. Roles are assigned to users. If the user's role has the appropriate permissions to access an object, then that user is granted access to the object. FHIR readily enables RBAC, as FHIR Resources are object types and the CRUDE events (the FHIR equivalent to permissions in the RBAC scheme) are operations on those objects. In Attribute-Based Access Control (ABAC), a user requests to perform operations on objects. That user's access request is granted or denied based on a set of access control policies that are specified in terms of attributes and conditions. FHIR readily enables ABAC, as instances of a Resource in FHIR (again, Resources are object types) can have attributes associated with them. These attributes include security tags, environment conditions, and a host of user and object characteristics, which are the same attributes as those used in ABAC. Attributes help define the access control policies that determine the operations a user may perform on a Resource (in FHIR) or object (in ABAC). For example, a tag (or attribute) may specify that the identified Resource (object) is not to be further disclosed without explicit consent from the patient.)

• ValueSet: SecurityRoleType (This example FHIR value set is comprised of example Actor Type codes, which can be used to value FHIR agents, actors, and other role elements such as those specified in financial transactions. The FHIR Actor value set is based on DICOM Audit Message, C402; ASTM Standard, E1762-95 [2013]; selected codes and derived actor roles from HL7 RoleClass OID 2.16.840.1.113883.5.110; HL7 Role Code 2.16.840.1.113883.5.111, including AgentRoleType; HL7 ParticipationType OID: 2.16.840.1.113883.5.90; and HL7 ParticipationFunction codes OID: 2.16.840.1.113883.5.88. This value set includes, by reference, role codes from external code systems: NUCC Health Care Provider Taxonomy OID: 2.16.840.1.113883.6.101; North American Industry Classification System [NAICS]OID: 2.16.840.1.113883.6.85; IndustryClassificationSystem 2.16.840.1.113883.1.11.16039; and US Census Occupation Code OID: 2.16.840.1.113883.6.243 for relevant recipient or custodian codes not included in this value set. If no source is indicated in the definition comments, then these are example FHIR codes. It can be extended with appropriate roles described by SNOMED as well as those described in the HL7 Role Based Access Control Catalog and the HL7 Healthcare (Security and Privacy) Access Control Catalog. In Role-Based Access Control (RBAC), permissions are operations on an object that a user wishes to access. Permissions are grouped into roles. A role characterizes the functions a user is allowed to perform. Roles are assigned to users. If the user's role has the appropriate permissions to access an object, then that user is granted access to the object. FHIR readily enables RBAC, as FHIR Resources are object types and the CRUDE events (the FHIR equivalent to permissions in the RBAC scheme) are operations on those objects. In Attribute-Based Access Control (ABAC), a user requests to perform operations on objects. That user's access request is granted or denied based on a set of access control policies that are specified in terms of attributes and conditions. FHIR readily enables ABAC, as instances of a Resource in FHIR (again, Resources are object types) can have attributes associated with them. These attributes include security tags, environment conditions, and a host of user and object characteristics, which are the same attributes as those used in ABAC. Attributes help define the access control policies that determine the operations a user may perform on a Resource (in FHIR) or object (in ABAC). For example, a tag (or attribute) may specify that the identified Resource (object) is not to be further disclosed without explicit consent from the patient.)

• ValueSet: v3 Code System ParticipationFunction (This code is used to specify the exact function an actor had in a service in all necessary detail. This domain may include local extensions (CWE).)

4.4.2.88.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition

1 | (_AuthorizedParticipationFunction) Abstract | | This code is used to specify the exact function an actor is authorized to have in a service in all necessary detail.

2 | (_AuthorizedReceiverParticipationFunction) Abstract | | This code is used to specify the exact function an actor is authorized to have as a receiver of information that is the subject of a consent directive or consent override.

3 | AUCG | caregiver information receiver | Description:Caregiver authorized to receive patient health information.

3 | AULR | legitimate relationship information receiver | Description:Provider with legitimate relationship authorized to receive patient health information.

3 | AUTM | care team information receiver | Description:Member of care team authorized to receive patient health information.

3 | AUWA | work area information receiver | Description:Entities within specified work area authorized to receive patient health information.

2 | (_ConsenterParticipationFunction) Abstract | | This code is used to specify the exact function an actor is authorized to have in authoring a consent directive.

3 | GRDCON | legal guardian consent author | Description:Legal guardian of the subject of consent authorized to author a consent directive for the subject of consent.

3 | POACON | healthcare power of attorney consent author | Description:Person authorized with healthcare power of attorney to author a consent directive for the subject of consent.

3 | PRCON | personal representative consent author | Description:Personal representative of the subject of consent authorized to author a consent directive for the subject of consent.

3 | PROMSK | authorized provider masking author | Definition:Provider authorized to mask information to protect the patient, a third party, or to ensure that the provider has consulted with the patient prior to release of this information.

3 | SUBCON | subject of consent author | Description:Subject of consent authorized to author a consent directive.

2 | (_OverriderParticipationFunction) Abstract | | This code is used to specify the exact function an actor is authorized to have in authoring a consent override.

3 | AUCOV | consent overrider | Description:Entity authorized to override a consent directive.

3 | AUEMROV | emergency overrider | Description:Entity authorized to override a consent directive or privacy policy in an emergency.

1 | (_CoverageParticipationFunction) Abstract | | Definition: Set of codes indicating the manner in which sponsors, underwriters, and payers participate in a policy or program.

2 | (_PayorParticipationFunction) Abstract | | Definition: Set of codes indicating the manner in which payors participate in a policy or program.</

3 | CLMADJ | claims adjudication | Definition: Manages all operations required to adjudicate fee for service claims or managed care encounter reports.

3 | ENROLL | enrollment broker | Definition: Managing the enrollment of covered parties.

3 | FFSMGT | ffs management | Definition: Managing all operations required to administer a fee for service or indemnity health plan including enrolling covered parties and providing customer service, provider contracting, claims payment, care management and utilization review.

3 | MCMGT | managed care management | Definition: Managing all operations required to administer a managed care plan including enrolling covered parties and providing customer service,, provider contracting, claims payment, care management and utilization review.

3 | PROVMGT | provider management | Definition: Managing provider contracting, provider services, credentialing, profiling, performance measures, and ensuring network adequacy.

3 | UMGT | utilization management | Definition: Managing utilization of services by ensuring that providers adhere to, e.g., payeraTMs clinical protocols for medical appropriateness and standards of medical necessity. May include management of authorizations for services and referrals.

2 | (_SponsorParticipationFunction) Abstract | | Definition: Set of codes indicating the manner in which sponsors participate in a policy or program. NOTE: use only when the Sponsor is not further specified with a SponsorRoleType as being either a fully insured sponsor or a self insured sponsor.

3 | FULINRD | fully insured | Definition: Responsibility taken by a sponsor to contract with one or more underwriters for the assumption of full responsibility for the risk and administration of a policy or program.

3 | SELFINRD | self insured | Definition: Responsibility taken by a sponsor to organize the underwriting of risk and administration of a policy or program.

2 | (_UnderwriterParticipationFunction) Abstract | | Definition: Set of codes indicating the manner in which underwriters participate in a policy or program.

3 | PAYORCNTR | payor contracting | Definition: Contracting for the provision and administration of health services to payors while retaining the risk for coverage. Contracting may be for all provision and administration; or for provision of certain types of services; for provision of services by region; and by types of administration, e.g., claims adjudication, enrollment, provider management, and utilization management. Typically done by underwriters for sponsors who need coverage provided to covered parties in multiple regions. The underwriter may act as the payor in some, but not all of the regions in which coverage is provided.

3 | REINS | reinsures | Definition: Underwriting reinsurance for another underwriter for the policy or program.

3 | RETROCES | retrocessionaires | Definition: Underwriting reinsurance for another reinsurer.

3 | SUBCTRT | subcontracting risk | Definition: Delegating risk for a policy or program to one or more subcontracting underwriters, e.g., a major health insurer may delegate risk for provision of coverage under a national health plan to other underwriters by region .

3 | UNDERWRTNG | underwriting | Definition: Provision of underwriting analysis for another underwriter without assumption of risk.

1 | ADMPHYS | admitting physician | A physician who admitted a patient to a hospital or other care unit that is the context of this service.

1 | ANEST | anesthesist | In a typical anesthesia setting an anesthesiologist or anesthesia resident in charge of the anesthesia and life support, but only a witness to the surgical procedure itself. To clarify responsibilities anesthesia should always be represented as a separate service related to the surgery.

1 | ANRS | anesthesia nurse | In a typical anesthesia setting the nurse principally assisting the anesthesiologist during the critical periods.

1 | ASSEMBLER | assembly software | A device that operates independently of an author on custodian's algorithms for data extraction of existing information for purpose of generating a new artifact. UsageConstraint: ASSEMBLER ParticipationFunction should be used with DEV (device) ParticipationType.
1 | ATTPHYS | attending physician | A physician who is primarily responsible for a patient during the hospitalization, which is the context of the service.
1 | COMPOSER | composer software | A device used by an author to record new information, which may also be used by the author to select existing information for aggregation with newly recorded information for the purpose of generating a new artifact. UsageConstraint: COMPOSER ParticipationFunction should be used with DEV (device) ParticipationType. Usage Note: This code will enable implementers to more specifically represent the manner in which a Device participated in and facilitated the generation of a CDA Clinical Document or a CDA Entry by the responsible Author, which is comprised of the Author's newly entered content, and may include the pre-existing content selected by the Author, for the purpose of establishing the provenance and accountability for these acts.
1 | DISPHYS | discharging physician | A physician who discharged a patient from a hospital or other care unit that is the context of this service.
1 | FASST | first assistant surgeon | In a typical surgery setting the assistant facing the primary surgeon. The first assistant performs parts of the operation and assists in others (e.g., incision, approach, electrocoutering, ligatures, sutures).
1 | MDWF | midwife | A person (usually female) helping a woman deliver a baby. Responsibilities vary locally, ranging from a mere optional assistant to a full required participant, responsible for (normal) births and pre- and post-natal care for both mother and baby.
1 | NASST | nurse assistant | In a typical surgery setting the non-sterile nurse handles material supply from the stock, forwards specimen to pathology, and helps with other non-sterile tasks (e.g., phone calls, etc.).
1 | PCP | primary care physician | The healthcare provider that holds primary responsibility for the overall care of a patient.
1 | PRISURG | primary surgeon | In a typical surgery setting the primary performing surgeon.
1 | REVIEWER | reviewer | A verifier who is accountable for reviewing and asserting that the verification of an Act complies with jurisdictional or organizational policy. UsageConstraint: UsageConstraint: Specifies the exact function that an actor is authorized to have as a verifier of an Act. Connotes that a specialized verifier asserts compliance for veracity of the review per jurisdictional or organizational policy. E.g., The Provider who takes responsibility for authenticity of a record submitted to a payer. REVIEW ParticipationFunction should be used with VFR (verifier)
1 | RNDPHYS | rounding physician | A physician who made rounds on a patient in a hospital or other care center.
1 | SASST | second assistant surgeon | In a typical surgery setting the assistant who primarily holds the hooks.
1 | SNRS | scrub nurse | In a typical surgery setting the nurse in charge of the instrumentation.
1 | TASST | third assistant | In a typical surgery setting there is rarely a third assistant (e.g., in some Hip operations the third assistant postures the affected leg).

4.4.2.89 HL7 v3 Code System ParticipationSignature

URL: https://hl7.org/fhir/R4/v3/ParticipationSignature/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-ParticipationSignature) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-ParticipationSignature
Version: | 2018-08-12
Name: | v3.ParticipationSignature
Title: | v3 Code System ParticipationSignature
Definition: | A set of codes specifying whether and how the participant has attested his participation through a signature and or whether such a signature is needed. Examples: A surgical Procedure act object (representing a procedure report) requires a signature of the performing and responsible surgeon, and possibly other participants. (See also: Participation.signatureText.)
OID: | 2.16.840.1.113883.5.89 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System ParticipationSignature (A set of codes specifying whether and how the participant has attested his participation through a signature and or whether such a signature is needed. Examples: A surgical Procedure act object (representing a procedure report) requires a signature of the performing and responsible surgeon, and possibly other participants. (See also: Participation.signatureText.))

4.4.2.89.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | I | intended | The particpant intends to provide a signature.
1 | S | signed | Signature has been affixed, either written on file, or electronic (incl. digital) signature in Participation.signatureText.
1 | X | required | A signature for the service is required of this actor.

4.4.2.90 HL7 v3 Code System ParticipationType

URL: https://hl7.org/fhir/R4/v3/ParticipationType/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-ParticipationType) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-ParticipationType
Version: | 2018-08-12
Name: | v3.ParticipationType
Title: | v3 Code System ParticipationType
Definition: | **** MISSING DEFINITIONS ****
OID: | 2.16.840.1.113883.5.90 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: Participant type (This value set defines a set of codes that can be used to indicate how an individual participates in an encounter.)

• ValueSet: v3 Code System ParticipationType (**** MISSING DEFINITIONS ****)

• ValueSet: V3 Value SetParticipationVerifier (A person who verifies the correctness and appropriateness of the service (plan, order, event, etc.) and hence takes on accountability.)

• ValueSet: Provenance Event History Agent Role Codes (Types of roles indicating how a particular agent was involved with the creation or modification of a resource for use when exposing event history)

• ValueSet: ImagingStudy series performer function (Performer function of an agent in an imaging study series)

• ValueSet: V3 Value SetParticipationTargetSubject (The principle target that the service acts on. E.g. the patient in physical examination, a specimen in a lab observation. May also be a patient's family member (teaching) or a device or room (cleaning, disinfecting, housekeeping). Note: not all direct targets are subjects, consumables, and devices used as tools for a service are not subjects. However, a device may be a subject of a maintenance service.)

• ValueSet: Focal Subject Codes (Example value set composed from SNOMED CT and HL7 V3 codes for observation targets such as donor, fetus or spouse. As use cases are discovered, more values may be added.)

• ValueSet: ParticipationRoleType (This FHIR value set is comprised of Actor participation Type codes, which can be used to value FHIR agents, actors, and other role elements. The FHIR Actor participation type value set is based on DICOM Audit Message, C402; ASTM Standard, E1762-95 [2013]; selected codes and derived actor roles from HL7 RoleClass OID 2.16.840.1.113883.5.110; HL7 Role Code 2.16.840.1.113883.5.111, including AgentRoleType; HL7 ParticipationType OID: 2.16.840.1.113883.5.90; and HL7 ParticipationFunction codes OID: 2.16.840.1.113883.5.88. This value set includes, by reference, role codes from external code systems: NUCC Health Care Provider Taxonomy OID: 2.16.840.1.113883.6.101; North American Industry Classification System [NAICS]OID: 2.16.840.1.113883.6.85; IndustryClassificationSystem 2.16.840.1.113883.1.11.16039; and US Census Occupation Code OID: 2.16.840.1.113883.6.243 for relevant recipient or custodian codes not included in this value set. If no source is indicated in the definition comments, then these are example FHIR codes.)

• ValueSet: ParticipationRoleType (This FHIR value set is comprised of Actor participation Type codes, which can be used to value FHIR agents, actors, and other role elements. The FHIR Actor participation type value set is based on DICOM Audit Message, C402; ASTM Standard, E1762-95 [2013]; selected codes and derived actor roles from HL7 RoleClass OID 2.16.840.1.113883.5.110; HL7 Role Code 2.16.840.1.113883.5.111, including AgentRoleType; HL7 ParticipationType OID: 2.16.840.1.113883.5.90; and HL7 ParticipationFunction codes OID: 2.16.840.1.113883.5.88. This value set includes, by reference, role codes from external code systems: NUCC Health Care Provider Taxonomy OID: 2.16.840.1.113883.6.101; North American Industry Classification System [NAICS]OID: 2.16.840.1.113883.6.85; IndustryClassificationSystem 2.16.840.1.113883.1.11.16039; and US Census Occupation Code OID: 2.16.840.1.113883.6.243 for relevant recipient or custodian codes not included in this value set. If no source is indicated in the definition comments, then these are example FHIR codes.)

• ValueSet: SecurityRoleType (This example FHIR value set is comprised of example Actor Type codes, which can be used to value FHIR agents, actors, and other role elements such as those specified in financial transactions. The FHIR Actor value set is based on DICOM Audit Message, C402; ASTM Standard, E1762-95 [2013]; selected codes and derived actor roles from HL7 RoleClass OID 2.16.840.1.113883.5.110; HL7 Role Code 2.16.840.1.113883.5.111, including AgentRoleType; HL7 ParticipationType OID: 2.16.840.1.113883.5.90; and HL7 ParticipationFunction codes OID: 2.16.840.1.113883.5.88. This value set includes, by reference, role codes from external code systems: NUCC Health Care Provider Taxonomy OID: 2.16.840.1.113883.6.101; North American Industry Classification System [NAICS]OID: 2.16.840.1.113883.6.85; IndustryClassificationSystem 2.16.840.1.113883.1.11.16039; and US Census Occupation Code OID: 2.16.840.1.113883.6.243 for relevant recipient or custodian codes not included in this value set. If no source is indicated in the definition comments, then these are example FHIR codes. It can be extended with appropriate roles described by SNOMED as well as those described in the HL7 Role Based Access Control Catalog and the HL7 Healthcare (Security and Privacy) Access Control Catalog. In Role-Based Access Control (RBAC), permissions are operations on an object that a user wishes to access. Permissions are grouped into roles. A role characterizes the functions a user is allowed to perform. Roles are assigned to users. If the user's role has the appropriate permissions to access an object, then that user is granted access to the object. FHIR readily enables RBAC, as FHIR Resources are object types and the CRUDE events (the FHIR

equivalent to permissions in the RBAC scheme) are operations on those objects. In Attribute-Based Access Control (ABAC), a user requests to perform operations on objects. That user's access request is granted or denied based on a set of access control policies that are specified in terms of attributes and conditions. FHIR readily enables ABAC, as instances of a Resource in FHIR (again, Resources are object types) can have attributes associated with them. These attributes include security tags, environment conditions, and a host of user and object characteristics, which are the same attributes as those used in ABAC. Attributes help define the access control policies that determine the operations a user may perform on a Resource (in FHIR) or object (in ABAC). For example, a tag (or attribute) may specify that the identified Resource (object) is not to be further disclosed without explicit consent from the patient.)

• ValueSet: V3 Value SetParticipationTargetLocation (The facility where the service is done. May be a static building (or room therein) or a moving location (e.g., ambulance, helicopter, aircraft, train, truck, ship, etc.))

• ValueSet: Provenance activity type (This value set contains representative Activity Type codes, which includes codes from the HL7 DocumentCompletion, ActStatus, and DataOperations code system, W3C PROV-DM and PROV-N concepts and display names, several HL7 Lifecycle Event codes for which there are agreed upon definitions, and non-duplicated codes from the HL7 Security and Privacy Ontology Operations codes.)

• ValueSet: V3 Value SetParticipationPhysicalPerformer (A person who actually and principally carries out the action. Need not be the principal responsible actor, e.g. a surgery resident operating under supervision of attending surgeon, and may be the patient in self-care, e.g. fingerstick blood sugar. The traditional order filler is a performer. This information should accompany every service event.)

• ValueSet: V3 Value SetParticipationInformationGenerator (Parties that may or should contribute or have contributed information to the Act. Such information includes information leading to the decision to perform the Act and how to perform the Act (e.g., consultant), information that the Act itself seeks to reveal (e.g., informant of clinical history), or information about what Act was performed (e.g., informant witness).)

• ValueSet: Participant type (This value set defines a set of codes that can be used to indicate how an individual participates in an encounter.)

• ValueSet: V3 Value SetParticipationIndirectTarget (Target that is not substantially present in the act and which is not directly affected by the act, but which will be a focus of the record or documentation of the act.)

• ValueSet: SecurityRoleType (This example FHIR value set is comprised of example Actor Type codes, which can be used to value FHIR agents, actors, and other role elements such as those specified in financial transactions. The FHIR Actor value set is based on DICOM Audit Message, C402; ASTM Standard, E1762-95 [2013]; selected codes and derived actor roles from HL7 RoleClass OID 2.16.840.1.113883.5.110; HL7 Role Code 2.16.840.1.113883.5.111, including AgentRoleType; HL7 ParticipationType OID: 2.16.840.1.113883.5.90; and HL7 ParticipationFunction codes OID: 2.16.840.1.113883.5.88. This value set includes, by reference, role codes from external code systems: NUCC Health Care Provider Taxonomy OID: 2.16.840.1.113883.6.101; North American Industry Classification System [NAICS]OID: 2.16.840.1.113883.6.85; IndustryClassificationSystem 2.16.840.1.113883.1.11.16039; and US Census Occupation Code OID: 2.16.840.1.113883.6.243 for relevant recipient or custodian codes not included in this value set. If no source is indicated in the definition comments, then these are example FHIR codes. It can be extended with appropriate roles described by SNOMED as well as those described in the HL7 Role Based Access Control Catalog and the HL7 Healthcare (Security and Privacy) Access Control Catalog. In Role-Based Access Control (RBAC), permissions are operations on an object that a user wishes to access. Permissions are grouped into roles. A role characterizes the functions a user is allowed to perform. Roles are assigned to users. If the user's role has the appropriate permissions to access an object, then that user is granted access to the object. FHIR readily enables RBAC, as FHIR Resources are object types and the CRUDE events (the FHIR equivalent to permissions in the RBAC scheme) are operations on those objects. In Attribute-Based Access Control (ABAC), a user requests to perform operations on objects. That user's access request is granted or denied based on a set of access control policies that are specified in terms of attributes and conditions. FHIR readily enables ABAC, as instances of a Resource in FHIR (again, Resources are object types) can have attributes associated with them. These attributes include security tags, environment conditions, and a host of user and object characteristics, which are the same attributes as those used in ABAC. Attributes help define the access control policies that determine the operations a user may perform on a Resource (in FHIR) or object (in ABAC). For example, a tag (or attribute) may specify that the identified Resource (object) is not to be further disclosed without explicit consent from the patient.)

• ValueSet: Provenance activity type (This value set contains representative Activity Type codes, which includes codes from the HL7 DocumentCompletion, ActStatus, and DataOperations code system, W3C PROV-DM and PROV-N concepts and display names, several HL7 Lifecycle Event codes for which there are agreed upon definitions, and non-duplicated codes from the HL7 Security and Privacy Ontology Operations codes.)

• ValueSet: V3 Value SetParticipationInformationTranscriber (An entity entering the data into the originating system. The data entry entity is collected optionally for internal quality control purposes. This includes the transcriptionist for dictated text transcribed into electronic form.)

• ValueSet: ImagingStudy series performer function (Performer function of an agent in an imaging study series)

• ValueSet: Focal Subject Codes (Example value set composed from SNOMED CT and HL7 V3 codes for observation targets such as donor, fetus or spouse. As use cases are discovered, more values may be added.)

• ValueSet: Provenance Event History Agent Role Codes (Types of roles indicating how a particular agent was involved with the creation or modification of a resource for use when exposing event history)

• ValueSet: V3 Value SetParticipationTargetDirect (Target that is substantially present in the service and which is directly affected by the service action (includes consumed material, devices, etc.).)

4.4.2.90.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition

1 | PART | Participation | Indicates that the target of the participation is involved in some manner in the act, but does not qualify how.

2 | (_ParticipationAncillary) Abstract | | Participations related, but not primary to an act. The Referring, Admitting, and Discharging practitioners must be the same person as those authoring the ControlAct event for their respective trigger events.

3 | ADM | admitter | The practitioner who is responsible for admitting a patient to a patient encounter.

3 | ATND | attender | The practitioner that has responsibility for overseeing a patient's care during a patient encounter.

3 | CALLBCK | callback contact | A person or organization who should be contacted for follow-up questions about the act in place of the author.

3 | CON | consultant | An advisor participating in the service by performing evaluations and making recommendations.

3 | DIS | discharger | The practitioner who is responsible for the discharge of a patient from a patient encounter.

3 | ESC | escort | Only with Transportation services. A person who escorts the patient.

3 | REF | referrer | A person having referred the subject of the service to the performer (referring physician). Typically, a referring physician will receive a report.

2 | (_ParticipationInformationGenerator) Abstract | | Parties that may or should contribute or have contributed information to the Act. Such information includes information leading to the decision to perform the Act and how to perform the Act (e.g., consultant), information that the Act itself seeks to reveal (e.g., informant of clinical history), or information about what Act was performed (e.g., informant witness).

3 | AUT | author (originator) | Definition: A party that originates the Act and therefore has responsibility for the information given in the Act and ownership of this Act. Example: the report writer, the person writing the act definition, the guideline author, the placer of an order, the EKG cart (device) creating a report etc. Every Act should have an author. Authorship is regardless of mood always actual authorship. Examples of such policies might include: The author and anyone they explicitly delegate may update the report; All administrators within the same clinic may cancel and reschedule appointments created by other administrators within that clinic; A party that is neither an author nor a party who is extended authorship maintenance rights by policy, may only amend, reverse, override, replace, or follow up in other ways on this Act, whereby the Act remains intact and is linked to another Act authored by that other party.

3 | INF | informant | A source of reported information (e.g., a next of kin who answers questions about the patient's history). For history questions, the patient is logically an informant, yet the informant of history questions is implicitly the subject.

3 | TRANS | Transcriber | An entity entering the data into the originating system. The data entry entity is collected optionally for internal quality control purposes. This includes the transcriptionist for dictated text transcribed into electronic form.

4 | ENT | data entry person | A person entering the data into the originating system. The data entry person is collected optionally for internal quality control purposes. This includes the transcriptionist for dictated text.

3 | WIT | witness | Only with service events. A person witnessing the action happening without doing anything. A witness is not necessarily aware, much less approves of anything stated in the service event. Example for a witness is students watching an operation or an advanced directive witness.

2 | CST | custodian | An entity (person, organization or device) that is in charge of maintaining the information of this act (e.g., who maintains the report or the master service catalog item, etc.).

2 | DIR | direct target | Target participant that is substantially present in the act and which is directly involved in the action (includes consumed material, devices, etc.).

3 | ALY | analyte | The target of an Observation action. Links an observation to a Role whose player is the substance or most specific component entity (material, micro-organism, etc.) being measured within the subject. Examples: A "plasma porcelain substance concentration" has analyte a Role with player substance Entity "porcelain". UsageNotes: The Role that this participation connects to may be any Role whose player is that substance measured. Very often, the scoper may indicate the system in which the component is being measured. E.g., for "plasma porcelain" the scoper could be "Plasma".

3 | BBY | baby | In an obstetric service, the baby.

3 | CAT | catalyst | The catalyst of a chemical reaction, such as an enzyme or a platinum surface. In biochemical reactions, connects the enzyme with the molecular interaction

3 | CSM | consumable | Participant material that is taken up, diminished, altered, or disappears in the act.

4 | TPA | therapeutic agent | Something incorporated in the subject of a therapy service to achieve a physiologic effect (e.g., heal, relieve, provoke a condition, etc.) on the subject. In an administration service the therapeutic agent is a consumable, in a preparation or dispense service, it is a product. Thus, consumable or product must be specified in accordance with the kind of service.

3 | DEV | device | Participant used in performing the act without being substantially affected by the act (i.e. durable or inert with respect to that particular service). Examples: monitoring equipment, tools, but also access/drainage lines, prostheses, pace maker, etc.

4 | NRD | non-reuseable device | A device that changes ownership due to the service, e.g., a pacemaker, a prosthesis, an insulin injection equipment (pen), etc. Such material may need to be restocked after he service.

4 | RDV | reusable device | A device that does not change ownership due to the service, i.e., a surgical instrument or tool or an endoscope. The distinction between reuseable and non-reuseable must be made in order to know whether material must be re-stocked.

3 | DON | donor | In some organ transplantation services and rarely in transfusion services a donor will be a target participant in the service. However, in most cases transplantation is decomposed in three services: explantation, transport, and implantation.

The identity of the donor (recipient) is often irrelevant for the explantation (implantation) service.

3 | EXPAGNT | ExposureAgent | Description: The entity playing the associated role is the physical (including energy), chemical or biological substance that is participating in the exposure. For example in communicable diseases, the associated playing entity is the disease causing pathogen.

3 | EXPART | ExposureParticipation | Description:Direct participation in an exposure act where it is unknown that the participant is the source or subject of the exposure. If the participant is known to be the contact of an exposure then the SBJ participation type should be used. If the participant is known to be the source then the EXSRC participation type should be used.

4 | EXPTRGT | ExposureTarget | Description: The entity playing the associated role is the target (contact) of exposure.

4 | EXSRC | ExposureSource | Description:The entity playing the associated role is the source of exposure.

3 | PRD | product | Participant material that is brought forth (produced) in the act (e.g., specimen in a specimen collection, access or drainage in a placement service, medication package in a dispense service). It does not matter whether the material produced had existence prior to the service, or whether it is created in the service (e.g., in supply services the product is taken from a stock).

3 | SBJ | subject | The principle target on which the action happens. Examples: The patient in physical examination, a specimen in a lab observation. May also be a patient's family member (teaching) or a device or room (cleaning, disinfecting, housekeeping). UsageNotes: Not all direct targets are subjects. Consumables and devices used as tools for an act are not subjects. However, a device may be a subject of a maintenance action.

4 | SPC | specimen | The subject of non-clinical (e.g. laboratory) observation services is a specimen.

2 | IND | indirect target | Target that is not substantially present in the act and which is not directly affected by the act, but which will be a focus of the record or documentation of the act.

3 | BEN | beneficiary | Target on behalf of whom the service happens, but that is not necessarily present in the service. Can occur together with direct target to indicate that a target is both, as in the case where the patient is the indirect beneficiary of a service rendered to a family member, e.g. counseling or given home care instructions. This concept includes a participant, such as a covered party, who derives benefits from a service act covered by a coverage act. Note that the semantic role of the intended recipient who benefits from the happening denoted by the verb in the clause. Thus, a patient who has no coverage under a policy or program may be a beneficiary of a health service while not being the beneficiary of coverage for that service.

3 | CAGNT | causative agent | Definition: A factor, such as a microorganism, chemical substance, or form of radiation, whose presence, excessive presence, or (in deficiency diseases) relative absence is essential, in whole or in part, for the occurrence of a condition. Constraint: The use of this participation is limited to observations.

3 | COV | coverage target | The target participation for an individual in a health care coverage act in which the target role is either the policy holder of the coverage, or a covered party under the coverage.

3 | GUAR | guarantor party | The target person or organization contractually recognized by the issuer as a participant who has assumed fiscal responsibility for another personaTMs financial obligations by guaranteeing to pay for amounts owed to a particular account Example:The subscriber of the patientaTMs health insurance policy signs a contract with the provider to be fiscally responsible for the patient billing account balance amount owed.

3 | HLD | holder | Participant who posses an instrument such as a financial contract (insurance policy) usually based on some agreement with the author.

3 | RCT | record target | The record target indicates whose medical record holds the documentation of this act. This is especially important when the subject of a service is not the patient himself.

3 | RCV | receiver | The person (or organization) who receives the product of an Act.

2 | IRCP | information recipient | A party, who may or should receive or who has recieved the Act or subsequent or derivative information of that Act. Information recipient is inert, i.e., independent of mood." Rationale: this is a generalization of a too diverse family that the definition can't be any more specific, and the concept is abstract so one of the specializations should be used.

3 | NOT | ugent notification contact | An information recipient to notify for urgent matters about this Act. (e.g., in a laboratory order, critical results are being called by phone right away, this is the contact to call; or for an inpatient encounter, a next of kin to notify when the patient becomes critically ill).

3 | PRCP | primary information recipient | Information recipient to whom an act statement is primarily directed. E.g., a primary care provider receiving a discharge letter from a hospitalist, a health department receiving information on a suspected case of infectious disease. Multiple of these participations may exist on the same act without requiring that recipients be ranked as primary vs. secondary.

3 | REFB | Referred By | A participant (e.g. provider) who has referred the subject of an act (e.g. patient). Typically, a referred by participant will provide a report (e.g. referral).

3 | REFT | Referred to | The person who receives the patient

3 | TRC | tracker | A secondary information recipient, who receives copies (e.g., a primary care provider receiving copies of results as ordered by specialist).

2 | LOC | location | The facility where the service is done. May be a static building (or room therein) or a moving location (e.g., ambulance, helicopter, aircraft, train, truck, ship, etc.)

3 | DST | destination | The destination for services. May be a static building (or room therein) or a movable facility (e.g., ship).

3 | ELOC | entry location | A location where data about an Act was entered.

3 | ORG | origin | The location of origin for services. May be a static building (or room therein) or a movable facility (e.g., ship).

3 | RML | remote | Some services take place at multiple concurrent locations (e.g., telemedicine, telephone consultation). The location where the principal performing actor is located is taken as the primary location (LOC) while the other location(s) are considered "remote."

3 | VIA | via | For services, an intermediate location that specifies a path between origin an destination.

2 | PRF | performer | Definition: A person, non-person living subject, organization or device that who actually and principally carries out the action. Device should only be assigned as a performer in circumstances where the device is performing independent of human intervention. Need not be the principal responsible actor. Exampe: A surgery resident operating under supervision of attending surgeon, a search and rescue dog locating survivors, an electronic laboratory analyzer or the laboratory discipline requested to perform a laboratory test. The performer may also be the patient in self-care, e.g. fingerstick blood sugar. The traditional order filler is a performer. This information should accompany every service event. Note: that existing HL7 designs assign an organization as the playing entity of the Role that is the performer. These designs should be revised in subsequent releases to make this the scooping entity for the role involved.

3 | DIST | distributor | Distributes material used in or generated during the act.

3 | PPRF | primary performer | The principal or primary performer of the act.

3 | SPRF | secondary performer | A person assisting in an act through his substantial presence and involvement This includes: assistants, technicians, associates, or whatever the job titles may be.

2 | RESP | responsible party | The person or organization that has primary responsibility for the act. The responsible party is not necessarily present in an action, but is accountable for the action through the power to delegate, and the duty to review actions with the performing actor after the fact. This responsibility may be ethical, legal, contractual, fiscal, or fiduciary in nature. Example: A person who is the head of a biochemical laboratory; a sponsor for a policy or government program.
2 | VRF | verifier | A person who verifies the correctness and appropriateness of the service (plan, order, event, etc.) and hence takes on accountability.
3 | AUTHEN | authenticator | A verifier who attests to the accuracy of an act, but who does not have privileges to legally authenticate the act. An example would be a resident physician who sees a patient and dictates a note, then later signs it. Their signature constitutes an authentication.
3 | LA | legal authenticator | A verifier who legally authenticates the accuracy of an act. An example would be a staff physician who sees a patient and dictates a note, then later signs it. Their signature constitutes a legal authentication.

4.4.2.91 HL7 v3 Code System PaymentTerms

URL: https://hl7.org/fhir/R4/v3/PaymentTerms/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-PaymentTerms) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-PaymentTerms
Version: | 2018-08-12
Name: | v3.PaymentTerms
Title: | v3 Code System PaymentTerms
Definition: | Describes payment terms for a financial transaction, used in an invoice. This is typically expressed as a responsibility of the acceptor or payor of an invoice.
OID: | 2.16.840.1.113883.5.91 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System PaymentTerms (Describes payment terms for a financial transaction, used in an invoice. This is typically expressed as a responsibility of the acceptor or payor of an invoice.)

4.4.2.91.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | COD | Cash on Delivery | Payment in full for products and/or services is required as soon as the service is performed or goods delivered.
1 | N30 | Net 30 days | Payment in full for products and/or services is required 30 days from the time the service is performed or goods delivered.
1 | N60 | Net 60 days | Payment in full for products and/or services is required 60 days from the time the service is performed or goods delivered.
1 | N90 | Net 90 days | Payment in full for products and/or services is required 90 days from the time the service is performed or goods delivered.

4.4.2.93 HL7 v3 Code System PersonDisabilityType

URL: https://hl7.org/fhir/R4/v3/PersonDisabilityType/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-PersonDisabilityType) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-PersonDisabilityType
Version: | 2018-08-12
Name: | v3.PersonDisabilityType
Title: | v3 Code System PersonDisabilityType
Definition: | A code identifying a person's disability.
OID: | 2.16.840.1.113883.5.93 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System PersonDisabilityType (A code identifying a person's disability.)

4.4.2.93.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | 1 | Vision impaired | Vision impaired
1 | 2 | Hearing impaired | Hearing impaired
1 | 3 | Speech impaired | Speech impaired
1 | 4 | Mentally impaired | Mentally impaired
1 | 5 | Mobility impaired

2 | CB | Requires crib | A crib is required to move the person
2 | CR | Requires crutches | Person requires crutches to ambulate
2 | G | Requires gurney | A gurney is required to move the person
2 | WC | Requires wheelchair | Person requires a wheelchair to be ambulatory
2 | WK | Requires walker | Person requires a walker to ambulate

4.4.2.100 HL7 v3 Code System ProcessingID

URL: https://hl7.org/fhir/R4/v3/ProcessingID/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-ProcessingID) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-ProcessingID
Version: | 2018-08-12
Name: | v3.ProcessingID
Title: | v3 Code System ProcessingID
Definition: | Codes used to specify whether a message is part of a production, training, or debugging system.
OID: | 2.16.840.1.113883.5.100 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System ProcessingID (Codes used to specify whether a message is part of a production, training, or debugging system.)

4.4.2.100.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | D | Debugging | Identifies debugging type of processing.
1 | P | Production | Identifies production type of processing.
1 | T | Training | Identifies training type of processing.

4.4.2.101 HL7 v3 Code System ProcessingMode

URL: https://hl7.org/fhir/R4/v3/ProcessingMode/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-ProcessingMode) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-ProcessingMode
Version: | 2018-08-12
Name: | v3.ProcessingMode
Title: | v3 Code System ProcessingMode
Definition: | **** MISSING DEFINITIONS ****
OID: | 2.16.840.1.113883.5.101 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System ProcessingMode (**** MISSING DEFINITIONS ****)

4.4.2.101.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | A | Archive | Identifies archive mode of processing.
1 | I | Initial load | Identifies initial load mode of processing.
1 | R | Restore from archive | Identifies restore mode of processing.
1 | T | Current processing | Identifies on-line mode of processing.

4.4.2.102 HL7 v3 Code System QueryPriority

URL: https://hl7.org/fhir/R4/v3/QueryPriority/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-QueryPriority) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-QueryPriority
Version: | 2018-08-12
Name: | v3.QueryPriority
Title: | v3 Code System QueryPriority

Definition: | **** MISSING DEFINITIONS ****
OID: | 2.16.840.1.113883.5.102 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System QueryPriority (**** MISSING DEFINITIONS ****)

4.4.2.102.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | D | Deferred | Query response is deferred.
1 | I | Immediate | Query response is immediate.

4.4.2.103 HL7 v3 Code System QueryStatusCode

URL: https://hl7.org/fhir/R4/v3/QueryStatusCode/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-QueryStatusCode) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-QueryStatusCode
Version: | 2018-08-12
Name: | v3.QueryStatusCode
Title: | v3 Code System QueryStatusCode
Definition: | A code specifying the state of the Query.
OID: | 2.16.840.1.113883.5.103 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System QueryStatusCode (A code specifying the state of the Query.)

4.4.2.103.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | aborted | aborted | Query status aborted
1 | deliveredResponse | deliveredResponse | Query Status delivered response
1 | executing | executing | Query Status executing
1 | new | new | Query Status new
1 | waitContinuedQueryResponse | waitContinuedQueryResponse | Query Status wait continued

4.4.2.104 HL7 v3 Code System Race

URL: https://hl7.org/fhir/R4/v3/Race/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-Race) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-Race
Version: | 2018-08-12
Name: | v3.Race
Title: | v3 Code System Race
Definition: | In the United States, federal standards for classifying data on race determine the categories used by federal agencies and exert a strong influence on categorization by state and local agencies and private sector organizations. The federal standards do not conceptually define race, and they recognize the absence of an anthropological or scientific basis for racial classification. Instead, the federal standards acknowledge that race is a social-political construct in which an individual's own identification with one more race categories is preferred to observer identification. The standards use a variety of features to define five minimum race categories. Among these features are descent from "the original peoples" of a specified region or nation. The minimum race categories are American Indian or Alaska Native, Asian, Black or African American, Native Hawaiian or Other Pacific Islander, and White. The federal standards stipulate that race data need not be limited to the five minimum categories, but any expansion must be collapsible to those categories.
OID: | 2.16.840.1.113883.5.104 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v2 ETHNIC GROUP (FHIR Value set/code system definition for HL7 v2 table 0005 (ETHNIC GROUP))

• ValueSet: v3 Code System Race (In the United States, federal standards for classifying data on race determine the categories used by federal agencies and exert a strong influence on categorization by state and local agencies and private sector organizations. The federal standards do not conceptually define race, and they recognize the absence of an anthropological or scientific basis for racial classification. Instead, the federal standards acknowledge that race is a social-political construct in which an individual's own

identification with one more race categories is preferred to observer identification. The standards use a variety of features to define five minimum race categories. Among these features are descent from "the original peoples" of a specified region or nation. The minimum race categories are American Indian or Alaska Native, Asian, Black or African American, Native Hawaiian or Other Pacific Islander, and White. The federal standards stipulate that race data need not be limited to the five minimum categories, but any expansion must be collapsible to those categories.)

4.4.2.104.1 Code System Content

Release Date: 2018-08-12

Level	Code	Display	Definition
1	1002-5	American Indian or Alaska Native	
2	1004-1	American Indian	
3	1006-6	Abenaki	
3	1008-2	Algonquian	
3	1010-8	Apache	
4	1011-6	Chiricahua	
4	1012-4	Fort Sill Apache	
4	1013-2	Jicarilla Apache	
4	1014-0	Lipan Apache	
4	1015-7	Mescalero Apache	
4	1016-5	Oklahoma Apache	
4	1017-3	Payson Apache	
4	1018-1	San Carlos Apache	
4	1019-9	White Mountain Apache	
3	1021-5	Arapaho	
4	1022-3	Northern Arapaho	
4	1023-1	Southern Arapaho	
4	1024-9	Wind River Arapaho	
3	1026-4	Arikara	
3	1028-0	Assiniboine	
3	1030-6	Assiniboine Sioux	
4	1031-4	Fort Peck Assiniboine Sioux	
3	1033-0	Bannock	
3	1035-5	Blackfeet	
3	1037-1	Brotherton	
3	1039-7	Burt Lake Band	
3	1041-3	Caddo	
4	1042-1	Oklahoma Cado	
3	1044-7	Cahuilla	
4	1045-4	Agua Caliente Cahuilla	
4	1046-2	Augustine	
4	1047-0	Cabazon	
4	1048-8	Los Coyotes	
4	1049-6	Morongo	
4	1050-4	Santa Rosa Cahuilla	
4	1051-2	Torres-Martinez	
3	1053-8	California Tribes	
4	1054-6	Cahto	
4	1055-3	Chimariko	
4	1056-1	Coast Miwok	
4	1057-9	Digger	
4	1058-7	Kawaiisu	
4	1059-5	Kern River	
4	1060-3	Mattole	
4	1061-1	Red Wood	
4	1062-9	Santa Rosa	
4	1063-7	Takelma	
4	1064-5	Wappo	
4	1065-2	Yana	
4	1066-0	Yuki	
3	1068-6	Canadian and Latin American Indian	
4	1069-4	Canadian Indian	
4	1070-2	Central American Indian	
4	1071-0	French American Indian	
4	1072-8	Mexican American Indian	
4	1073-6	South American Indian	
3	1074-4	Spanish American Indian	
3	1076-9	Catawba	
4	1741-8	Alatna	
4	1742-6	Alexander	
4	1743-4	Allakaket	

4 | 1744-2 | Alanvik
4 | 1745-9 | Anvik
4 | 1746-7 | Arctic
4 | 1747-5 | Beaver
4 | 1748-3 | Birch Creek
4 | 1749-1 | Cantwell
4 | 1750-9 | Chalkyitsik
4 | 1751-7 | Chickaloon
4 | 1752-5 | Chistochina
4 | 1753-3 | Chitina
4 | 1754-1 | Circle
4 | 1755-8 | Cook Inlet
4 | 1756-6 | Copper Center
4 | 1757-4 | Copper River
4 | 1758-2 | Dot Lake
4 | 1759-0 | Doyon
4 | 1760-8 | Eagle
4 | 1761-6 | Eklutna
4 | 1762-4 | Evansville
4 | 1763-2 | Fort Yukon
4 | 1764-0 | Gakona
4 | 1765-7 | Galena
4 | 1766-5 | Grayling
4 | 1767-3 | Gulkana
4 | 1768-1 | Healy Lake
4 | 1769-9 | Holy Cross
4 | 1770-7 | Hughes
4 | 1771-5 | Huslia
4 | 1772-3 | Iliamna
4 | 1773-1 | Kaltag
4 | 1774-9 | Kluti Kaah
4 | 1775-6 | Knik
4 | 1776-4 | Koyukuk
4 | 1777-2 | Lake Minchumina
4 | 1778-0 | Lime
4 | 1779-8 | Mcgrath
4 | 1780-6 | Manley Hot Springs
4 | 1781-4 | Mentasta Lake
4 | 1782-2 | Minto
4 | 1783-0 | Nenana
4 | 1784-8 | Nikolai
4 | 1785-5 | Ninilchik
4 | 1786-3 | Nondalton
4 | 1787-1 | Northway
4 | 1788-9 | Nulato
4 | 1789-7 | Pedro Bay
4 | 1790-5 | Rampart
4 | 1791-3 | Ruby
4 | 1792-1 | Salamatof
4 | 1793-9 | Seldovia
4 | 1794-7 | Slana
4 | 1795-4 | Shageluk
4 | 1796-2 | Stevens
4 | 1797-0 | Stony River
4 | 1798-8 | Takotna
4 | 1799-6 | Tanacross
4 | 1800-2 | Tanaina
4 | 1801-0 | Tanana
4 | 1802-8 | Tanana Chiefs
4 | 1803-6 | Tazlina
4 | 1804-4 | Telida
4 | 1805-1 | Tetlin
4 | 1806-9 | Tok
4 | 1807-7 | Tyonek
4 | 1808-5 | Venetie
4 | 1809-3 | Wiseman
3 | 1078-5 | Cayuse
3 | 1080-1 | Chehalis
3 | 1082-7 | Chemakuan
4 | 1083-5 | Hoh
4 | 1084-3 | Quileute
3 | 1086-8 | Chemehuevi

3 | 1088-4 | Cherokee
4 | 1089-2 | Cherokee Alabama
4 | 1090-0 | Cherokees of Northeast Alabama
4 | 1091-8 | Cherokees of Southeast Alabama
4 | 1092-6 | Eastern Cherokee
4 | 1093-4 | Echota Cherokee
4 | 1094-2 | Etowah Cherokee
4 | 1095-9 | Northern Cherokee
4 | 1096-7 | Tuscola
4 | 1097-5 | United Keetowah Band of Cherokee
4 | 1098-3 | Western Cherokee
3 | 1100-7 | Cherokee Shawnee
3 | 1102-3 | Cheyenne
4 | 1103-1 | Northern Cheyenne
4 | 1104-9 | Southern Cheyenne
3 | 1106-4 | Cheyenne-Arapaho
3 | 1108-0 | Chickahominy
4 | 1109-8 | Eastern Chickahominy
4 | 1110-6 | Western Chickahominy
3 | 1112-2 | Chickasaw
3 | 1114-8 | Chinook
4 | 1115-5 | Clatsop
4 | 1116-3 | Columbia River Chinook
4 | 1117-1 | Kathlamet
4 | 1118-9 | Upper Chinook
4 | 1119-7 | Wakiakum Chinook
4 | 1120-5 | Willapa Chinook
4 | 1121-3 | Wishram
3 | 1123-9 | Chippewa
4 | 1124-7 | Bad River
4 | 1125-4 | Bay Mills Chippewa
4 | 1126-2 | Bois Forte
4 | 1127-0 | Burt Lake Chippewa
4 | 1128-8 | Fond du Lac
4 | 1129-6 | Grand Portage
4 | 1130-4 | Grand Traverse Band of Ottawa-Chippewa
4 | 1131-2 | Keweenaw
4 | 1132-0 | Lac Courte Oreilles
4 | 1133-8 | Lac du Flambeau
4 | 1134-6 | Lac Vieux Desert Chippewa
4 | 1135-3 | Lake Superior
4 | 1136-1 | Leech Lake
4 | 1137-9 | Little Shell Chippewa
4 | 1138-7 | Mille Lacs
4 | 1139-5 | Minnesota Chippewa
4 | 1140-3 | Ontonagon
4 | 1141-1 | Red Cliff Chippewa
4 | 1142-9 | Red Lake Chippewa
4 | 1143-7 | Saginaw Chippewa
4 | 1144-5 | St. Croix Chippewa
4 | 1145-2 | Sault Ste. Marie Chippewa
4 | 1146-0 | Sokoagon Chippewa
4 | 1147-8 | Turtle Mountain
4 | 1148-6 | White Earth
3 | 1150-2 | Chippewa Cree
4 | 1151-0 | Rocky Boy's Chippewa Cree
3 | 1153-6 | Chitimacha
3 | 1155-1 | Choctaw
4 | 1156-9 | Clifton Choctaw
4 | 1157-7 | Jena Choctaw
4 | 1158-5 | Mississippi Choctaw
4 | 1159-3 | Mowa Band of Choctaw
4 | 1160-1 | Oklahoma Choctaw
3 | 1162-7 | Chumash
4 | 1163-5 | Santa Ynez
3 | 1165-0 | Clear Lake
3 | 1167-6 | Coeur D'Alene
3 | 1169-2 | Coharie
3 | 1171-8 | Colorado River
3 | 1173-4 | Colville
3 | 1175-9 | Comanche
4 | 1176-7 | Oklahoma Comanche

3 | 1178-3 | Coos, Lower Umpqua, Siuslaw
3 | 1180-9 | Coos
3 | 1182-5 | Coquilles
3 | 1184-1 | Costanoan
3 | 1186-6 | Coushatta
4 | 1187-4 | Alabama Coushatta
3 | 1189-0 | Cowlitz
3 | 1191-6 | Cree
3 | 1193-2 | Creek
4 | 1194-0 | Alabama Creek
4 | 1195-7 | Alabama Quassarte
4 | 1196-5 | Eastern Creek
4 | 1197-3 | Eastern Muscogee
4 | 1198-1 | Kialegee
4 | 1199-9 | Lower Muscogee
4 | 1200-5 | Machis Lower Creek Indian
4 | 1201-3 | Poarch Band
4 | 1202-1 | Principal Creek Indian Nation
4 | 1203-9 | Star Clan of Muscogee Creeks
4 | 1204-7 | Thlopthlocco
4 | 1205-4 | Tuckabachee
3 | 1207-0 | Croatan
3 | 1209-6 | Crow
3 | 1211-2 | Cupeno
4 | 1212-0 | Agua Caliente
3 | 1214-6 | Delaware
4 | 1215-3 | Eastern Delaware
4 | 1216-1 | Lenni-Lenape
4 | 1217-9 | Munsee
4 | 1218-7 | Oklahoma Delaware
4 | 1219-5 | Rampough Mountain
4 | 1220-3 | Sand Hill
3 | 1222-9 | Diegueno
4 | 1223-7 | Campo
4 | 1224-5 | Capitan Grande
4 | 1225-2 | Cuyapaipe
4 | 1226-0 | La Posta
4 | 1227-8 | Manzanita
4 | 1228-6 | Mesa Grande
4 | 1229-4 | San Pasqual
4 | 1230-2 | Santa Ysabel
4 | 1231-0 | Sycuan
3 | 1233-6 | Eastern Tribes
4 | 1234-4 | Attacapa
4 | 1235-1 | Biloxi
4 | 1236-9 | Georgetown
4 | 1237-7 | Moor
4 | 1238-5 | Nansemond
4 | 1239-3 | Natchez
4 | 1240-1 | Nausu Waiwash
4 | 1241-9 | Nipmuc
4 | 1242-7 | Paugussett
4 | 1243-5 | Pocomoke Acohonock
4 | 1244-3 | Southeastern Indians
4 | 1245-0 | Susquehanock
4 | 1246-8 | Tunica Biloxi
4 | 1247-6 | Waccamaw-Siousan
4 | 1248-4 | Wicomico
3 | 1250-0 | Esselen
3 | 1252-6 | Fort Belknap
3 | 1254-2 | Fort Berthold
3 | 1256-7 | Fort Mcdowell
3 | 1258-3 | Fort Hall
3 | 1260-9 | Gabrieleno
3 | 1262-5 | Grand Ronde
3 | 1264-1 | Gros Ventres
4 | 1265-8 | Atsina
3 | 1267-4 | Haliwa
3 | 1269-0 | Hidatsa
3 | 1271-6 | Hoopa
4 | 1272-4 | Trinity
4 | 1273-2 | Whilkut

3 | 1275-7 | Hoopa Extension
3 | 1277-3 | Houma
3 | 1279-9 | Inaja-Cosmit
3 | 1281-5 | Iowa
4 | 1282-3 | Iowa of Kansas-Nebraska
4 | 1283-1 | Iowa of Oklahoma
3 | 1285-6 | Iroquois
4 | 1286-4 | Cayuga
4 | 1287-2 | Mohawk
4 | 1288-0 | Oneida
4 | 1289-8 | Onondaga
4 | 1290-6 | Seneca
4 | 1291-4 | Seneca Nation
4 | 1292-2 | Seneca-Cayuga
4 | 1293-0 | Tonawanda Seneca
4 | 1294-8 | Tuscarora
4 | 1295-5 | Wyandotte
3 | 1297-1 | Juaneno
3 | 1299-7 | Kalispel
3 | 1301-1 | Karuk
3 | 1303-7 | Kaw
3 | 1305-2 | Kickapoo
4 | 1306-0 | Oklahoma Kickapoo
4 | 1307-8 | Texas Kickapoo
3 | 1309-4 | Kiowa
4 | 1310-2 | Oklahoma Kiowa
3 | 1312-8 | Klallam
4 | 1313-6 | Jamestown
4 | 1314-4 | Lower Elwha
4 | 1315-1 | Port Gamble Klallam
3 | 1317-7 | Klamath
3 | 1319-3 | Konkow
3 | 1321-9 | Kootenai
3 | 1323-5 | Lassik
3 | 1325-0 | Long Island
4 | 1326-8 | Matinecock
4 | 1327-6 | Montauk
4 | 1328-4 | Poospatuck
4 | 1329-2 | Setauket
3 | 1331-8 | Luiseno
4 | 1332-6 | La Jolla
4 | 1333-4 | Pala
4 | 1334-2 | Pauma
4 | 1335-9 | Pechanga
4 | 1336-7 | Soboba
4 | 1337-5 | Twenty-Nine Palms
4 | 1338-3 | Temecula
3 | 1340-9 | Lumbee
3 | 1342-5 | Lummi
3 | 1344-1 | Maidu
4 | 1345-8 | Mountain Maidu
4 | 1346-6 | Nishinam
3 | 1348-2 | Makah
3 | 1350-8 | Maliseet
3 | 1352-4 | Mandan
3 | 1354-0 | Mattaponi
3 | 1356-5 | Menominee
3 | 1358-1 | Miami
4 | 1359-9 | Illinois Miami
4 | 1360-7 | Indiana Miami
4 | 1361-5 | Oklahoma Miami
3 | 1363-1 | Miccosukee
3 | 1365-6 | Micmac
4 | 1366-4 | Aroostook
3 | 1368-0 | Mission Indians
3 | 1370-6 | Miwok
3 | 1372-2 | Modoc
3 | 1374-8 | Mohegan
3 | 1376-3 | Mono
3 | 1378-9 | Nanticoke
3 | 1380-5 | Narragansett
3 | 1382-1 | Navajo

4 | 1383-9 | Alamo Navajo
4 | 1384-7 | Canoncito Navajo
4 | 1385-4 | Ramah Navajo
3 | 1387-0 | Nez Perce
3 | 1389-6 | Nomalaki
3 | 1391-2 | Northwest Tribes
4 | 1392-0 | Alsea
4 | 1393-8 | Celilo
4 | 1394-6 | Columbia
4 | 1395-3 | Kalapuya
4 | 1396-1 | Molala
4 | 1397-9 | Talakamish
4 | 1398-7 | Tenino
4 | 1399-5 | Tillamook
4 | 1400-1 | Wenatchee
4 | 1401-9 | Yahooskin
3 | 1403-5 | Omaha
3 | 1405-0 | Oregon Athabaskan
3 | 1407-6 | Osage
3 | 1409-2 | Otoe-Missouria
3 | 1411-8 | Ottawa
4 | 1412-6 | Burt Lake Ottawa
4 | 1413-4 | Michigan Ottawa
4 | 1414-2 | Oklahoma Ottawa
3 | 1416-7 | Paiute
4 | 1417-5 | Bishop
4 | 1418-3 | Bridgeport
4 | 1419-1 | Burns Paiute
4 | 1420-9 | Cedarville
4 | 1421-7 | Fort Bidwell
4 | 1422-5 | Fort Independence
4 | 1423-3 | Kaibab
4 | 1424-1 | Las Vegas
4 | 1425-8 | Lone Pine
4 | 1426-6 | Lovelock
4 | 1427-4 | Malheur Paiute
4 | 1428-2 | Moapa
4 | 1429-0 | Northern Paiute
4 | 1430-8 | Owens Valley
4 | 1431-6 | Pyramid Lake
4 | 1432-4 | San Juan Southern Paiute
4 | 1433-2 | Southern Paiute
4 | 1434-0 | Summit Lake
4 | 1435-7 | Utu Utu Gwaitu Paiute
4 | 1436-5 | Walker River
4 | 1437-3 | Yerington Paiute
3 | 1439-9 | Pamunkey
3 | 1441-5 | Passamaquoddy
4 | 1442-3 | Indian Township
4 | 1443-1 | Pleasant Point Passamaquoddy
3 | 1445-6 | Pawnee
4 | 1446-4 | Oklahoma Pawnee
3 | 1448-0 | Penobscot
3 | 1450-6 | Peoria
4 | 1451-4 | Oklahoma Peoria
3 | 1453-0 | Pequot
4 | 1454-8 | Marshantucket Pequot
3 | 1456-3 | Pima
4 | 1457-1 | Gila River Pima-Maricopa
4 | 1458-9 | Salt River Pima-Maricopa
3 | 1460-5 | Piscataway
3 | 1462-1 | Pit River
3 | 1464-7 | Pomo
4 | 1465-4 | Central Pomo
4 | 1466-2 | Dry Creek
4 | 1467-0 | Eastern Pomo
4 | 1468-8 | Kashia
4 | 1469-6 | Northern Pomo
4 | 1470-4 | Scotts Valley
4 | 1471-2 | Stonyford
4 | 1472-0 | Sulphur Bank
3 | 1474-6 | Ponca

4 | 1475-3 | Nebraska Ponca
4 | 1476-1 | Oklahoma Ponca
3 | 1478-7 | Potawatomi
4 | 1479-5 | Citizen Band Potawatomi
4 | 1480-3 | Forest County
4 | 1481-1 | Hannahville
4 | 1482-9 | Huron Potawatomi
4 | 1483-7 | Pokagon Potawatomi
4 | 1484-5 | Prairie Band
4 | 1485-2 | Wisconsin Potawatomi
3 | 1487-8 | Powhatan
3 | 1489-4 | Pueblo
4 | 1490-2 | Acoma
4 | 1491-0 | Arizona Tewa
4 | 1492-8 | Cochiti
4 | 1493-6 | Hopi
4 | 1494-4 | Isleta
4 | 1495-1 | Jemez
4 | 1496-9 | Keres
4 | 1497-7 | Laguna
4 | 1498-5 | Nambe
4 | 1499-3 | Picuris
4 | 1500-8 | Piro
4 | 1501-6 | Pojoaque
4 | 1502-4 | San Felipe
4 | 1503-2 | San Ildefonso
4 | 1504-0 | San Juan Pueblo
4 | 1505-7 | San Juan De
4 | 1506-5 | San Juan
4 | 1507-3 | Sandia
4 | 1508-1 | Santa Ana
4 | 1509-9 | Santa Clara
4 | 1510-7 | Santo Domingo
4 | 1511-5 | Taos
4 | 1512-3 | Tesuque
4 | 1513-1 | Tewa
4 | 1514-9 | Tigua
4 | 1515-6 | Zia
4 | 1516-4 | Zuni
3 | 1518-0 | Puget Sound Salish
4 | 1519-8 | Duwamish
4 | 1520-6 | Kikiallus
4 | 1521-4 | Lower Skagit
4 | 1522-2 | Muckleshoot
4 | 1523-0 | Nisqually
4 | 1524-8 | Nooksack
4 | 1525-5 | Port Madison
4 | 1526-3 | Puyallup
4 | 1527-1 | Samish
4 | 1528-9 | Sauk-Suiattle
4 | 1529-7 | Skokomish
4 | 1530-5 | Skykomish
4 | 1531-3 | Snohomish
4 | 1532-1 | Snoqualmie
4 | 1533-9 | Squaxin Island
4 | 1534-7 | Steilacoom
4 | 1535-4 | Stillaguamish
4 | 1536-2 | Suquamish
4 | 1537-0 | Swinomish
4 | 1538-8 | Tulalip
4 | 1539-6 | Upper Skagit
3 | 1541-2 | Quapaw
3 | 1543-8 | Quinault
3 | 1545-3 | Rappahannock
3 | 1547-9 | Reno-Sparks
3 | 1549-5 | Round Valley
3 | 1551-1 | Sac and Fox
4 | 1552-9 | Iowa Sac and Fox
4 | 1553-7 | Missouri Sac and Fox
4 | 1554-5 | Oklahoma Sac and Fox
3 | 1556-0 | Salinan
3 | 1558-6 | Salish

3 | 1560-2 | Salish and Kootenai
3 | 1562-8 | Schaghticoke
3 | 1564-4 | Scott Valley
3 | 1566-9 | Seminole
4 | 1567-7 | Big Cypress
4 | 1568-5 | Brighton
4 | 1569-3 | Florida Seminole
4 | 1570-1 | Hollywood Seminole
4 | 1571-9 | Oklahoma Seminole
3 | 1573-5 | Serrano
4 | 1574-3 | San Manual
3 | 1576-8 | Shasta
3 | 1578-4 | Shawnee
4 | 1579-2 | Absentee Shawnee
4 | 1580-0 | Eastern Shawnee
3 | 1582-6 | Shinnecock
3 | 1584-2 | Shoalwater Bay
3 | 1586-7 | Shoshone
4 | 1587-5 | Battle Mountain
4 | 1588-3 | Duckwater
4 | 1589-1 | Elko
4 | 1590-9 | Ely
4 | 1591-7 | Goshute
4 | 1592-5 | Panamint
4 | 1593-3 | Ruby Valley
4 | 1594-1 | Skull Valley
4 | 1595-8 | South Fork Shoshone
4 | 1596-6 | Te-Moak Western Shoshone
4 | 1597-4 | Timbi-Sha Shoshone
4 | 1598-2 | Washakie
4 | 1599-0 | Wind River Shoshone
4 | 1600-6 | Yomba
3 | 1602-2 | Shoshone Paiute
4 | 1603-0 | Duck Valley
4 | 1604-8 | Fallon
4 | 1605-5 | Fort McDermitt
3 | 1607-1 | Siletz
3 | 1609-7 | Sioux
4 | 1610-5 | Blackfoot Sioux
4 | 1611-3 | Brule Sioux
4 | 1612-1 | Cheyenne River Sioux
4 | 1613-9 | Crow Creek Sioux
4 | 1614-7 | Dakota Sioux
4 | 1615-4 | Flandreau Santee
4 | 1616-2 | Fort Peck
4 | 1617-0 | Lake Traverse Sioux
4 | 1618-8 | Lower Brule Sioux
4 | 1619-6 | Lower Sioux
4 | 1620-4 | Mdewakanton Sioux
4 | 1621-2 | Miniconjou
4 | 1622-0 | Oglala Sioux
4 | 1623-8 | Pine Ridge Sioux
4 | 1624-6 | Pipestone Sioux
4 | 1625-3 | Prairie Island Sioux
4 | 1626-1 | Prior Lake Sioux
4 | 1627-9 | Rosebud Sioux
4 | 1628-7 | Sans Arc Sioux
4 | 1629-5 | Santee Sioux
4 | 1630-3 | Sisseton-Wahpeton
4 | 1631-1 | Sisseton Sioux
4 | 1632-9 | Spirit Lake Sioux
4 | 1633-7 | Standing Rock Sioux
4 | 1634-5 | Teton Sioux
4 | 1635-2 | Two Kettle Sioux
4 | 1636-0 | Upper Sioux
4 | 1637-8 | Wahpekute Sioux
4 | 1638-6 | Wahpeton Sioux
4 | 1639-4 | Wazhaza Sioux
4 | 1640-2 | Yankton Sioux
4 | 1641-0 | Yanktonai Sioux
3 | 1643-6 | Siuslaw
3 | 1645-1 | Spokane

3 | 1647-7 | Stewart
3 | 1649-3 | Stockbridge
3 | 1651-9 | Susanville
3 | 1653-5 | Tohono O'Odham
4 | 1654-3 | Ak-Chin
4 | 1655-0 | Gila Bend
4 | 1656-8 | San Xavier
4 | 1657-6 | Sells
3 | 1659-2 | Tolowa
3 | 1661-8 | Tonkawa
3 | 1663-4 | Tygh
3 | 1665-9 | Umatilla
3 | 1667-5 | Umpqua
4 | 1668-3 | Cow Creek Umpqua
3 | 1670-9 | Ute
4 | 1671-7 | Allen Canyon
4 | 1672-5 | Uintah Ute
4 | 1673-3 | Ute Mountain Ute
3 | 1675-8 | Wailaki
3 | 1677-4 | Walla-Walla
3 | 1679-0 | Wampanoag
4 | 1680-8 | Gay Head Wampanoag
4 | 1681-6 | Mashpee Wampanoag
3 | 1683-2 | Warm Springs
3 | 1685-7 | Wascopum
3 | 1687-3 | Washoe
4 | 1688-1 | Alpine
4 | 1689-9 | Carson
4 | 1690-7 | Dresslerville
3 | 1692-3 | Wichita
3 | 1694-9 | Wind River
3 | 1696-4 | Winnebago
4 | 1697-2 | Ho-chunk
4 | 1698-0 | Nebraska Winnebago
3 | 1700-4 | Winnemucca
3 | 1702-0 | Wintun
3 | 1704-6 | Wiyot
4 | 1705-3 | Table Bluff
3 | 1707-9 | Yakama
3 | 1709-5 | Yakama Cowlitz
3 | 1711-1 | Yaqui
4 | 1712-9 | Barrio Libre
4 | 1713-7 | Pascua Yaqui
3 | 1715-2 | Yavapai Apache
3 | 1717-8 | Yokuts
4 | 1718-6 | Chukchansi
4 | 1719-4 | Tachi
4 | 1720-2 | Tule River
3 | 1722-8 | Yuchi
3 | 1724-4 | Yuman
4 | 1725-1 | Cocopah
4 | 1726-9 | Havasupai
4 | 1727-7 | Hualapai
4 | 1728-5 | Maricopa
4 | 1729-3 | Mohave
4 | 1730-1 | Quechan
4 | 1731-9 | Yavapai
3 | 1732-7 | Yurok
4 | 1733-5 | Coast Yurok
2 | 1735-0 | Alaska Native
3 | 1737-6 | Alaska Indian
4 | 1739-2 | Alaskan Athabascan
5 | 1740-0 | Ahtna
4 | 1811-9 | Southeast Alaska
5 | 1813-5 | Tlingit-Haida
6 | 1814-3 | Angoon
6 | 1815-0 | Central Council of Tlingit and Haida Tribes
6 | 1816-8 | Chilkat
6 | 1817-6 | Chilkoot
6 | 1818-4 | Craig
6 | 1819-2 | Douglas
6 | 1820-0 | Haida

6 | 1821-8 | Hoonah
6 | 1822-6 | Hydaburg
6 | 1823-4 | Kake
6 | 1824-2 | Kasaan
6 | 1825-9 | Kenaitze
6 | 1826-7 | Ketchikan
6 | 1827-5 | Klawock
6 | 1828-3 | Pelican
6 | 1829-1 | Petersburg
6 | 1830-9 | Saxman
6 | 1831-7 | Sitka
6 | 1832-5 | Tenakee Springs
6 | 1833-3 | Tlingit
6 | 1834-1 | Wrangell
6 | 1835-8 | Yakutat
5 | 1837-4 | Tsimshian
6 | 1838-2 | Metlakatla
3 | 1840-8 | Eskimo
4 | 1842-4 | Greenland Eskimo
4 | 1844-0 | Inupiat Eskimo
5 | 1845-7 | Ambler
5 | 1846-5 | Anaktuvuk
5 | 1847-3 | Anaktuvuk Pass
5 | 1848-1 | Arctic Slope Inupiat
5 | 1849-9 | Arctic Slope Corporation
5 | 1850-7 | Atqasuk
5 | 1851-5 | Barrow
5 | 1852-3 | Bering Straits Inupiat
5 | 1853-1 | Brevig Mission
5 | 1854-9 | Buckland
5 | 1855-6 | Chinik
5 | 1856-4 | Council
5 | 1857-2 | Deering
5 | 1858-0 | Elim
5 | 1859-8 | Golovin
5 | 1860-6 | Inalik Diomede
5 | 1861-4 | Inupiaq
5 | 1862-2 | Kaktovik
5 | 1863-0 | Kawerak
5 | 1864-8 | Kiana
5 | 1865-5 | Kivalina
5 | 1866-3 | Kobuk
5 | 1867-1 | Kotzebue
5 | 1868-9 | Koyuk
5 | 1869-7 | Kwiguk
5 | 1870-5 | Mauneluk Inupiat
5 | 1871-3 | Nana Inupiat
5 | 1872-1 | Noatak
5 | 1873-9 | Nome
5 | 1874-7 | Noorvik
5 | 1875-4 | Nuiqsut
5 | 1876-2 | Point Hope
5 | 1877-0 | Point Lay
5 | 1878-8 | Selawik
5 | 1879-6 | Shaktoolik
5 | 1880-4 | Shishmaref
5 | 1881-2 | Shungnak
5 | 1882-0 | Solomon
5 | 1883-8 | Teller
5 | 1884-6 | Unalakleet
5 | 1885-3 | Wainwright
5 | 1886-1 | Wales
5 | 1887-9 | White Mountain
5 | 1888-7 | White Mountain Inupiat
5 | 1889-5 | Mary's Igloo
4 | 1891-1 | Siberian Eskimo
5 | 1892-9 | Gambell
5 | 1893-7 | Savoonga
5 | 1894-5 | Siberian Yupik
4 | 1896-0 | Yupik Eskimo
5 | 1897-8 | Akiachak
5 | 1898-6 | Akiak

5 | 1899-4 | Alakanuk
5 | 1900-0 | Aleknagik
5 | 1901-8 | Andreafsky
5 | 1902-6 | Aniak
5 | 1903-4 | Atmautluak
5 | 1904-2 | Bethel
5 | 1905-9 | Bill Moore's Slough
5 | 1906-7 | Bristol Bay Yupik
5 | 1907-5 | Calista Yupik
5 | 1908-3 | Chefornak
5 | 1909-1 | Chevak
5 | 1910-9 | Chuathbaluk
5 | 1911-7 | Clark's Point
5 | 1912-5 | Crooked Creek
5 | 1913-3 | Dillingham
5 | 1914-1 | Eek
5 | 1915-8 | Ekuk
5 | 1916-6 | Ekwok
5 | 1917-4 | Emmonak
5 | 1918-2 | Goodnews Bay
5 | 1919-0 | Hooper Bay
5 | 1920-8 | Iqurmuit (Russian Mission)
5 | 1921-6 | Kalskag
5 | 1922-4 | Kasigluk
5 | 1923-2 | Kipnuk
5 | 1924-0 | Koliganek
5 | 1925-7 | Kongiganak
5 | 1926-5 | Kotlik
5 | 1927-3 | Kwethluk
5 | 1928-1 | Kwigillingok
5 | 1929-9 | Levelock
5 | 1930-7 | Lower Kalskag
5 | 1931-5 | Manokotak
5 | 1932-3 | Marshall
5 | 1933-1 | Mekoryuk
5 | 1934-9 | Mountain Village
5 | 1935-6 | Naknek
5 | 1936-4 | Napaumute
5 | 1937-2 | Napakiak
5 | 1938-0 | Napaskiak
5 | 1939-8 | Newhalen
5 | 1940-6 | New Stuyahok
5 | 1941-4 | Newtok
5 | 1942-2 | Nightmute
5 | 1943-0 | Nunapitchukv
5 | 1944-8 | Oscarville
5 | 1945-5 | Pilot Station
5 | 1946-3 | Pitkas Point
5 | 1947-1 | Platinum
5 | 1948-9 | Portage Creek
5 | 1949-7 | Quinhagak
5 | 1950-5 | Red Devil
5 | 1951-3 | St. Michael
5 | 1952-1 | Scammon Bay
5 | 1953-9 | Sheldon's Point
5 | 1954-7 | Sleetmute
5 | 1955-4 | Stebbins
5 | 1956-2 | Togiak
5 | 1957-0 | Toksook
5 | 1958-8 | Tulukskak
5 | 1959-6 | Tuntutuliak
5 | 1960-4 | Tununak
5 | 1961-2 | Twin Hills
5 | 1962-0 | Georgetown
5 | 1963-8 | St. Mary's
5 | 1964-6 | Umkumiate
3 | 1966-1 | Aleut
4 | 1968-7 | Alutiiq Aleut
5 | 1969-5 | Tatitlek
5 | 1970-3 | Ugashik
4 | 1972-9 | Bristol Bay Aleut
5 | 1973-7 | Chignik

5 | 1974-5 | Chignik Lake
5 | 1975-2 | Egegik
5 | 1976-0 | Igiugig
5 | 1977-8 | Ivanof Bay
5 | 1978-6 | King Salmon
5 | 1979-4 | Kokhanok
5 | 1980-2 | Perryville
5 | 1981-0 | Pilot Point
5 | 1982-8 | Port Heiden
4 | 1984-4 | Chugach Aleut
5 | 1985-1 | Chenega
5 | 1986-9 | Chugach Corporation
5 | 1987-7 | English Bay
5 | 1988-5 | Port Graham
4 | 1990-1 | Eyak
4 | 1992-7 | Koniag Aleut
5 | 1993-5 | Akhiok
5 | 1994-3 | Agdaagux
5 | 1995-0 | Karluk
5 | 1996-8 | Kodiak
5 | 1997-6 | Larsen Bay
5 | 1998-4 | Old Harbor
5 | 1999-2 | Ouzinkie
5 | 2000-8 | Port Lions
4 | 2002-4 | Sugpiaq
4 | 2004-0 | Suqpigaq
4 | 2006-5 | Unangan Aleut
5 | 2007-3 | Akutan
5 | 2008-1 | Aleut Corporation
5 | 2009-9 | Aleutian
5 | 2010-7 | Aleutian Islander
5 | 2011-5 | Atka
5 | 2012-3 | Belkofski
5 | 2013-1 | Chignik Lagoon
5 | 2014-9 | King Cove
5 | 2015-6 | False Pass
5 | 2016-4 | Nelson Lagoon
5 | 2017-2 | Nikolski
5 | 2018-0 | Pauloff Harbor
5 | 2019-8 | Qagan Toyagungin
5 | 2020-6 | Qawalangin
5 | 2021-4 | St. George
5 | 2022-2 | St. Paul
5 | 2023-0 | Sand Point
5 | 2024-8 | South Naknek
5 | 2025-5 | Unalaska
5 | 2026-3 | Unga
1 | 2028-9 | Asian
2 | 2029-7 | Asian Indian
2 | 2030-5 | Bangladeshi
2 | 2031-3 | Bhutanese
2 | 2032-1 | Burmese
2 | 2033-9 | Cambodian
2 | 2034-7 | Chinese
2 | 2035-4 | Taiwanese
2 | 2036-2 | Filipino
2 | 2037-0 | Hmong
2 | 2038-8 | Indonesian
2 | 2039-6 | Japanese
2 | 2040-4 | Korean
2 | 2041-2 | Laotian
2 | 2042-0 | Malaysian
2 | 2043-8 | Okinawan
2 | 2044-6 | Pakistani
2 | 2045-3 | Sri Lankan
2 | 2046-1 | Thai
2 | 2047-9 | Vietnamese
2 | 2048-7 | Iwo Jiman
2 | 2049-5 | Maldivian
2 | 2050-3 | Nepalese
2 | 2051-1 | Singaporean
2 | 2052-9 | Madagascar

1 | 2054-5 | Black or African American
2 | 2056-0 | Black
2 | 2058-6 | African American
2 | 2060-2 | African
3 | 2061-0 | Botswanan
3 | 2062-8 | Ethiopian
3 | 2063-6 | Liberian
3 | 2064-4 | Namibian
3 | 2065-1 | Nigerian
3 | 2066-9 | Zairean
2 | 2067-7 | Bahamian
2 | 2068-5 | Barbadian
2 | 2069-3 | Dominican
2 | 2070-1 | Dominica Islander
2 | 2071-9 | Haitian
2 | 2072-7 | Jamaican
2 | 2073-5 | Tobagoan
2 | 2074-3 | Trinidadian
2 | 2075-0 | West Indian
1 | 2076-8 | Native Hawaiian or Other Pacific Islander
2 | 2078-4 | Polynesian
3 | 2079-2 | Native Hawaiian
3 | 2080-0 | Samoan
3 | 2081-8 | Tahitian
3 | 2082-6 | Tongan
3 | 2083-4 | Tokelauan
2 | 2085-9 | Micronesian
3 | 2086-7 | Guamanian or Chamorro
3 | 2087-5 | Guamanian
3 | 2088-3 | Chamorro
3 | 2089-1 | Mariana Islander
3 | 2090-9 | Marshallese
3 | 2091-7 | Palauan
3 | 2092-5 | Carolinian
3 | 2093-3 | Kosraean
3 | 2094-1 | Pohnpeian
3 | 2095-8 | Saipanese
3 | 2096-6 | Kiribati
3 | 2097-4 | Chuukese
3 | 2098-2 | Yapese
2 | 2100-6 | Melanesian
3 | 2101-4 | Fijian
3 | 2102-2 | Papua New Guinean
3 | 2103-0 | Solomon Islander
3 | 2104-8 | New Hebrides
2 | 2500-7 | Other Pacific Islander | Note that this term remains in the table for completeness, even though within HL7, the notion of Other code is deprecated.
1 | 2106-3 | White
2 | 2108-9 | European
3 | 2109-7 | Armenian
3 | 2110-5 | English
3 | 2111-3 | French
3 | 2112-1 | German
3 | 2113-9 | Irish
3 | 2114-7 | Italian
3 | 2115-4 | Polish
3 | 2116-2 | Scottish
2 | 2118-8 | Middle Eastern or North African
3 | 2119-6 | Assyrian
3 | 2120-4 | Egyptian
3 | 2121-2 | Iranian
3 | 2122-0 | Iraqi
3 | 2123-8 | Lebanese
3 | 2124-6 | Palestinian
3 | 2125-3 | Syrian
3 | 2126-1 | Afghanistani
3 | 2127-9 | Israeili
2 | 2129-5 | Arab
1 | 2131-1 | Other Race | Note that this term remains in the table for completeness, even though within HL7, the notion of Other code is deprecated.

4.4.2.105 HL7 v3 Code System RelationalOperator

URL: https://hl7.org/fhir/R4/v3/RelationalOperator/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-RelationalOperator) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-RelationalOperator
Version: | 2018-08-12
Name: | v3.RelationalOperator
Title: | v3 Code System RelationalOperator
Definition: | **** MISSING DEFINITIONS ****
OID: | 2.16.840.1.113883.5.105 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System RelationalOperator (**** MISSING DEFINITIONS ****)

4.4.2.105.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | CT | Contains | Specified set of things includes value being evaluated.
1 | EQ | Equal | Equal condition applied to comparisons.
1 | GE | Greater than or equal | Greater than or equal condition applied to comparisons.
1 | GN | Generic | A generic comparison selects a record for inclusion in the response if the beginning of the designated element value matches the select string.
1 | GT | Greater than | Greater than condition applied to comparisons.
1 | LE | Less than or equal | Less than or equal condition applied to comparisons.
1 | LT | Less than | Less than condition applied to comparisons.
1 | NE | Not Equal | Not equal condition applied to comparisons.

4.4.2.106 HL7 v3 Code System RelationshipConjunction

URL: https://hl7.org/fhir/R4/v3/RelationshipConjunction/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-RelationshipConjunction) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-RelationshipConjunction
Version: | 2018-08-12
Name: | v3.RelationshipConjunction
Title: | v3 Code System RelationshipConjunction
Definition: | **** MISSING DEFINITIONS ****
OID: | 2.16.840.1.113883.5.106 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System RelationshipConjunction (**** MISSING DEFINITIONS ****)

4.4.2.106.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | AND | and | This condition must be true.
1 | OR | or | At least one of the condition among all OR conditions must be true.
1 | XOR | exclusive or | One and only one of the XOR conditions must be true.

4.4.2.107 HL7 v3 Code System RoleLinkType

URL: https://hl7.org/fhir/R4/v3/RoleLinkType/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-RoleLinkType) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-RoleLinkType
Version: | 2018-08-12
Name: | v3.RoleLinkType
Title: | v3 Code System RoleLinkType
Definition: | **** MISSING DEFINITIONS ****
OID: | 2.16.840.1.113883.5.107 (for OID based terminology systems)

Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System RoleLinkType (**** MISSING DEFINITIONS ****)

4.4.2.107.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | REL | related | An action taken with respect to a subject Entity by a regulatory or authoritative body with supervisory capacity over that entity. The action is taken in response to behavior by the subject Entity that body finds to be undesirable. Suspension, license restrictions, monetary fine, letter of reprimand, mandated training, mandated supervision, etc.Examples:
2 | BACKUP | is backup for | This relationship indicates the source Role is available to the target Role as a backup. An entity in a backup role will be available as a substitute or replacement in the event that the entity assigned the role is unavailable. In medical roles where it is critical that the function be performed and there is a possibility that the individual assigned may be ill or otherwise indisposed, another individual is assigned to cover for the individual originally assigned the role. A backup may be required to be identified, but unless the backup is actually used, he/she would not assume the assigned entity role.
2 | CONT | has contact | This relationship indicates the target Role provides or receives information regarding the target role. For example, AssignedEntity is a contact for a ServiceDeliveryLocation.
2 | DIRAUTH | has direct authority over | The source Role has direct authority over the target role in a chain of authority.
2 | IDENT | Identification | Description: The source role provides identification for the target role. The source role must be IDENT. The player entity of the source role is constrained to be the same as (i.e. the equivalent of, or equal to) the player of the target role if present. If the player is absent from the source role, then it is assumed to be the same as the player of the target role.
2 | INDAUTH | has indirect authority over | The source Role has indirect authority over the target role in a chain of authority.
2 | PART | has part | The target Role is part of the Source Role.
2 | REPL | replaces | This relationship indicates that the source Role replaces (or subsumes) the target Role. Allows for new identifiers and/or new effective time for a registry entry or a certification, etc.

4.4.2.108 HL7 v3 Code System ResponseLevel

URL: https://hl7.org/fhir/R4/v3/ResponseLevel/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-ResponseLevel) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-ResponseLevel
Version: | 2018-08-12
Name: | v3.ResponseLevel
Title: | v3 Code System ResponseLevel
Definition: | Specifies whether a response is expected from the addressee of this interaction and what level of detail that response should include
OID: | 2.16.840.1.113883.5.108 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System ResponseLevel (Specifies whether a response is expected from the addressee of this interaction and what level of detail that response should include)

4.4.2.108.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | C | completion | Respond with exceptions and a notification of completion
1 | D | detail | Respond with exceptions, completion, modifications and include more detail information (if applicable)
1 | E | exception | Respond with exceptions only
1 | F | confirmation | Respond with exceptions, completion, and modification with detail (as above), and send positive confirmations even if no modifications are being made.
1 | N | message-control | Respond only with message level acknowledgements, i.e., only notify acceptance or rejection of the message, do not include any application-level detail
1 | R | modification | Respond with exceptions, completions and modifications or revisions done before completion
1 | X | none | Do not send any kind of response

4.4.2.109 HL7 v3 Code System ResponseModality

URL: https://hl7.org/fhir/R4/v3/ResponseModality/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-ResponseModality) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-ResponseModality
Version: | 2018-08-12
Name: | v3.ResponseModality
Title: | v3 Code System ResponseModality

Definition: | Defines the timing and grouping of the response instances. OpenIssue: Description copied from Concept Domain of same name. Must be verified.
OID: | 2.16.840.1.113883.5.109 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: v3 Code System ResponseModality (Defines the timing and grouping of the response instances. OpenIssue: Description copied from Concept Domain of same name. Must be verified.)

4.4.2.109.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition
1 | B | Batch | Query response to be sent as an HL7 Batch.
1 | R | Real Time | Query response to occur in real time.
1 | T | Bolus | Query response to sent as a series of responses at the same time without the use of batch formatting.

4.4.2.110 HL7 v3 Code System RoleClass

URL: https://hl7.org/fhir/R4/v3/RoleClass/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-RoleClass) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-RoleClass
Version: | 2018-08-12
Name: | v3.RoleClass
Title: | v3 Code System RoleClass
Definition: | Codes for the Role class hierarchy. The values in this hierarchy, represent a Role which is an association or relationship between two entities - the entity that plays the role and the entity that scopes the role. Roles names are derived from the name of the playing entity in that role. The role hierarchy stems from three core concepts, or abstract domains: RoleClassOntological is an abstract domain that collects roles in which the playing entity is defined or specified by the scoping entity. RoleClassPartitive collects roles in which the playing entity is in some sense a "part" of the scoping entity. RoleClassAssociative collects all of the remaining forms of association between the playing entity and the scoping entity. This set of roles is further partitioned between: RoleClassPassive which are roles in which the playing entity is used, known, treated, handled, built, or destroyed, etc. under the auspices of the scoping entity. The playing entity is passive in these roles in that the role exists without an agreement from the playing entity. RoleClassMutualRelationship which are relationships based on mutual behavior of the two entities. The basis of these relationship may be formal agreements or they may bede facto behavior. Thus, this sub-domain is further divided into: RoleClassRelationshipFormal in which the relationship is formally defined, frequently by a contract or agreement. Personal relationship which inks two people in a personal relationship. The hierarchy discussed above is represented In the current vocabulary tables as a set of abstract domains, with the exception of the "Personal relationship" which is a leaf concept. OpenIssue: Description copied from Concept Domain of same name. Must be verified.
OID: | 2.16.840.1.113883.5.110 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: V3 Value SetRoleClassSpecimen (A role played by a material entity that is a specimen for an act. It is scoped by the source of the specimen.)

• ValueSet: V3 Value SetRoleClassAssociative (A general association between two entities that is neither partitive nor ontological.)

• ValueSet: ParticipationRoleType (This FHIR value set is comprised of Actor participation Type codes, which can be used to value FHIR agents, actors, and other role elements. The FHIR Actor participation type value set is based on DICOM Audit Message, C402; ASTM Standard, E1762-95 [2013]; selected codes and derived actor roles from HL7 RoleClass OID 2.16.840.1.113883.5.110; HL7 Role Code 2.16.840.1.113883.5.111, including AgentRoleType; HL7 ParticipationType OID: 2.16.840.1.113883.5.90; and HL7 ParticipationFunction codes OID: 2.16.840.1.113883.5.88. This value set includes, by reference, role codes from external code systems: NUCC Health Care Provider Taxonomy OID: 2.16.840.1.113883.6.101; North American Industry Classification System [NAICS]OID: 2.16.840.1.113883.6.85; IndustryClassificationSystem 2.16.840.1.113883.1.11.16039; and US Census Occupation Code OID: 2.16.840.1.113883.6.243 for relevant recipient or custodian codes not included in this value set. If no source is indicated in the definition comments, then these are example FHIR codes.)

• ValueSet: V3 Value SetRoleClassAgent (An entity (player) that acts or is authorized to act on behalf of another entity (scoper).)

• ValueSet: ParticipationRoleType (This FHIR value set is comprised of Actor participation Type codes, which can be used to value FHIR agents, actors, and other role elements. The FHIR Actor participation type value set is based on DICOM Audit Message, C402; ASTM Standard, E1762-95 [2013]; selected codes and derived actor roles from HL7 RoleClass OID 2.16.840.1.113883.5.110; HL7 Role Code 2.16.840.1.113883.5.111, including AgentRoleType; HL7 ParticipationType OID: 2.16.840.1.113883.5.90; and HL7 ParticipationFunction codes OID: 2.16.840.1.113883.5.88. This value set includes, by reference, role codes from external code systems: NUCC Health Care Provider Taxonomy OID: 2.16.840.1.113883.6.101; North American Industry Classification

System [NAICS]OID: 2.16.840.1.113883.6.85; IndustryClassificationSystem 2.16.840.1.113883.1.11.16039; and US Census Occupation Code OID: 2.16.840.1.113883.6.243 for relevant recipient or custodian codes not included in this value set. If no source is indicated in the definition comments, then these are example FHIR codes.)

• ValueSet: v3 Code System RoleClass (Codes for the Role class hierarchy. The values in this hierarchy, represent a Role which is an association or relationship between two entities - the entity that plays the role and the entity that scopes the role. Roles names are derived from the name of the playing entity in that role. The role hierarchy stems from three core concepts, or abstract domains: RoleClassOntological is an abstract domain that collects roles in which the playing entity is defined or specified by the scoping entity. RoleClassPartitive collects roles in which the playing entity is in some sense a "part" of the scoping entity. RoleClassAssociative collects all of the remaining forms of association between the playing entity and the scoping entity. This set of roles is further partitioned between: RoleClassPassive which are roles in which the playing entity is used, known, treated, handled, built, or destroyed, etc. under the auspices of the scoping entity. The playing entity is passive in these roles in that the role exists without an agreement from the playing entity. RoleClassMutualRelationship which are relationships based on mutual behavior of the two entities. The basis of these relationship may be formal agreements or they may bede facto behavior. Thus, this sub-domain is further divided into: RoleClassRelationshipFormal in which the relationship is formally defined, frequently by a contract or agreement. Personal relationship which inks two people in a personal relationship. The hierarchy discussed above is represented In the current vocabulary tables as a set of abstract domains, with the exception of the "Personal relationship" which is a leaf concept. OpenIssue: Description copied from Concept Domain of same name. Must be verified.)

• ValueSet: SecurityRoleType (This example FHIR value set is comprised of example Actor Type codes, which can be used to value FHIR agents, actors, and other role elements such as those specified in financial transactions. The FHIR Actor value set is based on DICOM Audit Message, C402; ASTM Standard, E1762-95 [2013]; selected codes and derived actor roles from HL7 RoleClass OID 2.16.840.1.113883.5.110; HL7 Role Code 2.16.840.1.113883.5.111, including AgentRoleType; HL7 ParticipationType OID: 2.16.840.1.113883.5.90; and HL7 ParticipationFunction codes OID: 2.16.840.1.113883.5.88. This value set includes, by reference, role codes from external code systems: NUCC Health Care Provider Taxonomy OID: 2.16.840.1.113883.6.101; North American Industry Classification System [NAICS]OID: 2.16.840.1.113883.6.85; IndustryClassificationSystem 2.16.840.1.113883.1.11.16039; and US Census Occupation Code OID: 2.16.840.1.113883.6.243 for relevant recipient or custodian codes not included in this value set. If no source is indicated in the definition comments, then these are example FHIR codes. It can be extended with appropriate roles described by SNOMED as well as those described in the HL7 Role Based Access Control Catalog and the HL7 Healthcare (Security and Privacy) Access Control Catalog. In Role-Based Access Control (RBAC), permissions are operations on an object that a user wishes to access. Permissions are grouped into roles. A role characterizes the functions a user is allowed to perform. Roles are assigned to users. If the user's role has the appropriate permissions to access an object, then that user is granted access to the object. FHIR readily enables RBAC, as FHIR Resources are object types and the CRUDE events (the FHIR equivalent to permissions in the RBAC scheme) are operations on those objects. In Attribute-Based Access Control (ABAC), a user requests to perform operations on objects. That user's access request is granted or denied based on a set of access control policies that are specified in terms of attributes and conditions. FHIR readily enables ABAC, as instances of a Resource in FHIR (again, Resources are object types) can have attributes associated with them. These attributes include security tags, environment conditions, and a host of user and object characteristics, which are the same attributes as those used in ABAC. Attributes help define the access control policies that determine the operations a user may perform on a Resource (in FHIR) or object (in ABAC). For example, a tag (or attribute) may specify that the identified Resource (object) is not to be further disclosed without explicit consent from the patient.)

• ValueSet: V3 Value SetRoleClassManufacturedProduct (Scoped by the manufacturer)

• ValueSet: SecurityRoleType (This example FHIR value set is comprised of example Actor Type codes, which can be used to value FHIR agents, actors, and other role elements such as those specified in financial transactions. The FHIR Actor value set is based on DICOM Audit Message, C402; ASTM Standard, E1762-95 [2013]; selected codes and derived actor roles from HL7 RoleClass OID 2.16.840.1.113883.5.110; HL7 Role Code 2.16.840.1.113883.5.111, including AgentRoleType; HL7 ParticipationType OID: 2.16.840.1.113883.5.90; and HL7 ParticipationFunction codes OID: 2.16.840.1.113883.5.88. This value set includes, by reference, role codes from external code systems: NUCC Health Care Provider Taxonomy OID: 2.16.840.1.113883.6.101; North American Industry Classification System [NAICS]OID: 2.16.840.1.113883.6.85; IndustryClassificationSystem 2.16.840.1.113883.1.11.16039; and US Census Occupation Code OID: 2.16.840.1.113883.6.243 for relevant recipient or custodian codes not included in this value set. If no source is indicated in the definition comments, then these are example FHIR codes. It can be extended with appropriate roles described by SNOMED as well as those described in the HL7 Role Based Access Control Catalog and the HL7 Healthcare (Security and Privacy) Access Control Catalog. In Role-Based Access Control (RBAC), permissions are operations on an object that a user wishes to access. Permissions are grouped into roles. A role characterizes the functions a user is allowed to perform. Roles are assigned to users. If the user's role has the appropriate permissions to access an object, then that user is granted access to the object. FHIR readily enables RBAC, as FHIR Resources are object types and the CRUDE events (the FHIR

equivalent to permissions in the RBAC scheme) are operations on those objects. In Attribute-Based Access Control (ABAC), a user requests to perform operations on objects. That user's access request is granted or denied based on a set of access control policies that are specified in terms of attributes and conditions. FHIR readily enables ABAC, as instances of a Resource in FHIR (again, Resources are object types) can have attributes associated with them. These attributes include security tags, environment conditions, and a host of user and object characteristics, which are the same attributes as those used in ABAC. Attributes help define the access control policies that determine the operations a user may perform on a Resource (in FHIR) or object (in ABAC). For example, a tag (or attribute) may specify that the identified Resource (object) is not to be further disclosed without explicit consent from the patient.)

• ValueSet: V3 Value SetRoleClassServiceDeliveryLocation (A role played by a place at which services may be provided.)

• ValueSet: V3 Value SetRoleClassMutualRelationship (A relationship that is based on mutual behavior of the two Entities as being related. The basis of such relationship may be agreements (e.g., spouses, contract parties) or they may bede facto behavior (e.g. friends) or may be an incidental involvement with each other (e.g. parties over a dispute, siblings, children).)

• ValueSet: V3 Value SetRoleClassPartitive (An association between two Entities where the playing Entity is considered in some way "part" of the scoping Entity, e.g., as a member, component, ingredient, or content. Being "part" in the broadest sense of the word can mean anything from being an integral structural component to a mere incidental temporary association of a playing Entity with a (generally larger) scoping Entity.)

• ValueSet: V3 Value SetRoleClassRoot (Corresponds to the Role class)

• ValueSet: V3 Value SetRoleClassPassive (An association for a playing Entity that is used, known, treated, handled, built, or destroyed, etc. under the auspices of the scoping Entity. The playing Entity is passive in these roles (even though it may be active in other roles), in the sense that the kinds of things done to it in this role happen without an agreement from the playing Entity.)

• ValueSet: V3 Value SetRoleClassRelationshipFormal (A relationship between two entities that is formally recognized, frequently by a contract or similar agreement.)

4.4.2.110.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition

1 | ROL | role | Corresponds to the Role class

2 | (_RoleClassAssociative) Abstract | | A general association between two entities that is neither partitive nor ontological.

3 | (_RoleClassMutualRelationship) Abstract | | A relationship that is based on mutual behavior of the two Entities as being related. The basis of such relationship may be agreements (e.g., spouses, contract parties) or they may be de facto behavior (e.g. friends) or may be an incidental involvement with each other (e.g. parties over a dispute, siblings, children).

4 | (_RoleClassRelationshipFormal) Abstract | | A relationship between two entities that is formally recognized, frequently by a contract or similar agreement.

5 | AFFL | affiliate | Player of the Affiliate role has a business/professional relationship with scoper. Player and scoper may be persons or organization. The Affiliate relationship does not imply membership in a group, nor does it exist for resource scheduling purposes. Example: A healthcare provider is affiliated with another provider as a business associate.

5 | AGNT | agent | An entity (player) that acts or is authorized to act on behalf of another entity (scoper).

6 | ASSIGNED | assigned entity | An agent role in which the agent is an Entity acting in the employ of an organization. The focus is on functional role on behalf of the organization, unlike the Employee role where the focus is on the 'Human Resources' relationship between the employee and the organization.

7 | COMPAR | commissioning party | An Entity that is authorized to issue or instantiate permissions, privileges, credentials or other formal/legal authorizations.

7 | SGNOFF | signing authority or officer | The role of a person (player) who is the officer or signature authority for of a scoping entity, usually an organization (scoper).

6 | CON | contact | A person or an organization (player) which provides or receives information regarding another entity (scoper). Examples; patient NOK and emergency contacts; guarantor contact; employer contact.

7 | ECON | emergency contact | An entity to be contacted in the event of an emergency.

7 | NOK | next of kin | An individual designated for notification as the next of kin for a given entity.

6 | GUARD | guardian | Guardian of a ward

5 | CIT | citizen | Citizen of apolitical entity

5 | COVPTY | covered party | A role class played by a person who receives benefit coverage under the terms of a particular insurance policy. The underwriter of that policy is the scoping entity. The covered party receives coverage because of some contractual or other relationship with the holder of that policy. Discussion:This reason for coverage is captured in 'Role.code' and a relationship link with type code of indirect authority should be included using the policy holder role as the source, and the covered party role as the target. Note that a particular policy may cover several individuals one of whom may be, but need not be, the policy holder. Thus the notion of covered party is a role that is distinct from that of the policy holder.

6 | CLAIM | claimant | Description: A role played by a party making a claim for coverage under a policy or program. A claimant must be either a person or organization, or a group of persons or organizations. A claimant is not a named insured or a program eligible. Discussion: With respect to liability insurance such as property and casualty insurance, a claimant must file a claim requesting indemnification for a loss that the claimant considers covered under the policy of a named insured. The claims adjuster for the policy underwriter will review the claim to determine whether the loss meets the benefit coverage criteria under a policy, and base any indemnification or coverage payment on that review. If a third party is liable in whole or part for the loss, the underwriter may pursue third party liability recovery. A claimant may be involved in civil or criminal legal proceedings involving claims against a defendant party that is indemnified by an insurance policy or to protest the finding of a claims adjustor. With

respect to life insurance, a beneficiary designated by a named insured becomes a claimant of the proceeds of coverage, as in the case of a life insurance policy. However, a claimant for coverage under life insurance is not necessarily a designated beneficiary. Note: A claimant is not a named insured. However, a named insured may make a claim under a policy, e.g., an insured driver may make a claim for an injury under his or her comprehensive automobile insurance policy. Similarly, a program eligible may make a claim under program, e.g., an unemployed worker may claim benefits under an unemployment insurance program, but parties playing these covered party role classes are not, for purposes of this vocabulary and in an effort to clearly distinguish role classes, considered claimants. In the case of a named insured making a claim, a role type code INSCLM (insured claimant) subtypes the class to indicate that either a named insured or an individual insured has filed a claim for a loss. In the case of a program eligible, a role type code INJWKR (injured worker) subtypes the class to indicate that the covered party in a workers compensation program is an injured worker, and as such, has filed a "claim" under the program for benefits. Likewise, a covered role type code UNEMP (unemployed worker) subtypes the program eligible class to indicate that the covered party in an unemployment insurance program has filed a claim for unemployment benefits. Example: A claimant under automobile policy that is not the named insured.

6 | NAMED | named insured | Description: A role played by a party to an insurance policy to which the insurer agrees to indemnify for losses, provides benefits for, or renders services. A named insured may be either a person, non-person living subject, or an organization, or a group of persons, non-person living subjects, or organizations. Discussion: The coded concept NAMED should not be used where a more specific child concept in this Specializable value set applies. In some cases, the named insured may not be the policy holder, e.g., where a policy holder purchases life insurance policy in which another party is the named insured and the policy holder is the beneficiary of the policy. Note: The party playing the role of a named insured is not a claimant in the sense conveyed by the RoleClassCoveredParty CLAIM (claimant). However, a named insured may make a claim under a policy, e.g., e.g., a party that is the named insured and policy holder under a comprehensive automobile insurance policy may become the claimant for coverage under that policy e.g., if injured in an automobile accident and there is no liable third party. In the case of a named insured making a claim, a role type code INSCLM (insured claimant) subtypes the class to indicate that a named insured has filed a claim for a loss. Example: The named insured under a comprehensive automobile, disability, or property and casualty policy that is the named insured and may or may not be the policy holder.

7 | DEPEN | dependent | Description: A role played by a person covered under a policy or program based on an association with a subscriber, which is recognized by the policy holder. Note: The party playing the role of a dependent is not a claimant in the sense conveyed by the RoleClassCoveredParty CLAIM (claimant). However, a dependent may make a claim under a policy, e.g., a dependent under a health insurance policy may become the claimant for coverage under that policy for wellness examines or if injured and there is no liable third party. In the case of a dependent making a claim, a role type code INSCLM (insured claimant) subtypes the class to indicate that the dependent has filed a claim for services covered under the health insurance policy. Example: The dependent has an association with the subscriber such as a financial dependency or personal relationship such as that of a spouse, or a natural or adopted child. The policy holder may be required by law to recognize certain associations or may have discretion about the associations. For example, a policy holder may dictate the criteria for the dependent status of adult children who are students, such as requiring full time enrollment, or may recognize domestic partners as dependents. Under certain circumstances, the dependent may be under the indirect authority of a responsible party acting as a surrogate for the subscriber, for example, if the subscriber is differently abled or deceased, a guardian ad Lidem or estate executor may be appointed to assume the subscriberaTMs legal standing in the relationship with the dependent.

7 | INDIV | individual | Description: A role played by a party covered under a policy as the policy holder. An individual may be either a person or an organization. Note: The party playing the role of an individual insured is not a claimant in the sense conveyed by the RoleClassCoveredParty CLAIM (claimant). However, a named insured may make a claim under a policy, e.g., a party that is the named insured and policy holder under a comprehensive automobile insurance policy may become the claimant for coverage under that policy if injured in an automobile accident and there is no liable third party. In the case of an individual insured making a claim, a role type code INSCLM (insured claimant) subtypes the class to indicate that an individual insured has filed a claim for a loss. Example: The individual insured under a comprehensive automobile, disability, or property and casualty policy that is the policy holder.

7 | SUBSCR | subscriber | Description: A role played by a person covered under a policy based on association with a sponsor who is the policy holder, and whose association may provide for the eligibility of dependents for coverage. Discussion: The policy holder holds the contract with the policy or program underwriter. The subscriber holds a certificate of coverage under the contract. In legal proceedings concerning the policy or program, the terms of the contract takes precedence over the terms of the certificate of coverage if there are any inconsistencies. Note: The party playing the role of a subscriber is not a claimant in the sense conveyed by the RoleClassCoveredParty CLAIM (claimant). However, a subscriber may make a claim under a policy, e.g., a subscriber under a health insurance policy may become the claimant for coverage under that policy for wellness examines or if injured and there is no liable third party. In the case of a subscriber making a claim, a role type code INSCLM (insured claimant) subtypes the class to indicate that the subscriber has filed a claim for services covered under the health insurance policy. Example: An employee or a member of an association.

6 | PROG | program eligible | Description: A role played by a party that meets the eligibility criteria for coverage under a program. A program eligible may be either a person, non-person living subject, or an organization, or a group of persons, non-person living subjects, or organizations. Discussion: A program as typically government administered coverage for parties determined eligible under the terms of the program. Note: The party playing a program eligible is not a claimant in the sense conveyed by the RoleClassCoveredParty CLAIM (claimant). However a program eligible may make a claim under program, e.g., an unemployed worker may claim benefits under an unemployment insurance program, but parties playing these covered party role classes are not, for purposes of this vocabulary and in an effort to clearly distinguish role classes, considered claimants. In the case of a program eligible, a role type code INJWKR (injured worker) subtypes the class to indicate that the covered party in a workers compensation program is an injured worker, and as such, has filed a "claim" under the program for benefits. Likewise, a covered role type code UNEMP (unemployed worker) subtypes the program eligible class to indicate that the covered party in an unemployment insurance program has filed a claim for unemployment benefits. Example: A party meeting eligibility criteria related to health or financial status, e.g., in the U.S., persons meeting health, demographic, or financial criteria established by state and federal law are eligible for Medicaid.

5 | CRINV | clinical research investigator | A role played by a provider, always a person, who has agency authority from a Clinical Research Sponsor to direct the conduct of a clinical research trial or study on behalf of the sponsor.

5 | CRSPNSR | clinical research sponsor | A role played by an entity, usually an organization, that is the sponsor of a clinical research trial or study. The sponsor commissions the study, bears the expenses, is responsible for satisfying all legal requirements concerning subject safety and privacy, and is generally responsible for collection, storage and analysis of the data generated during the trial. No scoper is necessary, as a clinical research sponsor undertakes the role on its own authority and declaration. Clinical research sponsors are usually educational or other research organizations, government agencies or biopharmaceutical companies.

5 | EMP | employee | A relationship between a person or organization and a person or organization formed for the purpose of exchanging work for compensation. The purpose of the role is to identify the type of relationship the employee has to the employer, rather than the nature of the work actually performed. (Contrast with AssignedEntity.)

6 | MIL | military person | A role played by a member of a military service. Scoper is the military service (e.g. Army, Navy, Air Force, etc.) or, more specifically, the unit (e.g. Company C, 3rd Battalion, 4th Division, etc.)

5 | GUAR | guarantor | A person or organization (player) that serves as a financial guarantor for another person or organization (scoper).

5 | INVSBJ | Investigation Subject | An entity that is the subject of an investigation. This role is scoped by the party responsible for the investigation.

6 | CASEBJ | Case Subject | A person, non-person living subject, or place that is the subject of an investigation related to a notifiable condition (health circumstance that is reportable within the applicable public health jurisdiction)

6 | RESBJ | research subject | Definition:Specifies the administrative functionality within a formal experimental design for which the ResearchSubject role was established. Examples: Screening - role is used for pre-enrollment evaluation portion of the design; enrolled - role is used for subjects admitted to the experimental portion of the design.

5 | LIC | licensed entity | A relationship in which the scoper certifies the player (e. g. a medical care giver, a medical device or a provider organization) to perform certain activities that fall under the jurisdiction of the scoper (e.g., a health authority licensing healthcare providers, a medical quality authority certifying healthcare professionals).

6 | NOT | notary public

6 | PROV | healthcare provider | An Entity (player) that is authorized to provide health care services by some authorizing agency (scoper).

5 | PAT | patient | A Role of a LivingSubject (player) as an actual or potential recipient of health care services from a healthcare provider organization (scoper). Usage Note: Communication about relationships between patients and specific healthcare practitioners (people) is not done via scoper. Instead this is generally done using the CareProvision act. This allows linkage between patient and a particular healthcare practitioner role and also allows description of the type of care involved in the relationship.

5 | PAYEE | payee | The role of an organization or individual designated to receive payment for a claim against a particular coverage. The scoping entity is the organization that is the submitter of the invoice in question.

5 | PAYOR | invoice payor | The role of an organization that undertakes to accept claims invoices, assess the coverage or payments due for those invoices and pay to the designated payees for those invoices. This role may be either the underwriter or a third-party organization authorized by the underwriter. The scoping entity is the organization that underwrites the claimed coverage.

5 | POLHOLD | policy holder | A role played by a person or organization that holds an insurance policy. The underwriter of that policy is the scoping entity. Discussion:The identifier of the policy is captured in 'Role.id' when the Role is a policy holder. A particular policy may cover several individuals one of whom may be, but need not be, the policy holder. Thus the notion of covered party is a role that is distinct from that of the policy holder.

5 | QUAL | qualified entity | An entity (player) that has been recognized as having certain training/experience or other characteristics that would make said entity an appropriate performer for a certain activity. The scoper is an organization that educates or qualifies entities.

5 | SPNSR | coverage sponsor | A role played by an entity, usually an organization that is the sponsor of an insurance plan or a health program. A sponsor is the party that is ultimately accountable for the coverage by employment contract or by law. A sponsor can be an employer, union, government agency, or association. Fully insured sponsors establish the terms of the plan and contract with health insurance plans to assume the risk and to administer the plan. Self-insured sponsors delegate coverage administration, but not risk, to third-party administrators. Program sponsors designate services to be covered in accordance with statute. Program sponsors may administer the coverage themselves, delegate coverage administration, but not risk to third-party administrators, or contract with health insurance plans to assume the risk and administrator a program. Sponsors qualify individuals who may become a policy holder of the plan; where the sponsor is the policy holder, who may become a subscriber or a dependent to a policy under the plan; or where the sponsor is a government agency, who may become program eligibles under a program. The sponsor role may be further qualified by the SponsorRole.code. Entities playing the sponsor role may also play the role of a Coverage Administrator. Example: An employer, union, government agency, or association.

5 | STD | student | A role played by an individual who is a student of a school, which is the scoping entity.

5 | UNDWRT | underwriter | A role played by a person or an organization. It is the party that accepts fiscal responsibility for insurance plans and the policies created under those plans; administers and accepts fiscal responsibility for a program that provides coverage for services to eligible individuals; and/or has the responsibility to assess the merits of each risk and decide a suitable premium for accepting all or part of the risk. If played by an organization, this role may be further specified by an appropriate RoleCode. Example: A health insurer; Medicaid Program; Lloyd's of London

4 | CAREGIVER | caregiver | A person responsible for the primary care of a patient at home.

4 | PRS | personal relationship | Links two entities with classCode PSN (person) in a personal relationship. The character of the relationship must be defined by a PersonalRelationshipRoleType code. The player and scoper are determined by PersonalRelationshipRoleType code as well.

4 | SELF | self | The "same" roleclass asserts an identity between playing and scoping entities: that they are in fact instances of the same entity and, in the case of discrepancies (e.g different DOB, gender), that one or both are in error. Usage: playing and scoping entities must have same classcode, but need not have identical attributes or values. Example: a provider registry maintains sets of conflicting demographic data for what is reported to be the same individual.

3 | (_RoleClassPassive) Abstract | | An association for a playing Entity that is used, known, treated, handled, built, or destroyed, etc. under the auspices of the scoping Entity. The playing Entity is passive in these roles (even though it may be active in other roles), in the sense that the kinds of things done to it in this role happen without an agreement from the playing Entity.

4 | ACCESS | access | A role in which the playing entity (material) provides access to another entity. The principal use case is intravenous (or other bodily) access lines that preexist and need to be referred to for medication routing instructions.

4 | ADJY | adjacency | A physical association whereby two Entities are in some (even lose) spatial relationship with each other such that they touch each other in some way. Examples: the colon is connected (and therefore adjacent) to the jejunum; the colon is adjacent to the liver (even if not actually connected.) UsageConstraints: Adjacency is in principle a symmetrical connection, but scoper and player of the role should, where applicable, be assigned to have scoper be the larger, more central Entity and player the smaller, more distant, appendage.

5 | CONC | connection | An adjacency of two Entities held together by a bond which attaches to each of the two entities. Examples: biceps brachii muscle connected to the radius bone, port 3 on a network switch connected to port 5 on a patch panel. UsageConstraints: See Adjacency for the assignment of scoper (larger, more central) and player (smaller, more distant).

6 | BOND | molecular bond | A connection between two atoms of a molecule. Examples: double bond between first and second C in ethane, peptide bond between two amino-acid, disulfide bridge between two proteins, chelate and ion associations, even the much weaker van-der-Waals bonds can be considered molecular bonds. UsageConstraints: See connection and adjacency for the assignment of player and scoper.

6 | CONY | continuity | A connection between two regional parts. Examples: the connection between ascending aorta and the aortic arc, connection between descending colon and sigmoid. UsageConstraints: See connection and adjacency for the assignment of player and scoper.
4 | ADMM | Administerable Material | A material (player) that can be administered to an Entity (scoper).
4 | BIRTHPL | birthplace | Relates a place (playing Entity) as the location where a living subject (scoping Entity) was born.
4 | DEATHPLC | place of death | Definition: Relates a place (playing Entity) as the location where a living subject (scoping Entity) died.
4 | DST | distributed material | A material (player) distributed by a distributor (scoper) who functions between a manufacturer and a buyer or retailer.
5 | RET | retailed material | Material (player) sold by a retailer (scoper), who also give advice to prospective buyers.
4 | EXLOC | event location | A role played by a place at which the location of an event may be recorded.
5 | SDLOC | service delivery location | A role played by a place at which services may be provided.
6 | DSDLOC | dedicated service delivery location | A role of a place (player) that is intended to house the provision of services. Scoper is the Entity (typically Organization) that provides these services. This is not synonymous with "ownership."
6 | ISDLOC | incidental service delivery location | A role played by a place at which health care services may be provided without prior designation or authorization.
4 | EXPR | exposed entity | A role played by an entity that has been exposed to a person or animal suffering a contagious disease, or with a location from which a toxin has been distributed. The player of the role is normally a person or animal, but it is possible that other entity types could become exposed. The role is scoped by the source of the exposure, and it is quite possible for a person playing the role of exposed party to also become the scoper a role played by another person. That is to say, once a person has become infected, it is possible, perhaps likely, for that person to infect others. Management of exposures and tracking exposed parties is a key function within public health, and within most public health contexts - exposed parties are known as "contacts."
4 | HLD | held entity | Entity that is currently in the possession of a holder (scoper), who holds, or uses it, usually based on some agreement with the owner.
4 | HLTHCHRT | health chart | The role of a material (player) that is the physical health chart belonging to an organization (scoper).
4 | IDENT | identified entity | A role in which the scoping entity designates an identifier for a playing entity.
4 | MANU | manufactured product | Scoped by the manufacturer
5 | THER | therapeutic agent | A manufactured material (player) that is used for its therapeutic properties. The manufacturer is the scoper.
4 | MNT | maintained entity | An entity (player) that is maintained by another entity (scoper). This is typical role held by durable equipment. The scoper assumes responsibility for proper operation, quality, and safety.
4 | OWN | owned entity | An Entity (player) for which someone (scoper) is granted by law the right to call the material (player) his own. This entitles the scoper to make decisions about the disposition of that material.
4 | RGPR | regulated product | A product regulated by some governmentatl orgnization. The role is played by Material and scoped by Organization. Rationale: To support an entity clone used to identify the NDC number for a drug product.
4 | TERR | territory of authority | Relates a place entity (player) as the region over which the scoper (typically an Organization) has certain authority (jurisdiction). For example, the Calgary Regional Health Authority (scoper) has authority over the territory "Region 4 of Alberta" (player) in matters of health.
4 | USED | used entity | Description:An entity (player) that is used by another entity (scoper)
4 | WRTE | warranted product | A role a product plays when a guarantee is given to the purchaser by the seller (scoping entity) stating that the product is reliable and free from known defects and that the seller will repair or replace defective parts within a given time limit and under certain conditions.
2 | (_RoleClassOntological) Abstract | | A relationship in which the scoping Entity defines or specifies what the playing Entity is. Thus, the player's "being" (Greek: ontos) is specified.
3 | EQUIV | equivalent entity | Description: Specifies the player Entity (the equivalent Entity) as an Entity that is considered to be equivalent to a reference Entity (scoper). The equivalence is in principle a symmetric relationship, however, it is expected that the scoper is a reference entity which serves as reference entity for multiple different equivalent entities. Examples: An innovator's medicine formulation is the reference for "generics", i.e., formulations manufactured differently but having been proven to be biologically equivalent to the reference medicine. Another example is a reference ingredient that serves as basis for quantity specifications (basis of strength, e.g., metoprolol succinate specified in terms of metoprolol tartrate.)
4 | SAME | same | The "same" role asserts an identity between playing and scoping entities, i.e., that they are in fact two records of the same entity instance, and, in the case of discrepancies (e.g different DOB, gender), that one or both are in error. Usage: playing and scoping entities must have same classCode, but need not have identical attributes or values. Example: a provider registry maintains sets of conflicting demographic data for what is reported to be the same individual.
4 | SUBY | subsumed by | Relates a prevailing record of an Entity (scoper) with another record (player) that it subsumes. Examples: Show a correct new Person object (scoper) that subsumes one or more duplicate Person objects that had accidentally been created for the same physical person. Constraints: Both the player and scoper must have the same classCode.
3 | GEN | has generalization | Relates a specialized material concept (player) to its generalization (scoper).
4 | GRIC | has generic | A special link between pharmaceuticals indicating that the target (scoper) is a generic for the source (player).
3 | INST | instance | An individual piece of material (player) instantiating a class of material (scoper).
3 | SUBS Deprecated | subsumer | An entity that subsumes the identity of another. Used in the context of merging documented entity instances. Both the player and scoper must have the same classCode. The use of this code is deprecated in favor of the term SUBY which is its inverse and is more ontologically correct.
2 | (_RoleClassPartitive) Abstract | | An association between two Entities where the playing Entity is considered in some way "part" of the scoping Entity, e.g., as a member, component, ingredient, or content. Being "part" in the broadest sense of the word can mean anything from being an integral structural component to a mere incidental temporary association of a playing Entity with a (generally larger) scoping Entity.
3 | CONT | content | Relates a material as the content (player) to a container (scoper). Unlike ingredients, the content and a container remain separate (not mixed) and the content can be removed from the container. A content is not part of an empty container.
3 | EXPAGTCAR | exposure agent carrier | An exposure agent carrier is an entity that is capable of conveying an exposure agent from one entity to another. The scoper of the role must be the exposure agent (e.g., pathogen).
4 | EXPVECTOR | exposure vector | Description: A vector is a living subject that carries an exposure agent. The vector does not cause the disease itself, but exposes targets to the exposure agent. A mosquito carrying malaria is an example of a vector. The

scoper of the role must be the exposure agent (e.g., pathogen).
4 | FOMITE | fomite | Description: A fomite is a non-living entity that is capable of conveying exposure agent from one entity to another. A doorknob contaminated with a Norovirus is an example of a fomite. Anyone touching the doorknob would be exposed to the virus. The scoper of the role must be the exposure agent (e.g., pathogen).
3 | INGR | ingredient | Relates a component (player) to a mixture (scoper). E.g., Glucose and Water are ingredients of D5W, latex may be an ingredient in a tracheal tube.
4 | ACTI | active ingredient | Definition: a therapeutically active ingredient (player) in a mixture (scoper), where the mixture is typically a manufactured pharmaceutical. It is unknown if the quantity of such an ingredient is expressed precisely in terms of the playing ingredient substance, or, if it is specified in terms of a closely related substance (active moiety or reference substance).
5 | ACTIB | active ingredient - basis of strength | Description: Active ingredient, where the ingredient substance (player) is itself the "basis of strength", i.e., where the Role.quantity specifies exactly the quantity of the player substance in the medicine formulation. Examples: Lopressor 50 mg actually contains 50 mg of metoprolol succinate, even though the active moiety is metoprolol, but also: Tenormin 50 mg contain 50 mg of atenolol, as free base, i.e., where the active ingredient atenolol is also the active moiety.
5 | ACTIM | active ingredient - moiety is basis of strength | Description: Active ingredient, where not the ingredient substance (player), but itaTMs active moiety is the "basis of strength", i.e., where the Role.quantity specifies the quantity of the player substance's active moiety in the medicine formulation. Examples: 1 mL of Betopic 5mg/mL eye drops contains 5.6 mg betaxolol hydrochloride equivalent to betaxolol base 5 mg.
5 | ACTIR | active ingredient - reference substance is basis of strength | Description: Active ingredient, where not the ingredient substance (player) but another reference substance with the same active moiety, is the "basis of strength", i.e., where the Role.quantity specifies the quantity of a reference substance, similar but different from the player substance's in the medicine formulation. Examples: Toprol-XL 50 mg contains 47.5 mg of metoprolol succinate equivalent to 50 mg of metoprolol tartrate.
4 | ADJV | adjuvant | A component (player) added to enhance the action of an active ingredient (scoper) (in the manner of a catalyst) but which has no active effect in and of itself. Such ingredients are significant in defining equivalence of products in a way that inactive ingredients are not.
4 | ADTV | additive | An ingredient (player) that is added to a base (scoper), that amounts to a minor part of the overall mixture.
4 | BASE | base | A base ingredient (player) is what comprises the major part of a mixture (scoper). E.g., Water in most i.v. solutions, or Vaseline in salves. Among all ingredients of a material, there should be only one base. A base substance can, in turn, be a mixture.
4 | CNTM | contaminant ingredient | An ingredient whose presence is not intended but may not be reasonably avoided given the circumstances of the mixture's nature or origin.
4 | IACT | inactive ingredient | An ingredient which is not considered therapeutically active, e.g., colors, flavors, stabilizers, or preservatives, fillers, or structural components added to an active ingredient in order to facilitate administration of the active ingredient but without being considered therapeutically active. An inactive ingredient need not be biologically inert, e.g., might be active as an allergen or might have a pleasant taste, but is not an essential constituent delivering the therapeutic effect.
5 | COLR | color additive | A substance (player) influencing the optical aspect of material (scoper).
5 | FLVR | flavor additive | A substance (player) added to a mixture (scoper) to make it taste a certain way. In food the use is obvious, in pharmaceuticals flavors can hide disgusting taste of the active ingredient (important in pediatric treatments).
5 | PRSV | preservative | A substance (player) added to a mixture (scoper) to prevent microorganisms (fungi, bacteria) to spoil the mixture.
5 | STBL | stabilizer | A stabilizer (player) added to a mixture (scoper) in order to prevent the molecular disintegration of the main substance.
4 | MECH | mechanical ingredient | An ingredient (player) of a medication (scoper) that is inseparable from the active ingredients, but has no intended chemical or pharmaceutical effect itself, but which may have some systemic effect on the patient. An example is a collagen matrix used as a base for transplanting skin cells. The collagen matrix can be left permanently in the graft site. Because it is of bovine origin, the patient may exhibit allergies or may have cultural objections to its use.
3 | LOCE | located entity | Relates an entity (player) to a location (scoper) at which it is present in some way. This presence may be limited in time.
4 | STOR | stored entity | Relates an entity (player) (e.g. a device) to a location (scoper) at which it is normally found or stored when not used.
3 | MBR | member | A role played by an entity that is a member of a group. The group provides the scope for this role. Among other uses, groups as used in insurance (groups of covered individuals) and in scheduling where resources may be grouped for scheduling and logistical purposes.
3 | PART | part | Definition: an association between two Entities where the playing Entity (the part) is a component of the whole (scoper) in the sense of an integral structural component, that is distinct from other parts in the same whole, has a distinct function in the whole, and, as an effect, the full integrity of the whole depends (to some degree) on the presence of this part, even though the part may often be separable from the whole. Discussion: Part is defined in opposition to (a) ingredient (not separable), (b) content (not a functional component), and (c) member (not functionally distinct from other members).
4 | ACTM | active moiety | The molecule or ion that is responsible for the intended pharmacological action of the drug substance, excluding those appended or associated parts of the molecule that make the molecule an ester, salt (including a salt with hydrogen or coordination bonds), or other noncovalent derivative (such as a complex, chelate, or clathrate). Examples: heparin-sodium and heparin-potassium have the same active moiety, heparin; the active moiety of morphine-hydrochloride is morphine.
3 | SPEC | specimen | A role played by a material entity that is a specimen for an act. It is scoped by the source of the specimen.
4 | ALQT | aliquot | A portion (player) of an original or source specimen (scoper) used for testing or transportation.
4 | ISLT | isolate | A microorganism that has been isolated from other microorganisms or a source matrix.
1 | CHILD | child | The player of the role is a child of the scoping entity, in a generic sense.
1 | CRED | credentialed entity | A role played by an entity that receives credentials from the scoping entity.
1 | NURPRAC | nurse practitioner
1 | NURS | nurse
1 | PA | physician assistant
1 | PHYS | physician

4.4.2.111 HL7 v3 Code System RoleCode

URL: https://hl7.org/fhir/R4/v3/RoleCode/cs.html
Vocabulary Work Group | Maturity Level: N/A | External | Use Context: Any

This code system (http://terminology.hl7.org/CodeSystem/v3-RoleCode) is defined as part of HL7 v3.

Summary

Defining URL: | http://terminology.hl7.org/CodeSystem/v3-RoleCode
Version: | 2018-08-12
Name: | v3.RoleCode
Title: | v3 Code System RoleCode
Definition: | A set of codes further specifying the kind of Role; specific classification codes for further qualifying RoleClass codes.
OID: | 2.16.840.1.113883.5.111 (for OID based terminology systems)
Source Resource | XML / JSON

This Code system is used in the following value sets:

• ValueSet: Parent Relationship Codes (The value set includes the v3 RoleCode PRN (parent), TWIN (twin) and all of their specializations. It covers the relationships needed to establish genetic pedigree relationships between family members.)

• ValueSet: Parent Relationship Codes (The value set includes the v3 RoleCode PRN (parent), TWIN (twin) and all of their specializations. It covers the relationships needed to establish genetic pedigree relationships between family members.)

• ValueSet: V3 Value SetFamilyMember (A relationship between two people characterizing their "familial" relationship)

• ValueSet: V3 Value SetMilitaryRoleType (Definition: A person playing the role of program eligible under a program based on military status. Discussion: This CoveredPartyRoleType.code is typically used when the CoveredPartyRole class code is either "program eligible" or "subscriber" and the person's status as a member of the military meets jurisdictional or program criteria)

• ValueSet: Focal Subject Codes (Example value set composed from SNOMED CT and HL7 V3 codes for observation targets such as donor, fetus or spouse. As use cases are discovered, more values may be added.)

• ValueSet: ParticipationRoleType (This FHIR value set is comprised of Actor participation Type codes, which can be used to value FHIR agents, actors, and other role elements. The FHIR Actor participation type value set is based on DICOM Audit Message, C402; ASTM Standard, E1762-95 [2013]; selected codes and derived actor roles from HL7 RoleClass OID 2.16.840.1.113883.5.110; HL7 Role Code 2.16.840.1.113883.5.111, including AgentRoleType; HL7 ParticipationType OID: 2.16.840.1.113883.5.90; and HL7 ParticipationFunction codes OID: 2.16.840.1.113883.5.88. This value set includes, by reference, role codes from external code systems: NUCC Health Care Provider Taxonomy OID: 2.16.840.1.113883.6.101; North American Industry Classification System [NAICS]OID: 2.16.840.1.113883.6.85; IndustryClassificationSystem 2.16.840.1.113883.1.11.16039; and US Census Occupation Code OID: 2.16.840.1.113883.6.243 for relevant recipient or custodian codes not included in this value set. If no source is indicated in the definition comments, then these are example FHIR codes.)

• ValueSet: ParticipationRoleType (This FHIR value set is comprised of Actor participation Type codes, which can be used to value FHIR agents, actors, and other role elements. The FHIR Actor participation type value set is based on DICOM Audit Message, C402; ASTM Standard, E1762-95 [2013]; selected codes and derived actor roles from HL7 RoleClass OID 2.16.840.1.113883.5.110; HL7 Role Code 2.16.840.1.113883.5.111, including AgentRoleType; HL7 ParticipationType OID: 2.16.840.1.113883.5.90; and HL7 ParticipationFunction codes OID: 2.16.840.1.113883.5.88. This value set includes, by reference, role codes from external code systems: NUCC Health Care Provider Taxonomy OID: 2.16.840.1.113883.6.101; North American Industry Classification System [NAICS]OID: 2.16.840.1.113883.6.85; IndustryClassificationSystem 2.16.840.1.113883.1.11.16039; and US Census Occupation Code OID: 2.16.840.1.113883.6.243 for relevant recipient or custodian codes not included in this value set. If no source is indicated in the definition comments, then these are example FHIR codes.)

• ValueSet: SecurityRoleType (This example FHIR value set is comprised of example Actor Type codes, which can be used to value FHIR agents, actors, and other role elements such as those specified in financial transactions. The FHIR Actor value set is based on DICOM Audit Message, C402; ASTM Standard, E1762-95 [2013]; selected codes and derived actor roles from HL7 RoleClass OID 2.16.840.1.113883.5.110; HL7 Role Code 2.16.840.1.113883.5.111, including AgentRoleType; HL7 ParticipationType OID: 2.16.840.1.113883.5.90; and HL7 ParticipationFunction codes OID: 2.16.840.1.113883.5.88. This value set includes, by reference, role codes from external code systems: NUCC Health Care Provider Taxonomy OID: 2.16.840.1.113883.6.101; North American Industry Classification System [NAICS]OID: 2.16.840.1.113883.6.85; IndustryClassificationSystem 2.16.840.1.113883.1.11.16039; and US Census Occupation Code OID: 2.16.840.1.113883.6.243 for relevant recipient or custodian codes not included in this value set. If no source is indicated in the definition comments, then these are example FHIR codes. It can be extended with appropriate roles described by SNOMED as well as those described in the HL7 Role Based Access Control Catalog and the HL7 Healthcare (Security and Privacy) Access Control Catalog. In Role-Based Access Control (RBAC), permissions are operations on an object that a user wishes to access. Permissions are grouped into roles. A role characterizes the functions a user is allowed to perform. Roles are assigned to users. If the user's role has the appropriate permissions to access an object, then that user is granted access to the object. FHIR readily enables RBAC, as FHIR Resources are object types and the CRUDE events (the FHIR

equivalent to permissions in the RBAC scheme) are operations on those objects. In Attribute-Based Access Control (ABAC), a user requests to perform operations on objects. That user's access request is granted or denied based on a set of access control policies that are specified in terms of attributes and conditions. FHIR readily enables ABAC, as instances of a Resource in FHIR (again, Resources are object types) can have attributes associated with them. These attributes include security tags, environment conditions, and a host of user and object characteristics, which are the same attributes as those used in ABAC. Attributes help define the access control policies that determine the operations a user may perform on a Resource (in FHIR) or object (in ABAC). For example, a tag (or attribute) may specify that the identified Resource (object) is not to be further disclosed without explicit consent from the patient.)

• ValueSet: SecurityRoleType (This example FHIR value set is comprised of example Actor Type codes, which can be used to value FHIR agents, actors, and other role elements such as those specified in financial transactions. The FHIR Actor value set is based on DICOM Audit Message, C402; ASTM Standard, E1762-95 [2013]; selected codes and derived actor roles from HL7 RoleClass OID 2.16.840.1.113883.5.110; HL7 Role Code 2.16.840.1.113883.5.111, including AgentRoleType; HL7 ParticipationType OID: 2.16.840.1.113883.5.90; and HL7 ParticipationFunction codes OID: 2.16.840.1.113883.5.88. This value set includes, by reference, role codes from external code systems: NUCC Health Care Provider Taxonomy OID: 2.16.840.1.113883.6.101; North American Industry Classification System [NAICS]OID: 2.16.840.1.113883.6.85; IndustryClassificationSystem 2.16.840.1.113883.1.11.16039; and US Census Occupation Code OID: 2.16.840.1.113883.6.243 for relevant recipient or custodian codes not included in this value set. If no source is indicated in the definition comments, then these are example FHIR codes. It can be extended with appropriate roles described by SNOMED as well as those described in the HL7 Role Based Access Control Catalog and the HL7 Healthcare (Security and Privacy) Access Control Catalog. In Role-Based Access Control (RBAC), permissions are operations on an object that a user wishes to access. Permissions are grouped into roles. A role characterizes the functions a user is allowed to perform. Roles are assigned to users. If the user's role has the appropriate permissions to access an object, then that user is granted access to the object. FHIR readily enables RBAC, as FHIR Resources are object types and the CRUDE events (the FHIR equivalent to permissions in the RBAC scheme) are operations on those objects. In Attribute-Based Access Control (ABAC), a user requests to perform operations on objects. That user's access request is granted or denied based on a set of access control policies that are specified in terms of attributes and conditions. FHIR readily enables ABAC, as instances of a Resource in FHIR (again, Resources are object types) can have attributes associated with them. These attributes include security tags, environment conditions, and a host of user and object characteristics, which are the same attributes as those used in ABAC. Attributes help define the access control policies that determine the operations a user may perform on a Resource (in FHIR) or object (in ABAC). For example, a tag (or attribute) may specify that the identified Resource (object) is not to be further disclosed without explicit consent from the patient.)

• ValueSet: V3 Value SetServiceDeliveryLocationRoleType (A role of a place that further classifies the setting (e.g., accident site, road side, work site, community location) in which services are delivered.)

• ValueSet: Sibling Relationship Codes (The value set includes the v3 RoleCode SIB (sibling) and all of its specializations. It covers the relationships needed to establish genetic pedigree relationships between family members.)

• ValueSet: Sibling Relationship Codes (The value set includes the v3 RoleCode SIB (sibling) and all of its specializations. It covers the relationships needed to establish genetic pedigree relationships between family members.)

• ValueSet: Patient relationship type (A set of codes that can be used to indicate the relationship between a Patient and a Related Person.)

• ValueSet: v3 Code System RoleCode (A set of codes further specifying the kind of Role; specific classification codes for further qualifying RoleClass codes.)

• ValueSet: V3 Value SetPersonalRelationshipRoleType (Types of personal relationships between two living subjects. Example: Parent, sibling, unrelated friend, neighbor)

• ValueSet: Patient relationship type (A set of codes that can be used to indicate the relationship between a Patient and a Related Person.)

• ValueSet: Focal Subject Codes (Example value set composed from SNOMED CT and HL7 V3 codes for observation targets such as donor, fetus or spouse. As use cases are discovered, more values may be added.)

4.4.2.111.1 Code System Content

Release Date: 2018-08-12

Level | Code | Display | Definition

1 | (_AffiliationRoleType) Abstract | | Concepts characterizing the type of association formed by player and scoper when there is a recognized Affiliate role by which the two parties are related. Examples: Business Partner, Business Associate, Colleague

2 | (_AgentRoleType) Abstract | | Parties that may or should contribute or have contributed to an Act.

3 | AMENDER | amender | An entity which corrected, edited, or amended pre-existing information.

3 | CLASSIFIER | classifier | An individual authorized to assign an original classification to information, including compilations of unclassified information, based on a determination that the information requires protection against unauthorized disclosure. The individual marks the information with immutable, computable, and human readable security labels in accordance with applicable

security labeling policies. The labeling policies provide instructions on whether and if so how the security labels may be later reclassified [i.e., upgraded, downgraded, used in derivative classification, or declassified] in a manner that preserves the overridden original classification binding and provenance.

3 | CONSENTER | consenter | An entity or an entity's delegatee who is the grantee in an agreement such as a consent for services, advanced directive, or a privacy consent directive in accordance with jurisdictional, organizational, or patient policy.

3 | CONSWIT | consent witness | An entity which has witnessed and attests to observing another entity being counseled about an agreement such as a consent for services, advanced directive, or a privacy consent directive.

3 | COPART | co-participant | An entity which participates in the generation of and attest to veracity of content, but is not an author or coauthor. For example a surgeon who is required by institutional, regulatory, or legal rules to sign an operative report, but who was not involved in the authorship of that report.

3 | DECLASSIFIER | declassifier | An individual which is authorized to declassify information based on a determination that the information no longer requires protection against unauthorized disclosure. The individual marks the information being declassified using computable and human readable security labels indicating that this is copy of previously classified information is unclassified in accordance with applicable security labeling policies. The labeling policies provide instructions on whether and if so how the security labels may be later reclassified [i.e., upgraded or used in derivative classification] in a manner that preserves the overridden original classification binding and provenance.

3 | DELEGATEE | delegatee | A party to whom some right or authority is granted by a delegator.

3 | DELEGATOR | delegator | A party that grants all or some portion its right or authority to another party.

3 | DOWNGRDER | downgrader | An individual authorized to lower the classification level of labeled content and provide rationale for doing so as directed by a classification guide.

3 | DRIVCLASSIFIER | derivative classifier | An individual who is only authorized to classify reproduced, extracted, or summarized classified information, or compile classified and unclassified information by applying classification markings derived from source material or as directed by a classification guide.

3 | GRANTEE | grantee | An entity which accepts certain rights or authority from a grantor.

3 | GRANTOR | grantor | An entity which agrees to confer certain rights or authority to a grantee.

3 | INTPRTER | interpreter | An entity which converts spoken or written language into the language of key participants in an event such as when a provider is obtaining a patient's consent to treatment or permission to disclose information.

3 | REVIEWER | reviewer | An entity authorized to filter information according to approved criteria.

3 | VALIDATOR | validator | An entity authorized to validate information for inclusion in a record.

2 | (_CoverageSponsorRoleType) Abstract | | Description:Codes that indicate a specific type of sponsor. Used when the sponsor's role is only either as a fully insured sponsor or only as a self-insured sponsor. NOTE: Where a sponsor may be either, use the SponsorParticipationFunction.code (fully insured or self insured) to indicate the type of responsibility. (CO6-0057)

3 | FULLINS | Fully insured coverage sponsor | Description:An employer or organization that contracts with an underwriter to assumes the financial risk and administrative responsibility for coverage of health services for covered parties.

3 | SELFINS | Self insured coverage sponsor | Description:An employer or organization that assumes the financial risk and administrative responsibility for coverage of health services for covered parties.

2 | (_PayorRoleType) Abstract | | Description:PayorRoleType for a particular type of policy or program benefit package or plan where more detail about the coverage administration role of the Payor is required. The functions performed by a Payor qualified by a PayorRoleType may be specified by the PayorParticpationFunction value set. Examples:A Payor that is a TPA may administer a managed care plan without underwriting the risk.

3 | ENROLBKR | Enrollment Broker | Description:A payor that is responsible for functions related to the enrollment of covered parties.

3 | TPA | Third party administrator | Description:Third party administrator (TPA) is a payor organization that processes health care claims without carrying insurance risk. Third party administrators are prominent players in the managed care industry and have the expertise and capability to administer all or a portion of the claims process. They are normally contracted by a health insurer or self-insuring companies to administer services, including claims administration, premium collection, enrollment and other administrative activities. Self-insured employers often contract with third party administrator to handle their insurance functions. Insurance companies oftentimes outsource the claims, utilization review or membership functions to a TPA. Sometimes TPAs only manage provider networks, only claims or only utilization review. While some third-party administrators may operate as units of insurance companies, they are often independent. However, hospitals or provider organizations desiring to set up their own health plans will often outsource certain responsibilities to TPAs. TPAs may perform one or several payor functions, specified by the PayorParticipationFunction value set, such as provider management, enrollment, utilization management, and fee for service claims adjudication management.

3 | UMO | Utilization management organization | Description:A payor that is responsible for review and case management of health services covered under a policy or program.

2 | RESPRSN | responsible party | The role played by a party who has legal responsibility for another party.

3 | EXCEST | executor of estate | The role played by a person acting as the estate executor for a deceased subscriber or policyholder who was the responsible party

3 | GUADLTM | guardian ad lidem | The role played by a person appointed by the court to look out for the best interests of a minor child during the course of legal proceedings.

3 | GUARD | guardian | The role played by a person or institution legally empowered with responsibility for the care of a ward.

3 | POWATT | power of attorney | A relationship between two people in which one person authorizes another to act for him in a manner which is a legally binding upon the person giving such authority as if he or she personally were to do the acts.

4 | DPOWATT | durable power of attorney | A relationship between two people in which one person authorizes another, usually a family member or relative, to act for him or her in a manner which is a legally binding upon the person giving such authority as if he or she personally were to do the acts that is often limited in the kinds of powers that can be assigned. Unlike ordinary powers of attorney, durable powers can survive for long periods of time, and again, unlike standard powers of attorney, durable powers can continue after incompetency.

4 | HPOWATT | healthcare power of attorney | A relationship between two people in which one person authorizes another to act for him or her in a manner which is a legally binding upon the person giving such authority as if he or she personally were to do the acts that continues (by its terms) to be effective even though the grantor has become mentally incompetent after signing the document.

4 | SPOWATT | special power of attorney | A relationship between two people in which one person authorizes another to act for him or her in a manner which is a legally binding upon the person giving such authority as if he or she personally were to do the acts that is often limited in the kinds of powers that can be assigned.

1 | (_AssignedRoleType) Abstract

2 | (_AssignedNonPersonLivingSubjectRoleType) Abstract | | Description:A role type that is used to further qualify a non-person subject playing a role where the role class attribute is set to RoleClass AssignedEntity
3 | ASSIST | Assistive non-person living subject | Description:Dogs trained to assist the ill or physically challenged.
3 | BIOTH | Biotherapeutic non-person living subject | Description:Animals, including fish and insects, and microorganisms which may participate as assigned entities in biotherapies.
4 | ANTIBIOT | Antibiotic | Description:Non-person living subject used as antibiotic. Examples:Bacteriophage, is a virus that infects bacteria.
4 | DEBR | Debridement | Description:Maggots raised for biodebridement. Discussion: Maggot Debridement Therapy is the medical use of live maggots for cleaning non-healing wounds. Examples:Removal of burnt skin.
3 | CCO | Clinical Companion | Description:Companion animals, such as dogs, cats, and rabbits, which may be provided to patients to improve general mood, decrease depression and loneliness, and distract from stress-inducing concerns to improve quality of life.
3 | SEE | Seeing | Description:Dogs trained to assist persons who are seeing impaired or blind.
3 | SNIFF | Sniffing | Description:Dogs trained or having the ability to detect imminent seizures or cancers in humans, probably as a result of volatile chemical (odors) given off by the malignancy of the host.
1 | (_CertifiedEntityType) Abstract | | Defines types of certifications for all entities
1 | _CitizenRoleType | CitizenRoleType | A role type used to qualify a person's legal status within a country or nation.
2 | CAS | asylum seeker | A person who has fled his or her home country to find a safe place elsewhere.
3 | CASM | single minor asylum seeker | A person who is someone of below legal age who has fled his or her home country, without his or her parents, to find a safe place elsewhere at time of categorization.
2 | CN | national | A person who is legally recognized as a member of a nation or country, with associated rights and obligations.
2 | CNRP | non-country member without residence permit | A foreigner who is present in a country (which is foreign to him/her) unlawfully or without the country's authorization (may be called an illegal alien).
3 | CNRPM | non-country member minor without residence permit | A person who is below legal age present in a country, without his or her parents, (which is foreign to him/her) unlawfully or without the country's authorization.
2 | CPCA | permit card applicant | A non-country member admitted to the territory of a nation or country as a non-resident for an explicit purpose.
2 | CRP | non-country member with residence permit | A foreigner who is a resident of the country but does not have citizenship.
3 | CRPM | non-country member minor with residence permit | A person who is a resident below legal age of the country without his or her parents and does not have citizenship.
1 | (_ContactRoleType) Abstract | | Types of contact for Role code "CON"
2 | (_AdministrativeContactRoleType) Abstract | | A contact role used for business and/or administrative purposes.
3 | BILL | Billing Contact | A contact role used to identify a person within a Provider organization that can be contacted for billing purposes (e.g. about the content of the Invoice).
3 | ORG | organizational contact | A contact for an organization for administrative purposes. Contact role specifies a person acting as a liason for the organization. Example: HR Department representative.
3 | PAYOR | Payor Contact | A contact role used to identify a person within a Payor organization to whom this communication is addressed.
2 | ECON Deprecated | emergency contact | A contact designated for contact in emergent situations.
2 | NOK Deprecated | next of kin | Played by an individual who is designated as the next of kin for another individual which scopes the role.
1 | (_IdentifiedEntityType) Abstract | | Definition: A code representing the type of identifier that has been assigned to the identified entity (IDENT). Examples: Example values include Social Insurance Number, Product Catalog ID, Product Model Number.
2 | (_LocationIdentifiedEntityRoleCode) Abstract | | Description:Describes types of identifiers other than the primary location registry identifier for a service delivery location. Identifiers may be assigned by a local service delivery organization, a formal body capable of accrediting the location for the capability to provide specific services or the identifier may be assigned at a jurisdictional level.
3 | ACHFID | accreditation location identifier | Description:Identifier assigned to a location by the organization responsible for accrediting the location.
3 | JURID | jurisdiction location identifier | Description:Identifier assigned to a location by a jurisdiction.
3 | LOCHFID | local location identifier | Description:Identifier assigned to a location by a local party (which could be the facility itself or organization overseeing a group of facilities).
1 | (_LivingSubjectProductionClass) Abstract | | Code indicating the primary use for which a living subject is bred or grown
2 | BF | Beef | Cattle used for meat production
2 | BL | Broiler | Chickens raised for meat
2 | BR | Breeder | Breeding/genetic stock
2 | CO | Companion | Companion animals
2 | DA | Dairy | Milk production
2 | DR | Draft | Draft animals
2 | DU | Dual | Dual purpose. Defined purposes based on species and breed
2 | FI | Fiber | Animals raised for their fur, hair or skins
2 | LY | Layer | Chickens raised for egg production
2 | MT | Meat | Animals raised for meat production
2 | MU | Multiplier | Poultry flocks used for chick/poult production
2 | PL | Pleasure | Animals rasied for recreation
2 | RC | Racing | Animals raised for racing perfomance
2 | SH | Show | Animals raised for shows
2 | VL | Veal | Cattle raised for veal meat production. Implicit is the husbandry method.
2 | WL | Wool | Sheep, goats and other mammals raised for their fiber
2 | WO | Working | Animals used to perform work
1 | (_MedicationGeneralizationRoleType) Abstract | | Identifies the specific hierarchical relationship between the playing and scoping medications. Examples: Generic, Generic Formulation, Therapeutic Class, etc.
2 | DC | therapeutic class | Description:A categorization of medicinal products by their therapeutic properties and/or main therapeutic use.

2 | GD | generic drug | Relates a manufactured drug product to the non-proprietary (generic) representation of its ingredients independent of strength, and form. The scoping entity identifies a unique combination of medicine ingredients; sometimes referred to as "ingredient set".
3 | GDF | generic drug form | Relates a manufactured drug product to the non-proprietary (generic) representation of its ingredients and dose form, independent of strength of the ingredients. The scoping entity identifies a unique combination of medicine ingredients in a specific dose form.
3 | GDS | generic drug strength | Relates a manufactured drug product to the non-proprietary (generic) representation of is ingredients with their strength. The scoping entity identifies a unique combination of medicine ingredients with their strength.
3 | GDSF | generic drug strength form | Relates a manufactured drug product to the non-proprietary (generic) representation of its ingredients with their strength in a specific dose form. The scoping entity identifies a unique combination of medicine ingredients with their strength in a single dose form.
2 | MGDSF | manufactured drug strength form | Relates a manufactured drug product to the non-proprietary (generic) representation of its ingredients with their strength in a specific dose form. The scoping entity identifies a unique combination of medicine ingredients with their strength in a single dose form.
1 | (_MemberRoleType) Abstract | | Types of membership for Role code "MBR"
2 | TRB | Tribal Member | A person who is a member of a tribe.
1 | (_PersonalRelationshipRoleType) Abstract
2 | FAMMEMB | family member | A relationship between two people characterizing their "familial" relationship
3 | CHILD | child | The player of the role is a child of the scoping entity.
4 | CHLDADOPT | adopted child | The player of the role is a child taken into a family through legal means and raised by the scoping person (parent) as his or her own child.
5 | DAUADOPT | adopted daughter | The player of the role is a female child taken into a family through legal means and raised by the scoping person (parent) as his or her own child.
5 | SONADOPT | adopted son | The player of the role is a male child taken into a family through legal means and raised by the scoping person (parent) as his or her own child.
4 | CHLDFOST | foster child | The player of the role is a child receiving parental care and nurture from the scoping person (parent) but not related to him or her through legal or blood ties.
5 | DAUFOST | foster daughter | The player of the role is a female child receiving parental care and nurture from the scoping person (parent) but not related to him or her through legal or blood ties.
5 | SONFOST | foster son | The player of the role is a male child receiving parental care and nurture from the scoping person (parent) but not related to him or her through legal or blood ties.
4 | DAUC | daughter | Description: The player of the role is a female child (of any type) of scoping entity (parent)
5 | DAUADOPT
5 | DAUFOST
5 | DAU | natural daughter | The player of the role is a female offspring of the scoping entity (parent).
5 | STPDAU | stepdaughter | The player of the role is a daughter of the scoping person's spouse by a previous union.
4 | NCHILD | natural child | The player of the role is an offspring of the scoping entity as determined by birth.
5 | DAU
5 | SON | natural son | The player of the role is a male offspring of the scoping entity (parent).
4 | SONC | son | Description: The player of the role is a male child (of any type) of scoping entity (parent)
5 | SONADOPT
5 | SONFOST
5 | SON
5 | STPSON | stepson | The player of the role is a son of the scoping person's spouse by a previous union.
4 | STPCHLD | step child | The player of the role is a child of the scoping person's spouse by a previous union.
5 | STPDAU
5 | STPSON
3 | EXT | extended family member | Description: A family member not having an immediate genetic or legal relationship e.g. Aunt, cousin, great grandparent, grandchild, grandparent, niece, nephew or uncle.
4 | AUNT | aunt | The player of the role is a sister of the scoping person's mother or father.
5 | MAUNT | maternal aunt | Description:The player of the role is a biological sister of the scoping person's biological mother.
5 | PAUNT | paternal aunt | Description:The player of the role is a biological sister of the scoping person's biological father.
4 | COUSN | cousin | The player of the role is a relative of the scoping person descended from a common ancestor, such as a grandparent, by two or more steps in a diverging line.
5 | MCOUSN | maternal cousin | Description:The player of the role is a biological relative of the scoping person descended from a common ancestor on the player's mother's side, such as a grandparent, by two or more steps in a diverging line.
5 | PCOUSN | paternal cousin | Description:The player of the role is a biological relative of the scoping person descended from a common ancestor on the player's father's side, such as a grandparent, by two or more steps in a diverging line.
4 | GGRPRN | great grandparent | The player of the role is a parent of the scoping person's grandparent.
5 | GGRFTH | great grandfather | The player of the role is the father of the scoping person's grandparent.
6 | MGGRFTH | maternal great-grandfather | Description:The player of the role is the biological father of the scoping person's biological mother's parent.
6 | PGGRFTH | paternal great-grandfather | Description:The player of the role is the biological father of the scoping person's biological father's parent.
5 | GGRMTH | great grandmother | The player of the role is the mother of the scoping person's grandparent.
6 | MGGRMTH | maternal great-grandmother | Description:The player of the role is the biological mother of the scoping person's biological mother's parent.
6 | PGGRMTH | paternal great-grandmother | Description:The player of the role is the biological mother of the scoping person's biological father's parent.
5 | MGGRPRN | maternal great-grandparent | Description:The player of the role is a biological parent of the scoping person's biological mother's parent.
6 | MGGRFTH
6 | MGGRMTH
5 | PGGRPRN | paternal great-grandparent | Description:The player of the role is a biological parent of the scoping person's biological father's parent.

6 | PGGRFTH
6 | PGGRMTH
4 | GRNDCHILD | grandchild | The player of the role is a child of the scoping person's son or daughter.
5 | GRNDDAU | granddaughter | The player of the role is a daughter of the scoping person's son or daughter.
5 | GRNDSON | grandson | The player of the role is a son of the scoping person's son or daughter.
4 | GRPRN | grandparent | The player of the role is a parent of the scoping person's mother or father.
5 | GRFTH | grandfather | The player of the role is the father of the scoping person's mother or father.
6 | MGRFTH | maternal grandfather | Description:The player of the role is the biological father of the scoping person's biological mother.
6 | PGRFTH | paternal grandfather | Description:The player of the role is the biological father of the scoping person's biological father.
5 | GRMTH | grandmother | The player of the role is the mother of the scoping person's mother or father.
6 | MGRMTH | maternal grandmother | Description:The player of the role is the biological mother of the scoping person's biological mother.
6 | PGRMTH | paternal grandmother | Description:The player of the role is the biological mother of the scoping person's biological father.
5 | MGRPRN | maternal grandparent | Description:The player of the role is the biological parent of the scoping person's biological mother.
6 | MGRFTH
6 | MGRMTH
5 | PGRPRN | paternal grandparent | Description:The player of the role is the biological parent of the scoping person's biological father.
6 | PGRFTH
6 | PGRMTH
4 | INLAW | inlaw | A relationship between an individual and a member of their spousal partner's immediate family.
5 | CHLDINLAW | child-in-law | The player of the role is the spouse of scoping person's child.
6 | DAUINLAW | daughter in-law | The player of the role is the wife of scoping person's son.
6 | SONINLAW | son in-law | The player of the role is the husband of scoping person's daughter.
5 | PRNINLAW | parent in-law | The player of the role is the parent of scoping person's husband or wife.
6 | FTHINLAW | father-in-law | The player of the role is the father of the scoping person's husband or wife.
6 | MTHINLAW | mother-in-law | The player of the role is the mother of the scoping person's husband or wife.
5 | SIBINLAW | sibling in-law | The player of the role is: (1) a sibling of the scoping person's spouse, or (2) the spouse of the scoping person's sibling, or (3) the spouse of a sibling of the scoping person's spouse.
6 | BROINLAW | brother-in-law | The player of the role is: (1) a brother of the scoping person's spouse, or (2) the husband of the scoping person's sister, or (3) the husband of a sister of the scoping person's spouse.
6 | SISINLAW | sister-in-law | The player of the role is: (1) a sister of the scoping person's spouse, or (2) the wife of the scoping person's brother, or (3) the wife of a brother of the scoping person's spouse.
4 | NIENEPH | niece/nephew | The player of the role is a child of scoping person's brother or sister or of the brother or sister of the scoping person's spouse.
5 | NEPHEW | nephew | The player of the role is a son of the scoping person's brother or sister or of the brother or sister of the scoping person's spouse.
5 | NIECE | niece | The player of the role is a daughter of the scoping person's brother or sister or of the brother or sister of the scoping person's spouse.
4 | UNCLE | uncle | The player of the role is a brother of the scoping person's mother or father.
5 | MUNCLE | maternal uncle | Description:The player of the role is a biological brother of the scoping person's biological mother.
5 | PUNCLE | paternal uncle | Description:The player of the role is a biological brother of the scoping person's biological father.
3 | PRN | parent | The player of the role is one who begets, gives birth to, or nurtures and raises the scoping entity (child).
4 | ADOPTP | adoptive parent | The player of the role (parent) has taken the scoper (child) into their family through legal means and raises them as his or her own child.
5 | ADOPTF | adoptive father | The player of the role (father) is a male who has taken the scoper (child) into their family through legal means and raises them as his own child.
5 | ADOPTM | adoptive mother | The player of the role (father) is a female who has taken the scoper (child) into their family through legal means and raises them as her own child.
4 | FTH | father | The player of the role is a male who begets or raises or nurtures the scoping entity (child).
5 | ADOPTF
5 | FTHFOST | foster father | The player of the role (parent) who is a male state-certified caregiver responsible for the scoper (child) who has been placed in the parent's care. The placement of the child is usually arranged through the government or a social-service agency, and temporary. The state, via a jurisdiction recognized child protection agency, stands as in loco parentis to the child, making all legal decisions while the foster parent is responsible for the day-to-day care of the specified child.
5 | NFTH | natural father | The player of the role is a male who begets the scoping entity (child).
6 | NFTHF | natural father of fetus | Indicates the biologic male parent of a fetus.
5 | STPFTH | stepfather | The player of the role is the husband of scoping person's mother and not the scoping person's natural father.
4 | MTH | mother | The player of the role is a female who conceives, gives birth to, or raises and nurtures the scoping entity (child).
5 | ADOPTM
5 | GESTM | gestational mother | The player is a female whose womb carries the fetus of the scoper. Generally used when the gestational mother and natural mother are not the same.
5 | MTHFOST | foster mother | The player of the role (parent) who is a female state-certified caregiver responsible for the scoper (child) who has been placed in the parent's care. The placement of the child is usually arranged through the government or a social-service agency, and temporary. The state, via a jurisdiction recognized child protection agency, stands as in loco parentis to the child, making all legal decisions while the foster parent is responsible for the day-to-day care of the specified child.
5 | NMTH | natural mother | The player of the role is a female who conceives or gives birth to the scoping entity (child).
6 | NMTHF | natural mother of fetus | The player is the biologic female parent of the scoping fetus.

5 | STPMTH | stepmother | The player of the role is the wife of scoping person's father and not the scoping person's natural mother.
4 | NPRN | natural parent
5 | NFTH
5 | NMTH
4 | PRNFOST | foster parent | The player of the role (parent) who is a state-certified caregiver responsible for the scoper (child) who has been placed in the parent's care. The placement of the child is usually arranged through the government or a social-service agency, and temporary. The state, via a jurisdiction recognized child protection agency, stands as in loco parentis to the child, making all legal decisions while the foster parent is responsible for the day-to-day care of the specified child.
5 | FTHFOST
5 | MTHFOST
4 | STPPRN | step parent | The player of the role is the spouse of the scoping person's parent and not the scoping person's natural parent.
5 | STPFTH
5 | STPMTH
3 | SIB | sibling | The player of the role shares one or both parents in common with the scoping entity.
4 | BRO | brother | The player of the role is a male sharing one or both parents in common with the scoping entity.
5 | HBRO | half-brother | The player of the role is a male related to the scoping entity by sharing only one biological parent.
5 | NBRO | natural brother | The player of the role is a male having the same biological parents as the scoping entity.
6 | TWINBRO | twin brother | The scoper was carried in the same womb as the male player and shares common biological parents.
7 | FTWINBRO | fraternal twin brother | The scoper was carried in the same womb as the male player and shares common biological parents but is the product of a distinct egg/sperm pair.
7 | ITWINBRO | identical twin brother | The male scoper is an offspring of the same egg-sperm pair as the male player.
5 | STPBRO | stepbrother | The player of the role is a son of the scoping person's stepparent.
4 | HSIB | half-sibling | The player of the role is related to the scoping entity by sharing only one biological parent.
5 | HBRO
5 | HSIS | half-sister | The player of the role is a female related to the scoping entity by sharing only one biological parent.
4 | NSIB | natural sibling | The player of the role has both biological parents in common with the scoping entity.
5 | NBRO
5 | NSIS | natural sister | The player of the role is a female having the same biological parents as the scoping entity.
6 | TWINSIS | twin sister | The scoper was carried in the same womb as the female player and shares common biological parents.
7 | FTWINSIS | fraternal twin sister | The scoper was carried in the same womb as the female player and shares common biological parents but is the product of a distinct egg/sperm pair.
7 | ITWINSIS | identical twin sister | The female scoper is an offspring of the same egg-sperm pair as the female player.
5 | TWIN | twin | The scoper and player were carried in the same womb and shared common biological parents.
6 | TWINBRO
6 | TWINSIS
6 | FTWIN | fraternal twin | The scoper and player were carried in the same womb and share common biological parents but are the product of distinct egg/sperm pairs.
7 | FTWINBRO
7 | FTWINSIS
6 | ITWIN | identical twin | The scoper and player are offspring of the same egg-sperm pair.
7 | ITWINBRO
7 | ITWINSIS
4 | SIS | sister | The player of the role is a female sharing one or both parents in common with the scoping entity.
5 | HSIS
5 | NSIS
5 | STPSIS | stepsister | The player of the role is a daughter of the scoping person's stepparent.
4 | STPSIB | step sibling | The player of the role is a child of the scoping person's stepparent.
5 | STPBRO
5 | STPSIS
3 | SIGOTHR | significant other | A person who is important to one's well being; especially a spouse or one in a similar relationship. (The player is the one who is important)
4 | DOMPART | domestic partner | The player of the role cohabits with the scoping person but is not the scoping person's spouse.
4 | FMRSPS | former spouse | Player of the role was previously joined to the scoping person in marriage and this marriage is now dissolved and inactive. Usage Note: This is significant to indicate as some jurisdictions have different legal requirements for former spouse to access the patient's record, from a general friend.
4 | SPS | spouse | The player of the role is a marriage partner of the scoping person.
5 | HUSB | husband | The player of the role is a man joined to a woman (scoping person) in marriage.
5 | WIFE | wife | The player of the role is a woman joined to a man (scoping person) in marriage.
2 | FRND | unrelated friend | The player of the role is a person who is known, liked, and trusted by the scoping person.
2 | NBOR | neighbor | The player of the role lives near or next to the scoping person.
2 | ONESELF | self | The relationship that a person has with his or her self.
2 | ROOM | Roommate | One who shares living quarters with the subject.
1 | (_PolicyOrProgramCoverageRoleType) Abstract | | Description: A role recognized through the eligibility of an identified party for benefits covered under an insurance policy or a program based on meeting eligibility criteria. Eligibility as a covered party may be conditioned on the party meeting criteria to qualify for coverage under a policy or program, which may be mandated by law. These criteria may be: The sole basis for coverage, e.g., being differently abled may qualify a person for disability coverage May more fully qualify a covered party role e.g, being differently abled may qualify an adult child as a dependent May impact the level of coverage for a covered party under a policy or program, e.g., being differently abled may qualify a program eligible for additional benefits. Discussion: The Abstract Value Set "CoverageRoleType", which was developed for use in the Canadian realm "pre-coordinate" coverage roles with other roles that a covered party must play in order to be eligible for coverage, e.g., "handicapped dependent". These role.codes may only be used with COVPTY to avoid overlapping concepts that would result from using them to specify the specializations of COVPTY, e.g., the role.class DEPEN should not be used with the role.code family

dependent because that relationship has overlapping concepts due to the role.code precoodination and is conveyed in FICO with the personal relationship role that has a PART role link to the covered party role. For the same reasons, the role.class DEPEN should not be used with the role.code HANDIC (handicapped dependent); the role.code DIFFABLE (differently abled) should be used instead. In summary, the coded concepts in the Abstract Value Set "CoveredPartyRoleType" can be "post-coordinated" with the "RoleClassCoveredParty" Abstract Value Set. Decoupling these concepts is intended to support an expansive range of covered party concepts and their semantic comparability.

2 | (_CoverageRoleType) Abstract | | Role recognized through the issuance of insurance coverage to an identified covered party who has this relationship with the policy holder such as the policy holder themselves (self), spouse, child, etc

3 | FAMDEP | family dependent | The player of the role is dependent of the scoping entity.

3 | HANDIC | handicapped dependent | Covered party is a dependent of the policy holder with a physical or mental disability causing a disadvantage that makes independent achievement unusually difficult.

3 | INJ | injured plaintiff | Covered party is an injured party with a legal claim for compensation against a policy holder under an insurance policy.

3 | SELF | self | Covered party is the policy holder. Also known as the subscriber.

3 | SPON | sponsored dependent | Covered party is an individual that the policy holder has assumed responsibility for, such as foster child or legal ward.

3 | STUD | student | Covered party to an insurance policy has coverage through full-time or part-time attendance at a recognized educational institution as defined by a particular insurance policy.

4 | FSTUD | full-time student | Covered party to an insurance policy has coverage through full-time attendance at a recognized educational institution as defined by a particular insurance policy.

4 | PSTUD | part-time student | Covered party to an insurance policy has coverage through part-time attendance at a recognized educational institution as defined by a particular insurance policy.

3 | ADOPT | adopted child | A child taken into one's family through legal means and raised as one's own child.

3 | GCHILD | grandchild | A child of one's son or daughter.

3 | GPARNT | grandparent | parent of a parent of the subject.

3 | NAT | natural child | A child as determined by birth.

3 | NIENE | niece/nephew | A child of one's brother or sister or of the brother or sister of one's spouse.

3 | PARNT | parent | One that begets or brings forth offspring or a person who brings up and cares for for another (Webster's Collegiate Dictionary)

3 | SPSE | spouse | A marriage partner; a husband or wife.

3 | STEP | step child | A child receiving parental care and nurture from a person who is related to them through marriage to their parent.

2 | (_CoveredPartyRoleType) Abstract | | A role recognized through the eligibility of an identified living subject for benefits covered under an insurance policy or a program. Eligibility as a covered party may be conditioned on a relationship with (1) the policy holder such as the policy holder who is covered as an individual under a poliy or as a party sponsored for coverage by the policy holder. Example:An employee as a subscriber; or (2) on being scoped another covered party such as the subscriber, as in the case of a dependent. Discussion: The Abstract Value Set "CoverageRoleType", which was developed for use in the Canadian realm "pre-coordinate" coverage roles with other roles that a covered party must play in order to be eligible for coverage, e.g., "handicapped dependent". Other codes in the Abstract Value Set CoveredPartyRoleType domain can be "post-coordinated" with the EligiblePartyRoleType codes to denote comparable concepts. Decoupling the concepts is intended to support a wider range of concepts and semantic comparability of coded concepts.

3 | (_ClaimantCoveredPartyRoleType) Abstract | | DescriptionA role recognized through the eligibility of a party play a claimant for benefits covered or provided under an insurance policy.

4 | CRIMEVIC | crime victim | Description: A person playing the role of program eligible under a program based on allegations of being the victim of a crime. Discussion: This CoveredPartyRoleType.code is typically used when the CoveredPartyRole class code is "program eligible" and the person's status as a crime victim meets jurisdictional or program criteria.

4 | INJWKR | injured worker | Description: A person playing the role of program eligible under a workers compensation program based on the filing of work-related injury claim. Discussion: This CoveredPartyRoleType.code is used when the CoveredPartyRole class code is either "program eligible", a "named insured", and "individual insured", or "dependent", and the person's status as differently abled or "handicapped" meets jurisdictional, policy, or program criteria.

3 | (_DependentCoveredPartyRoleType) Abstract | | Description: A role recognized through the eligibility of a party to play a dependent for benefits covered or provided under a health insurance policy because of an association with the subscriber that is recognized by the policy underwriter.

4 | COCBEN | continuity of coverage beneficiary | Description: A person playing the role of an individual insured with continuity of coverage under a policy which is being terminated based on loss of original status that was the basis for coverage. Criteria for qualifying for continuity of coverage may be set by law. Discussion: This CoveredPartyRoleType.code is typically used when the CoveredPartyRole class code is either "program eligible" or "subscriber" and the person's status as a continuity of coverage beneficiary meets jurisdictional or policy criteria.

4 | DIFFABL | differently abled | Description: A person playing the role of program eligible under a program based on meeting criteria for health or functional limitation set by law or by the program. Discussion: This CoveredPartyRoleType.code is typically used when the CoveredPartyRole class code is either "program eligible", "named insured", "individual insured", or "dependent", and the person's status as differently abled meets jurisdictional, policy, or program criteria.

4 | WARD | ward | Description: A person, who is a minor or is deemed incompetent, who plays the role of a program eligible where eligibility for coverage is based on meeting program eligibility criteria for status as a ward of a court or jurisdiction. Discussion: This CoveredPartyRoleType.code is typically used when the CoveredPartyRole class code is a "claimant", "program eligible", a "named insured", an "individual Insured" or a "dependent", and the person's status as a ward meets program or policy criteria. In the case of a ward covered under a program providing financial or health benefits, a governmental agency may take temporary custody of a minor or incompetent for his/her protection and care, e.g., if the ward is suffering from neglect or abuse, or has been in trouble with the law.

3 | (_IndividualInsuredPartyRoleType) Abstract | | A role recognized through the eligibility of a party to play an individual insured for benefits covered or provided under an insurance policy where the party is also the policy holder.

4 | COCBEN

4 | RETIREE | retiree | Description: A person playing the role of an individual insured under a policy based on meeting criteria for the employment status of retired set by law or the policy. Discussion: This CoveredPartyRoleType.code is typically used when the CoveredPartyRole class code is either "program eligible" or "subscriber" and the person's status as a retiree meets jurisdictional or policy criteria.

3 | (_ProgramEligiblePartyRoleType) Abstract | | Description:A role recognized through the eligibility of a party to play a program eligible for benefits covered or provided under a program.
4 | CRIMEVIC
4 | INJWKR
4 | DIFFABL
4 | WARD
4 | INDIG | member of an indigenous people | Description: A person playing the role of program eligible under a program based on aboriginal ancestry or as a member of an aboriginal community. Discussion: This CoveredPartyRoleType.code is typically used when the CoveredPartyRole class code is "program eligible" and the person's status as a member of an indigenous people meets jurisdictional or program criteria.
4 | MIL | military | Definition: A person playing the role of program eligible under a program based on military status. Discussion: This CoveredPartyRoleType.code is typically used when the CoveredPartyRole class code is either "program eligible" or "subscriber" and the person's status as a member of the military meets jurisdictional or program criteria
5 | ACTMIL | active duty military | Description: A person playing the role of program eligible under a program based on active military status. Discussion: This CoveredPartyRoleType.code is typically used when the CoveredPartyRole class code is either "program eligible" or "subscriber" and the persons status as active duty military meets jurisdictional or program criteria.
5 | RETMIL | retired military | Description: A person playing the role of program eligible under a program based on retired military status. Discussion: This CoveredPartyRoleType.code is typically used when the CoveredPartyRole class code is either "program eligible" or "subscriber" and the persons status as retired military meets jurisdictional or program criteria.
5 | VET | veteran | Description: A person playing the role of program eligible under a program based on status as a military veteran. Discussion: This CoveredPartyRoleType.code is typically used when the CoveredPartyRole class code is either "program eligible" or "subscriber" and the persons status as a veteran meets jurisdictional or program criteria.
3 | (_SubscriberCoveredPartyRoleType) Abstract | | Description: A role recognized through the eligibility of a party to play a subscriber for benefits covered or provided under a health insurance policy.
4 | COCBEN
4 | RETIREE
4 | MIL
1 | (_ResearchSubjectRoleBasis) Abstract | | Specifies the administrative functionality within a formal experimental design for which the ResearchSubject role was established. Examples: screening - role is used for pre-enrollment evaluation portion of the design; enrolled - role is used for subjects admitted to the active treatment portion of the design.
2 | ERL | enrollment | Definition:The specific role being played by a research subject participating in the active treatment or primary data collection portion of a research study.
2 | SCN | screening | Definition:The specific role being played by a research subject participating in the pre-enrollment evaluation portion of a research study.
1 | (_ServiceDeliveryLocationRoleType) Abstract | | A role of a place that further classifies the setting (e.g., accident site, road side, work site, community location) in which services are delivered.
2 | (_DedicatedServiceDeliveryLocationRoleType) Abstract | | A role of a place that further classifies a setting that is intended to house the provision of services.
3 | (_DedicatedClinicalLocationRoleType) Abstract | | A role of a place that further classifies the clinical setting (e.g., cardiology clinic, primary care clinic, rehabilitation hospital, skilled nursing facility) in which care is delivered during an encounter.
4 | DX | Diagnostics or therapeutics unit | A practice setting where diagnostic procedures or therapeutic interventions are performed
5 | CVDX | Cardiovascular diagnostics or therapeutics unit | A practice setting where cardiovascular diagnostic procedures or therapeutic interventions are performed (e.g., cardiac catheterization lab, echocardiography suite)
6 | CATH | Cardiac catheterization lab
6 | ECHO | Echocardiography lab
5 | GIDX | Gastroenterology diagnostics or therapeutics lab | A practice setting where GI procedures (such as endoscopies) are performed
6 | ENDOS | Endoscopy lab | (X12N 261QD0000N)
5 | RADDX | Radiology diagnostics or therapeutics unit | A practice setting where radiology services (diagnostic or therapeutic) are provided (X12N 261QR0200N)
6 | RADO | Radiation oncology unit | (X12N 261QX0203N)
6 | RNEU | Neuroradiology unit
4 | HOSP | Hospital | An acute care institution that provides medical, surgical, or psychiatric care and treatment for the sick or the injured.
5 | CHR | Chronic Care Facility | (1) A hospital including a physical plant and personnel that provides multidisciplinary diagnosis and treatment for diseases that have one or more of the following characteristics: is permanent; leaves residual disability; is caused by nonreversible pathological alteration; requires special training of the patient for rehabilitation; and/or may be expected to require a long period of supervision or care. In addition, patients require the safety, security, and shelter of these specialized inpatient or partial hospitalization settings. (2) A hospital that provides medical and skilled nursing services to patients with long-term illnesses who are not in an acute phase but who require an intensity of services not available in nursing homes
5 | GACH | Hospitals; General Acute Care Hospital | (X12N 282N00000N)
5 | MHSP | Military Hospital | A health care facility operated by the Department of Defense or other military operation.
5 | PSYCHF | Psychatric Care Facility | Healthcare facility that cares for patients with psychiatric illness(s).
5 | RH | Rehabilitation hospital | (X12N 283X00000N)
6 | RHAT | addiction treatment center | Description: A location that plays the role of delivering services which may include life training and/or social support to people with addictions.
6 | RHII | intellectual impairment center | Description: A location that plays the role of delivering services which may include adaptation, rehabilitation and social integration services for people with intellectual and/or pervasive development disorders such as autism or severe behaviour disorder.
6 | RHMAD | parents with adjustment difficulties center | Description: A location that plays the role of delivering services which may social support services for adolescents who are pregnant or have child and are experiencing adaptation issues/difficulties in their current or eventual parenting role.
6 | RHPI | physical impairment center | Description: A location that plays the role of delivering services which may include adaptation, rehabilitation and social integration services for people with physical impairments.

7 | RHPIH | physical impairment - hearing center | Description: A location that plays the role of delivering services for people with hearing impairments.
7 | RHPIMS | physical impairment - motor skills center | Description: A location that plays the role of delivering services for people with motor skill impairments.
7 | RHPIVS | physical impairment - visual skills center | Description: A location that plays the role of delivering services for people with visual skill impairments.
6 | RHYAD | youths with adjustment difficulties center | Description: A location that plays the role of delivering services which may include life training and/or social support services for the adaption, rehabilitation and social integration of youths with adjustment difficulties.
4 | HU | Hospital unit
5 | BMTU | Bone marrow transplant unit
5 | CCU | Coronary care unit
5 | CHEST | Chest unit | A specialty unit in hospital that focuses on chronic respirator patients and pulmonary failure
5 | EPIL | Epilepsy unit
5 | ER | Emergency room | The section of a health care facility for providing rapid treatment to victims of sudden illness or trauma.
6 | ETU | Emergency trauma unit
5 | HD | Hemodialysis unit
5 | HLAB | hospital laboratory | Description: A location that plays the role of delivering services which may include tests done based on clinical specimens to get health information about a patient as pertaining to the diagnosis, treatment and prevention of disease. Hospital laboratories may be further divided into specialized units such as Anatomic Pathology, Microbiology, and Biochemistry.
6 | INLAB | inpatient laboratory | Description: A location that plays the role of delivering services which may include tests are done on clinical specimens to get health information about a patient pertaining to the diagnosis, treatment, and prevention of disease for a hospital visit longer than one day.
6 | OUTLAB | outpatient laboratory | Description: A location that plays the role of delivering services which may include tests are done on clinical specimens to get health information about a patient pertaining to the diagnosis, treatment, and prevention of disease for same day visits.
5 | HRAD | radiology unit | Description: A location that plays the role of delivering services which may include the branch of medicine that uses ionizing and non-ionizing radiation to diagnose and treat diseases. The radiology unit may be further divided into subspecialties such as Imaging, Cardiovascular, Thoracic, and Ultrasound.
5 | HUSCS | specimen collection site | Description: A location that plays the role of delivering services which may include collecting specimens and/or samples from patients for laboratory testing purposes, but does not perform any tests or analysis functions.
5 | ICU | Intensive care unit
6 | PEDICU | Pediatric intensive care unit
7 | PEDNICU | Pediatric neonatal intensive care unit
5 | INPHARM | inpatient pharmacy | Description: A location that plays the role of delivering services which may include providing judicious, safe, efficacious, appropriate and cost effective use of medicines for treatment of patients for visits longer than one day. The distinction between inpatient pharmacies and retail (or outpatient) pharmacies is that they are part of a patient's continuity of care while staying in the hospital.
5 | MBL | medical laboratory | Description: A location that plays the role of delivering services which include biochemistry, hematology, microbiology, immunochemistry, and toxicology.
5 | NCCS | Neurology critical care and stroke unit
5 | NS | Neurosurgery unit
5 | OUTPHARM | outpatient pharmacy | Description: A location that plays the role of delivering services which may include providing judicious, safe, efficacious, appropriate and cost effective use of medicines for treatment of patients for outpatient visits and may also be used for discharge prescriptions.
5 | PEDU | Pediatric unit
6 | PEDICU
5 | PHU | Psychiatric hospital unit | (X12N 273R00000N)
5 | RHU | Rehabilitation hospital unit
5 | SLEEP | Sleep disorders unit | (X12N 261QA1200N)
4 | NCCF | Nursing or custodial care facility
5 | SNF | Skilled nursing facility | (X12N 314000000N)
4 | OF | Outpatient facility
5 | ALL | Allergy clinic
5 | AMPUT | Amputee clinic
5 | BMTC | Bone marrow transplant clinic
5 | BREAST | Breast clinic
5 | CANC | Child and adolescent neurology clinic
5 | CAPC | Child and adolescent psychiatry clinic
5 | CARD | Ambulatory Health Care Facilities; Clinic/Center; Rehabilitation: Cardiac Facilities
6 | PEDCARD | Pediatric cardiology clinic
5 | COAG | Coagulation clinic
5 | CRS | Colon and rectal surgery clinic
5 | DERM | Dermatology clinic
5 | ENDO | Endocrinology clinic
6 | PEDE | Pediatric endocrinology clinic
5 | ENT | Otorhinolaryngology clinic
5 | FMC | Family medicine clinic
5 | GI | Gastroenterology clinic
6 | PEDGI | Pediatric gastroenterology clinic
5 | GIM | General internal medicine clinic
5 | GYN | Gynecology clinic
5 | HEM | Hematology clinic

6 | PEDHEM | Pediatric hematology clinic
5 | HTN | Hypertension clinic
5 | IEC | Impairment evaluation center | Focuses on assessing disability
5 | INFD | Infectious disease clinic
6 | PEDID | Pediatric infectious disease clinic
5 | INV | Infertility clinic
5 | LYMPH | Lympedema clinic
5 | MGEN | Medical genetics clinic
5 | NEPH | Nephrology clinic
6 | PEDNEPH | Pediatric nephrology clinic
5 | NEUR | Neurology clinic
5 | OB | Obstetrics clinic
5 | OMS | Oral and maxillofacial surgery clinic
5 | ONCL | Medical oncology clinic
6 | PEDHO | Pediatric oncology clinic
5 | OPH | Opthalmology clinic
5 | OPTC | optometry clinic | Description: A location that plays the role of delivering services which may include examination, diagnosis, treatment, management, and prevention of diseases and disorders of the eye as well as prescribing and fitting appropriate corrective lenses (glasses or contact lenses) as needed. Optometry clinics may also provide tests for visual field screening, measuring intra-ocular pressure and ophthalmoscopy, as and when required.
5 | ORTHO | Orthopedics clinic
6 | HAND | Hand clinic
5 | PAINCL | Pain clinic | (X12N 261QP3300N)
5 | PC | Primary care clinic | (X12N 261QP2300N)
5 | PEDC | Pediatrics clinic
6 | PEDCARD
6 | PEDE
6 | PEDGI
6 | PEDHEM
6 | PEDID
6 | PEDNEPH
6 | PEDHO
6 | PEDRHEUM | Pediatric rheumatology clinic
5 | POD | Podiatry clinic | (X12N 261QP1100N)
5 | PREV | Preventive medicine clinic
5 | PROCTO | Proctology clinic
5 | PROFF | Provider's Office | Location where healthcare service was delivered, identified as the healthcare provider's practice office.
5 | PROS | Prosthodontics clinic
5 | PSI | Psychology clinic
5 | PSY | Psychiatry clinic
5 | RHEUM | Rheumatology clinic
6 | PEDRHEUM
5 | SPMED | Sports medicine clinic
5 | SU | Surgery clinic
6 | PLS | Plastic surgery clinic
6 | URO | Urology clinic
5 | TR | Transplant clinic
5 | TRAVEL | Travel and geographic medicine clinic
5 | WND | Wound clinic
4 | RTF | Residential treatment facility
5 | PRC | Pain rehabilitation center
5 | SURF | Substance use rehabilitation facility | (X12N 324500000N)
3 | (_DedicatedNonClinicalLocationRoleType) Abstract | | A role of a place that further classifies a setting that is intended to house the provision of non-clinical services.
4 | DADDR | Delivery Address | Location address where medical supplies were transported to for use.
4 | MOBL | Mobile Unit | Location (mobile) where healthcare service was delivered.
5 | AMB | Ambulance | Location (mobile) where healthcare service was delivered, identified specifically as an ambulance.
4 | PHARM | Pharmacy | Location where healthcare service was delivered, identified as a pharmacy.
5 | INPHARM
5 | OUTPHARM
2 | (_IncidentalServiceDeliveryLocationRoleType) Abstract
3 | ACC | accident site | Location of an accident where healthcare service was delivered, such as a roadside.
3 | COMM | Community Location | Community location where healthcare is delivered.
4 | CSC | community service center | Description: A location that plays the role of delivering services which may include providing front-line services to the population of a defined geographic area such as: healthcare services and social services, preventive or curative, rehabilitation or reintegration.
3 | PTRES | Patient's Residence | location where healthcare was delivered which is the residence of the Patient.
3 | SCHOOL | school | Location where healthcare service was delivered, identified as a school or educational facility.
3 | UPC | underage protection center | Description: A location that plays the role of delivering services which may include: social emergency services required for a young person as required under any jurisdictional youth laws, child placement, and family mediation in the defined geographical area the SDL is responsible for. It may provide expertise in a judiciary setting on child custody, adoption and biological history research.

3 | WORK | work site | Location where healthcare service was delivered, identified as a work place.
1 | (_SpecimenRoleType) Abstract
2 | C | Calibrator | A specimen used for initial calibration settings of an instrument
2 | G | Group | A set of patient samples in which the individuals of the group may or may not be identified.
2 | L | Pool | Aliquots of individual specimens combined to form a single specimen representing all of the included individuals.
2 | P | Patient | A specimen that has been collected from a patient.
2 | Q | Quality Control | A specimen specifically used to verify the sensitivity, specificity, accuracy or other perfomance parameter of a diagnostic test.
3 | B | Blind | Quality Control specimen submitted to the lab whose identity and composition is not known to the lab.
3 | E | Electronic QC | An electronically simulated QC specimen
3 | F | Filler Proficiency | Specimen used for testing proficiency of an organization performing testing (how does this differ from O?)
3 | O | Operator Proficiency | A specimen used for evaluation of operator proficiency (operator in what context?)
3 | V | Verifying | A specimen used for periodic calibration checks of instruments
2 | R | Replicate | A portion of an original patent sample that is tested at the same time as the original sample
1 | CLAIM | claimant | A party that makes a claim for coverage under a policy.
1 | communityLaboratory | Community Laboratory
1 | GT | Guarantor | An individual or organization that makes or gives a promise, assurance, pledge to pay or has paid the healthcare service provider.
1 | homeHealth | Home Health
1 | laboratory | Laboratory
1 | pathologist | Pathologist
1 | PH | Policy Holder | Policy holder for the insurance policy.
1 | phlebotomist | Phlebotomist
1 | PROG | program eligible | A party that meets the eligibility criteria for coverage under a program.
1 | PT | Patient | The recipient for the service(s) and/or product(s) when they are not the covered party.
1 | subject | Self
1 | thirdParty | Third Party
1 | DEP
1 | DEPEN | dependent | A party covered under a policy based on association with a subscriber.
1 | FM | Family Member | A member of the covered party's family. This could be the spouse, a parent, a grand parent, a sibling, etc.
1 | INDIV | individual | A party covered under a policy as the policyholder.
1 | NAMED | named insured | A party to an insurance policy to whom the insurer agrees to indemnify for losses, provides benefits for, or renders services.
1 | PSYCHCF
1 | SUBSCR | subscriber | A party covered under a policy based on association with a sponsor who is the policy holder, and whose association may provide for the eligibility of dependents for coverage