

Week 18 Report

Session ID: 1 | Name: demo | Phase: EarlySprouting
Started: 2025-09-24 21:01:45.999471

Summary (plant-level)

Normal plants: 10
Abnormal plants: 3
Empty bags: 0
Defect rate (Abnormal/(Normal+Abnormal)): 23.08%


- Det #7: Seven ginger plants were detected. The single abnormal plant amidst six normal ones during rhizome maturation, under overcast conditions (25.1°C, 87% humidity, light 6 km/h wind, and no recent rain), suggests a local...
- Det #8: Six ginger plants were detected in this snapshot. The 2:4 ratio of abnormal to normal plants during Rhizome Maturation under overcast conditions (25.1°C, 87% humidity) suggests a

Detections

Det #7 — Abnormal — RhizomeMaturation — RhizomeMaturation

Time: 2025-09-24 21:04
Weather: Overcast Clouds | 25.1°C | 87% | 0.0 mm (1h) | 6 km/h

Seven ginger plants were detected. The single abnormal plant amidst six normal ones during rhizome maturation, under overcast conditions (25.1°C, 87% humidity, light 6 km/h wind, and no recent rain), suggests a localized issue rather than a widespread problem affecting the whole ginger patch. The favorable temperature and humidity generally support good growth, but the abnormal plant indicates a potential nutrient deficiency, pest infestation, or localized disease.




This anomaly warrants close inspection. The cause should be determined quickly to prevent its spread. The absence of recent rain might point to localized water stress if the plant isn't properly irrigated.

- Investigate: Examine the abnormal plant closely for signs of disease, pests, or nutrient deficiencies.
- Address Water: Ensure consistent moisture around the abnormal plant without overwatering.
- Monitor: Continue daily monitoring for changes in the abnormal plant and any spread of the problem.

Det #8 — Abnormal — RhizomeMaturation — RhizomeMaturation

Time: 2025-09-24 21:04
Weather: Overcast Clouds | 25.1°C | 87% | 0.0 mm (1h) | 6 km/h

Six ginger plants were detected in this snapshot. The 2:4 ratio of abnormal to normal plants during Rhizome Maturation under overcast conditions (25.1°C, 87% humidity) suggests a potential issue impacting rhizome development. While the lack of recent rain and moderate wind are positive, the high humidity may be contributing to fungal diseases or hindering proper rhizome growth, especially considering the high percentage of abnormal plants.



This imbalance warrants immediate attention to prevent further yield loss. The relatively warm temperature is beneficial, but the high humidity needs managing.

- Improve aeration: Increase spacing between plants or consider pruning lower leaves to improve airflow and reduce humidity around rhizomes.
- Monitor for disease: Closely inspect abnormal plants for signs of fungal infections and take appropriate action (e.g., targeted fungicide application if necessary).
- Assess soil drainage: Ensure the soil drains well to prevent waterlogging which can also contribute to poor rhizome development.