

Evaluation of Zernike Basis Functions

J. Schilling

September 28, 2021

The Zernike basis polynomials $R_{m+2j}^m(r)$ can be defined using a recurrence relation as follows for $j = 0, 1, 2, \dots$. For given n and m , j has to go up to a maximum value of $j_{\max} = (n - m)/2$. This requires that the difference $(n - m)$ is positive and even. The following recurrence relation is used:

$$R_{n+2}^m(r) = \frac{n+2}{(n+2)^2 - m^2} \left\{ \left[4(n+1)r^2 - \frac{(n+m)^2}{n} - \frac{(n-m+2)^2}{n+2} \right] R_n^m - \frac{n^2 - m^2}{n} R_{n-2}^m \right\}. \quad (1)$$

An index shift by 2 in n is performed:

$$R_n^m(r) = \frac{n}{n^2 - m^2} \left\{ \left[4(n-1)r^2 - \frac{(n-2+m)^2}{n-2} - \frac{(n-m)^2}{n} \right] \underbrace{R_{n-2}^m}_{=R_{j-1}^m} - \frac{(n-2)^2 - m^2}{n-2} \underbrace{R_{n-4}^m}_{=R_{j-2}^m} \right\}. \quad (2)$$

The first two elements are computed separately:

$$R_m^m(r) = r^m \quad : j = 0 \quad (3)$$

$$R_{m+2}^m(r) = (m+2)r^{m+2} - (m+1)r^m \quad : j = 1. \quad (4)$$

This method is described in Ref. [1].

A demo implementation of this “method” is provided in the file `Zernike.py` along with this document. In Fig. 1 a demo output from this method is shown.




Figure 1: Evaluated Zernike basis function using the method presented in this document.

References

- [1] E. C. Kintner, “On the Mathematical Properties of the Zernike Polynomials”, *Optica Acta: International Journal of Optics* **23**, 679–680 (1976).