
ASSIGNMENT

PROGRAMMING USING C

WEEK – 06

NAME : K.Venkateshwari
ROLL NO : 2116240801373
DEPT & SEC :ECE F

Week-6

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that $A[i] - A[j] = k$, $i \neq j$.

```
#include<stdio.h>
int main()
{
    int t,k,i,j,n,f;
    scanf("%d",&t);
    while(t--)
    {
        scanf("%d",&n);
        int A[n];
        for(i=0;i<n;i++)
        {
            scanf("%d",&a[i]);
        }
        scanf("%d",&k);
        f=0;
        for(i=0;i<n-1;i++)
        {
            for(i=0;i<n-1;)
            {
            }
        }
    }
}
```

Syntax Error(s)

```
__tester__.c: In function 'main':
__tester__.c:12:25: error: 'a' undeclared (first use in this function)
   12 |         scanf("%d",&a[i]);
      |                        ^
__tester__.c:12:25: note: each undeclared identifier is reported only once for each function it appears in
__tester__.c:19:9: error: expected expression before '}' token
   19 |     }
      |     ^
__tester__.c:9:13: error: unused variable 'A' [-Werror=unused-variable]
    9 |         int A[n];
      |         ^
__tester__.c:4:19: error: variable 'f' set but not used [-Werror=unused-but-set-variable]
    4 |         int t,k,i,j,n,f;
      |         ^
__tester__.c:4:15: error: unused variable 'j' [-Werror=unused-variable]
    4 |         int t,k,i,j,n,f;
      |         ^
cc1: all warnings being treated as errors
```

Sam loves chocolates and starts buying them on the 1st day of the year. Each day of the year, x , is numbered from 1 to Y . On days when x is odd, Sam will buy x chocolates; on days when x is even, Sam will not purchase any chocolates.

1	
---	--

Complete the code in the editor below. The program must return an array of m positive integers, one for each $\text{maxes}[i]$ representing the total number of elements $\text{nums}[j]$ satisfying $\text{nums}[j] \leq \text{maxes}[i]$ where $0 \leq j < n$ and $0 \leq i < m$, in the given order.

1	
---	--