

ELITE MATHEMATICS

Pearson Edexcel International A Level

Pure 2

Adaptive Classified

Complete Book

With Answers

Questions with full worked solutions

- 12 chapters · 474 pages
- Past papers up to January 2026

Contents

Complete Book — click a chapter to open it

01	Proof	p. 4
02	Polynomials	p. 39
03	Circles	p. 77
04	Binomial Expansion	p. 124
05	Arithmetic Sequences	p. 168
06	Geometric Sequences	p. 185
07	Sequences and Series	p. 206
08	Modelling with Sequences and Series	p. 231
09	Laws of Logarithms	p. 259
10	Trigonometric Equations	p. 307
11	Applications of Differentiation	p. 357
12	Integration	p. 410

PURE 2 | TOPIC 01

Proof

Your classified route

15

QUESTIONS

75

MARKS

4

FAMILIES

5

VARIANTS

DR ESLAM AHMED

CLASSIFIED BY IDEA

Move from recognition to method, then use the working space to complete the solution.

TOPIC 01 | FAMILY 01

Exhaustion/Cases

WHAT THIS FAMILY ASKS

Recognise the exhaustion/cases demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

V02 For modulo cases, use $3k, 3k+1, 3k+2$ or $4k, 4k+1, 4k+2, 4k+3$.

8 classified questions

36 total marks

T01

01_CASE

01_CASE_CORE

CHAPTER
ProofFAMILY
Exhaustion/casesVARIANT
core family move

5 marks

WMA12/01 JANUARY 2022

Question 10

Past-paper question

WMA12-01_2022_Jan_Q10

10. (i) Prove by counter example that the statement

“if p is a prime number then $2p + 1$ is also a prime number”

is not true.

(1)

(ii) Use proof by exhaustion to prove that if n is an integer then

$$5n^2 + n + 12$$

is always even.

(4)

(Total 5 marks)

T01 Proof

01_CASE Exhaustion/cases

01_CASE_CORE core family
move

WORKED SOLUTION

Question 10

WMA12/01 January 2022

WMA12-01_2022_Jan_Q10

1

Give a counterexample

Take $p = 7$, which is prime.

2

Show the statement fails

$2p + 1 = 2(7) + 1 = 15$, and 15 is not prime. So the statement is false.

3

Case one: n even

If n is even, let $n = 2k$. Then $5n^2 + n + 12 = 5(2k)^2 + 2k + 12 = 20k^2 + 2k + 12 = 2(10k^2 + k + 6)$, which is even.

4

Case two: n odd

If n is odd, let $n = 2k + 1$. Then $5n^2 + n + 12 = 5(2k + 1)^2 + (2k + 1) + 12$.

5

Simplify the odd case

$5(2k + 1)^2 + (2k + 1) + 12 = 20k^2 + 22k + 18 = 2(10k^2 + 11k + 9)$, which is even.

6

Conclude by exhaustion

Every integer is either even or odd, and both cases give an even result. Therefore $5n^2 + n + 12$ is always even.

FINAL ANSWER

WMA12-01_2022_Jan_Q10

(i) For $p = 7$, $2p + 1 = 15$, not prime. (ii) Proven.

T01

01_CASE

01_CASE_CORE

CHAPTER
Proof

FAMILY
Exhaustion/cases

VARIANT
core family move
3 marks

WMA12/01 OCTOBER 2022

Question 1

Past-paper question

WMA12-01_2022_Oct_Q01

1. Given that a , b and c are integers greater than 0 such that

- $c = b + 2$
- $a + b + c = 10$

Prove, by exhaustion, that the product of a , b and c is always even.

You may use the table below to illustrate your answer.

(3)

You may not need to use all rows of this table.

a	b	c
	1	
	2	

T01 Proof

01_CASE Exhaustion/cases

01_CASE_CORE core family
move

WORKED SOLUTION

Question 1

WMA12/01 October 2022

WMA12-01_2022_Oct_Q01

1

Use the two conditions

Since $c = b + 2$ and $a + b + c = 10$, substitute for c : $a + b + (b + 2) = 10$.

2

Write a in terms of b

This gives $a + 2b + 2 = 10$, so $a = 8 - 2b$.

3

List the possible b values

The integers a, b, c are all positive. Since $b > 0$ and $a = 8 - 2b > 0$, the only possible values are $b = 1, 2, 3$.

4

Exhaust the cases

If $b = 1$, then $a = 6$ and $c = 3$, so $abc = 18$. If $b = 2$, then $a = 4$ and $c = 4$, so $abc = 32$. If $b = 3$, then $a = 2$ and $c = 5$, so $abc = 30$.

5

Conclude the proof

The list contains every possible case, and in every case the product is even. Therefore abc is always even.

FINAL ANSWER

WMA12-01_2022_Oct_Q01

The possible triples are $(6, 1, 3)$, $(4, 2, 4)$ and $(2, 3, 5)$, giving products 18, 32 and 30. Each product is even, so abc is always even.

T01

01_CASE

01_CASE_CORE

CHAPTER
ProofFAMILY
Exhaustion/casesVARIANT
core family move

5 marks

WMA12/01 MAY/JUNE 2023

Question 8

Past-paper question

WMA12-01_2023_MayJune_Q08

8. (i) A student writes the following statement:

“When a and b are consecutive **prime** numbers, $a^2 + b^2$ is never a multiple of 10”

Prove by counter example that this statement is **not** true.

(2)

- (ii) Given that x and y are even integers greater than 0 and less than 6, prove by exhaustion, that

$$1 < x^2 - \frac{xy}{4} < 15$$

(3)

T01 Proof

01_CASE Exhaustion/cases

01_CASE_CORE core family
move

WORKED SOLUTION

Question 8

WMA12/01 May/June 2023

WMA12-01_2023_MayJune_Q08

1

Choose consecutive primes

7 and 11 are consecutive prime numbers.

2

Check the statement $7^2 + 11^2 = 49 + 121 = 170$, which is a multiple of 10.

3

Conclude the counterexample

This single counterexample proves the student's statement is not true.

4

List the possible even valuesThe even integers greater than 0 and less than 6 are 2 and 4. Therefore the possible ordered pairs are $(2, 2)$, $(2, 4)$, $(4, 2)$, $(4, 4)$.

5

Check x equals twoIf $(x, y) = (2, 2)$, then $x^2 - \frac{xy}{4} = 4 - 1 = 3$. If $(x, y) = (2, 4)$, then $x^2 - \frac{xy}{4} = 4 - 2 = 2$.

6

Check x equals fourIf $(x, y) = (4, 2)$, then $x^2 - \frac{xy}{4} = 16 - 2 = 14$. If $(x, y) = (4, 4)$, then $x^2 - \frac{xy}{4} = 16 - 4 = 12$.

7

Complete the exhaustionThe possible values are 3, 2, 14, 12, and each lies strictly between 1 and 15. Hence $1 < x^2 - \frac{xy}{4} < 15$ in every possible case.**FINAL ANSWER**

WMA12-01_2023_MayJune_Q08

(i) A counterexample is $a = 7, b = 11$, since $7^2 + 11^2 = 170$. (ii) Exhaustion proves the inequality for $(x, y) = (2, 2), (2, 4), (4, 2), (4, 4)$.

T01

01_CASE

01_CASE_CORE

CHAPTER
Proof

FAMILY
Exhaustion/cases

VARIANT
core family move
3 marks

WMA12/01 OCTOBER 2023

Question 1

Past-paper question

WMA12-01_2023_Oct_Q01

1. Given that a , b and c are integers greater than 0 such that
- $c = 3a + 1$
 - $a + b + c = 15$

prove, by exhaustion, that the product abc is always a multiple of 4

You may use the table below to illustrate your answer.

(3)

You may not need to use all rows of this table.

a	b	c	abc

T01 Proof

01_CASE Exhaustion/cases

01_CASE_CORE core family
move

WORKED SOLUTION

Question 1

WMA12/01 October 2023

WMA12-01_2023_Oct_Q01

1

Use the conditions

Since $c = 3a + 1$ and $a + b + c = 15$, substitute for c to get $a + b + 3a + 1 = 15$.

2

Write b in terms of a

This gives $b = 14 - 4a$.

3

List possible a values

The integers a, b, c are positive. Since $b = 14 - 4a > 0$, we need $a < 3.5$. Therefore $a = 1, 2, 3$.

4

Check each case

If $a = 1$, then $b = 10$ and $c = 4$, so $abc = 40$. If $a = 2$, then $b = 6$ and $c = 7$, so $abc = 84$. If $a = 3$, then $b = 2$ and $c = 10$, so $abc = 60$.

5

Conclude the exhaustion

The products 40, 84 and 60 are all multiples of 4. All possible cases have been checked, so abc is always a multiple of 4.

FINAL ANSWER

WMA12-01_2023_Oct_Q01

The only possible triples are $(1, 10, 4)$, $(2, 6, 7)$ and $(3, 2, 10)$, giving products 40, 84 and 60. Each is a multiple of 4, so abc is always a multiple of 4.

T01

01_CASE

01_CASE_CORE

CHAPTER
ProofFAMILY
Exhaustion/casesVARIANT
core family move

6 marks

WMA12/01 JANUARY 2024

Question 8

Past-paper question

WMA12-01_2024_Jan_Q08

8. (i) Use a counter example to show that the following statement is **false**

“ $n^2 + 3n + 1$ is prime for all $n \in \mathbb{N}$ ”

(2)

- (ii) Use algebra to prove by exhaustion that for all $n \in \mathbb{N}$

“ $n^2 - 2$ is **not** a multiple of 4”

(4)

T01 Proof

01_CASE Exhaustion/cases

01_CASE_CORE core family
move

WORKED SOLUTION

Question 8

WMA12/01 January 2024

WMA12-01_2024_Jan_Q08

1

Choose a counterexampleTake $n = 6$.

2

Evaluate the expression

$$n^2 + 3n + 1 = 6^2 + 3(6) + 1 = 36 + 18 + 1 = 55.$$

3

Conclude false

55 is not prime, so the statement is false.

4

Use residues modulo fourEvery natural number is one of $4k$, $4k + 1$, $4k + 2$ or $4k + 3$.

5

Check n equals $4k$ If $n = 4k$, then $n^2 - 2 = 16k^2 - 2 = 4(4k^2 - 1) + 2$, not a multiple of 4.

6

Check n equals $4k$ plus oneIf $n = 4k + 1$, then $n^2 - 2 = 16k^2 + 8k - 1 = 4(4k^2 + 2k - 1) + 3$, not a multiple of 4.

7

Check n equals $4k$ plus twoIf $n = 4k + 2$, then $n^2 - 2 = 16k^2 + 16k + 2 = 4(4k^2 + 4k) + 2$, not a multiple of 4.

8

Check n equals $4k$ plus threeIf $n = 4k + 3$, then $n^2 - 2 = 16k^2 + 24k + 7 = 4(4k^2 + 6k + 1) + 3$, not a multiple of 4.

9

Complete the proofAll possible cases leave remainder 2 or 3 when divided by 4, so $n^2 - 2$ is not a multiple of 4.**FINAL ANSWER**

WMA12-01_2024_Jan_Q08

(i) **Counterexample:** $n = 6$ gives 55, which is not prime. (ii) **Proven by considering** $n = 4k, 4k + 1, 4k + 2, 4k + 3$.

T01

01_CASE

01_CASE_CORE

CHAPTER
ProofFAMILY
Exhaustion/casesVARIANT
core family move

6 marks

WMA12/01 OCTOBER 2024

Question 11

Past-paper question

WMA12-01_2024_Oct_Q11

11. (i) Prove by counter example that the statement

“If n is a prime number then $3^n + 2$ is also a prime number.”

is false.

(2)

(ii) Use proof by exhaustion to prove that if m is an integer that is **not** divisible by 3, then

$$m^2 - 1$$

is divisible by 3

(4)

T01 Proof

01_CASE Exhaustion/cases

01_CASE_CORE core family
move

WORKED SOLUTION

Question 11

WMA12/01 October 2024

WMA12-01_2024_Oct_Q11

1

Choose a prime value

Take $n = 5$, which is prime.

2

Evaluate the expression

$3^5 + 2 = 243 + 2 = 245 = 5 \times 49$, so it is not prime. This disproves the statement.

3

List the possible forms of m

If integer m is not divisible by 3, then m has remainder 1 or 2 on division by 3.

4

Case one

If $m = 3q + 1$, then $m^2 - 1 = (3q + 1)^2 - 1 = 9q^2 + 6q = 3(3q^2 + 2q)$, which is divisible by 3.

5

Case two

If $m = 3q + 2$, then $m^2 - 1 = (3q + 2)^2 - 1 = 9q^2 + 12q + 3 = 3(3q^2 + 4q + 1)$, which is divisible by 3.

6

Conclude by exhaustion

Both possible cases for an integer not divisible by 3 have been checked, so $m^2 - 1$ is divisible by 3.

FINAL ANSWER

WMA12-01_2024_Oct_Q11

(i) **Counterexample:** $n = 5$ gives $3^5 + 2 = 245$, not prime. (ii) **Proven.**

T01

01_CASE

01_CASE_CORE

CHAPTER
ProofFAMILY
Exhaustion/casesVARIANT
core family move

4 marks

WMA12/01 MAY/JUNE 2025

Question 10

Past-paper question

WMA12-01_2025_MayJune_Q10

10: In this question you must show detailed reasoning.

Use algebra to prove by exhaustion that,

for all positive integers m that are **not** multiples of 3, the value of

$$m^2 + 3m + 2$$

is always a multiple of 3

(4)

T01 Proof

01_CASE Exhaustion/cases

01_CASE_CORE core family
move

WORKED SOLUTION

Question 10

WMA12/01 May/June 2025

WMA12-01_2025_MayJune_Q10

1

List the possible cases

A positive integer that is not a multiple of 3 must have the form $m = 3q + 1$ or $m = 3q + 2$.

2

Case one

If $m = 3q + 1$, then $m^2 + 3m + 2 = (3q + 1)^2 + 3(3q + 1) + 2 = 9q^2 + 15q + 6 = 3(3q^2 + 5q + 2)$.

3

Case two

If $m = 3q + 2$, then $m^2 + 3m + 2 = (3q + 2)^2 + 3(3q + 2) + 2 = 9q^2 + 21q + 12 = 3(3q^2 + 7q + 4)$.

4

Conclude by exhaustion

Both possible cases give a factor of 3, so $m^2 + 3m + 2$ is always a multiple of 3.

FINAL ANSWER**Proven.**

WMA12-01_2025_MayJune_Q10

T01

01_CASE

01_CASE_HV01

CHAPTER
Proof

FAMILY
Exhaustion/cases

VARIANT
For modulo cases, use
/(3k,3k+1,3k+2/) or
/(4k,4k+1,4k+2,4k+3/).
4 marks

WMA12/01 JANUARY 2023
Question 10

Past-paper question

WMA12-01_2023_Jan_Q10

10. A student was asked to prove by exhaustion that

if n is an integer then $2n^2 + n + 1$ is **not** divisible by 3

The start of the student's proof is shown in the box below.

Consider the case when $n = 3k$

$$2n^2 + n + 1 = 18k^2 + 3k + 1 = 3(6k^2 + k) + 1$$

which is not divisible by 3

Complete this proof.

(4)

T01 Proof

01_CASE Exhaustion/cases

01_CASE_HV01 For modulo cases, use $\wedge(3k, 3k+1, 3k+2)$ or $\wedge(4k, 4k+1, 4k+2, 4k+3)$.

WORKED SOLUTION

Question 10

WMA12/01 January 2023

WMA12-01_2023_Jan_Q10

1

Use the three possible formsEvery integer is exactly one of the forms $3k$, $3k + 1$ or $3k + 2$, where k is an integer.

2

First case already givenFor $n = 3k$, the expression becomes $3(6k^2 + k) + 1$, so it leaves remainder 1 when divided by 3.

3

Try n equals $3k$ plus 1If $n = 3k + 1$, then $2n^2 + n + 1 = 2(3k + 1)^2 + (3k + 1) + 1$.

4

Simplify the second case $2(9k^2 + 6k + 1) + 3k + 2 = 18k^2 + 15k + 4 = 3(6k^2 + 5k + 1) + 1$, which is not divisible by 3.

5

Try n equals $3k$ plus 2If $n = 3k + 2$, then $2n^2 + n + 1 = 2(3k + 2)^2 + (3k + 2) + 1$.

6

Simplify the third case $2(9k^2 + 12k + 4) + 3k + 3 = 18k^2 + 27k + 11 = 3(6k^2 + 9k + 3) + 2$, which is not divisible by 3.

7

Complete the exhaustionAll possible integer forms have been checked, and none gives a multiple of 3. Therefore $2n^2 + n + 1$ is not divisible by 3.**FINAL ANSWER**

WMA12-01_2023_Jan_Q10

For $n = 3k$, $n = 3k + 1$ and $n = 3k + 2$, the expression has remainders 1, 1 and 2 respectively when divided by 3. Therefore $2n^2 + n + 1$ is never divisible by 3.

TOPIC 01 | FAMILY 02

Disproof

WHAT THIS FAMILY ASKS

Recognise the disproof demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE**V01** core family move**2 classified questions**

13 total marks

T01

01_COUNTER

01_COUNTER_CORE

CHAPTER

Proof

FAMILY

Disproof

VARIANT

core family move

7 marks

WMA12/01 MAY/JUNE 2022

Question 3

Past-paper question

WMA12-01_2022_MayJune_Q03

3. (i) Show that the following statement is **false**:

“($n + 1$)³ – n ³ is prime for all $n \in \mathbb{N}$ ”

(2)

- (ii) Given that the points $A(1, 0)$, $B(3, -10)$ and $C(7, -6)$ lie on a circle, prove that AB is a diameter of this circle.

(5)

T01 Proof

01_COUNTER Disproof

01_COUNTER_CORE core
family move

WORKED SOLUTION

Question 3

WMA12/01 May/June 2022

WMA12-01_2022_MayJune_Q03

1

Choose a counterexampleTake $n = 5$.

2

Evaluate the expression

$$(5 + 1)^3 - 5^3 = 216 - 125 = 91.$$

3

Show it is not prime $91 = 7 \times 13$, so the statement is false.

4

Find the gradient CAFor $A(1, 0)$ and $C(7, -6)$, the gradient of CA is $\frac{0 - (-6)}{1 - 7} = -1$.

5

Find the gradient CBFor $B(3, -10)$ and $C(7, -6)$, the gradient of CB is $\frac{-10 - (-6)}{3 - 7} = 1$.

6

Show the right angleThe product of the gradients is -1 , so CA is perpendicular to CB and $\angle ACB = 90^\circ$.

7

Conclude the diameterAn angle of 90° at the circumference subtends a diameter, so AB is a diameter of the circle.**FINAL ANSWER**

WMA12-01_2022_MayJune_Q03

(i) Counterexample: $n = 5$ gives 91, not prime. (ii) Proven: AB is a diameter.

T01

01_COUNTER

01_COUNTER_CORE

CHAPTER

Proof

FAMILY

Disproof

VARIANT

core family move

6 marks

WMA12/01 MAY/JUNE 2024

Question 5

Past-paper question

WMA12-01_2024_MayJune_Q05

5. In this question you must show detailed reasoning.

- (a) Given that x and y are positive numbers such that

$$(x - y)^3 > x^3 - y^3$$

prove that

$$y > x$$

(4)

- (b) Using a counter example, show that the result in part (a) is not true for all real numbers.

(2)

T01 Proof

01_COUNTER Disproof

01_COUNTER_CORE core
family move

WORKED SOLUTION

Question 5

WMA12/01 May/June 2024

WMA12-01_2024_MayJune_Q05

1

Start from the inequality

$$(x - y)^3 > x^3 - y^3.$$

2

Expand the left side

$$x^3 - 3x^2y + 3xy^2 - y^3 > x^3 - y^3.$$

3

Cancel common terms

Subtract $x^3 - y^3$ from both sides to get $-3x^2y + 3xy^2 > 0$.

4

Factor

$$3xy(y - x) > 0.$$

5

Use positivity

Since x and y are positive, $3xy > 0$. Therefore $y - x > 0$, so $y > x$.

6

Give a real-number counterexample

Take $x = 1$ and $y = -1$. Then $(x - y)^3 = 2^3 = 8$ and $x^3 - y^3 = 1 - (-1) = 2$, so the given inequality is true.

7

Show the conclusion fails

But $y > x$ would mean $-1 > 1$, which is false. Therefore the result is not true for all real numbers.

FINAL ANSWER

WMA12-01_2024_MayJune_Q05

(a) **Proven.** (b) **For example, $x = 1, y = -1$ satisfies the inequality but not $y > x$.**

TOPIC 01 | FAMILY 03

Algebraic Divisibility

WHAT THIS FAMILY ASKS

Recognise the algebraic divisibility demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE**V01** core family move**3 classified questions**

18 total marks

T01

01_FACTOR

01_FACTOR_CORE

CHAPTER

Proof

FAMILY

Algebraic divisibility

VARIANT

core family move

5 marks

WMA12/01 MAY/JUNE 2021

Question 3

Past-paper question

WMA12-01_2021_MayJune_Q03

3. (i) Prove that for all single digit prime numbers, p ,

$$p^3 + p \text{ is a multiple of } 10$$

(2)

- (ii) Show, using algebra, that for $n \in \mathbb{N}$

$$(n + 1)^3 - n^3 \text{ is not a multiple of } 3$$

(3)

T01 Proof

01_FACTOR Algebraic
divisibility01_FACTOR_CORE core
family move

WORKED SOLUTION

Question 3

WMA12/01 May/June 2021

WMA12-01_2021_MayJune_Q03

1

List the cases

The single digit prime numbers are 2, 3, 5, and 7.

2

Check each prime $2^3 + 2 = 10$, $3^3 + 3 = 30$, $5^3 + 5 = 130$, and $7^3 + 7 = 350$.

3

Conclude part oneEach result is a multiple of 10, so $p^3 + p$ is a multiple of 10 for all single digit primes.

4

Expand the difference of cubes $(n + 1)^3 - n^3 = (n^3 + 3n^2 + 3n + 1) - n^3 = 3n^2 + 3n + 1$.

5

Factor the multiple of three $3n^2 + 3n + 1 = 3n(n + 1) + 1$.

6

Use the remainder $3n(n + 1)$ is a multiple of 3, so $3n(n + 1) + 1$ is one more than a multiple of 3. Therefore it is not a multiple of 3.**FINAL ANSWER**

WMA12-01_2021_MayJune_Q03

(i) **Proven by checking** $p = 2, 3, 5, 7$. (ii) **Proven:** $(n + 1)^3 - n^3$ **leaves remainder 1 on division by 3.**

T01

01_FACTOR

01_FACTOR_CORE

CHAPTER

Proof

FAMILY

Algebraic divisibility

VARIANT

core family move

5 marks

WMA12/01 JANUARY 2025

Question 8

Past-paper question

WMA12-01_2025_Jan_Q08

8. (i) A student states

“If x and y are irrational numbers, $x \neq y$, then xy is also irrational.”

Show, by counter example, that this statement is not always true.

(1)

- (ii) Prove, using algebra, that for all odd integers n , the value of the expression

$$n^3 + 3n + 2$$

is always even but never a multiple of 4

(4)

T01 Proof

01_FACTOR Algebraic
divisibility01_FACTOR_CORE core
family move

WORKED SOLUTION

Question 8

WMA12/01 January 2025

WMA12-01_2025_Jan_Q08

1

Give a counterexample

Let $x = \sqrt{2}$ and $y = \frac{1}{\sqrt{2}}$. Both are irrational and $x \neq y$, but $xy = 1$, which is rational.

2

Represent an odd integer

Let $n = 2q + 1$, where q is an integer.

3

Substitute into the expression

$$n^3 + 3n + 2 = (2q + 1)^3 + 3(2q + 1) + 2.$$

4

Expand and factor

$$n^3 + 3n + 2 = 8q^3 + 12q^2 + 12q + 6 = 2(4q^3 + 6q^2 + 6q + 3).$$

5

Show it is even

The expression has a factor of 2, so it is even.

6

Show it is not a multiple of four

The bracket $4q^3 + 6q^2 + 6q + 3$ is odd, because the first three terms are even and 3 is odd. Therefore the expression is 2 times an odd integer, so it is not divisible by 4.

FINAL ANSWER

WMA12-01_2025_Jan_Q08

(i) For example, $x = \sqrt{2}$ and $y = \frac{1}{\sqrt{2}}$ give $xy = 1$. (ii) Proven.

T01

01_FACTOR

01_FACTOR_CORE

CHAPTER

Proof

FAMILY

Algebraic divisibility

VARIANT

core family move

8 marks

WMA12/01 JANUARY 2026

Question 6

Past-paper question

WMA12-01_2026_Jan_Q06

6. (i) Given that p and q are consecutive odd numbers, where $p > q > 0$, prove that

$$p^2 - q^2$$

is a multiple of 8

(4)

- (ii) The curve C has equation

$$y = x^3 + 12x^2 + 49x + 2$$

Prove that C has no stationary points.

(4)

T01 Proof

01_FACTOR Algebraic
divisibility01_FACTOR_CORE core
family move

WORKED SOLUTION

Question 6

WMA12/01 January 2026

WMA12-01_2026_Jan_Q06

1

Represent the consecutive odd numbers

Let $q = 2n - 1$ and $p = 2n + 1$, where n is a positive integer. Then $p > q > 0$ and the numbers are consecutive odd numbers.

2

Subtract the squares

$$p^2 - q^2 = (2n + 1)^2 - (2n - 1)^2.$$

3

Simplify the expression

$$p^2 - q^2 = (4n^2 + 4n + 1) - (4n^2 - 4n + 1) = 8n.$$

4

Conclude the proof

Since n is an integer, $8n$ is a multiple of 8, so $p^2 - q^2$ is a multiple of 8.

5

Differentiate the curve

For $y = x^3 + 12x^2 + 49x + 2$, $\frac{dy}{dx} = 3x^2 + 24x + 49$.

6

Show the gradient is never zero

$$3x^2 + 24x + 49 = 3(x + 4)^2 + 1. \text{ This is always positive for real } x.$$

7

State the conclusion

Since $\frac{dy}{dx}$ is never 0, the curve has no stationary points.

FINAL ANSWER

WMA12-01_2026_Jan_Q06

(i) **Proven.** (ii) **C has no stationary points.**

TOPIC 01 | FAMILY 04

Inequality Proof

WHAT THIS FAMILY ASKS

Recognise the inequality proof demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

2 classified questions

8 total marks

T01

01_INEQ

01_INEQ_CORE

CHAPTER
Proof

FAMILY
Inequality proof

VARIANT
core family move

4 marks

WMA12/01 JANUARY 2021

Question 5

Past-paper question

WMA12-01_2021_Jan_Q05

5. (i) Use algebra to prove that for all $x \geq 0$

$$3x + 1 \geq 2\sqrt{3x}$$

(3)

- (ii) Show that the following statement is not true.

“The sum of three consecutive prime numbers is always a multiple of 5”

(1)

T01 Proof

01_INEQ Inequality proof

01_INEQ_CORE core family
move

WORKED SOLUTION

Question 5

WMA12/01 January 2021

WMA12-01_2021_Jan_Q05

1

Start from a square

For $x \geq 0$, $\sqrt{3x}$ is real and $(\sqrt{3x} - 1)^2 \geq 0$.

2

Expand the square
$$(\sqrt{3x} - 1)^2 = 3x - 2\sqrt{3x} + 1 \geq 0.$$

3

Rearrange the inequality

Therefore $3x + 1 \geq 2\sqrt{3x}$, as required.

4

Give a counterexample

The consecutive prime numbers 5, 7, and 11 have sum $5 + 7 + 11 = 23$.

5

Complete the disproof

23 is not a multiple of 5, so the statement is not true.

FINAL ANSWER

WMA12-01_2021_Jan_Q05

(i) **Proven.** (ii) **For example, $5 + 7 + 11 = 23$, which is not a multiple of 5.**

T01

01_INEQ

01_INEQ_CORE

CHAPTER
Proof

FAMILY
Inequality proof

VARIANT
core family move

4 marks

WMA12/01 OCTOBER 2021

Question 9

Past-paper question

WMA12-01_2021_Oct_Q09

9. (a) Prove that for all positive values of x and y ,

$$\frac{x+y}{2} \geq \sqrt{xy} \quad (3)$$

- (b) Prove by counter-example that this inequality does not hold when x and y are both negative.

(1)

T01 Proof

01_INEQ Inequality proof

01_INEQ_CORE core family
move

WORKED SOLUTION

Question 9

WMA12/01 October 2021

WMA12-01_2021_Oct_Q09

1

Start with a squareFor positive x and y , both $\sqrt{x}$ and $\sqrt{y}$ are real, so $(\sqrt{x} - \sqrt{y})^2 \geq 0$.

2

Expand

$$(\sqrt{x} - \sqrt{y})^2 = x - 2\sqrt{xy} + y \geq 0.$$

3

RearrangeThus $x + y \geq 2\sqrt{xy}$, and dividing by 2 gives $\frac{x+y}{2} \geq \sqrt{xy}$.

4

Give a negative counterexampleTake $x = -1$ and $y = -1$. Then $\frac{x+y}{2} = -1$, while $\sqrt{xy} = \sqrt{1} = 1$.

5

ConcludeSince $-1 \geq 1$ is false, the inequality does not hold for all negative x and y .**FINAL ANSWER**

WMA12-01_2021_Oct_Q09

(a) **Proven.** (b) **For example, $x = y = -1$ gives $-1 \not\geq 1$.**

PURE MATHEMATICS 2

ADAPTIVE ALL CLASSIFIED

Presentation Edition | Solutions

CLASSIFIED BY IDEA | ROOM TO THINK

Every demand appears clearly, with practical space for the mathematics.

Dr Eslam Ahmed

Elite Mathematics

ElitelGCSE.com | 011 2000 9622

PURE 2

PURE 2 | TOPIC 02

Polynomials

Your classified route

15

QUESTIONS

117

MARKS

4

FAMILIES

4

VARIANTS

DR ESLAM AHMED

CLASSIFIED BY IDEA

Move from recognition to method, then use the working space to complete the solution.

TOPIC 02 | FAMILY 01

Polynomial Division

WHAT THIS FAMILY ASKS

Recognise the polynomial division demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

2 classified questions

19 total marks

T02

02_DIV

02_DIV_CORE

CHAPTER

Polynomials

FAMILY

Polynomial division

VARIANT

core family move

12 marks

WMA12/01 OCTOBER 2025

Question 4

Past-paper question

WMA12-01_2025_Oct_Q04

4.

In this question you must show all stages of your working.
Solutions relying on calculator technology are not acceptable.

$$f(x) = 4x^3 + 13x^2 - 10x + 8$$

(a) When $f(x)$ is divided by $(x - 2)$ the remainder is R and the quotient is $Q(x)$.

(i) Find $Q(x)$.

(ii) Find R .

(4)

(b) (i) Use the factor theorem to show that $(x + 4)$ is a factor of $f(x)$.

(ii) Hence prove, using algebra, that the equation $f(x) = 0$ has only one real solution.

(5)

(c) Find the range of values of x for which $f(x)$ is decreasing.

(3)

T02 Polynomials

02_DIV Polynomial division

02_DIV_CORE core family
move

WORKED SOLUTION

Question 4

WMA12/01 October 2025

WMA12-01_2025_Oct_Q04

1

Divide by x minus two

$$4x^3 + 13x^2 - 10x + 8 = (x - 2)(4x^2 + 21x + 32) + 72.$$

2

State Q and R

$$\text{Hence } Q(x) = 4x^2 + 21x + 32 \text{ and } R = 72.$$

3

Use the factor theorem

$$f(-4) = 4(-4)^3 + 13(-4)^2 - 10(-4) + 8 = -256 + 208 + 40 + 8 = 0.$$

4

Show the factor

Since $f(-4) = 0$, $(x + 4)$ is a factor of $f(x)$.

5

Factor the cubic

$$\text{Dividing by } (x + 4) \text{ gives } f(x) = (x + 4)(4x^2 - 3x + 2).$$

6

Check the quadratic

The discriminant of $4x^2 - 3x + 2$ is $(-3)^2 - 4(4)(2) = 9 - 32 = -23 < 0$, so this quadratic has no real roots.

7

Conclude the real solution

Therefore $f(x) = 0$ has only one real solution, $x = -4$.

8

Differentiate

$$f'(x) = 12x^2 + 26x - 10 = 2(3x - 1)(2x + 5).$$

9

Find where f is decreasing

$f(x)$ is decreasing when $f'(x) < 0$, which is between the roots. Hence $-\frac{5}{2} < x < \frac{1}{3}$.

FINAL ANSWER

WMA12-01_2025_Oct_Q04

(a)(i) $Q(x) = 4x^2 + 21x + 32$. (a)(ii) $R = 72$. (b) **Proven; only real solution is**
 $x = -4$. (c) $-\frac{5}{2} < x < \frac{1}{3}$.

T02

02_DIV

02_DIV_CORE

CHAPTER

Polynomials

FAMILY

Polynomial division

VARIANT

core family move

7 marks

WMA12/01 JANUARY 2026

Question 3

Past-paper question

WMA12-01_2026_Jan_Q03

3. (i) Given that

- $f(x) = 4x^3 + 6x + k$, where k is a constant
- $(x + 2)$ is a factor of $f(x)$

find the remainder when $f(x)$ is divided by $(x - 5)$

(4)

(ii) Find the remainder, R , and the quotient, $Q(x)$, when

$$6x^3 - 15x^2 - 21x + 8$$

is divided by $(2x + 3)$

(3)

T02 Polynomials

02_DIV Polynomial division

02_DIV_CORE core family
move

WORKED SOLUTION

Question 3

WMA12/01 January 2026

WMA12-01_2026_Jan_Q03

1

Use the factor theorem

Since $(x + 2)$ is a factor of $f(x) = 4x^3 + 6x + k$, we have $f(-2) = 0$.

2

Find k

$4(-2)^3 + 6(-2) + k = 0$, so $-32 - 12 + k = 0$ and $k = 44$.

3

Find the required remainder

The remainder on division by $(x - 5)$ is $f(5) = 4(5)^3 + 6(5) + 44 = 574$.

4

Set up the division

Write $6x^3 - 15x^2 - 21x + 8 = (2x + 3)(Ax^2 + Bx + C) + R$.

5

Match coefficients

Comparing coefficients gives $A = 3$, $B = -12$, and $C = \frac{15}{2}$.

6

Find the remainder

Then $(2x + 3)\left(3x^2 - 12x + \frac{15}{2}\right) = 6x^3 - 15x^2 - 21x + \frac{45}{2}$, so $R = 8 - \frac{45}{2} = -\frac{29}{2}$.

FINAL ANSWER

WMA12-01_2026_Jan_Q03

(i) 574. (ii) $Q(x) = 3x^2 - 12x + \frac{15}{2}$, $R = -\frac{29}{2}$.

TOPIC 02 | FAMILY 02

Factor Theorem

WHAT THIS FAMILY ASKS

Recognise the factor theorem demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE**V01** core family move**7 classified questions**

58 total marks

T02

02_FACT

02_FACT_CORE

CHAPTER
Polynomials

FAMILY
Factor theorem

VARIANT
core family move

8 marks

WMA12/01 JANUARY 2020

Question 3

Past-paper question

WMA12-01_2020_Jan_Q03

3. $f(x) = 6x^3 + 17x^2 + 4x - 12$

(a) Use the factor theorem to show that $(2x + 3)$ is a factor of $f(x)$.

(2)

(b) Hence, using algebra, write $f(x)$ as a product of three linear factors.

(4)

(c) Solve, for $\frac{\pi}{2} < \theta < \pi$, the equation

$$6 \tan^3 \theta + 17 \tan^2 \theta + 4 \tan \theta - 12 = 0$$

giving your answers to 3 significant figures.

(2)

T02 Polynomials

02_FACT Factor theorem

02_FACT_CORE core family
move

WORKED SOLUTION

Question 3

WMA12/01 January 2020

WMA12-01_2020_Jan_Q03

1

Use the factor theorem

For $(2x + 3)$, use $x = -\frac{3}{2}$. Substituting gives $f\left(-\frac{3}{2}\right) = 6\left(-\frac{3}{2}\right)^3 + 17\left(-\frac{3}{2}\right)^2 + 4\left(-\frac{3}{2}\right) - 12 = 0$.

2

State the factor result

Since $f\left(-\frac{3}{2}\right) = 0$, $(2x + 3)$ is a factor of $f(x)$.

3

Divide by the known factor

$$6x^3 + 17x^2 + 4x - 12 = (2x + 3)(3x^2 + 4x - 4).$$

4

Factorise the quadratic

$$3x^2 + 4x - 4 = (3x - 2)(x + 2), \text{ so } f(x) = (2x + 3)(3x - 2)(x + 2).$$

5

Use the factorisation for tan theta

Let $t = \tan \theta$. Then $(2t + 3)(3t - 2)(t + 2) = 0$, so $t = -\frac{3}{2}$, $t = \frac{2}{3}$, or $t = -2$.

6

Choose the roots in the interval

For $\frac{\pi}{2} < \theta < \pi$, $\tan \theta$ is negative, so use $\tan \theta = -\frac{3}{2}$ and $\tan \theta = -2$.

7

Find theta

This gives $\theta = \pi - \tan^{-1}\left(\frac{3}{2}\right) = 2.16$ and $\theta = \pi - \tan^{-1}(2) = 2.03$, to 3 significant figures.

FINAL ANSWER

WMA12-01_2020_Jan_Q03

(a) **Proven.** (b) $f(x) = (2x + 3)(3x - 2)(x + 2)$. (c) $\theta = 2.03, 2.16$ **radians.**

T02

02_FACT

02_FACT_CORE

CHAPTER
Polynomials

FAMILY
Factor theorem

VARIANT
core family move

8 marks

WMA12/01 OCTOBER 2021

Question 4

Past-paper question

WMA12-01_2021_Oct_Q04

4.

$$f(x) = (x^2 - 2)(2x - 3) - 21$$

- (a) State the value of the remainder when $f(x)$ is divided by $(2x - 3)$

(1)

- (b) Use the factor theorem to show that $(x - 3)$ is a factor of $f(x)$

(2)

- (c) Hence,

- (i) factorise $f(x)$

- (ii) show that the equation $f(x) = 0$ has only one real root.

(5)

T02 Polynomials

02_FACT Factor theorem

02_FACT_CORE core family
move

WORKED SOLUTION

Question 4

WMA12/01 October 2021

WMA12-01_2021_Oct_Q04

1

Use the remainder theoremDividing by $(2x - 3)$ means use $x = \frac{3}{2}$.

2

Find the remainder

$$f\left(\frac{3}{2}\right) = \left(\frac{9}{4} - 2\right)(0) - 21 = -21.$$

3

Check x equals three

$$f(3) = (9 - 2)(6 - 3) - 21 = 7 \cdot 3 - 21 = 0.$$

4

Use the factor theoremSince $f(3) = 0$, $(x - 3)$ is a factor of $f(x)$.

5

Expand f

$$f(x) = (x^2 - 2)(2x - 3) - 21 = 2x^3 - 3x^2 - 4x - 15.$$

6

FactoriseDividing by $(x - 3)$ gives $2x^2 + 3x + 5$, so $f(x) = (x - 3)(2x^2 + 3x + 5)$.

7

Check the quadratic rootsFor $2x^2 + 3x + 5$, the discriminant is $3^2 - 4(2)(5) = 9 - 40 = -31 < 0$.

8

ConcludeThe quadratic has no real roots, so $f(x) = 0$ has only one real root, $x = 3$.**FINAL ANSWER**

WMA12-01_2021_Oct_Q04

(a) -21 . (b) **Proven.** (c)(i) $f(x) = (x - 3)(2x^2 + 3x + 5)$. (c)(ii) **Only real root:**
 $x = 3$.

T02

02_FACT

02_FACT_CORE

CHAPTER
Polynomials

FAMILY
Factor theorem

VARIANT
core family move

9 marks

WMA12/01 JANUARY 2023

Question 5

Past-paper question

WMA12-01_2023_Jan_Q05

5.
$$f(x) = x^3 + (p + 3)x^2 - x + q$$

where p and q are constants and $p > 0$

Given that $(x - 3)$ is a factor of $f(x)$

(a) show that

$$9p + q = -51 \quad (2)$$

Given also that when $f(x)$ is divided by $(x + p)$ the remainder is 9

(b) show that

$$3p^2 + p + q - 9 = 0 \quad (2)$$

(c) Hence find the value of p and the value of q .

(3)

(d) Hence find a quadratic expression $g(x)$ such that

$$f(x) = (x - 3)g(x) \quad (2)$$

T02 Polynomials

02_FACT Factor theorem

02_FACT_CORE core family
move

WORKED SOLUTION

Question 5

WMA12/01 January 2023

WMA12-01_2023_Jan_Q05

1

Use the factor theoremSince $(x - 3)$ is a factor, $f(3) = 0$.

2

Substitute x equals three $27 + 9(p + 3) - 3 + q = 0$, so $51 + 9p + q = 0$.

3

Show the first relationTherefore $9p + q = -51$.

4

Use the remainder theoremDividing by $(x + p)$ means substituting $x = -p$. The remainder is $f(-p)$.

5

Substitute x equals negative p $f(-p) = -p^3 + (p + 3)p^2 + p + q = 3p^2 + p + q$.

6

Show the second relationThe remainder is 9, so $3p^2 + p + q = 9$, hence $3p^2 + p + q - 9 = 0$.

7

Eliminate qFrom $9p + q = -51$, $q = -51 - 9p$. Substitute into $3p^2 + p + q - 9 = 0$.

8

Solve for p $3p^2 + p - 51 - 9p - 9 = 0$, so $3p^2 - 8p - 60 = 0$. This gives $p = 6$ or $p = -\frac{10}{3}$.

9

Use p greater than zeroSince $p > 0$, $p = 6$. Then $q = -51 - 9(6) = -105$.

10

Form f with p and q $f(x) = x^3 + 9x^2 - x - 105$.

11

Divide by x minus three $x^3 + 9x^2 - x - 105 = (x - 3)(x^2 + 12x + 35)$, so $g(x) = x^2 + 12x + 35$.**FINAL ANSWER**

WMA12-01_2023_Jan_Q05

(a) **Proven.** (b) **Proven.** (c) $p = 6, q = -105$. (d) $g(x) = x^2 + 12x + 35$.

DR ESLAM AHMED

T02

02_FACT

02_FACT_CORE

CHAPTER
Polynomials

FAMILY
Factor theorem

VARIANT
core family move

9 marks

WMA12/01 OCTOBER 2023

Question 4

Past-paper question

WMA12-01_2023_Oct_Q04

4. In this question you must show all stages of your working.
Solutions relying on calculator technology are not acceptable.

$$f(x) = 4x^3 + ax^2 - 29x + b$$

where a and b are constants.

Given that $(2x + 1)$ is a factor of $f(x)$,

- (a) show that

$$a + 4b = -56 \quad (2)$$

Given also that when $f(x)$ is divided by $(x - 2)$ the remainder is -25

- (b) find a second simplified equation linking a and b . (2)

- (c) Hence, using algebra and showing your working,

(i) find the value of a and the value of b ,

(ii) fully factorise $f(x)$. (5)

T02 Polynomials

02_FACT Factor theorem

02_FACT_CORE core family
move

WORKED SOLUTION

Question 4

WMA12/01 October 2023

WMA12-01_2023_Oct_Q04

1

Use the factor theorem

Since $(2x + 1)$ is a factor, $x = -\frac{1}{2}$ is a root, so $f\left(-\frac{1}{2}\right) = 0$.

2

Substitute x equals negative half

$$4\left(-\frac{1}{2}\right)^3 + a\left(-\frac{1}{2}\right)^2 - 29\left(-\frac{1}{2}\right) + b = 0.$$

3

Show the first relation

$-\frac{1}{2} + \frac{a}{4} + \frac{29}{2} + b = 0$. Multiplying by 4 gives $a + 4b + 56 = 0$, so $a + 4b = -56$.

4

Use the remainder theorem

Dividing by $(x - 2)$ gives remainder $f(2)$.

5

Substitute x equals two

$$f(2) = 4(8) + 4a - 58 + b = 4a + b - 26.$$

6

Find the second relation

The remainder is -25 , so $4a + b - 26 = -25$, hence $4a + b = 1$.

7

Solve the simultaneous equations

Solve $a + 4b = -56$ and $4a + b = 1$. Multiplying the second equation by 4 gives $16a + 4b = 4$.

8

Find a and b

Subtracting $a + 4b = -56$ from $16a + 4b = 4$ gives $15a = 60$, so $a = 4$. Then $16 + b = 1$, so $b = -15$.

9

Write f(x)

Therefore $f(x) = 4x^3 + 4x^2 - 29x - 15$.

10

Factorise

Using the known factors $(2x + 1)$ and $(x - 2)$, division gives $f(x) = (2x + 1)(x - 2)(2x + 15)$.

FINAL ANSWER

WMA12-01_2023_Oct_Q04

(a) **Proven.** (b) $4a + b = 1$. (c)(i) $a = 4, b = -15$. (c)(ii) $f(x) = (2x + 1)(x - 2)(2x + 15)$.

DR ESLAM AHMED

T02

02_FACT

02_FACT_CORE

CHAPTER
Polynomials

FAMILY
Factor theorem

VARIANT
core family move

8 marks

WMA12/01 MAY/JUNE 2024

Question 4

Past-paper question

WMA12-01_2024_MayJune_Q04

4. $f(x) = (x - 2)(2x^2 + 5x + k) + 21$

where k is a constant.

- (a) State the remainder when $f(x)$ is divided by $(x - 2)$ (1)

Given that $(2x - 1)$ is a factor of $f(x)$

- (b) show that $k = 11$ (2)

(c) Hence

- (i) fully factorise $f(x)$,
(ii) find the number of real solutions of the equation

$$f(x) = 0$$

giving a reason for your answer. (5)

T02 Polynomials

02_FACT Factor theorem

02_FACT_CORE core family
move

WORKED SOLUTION

Question 4

WMA12/01 May/June 2024

WMA12-01_2024_MayJune_Q04

1

Use the remainder theoremThe remainder on division by $(x - 2)$ is $f(2)$.

2

Find f of two

$$f(2) = (2 - 2)(2(2)^2 + 5(2) + k) + 21 = 21.$$

3

Use the factor theoremSince $(2x - 1)$ is a factor, $x = \frac{1}{2}$ is a root, so $f\left(\frac{1}{2}\right) = 0$.

4

Substitute x equals half

$$\left(\frac{1}{2} - 2\right) \left(2\left(\frac{1}{2}\right)^2 + 5\left(\frac{1}{2}\right) + k\right) + 21 = 0.$$

5

Solve for k

$$-\frac{3}{2}(3 + k) + 21 = 0. \text{ Multiplying by 2 gives } -3(k + 3) + 42 = 0, \text{ so } k = 11.$$

6

Write f with k equals eleven

$$f(x) = (x - 2)(2x^2 + 5x + 11) + 21 = 2x^3 + x^2 + x - 1.$$

7

Factorise

$$2x^3 + x^2 + x - 1 = (2x - 1)(x^2 + x + 1).$$

8

Count real solutions

The quadratic $x^2 + x + 1$ has discriminant $1 - 4 = -3$, so it has no real roots. Therefore $f(x) = 0$ has one real solution.

FINAL ANSWER

WMA12-01_2024_MayJune_Q04

(a) 21. (b) $k = 11$. (c)(i) $(2x - 1)(x^2 + x + 1)$. (c)(ii) **One real solution.**

T02

02_FACT

02_FACT_CORE

CHAPTER
Polynomials

FAMILY
Factor theorem

VARIANT
core family move

7 marks

WMA12/01 OCTOBER 2024

Question 3

Past-paper question

WMA12-01_2024_Oct_Q03

3.

$$f(x) = 2x^3 - x^2 + Ax + B$$

where A and B are integers.

Given that when $f(x)$ is divided by $(x + 3)$ the remainder is 55

(a) show that

$$3A - B = -118 \quad (2)$$

Given also that $(2x - 5)$ is a factor of $f(x)$,

(b) find the value of A and the value of B . (3)

(c) Hence find the quotient when $f(x)$ is divided by $(x - 7)$ (2)

T02 Polynomials

02_FACT Factor theorem

02_FACT_CORE core family
move

WORKED SOLUTION

Question 3

WMA12/01 October 2024

WMA12-01_2024_Oct_Q03

1

Use the remainder theoremDividing by $x + 3$ leaves remainder 55, so $f(-3) = 55$.

2

Show the given equation $f(-3) = 2(-3)^3 - (-3)^2 - 3A + B = -63 - 3A + B$. Therefore $-63 - 3A + B = 55$, so $3A - B = -118$.

3

Use the factor theoremSince $2x - 5$ is a factor, $x = \frac{5}{2}$ is a root and $f\left(\frac{5}{2}\right) = 0$.

4

Form a second equation $2\left(\frac{5}{2}\right)^3 - \left(\frac{5}{2}\right)^2 + \frac{5}{2}A + B = 0$, so $25 + \frac{5}{2}A + B = 0$. Multiplying by 2 gives $5A + 2B = -50$.

5

Solve simultaneouslyFrom $3A - B = -118$, $B = 3A + 118$. Substitute into $5A + 2B = -50$ to get $11A = -286$, so $A = -26$ and $B = 40$.

6

Write the polynomial $f(x) = 2x^3 - x^2 - 26x + 40$.

7

Divide by x minus seven $2x^3 - x^2 - 26x + 40 = (x - 7)(2x^2 + 13x + 65) + 495$, so the quotient is $2x^2 + 13x + 65$.**FINAL ANSWER**

WMA12-01_2024_Oct_Q03

(a) **Proven.** (b) $A = -26$, $B = 40$. (c) $2x^2 + 13x + 65$.

T02

02_FACT

02_FACT_CORE

CHAPTER
Polynomials

FAMILY
Factor theorem

VARIANT
core family move

9 marks

WMA12/01 JANUARY 2025

Question 5

Past-paper question

WMA12-01_2025_Jan_Q05

5.

$$f(x) = 3x^3 + ax^2 - 10x + b$$

where a and b are constants.

Given that $(3x - 4)$ is a factor of $f(x)$,

(a) show that $16a + 9b = 56$

(2)

Given further that when $f(x)$ is divided by $(x - 2)$ the remainder is b ,

(b) find the value of a and the value of b .

(4)

(c) Hence, using algebra, fully factorise $f(x)$.

(3)

T02 Polynomials

02_FACT Factor theorem

02_FACT_CORE core family
move

WORKED SOLUTION

Question 5

WMA12/01 January 2025

WMA12-01_2025_Jan_Q05

1

Use the factor theorem

Since $3x - 4$ is a factor, $x = \frac{4}{3}$ is a root and $f\left(\frac{4}{3}\right) = 0$.

2

Show the given equation

$3\left(\frac{4}{3}\right)^3 + a\left(\frac{4}{3}\right)^2 - 10\left(\frac{4}{3}\right) + b = 0$, so $\frac{64}{9} + \frac{16a}{9} - \frac{120}{9} + b = 0$. Multiplying by 9 gives $16a + 9b = 56$.

3

Use the remainder condition

When dividing by $x - 2$, the remainder is $f(2)$. Given that the remainder is b , $f(2) = b$.

4

Find a

$f(2) = 24 + 4a - 20 + b = 4 + 4a + b$. Thus $4 + 4a + b = b$, so $a = -1$.

5

Find b

Substitute $a = -1$ into $16a + 9b = 56$: $-16 + 9b = 56$, so $b = 8$.

6

Write f of x

$f(x) = 3x^3 - x^2 - 10x + 8$.

7

Find another factor

$f(1) = 3 - 1 - 10 + 8 = 0$, so $(x - 1)$ is a factor.

8

Fully factorise

Dividing by $(x - 1)$ gives $3x^2 + 2x - 8 = (3x - 4)(x + 2)$. Hence $f(x) = (x - 1)(x + 2)(3x - 4)$.

FINAL ANSWER

WMA12-01_2025_Jan_Q05

(a) **Proven.** (b) $a = -1, b = 8$. (c) $f(x) = (x - 1)(x + 2)(3x - 4)$.

TOPIC 02 | FAMILY 03

Remainder Theorem

WHAT THIS FAMILY ASKS

Recognise the remainder theorem demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE**V01** core family move**5 classified questions**

34 total marks

T02

02_REMA

02_REMA_CORE

CHAPTER
Polynomials

FAMILY
Remainder theorem

VARIANT
core family move

8 marks

WMA12/01 OCTOBER 2019

Question 4

Past-paper question

WMA12-01_2019_Oct_Q04

4. $f(x) = (x - 3)(3x^2 + x + a) - 35$ where a is a constant

(a) State the remainder when $f(x)$ is divided by $(x - 3)$.

(1)

Given $(3x - 2)$ is a factor of $f(x)$,

(b) show that $a = -17$

(2)

(c) Using algebra and showing each step of your working, fully factorise $f(x)$.

(5)

T02 Polynomials

02_REMA Remainder
theorem02_REMA_CORE core family
move

WORKED SOLUTION

Question 4

WMA12/01 October 2019

WMA12-01_2019_Oct_Q04

1

Use the remainder theorem

When $f(x)$ is divided by $(x - 3)$, the remainder is $f(3)$. Since $f(3) = (3 - 3)(3 \cdot 3^2 + 3 + a) - 35$, the remainder is -35 .

2

Use the given factor

Since $(3x - 2)$ is a factor, $x = \frac{2}{3}$ is a root and $f\left(\frac{2}{3}\right) = 0$.

3

Solve for a

$0 = \left(\frac{2}{3} - 3\right) \left(3 \left(\frac{2}{3}\right)^2 + \frac{2}{3} + a\right) - 35 = -\frac{7}{3}(a + 2) - 35$. Hence $-7(a + 2) = 105$, so $a = -17$.

4

Substitute a into f(x)

$f(x) = (x - 3)(3x^2 + x - 17) - 35 = 3x^3 - 8x^2 - 20x + 16$.

5

Divide by the known factor

Dividing $3x^3 - 8x^2 - 20x + 16$ by $(3x - 2)$ gives $x^2 - 2x - 8$.

6

Factorise fully

$x^2 - 2x - 8 = (x + 2)(x - 4)$, so $f(x) = (3x - 2)(x + 2)(x - 4)$.

FINAL ANSWER

WMA12-01_2019_Oct_Q04

(a) -35 . (b) $a = -17$. (c) $f(x) = (3x - 2)(x + 2)(x - 4)$.

T02

02_REMA

02_REMA_CORE

CHAPTER
Polynomials

FAMILY
Remainder theorem

VARIANT
core family move

6 marks

WMA12/01 JANUARY 2021

Question 1

Past-paper question

WMA12-01_2021_Jan_Q01

1.
$$f(x) = x^4 + ax^3 - 3x^2 + bx + 5$$

where a and b are constants.

When $f(x)$ is divided by $(x + 1)$, the remainder is 4

(a) Show that $a + b = -1$

(2)

When $f(x)$ is divided by $(x - 2)$, the remainder is -23

(b) Find the value of a and the value of b .

(4)

T02 Polynomials

02_REMA Remainder
theorem02_REMA_CORE core family
move

WORKED SOLUTION

Question 1

WMA12/01 January 2021

WMA12-01_2021_Jan_Q01

1

Use the first remainder

Dividing by $(x + 1)$ means use $x = -1$. Hence $f(-1) = 1 - a - 3 - b + 5 = 3 - a - b$.

2

Show the required relation

The remainder is 4, so $3 - a - b = 4$. Therefore $a + b = -1$.

3

Use the second remainder

Dividing by $(x - 2)$ means use $x = 2$. Hence $f(2) = 16 + 8a - 12 + 2b + 5 = 9 + 8a + 2b$.

4

Form the second equation

The remainder is -23 , so $9 + 8a + 2b = -23$, which gives $4a + b = -16$.

5

Solve simultaneously

Subtract $a + b = -1$ from $4a + b = -16$ to get $3a = -15$, so $a = -5$.

6

Find b

Using $a + b = -1$, $-5 + b = -1$, hence $b = 4$.

FINAL ANSWER

WMA12-01_2021_Jan_Q01

(a) **Proven:** $a + b = -1$. (b) $a = -5$, $b = 4$.

T02

02_REMA

02_REMA_CORE

CHAPTER
Polynomials

FAMILY
Remainder theorem

VARIANT
core family move

8 marks

WMA12/01 JANUARY 2022

Question 5

Past-paper question

WMA12-01_2022_Jan_Q05

5. $f(x) = 3x^3 + Ax^2 + Bx - 10$

where A and B are integers.

Given that

- when $f(x)$ is divided by $(x - 1)$ the remainder is k
- when $f(x)$ is divided by $(x + 1)$ the remainder is $-10k$
- k is a constant

(a) show that

$$11A + 9B = 83$$

(3)

Given also that $(3x - 2)$ is a factor of $f(x)$,

(b) find the value of A and the value of B .

(3)

(c) Hence find the quadratic expression $g(x)$ such that

$$f(x) = (3x - 2)g(x)$$

(2)

T02 Polynomials

02_REMA Remainder
theorem02_REMA_CORE core family
move

WORKED SOLUTION

Question 5

WMA12/01 January 2022

WMA12-01_2022_Jan_Q05

1

Use x equals one

$$f(1) = 3 + A + B - 10 = A + B - 7, \text{ so } k = A + B - 7.$$

2

Use x equals negative one

$$f(-1) = -3 + A - B - 10 = A - B - 13.$$

3

Use the second remainder relation

$$\text{Since } f(-1) = -10k, A - B - 13 = -10(A + B - 7).$$

4

Show the relation

$$\text{Rearranging gives } 11A + 9B = 83.$$

5

Use the factor

$$\text{Since } (3x - 2) \text{ is a factor, } f\left(\frac{2}{3}\right) = 0.$$

6

Form the second equation

$$3\left(\frac{2}{3}\right)^3 + A\left(\frac{2}{3}\right)^2 + B\left(\frac{2}{3}\right) - 10 = 0. \text{ Multiplying by 9 gives } 2A + 3B = 41.$$

7

Solve for A and B

$$\text{Solving } 11A + 9B = 83 \text{ with } 2A + 3B = 41 \text{ gives } A = -8 \text{ and } B = 19.$$

8

Divide by the factor

$$\text{With these values, } f(x) = 3x^3 - 8x^2 + 19x - 10 = (3x - 2)(x^2 - 2x + 5).$$

FINAL ANSWER

WMA12-01_2022_Jan_Q05

$$(a) \text{ Proven: } 11A + 9B = 83. \quad (b) A = -8, B = 19. \quad (c) g(x) = x^2 - 2x + 5.$$

T02

02_REMA

02_REMA_CORE

CHAPTER
Polynomials

FAMILY
Remainder theorem

VARIANT
core family move

3 marks

WMA12/01 JANUARY 2024

Question 1

Past-paper question

WMA12-01_2024_Jan_Q01

1. $f(x) = ax^3 + 3x^2 - 8x + 2$ where a is a constant

Given that when $f(x)$ is divided by $(x - 2)$ the remainder is 3, find the value of a .

(3)

DR ESLAM AHMED

T02 Polynomials

02_REMA Remainder
theorem02_REMA_CORE core family
move

WORKED SOLUTION

Question 1

WMA12/01 January 2024

WMA12-01_2024_Jan_Q01

1

Use the remainder theoremDividing by $(x - 2)$ means the remainder is $f(2)$.

2

Substitute x equals two

$$f(2) = a(2)^3 + 3(2)^2 - 8(2) + 2 = 8a + 12 - 16 + 2 = 8a - 2.$$

3

Use the given remainderThe remainder is 3, so $8a - 2 = 3$.

4

Solve for a

$$8a = 5, \text{ hence } a = \frac{5}{8}.$$

FINAL ANSWER

$$a = \frac{5}{8}.$$

WMA12-01_2024_Jan_Q01

T02

02_REMA

02_REMA_CORE

CHAPTER
Polynomials

FAMILY
Remainder theorem

VARIANT
core family move

9 marks

WMA12/01 MAY/JUNE 2025

Question 4

Past-paper question

WMA12-01_2025_MayJune_Q04

4: The function $f(x)$ is defined by

$$f(x) = ax^3 + bx^2 + 5x - 3$$

where a and b are constants.

Given that $(x + 3)$ is a factor of $f(x)$,

(a) show that

$$b = 3a + 2$$

(2)

Given further that when $f(x)$ is divided by $(2x - 1)$ the remainder is $\frac{7}{4}$

(b) find the value of a and the value of b .

(4)

(c) Using algebra, find the quotient and the remainder when $f(x)$ is divided by $(x - 2)$

(3)

T02 Polynomials

02_REMA Remainder
theorem02_REMA_CORE core family
move

WORKED SOLUTION

Question 4

WMA12/01 May/June 2025

WMA12-01_2025_MayJune_Q04

1

Use the factor theoremSince $x + 3$ is a factor, $f(-3) = 0$.

2

Show the relation $f(-3) = -27a + 9b - 15 - 3 = 0$, so $9b = 27a + 18$ and therefore $b = 3a + 2$.

3

Use the remainder theoremWhen dividing by $2x - 1$, substitute $x = \frac{1}{2}$. The remainder is $\frac{7}{4}$, so $f\left(\frac{1}{2}\right) = \frac{7}{4}$.

4

Form the second equation $\frac{a}{8} + \frac{b}{4} + \frac{5}{2} - 3 = \frac{7}{4}$. Multiplying by 8 gives $a + 2b - 4 = 14$, so $a + 2b = 18$.

5

Solve for a and bSubstitute $b = 3a + 2$ into $a + 2b = 18$: $a + 6a + 4 = 18$, so $a = 2$ and $b = 8$.

6

Write f of x $f(x) = 2x^3 + 8x^2 + 5x - 3$.

7

Divide by x minus two $2x^3 + 8x^2 + 5x - 3 = (x - 2)(2x^2 + 12x + 29) + 55$.

8

State the resultThe quotient is $2x^2 + 12x + 29$ and the remainder is 55.**FINAL ANSWER**

WMA12-01_2025_MayJune_Q04

(a) **Proven.** (b) $a = 2, b = 8$. (c) **Quotient** $2x^2 + 12x + 29$, **remainder** 55.

TOPIC 02 | FAMILY 04

Roots/Factors

WHAT THIS FAMILY ASKS

Recognise the roots/factors demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

1 classified questions

6 total marks

T02

02_ROOT

02_ROOT_CORE

CHAPTER
Polynomials

FAMILY
Roots/factors

VARIANT
core family move

6 marks

WMA12/01 MAY/JUNE 2023

Question 2

Past-paper question

WMA12-01_2023_MayJune_Q02

2.

In this question you must show all stages of your working.

Solutions relying on calculator technology are not acceptable.

$$f(x) = 4x^3 - 8x^2 + 5x + a$$

where a is a constant.

Given that $(2x - 3)$ is a factor of $f(x)$,

(a) use the factor theorem to show that $a = -3$

(2)

(b) Hence show that the equation $f(x) = 0$ has only one real root.

(4)

T02 Polynomials

02_ROOT Roots/factors

02_ROOT_CORE core family
move

WORKED SOLUTION

Question 2

WMA12/01 May/June 2023

WMA12-01_2023_MayJune_Q02

1

Use the factor theorem

Since $(2x - 3)$ is a factor, $x = \frac{3}{2}$ is a root, so $f\left(\frac{3}{2}\right) = 0$.

2

Substitute the root

$$4\left(\frac{3}{2}\right)^3 - 8\left(\frac{3}{2}\right)^2 + 5\left(\frac{3}{2}\right) + a = 0.$$

3

Find a

$$13.5 - 18 + 7.5 + a = 0, \text{ so } 3 + a = 0 \text{ and } a = -3.$$

4

Write the polynomial

$$\text{With } a = -3, f(x) = 4x^3 - 8x^2 + 5x - 3.$$

5

Factor using the known factor

$$4x^3 - 8x^2 + 5x - 3 = (2x - 3)(2x^2 - x + 1).$$

6

Check the quadratic

$$\text{The discriminant of } 2x^2 - x + 1 \text{ is } (-1)^2 - 4(2)(1) = 1 - 8 = -7.$$

7

Conclude about real roots

Since the quadratic has negative discriminant, it has no real roots. Therefore $f(x) = 0$ has only the one real root from $2x - 3 = 0$, namely $x = \frac{3}{2}$.

FINAL ANSWER

WMA12-01_2023_MayJune_Q02

(a) $a = -3$. (b) $f(x) = 0$ has only one real root, $x = \frac{3}{2}$.

PURE MATHEMATICS 2

ADAPTIVE ALL CLASSIFIED

Presentation Edition | Solutions

CLASSIFIED BY IDEA | ROOM TO THINK

Every demand appears clearly, with practical space for the mathematics.

Dr Eslam Ahmed

Elite Mathematics

ElitelGCSE.com | 011 2000 9622

PURE 2

PURE 2 | TOPIC 03

Circles

Your classified route

19

QUESTIONS

154

MARKS

4

FAMILIES

5

VARIANTS

DR ESLAM AHMED

CLASSIFIED BY IDEA

Move from recognition to method, then use the working space to complete the solution.

TOPIC 03 | FAMILY 01

Circle Equation

WHAT THIS FAMILY ASKS

Recognise the circle equation demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

3 classified questions

26 total marks

T03

03_CENTRE

03_CENTRE_CORE

CHAPTER
Circles

FAMILY
Circle equation

VARIANT
core family move

12 marks

WMA12/01 OCTOBER 2022

Question 9

Past-paper question

WMA12-01_2022_Oct_Q09

9.

In this question you must show detailed reasoning.

Solutions relying entirely on calculator technology are not acceptable.




Figure 3

Figure 3 shows

- the curve C_1 with equation $y = x^3 - 5x^2 + 3x + 14$
- the circle C_2 with centre T

The point T is the minimum turning point of C_1

Using Figure 3 and calculus,

- (a) find the coordinates of T

(3)

The curve C_1 intersects the circle C_2 at the point A with x coordinate 2

- (b) Find an equation of the circle C_2

(3)

The line l shown in Figure 3, is the tangent to circle C_2 at A

- (c) Show that an equation of l is

$$y = \frac{1}{3}x + \frac{22}{3}$$

(3)

The region R , shown shaded in Figure 3, is bounded by C_1 , l and the y -axis.

- (d) Find the exact area of R .

(3)

T03 Circles

03_CENTRE Circle equation

03_CENTRE_CORE core
family move

WORKED SOLUTION

Question 9

WMA12/01 October 2022

WMA12-01_2022_Oct_Q09

1

Differentiate the cubicFor $y = x^3 - 5x^2 + 3x + 14$, $\frac{dy}{dx} = 3x^2 - 10x + 3$.

2

Find stationary x values $3x^2 - 10x + 3 = 0$ factors as $(3x - 1)(x - 3) = 0$, so $x = \frac{1}{3}$ or $x = 3$.

3

Identify the minimum $\frac{d^2y}{dx^2} = 6x - 10$. At $x = 3$, this is positive, so the minimum turning point has $x = 3$.

4

Find TSubstitute $x = 3$: $y = 27 - 45 + 9 + 14 = 5$. Hence $T = (3, 5)$.

5

Find AAt $x = 2$ on C_1 , $y = 8 - 20 + 6 + 14 = 8$, so $A = (2, 8)$.

6

Find the circle radiusThe centre of C_2 is $T = (3, 5)$ and it passes through $A = (2, 8)$. Thus $r^2 = (2 - 3)^2 + (8 - 5)^2 = 10$.

7

Write the circle equationTherefore C_2 has equation $(x - 3)^2 + (y - 5)^2 = 10$.

8

Find the tangent gradientThe radius TA has gradient $\frac{8-5}{2-3} = -3$, so the tangent at A has gradient $\frac{1}{3}$.

9

Write the tangent equationUsing $A = (2, 8)$, $y - 8 = \frac{1}{3}(x - 2)$, so $y = \frac{1}{3}x + \frac{22}{3}$.

10

Set up the areaFrom $x = 0$ to $x = 2$, the cubic is above the line. The required area is $\int_0^2 \left[x^3 - 5x^2 + 3x + 14 - \left(\frac{1}{3}x + \frac{22}{3} \right) \right] dx$.

11

Simplify the integrand

The integrand is $x^3 - 5x^2 + \frac{8}{3}x + \frac{20}{3}$.

12

Integrate

$$\left[\frac{x^4}{4} - \frac{5x^3}{3} + \frac{4x^2}{3} + \frac{20x}{3} \right]_0^2 = 4 - \frac{40}{3} + \frac{16}{3} + \frac{40}{3} = \frac{28}{3}.$$

FINAL ANSWER

WMA12-01_2022_Oct_Q09

(a) $T = (3, 5)$. (b) $(x - 3)^2 + (y - 5)^2 = 10$. (c) $y = \frac{1}{3}x + \frac{22}{3}$. (d) $\frac{28}{3}$.

DR ESLAM AHMED

T03

03_CENTRE

03_CENTRE_CORE

CHAPTER
Circles

FAMILY
Circle equation

VARIANT
core family move

6 marks

WMA12/01 JANUARY 2024

Question 3

Past-paper question

WMA12-01_2024_Jan_Q03

3. The circle C

- has centre $A(3, 5)$
- passes through the point $B(8, -7)$

(a) Find an equation for C .

(3)

The points M and N lie on C such that MN is a chord of C .

Given that MN

- lies above the x -axis
- is parallel to the x -axis
- has length $4\sqrt{22}$

(b) find an equation for the line passing through points M and N .

(3)

T03 Circles

03_CENTRE Circle equation

03_CENTRE_CORE core
family move

WORKED SOLUTION

Question 3

WMA12/01 January 2024

WMA12-01_2024_Jan_Q03

1

Find the radius squared

The centre is $A(3, 5)$ and the circle passes through $B(8, -7)$, so $r^2 = (8 - 3)^2 + (-7 - 5)^2 = 25 + 144 = 169$.

2

Write the circle equation

Hence C has equation $(x - 3)^2 + (y - 5)^2 = 169$.

3

Use the horizontal chord

Since MN is parallel to the x -axis, its equation has the form $y = c$.

4

Use half the chord length

The chord length is $4\sqrt{22}$, so the half-length is $2\sqrt{22}$.

5

Relate chord length to the circle

At height $y = c$, the horizontal half-length satisfies $169 - (c - 5)^2 = (2\sqrt{22})^2 = 88$.

6

Solve for c

$(c - 5)^2 = 81$, so $c = 14$ or $c = -4$.

7

Use the position above the x -axis

The chord lies above the x -axis, so $c = 14$. Therefore the line through M and N is $y = 14$.

FINAL ANSWER

WMA12-01_2024_Jan_Q03

(a) $(x - 3)^2 + (y - 5)^2 = 169$. (b) $y = 14$.

T03

03_CENTRE

03_CENTRE_CORE

CHAPTER
Circles

FAMILY
Circle equation

VARIANT
core family move

8 marks

WMA12/01 JANUARY 2025

Question 6

Past-paper question

WMA12-01_2025_Jan_Q06

6.




Figure 2

The point $A(-1, 2)$ and the point $B(11, 6)$ both lie on a circle with centre P .

The point M is the midpoint of AB .

Given that the line l passes through M and P , as shown in Figure 2,

- (a) find an equation for l , giving your answer in the form $y = mx + c$, where m and c are constants.

(4)

Given that P has coordinates $(7, k)$, where k is a constant,

- (b) find the value of k ,

(1)

- (c) find an equation for the circle.

(3)

T03 Circles

03_CENTRE Circle equation

03_CENTRE_CORE core
family move

WORKED SOLUTION

Question 6

WMA12/01 January 2025

WMA12-01_2025_Jan_Q06

1

Find the midpointFor $A(-1, 2)$ and $B(11, 6)$, the midpoint is $M\left(\frac{-1+11}{2}, \frac{2+6}{2}\right) = (5, 4)$.

2

Find the gradient of ABThe gradient of AB is $\frac{6-2}{11-(-1)} = \frac{4}{12} = \frac{1}{3}$.

3

Use the perpendicular bisectorThe centre lies on the perpendicular bisector of the chord AB , so line l has gradient -3 and passes through $M(5, 4)$.

4

Find line l $y - 4 = -3(x - 5)$, so $y = -3x + 19$.

5

Use the centre coordinatesSince $P = (7, k)$ lies on l , $k = -3(7) + 19 = -2$.

6

Find the radius squaredThe centre is $(7, -2)$. Using point A , $r^2 = (7 - (-1))^2 + (-2 - 2)^2 = 8^2 + (-4)^2 = 80$.

7

Write the circle equationTherefore the circle is $(x - 7)^2 + (y + 2)^2 = 80$.**FINAL ANSWER**

WMA12-01_2025_Jan_Q06

(a) $y = -3x + 19$. (b) $k = -2$. (c) $(x - 7)^2 + (y + 2)^2 = 80$.

TOPIC 03 | FAMILY 02

Line-Circle Intersections

WHAT THIS FAMILY ASKS

Recognise the line-circle intersections demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 Tangency means the quadratic has one repeated root.

1 classified questions

12 total marks

T03

03_MEET

03_MEET_HV01

CHAPTER
CirclesFAMILY
Line-circle intersectionsVARIANT
Tangency means the
quadratic has one repeated
root.

12 marks

WMA12/01 MAY/JUNE 2022

Question 10

Past-paper question

WMA12-01_2022_MayJune_Q10

10. The circle C has centre $X(3, 5)$ and radius r

The line l has equation $y = 2x + k$, where k is a constant.

- (a) Show that l and C intersect when

$$5x^2 + (4k - 26)x + k^2 - 10k + 34 - r^2 = 0 \quad (3)$$

Given that l is a tangent to C ,

- (b) show that $5r^2 = (k + p)^2$, where p is a constant to be found. (3)




Figure 2

The line l

- cuts the y -axis at the point A
- touches the circle C at the point B

as shown in Figure 2.

Given that $AB = 2r$

- (c) find the value of k (6)

(Total 12 marks)

T03 Circles

03_MEET Line-circle intersections

03_MEET_HV01 Tangency means the quadratic has one repeated root.

WORKED SOLUTION

Question 10

WMA12/01 May/June 2022

WMA12-01_2022_MayJune_Q10

1

Write the circle equationThe circle has centre $(3, 5)$ and radius r , so $(x - 3)^2 + (y - 5)^2 = r^2$.

2

Substitute the lineOn the line, $y = 2x + k$. Substitution gives $(x - 3)^2 + (2x + k - 5)^2 = r^2$.

3

Expand

$$x^2 - 6x + 9 + 4x^2 + 4kx - 20x + k^2 - 10k + 25 - r^2 = 0.$$

4

Show the intersection equation

$$\text{Collecting terms gives } 5x^2 + (4k - 26)x + k^2 - 10k + 34 - r^2 = 0.$$

5

Use tangency

For a tangent, the quadratic has one repeated root, so its discriminant is zero.

6

Set the discriminant to zero

$$(4k - 26)^2 - 20(k^2 - 10k + 34 - r^2) = 0.$$

7

SimplifyThis reduces to $20r^2 - 4k^2 - 8k - 4 = 0$, hence $5r^2 = (k + 1)^2$. Therefore $p = 1$.

8

Find the tangent point x coordinate

$$\text{For the repeated root, } x_B = -\frac{4k-26}{2 \cdot 5} = \frac{13-2k}{5}.$$

9

Use distance ABThe point A is $(0, k)$ and B lies on a line of gradient 2, so $AB = \sqrt{x_B^2 + (2x_B)^2} = \sqrt{5}x_B$.

10

Use the radius relationFrom $5r^2 = (k + 1)^2$, $r = \frac{k+1}{\sqrt{5}}$ in the shown configuration.

11

Apply AB equals two r

$$\sqrt{5} \left(\frac{13-2k}{5} \right) = 2 \left(\frac{k+1}{\sqrt{5}} \right).$$

12

Solve for k

Multiplying by $\sqrt{5}$ gives $13 - 2k = 2k + 2$, so $k = \frac{11}{4}$.

FINAL ANSWER

WMA12-01_2022_MayJune_Q10

(a) **Proven.** (b) $5r^2 = (k+1)^2$, so $p = 1$. (c) $k = \frac{11}{4}$.

DR ESLAM AHMED

TOPIC 03 | FAMILY 03

Radius/Distance

WHAT THIS FAMILY ASKS

Recognise the radius/distance demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

7 classified questions

49 total marks

T03

03_RADIUS

03_RADIUS_CORE

CHAPTER
Circles

FAMILY
Radius/distance

VARIANT
core family move

8 marks

WMA12/01 JANUARY 2020

Question 6

Past-paper question

WMA12-01_2020_Jan_Q06

6. The circle C has equation

$$x^2 + y^2 + 6x - 4y - 14 = 0$$

- (a) Find

- (i) the coordinates of the centre of C ,
- (ii) the exact radius of C .

(3)

The line with equation $y = k$, where k is a constant, is a tangent to C .

- (b) Find the possible values of k .

(2)

The line with equation $y = p$, where p is a negative constant, is a chord of C .

Given that the length of this chord is 4 units,

- (c) find the value of p .

(3)

T03 Circles

03_RADIUS Radius/distance

03_RADIUS_CORE core
family move

WORKED SOLUTION

Question 6

WMA12/01 January 2020

WMA12-01_2020_Jan_Q06

1

Complete the square

$$x^2 + 6x = (x + 3)^2 - 9 \text{ and } y^2 - 4y = (y - 2)^2 - 4.$$

2

Find the centre and radius

The circle becomes $(x + 3)^2 + (y - 2)^2 = 27$. Hence the centre is $(-3, 2)$ and the radius is $\sqrt{27} = 3\sqrt{3}$.

3

Use horizontal tangents

A horizontal tangent has equation $y = k$. The possible tangent levels are one radius above and below the centre, so $k = 2 \pm 3\sqrt{3}$.

4

Use half the chord length

The chord has length 4, so half the chord is 2. If the chord is $y = p$, its perpendicular distance from the centre is $|p - 2|$.

5

Apply Pythagoras

$$(p - 2)^2 + 2^2 = (3\sqrt{3})^2 = 27, \text{ so } (p - 2)^2 = 23.$$

6

Choose the negative value

Since p is negative, $p = 2 - \sqrt{23}$.

FINAL ANSWER

WMA12-01_2020_Jan_Q06

(a)(i) $(-3, 2)$. (a)(ii) $3\sqrt{3}$. (b) $k = 2 \pm 3\sqrt{3}$. (c) $p = 2 - \sqrt{23}$.

T03

03_RADIUS

03_RADIUS_CORE

CHAPTER
Circles

FAMILY
Radius/distance

VARIANT
core family move
8 marks

WMA12/01 OCTOBER 2021

Question 6

Past-paper question

WMA12-01_2021_Oct_Q06

6. (i) The circle C_1 has equation

$$x^2 + y^2 + 10x - 12y = k \quad \text{where } k \text{ is a constant}$$

- (a) Find the coordinates of the centre of C_1 (2)

- (b) State the possible range in values for k . (2)

- (ii) The point $P(p, 0)$, the point $Q(-2, 10)$ and the point $R(8, -14)$ lie on a different circle, C_2

Given that

- p is a positive constant
- QR is a diameter of C_2

find the exact value of p .

(4)

T03 Circles

03_RADIUS Radius/distance

03_RADIUS_CORE core
family move

WORKED SOLUTION

Question 6

WMA12/01 October 2021

WMA12-01_2021_Oct_Q06

1

Complete the square

$$x^2 + 10x = (x + 5)^2 - 25 \text{ and } y^2 - 12y = (y - 6)^2 - 36.$$

2

Write centre-radius form

$$\text{The circle becomes } (x + 5)^2 + (y - 6)^2 = k + 61.$$

3

Read the centre

The centre is $(-5, 6)$.

4

State the range for k

For a circle with positive radius, $k + 61 > 0$, so $k > -61$.

5

Use the diameter property

If QR is a diameter and P lies on the circle, then $\angle QPR = 90^\circ$.

6

Use perpendicular vectors

So $(Q - P) \cdot (R - P) = 0$. With $P = (p, 0)$, $Q = (-2, 10)$, and $R = (8, -14)$, this gives $(-2 - p, 10) \cdot (8 - p, -14) = 0$.

7

Form the quadratic

$$(-2 - p)(8 - p) - 140 = 0, \text{ so } p^2 - 6p - 156 = 0.$$

8

Choose the positive value

$$p = \frac{6 \pm \sqrt{660}}{2} = 3 \pm \sqrt{165}. \text{ Since } p \text{ is positive, } p = 3 + \sqrt{165}.$$

FINAL ANSWER

WMA12-01_2021_Oct_Q06

(i)(a) $(-5, 6)$. (i)(b) $k > -61$. (ii) $p = 3 + \sqrt{165}$.

T03

03_RADIUS

03_RADIUS_CORE

CHAPTER
Circles

FAMILY
Radius/distance

VARIANT
core family move

6 marks

WMA12/01 MAY/JUNE 2024

Question 7

Past-paper question

WMA12-01_2024_MayJune_Q07

7. The circle C_1 has equation

$$x^2 + y^2 + 8x - 10y = 29$$

- (a) (i) Find the coordinates of the centre of C_1
(ii) Find the exact value of the radius of C_1

(3)

In part (b) you must show detailed reasoning.

The circle C_2 has equation

$$(x - 5)^2 + (y + 8)^2 = 52$$

- (b) Prove that the circles C_1 and C_2 neither touch nor intersect.

(3)

T03 Circles

03_RADIUS Radius/distance

03_RADIUS_CORE core
family move

WORKED SOLUTION

Question 7

WMA12/01 May/June 2024

WMA12-01_2024_MayJune_Q07

1

Complete the square $x^2 + y^2 + 8x - 10y = 29$ becomes $(x + 4)^2 - 16 + (y - 5)^2 - 25 = 29$.

2

Write circle C1 in standard form $(x + 4)^2 + (y - 5)^2 = 70$.

3

Read centre and radiusThe centre of C_1 is $(-4, 5)$ and its radius is $\sqrt{70}$.

4

Read C2 C_2 has equation $(x - 5)^2 + (y + 8)^2 = 52$, so its centre is $(5, -8)$ and its radius is $\sqrt{52} = 2\sqrt{13}$.

5

Find distance between centresThe distance between $(-4, 5)$ and $(5, -8)$ is $\sqrt{9^2 + (-13)^2} = \sqrt{250} = 5\sqrt{10}$.

6

Compare with radii $\sqrt{70} + 2\sqrt{13} = 15.58 \dots$, while $5\sqrt{10} = 15.81 \dots$

7

Conclude

The distance between centres is greater than the sum of the radii, so the circles are separate. Therefore they neither touch nor intersect.

FINAL ANSWER

WMA12-01_2024_MayJune_Q07

(a)(i) $(-4, 5)$. (a)(ii) $\sqrt{70}$. (b) **The circles neither touch nor intersect.**

T03

03_RADIUS

03_RADIUS_CORE

CHAPTER
Circles

FAMILY
Radius/distance

VARIANT
core family move
5 marks

WMA12/01 MAY/JUNE (R) 2024

Question 3

Past-paper question

WMA12-01_2024_MayJuneR_Q03

3. A circle has equation

$$x^2 + y^2 + 8x - 14y - 79 = 0$$

(a) Find

- (i) the coordinates of the centre of the circle,
- (ii) the radius of the circle.

(3)

Given that P is the point on the circle that is nearest the origin O ,

(b) find the exact length of OP

(2)

T03 Circles

03_RADIUS Radius/distance

03_RADIUS_CORE core
family move

WORKED SOLUTION

Question 3

WMA12/01 May/June (R) 2024

WMA12-01_2024_MayJuneR_Q03

1

Complete the square $x^2 + y^2 + 8x - 14y - 79 = 0$ becomes $(x + 4)^2 - 16 + (y - 7)^2 - 49 - 79 = 0$.

2

Write standard form $(x + 4)^2 + (y - 7)^2 = 144$.

3

Read centre and radiusThe centre is $(-4, 7)$ and the radius is 12.

4

Find distance from centre to originThe distance from $(-4, 7)$ to $O(0, 0)$ is $\sqrt{(-4)^2 + 7^2} = \sqrt{65}$.

5

Use origin inside the circleSince $\sqrt{65} < 12$, the origin is inside the circle.

6

Find nearest distance to the circleThe nearest point on the circle lies along the line from the centre to the origin, so $OP = 12 - \sqrt{65}$.**FINAL ANSWER**

WMA12-01_2024_MayJuneR_Q03

 $(a)(i) (-4, 7).$ $(a)(ii) 12.$ $(b) 12 - \sqrt{65}.$

T03

03_RADIUS

03_RADIUS_CORE

CHAPTER
Circles

FAMILY
Radius/distance

VARIANT
core family move

7 marks

WMA12/01 MAY/JUNE 2025

Question 2

Past-paper question

WMA12-01_2025_MayJune_Q02

2: The line joining the points $(-2, 5)$ and $(4, 15)$ is the diameter of a circle C .

(a) Find an equation for C .

(5)

(b) Hence find the exact coordinates of the point on C that is nearest the x -axis.

(2)

DR ESLAM AHMED

T03 Circles

03_RADIUS Radius/distance

03_RADIUS_CORE core
family move

WORKED SOLUTION

Question 2

WMA12/01 May/June 2025

WMA12-01_2025_MayJune_Q02

1

Find the centre

The centre is the midpoint of $(-2, 5)$ and $(4, 15)$, so it is $\left(\frac{-2+4}{2}, \frac{5+15}{2}\right) = (1, 10)$.

2

Find the radius squared

Using $(-2, 5)$, $r^2 = (-2 - 1)^2 + (5 - 10)^2 = 9 + 25 = 34$.

3

Write the circle equation

Therefore C has equation $(x - 1)^2 + (y - 10)^2 = 34$.

4

Find the nearest point to the x axis

The nearest point to the x -axis is vertically below the centre by one radius.

5

Write the coordinates

The radius is $\sqrt{34}$, so the point is $(1, 10 - \sqrt{34})$.

FINAL ANSWER

WMA12-01_2025_MayJune_Q02

(a) $(x - 1)^2 + (y - 10)^2 = 34$. (b) $(1, 10 - \sqrt{34})$.

T03

03_RADIUS

03_RADIUS_CORE

CHAPTER
CirclesFAMILY
Radius/distanceVARIANT
core family move

9 marks

WMA12/01 OCTOBER 2025

Question 6

Past-paper question

WMA12-01_2025_Oct_Q06

6.




Figure 2

The points $A(-2, 18)$ and $B(14, 6)$ lie on a circle C_1 as shown in Figure 2.

Given that AB is a diameter of the circle C_1

(a) find an equation for C_1 making your method clear.

(5)

A circle C_2 has its centre at the origin.

Given that circles C_1 and C_2 touch,

(b) find possible equations for C_2

(4)

T03 Circles

03_RADIUS Radius/distance

03_RADIUS_CORE core
family move

WORKED SOLUTION

Question 6

WMA12/01 October 2025

WMA12-01_2025_Oct_Q06

1

Find the centre of C one

Since AB is a diameter, the centre is the midpoint of $A(-2, 18)$ and $B(14, 6)$. This is $(6, 12)$.

2

Find the radius squared

Using $A(-2, 18)$, $r^2 = (-2 - 6)^2 + (18 - 12)^2 = 64 + 36 = 100$.

3

Write C one

Therefore C_1 has equation $(x - 6)^2 + (y - 12)^2 = 100$.

4

Find the centre distance

The distance from the origin to $(6, 12)$ is $\sqrt{6^2 + 12^2} = 6\sqrt{5}$.

5

Find possible radii for C two

Since C_1 has radius 10, a circle centred at the origin can touch it with radius $6\sqrt{5} - 10$ or $6\sqrt{5} + 10$.

6

Write the equations

Thus C_2 can be $x^2 + y^2 = (6\sqrt{5} - 10)^2 = 280 - 120\sqrt{5}$ or $x^2 + y^2 = (6\sqrt{5} + 10)^2 = 280 + 120\sqrt{5}$.

FINAL ANSWER

WMA12-01_2025_Oct_Q06

(a) $(x - 6)^2 + (y - 12)^2 = 100$. (b) $x^2 + y^2 = 280 - 120\sqrt{5}$ or $x^2 + y^2 = 280 + 120\sqrt{5}$.

T03

03_RADIUS

03_RADIUS_CORE

CHAPTER
Circles

FAMILY
Radius/distance

VARIANT
core family move

6 marks

WMA12/01 JANUARY 2026

Question 5

Past-paper question

WMA12-01_2026_Jan_Q05

5. The circle C_1 has equation

$$x^2 + y^2 - 6x + 5y - 41 = 0$$

(a) Find

- (i) the coordinates of the centre of C_1
- (ii) the radius of C_1

(3)

The circle C_2 has

- centre $(-k, 0)$ where k is a positive constant
- radius 5

Given that circles C_1 and C_2 touch

(b) find the exact value of k .

(3)

T03 Circles

03_RADIUS Radius/distance

03_RADIUS_CORE core
family move

WORKED SOLUTION

Question 5

WMA12/01 January 2026

WMA12-01_2026_Jan_Q05

1

Complete the square

$$x^2 - 6x = (x - 3)^2 - 9 \text{ and } y^2 + 5y = \left(y + \frac{5}{2}\right)^2 - \frac{25}{4}.$$

2

Write the circle in centre-radius form

$$\text{The equation becomes } (x - 3)^2 + \left(y + \frac{5}{2}\right)^2 = \frac{225}{4}.$$

3

Read the centre and radius

$$\text{The centre is } \left(3, -\frac{5}{2}\right) \text{ and the radius is } \sqrt{\frac{225}{4}} = \frac{15}{2}.$$

4

Use the touching condition

The centre of C_2 is $(-k, 0)$ and its radius is 5. Since the circles touch externally, the distance between centres is $\frac{15}{2} + 5 = \frac{25}{2}$.

5

Solve for k

$$(k+3)^2 + \left(\frac{5}{2}\right)^2 = \left(\frac{25}{2}\right)^2, \text{ so } (k+3)^2 = 150. \text{ Since } k > 0, k+3 = 5\sqrt{6}, \text{ hence } k = 5\sqrt{6} - 3.$$

FINAL ANSWER

WMA12-01_2026_Jan_Q05

$$(a)(i) \left(3, -\frac{5}{2}\right). \quad (a)(ii) \frac{15}{2}. \quad (b) k = 5\sqrt{6} - 3.$$

TOPIC 03 | FAMILY 04

Tangents

WHAT THIS FAMILY ASKS

Recognise the tangents demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

V02 If the radius is vertical, the tangent is horizontal.

8 classified questions

67 total marks

T03

03_TANG

03_TANG_CORE

CHAPTER
CirclesFAMILY
TangentsVARIANT
core family move

7 marks

WMA12/01 OCTOBER 2019

Question 6

Past-paper question

WMA12-01_2019_Oct_Q06

6.




Figure 1

Figure 1 shows a sketch of a circle C with centre $N(4, -1)$.

The line l with equation $y = \frac{1}{2}x$ is a tangent to C at the point P .

Find

(a) the equation of line PN in the form $y = mx + c$, where m and c are constants, (2)

(b) the equation of C . (5)

T03 Circles

03_TANG Tangents

03_TANG_CORE core family
move

WORKED SOLUTION

Question 6

WMA12/01 October 2019

WMA12-01_2019_Oct_Q06

1

Use perpendicular gradients

The tangent has gradient $\frac{1}{2}$, so the radius PN has gradient -2 .

2

Find the equation of PN

Line PN passes through $N(4, -1)$, so $y + 1 = -2(x - 4)$. Therefore $y = -2x + 7$.

3

Find point P

At P , the tangent $y = \frac{1}{2}x$ meets the radius $y = -2x + 7$. Thus $\frac{1}{2}x = -2x + 7$, so $x = \frac{14}{5}$ and $y = \frac{7}{5}$.

4

Find the radius squared

$$r^2 = \left(\frac{14}{5} - 4\right)^2 + \left(\frac{7}{5} + 1\right)^2 = \left(-\frac{6}{5}\right)^2 + \left(\frac{12}{5}\right)^2 = \frac{36}{5}.$$

5

Write the circle equation

With centre $(4, -1)$ and radius squared $\frac{36}{5}$, the circle is $(x - 4)^2 + (y + 1)^2 = \frac{36}{5}$.

FINAL ANSWER

WMA12-01_2019_Oct_Q06

$$(a) y = -2x + 7. \quad (b) (x - 4)^2 + (y + 1)^2 = \frac{36}{5}.$$

T03

03_TANG

03_TANG_CORE

CHAPTER
Circles

FAMILY
Tangents

VARIANT
core family move

10 marks

WMA12/01 JANUARY 2021

Question 9

Past-paper question

WMA12-01_2021_Jan_Q09

9. A circle C has equation

$$(x - k)^2 + (y - 2k)^2 = k + 7$$

where k is a positive constant.

- (a) Write down, in terms of k ,

- (i) the coordinates of the centre of C ,
- (ii) the radius of C .

(2)

Given that the point $P(2, 3)$ lies on C

- (b) (i) show that $5k^2 - 17k + 6 = 0$

- (ii) hence find the possible values of k .

(3)

The tangent to the circle at P intersects the x -axis at point T .

Given that $k < 2$

- (c) calculate the exact area of triangle OPT .

(5)

T03 Circles

03_TANG Tangents

03_TANG_CORE core family
move

WORKED SOLUTION

Question 9

WMA12/01 January 2021

WMA12-01_2021_Jan_Q09

1

Read the centre and radius

From $(x - k)^2 + (y - 2k)^2 = k + 7$, the centre is $(k, 2k)$ and the radius is $\sqrt{k + 7}$.

2

Substitute point P

Since $P(2, 3)$ lies on the circle, $(2 - k)^2 + (3 - 2k)^2 = k + 7$.

3

Show the quadratic

Expanding gives $k^2 - 4k + 4 + 4k^2 - 12k + 9 = k + 7$, hence $5k^2 - 17k + 6 = 0$.

4

Solve for k

$5k^2 - 17k + 6 = (5k - 2)(k - 3)$, so $k = \frac{2}{5}$ or $k = 3$.

5

Use the condition k less than two

For the tangent calculation, $k < 2$, so the centre is $(\frac{2}{5}, \frac{4}{5})$.

6

Find the tangent gradient

The radius from the centre to P has gradient $\frac{3 - 4/5}{2 - 2/5} = \frac{11}{8}$, so the tangent gradient is $-\frac{8}{11}$.

7

Find the x intercept

The tangent is $y - 3 = -\frac{8}{11}(x - 2)$. Setting $y = 0$ gives $x = \frac{49}{8}$, so $T = (\frac{49}{8}, 0)$.

8

Calculate the area

Triangle OPT has base $\frac{49}{8}$ on the x -axis and height 3, so its area is $\frac{1}{2} \cdot \frac{49}{8} \cdot 3 = \frac{147}{16}$.

FINAL ANSWER

WMA12-01_2021_Jan_Q09

(a)(i) $(k, 2k)$. (a)(ii) $\sqrt{k + 7}$. (b)(ii) $k = 3$ or $\frac{2}{5}$. (c) $\frac{147}{16}$.

T03

03_TANG

03_TANG_CORE

CHAPTER
CirclesFAMILY
TangentsVARIANT
core family move

8 marks

WMA12/01 JANUARY 2022

Question 6

Past-paper question

WMA12-01_2022_Jan_Q06

6.




Figure 1

The points $P(23, 14)$, $Q(15, -30)$ and $R(-7, -26)$ lie on the circle C , as shown in Figure 1.

(a) Show that $\angle PQR = 90^\circ$ (2)

(b) Hence, or otherwise, find

(i) the centre of C ,

(ii) the radius of C . (3)

Given that the point S lies on C such that the distance QS is greatest,

(c) find an equation of the tangent to C at S , giving your answer in the form $ax + by + c = 0$, where a , b and c are integers to be found. (3)

T03 Circles

03_TANG Tangents

03_TANG_CORE core family
move

WORKED SOLUTION

Question 6

WMA12/01 January 2022

WMA12-01_2022_Jan_Q06

1

Find the gradient QPThe gradient of QP is $\frac{14 - (-30)}{23 - 15} = \frac{44}{8} = \frac{11}{2}$.

2

Find the gradient QRThe gradient of QR is $\frac{-26 - (-30)}{-7 - 15} = \frac{4}{-22} = -\frac{2}{11}$.

3

Show the right angleThe product of the two gradients is -1 , so QP is perpendicular to QR and $\angle PQR = 90^\circ$.

4

Use the diameterSince the angle at Q is a right angle, PR is a diameter of the circle.

5

Find the centreThe centre is the midpoint of $P(23, 14)$ and $R(-7, -26)$, so it is $(8, -6)$.

6

Find the radius $PR = \sqrt{30^2 + 40^2} = 50$, so the radius is 25.

7

Find the farthest point from QThe greatest distance from Q occurs at the point opposite Q on the diameter through the centre. Thus $S = 2(8, -6) - (15, -30) = (1, 18)$.

8

Find the tangent gradientThe radius from the centre to S has gradient $\frac{18 - (-6)}{1 - 8} = -\frac{24}{7}$, so the tangent gradient is $\frac{7}{24}$.

9

Write the tangent equation $y - 18 = \frac{7}{24}(x - 1)$, which rearranges to $7x - 24y + 425 = 0$.**FINAL ANSWER**

WMA12-01_2022_Jan_Q06

(a) **Proven:** $\angle PQR = 90^\circ$. (b)(i) $(8, -6)$. (b)(ii) 25. (c) $7x - 24y + 425 = 0$.

T03

03_TANG

03_TANG_CORE

CHAPTER
Circles

FAMILY
Tangents

VARIANT
core family move

10 marks

WMA12/01 JANUARY 2023

Question 6

Past-paper question

WMA12-01_2023_Jan_Q06

6. The circle C has equation

$$x^2 + y^2 + 8x - 4y = 0$$

- (a) Find

- (i) the coordinates of the centre of C ,
- (ii) the exact radius of C .

(3)

The point P lies on C .

Given that the tangent to C at P has equation $x + 2y + 10 = 0$

- (b) find the coordinates of P

(4)

- (c) Find the equation of the normal to C at P , giving your answer in the form $y = mx + c$ where m and c are integers to be found.

(3)

T03 Circles

03_TANG Tangents

03_TANG_CORE core family
move

WORKED SOLUTION

Question 6

WMA12/01 January 2023

WMA12-01_2023_Jan_Q06

1

Complete the square $x^2 + y^2 + 8x - 4y = 0$ becomes $(x + 4)^2 - 16 + (y - 2)^2 - 4 = 0$.

2

Write the circle formSo $(x + 4)^2 + (y - 2)^2 = 20$.

3

Read the centre and radiusThe centre is $(-4, 2)$ and the radius is $\sqrt{20} = 2\sqrt{5}$.

4

Use the tangent gradientThe tangent $x + 2y + 10 = 0$ has gradient $-\frac{1}{2}$.

5

Find the radius gradient

The radius to the point of contact is perpendicular to the tangent, so its gradient is 2.

6

Write the normal through the centreThe radius line through the centre $(-4, 2)$ is $y - 2 = 2(x + 4)$, so $y = 2x + 10$.

7

Intersect with the tangentSubstitute $y = 2x + 10$ into $x + 2y + 10 = 0$: $x + 2(2x + 10) + 10 = 0$.

8

Find P $5x + 30 = 0$, so $x = -6$. Then $y = 2(-6) + 10 = -2$, hence $P = (-6, -2)$.

9

State the normalThe normal at P is the radius line already found, so its equation is $y = 2x + 10$.**FINAL ANSWER**

WMA12-01_2023_Jan_Q06

(a)(i) $(-4, 2)$. (a)(ii) $2\sqrt{5}$. (b) $P = (-6, -2)$. (c) $y = 2x + 10$.

T03

03_TANG

03_TANG_CORE

CHAPTER
Circles

FAMILY
Tangents

VARIANT
core family move

7 marks

WMA12/01 MAY/JUNE 2023

Question 3

Past-paper question

WMA12-01_2023_MayJune_Q03

3. A circle C has centre $(2, 5)$

Given that the point $P(8, -3)$ lies on C

- (a) (i) find the radius of C

- (ii) find an equation for C

(3)

- (b) Find the equation of the tangent to C at P giving your answer in the form $ax + by + c = 0$ where a , b and c are integers to be found.

(4)

T03 Circles

03_TANG Tangents

03_TANG_CORE core family
move

WORKED SOLUTION

Question 3

WMA12/01 May/June 2023

WMA12-01_2023_MayJune_Q03

1

Use the distance formulaThe centre is $(2, 5)$ and $P = (8, -3)$. The radius is $\sqrt{(8-2)^2 + (-3-5)^2}$.

2

Find the radius

$$r = \sqrt{6^2 + (-8)^2} = \sqrt{100} = 10.$$

3

Write the circle equationWith centre $(2, 5)$ and radius 10, the circle is $(x-2)^2 + (y-5)^2 = 100$.

4

Find the radius gradientThe gradient of the radius from the centre to P is $\frac{-3-5}{8-2} = \frac{-8}{6} = -\frac{4}{3}$.

5

Find the tangent gradientThe tangent is perpendicular to the radius, so its gradient is $\frac{3}{4}$.

6

Use point PThe tangent through $(8, -3)$ is $y + 3 = \frac{3}{4}(x - 8)$.

7

Put into integer formMultiplying by 4 gives $4y + 12 = 3x - 24$, hence $3x - 4y - 36 = 0$.**FINAL ANSWER**

WMA12-01_2023_MayJune_Q03

 $(a)(i)$ 10. $(a)(ii)$ $(x-2)^2 + (y-5)^2 = 100$. (b) $3x - 4y - 36 = 0$.

T03

03_TANG

03_TANG_CORE

CHAPTER
CirclesFAMILY
TangentsVARIANT
core family move

8 marks

WMA12/01 OCTOBER 2023

Question 7

Past-paper question

WMA12-01_2023_Oct_Q07

7.




Figure 2

Figure 2 shows a sketch of

- the circle C with centre $X(4, -3)$
- the line l with equation $y = \frac{5}{2}x - \frac{55}{2}$

Given that l is the tangent to C at the point N ,(a) show that an equation for the straight line passing through X and N is

$$2x + 5y + 7 = 0$$

(3)

(b) Hence find

- the coordinates of N ,
- an equation for C .

(5)

T03 Circles

03_TANG Tangents

03_TANG_CORE core family
move

WORKED SOLUTION

Question 7

WMA12/01 October 2023

WMA12-01_2023_Oct_Q07

1

Use perpendicular gradients

The tangent l has gradient $\frac{5}{2}$, so the radius XN is perpendicular to it and has gradient $-\frac{2}{5}$.

2

Use the centre

The line through $X(4, -3)$ with gradient $-\frac{2}{5}$ is $y + 3 = -\frac{2}{5}(x - 4)$.

3

Show the required line

Multiplying by 5 gives $5y + 15 = -2x + 8$, so $2x + 5y + 7 = 0$.

4

Intersect the radius and tangent

Use $y = \frac{5}{2}x - \frac{55}{2}$ in $2x + 5y + 7 = 0$.

5

Find x coordinate

$2x + 5\left(\frac{5}{2}x - \frac{55}{2}\right) + 7 = 0$. Multiplying by 2 gives $4x + 25x - 275 + 14 = 0$, so $29x = 261$ and $x = 9$.

6

Find y coordinate

$y = \frac{5}{2}(9) - \frac{55}{2} = \frac{45-55}{2} = -5$, so $N = (9, -5)$.

7

Find the radius squared

$XN^2 = (9 - 4)^2 + (-5 + 3)^2 = 5^2 + (-2)^2 = 29$.

8

Write the circle equation

With centre $(4, -3)$ and radius squared 29, the circle is $(x - 4)^2 + (y + 3)^2 = 29$.

FINAL ANSWER

WMA12-01_2023_Oct_Q07

(a) **Proven.** (b)(i) $N = (9, -5)$. (b)(ii) $(x - 4)^2 + (y + 3)^2 = 29$.

T03

03_TANG

03_TANG_CORE

CHAPTER
Circles

FAMILY
Tangents

VARIANT
core family move

10 marks

WMA12/01 OCTOBER 2024

Question 10

Past-paper question

WMA12-01_2024_Oct_Q10

10. The circle C has equation

$$x^2 + y^2 + 4x - 30y + 209 = 0$$

(a) Find

- (i) the coordinates of the centre of C ,
- (ii) the exact value of the radius of C .

(3)

The line L has equation $y = mx + 1$, where m is a constant.

Given that L is the tangent to C at the point P ,

(b) show that

$$2m^2 - 7m - 22 = 0$$

(3)

(c) Hence find the possible pairs of coordinates of P .

(4)

T03 Circles

03_TANG Tangents

03_TANG_CORE core family
move

WORKED SOLUTION

Question 10

WMA12/01 October 2024

WMA12-01_2024_Oct_Q10

1

Complete the square $x^2 + y^2 + 4x - 30y + 209 = 0$ becomes $(x + 2)^2 + (y - 15)^2 = 20$.

2

Read the centre and radiusThe centre is $(-2, 15)$ and the radius is $\sqrt{20} = 2\sqrt{5}$.

3

Use the tangent distanceThe line $y = mx + 1$ can be written as $mx - y + 1 = 0$. Its distance from $(-2, 15)$ must be $2\sqrt{5}$.

4

Set up the equation for m $\frac{|m(-2) - 15 + 1|}{\sqrt{m^2 + 1}} = 2\sqrt{5}$, so $\frac{|-2m - 14|}{\sqrt{m^2 + 1}} = 2\sqrt{5}$.

5

Show the quadraticSquaring gives $4(m+7)^2 = 20(m^2+1)$. Hence $(m+7)^2 = 5(m^2+1)$, so $2m^2 - 7m - 22 = 0$.

6

Find the possible gradients $2m^2 - 7m - 22 = 0$ gives $m = -2$ or $m = \frac{11}{2}$.

7

Use perpendicular radii

A tangent is perpendicular to the radius at the point of contact.

8

Find the first pointFor $m = -2$, the radius gradient is $\frac{1}{2}$. The radius through the centre is $y - 15 = \frac{1}{2}(x + 2)$. Solving with $y = -2x + 1$ gives $x = -6$ and $y = 13$.

9

Find the second pointFor $m = \frac{11}{2}$, the radius gradient is $-\frac{2}{11}$. The radius through the centre is $y - 15 = -\frac{2}{11}(x + 2)$. Solving with $y = \frac{11}{2}x + 1$ gives $x = \frac{12}{5}$ and $y = \frac{71}{5}$.

FINAL ANSWER

WMA12-01_2024_Oct_Q10

(a)(i) $(-2, 15)$. (a)(ii) $2\sqrt{5}$. (b) **Proven.** (c) $(-6, 13)$ and $(\frac{12}{5}, \frac{71}{5})$.

DR ESLAM AHMED

T03

03_TANG

03_TANG_HV01

CHAPTER
CirclesFAMILY
Tangents

VARIANT

If the radius is vertical, the
tangent is horizontal.

7 marks

WMA12/01 MAY/JUNE 2021

Question 6

Past-paper question

WMA12-01_2021_MayJune_Q06

6. A circle has equation

$$x^2 - 6x + y^2 + 8y + k = 0$$

where k is a positive constant.

Given that the x -axis is a tangent to this circle,

- (a) find the value of k .

(3)

The circle meets the coordinate axes at the points R , S and T .

- (b) Find the exact area of the triangle RST .

(4)

T03 Circles

03_TANG Tangents

03_TANG_HV01 If the radius is vertical, the tangent is horizontal.

WORKED SOLUTION

Question 6

WMA12/01 May/June 2021

WMA12-01_2021_MayJune_Q06

1

Complete the square

$$x^2 - 6x = (x - 3)^2 - 9 \text{ and } y^2 + 8y = (y + 4)^2 - 16.$$

2

Write centre-radius form

$$\text{The circle becomes } (x - 3)^2 + (y + 4)^2 = 25 - k.$$

3

Use the tangent condition

The centre is $(3, -4)$. Since the x -axis is tangent, the radius is the distance from the centre to the x -axis, which is 4.

4

Find k

$$\text{Thus } 25 - k = 4^2 = 16, \text{ so } k = 9.$$

5

Find the intercepts

With $k = 9$, the circle is $(x - 3)^2 + (y + 4)^2 = 16$. On the x -axis, $y = 0$, giving $x = 3$.

6

Find the y -axis points

$$\text{On the } y\text{-axis, } x = 0, \text{ so } 9 + (y + 4)^2 = 16. \text{ Hence } y = -4 \pm \sqrt{7}.$$

7

Calculate the triangle area

The vertical base on the y -axis has length $2\sqrt{7}$, and its perpendicular distance to $(3, 0)$ is 3. The area is $\frac{1}{2}(2\sqrt{7})(3) = 3\sqrt{7}$.

FINAL ANSWER

$$(a) k = 9. \quad (b) 3\sqrt{7}.$$

WMA12-01_2021_MayJune_Q06

PURE MATHEMATICS 2

ADAPTIVE ALL CLASSIFIED

Presentation Edition | Solutions

CLASSIFIED BY IDEA | ROOM TO THINK

Every demand appears clearly, with practical space for the mathematics.

Dr Eslam Ahmed

Elite Mathematics

ElitelGCSE.com | 011 2000 9622

PURE 2

PURE 2 | TOPIC 04

Binomial Expansion

Your classified route

19

QUESTIONS

124

MARKS

3

FAMILIES

3

VARIANTS

DR ESLAM AHMED

CLASSIFIED BY IDEA

Move from recognition to method, then use the working space to complete the solution.

TOPIC 04 | FAMILY 01

Approximation

WHAT THIS FAMILY ASKS

Recognise the approximation demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

1 classified questions

8 total marks

T04

04_APPROX

04_APPROX_CORE

CHAPTER
Binomial Expansion

FAMILY
Approximation

VARIANT
core family move

8 marks

WMA12/01 MAY/JUNE 2021

Question 4

Past-paper question

WMA12-01_2021_MayJune_Q04

4. (a) Find, in ascending powers of x , up to and including the term in x^3 , the binomial expansion of

$$\left(2 + \frac{x}{8}\right)^{13}$$

fully simplifying each coefficient.

(4)

- (b) Use the answer to part (a) to find an approximation for 2.0125^{13}

Give your answer to 3 decimal places.

(3)

Without calculating 2.0125^{13}

- (c) state, with a reason, whether the answer to part (b) is an overestimate or an underestimate.

(1)

T04 Binomial Expansion

04_APPROX Approximation

04_APPROX_CORE core
family move

WORKED SOLUTION

Question 4

WMA12/01 May/June 2021

WMA12-01_2021_MayJune_Q04

1

Write the binomial terms

$$(2 + \frac{x}{8})^{13} = 2^{13} + \binom{13}{1} 2^{12} \frac{x}{8} + \binom{13}{2} 2^{11} (\frac{x}{8})^2 + \binom{13}{3} 2^{10} (\frac{x}{8})^3 + \dots$$

2

Simplify the expansion

This gives $8192 + 6656x + 2496x^2 + 572x^3 + \dots$.

3

Match the value

Since $2.0125 = 2 + 0.0125 = 2 + \frac{0.1}{8}$, use $x = 0.1$ in the expansion.

4

Calculate the approximation

$$8192 + 6656(0.1) + 2496(0.1)^2 + 572(0.1)^3 = 8883.132.$$

5

Decide the direction

The next terms in the expansion are positive because $x = 0.1 > 0$, so stopping after the x^3 term gives an underestimate.

FINAL ANSWER

WMA12-01_2021_MayJune_Q04

(a) $8192 + 6656x + 2496x^2 + 572x^3$. (b) 8883.132. (c) **Underestimate.**

TOPIC 04 | FAMILY 02

Coefficients

WHAT THIS FAMILY ASKS

Recognise the coefficients demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

8 classified questions

50 total marks

T04

04_COEFF

04_COEFF_CORE

CHAPTER
Binomial Expansion

FAMILY
Coefficients

VARIANT
core family move

8 marks

WMA12/01 JANUARY 2021

Question 4

Past-paper question

WMA12-01_2021_Jan_Q04

4. (a) Find the first three terms, in ascending powers of x , of the binomial expansion of

$$(2 + px)^6$$

where p is a constant. Give each term in simplest form.

(4)

Given that in the expansion of

$$\left(3 - \frac{1}{2}x\right)(2 + px)^6$$

the coefficient of x^2 is $-\frac{3}{4}$

- (b) find the possible values of p .

(4)

T04 Binomial Expansion

04_COEFF Coefficients

04_COEFF_CORE core family
move

WORKED SOLUTION

Question 4

WMA12/01 January 2021

WMA12-01_2021_Jan_Q04

1

Write the first binomial terms

$$(2 + px)^6 = 2^6 + \binom{6}{1}2^5(px) + \binom{6}{2}2^4(px)^2 + \dots$$

2

Simplify the expansion

The first three terms are $64 + 192px + 240p^2x^2$.

3

Find the coefficient of x squared

In $(3 - \frac{1}{2}x)(2 + px)^6$, the x^2 coefficient is $3(240p^2) - \frac{1}{2}(192p)$.

4

Form the equation

$$\text{So } 720p^2 - 96p = -\frac{3}{4}.$$

5

Solve the quadratic

Multiplying by 4 gives $2880p^2 - 384p + 3 = 0$, or $960p^2 - 128p + 1 = 0$.

6

State the values of p

$$p = \frac{128 \pm 112}{1920}, \text{ so } p = \frac{1}{8} \text{ or } p = \frac{1}{120}.$$

FINAL ANSWER

WMA12-01_2021_Jan_Q04

(a) $64 + 192px + 240p^2x^2$. (b) $p = \frac{1}{8}$ or $p = \frac{1}{120}$.

T04

04_COEFF

04_COEFF_CORE

CHAPTER
Binomial Expansion

FAMILY
Coefficients

VARIANT
core family move

6 marks

WMA12/01 OCTOBER 2021

Question 1

Past-paper question

WMA12-01_2021_Oct_Q01

1. The first three terms, in ascending powers of x , of the binomial expansion of $(1 + kx)^{16}$ are

$$1, -4x \text{ and } px^2$$

where k and p are constants.

- (a) Find, in simplest form,

(i) the value of k

(ii) the value of p

(3)

$$g(x) = \left(2 + \frac{16}{x}\right)(1 + kx)^{16}$$

Using the value of k found in part (a),

- (b) find the term in x^2 in the expansion of $g(x)$.

(3)

T04 Binomial Expansion

04_COEFF Coefficients

04_COEFF_CORE core family
move

WORKED SOLUTION

Question 1

WMA12/01 October 2021

WMA12-01_2021_Oct_Q01

1

Compare the linear termIn $(1 + kx)^{16}$, the coefficient of x is $16k$.

2

Find k Since the coefficient of x is -4 , $16k = -4$, so $k = -\frac{1}{4}$.

3

Find p The coefficient of x^2 is $\binom{16}{2}k^2 = 120\left(\frac{1}{16}\right) = \frac{15}{2}$.

4

Find the needed binomial termsFor $k = -\frac{1}{4}$, the x^2 coefficient in $(1 + kx)^{16}$ is $\frac{15}{2}$ and the x^3 coefficient is $\binom{16}{3}\left(-\frac{1}{4}\right)^3 = -\frac{35}{4}$.

5

Build the x squared term in g In $g(x) = \left(2 + \frac{16}{x}\right)(1 + kx)^{16}$, the x^2 term comes from $2\left(\frac{15}{2}x^2\right)$ and $\frac{16}{x}\left(-\frac{35}{4}x^3\right)$.

6

SimplifyThe required term is $15x^2 - 140x^2 = -125x^2$.**FINAL ANSWER**

WMA12-01_2021_Oct_Q01

 $(a)(i) k = -\frac{1}{4}$. $(a)(ii) p = \frac{15}{2}$. $(b) -125x^2$.

T04

04_COEFF

04_COEFF_CORE

CHAPTER
Binomial Expansion

FAMILY
Coefficients

VARIANT
core family move

7 marks

WMA12/01 JANUARY 2022

Question 3

Past-paper question

WMA12-01_2022_Jan_Q03

3. (a) Find the first 4 terms, in ascending powers of x , of the binomial expansion of

$$\left(2 - \frac{kx}{4}\right)^8$$

where k is a non-zero constant. Give each term in simplest form.

(4)

$$f(x) = (5 - 3x)\left(2 - \frac{kx}{4}\right)^8$$

In the expansion of $f(x)$, the constant term is 3 times the coefficient of x .

- (b) Find the value of k .

(3)

T04 Binomial Expansion

04_COEFF Coefficients

04_COEFF_CORE core family
move

WORKED SOLUTION

Question 3

WMA12/01 January 2022

WMA12-01_2022_Jan_Q03

1

Write the first terms

$$\left(2 - \frac{kx}{4}\right)^8 = 2^8 + \binom{8}{1}2^7\left(-\frac{kx}{4}\right) + \binom{8}{2}2^6\left(-\frac{kx}{4}\right)^2 + \binom{8}{3}2^5\left(-\frac{kx}{4}\right)^3 + \dots$$

2

Simplify the expansion

The first four terms are $256 - 256kx + 112k^2x^2 - 28k^3x^3$.

3

Find the constant term in f

For $f(x) = (5 - 3x)\left(2 - \frac{kx}{4}\right)^8$, the constant term is $5(256) = 1280$.

4

Find the coefficient of x

The coefficient of x is $5(-256k) - 3(256) = -1280k - 768$.

5

Use the condition

The constant term is 3 times the coefficient of x , so $1280 = 3(-1280k - 768)$.

6

Solve for k

$$1280 = -3840k - 2304, \text{ hence } k = -\frac{3584}{3840} = -\frac{14}{15}.$$

FINAL ANSWER

WMA12-01_2022_Jan_Q03

$$(a) 256 - 256kx + 112k^2x^2 - 28k^3x^3. \quad (b) k = -\frac{14}{15}.$$

T04

04_COEFF

04_COEFF_CORE

CHAPTER

Binomial Expansion

FAMILY

Coefficients

VARIANT

core family move

7 marks

WMA12/01 OCTOBER 2022

Question 2

Past-paper question

WMA12-01_2022_Oct_Q02

2. A curve C has equation $y = f(x)$ where

$$f(x) = (2 - kx)^5$$

and k is a constant.

Given that when $f(x)$ is divided by $(4x - 5)$ the remainder is $\frac{243}{32}$

- (a) show that $k = \frac{2}{5}$

(2)

- (b) Find the first three terms, in ascending powers of x , of the binomial expansion of

$$\left(2 - \frac{2}{5}x\right)^5$$

giving each term in simplest form.

(3)

Using the solution to part (b) and making your method clear,

- (c) find the gradient of C at the point where $x = 0$

(2)

T04 Binomial Expansion

04_COEFF Coefficients

04_COEFF_CORE core family
move

WORKED SOLUTION

Question 2

WMA12/01 October 2022

WMA12-01_2022_Oct_Q02

1

Use the remainder theoremDividing by $4x - 5$ means the remainder is $f\left(\frac{5}{4}\right)$.

2

Substitute x equals five quarters

$$f\left(\frac{5}{4}\right) = \left(2 - \frac{5k}{4}\right)^5 = \frac{243}{32}.$$

3

Take the fifth rootSince $\frac{243}{32} = \left(\frac{3}{2}\right)^5$, we have $2 - \frac{5k}{4} = \frac{3}{2}$.

4

Find k

$$\frac{5k}{4} = \frac{1}{2}, \text{ so } k = \frac{2}{5}.$$

5

Expand the binomialWith $k = \frac{2}{5}$, $f(x) = \left(2 - \frac{2}{5}x\right)^5$.

6

Find the first three terms

$$2^5 + 5(2^4)\left(-\frac{2}{5}x\right) + 10(2^3)\left(-\frac{2}{5}x\right)^2 = 32 - 32x + \frac{64}{5}x^2.$$

7

Use the coefficient of x For a polynomial, the gradient at $x = 0$ is the coefficient of x in the expansion. Therefore $f'(0) = -32$.**FINAL ANSWER**

WMA12-01_2022_Oct_Q02

$$(a) k = \frac{2}{5}. \quad (b) 32 - 32x + \frac{64}{5}x^2. \quad (c) -32.$$

T04

04_COEFF

04_COEFF_CORE

CHAPTER
Binomial Expansion

FAMILY
Coefficients

VARIANT
core family move

7 marks

WMA12/01 MAY/JUNE 2023

Question 4

Past-paper question

WMA12-01_2023_MayJune_Q04

4. The binomial expansion, in ascending powers of x , of

$$(3 + px)^5$$

where p is a constant, can be written in the form

$$A + Bx + Cx^2 + Dx^3 \dots$$

where A , B , C and D are constants.

- (a) Find the value of A

(1)

Given that

- $B = 18D$
- $p < 0$

- (b) find

- (i) the value of p
- (ii) the value of C

(6)

T04 Binomial Expansion

04_COEFF Coefficients

04_COEFF_CORE core family
move

WORKED SOLUTION

Question 4

WMA12/01 May/June 2023

WMA12-01_2023_MayJune_Q04

1

Find the constant termIn $(3 + px)^5$, the constant term is $3^5 = 243$, so $A = 243$.

2

Find BThe coefficient of x is $5(3^4)p = 405p$, so $B = 405p$.

3

Find DThe coefficient of x^3 is $\binom{5}{3}(3^2)p^3 = 10 \cdot 9p^3 = 90p^3$, so $D = 90p^3$.

4

Use B equals 18D $405p = 18(90p^3) = 1620p^3$.

5

Solve for pSince $p < 0$, $p \neq 0$. Divide by $405p$ to get $1 = 4p^2$, so $p = \pm \frac{1}{2}$. Therefore $p = -\frac{1}{2}$.

6

Find CThe coefficient of x^2 is $\binom{5}{2}(3^3)p^2 = 10 \cdot 27 \cdot \frac{1}{4} = \frac{135}{2}$.**FINAL ANSWER**

WMA12-01_2023_MayJune_Q04

 $(a) A = 243.$ $(b)(i) p = -\frac{1}{2}.$ $(b)(ii) C = \frac{135}{2}.$

T04

04_COEFF

04_COEFF_CORE

CHAPTER
Binomial Expansion

FAMILY
Coefficients

VARIANT
core family move

3 marks

WMA12/01 JANUARY 2024

Question 2

Past-paper question

WMA12-01_2024_Jan_Q02

2. Find the coefficient of the term in x^7 of the binomial expansion of

$$\left(\frac{3}{8} + 4x\right)^{12}$$

giving your answer in simplest form.

(3)

T04 Binomial Expansion

04_COEFF Coefficients

04_COEFF_CORE core family
move

WORKED SOLUTION

Question 2

WMA12/01 January 2024

WMA12-01_2024_Jan_Q02

1

Choose the x power

In $\left(\frac{3}{8} + 4x\right)^{12}$, the term in x^7 comes from choosing $(4x)^7$ and $\left(\frac{3}{8}\right)^5$.

2

Write the term

The coefficient is $\binom{12}{7} \left(\frac{3}{8}\right)^5 4^7$.

3

Simplify the powers

$4^7 = 2^{14}$ and $8^5 = 2^{15}$, so $\left(\frac{3}{8}\right)^5 4^7 = \frac{3^5}{2}$.

4

Evaluate

$\binom{12}{7} = 792$, so the coefficient is $792 \cdot \frac{243}{2} = 96228$.

FINAL ANSWER

96228.

WMA12-01_2024_Jan_Q02

T04

04_COEFF

04_COEFF_CORE

CHAPTER
Binomial Expansion

FAMILY
Coefficients

VARIANT
core family move

6 marks

WMA12/01 OCTOBER 2025

Question 2

Past-paper question

WMA12-01_2025_Oct_Q02

2. The first 4 terms, in ascending powers of x , in the binomial expansion of

$$(1 + px)^{10}$$

are

$$1 + 15x + qx^2 + rx^3$$

where p , q and r are constants.

Find the value of p , the value of q and the value of r .

(6)

T04 Binomial Expansion

04_COEFF Coefficients

04_COEFF_CORE core family
move

WORKED SOLUTION

Question 2

WMA12/01 October 2025

WMA12-01_2025_Oct_Q02

1

Write the first four terms

$$(1 + px)^{10} = 1 + 10px + \binom{10}{2}p^2x^2 + \binom{10}{3}p^3x^3 + \dots$$

2

Compare the x coefficient

Since the coefficient of x is 15, $10p = 15$, so $p = \frac{3}{2}$.

3

Find q

$$q = \binom{10}{2}p^2 = 45 \left(\frac{3}{2}\right)^2 = \frac{405}{4}.$$

4

Find r

$$r = \binom{10}{3}p^3 = 120 \left(\frac{3}{2}\right)^3 = 405.$$

FINAL ANSWER

$$p = \frac{3}{2}, q = \frac{405}{4}, r = 405.$$

WMA12-01_2025_Oct_Q02

T04

04_COEFF

04_COEFF_CORE

CHAPTER
Binomial Expansion

FAMILY
Coefficients

VARIANT
core family move

6 marks

WMA12/01 JANUARY 2026

Question 1

Past-paper question

WMA12-01_2026_Jan_Q01

1. (a) Find the first 4 terms, in ascending powers of x , of the binomial expansion of

$$(3 + kx)^7$$

where k is a non-zero constant. Give each term in simplest form.

(4)

Given that, in this expansion, the coefficient of x^2 is 3 times the coefficient of x^3

- (b) find, in simplest form, the value of k .

(2)

T04 Binomial Expansion

04_COEFF Coefficients

04_COEFF_CORE core family
move

WORKED SOLUTION

Question 1

WMA12/01 January 2026

WMA12-01_2026_Jan_Q01

1

Use the binomial expansionFor $(3 + kx)^7$, the first four terms are $3^7 + \binom{7}{1}3^6(kx) + \binom{7}{2}3^5(kx)^2 + \binom{7}{3}3^4(kx)^3$.

2

Simplify the coefficientsThis gives $2187 + 5103kx + 5103k^2x^2 + 2835k^3x^3$.

3

Compare the required coefficientsThe coefficient of x^2 is $5103k^2$ and the coefficient of x^3 is $2835k^3$.

4

Solve for kSince $5103k^2 = 3(2835k^3)$ and $k \neq 0$, divide by k^2 to get $5103 = 8505k$, hence $k = \frac{5103}{8505} = \frac{3}{5}$.**FINAL ANSWER**

WMA12-01_2026_Jan_Q01

(a) $2187 + 5103kx + 5103k^2x^2 + 2835k^3x^3$. (b) $k = \frac{3}{5}$.

TOPIC 04 | FAMILY 03

Expansion Terms

WHAT THIS FAMILY ASKS

Recognise the expansion terms demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

10 classified questions

66 total marks

T04

04_TERM

04_TERM_CORE

CHAPTER
Binomial Expansion

FAMILY
Expansion terms

VARIANT
core family move

6 marks

WMA12/01 OCTOBER 2019

Question 3

Past-paper question

WMA12-01_2019_Oct_Q03

3. (a) Find the first 4 terms, in ascending powers of x , in the binomial expansion of

$$\left(1 + \frac{x}{4}\right)^{12}$$

giving each coefficient in its simplest form.

(3)

- (b) Find the term independent of x in the expansion of

$$\left(\frac{x^2 + 8}{x^5}\right)\left(1 + \frac{x}{4}\right)^{12}$$

(3)

T04 Binomial Expansion

04_TERM Expansion terms

04_TERM_CORE core family
move

WORKED SOLUTION

Question 3

WMA12/01 October 2019

WMA12-01_2019_Oct_Q03

1

Write the binomial terms

$$(1 + \frac{x}{4})^{12} = 1 + \binom{12}{1}\frac{x}{4} + \binom{12}{2}(\frac{x}{4})^2 + \binom{12}{3}(\frac{x}{4})^3 + \dots$$

2

Simplify the first four terms

This gives $1 + 3x + \frac{33}{8}x^2 + \frac{55}{16}x^3$.

3

Rewrite the prefactor

$$\frac{x^2+8}{x^5} = x^{-3} + 8x^{-5}.$$

4

Find the constant contributions

The x^{-3} part needs the x^3 coefficient, which is $\frac{55}{16}$. The $8x^{-5}$ part needs the x^5 coefficient, which is $8\binom{12}{5}(\frac{1}{4})^5 = 8 \cdot \frac{99}{128} = \frac{99}{16}$.

5

Add the constant terms

The term independent of x is $\frac{55}{16} + \frac{99}{16} = \frac{154}{16} = \frac{77}{8}$.

FINAL ANSWER

WMA12-01_2019_Oct_Q03

(a) $1 + 3x + \frac{33}{8}x^2 + \frac{55}{16}x^3$. (b) $\frac{77}{8}$.

T04

04_TERM

04_TERM_CORE

CHAPTER
Binomial Expansion

FAMILY
Expansion terms

VARIANT
core family move

7 marks

WMA12/01 JANUARY 2020

Question 2

Past-paper question

WMA12-01_2020_Jan_Q02

2. One of the terms in the binomial expansion of $(3 + ax)^6$, where a is a constant, is $540x^4$

(a) Find the possible values of a .

(4)

(b) Hence find the term independent of x in the expansion of

$$\left(\frac{1}{81} + \frac{1}{x^6}\right)(3 + ax)^6$$

(3)

T04 Binomial Expansion

04_TERM Expansion terms

04_TERM_CORE core family
move

WORKED SOLUTION

Question 2

WMA12/01 January 2020

WMA12-01_2020_Jan_Q02

1

Identify the x to the fourth termIn $(3 + ax)^6$, the x^4 term is $\binom{6}{4}3^2(ax)^4$.

2

Form the equation for a $\binom{6}{4}3^2a^4x^4 = 15 \cdot 9a^4x^4 = 135a^4x^4$. Since this term is $540x^4$, $135a^4 = 540$.

3

Solve for a $a^4 = 4$, so the real possible values are $a = \sqrt{2}$ or $a = -\sqrt{2}$.

4

Find the constant contribution from 1 over 81The constant term from $\frac{1}{81}(3 + ax)^6$ is $\frac{1}{81} \cdot 3^6 = 9$.

5

Find the constant contribution from x to the minus sixThe x^6 term in $(3 + ax)^6$ is a^6x^6 . Since $a = \pm\sqrt{2}$, $a^6 = 8$, so $\frac{1}{x^6} \cdot a^6x^6$ contributes 8.

6

Add the constant termsThe term independent of x is $9 + 8 = 17$.**FINAL ANSWER**(a) $a = \pm\sqrt{2}$. (b) 17.

WMA12-01_2020_Jan_Q02

T04

04_TERM

04_TERM_CORE

CHAPTER
Binomial Expansion

FAMILY
Expansion terms

VARIANT
core family move

4 marks

WMA12/01 MAY/JUNE 2022

Question 1

Past-paper question

WMA12-01_2022_MayJune_Q01

1. Find the first four terms, in ascending powers of x , of the binomial expansion of

$$\left(2 + \frac{3}{8}x\right)^{10}$$

Give each coefficient as an integer.

(4)

T04 Binomial Expansion

04_TERM Expansion terms

04_TERM_CORE core family
move

WORKED SOLUTION

Question 1

WMA12/01 May/June 2022

WMA12-01_2022_MayJune_Q01

1

Write the binomial terms

$$(2 + \frac{3}{8}x)^{10} = 2^{10} + \binom{10}{1}2^9\left(\frac{3}{8}x\right) + \binom{10}{2}2^8\left(\frac{3}{8}x\right)^2 + \binom{10}{3}2^7\left(\frac{3}{8}x\right)^3 + \dots$$

2

Simplify the constant and linear terms

$$2^{10} = 1024 \text{ and } \binom{10}{1}2^9\left(\frac{3}{8}\right)x = 1920x.$$

3

Simplify the quadratic term

$$\binom{10}{2}2^8\left(\frac{3}{8}\right)^2x^2 = 1620x^2.$$

4

Simplify the cubic term

$$\binom{10}{3}2^7\left(\frac{3}{8}\right)^3x^3 = 810x^3.$$

5

State the expansion

$$\text{The first four terms are } 1024 + 1920x + 1620x^2 + 810x^3.$$

FINAL ANSWER

WMA12-01_2022_MayJune_Q01

$$1024 + 1920x + 1620x^2 + 810x^3.$$

T04

04_TERM

04_TERM_CORE

CHAPTER
Binomial Expansion

FAMILY
Expansion terms

VARIANT
core family move

7 marks

WMA12/01 JANUARY 2023
Question 3

Past-paper question

WMA12-01_2023_Jan_Q03

3.
$$f(x) = \left(2 + \frac{kx}{8}\right)^7 \quad \text{where } k \text{ is a non-zero constant}$$

- (a) Find the first 4 terms, in ascending powers of x , of the binomial expansion of $f(x)$.
Give each term in simplest form.

(4)

Given that, in the binomial expansion of $f(x)$, the coefficients of x , x^2 and x^3 are the first 3 terms of an arithmetic progression,

- (b) find, using algebra, the possible values of k .

(Solutions relying entirely on calculator technology are not acceptable.)

(3)

T04 Binomial Expansion

04_TERM Expansion terms

04_TERM_CORE core family
move

WORKED SOLUTION

Question 3

WMA12/01 January 2023

WMA12-01_2023_Jan_Q03

1

Start the binomial expansion

$\left(2 + \frac{kx}{8}\right)^7$ begins with $2^7 + 7(2^6)\left(\frac{kx}{8}\right) + \binom{7}{2}(2^5)\left(\frac{kx}{8}\right)^2 + \binom{7}{3}(2^4)\left(\frac{kx}{8}\right)^3$.

2

Simplify the terms

This gives $128 + 56kx + \frac{21}{2}k^2x^2 + \frac{35}{32}k^3x^3$.

3

Use the arithmetic progression condition

The coefficients of x , x^2 and x^3 are $56k$, $\frac{21}{2}k^2$ and $\frac{35}{32}k^3$.

4

Set middle twice

For three terms in an arithmetic progression, twice the middle term equals the sum of the outer terms: $21k^2 = 56k + \frac{35}{32}k^3$.

5

Use k non-zero

Since $k \neq 0$, divide by k and multiply by 32: $672k = 1792 + 35k^2$.

6

Solve the quadratic

So $35k^2 - 672k + 1792 = 0$. The discriminant is $672^2 - 4(35)(1792) = 200704 = 448^2$.

7

Find the possible values

$k = \frac{672 \pm 448}{70}$, giving $k = 16$ or $k = \frac{16}{5}$.

FINAL ANSWER

WMA12-01_2023_Jan_Q03

(a) $128 + 56kx + \frac{21}{2}k^2x^2 + \frac{35}{32}k^3x^3$. (b) $k = \frac{16}{5}$ or $k = 16$.

T04

04_TERM

04_TERM_CORE

CHAPTER
Binomial Expansion

FAMILY
Expansion terms

VARIANT
core family move

12 marks

WMA12/01 OCTOBER 2023

Question 10

Past-paper question

WMA12-01_2023_Oct_Q10

10. (i) (a) Find, in ascending powers of x , the 2nd, 3rd and 5th terms of the binomial expansion of

$$(3 + 2x)^6 \quad (3)$$

For a particular value of x , these three terms form consecutive terms in a geometric series.

- (b) Find this value of x . (3)

- (ii) In a **different** geometric series,

- the first term is $\sin^2 \theta$
- the common ratio is $2 \cos \theta$
- the sum to infinity is $\frac{8}{5}$

- (a) Show that

$$5 \cos^2 \theta - 16 \cos \theta + 3 = 0 \quad (3)$$

- (b) Hence find the exact value of the 2nd term in the series. (3)

T04 Binomial Expansion

04_TERM Expansion terms

04_TERM_CORE core family
move

WORKED SOLUTION

Question 10

WMA12/01 October 2023

WMA12-01_2023_Oct_Q10

1

Find the second term

The 2nd term of $(3 + 2x)^6$ is $\binom{6}{1}3^5(2x) = 6 \cdot 243 \cdot 2x = 2916x$.

2

Find the third term

The 3rd term is $\binom{6}{2}3^4(2x)^2 = 15 \cdot 81 \cdot 4x^2 = 4860x^2$.

3

Find the fifth term

The 5th term is $\binom{6}{4}3^2(2x)^4 = 15 \cdot 9 \cdot 16x^4 = 2160x^4$.

4

Use the geometric condition

For three consecutive terms in a geometric series, the middle term squared equals the product of the two outer terms.

5

Set up the equation

$$(4860x^2)^2 = (2916x)(2160x^4).$$

6

Solve for x

For the non-zero geometric case, divide by x^4 : $4860^2 = 2916 \cdot 2160x$, so $x = \frac{4860^2}{2916 \cdot 2160} = \frac{15}{4}$.

7

Use the sum to infinity

In the different geometric series, first term $= \sin^2 \theta$, common ratio $= 2 \cos \theta$, and $S_\infty = \frac{8}{5}$.

8

Form the equation

$$\frac{\sin^2 \theta}{1 - 2 \cos \theta} = \frac{8}{5}.$$

9

Use sine squared

$5 \sin^2 \theta = 8(1 - 2 \cos \theta)$. With $\sin^2 \theta = 1 - \cos^2 \theta$, this becomes $5(1 - \cos^2 \theta) = 8 - 16 \cos \theta$.

10

Show the required equationRearranging gives $5 \cos^2 \theta - 16 \cos \theta + 3 = 0$.

11

Solve for cosine $5 \cos^2 \theta - 16 \cos \theta + 3 = 0$ gives $\cos \theta = 3$ or $\cos \theta = \frac{1}{5}$. The possible value is $\cos \theta = \frac{1}{5}$.

12

Find the second termThe 2nd term is first term times common ratio: $\sin^2 \theta (2 \cos \theta) = \left(1 - \frac{1}{25}\right) \cdot \frac{2}{5} = \frac{24}{25} \cdot \frac{2}{5} = \frac{48}{125}$.**FINAL ANSWER**

WMA12-01_2023_Oct_Q10

(i)(a) $2916x, 4860x^2, 2160x^4$. (i)(b) $x = \frac{15}{4}$. (ii)(a) **Proven.** (ii)(b) $\frac{48}{125}$.

DR ESLAM AHMED

T04

04_TERM

04_TERM_CORE

CHAPTER
Binomial Expansion

FAMILY
Expansion terms

VARIANT
core family move

5 marks

WMA12/01 MAY/JUNE 2024

Question 1

Past-paper question

WMA12-01_2024_MayJune_Q01

1. (a) Find the first four terms, in ascending powers of x , of the binomial expansion of

$$\left(1 - \frac{1}{6}x\right)^9$$

giving each term in simplest form.

(3)

- (b) Hence find the coefficient of x^3 in the expansion of

$$(10x + 3)\left(1 - \frac{1}{6}x\right)^9$$

giving the answer in simplest form.

(2)

T04 Binomial Expansion

04_TERM Expansion terms

04_TERM_CORE core family
move

WORKED SOLUTION

Question 1

WMA12/01 May/June 2024

WMA12-01_2024_MayJune_Q01

1

Expand the first terms

$$\left(1 - \frac{1}{6}x\right)^9 = 1 + 9\left(-\frac{1}{6}x\right) + \binom{9}{2}\left(-\frac{1}{6}x\right)^2 + \binom{9}{3}\left(-\frac{1}{6}x\right)^3 + \dots$$

2

Simplify

The first four terms are $1 - \frac{3}{2}x + x^2 - \frac{7}{18}x^3$.

3

Identify contributions to x^3 cubed

In $(10x + 3)\left(1 - \frac{1}{6}x\right)^9$, the x^3 coefficient comes from 3 times the x^3 coefficient and $10x$ times the x^2 coefficient.

4

Calculate the coefficient

$$3\left(-\frac{7}{18}\right) + 10(1) = -\frac{7}{6} + 10 = \frac{53}{6}.$$

FINAL ANSWER

WMA12-01_2024_MayJune_Q01

$$(a) 1 - \frac{3}{2}x + x^2 - \frac{7}{18}x^3. \quad (b) \frac{53}{6}.$$

T04

04_TERM

04_TERM_CORE

CHAPTER
Binomial Expansion

FAMILY
Expansion terms

VARIANT
core family move

7 marks

WMA12/01 MAY/JUNE (R) 2024

Question 4

Past-paper question

WMA12-01_2024_MayJuneR_Q04

4. (a) Find the first 4 terms, in ascending powers of x , of the binomial expansion of

$$(3 + 2x)^6$$

giving each coefficient in simplest form.

(4)

- (b) Hence find the coefficient of x^2 in the expansion of

$$\left(2x^2 - \frac{1}{6x}\right)(3 + 2x)^6$$

(3)

T04 Binomial Expansion

04_TERM Expansion terms

04_TERM_CORE core family
move

WORKED SOLUTION

Question 4

WMA12/01 May/June (R) 2024

WMA12-01_2024_MayJuneR_Q04

1

Write the expansion terms

$$(3 + 2x)^6 = 3^6 + \binom{6}{1}3^5(2x) + \binom{6}{2}3^4(2x)^2 + \binom{6}{3}3^3(2x)^3 + \dots$$

2

Simplify the first four terms

This gives $729 + 2916x + 4860x^2 + 4320x^3$.

3

Identify x squared contributions

In $(2x^2 - \frac{1}{6x})(3 + 2x)^6$, the x^2 term comes from $2x^2$ times the constant term and $-\frac{1}{6x}$ times the x^3 term.

4

Calculate the coefficient

The coefficient is $2(729) - \frac{1}{6}(4320) = 1458 - 720 = 738$.

FINAL ANSWER

WMA12-01_2024_MayJuneR_Q04

(a) $729 + 2916x + 4860x^2 + 4320x^3$. (b) 738.

T04

04_TERM

04_TERM_CORE

CHAPTER
Binomial Expansion

FAMILY
Expansion terms

VARIANT
core family move

7 marks

WMA12/01 OCTOBER 2024

Question 5

Past-paper question

WMA12-01_2024_Oct_Q05

5. (a) Find, in terms of a , the first 3 terms, in ascending powers of x , of the binomial expansion of

$$(2 + ax)^6$$

where a is a non-zero constant. Give each term in simplest form.

(3)

$$f(x) = \left(3 + \frac{1}{x}\right)^2 (2 + ax)^6$$

Given that the constant term in the expansion of $f(x)$ is 576

- (b) find the value of a .

(4)

T04 Binomial Expansion

04_TERM Expansion terms

04_TERM_CORE core family
move

WORKED SOLUTION

Question 5

WMA12/01 October 2024

WMA12-01_2024_Oct_Q05

1

Write the first three binomial terms

$$(2 + ax)^6 = 2^6 + 6(2^5)(ax) + \binom{6}{2}(2^4)(ax)^2 + \dots$$

2

Simplify the terms

The first three terms are $64 + 192ax + 240a^2x^2$.

3

Expand the other factor

$$\left(3 + \frac{1}{x}\right)^2 = 9 + 6x^{-1} + x^{-2}.$$

4

Collect constant contributions

The constant term in $\left(3 + \frac{1}{x}\right)^2 (2 + ax)^6$ is $9(64) + 6(192a) + 240a^2$.

5

Use the given constant term

$$576 + 1152a + 240a^2 = 576, \text{ so } 240a^2 + 1152a = 0.$$

6

Solve for a

$$48a(5a + 24) = 0. \text{ Since } a \neq 0, 5a + 24 = 0, \text{ so } a = -\frac{24}{5}.$$

FINAL ANSWER

WMA12-01_2024_Oct_Q05

$$(a) 64 + 192ax + 240a^2x^2. \quad (b) a = -\frac{24}{5}.$$

T04

04_TERM

04_TERM_CORE

CHAPTER
Binomial Expansion

FAMILY
Expansion terms

VARIANT
core family move

5 marks

WMA12/01 JANUARY 2025
Question 2

Past-paper question

WMA12-01_2025_Jan_Q02

2. (a) Find the first 4 terms, in ascending powers of x , of the binomial expansion of

$$(2 - 5x)^8$$

giving each term in simplest form.

(4)

This expansion is to be used to find an approximation for 2.05^8

- (b) State the value of x that should be used.

(There is no need to carry out this calculation.)

(1)

T04 Binomial Expansion

04_TERM Expansion terms

04_TERM_CORE core family
move

WORKED SOLUTION

Question 2

WMA12/01 January 2025

WMA12-01_2025_Jan_Q02

1

Write the first four terms

$$(2 - 5x)^8 = 2^8 + 8(2^7)(-5x) + \binom{8}{2}(2^6)(-5x)^2 + \binom{8}{3}(2^5)(-5x)^3 + \dots$$

2

Simplify each term

The first four terms are $256 - 5120x + 44800x^2 - 224000x^3$.

3

Connect the expansion to 2.05

To use $(2 - 5x)^8$ for 2.05^8 , set $2 - 5x = 2.05$.

4

Find x

$$-5x = 0.05, \text{ so } x = -0.01.$$

FINAL ANSWER

WMA12-01_2025_Jan_Q02

$$(a) 256 - 5120x + 44800x^2 - 224000x^3. \quad (b) x = -0.01.$$

T04

04_TERM

04_TERM_CORE

CHAPTER
Binomial Expansion

FAMILY
Expansion terms

VARIANT
core family move

6 marks

WMA12/01 MAY/JUNE 2025

Question 1

Past-paper question

WMA12-01_2025_MayJune_Q01

- 1: (a) Find the first 4 terms, in ascending powers of x , of the binomial expansion of

$$(1 - 4x)^7$$

giving each term in simplest form.

(4)

In the series expansion of

$$(5 + kx)(1 - 4x)^7$$

where k is a constant

the coefficient of the term in x^2 is 1316

- (b) Use the answer to part (a) to find the value of k .

(2)

T04 Binomial Expansion

04_TERM Expansion terms

04_TERM_CORE core family
move

WORKED SOLUTION

Question 1

WMA12/01 May/June 2025

WMA12-01_2025_MayJune_Q01

1

Write the binomial terms

$$(1 - 4x)^7 = 1 + 7(-4x) + \binom{7}{2}(-4x)^2 + \binom{7}{3}(-4x)^3 + \dots$$

2

Simplify

The first four terms are $1 - 28x + 336x^2 - 2240x^3$.

3

Find the x squared terms

In $(5 + kx)(1 - 4x)^7$, the x^2 coefficient comes from $5(336x^2)$ and $(kx)(-28x)$.

4

Use the given coefficient

$$5(336) - 28k = 1316, \text{ so } 1680 - 28k = 1316.$$

5

Find k

$$28k = 364, \text{ hence } k = 13.$$

FINAL ANSWER

WMA12-01_2025_MayJune_Q01

$$(a) 1 - 28x + 336x^2 - 2240x^3. \quad (b) k = 13.$$

PURE MATHEMATICS 2

ADAPTIVE ALL CLASSIFIED

Presentation Edition | Solutions

CLASSIFIED BY IDEA | ROOM TO THINK

Every demand appears clearly, with practical space for the mathematics.

Dr Eslam Ahmed

Elite Mathematics

ElitelGCSE.com | 011 2000 9622

PURE 2

PURE 2 | TOPIC 05

Arithmetic Sequences

Your classified route

6

QUESTIONS

41

MARKS

2

FAMILIES

3

VARIANTS

DR ESLAM AHMED

CLASSIFIED BY IDEA

Move from recognition to method, then use the working space to complete the solution.

TOPIC 05 | FAMILY 01

Unknown Terms

WHAT THIS FAMILY ASKS

Recognise the unknown terms demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 Use simultaneous equations when two term positions are given.

2 classified questions

14 total marks

T05

05_SOLVE

05_SOLVE_HV02

CHAPTER
Arithmetic Sequences

FAMILY
Unknown terms

VARIANT
Use simultaneous equations
when two term positions are
given.

7 marks

WMA12/01 OCTOBER 2023

Question 8

Past-paper question

WMA12-01_2023_Oct_Q08

8. In a large theatre there are n rows of seats, where n is a constant.

The number of seats in the first row is a , where a is a constant.

In each subsequent row there are 4 more seats than in the previous row so that

- in the 2nd row there are $(a + 4)$ seats
- in the 3rd row there are $(a + 8)$ seats
- the number of seats in each row form an **arithmetic** sequence

Given that the **total** number of seats in the first 10 rows is 360

- (a) find the value of a .

(2)

Given also that the total number of seats in the n rows is 2146

- (b) show that

$$n^2 + 8n - 1073 = 0$$

(2)

- (c) Hence

- (i) state the number of rows of seats in the theatre,
- (ii) find the maximum number of seats in any one row.

(3)

T05 Arithmetic Sequences

05_SOLVE Unknown terms

05_SOLVE_HV02 Use simultaneous equations when two term positions are given.

WORKED SOLUTION

Question 8

WMA12/01 October 2023

WMA12-01_2023_Oct_Q08

1

Use the arithmetic sum formulaThe first term is a and the common difference is 4.

2

Use the first 10 rows

$$S_{10} = \frac{10}{2} (2a + 9 \cdot 4) = 360.$$

3

Find a

$$5(2a + 36) = 360, \text{ so } 10a + 180 = 360 \text{ and } a = 18.$$

4

Write S_n

$$S_n = \frac{n}{2} (2(18) + (n-1)4) = \frac{n}{2} (4n + 32) = 2n^2 + 16n.$$

5

Use the total number of seats

$$\text{Since the total is 2146, } 2n^2 + 16n = 2146.$$

6

Show the equation

$$\text{Dividing by 2 and rearranging gives } n^2 + 8n - 1073 = 0.$$

7

Solve for n

$$n = \frac{-8 \pm \sqrt{8^2 + 4(1073)}}{2} = \frac{-8 \pm 66}{2}, \text{ so } n = 29 \text{ or } n = -37.$$

8

Choose the row count

The number of rows must be positive, so there are 29 rows.

9

Find the maximum row size

$$\text{The largest row is the last row: } 18 + (29-1)4 = 18 + 112 = 130 \text{ seats.}$$

FINAL ANSWER

WMA12-01_2023_Oct_Q08

(a) $a = 18$. (b) **Proven.** (c)(i) **29 rows.** (c)(ii) **130 seats.**

T05

05_SOLVE

05_SOLVE_HV02

CHAPTER
Arithmetic Sequences

FAMILY
Unknown terms

VARIANT
Use simultaneous equations
when two term positions are
given.

7 marks

WMA12/01 MAY/JUNE 2024

Question 2

Past-paper question

WMA12-01_2024_MayJune_Q02

2. **In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.**

In an arithmetic series,

- the sixth term is 2
- the sum of the first ten terms is -80

For this series,

- (a) find the value of the first term and the value of the common difference.

(4)

- (b) Hence find the smallest value of n for which

$$S_n > 8000$$

(3)

T05 Arithmetic Sequences

05_SOLVE Unknown terms

05_SOLVE_HV02 Use simultaneous equations when two term positions are given.

WORKED SOLUTION

Question 2

WMA12/01 May/June 2024

WMA12-01_2024_MayJune_Q02

1

Use the sixth termFor an arithmetic series, $u_6 = a + 5d = 2$.

2

Use the sum of ten terms $S_{10} = \frac{10}{2}(2a + 9d) = -80$, so $2a + 9d = -16$.

3

Solve simultaneouslyFrom $a + 5d = 2$, $a = 2 - 5d$. Substitute into $2a + 9d = -16$.

4

Find a and d $2(2 - 5d) + 9d = -16$, so $4 - d = -16$ and $d = 20$. Then $a = 2 - 5(20) = -98$.

5

Write S_n $S_n = \frac{n}{2}(2a + (n-1)d) = \frac{n}{2}(-196 + 20n - 20) = 10n^2 - 108n$.

6

Set up the inequalityWe need $10n^2 - 108n > 8000$.

7

Solve the boundary $10n^2 - 108n - 8000 = 0$ gives $5n^2 - 54n - 4000 = 0$.

8

Find the positive threshold $n = \frac{54 + \sqrt{82916}}{10} = 34.195 \dots$

9

Choose the smallest integerTherefore the smallest integer value with $S_n > 8000$ is $n = 35$.**FINAL ANSWER**

WMA12-01_2024_MayJune_Q02

(a) First term = -98 , common difference = 20 . (b) $n = 35$.

TOPIC 05 | FAMILY 02

Series Sum

WHAT THIS FAMILY ASKS

Recognise the series sum demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

V02 If last term is unknown, use $l = a + (n-1)d$ first.

4 classified questions

27 total marks

T05

05_SUM

05_SUM_CORE

CHAPTER

Arithmetic Sequences

FAMILY

Series sum

VARIANT

core family move

13 marks

WMA12/01 MAY/JUNE (R) 2024

Question 8

Past-paper question

WMA12-01_2024_MayJuneR_Q08

8. (i) (a) In an **arithmetic** series the first term is a and the common difference is d .

Show that

$$S_n = \frac{n}{2} \{2a + (n-1)d\} \quad (3)$$

- (b) Hence find

$$900 + 892 + 884 + \dots + 500 \quad (3)$$

- (ii) Given that the first three terms of a **geometric** series are

$$k+4 \quad k-2 \quad 11-k$$

where k is a constant,

- (a) show that

$$2k^2 - 11k - 40 = 0 \quad (3)$$

Given also that this series is convergent,

- (b) find the value of S_∞ (4)

T05 Arithmetic Sequences

05_SUM Series sum

05_SUM_CORE core family
move

WORKED SOLUTION

Question 8

WMA12/01 May/June (R) 2024

WMA12-01_2024_MayJuneR_Q08

1

Write the first and last termsThe first term is a and the n th term is $a + (n - 1)d$.

2

Use the arithmetic sum formula $S_n = \frac{n}{2}(\text{first term} + \text{last term})$.

3

Show the formulaTherefore $S_n = \frac{n}{2}\{a + [a + (n - 1)d]\} = \frac{n}{2}\{2a + (n - 1)d\}$.

4

Identify the arithmetic seriesFor $900 + 892 + 884 + \dots + 500$, the first term is 900, the common difference is -8 , and the last term is 500.

5

Find the number of terms $500 = 900 + (n - 1)(-8)$, so $n - 1 = 50$ and $n = 51$.

6

Find the sum $S = \frac{51}{2}(900 + 500) = 35700$.

7

Use the geometric conditionFor $k + 4$, $k - 2$, $11 - k$ to be consecutive geometric terms, $(k - 2)^2 = (k + 4)(11 - k)$.

8

Show the quadratic $k^2 - 4k + 4 = -k^2 + 7k + 44$, so $2k^2 - 11k - 40 = 0$.

9

Solve for k $2k^2 - 11k - 40 = 0$ gives $k = 8$ or $k = -\frac{5}{2}$.

10

Use convergenceIf $k = 8$, the terms are 12, 6, 3 and the common ratio is $\frac{1}{2}$, so the series is convergent. If $k = -\frac{5}{2}$, the common ratio is -3 , so it is not convergent.

11

Find S_{∞}

Use $k = 8$: $S_{\infty} = \frac{12}{1-1/2} = 24$.

FINAL ANSWER

WMA12-01_2024_MayJuneR_Q08

(i)(a) **Proven.** (i)(b) 35700. (ii)(a) **Proven.** (ii)(b) $S_{\infty} = 24$.

DR ESLAM AHMED

T05

05_SUM

05_SUM_CORE

CHAPTER
Arithmetic Sequences

FAMILY
Series sum

VARIANT
core family move

4 marks

WMA12/01 JANUARY 2025
Question 1

Past-paper question

WMA12-01_2025_Jan_Q01

1. The arithmetic series S is given by

$$S = 2 + 5 + 8 + 11 + \dots + 254$$

Find

- (a) the number of terms in the series,

(2)

- (b) the sum of the series.

(2)

T05 Arithmetic Sequences

05_SUM Series sum

05_SUM_CORE core family
move

WORKED SOLUTION

Question 1

WMA12/01 January 2025

WMA12-01_2025_Jan_Q01

1

Identify the arithmetic series

The first term is 2, the common difference is 3, and the last term is 254.

2

Find the number of terms

Use $254 = 2 + (n - 1)3$. Then $252 = 3(n - 1)$, so $n - 1 = 84$ and $n = 85$.

3

Use the sum formula

$S_n = \frac{n}{2}(a + l)$, where a is the first term and l is the last term.

4

Find the sum

$$S = \frac{85}{2}(2 + 254) = 85 \times 128 = 10880.$$

FINAL ANSWER

(a) 85. (b) 10880.

WMA12-01_2025_Jan_Q01

T05

05_SUM

05_SUM_CORE

CHAPTER

Arithmetic Sequences

FAMILY

Series sum

VARIANT

core family move

4 marks

WMA12/01 OCTOBER 2025

Question 1

Past-paper question

WMA12-01_2025_Oct_Q01

1. An arithmetic series starts

$$78 + 75 + 72 + \dots$$

Find

- (a) the 18th term in the series,

(2)

- (b) the sum of the first 100 terms of the series.

(2)

T05 Arithmetic Sequences

05_SUM Series sum

05_SUM_CORE core family
move

WORKED SOLUTION

Question 1

WMA12/01 October 2025

WMA12-01_2025_Oct_Q01

1

Identify the sequence

The first term is 78 and the common difference is $75 - 78 = -3$.

2

Find the eighteenth term

$$u_{18} = 78 + 17(-3) = 27.$$

3

Use the sum formula

$$S_{100} = \frac{100}{2}\{2(78) + 99(-3)\}.$$

4

Evaluate the sum

$$S_{100} = 50(156 - 297) = 50(-141) = -7050.$$

FINAL ANSWER

(a) 27. (b) -7050.

WMA12-01_2025_Oct_Q01

T05

05_SUM

05_SUM_HV01

CHAPTER

Arithmetic Sequences

FAMILY

Series sum

VARIANT

If last term is unknown, use
 $(l=a+(n-1)d)$ first.

6 marks

WMA12/01 OCTOBER 2021

Question 5

Past-paper question

WMA12-01_2021_Oct_Q05

5. A company that owned a silver mine

- extracted 480 tonnes of silver from the mine in year 1
- extracted 465 tonnes of silver from the mine in year 2
- extracted 450 tonnes of silver from the mine in year 3

and so on, forming an arithmetic sequence.

(a) Find the mass of silver extracted in year 14

(2)

After a total of 7770 tonnes of silver was extracted, the company stopped mining.

Given that this occurred at the end of year N ,

(b) show that

$$N^2 - 65N + 1036 = 0$$

(3)

(c) Hence, state the value of N .

(1)

T05 Arithmetic Sequences

05_SUM Series sum

05_SUM_HV01 If last term is unknown, use $l = a + (n-1)d$ first.

WORKED SOLUTION

Question 5

WMA12/01 October 2021

WMA12-01_2021_Oct_Q05

1

Identify the arithmetic sequence

The first term is 480 and the common difference is $465 - 480 = -15$.

2

Find year fourteen

$u_{14} = 480 + 13(-15) = 285$, so 285 tonnes were extracted in year 14.

3

Set up the total

The total after N years is $S_N = \frac{N}{2}\{2(480) + (N-1)(-15)\}$.

4

Use the total extraction

$\frac{N}{2}(975 - 15N) = 7770$.

5

Show the quadratic

Multiplying by 2 and rearranging gives $15N^2 - 975N + 15540 = 0$, so $N^2 - 65N + 1036 = 0$.

6

Solve and choose N

$N = 28$ or 37 . Since extraction amounts would become negative before year 37, the stopping year is $N = 28$.

FINAL ANSWER

WMA12-01_2021_Oct_Q05

(a) 285 tonnes. (b) Proven: $N^2 - 65N + 1036 = 0$. (c) $N = 28$.

PURE MATHEMATICS 2

ADAPTIVE ALL CLASSIFIED

Presentation Edition | Solutions

CLASSIFIED BY IDEA | ROOM TO THINK

Every demand appears clearly, with practical space for the mathematics.

Dr Eslam Ahmed

Elite Mathematics

ElitelGCSE.com | 011 2000 9622

PURE 2

PURE 2 | TOPIC 06

Geometric Sequences

Your classified route

7

QUESTIONS

55

MARKS

3

FAMILIES

5

VARIANTS

DR ESLAM AHMED

CLASSIFIED BY IDEA

Move from recognition to method, then use the working space to complete the solution.

TOPIC 06 | FAMILY 01

Infinite Sum

WHAT THIS FAMILY ASKS

Recognise the infinite sum demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

V02 State the convergence condition before using the formula.

3 classified questions

24 total marks

T06

06_INFTY

06_INFTY_CORE

CHAPTER
Geometric Sequences

FAMILY
Infinite sum

VARIANT
core family move

8 marks

WMA12/01 MAY/JUNE 2022

Question 6

Past-paper question

WMA12-01_2022_MayJune_Q06

6. In a geometric sequence $u_1, u_2, u_3, \dots$

- the common ratio is r
- $u_2 + u_3 = 6$
- $u_4 = 8$

(a) Show that r satisfies

$$3r^2 - 4r - 4 = 0$$

(3)

Given that the geometric sequence has a sum to infinity,

(b) find u_1

(3)

(c) find S_∞

(2)

T06 Geometric Sequences

06_INFTY Infinite sum

06_INFTY_CORE core family
move

WORKED SOLUTION

Question 6

WMA12/01 May/June 2022

WMA12-01_2022_MayJune_Q06

1

Write the terms

$$u_2 = u_1 r, u_3 = u_1 r^2, \text{ and } u_4 = u_1 r^3.$$

2

Use u four

$$\text{Since } u_4 = 8, u_1 r^3 = 8, \text{ so } u_1 r = \frac{8}{r^2}.$$

3

Use u two plus u three

$$u_2 + u_3 = u_1 r(1 + r) = 6.$$

4

Show the quadratic

$$\text{Substitute } u_1 r = \frac{8}{r^2} \text{ to get } \frac{8}{r^2}(1 + r) = 6. \text{ Hence } 8 + 8r = 6r^2, \text{ so } 3r^2 - 4r - 4 = 0.$$

5

Solve for r

$$3r^2 - 4r - 4 = 0 \text{ gives } r = 2 \text{ or } r = -\frac{2}{3}.$$

6

Choose the sum-to-infinity ratio

$$\text{A sum to infinity requires } |r| < 1, \text{ so } r = -\frac{2}{3}.$$

7

Find u one

$$u_1 r^3 = 8, \text{ so } u_1 \left(-\frac{2}{3}\right)^3 = 8 \text{ and } u_1 = -27.$$

8

Find the infinite sum

$$S_\infty = \frac{u_1}{1-r} = \frac{-27}{1+2/3} = -\frac{81}{5}.$$

FINAL ANSWER

WMA12-01_2022_MayJune_Q06

$$(a) \text{ Proven: } 3r^2 - 4r - 4 = 0. \quad (b) u_1 = -27. \quad (c) S_\infty = -\frac{81}{5}.$$

T06

06_INFTY

06_INFTY_CORE

CHAPTER
Geometric Sequences

FAMILY
Infinite sum

VARIANT
core family move

6 marks

WMA12/01 JANUARY 2025

Question 4

Past-paper question

WMA12-01_2025_Jan_Q04

4. **In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.**

Given that, in a particular geometric series,

- the sum of the first three terms is 70.2
- the sum to infinity is 75

find, for this series,

(a) the common ratio,

(4)

(b) the first term.

(2)

T06 Geometric Sequences

06_INFTY Infinite sum

06_INFTY_CORE core family
move

WORKED SOLUTION

Question 4

WMA12/01 January 2025

WMA12-01_2025_Jan_Q04

1

Write the two sums

For first term a and common ratio r , $S_{\infty} = \frac{a}{1-r} = 75$ and $S_3 = \frac{a(1-r^3)}{1-r} = 70.2$.

2

Divide the equations

$\frac{S_3}{S_{\infty}} = 1 - r^3$, so $\frac{70.2}{75} = 1 - r^3$.

3

Find the ratio

$0.936 = 1 - r^3$, so $r^3 = 0.064$ and $r = 0.4$.

4

Find the first term

Use $\frac{a}{1-0.4} = 75$. Hence $a = 75(0.6) = 45$.

FINAL ANSWER

WMA12-01_2025_Jan_Q04

(a) $r = 0.4$. (b) 45.

T06

06_INFTY

06_INFTY_HV01

CHAPTER
Geometric Sequences

FAMILY
Infinite sum

VARIANT
State the convergence
condition before using the
formula.

10 marks

WMA12/01 OCTOBER 2021

Question 7

Past-paper question

WMA12-01_2021_Oct_Q07

7. (i) A geometric sequence has first term 4 and common ratio 6
Given that the n^{th} term is greater than 10^{100} , find the minimum possible value of n .
(3)

- (ii) A different geometric sequence has first term a and common ratio r .

Given that

- the second term of the sequence is -6
- the sum to infinity of the series is 25

- (a) show that

$$25r^2 - 25r - 6 = 0 \quad (3)$$

- (b) Write down the solutions of

$$25r^2 - 25r - 6 = 0 \quad (1)$$

Hence,

- (c) state the value of r , giving a reason for your answer,
(1)

- (d) find the sum of the first 4 terms of the series.
(2)

T06 Geometric Sequences

06_INFTY Infinite sum

06_INFTY_HV01 State the convergence condition before using the formula.

WORKED SOLUTION

Question 7

WMA12/01 October 2021

WMA12-01_2021_Oct_Q07

1

Write the n th term

The n th term of the first sequence is $4 \cdot 6^{n-1}$.

2

Use logarithms

We need $4 \cdot 6^{n-1} > 10^{100}$, so $n - 1 > \frac{100 - \log_{10} 4}{\log_{10} 6} = 127.736 \dots$

3

Choose the smallest integer

Thus n must be greater than 128.736..., so the minimum possible value is 129.

4

Use the second term

For the second sequence, $ar = -6$, so $a = -\frac{6}{r}$.

5

Use the sum to infinity

$\frac{a}{1-r} = 25$. Substituting $a = -\frac{6}{r}$ gives $-6 = 25r(1-r)$.

6

Show the quadratic

Rearranging gives $25r^2 - 25r - 6 = 0$.

7

Solve for r

$25r^2 - 25r - 6 = 0$ gives $r = \frac{6}{5}$ or $r = -\frac{1}{5}$.

8

Choose the valid ratio

A sum to infinity exists only when $|r| < 1$, so $r = -\frac{1}{5}$.

9

Find the first term

Since $ar = -6$, $a = 30$.

10

Sum four terms

$S_4 = 30 \frac{1 - (-1/5)^4}{1 - (-1/5)} = \frac{624}{25}$.

FINAL ANSWER

WMA12-01_2021_Oct_Q07

(i) $n = 129$. (ii)(a) **Proven.** (ii)(b) $r = \frac{6}{5}$ or $-\frac{1}{5}$. (ii)(c) $r = -\frac{1}{5}$. (ii)(d) $\frac{624}{25}$.

DR ESLAM AHMED

TOPIC 06 | FAMILY 02

Common Ratio

WHAT THIS FAMILY ASKS

Recognise the common ratio demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

1 classified questions

7 total marks

T06

06_RATIO

06_RATIO_CORE

CHAPTER
Geometric Sequences

FAMILY
Common ratio

VARIANT
core family move

7 marks

WMA12/01 OCTOBER 2022
Question 8

Past-paper question

WMA12-01_2022_Oct_Q08

8. A geometric sequence has first term a and common ratio r

Given that $S_{\infty} = 3a$

- (a) show that $r = \frac{2}{3}$

(2)

Given also that

$$u_2 - u_4 = 16$$

where u_k is the k^{th} term of this sequence,

- (b) find the value of S_{10} giving your answer to one decimal place.

(5)

T06 Geometric Sequences

06_RATIO Common ratio

06_RATIO_CORE core family
move

WORKED SOLUTION

Question 8

WMA12/01 October 2022

WMA12-01_2022_Oct_Q08

1

Use the sum to infinity formulaFor a geometric sequence, $S_{\infty} = \frac{a}{1-r}$. The question gives $S_{\infty} = 3a$.

2

Find the common ratio $\frac{a}{1-r} = 3a$. Since $u_2 - u_4 = 16$, $a \neq 0$, so $\frac{1}{1-r} = 3$ and $r = \frac{2}{3}$.

3

Use u_2 minus u_4 $u_2 - u_4 = ar - ar^3 = ar(1 - r^2) = 16$.

4

Substitute r $a\left(\frac{2}{3}\right)\left(1 - \frac{4}{9}\right) = 16$.

5

Solve for a $a \cdot \frac{2}{3} \cdot \frac{5}{9} = 16$, so $\frac{10a}{27} = 16$ and $a = \frac{216}{5} = 43.2$.

6

Find S_{10} $S_{10} = \frac{a(1-r^{10})}{1-r} = 3a\left(1 - \left(\frac{2}{3}\right)^{10}\right)$.

7

Evaluate $S_{10} = 3(216/5)(1 - (2/3)^{10}) = 92840/729 = 127.352\dots$, so $S_{10} = 127.4$ to 1 decimal place, as requested in the question.**FINAL ANSWER**

WMA12-01_2022_Oct_Q08

(a) $r = 2/3$. (b) $a = 43.2$; $S_{10} = 92840/729 = 127.352\dots$, so $S_{10} = 127.4$ to 1 d.p.

TOPIC 06 | FAMILY 03

Finite Sum

WHAT THIS FAMILY ASKS

Recognise the finite sum demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

V02 If a sum is given, solve the resulting exponential equation carefully.

3 classified questions

24 total marks

T06

06_SUM

06_SUM_CORE

CHAPTER
Geometric Sequences

FAMILY
Finite sum

VARIANT
core family move

8 marks

WMA12/01 OCTOBER 2025

Question 8

Past-paper question

WMA12-01_2025_Oct_Q08

8. (i) A geometric series has first term a and common ratio r .

Prove that the sum of the first n terms of this series S_n is given by

$$S_n = \frac{a(1-r^n)}{1-r} \quad (3)$$

- (ii) A liquid is to be stored in a barrel.

Due to evaporation, the volume of the liquid in the barrel at the end of each year is 8% less than the volume of the liquid in the barrel at the start of the year.

At the start of the first year, the barrel is filled with 150 litres of the liquid.

- (a) Show that the amount of the liquid in the barrel at the end of 6 years is approximately 91 litres. (2)

At the start of each year a new barrel is filled with 150 litres of the liquid so that, at the end of 40 years, there are 40 barrels containing the liquid.

- (b) Calculate the total amount of the liquid, to the nearest litre, in the 40 barrels at the end of 40 years. (3)

T06 Geometric Sequences

06_SUM Finite sum

06_SUM_CORE core family
move

WORKED SOLUTION

Question 8

WMA12/01 October 2025

WMA12-01_2025_Oct_Q08

1

Write S_n

$$S_n = a + ar + ar^2 + \dots + ar^{n-1}.$$

2

Multiply by r

$$rS_n = ar + ar^2 + ar^3 + \dots + ar^n.$$

3

Subtract

$$S_n - rS_n = a - ar^n = a(1 - r^n).$$

4

Rearrange

$$(1 - r)S_n = a(1 - r^n), \text{ so } S_n = \frac{a(1 - r^n)}{1 - r}.$$

5

Use the yearly multiplier

An 8% decrease each year means the multiplier is 0.92.

6

Show the amount after six years

After 6 years, the amount is $150(0.92)^6 = 90.953 \dots$, which is approximately 91 litres.

7

Set up the forty barrel total

At the end of 40 years, the amounts are $150(0.92), 150(0.92)^2, \dots, 150(0.92)^{40}$.

8

Sum the geometric sequence

$$\text{The total is } 150(0.92) \frac{1 - 0.92^{40}}{1 - 0.92} = 1663.581 \dots$$

9

Round the total

To the nearest litre, the total amount is 1664 litres.

FINAL ANSWER

WMA12-01_2025_Oct_Q08

(i) **Proven.** (ii)(a) **Approximately 91 litres.** (ii)(b) **1664 litres.**

T06

06_SUM

06_SUM_HV03

CHAPTER

Geometric Sequences

FAMILY

Finite sum

VARIANT

If a sum is given, solve the
resulting exponential
equation carefully.

7 marks

WMA12/01 JANUARY 2023

Question 7

Past-paper question

WMA12-01_2023_Jan_Q07

7. A geometric sequence has first term a and common ratio r , where $r > 0$

Given that

- the 3rd term is 20
- the 5th term is 12.8

- (a) show that $r = 0.8$

(1)

- (b) Hence find the value of a .

(2)

Given that the sum of the first n terms of this sequence is greater than 156

- (c) find the smallest possible value of n .

(Solutions based entirely on graphical or numerical methods are not acceptable.)

(4)

T06 Geometric Sequences

06_SUM Finite sum

06_SUM_HV03 If a sum is given, solve the resulting exponential equation carefully.

WORKED SOLUTION

Question 7

WMA12/01 January 2023

WMA12-01_2023_Jan_Q07

1

Write the given terms

For a geometric sequence, the third term is $ar^2 = 20$ and the fifth term is $ar^4 = 12.8$.

2

Divide the equations

$$\frac{ar^4}{ar^2} = \frac{12.8}{20}, \text{ so } r^2 = 0.64.$$

3

Use r positive

Since $r > 0$, $r = 0.8$.

4

Find a

Using $ar^2 = 20$, $a(0.8)^2 = 20$, so $a = \frac{20}{0.64} = 31.25$.

5

Write the sum formula

$$S_n = \frac{a(1-r^n)}{1-r} = \frac{31.25(1-0.8^n)}{0.2} = 156.25(1-0.8^n).$$

6

Set up the inequality

We need $156.25(1-0.8^n) > 156$.

7

Isolate the power

$$1 - 0.8^n > \frac{156}{156.25} = \frac{624}{625}, \text{ so } 0.8^n < \frac{1}{625}.$$

8

Use logarithms

$n \log(0.8) < \log\left(\frac{1}{625}\right)$. Since $\log(0.8) < 0$, dividing reverses the inequality: $n > \frac{\log(1/625)}{\log(0.8)} = 28.85 \dots$

9

Choose the smallest integer

The smallest possible integer value is therefore $n = 29$.

FINAL ANSWER

(a) $r = 0.8$. (b) $a = 31.25$. (c) $n = 29$.

WMA12-01_2023_Jan_Q07

T06

06_SUM

06_SUM_HV03

CHAPTER

Geometric Sequences

FAMILY

Finite sum

VARIANT

If a sum is given, solve the
resulting exponential
equation carefully.

9 marks

WMA12/01 MAY/JUNE 2023

Question 6

Past-paper question

WMA12-01_2023_MayJune_Q06

6. In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

A software developer released an app to download.

The numbers of downloads of the app each month, in thousands, for the first three months after the app was released were

$$2k - 15 \quad k \quad k + 4$$

where k is a constant.

Given that the numbers of downloads each month are modelled as a geometric series,

(a) show that $k^2 - 7k - 60 = 0$ (2)

(b) predict the number of downloads in the 4th month. (4)

The **total** number of all downloads of the app is predicted to exceed 3 million for the first time in the N th month.

(c) Calculate the value of N according to the model. (3)

T06 Geometric Sequences

06_SUM Finite sum

06_SUM_HV03 If a sum is given, solve the resulting exponential equation carefully.

WORKED SOLUTION

Question 6

WMA12/01 May/June 2023

WMA12-01_2023_MayJune_Q06

1

Use the geometric condition

For three consecutive terms of a geometric sequence, the middle term squared equals the product of the outer terms.

2

Set up the equation

So $k^2 = (2k - 15)(k + 4)$.

3

Show the required quadratic

Expanding gives $k^2 = 2k^2 - 7k - 60$, hence $k^2 - 7k - 60 = 0$.

4

Find k

$k^2 - 7k - 60 = (k - 12)(k + 5) = 0$, so $k = 12$ or $k = -5$. The download numbers must be positive, so $k = 12$.

5

Find the common ratio

The first three monthly values are 9, 12 and 16 thousand. The common ratio is $\frac{12}{9} = \frac{4}{3}$.

6

Predict month four

The 4th month is $16 \cdot \frac{4}{3} = \frac{64}{3}$ thousand downloads, which is about 21300 downloads.

7

Write the cumulative total

In thousands, the total after N months is $S_N = \frac{9\left(\left(\frac{4}{3}\right)^N - 1\right)}{\frac{4}{3} - 1} = 27\left(\left(\frac{4}{3}\right)^N - 1\right)$.

8

Compare with three million

Three million downloads is 3000 thousand, so we need $27\left(\left(\frac{4}{3}\right)^N - 1\right) > 3000$.

9

Solve the inequality

$\left(\frac{4}{3}\right)^N > \frac{1009}{9}$. Taking logarithms gives $N > \frac{\log(1009/9)}{\log(4/3)} = 16.405 \dots$

10

Choose the first month

The first integer month for which the total exceeds 3 million is $N = 17$.

FINAL ANSWER

WMA12-01_2023_MayJune_Q06

(a) **Proven.** (b) $\frac{64}{3}$ **thousand downloads, about 21300 downloads.** (c) $N = 17$.

DR ESLAM AHMED

PURE MATHEMATICS 2

ADAPTIVE ALL CLASSIFIED

Presentation Edition | Solutions

CLASSIFIED BY IDEA | ROOM TO THINK

Every demand appears clearly, with practical space for the mathematics.

Dr Eslam Ahmed

Elite Mathematics

ElitelGCSE.com | 011 2000 9622

PURE 2

PURE 2 | TOPIC 07

Sequences & Series

Your classified route

10

QUESTIONS

68

MARKS

2

FAMILIES

2

VARIANTS

DR ESLAM AHMED

CLASSIFIED BY IDEA

Move from recognition to method, then use the working space to complete the solution.

TOPIC 07 | FAMILY 01

Unknown Count

WHAT THIS FAMILY ASKS

Recognise the unknown count demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

4 classified questions

29 total marks

T07

07_MIX

07_MIX_CORE

CHAPTER

Sequences & Series

FAMILY

Unknown count

VARIANT

core family move

7 marks

WMA12/01 JANUARY 2021

Question 8

Past-paper question

WMA12-01_2021_Jan_Q08

8. A sequence $a_1, a_2, a_3, \dots$ is defined by

$$a_{n+1} = 2(a_n + 3)^2 - 7$$
$$a_1 = p - 3$$

where p is a constant.

- (a) Find an expression for a_2 in terms of p , giving your answer in simplest form.

(1)

Given that $\sum_{n=1}^3 a_n = p + 15$

- (b) find the possible values of a_2

(6)

T07 Sequences & Series

07_MIX Unknown count

07_MIX_CORE core family
move

WORKED SOLUTION

Question 8

WMA12/01 January 2021

WMA12-01_2021_Jan_Q08

1

Find a two

Since $a_1 = p - 3$, we have $a_1 + 3 = p$. Therefore $a_2 = 2(a_1 + 3)^2 - 7 = 2p^2 - 7$.

2

Find a three

$a_2 + 3 = 2p^2 - 4$, so $a_3 = 2(2p^2 - 4)^2 - 7 = 8(p^2 - 2)^2 - 7$.

3

Use the given sum

$a_1 + a_2 + a_3 = p + 15$.

4

Substitute the expressions

$(p - 3) + (2p^2 - 7) + 8(p^2 - 2)^2 - 7 = p + 15$.

5

Simplify

Cancelling p and expanding gives $8p^4 - 30p^2 = 0$, so $2p^2(4p^2 - 15) = 0$.

6

Find the possible a two values

If $p = 0$, then $a_2 = -7$. If $p^2 = \frac{15}{4}$, then $a_2 = 2 \cdot \frac{15}{4} - 7 = \frac{1}{2}$.

FINAL ANSWER

WMA12-01_2021_Jan_Q08

(a) $a_2 = 2p^2 - 7$. (b) $a_2 = -7$ or $\frac{1}{2}$.

T07

07_MIX

07_MIX_CORE

CHAPTER

Sequences & Series

FAMILY

Unknown count

VARIANT

core family move

5 marks

WMA12/01 OCTOBER 2021

Question 2

Past-paper question

WMA12-01_2021_Oct_Q02

2. A sequence is defined by

$$\begin{aligned}u_1 &= 6 \\ u_{n+1} &= ku_n + 3\end{aligned}$$

where k is a positive constant.

- (a) Find, in terms of k , an expression for u_3 (2)

Given that $\sum_{n=1}^3 u_n = 117$

- (b) find the value of k . (3)

T07 Sequences & Series

07_MIX Unknown count

07_MIX_CORE core family
move

WORKED SOLUTION

Question 2

WMA12/01 October 2021

WMA12-01_2021_Oct_Q02

1

Find u two

$$u_2 = ku_1 + 3 = 6k + 3.$$

2

Find u three

$$u_3 = ku_2 + 3 = k(6k + 3) + 3 = 6k^2 + 3k + 3.$$

3

Use the given sum

$$u_1 + u_2 + u_3 = 117, \text{ so } 6 + (6k + 3) + (6k^2 + 3k + 3) = 117.$$

4

Simplify the equation

$$6k^2 + 9k + 12 = 117, \text{ hence } 2k^2 + 3k - 35 = 0.$$

5

Solve for k

$$(2k - 7)(k + 5) = 0. \text{ Since } k \text{ is positive, } k = \frac{7}{2}.$$

FINAL ANSWER

$$(a) u_3 = 6k^2 + 3k + 3. \quad (b) k = \frac{7}{2}.$$

WMA12-01_2021_Oct_Q02

T07

07_MIX

07_MIX_CORE

CHAPTER

Sequences & Series

FAMILY

Unknown count

VARIANT

core family move

8 marks

WMA12/01 MAY/JUNE 2023

Question 11

Past-paper question

WMA12-01_2023_MayJune_Q11

11. A sequence $u_1, u_2, u_3, \dots$ is defined by

$$u_{n+1} = b - au_n$$

$$u_1 = 3$$

where a and b are constants.

(a) Find, in terms of a and b ,

(i) u_2

(ii) u_3

(2)

Given

- $\sum_{n=1}^3 u_n = 153$

- $b = a + 9$

(b) show that

$$a^2 - 5a - 66 = 0$$

(3)

(c) Hence find the larger possible value of u_2

(3)

T07 Sequences & Series

07_MIX Unknown count

07_MIX_CORE core family
move

WORKED SOLUTION

Question 11

WMA12/01 May/June 2023

WMA12-01_2023_MayJune_Q11

1

Find u_2 Using $u_{n+1} = b - au_n$ and $u_1 = 3$, we get $u_2 = b - a(3) = b - 3a$.

2

Find u_3 $u_3 = b - au_2 = b - a(b - 3a) = b - ab + 3a^2$.

3

Use the sum condition $u_1 + u_2 + u_3 = 153$, so $3 + (b - 3a) + (b - ab + 3a^2) = 153$.

4

Substitute b equals a plus nineUsing $b = a + 9$, the left side becomes $3 + (a + 9) - 3a + (a + 9) - a(a + 9) + 3a^2$.

5

SimplifyThis is $2a^2 - 10a + 21$, so $2a^2 - 10a + 21 = 153$.

6

Show the required equation $2a^2 - 10a - 132 = 0$. Dividing by 2 gives $a^2 - 5a - 66 = 0$.

7

Solve for a $a^2 - 5a - 66 = (a - 11)(a + 6) = 0$, so $a = 11$ or $a = -6$.

8

Find the corresponding u_2 valuesIf $a = 11$, then $b = 20$ and $u_2 = 20 - 3(11) = -13$. If $a = -6$, then $b = 3$ and $u_2 = 3 - 3(-6) = 21$.

9

Choose the larger valueThe larger possible value of u_2 is 21.**FINAL ANSWER**

WMA12-01_2023_MayJune_Q11

(a)(i) $u_2 = b - 3a$. (a)(ii) $u_3 = b - ab + 3a^2$. (b) **Proven.** (c) 21.

T07

07_MIX

07_MIX_CORE

CHAPTER

Sequences & Series

FAMILY

Unknown count

VARIANT

core family move

9 marks

WMA12/01 MAY/JUNE 2025

Question 7

Past-paper question

WMA12-01_2025_MayJune_Q07

- 7: **In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.**

- (i) A **geometric** series begins

$$10 + 8 + 6.4 + \dots$$

- (a) Find the sum to infinity of this series.

(2)

Given that the k th term of this series is less than 0.0005

- (b) use algebra to find the smallest possible value of k .

(3)

- (ii) An **arithmetic** series begins

$$850 + 843 + 836 + \dots$$

Given that the sum of the first n terms of this series is S_n
find the greatest possible value of S_n

(4)

T07 Sequences & Series

07_MIX Unknown count

07_MIX_CORE core family
move

WORKED SOLUTION

Question 7

WMA12/01 May/June 2025

WMA12-01_2025_MayJune_Q07

1

Identify the geometric sequenceThe first term is 10 and the common ratio is $\frac{8}{10} = 0.8$.

2

Find the sum to infinity

$$S_{\infty} = \frac{10}{1-0.8} = 50.$$

3

Write the k th termThe k th term is $10(0.8)^{k-1}$.

4

Use the inequality

$$10(0.8)^{k-1} < 0.0005, \text{ so } (0.8)^{k-1} < 0.00005.$$

5

Solve with logarithms

Taking logarithms gives $(k-1)\log(0.8) < \log(0.00005)$. Since $\log(0.8) < 0$, $k-1 > \frac{\log(0.00005)}{\log(0.8)} = 44.381 \dots$

6

Find the smallest integerThus $k > 45.381 \dots$, so the smallest possible value is $k = 46$.

7

Write the arithmetic sumFor $850 + 843 + 836 + \dots$, the first term is 850 and the common difference is -7 .

8

Form S_n

$$S_n = \frac{n}{2}\{2(850) + (n-1)(-7)\} = \frac{n}{2}(1707 - 7n).$$

9

Find the best integer n This quadratic has its maximum near $n = \frac{1707}{14} = 121.928 \dots$, so check $n = 122$.

10

Find the greatest sum

$$S_{122} = \frac{122}{2}(1707 - 854) = 61(853) = 52033.$$

FINAL ANSWER

WMA12-01_2025_MayJune_Q07

(i)(a) 50. (i)(b) 46. (ii) 52033.

DR ESLAM AHMED

TOPIC 07 | FAMILY 02

Sigma Sums

WHAT THIS FAMILY ASKS

Recognise the sigma sums demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

6 classified questions

39 total marks

T07

07_SIGMA

07_SIGMA_CORE

CHAPTER

Sequences & Series

FAMILY

Sigma sums

VARIANT

core family move

7 marks

WMA12/01 JANUARY 2020

Question 8

Past-paper question

WMA12-01_2020_Jan_Q08

8. (i) An arithmetic series has first term a and common difference d .

Prove that the sum to n terms of this series is

$$\frac{n}{2}\{2a + (n-1)d\} \quad (3)$$

- (ii) A sequence $u_1, u_2, u_3, \dots$ is given by

$$u_n = 5n + 3(-1)^n$$

Find the value of

(a) u_5 (1)

(b) $\sum_{n=1}^{59} u_n$ (3)

T07 Sequences & Series

07_SIGMA Sigma sums

07_SIGMA_CORE core family
move

WORKED SOLUTION

Question 8

WMA12/01 January 2020

WMA12-01_2020_Jan_Q08

1

Write the series forwards

$$S_n = a + (a + d) + (a + 2d) + \cdots + (a + (n - 1)d).$$

2

Write the series backwards

$$S_n = (a + (n - 1)d) + (a + (n - 2)d) + \cdots + a.$$

3

Add the two forms

Each pair adds to $2a + (n - 1)d$, and there are n pairs. Hence $2S_n = n\{2a + (n - 1)d\}$.

4

Complete the proof

Dividing by 2 gives $S_n = \frac{n}{2}\{2a + (n - 1)d\}$.

5

Find u_5

$$u_5 = 5(5) + 3(-1)^5 = 25 - 3 = 22.$$

6

Sum the sequence

$$\sum_{n=1}^{59} u_n = 5 \sum_{n=1}^{59} n + 3 \sum_{n=1}^{59} (-1)^n.$$

7

Calculate the total

$5 \sum_{n=1}^{59} n = 5 \cdot \frac{59 \cdot 60}{2} = 8850$, and $\sum_{n=1}^{59} (-1)^n = -1$. Therefore the total is $8850 - 3 = 8847$.

FINAL ANSWER

WMA12-01_2020_Jan_Q08

(i) **Proven.** (ii) (a) 22. (b) 8847.

T07

07_SIGMA

07_SIGMA_CORE

CHAPTER

Sequences & Series

FAMILY

Sigma sums

VARIANT

core family move

7 marks

WMA12/01 OCTOBER 2022

Question 3

Past-paper question

WMA12-01_2022_Oct_Q03

3. A sequence $a_1, a_2, a_3, \dots$ is defined by

$$a_n = \cos^2\left(\frac{n\pi}{3}\right)$$

Find the exact values of

(a) (i) a_1

(ii) a_2

(iii) a_3

(3)

- (b) Hence find the exact value of

$$\sum_{n=1}^{50} \left\{ n + \cos^2\left(\frac{n\pi}{3}\right) \right\}$$

You must make your method clear.

(4)

T07 Sequences & Series

07_SIGMA Sigma sums

07_SIGMA_CORE core family
move

WORKED SOLUTION

Question 3

WMA12/01 October 2022

WMA12-01_2022_Oct_Q03

1

Find the first term

$$a_1 = \cos^2\left(\frac{\pi}{3}\right) = \left(\frac{1}{2}\right)^2 = \frac{1}{4}.$$

2

Find the second term

$$a_2 = \cos^2\left(\frac{2\pi}{3}\right) = \left(-\frac{1}{2}\right)^2 = \frac{1}{4}.$$

3

Find the third term

$$a_3 = \cos^2(\pi) = (-1)^2 = 1.$$

4

Recognise the repeating pattern

The values of $\cos^2\left(\frac{n\pi}{3}\right)$ repeat every 3 terms as $\frac{1}{4}, \frac{1}{4}, 1$.

5

Sum the n part

$$\sum_{n=1}^{50} n = \frac{50 \cdot 51}{2} = 1275.$$

6

Sum the cosine part

There are 16 complete groups of 3 terms in the first 48 terms, then the first 2 terms repeat. Hence the cosine-square sum is $16\left(\frac{1}{4} + \frac{1}{4} + 1\right) + \frac{1}{4} + \frac{1}{4} = 24 + \frac{1}{2} = \frac{49}{2}$.

7

Add the two sums

$$\sum_{n=1}^{50} \left(n + \cos^2\left(\frac{n\pi}{3}\right)\right) = 1275 + \frac{49}{2} = \frac{2599}{2}.$$

FINAL ANSWER

WMA12-01_2022_Oct_Q03

$$(a) a_1 = \frac{1}{4}, a_2 = \frac{1}{4}, a_3 = 1. \quad (b) \frac{2599}{2}.$$

T07

07_SIGMA

07_SIGMA_CORE

CHAPTER

Sequences & Series

FAMILY

Sigma sums

VARIANT

core family move

5 marks

WMA12/01 OCTOBER 2023

Question 2

Past-paper question

WMA12-01_2023_Oct_Q02

2. A sequence $u_1, u_2, u_3, \dots$ is defined by

$$u_1 = 3$$

$$u_{n+1} = 2 - \frac{4}{u_n}$$

- (a) Find the value of u_2 , the value of u_3 and the value of u_4

(3)

- (b) Find the value of

$$\sum_{r=1}^{100} u_r$$

(2)

T07 Sequences & Series

07_SIGMA Sigma sums

07_SIGMA_CORE core family
move

WORKED SOLUTION

Question 2

WMA12/01 October 2023

WMA12-01_2023_Oct_Q02

1

Find u_2

$$u_2 = 2 - \frac{4}{u_1} = 2 - \frac{4}{3} = \frac{2}{3}.$$

2

Find u_3

$$u_3 = 2 - \frac{4}{u_2} = 2 - \frac{4}{2/3} = 2 - 6 = -4.$$

3

Find u_4

$$u_4 = 2 - \frac{4}{u_3} = 2 - \frac{4}{-4} = 3.$$

4

Notice the cycle

The sequence starts $3, \frac{2}{3}, -4, 3, \frac{2}{3}, -4, \dots$, so it repeats every 3 terms.

5

Sum one cycle

$$3 + \frac{2}{3} - 4 = -\frac{1}{3}.$$

6

Use 100 terms

There are 33 complete cycles in the first 99 terms, then one extra term $u_{100} = 3$.

7

Find the sum

$$\sum_{r=1}^{100} u_r = 33 \left(-\frac{1}{3} \right) + 3 = -11 + 3 = -8.$$

FINAL ANSWER

WMA12-01_2023_Oct_Q02

(a) $u_2 = \frac{2}{3}$, $u_3 = -4$, $u_4 = 3$. (b) -8 .

T07

07_SIGMA

07_SIGMA_CORE

CHAPTER

Sequences & Series

FAMILY

Sigma sums

VARIANT

core family move

8 marks

WMA12/01 JANUARY 2024

Question 5

Past-paper question

WMA12-01_2024_Jan_Q05

5. (i) Find the value of

$$\sum_{r=1}^{\infty} 6 \times (0.25)^r \quad (3)$$

- (ii) A sequence $u_1, u_2, u_3, \dots$ is defined by

$$\begin{aligned} u_1 &= 3 \\ u_{n+1} &= \frac{u_n - 3}{u_n - 2} \quad n \in \mathbb{N} \end{aligned}$$

- (a) Show that this sequence is periodic. (2)
- (b) State the order of this sequence. (1)
- (c) Hence find

$$\sum_{n=1}^{70} u_n \quad (2)$$

T07 Sequences & Series

07_SIGMA Sigma sums

07_SIGMA_CORE core family
move

WORKED SOLUTION

Question 5

WMA12/01 January 2024

WMA12-01_2024_Jan_Q05

1

Identify the geometric series $\sum_{r=1}^{\infty} 6(0.25)^r$ has first term $6(0.25) = 1.5$ and common ratio 0.25.

2

Sum to infinity

$$S_{\infty} = \frac{1.5}{1-0.25} = \frac{1.5}{0.75} = 2.$$

3

Find the first few terms

$$u_1 = 3. \text{ Then } u_2 = \frac{3-3}{3-2} = 0.$$

4

Continue the recurrence

$$u_3 = \frac{0-3}{0-2} = \frac{3}{2}, \text{ and } u_4 = \frac{\frac{3}{2}-3}{\frac{3}{2}-2} = 3.$$

5

Show periodicitySince $u_4 = 3 = u_1$, the terms repeat as $3, 0, \frac{3}{2}, 3, 0, \frac{3}{2}, \dots$

6

State the order

The repeating cycle has length 3, so the order is 3.

7

Sum one cycle

$$\text{One full cycle sums to } 3 + 0 + \frac{3}{2} = \frac{9}{2}.$$

8

Use 70 termsSince $70 = 23 \cdot 3 + 1$, the first 70 terms contain 23 complete cycles and one extra term $u_1 = 3$.

9

Find the required sum

$$\sum_{n=1}^{70} u_n = 23 \left(\frac{9}{2} \right) + 3 = \frac{207}{2} + \frac{6}{2} = \frac{213}{2}.$$

FINAL ANSWER

WMA12-01_2024_Jan_Q05

(i) 2. (ii)(a) **Periodic.** (ii)(b) **Order 3.** (ii)(c) $\frac{213}{2}$.

T07

07_SIGMA

07_SIGMA_CORE

CHAPTER

Sequences & Series

FAMILY

Sigma sums

VARIANT

core family move

5 marks

WMA12/01 MAY/JUNE (R) 2024

Question 1

Past-paper question

WMA12-01_2024_MayJuneR_Q01

1. The sequence $u_1, u_2, u_3, \dots$ satisfies

$$u_{n+2} = 3u_{n+1} - 2u_n$$

Given that

- $u_1 = 7$
- $u_3 = 4$

- (a) find the value of u_2

(2)

- (b) find $\sum_{r=1}^4 (u_r + 2r)$

(3)

T07 Sequences & Series

07_SIGMA Sigma sums

07_SIGMA_CORE core family
move

WORKED SOLUTION

Question 1

WMA12/01 May/June (R) 2024

WMA12-01_2024_MayJuneR_Q01

1

Use the recurrence at n equals one

$$u_3 = 3u_2 - 2u_1.$$

2

Substitute given values

$$4 = 3u_2 - 2(7), \text{ so } 4 = 3u_2 - 14.$$

3

Find u_2

$$3u_2 = 18, \text{ hence } u_2 = 6.$$

4

Find u_4

$$u_4 = 3u_3 - 2u_2 = 3(4) - 2(6) = 0.$$

5

Evaluate the sum

$$\sum_{r=1}^4 (u_r + 2r) = (7 + 2) + (6 + 4) + (4 + 6) + (0 + 8).$$

6

Add the terms

$$\text{The sum is } 9 + 10 + 10 + 8 = 37.$$

FINAL ANSWER

$$(a) u_2 = 6. \quad (b) 37.$$

WMA12-01_2024_MayJuneR_Q01

T07

07_SIGMA

07_SIGMA_CORE

CHAPTER

Sequences & Series

FAMILY

Sigma sums

VARIANT

core family move

7 marks

WMA12/01 OCTOBER 2024

Question 2

Past-paper question

WMA12-01_2024_Oct_Q02

2. A sequence of numbers $u_1, u_2, u_3, \dots$ is defined by

$$u_1 = 7$$

$$u_{n+1} = (-1)^n u_n + k$$

where k is a constant.

- (a) Show that $u_5 = 7$

(3)

Given that $\sum_{r=1}^4 u_r = 30$

- (b) find the value of k .

(2)

- (c) Hence find the value of $\sum_{r=1}^{150} u_r$

(2)

T07 Sequences & Series

07_SIGMA Sigma sums

07_SIGMA_CORE core family
move

WORKED SOLUTION

Question 2

WMA12/01 October 2024

WMA12-01_2024_Oct_Q02

1

Find the first few terms

$$u_1 = 7, u_2 = (-1)^1 u_1 + k = k - 7, u_3 = (-1)^2 u_2 + k = 2k - 7, \text{ and } u_4 = (-1)^3 u_3 + k = 7 - k.$$

2

Show u five

$$u_5 = (-1)^4 u_4 + k = (7 - k) + k = 7.$$

3

Use the given sum

$$\sum_{r=1}^4 u_r = 7 + (k - 7) + (2k - 7) + (7 - k) = 2k.$$

4

Find k

$$\text{Since } \sum_{r=1}^4 u_r = 30, \text{ we have } 2k = 30, \text{ so } k = 15.$$

5

Write one repeated block

With $k = 15$, the first four terms are 7, 8, 23, -8 , and their sum is 30. Since $u_5 = u_1$, the pattern repeats every four terms.

6

Sum 150 terms

There are 37 complete blocks in the first 150 terms, with the first two terms left over. Hence $\sum_{r=1}^{150} u_r = 37(30) + 7 + 8 = 1125$.

FINAL ANSWER

WMA12-01_2024_Oct_Q02

(a) Proven. (b) $k = 15$. (c) 1125.

PURE MATHEMATICS 2

ADAPTIVE ALL CLASSIFIED

Presentation Edition | Solutions

CLASSIFIED BY IDEA | ROOM TO THINK

Every demand appears clearly, with practical space for the mathematics.

Dr Eslam Ahmed

Elite Mathematics

ElitelGCSE.com | 011 2000 9622

PURE 2

PURE 2 | TOPIC 08

Modelling with Sequences & Series

Your classified route

10

QUESTIONS

83

MARKS

4

FAMILIES

5

VARIANTS

DR ESLAM AHMED

CLASSIFIED BY IDEA

Move from recognition to method, then use the working space to complete the solution.

TOPIC 08 | FAMILY 01

Model Fitting

WHAT THIS FAMILY ASKS

Recognise the model fitting demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

2 classified questions

17 total marks

T08

08_FIT

08_FIT_CORE

CHAPTER

Modelling with Sequences & Series

FAMILY

Model fitting

VARIANT

core family move

9 marks

WMA12/01 MAY/JUNE 2022

Question 9

Past-paper question

WMA12-01_2022_MayJune_Q09

9. A scientist is using carbon-14 dating to determine the age of some wooden items.

The equation for carbon-14 dating an item is given by

$$N = k\lambda^t$$

where

- N grams is the amount of carbon-14 **currently** present in the item
- k grams was the **initial** amount of carbon-14 present in the item
- t is the number of years since the item was made
- λ is a constant, with $0 < \lambda < 1$

- (a) Sketch the graph of N against t for $k = 1$

(2)

Given that it takes 5700 years for the amount of carbon-14 to reduce to half its initial value,

- (b) show that the value of the constant λ is 0.999878 to 6 decimal places.

(2)

Given that Item A

- is known to have had 15 grams of carbon-14 present initially
- is thought to be 3250 years old

- (c) calculate, to 3 significant figures, how much carbon-14 the equation predicts is currently in Item A .

(2)

Item B is known to have initially had 25 grams of carbon-14 present, but only 18 grams now remain.

- (d) Use algebra to calculate the age of Item B to the nearest 100 years.

(3)

T08 Modelling with
Sequences & Series

08_FIT Model fitting

08_FIT_CORE core family
move

WORKED SOLUTION

Question 9

WMA12/01 May/June 2022

WMA12-01_2022_MayJune_Q09

1

Sketch the model for k equals one

For $k = 1$, $N = \lambda^t$ with $0 < \lambda < 1$. The graph starts at $(0, 1)$, decreases as t increases, and tends towards 0.

2

Use the half-life information

After 5700 years, $N = \frac{k}{2}$, so $k\lambda^{5700} = \frac{k}{2}$.

3

Find lambda

$\lambda^{5700} = \frac{1}{2}$, hence $\lambda = \left(\frac{1}{2}\right)^{1/5700} = 0.999878 \dots$, so $\lambda = 0.999878$ to 6 decimal places.

4

Model item A

For item A, $k = 15$ and $t = 3250$, so $N = 15\lambda^{3250}$.

5

Calculate item A

$N = 15(0.9998784026 \dots)^{3250} = 10.103 \dots$, so the current amount is 10.1 g to 3 significant figures.

6

Set up item B

For item B, $18 = 25\lambda^t$, so $\lambda^t = \frac{18}{25}$.

7

Take logarithms

$t \log \lambda = \log \left(\frac{18}{25}\right)$, so $t = \frac{\log(18/25)}{\log \lambda}$.

8

Round the age

Using $\lambda = (1/2)^{1/5700}$ gives $t = 2701 \dots$, so the age is 2700 years to the nearest 100 years.

FINAL ANSWER

WMA12-01_2022_MayJune_Q09

(a) Decreasing exponential through $(0, 1)$. (b) $\lambda = 0.999878$.
(c) 10.1 g. (d) 2700 years.

T08

08_FIT

08_FIT_CORE

CHAPTER

Modelling with Sequences & Series

FAMILY

Model fitting

VARIANT

core family move

8 marks

WMA12/01 MAY/JUNE 2024

Question 10

Past-paper question

WMA12-01_2024_MayJune_Q10

10. **In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.**

The number of dormice and the number of voles on an island are being monitored.

Initially there are 2000 dormice on the island.

A model predicts that the number of dormice will increase by 3% each year, so that the numbers of dormice on the island at the end of each year form a geometric sequence.

- (a) Find, according to the model, the number of dormice on the island 6 years after monitoring began. Give your answer to 3 significant figures.

(2)

The number of voles on the island is being monitored over the same period of time.

Given that

- 4 years after monitoring began there were 3690 voles on the island
- 7 years after monitoring began there were 3470 voles on the island
- the number of voles on the island at the end of each year is modelled as a geometric sequence

- (b) find the equation of this model in the form

$$N = ab^t$$

where N is the number of voles, t years after monitoring began and a and b are constants. Give the value of a and the value of b to 2 significant figures.

(3)

When $t = T$, the number of dormice on the island is equal to the number of voles on the island.

- (c) Find, according to the models, the value of T , giving your answer to one decimal place.

(3)

T08 Modelling with
Sequences & Series

08_FIT Model fitting

08_FIT_CORE core family
move

WORKED SOLUTION

Question 10

WMA12/01 May/June 2024

WMA12-01_2024_MayJune_Q10

1

Model the dormice

The dormice model is $D = 2000(1.03)^t$.

2

Find the dormice after six years

$D = 2000(1.03)^6 = 2388.10 \dots$, so there are 2390 dormice to 3 significant figures.

3

Set up the vole model

Let the vole model be $N = ab^t$.

4

Use the two data points

At $t = 4$, $ab^4 = 3690$. At $t = 7$, $ab^7 = 3470$.

5

Find b

Dividing gives $b^3 = \frac{3470}{3690}$, so $b = 0.9797 \dots$, which is 0.98 to 2 significant figures.

6

Find a

$a = \frac{3690}{b^4} = 4005.18 \dots$, which is 4000 to 2 significant figures.

7

Write the vole model

So the model is approximately $N = 4000(0.98)^t$.

8

Set the two models equal

Use $2000(1.03)^T = ab^T$ with the unrounded constants from the vole model.

9

Solve using logarithms

$\left(\frac{1.03}{b}\right)^T = \frac{a}{2000}$, so $T = \frac{\log(a/2000)}{\log(1.03/b)}$.

10

Calculate T

Substituting $a = 4005.18 \dots$ and $b = 0.9797 \dots$ gives $T = 13.875 \dots$, so $T = 13.9$ to one decimal place.

FINAL ANSWER

WMA12-01_2024_MayJune_Q10

(a) 2390 **dormice.** (b) $N = 4000(0.98)^t$ **to 2 s.f. constants.** (c) $T = 13.9.$

DR ESLAM AHMED

TOPIC 08 | FAMILY 02

Interpretation

WHAT THIS FAMILY ASKS

Recognise the interpretation demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 For 'at least' or 'exceeds', test the boundary integer.

V02 If the model starts at $n=0$, translate n into calendar time.

3 classified questions

24 total marks

T08

08_INTERP

08_INTERP_HV01

CHAPTER

Modelling with Sequences & Series

FAMILY

Interpretation

VARIANT

For 'at least' or 'exceeds', test the boundary integer.**8 marks**

WMA12/01 JANUARY 2020

Question 5

Past-paper question

WMA12-01_2020_Jan_Q05

5. In this question you must show all stages of your working.

Solutions relying on calculator technology are not acceptable.

A colony of bees is being studied.

The number of bees in the colony at the start of the study was 30 000

Three years after the start of the study, the number of bees in the colony is 34 000

A model predicts that the number of bees in the colony will increase by $p\%$ each year, so that the number of bees in the colony at the end of each year of study forms a geometric sequence.

Assuming the model,

(a) find the value of p , giving your answer to 2 decimal places.

(3)

According to the model, at the end of N years of study the number of bees in the colony exceeds 75 000

(b) Find, showing all steps in your working, the smallest integer value of N .

(5)

T08 Modelling with
Sequences & Series

08_INTERP Interpretation

08_INTERP_HV01 For 'at
least' or 'exceeds', test the
boundary integer.

WORKED SOLUTION

Question 5

WMA12/01 January 2020

WMA12-01_2020_Jan_Q05

1

Set up the growth ratio

Let the yearly ratio be r . After 3 years, $30000r^3 = 34000$.

2

Find p

$r^3 = \frac{17}{15}$, so $r = \left(\frac{17}{15}\right)^{1/3} \approx 1.04257$. Therefore $p = (r - 1) \times 100 \approx 4.26$.

3

Set up the inequality

After N years the model gives $30000r^N$. We need $30000r^N > 75000$, so $r^N > 2.5$.

4

Take logarithms

$N \log r > \log(2.5)$, so $N > \frac{\log(2.5)}{\log r}$.

5

Find the smallest integer

Using $r \approx 1.04257$, $\frac{\log(2.5)}{\log r} \approx 21.98$. Hence the smallest integer value is $N = 22$.

FINAL ANSWER

WMA12-01_2020_Jan_Q05

(a) $p = 4.26$. (b) $N = 22$.

T08

08_INTERP

08_INTERP_HV01

CHAPTER

Modelling with Sequences & Series

FAMILY

Interpretation

VARIANT

For 'at least' or 'exceeds', test the boundary integer.

9 marks

WMA12/01 JANUARY 2022

Question 8

Past-paper question

WMA12-01_2022_Jan_Q08

8. A metal post is repeatedly hit in order to drive it into the ground.

Given that

- on the 1st hit, the post is driven 100 mm into the ground
- on the 2nd hit, the post is driven an **additional** 98 mm into the ground
- on the 3rd hit, the post is driven an **additional** 96 mm into the ground
- the **additional** distances the post travels on each subsequent hit form an arithmetic sequence

- (a) show that the post is driven an **additional** 62 mm into the ground with the 20th hit.

(1)

- (b) Find the **total distance** that the post has been driven into the ground after 20 hits.

(2)

Given that for each subsequent hit after the 20th hit

- the **additional** distances the post travels form a geometric sequence with common ratio r
- on the 22nd hit, the post is driven an **additional** 60 mm into the ground

- (c) find the value of r , giving your answer to 3 decimal places.

(2)

After a total of N hits, the post will have been driven more than 3 m into the ground.

- (d) Find, showing all steps in your working, the smallest possible value of N .

(4)

T08 Modelling with
Sequences & Series

08_INTERP Interpretation

08_INTERP_HV01 For 'at
least' or 'exceeds', test the
boundary integer.

WORKED SOLUTION

Question 8

WMA12/01 January 2022

WMA12-01_2022_Jan_Q08

1

Use the arithmetic rule

For the first 20 hits, the additional distances form an arithmetic sequence with first term 100 and common difference -2 .

2

Show the twentieth hit

The twentieth additional distance is $100 + 19(-2) = 62$ mm.

3

Sum the first twenty hits

$$S_{20} = \frac{20}{2}(100 + 62) = 1620 \text{ mm.}$$

4

Use the geometric information

After the twentieth hit, the distances follow the same ratio, so the twenty second distance is $62r^2$.

5

Find r

$$62r^2 = 60, \text{ hence } r = \sqrt{\frac{30}{31}} = 0.984 \text{ to 3 decimal places.}$$

6

Set up the remaining distance

More than 3 m means more than 3000 mm. After 20 hits, the remaining amount to exceed is $3000 - 1620 = 1380$ mm.

7

Sum the geometric tail

$$\text{If there are } m \text{ further hits, their total is } 62r + 62r^2 + \dots + 62r^m = 62r \frac{1-r^{m+1}}{1-r}.$$

8

Find the first m that works

Using $r = \sqrt{30/31}$, $62r \frac{1-r^{27}}{1-r}$ is not enough, but $62r \frac{1-r^{28}}{1-r}$ makes the total exceed 3000 mm.

9

State N

Therefore $m = 28$ further hits are needed after the first 20, so $N = 20 + 28 = 48$.

FINAL ANSWER

WMA12-01_2022_Jan_Q08

(a) **Proven:** 62 mm. (b) 1620 mm. (c) $r = 0.984$. (d) $N = 48$.

DR ESLAM AHMED

T08

08_INTERP

08_INTERP_HV02

CHAPTER

Modelling with Sequences & Series

FAMILY

Interpretation

VARIANT

If the model starts at $t(n=0)$,
translate $t(n)$ into calendar
time.

7 marks

WMA12/01 MAY/JUNE 2021

Question 1

Past-paper question

WMA12-01_2021_MayJune_Q01

1. Adina is saving money to buy a new computer. She saves £5 in week 1, £5.25 in week 2, £5.50 in week 3 and so on until she has enough money, in total, to buy the computer.

She decides to model her savings using either an arithmetic series or a geometric series.

Using the information given,

- (a) (i) state with a reason whether an arithmetic series or a geometric series should be used,
- (ii) write down an expression, in terms of n , for the amount, in pounds (£), saved in week n .

(3)

Given that the computer Adina wants to buy costs £350

- (b) find the number of weeks it will take for Adina to save enough money to buy the computer.

(4)

T08 Modelling with
Sequences & Series

08_INTERP Interpretation

08_INTERP_HV02 If the model
starts at $t=0$, translate $t(n)$
into calendar time.

WORKED SOLUTION

Question 1

WMA12/01 May/June 2021

WMA12-01_2021_MayJune_Q01

1

Choose the model

The weekly amounts are 5, 5.25, 5.50, and so on. The increase is a constant 0.25, so an arithmetic series should be used.

2

Write the n th amount

The first amount is 5 and the common difference is 0.25, so the amount saved in week n is $5 + 0.25(n - 1)$.

3

Set up the total

After n weeks, $S_n = \frac{n}{2}\{2(5) + (n - 1)(0.25)\}$.

4

Form the inequality

Adina needs $S_n \geq 350$, so $\frac{n}{2}(9.75 + 0.25n) \geq 350$.

5

Solve for n

This is $n^2 + 39n - 2800 \geq 0$. The positive root is 36.89 ..., so the first whole number that works is $n = 37$.

6

Check the week

After 36 weeks the total is below 350, and after 37 weeks it is 351.50, so it takes 37 weeks.

FINAL ANSWER

WMA12-01_2021_MayJune_Q01

(a)(i) **Arithmetic series.** (a)(ii) $5 + 0.25(n - 1)$. (b) **37 weeks.**

TOPIC 08 | FAMILY 03

Limiting Models

WHAT THIS FAMILY ASKS

Recognise the limiting models demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

2 classified questions

21 total marks

T08

08_LIMIT

08_LIMIT_CORE

CHAPTER

Modelling with Sequences & Series

FAMILY

Limiting models

VARIANT

core family move**11 marks**

WMA12/01 JANUARY 2021

Question 10

Past-paper question

WMA12-01_2021_Jan_Q10

10. In this question you must show detailed reasoning.

Owen wants to train for 12 weeks in preparation for running a marathon.

During the 12-week period he will run every Sunday and every Wednesday.

- On Sunday in week 1 he will run 15 km
- On Sunday in week 12 he will run 37 km

He considers two different 12-week training plans.

In training plan *A*, he will increase the distance he runs each Sunday by the same amount.

- (a) Calculate the distance he will run on Sunday in week 5 under training plan *A*.

(3)

In training plan *B*, he will increase the distance he runs each Sunday by the same percentage.

- (b) Calculate the distance he will run on Sunday in week 5 under training plan *B*.
Give your answer in km to one decimal place.

(3)

Owen will also run a fixed distance, x km, each Wednesday over the 12-week period.

Given that

- x is an integer
- the total distance that Owen will run on Sundays and Wednesdays over the 12 weeks will not exceed 360 km

- (c) (i) find the maximum value of x , if he uses training plan *A*,
(ii) find the maximum value of x , if he uses training plan *B*.

(5)**(Total 11 marks)**

T08 Modelling with
Sequences & Series

08_LIMIT Limiting models

08_LIMIT_CORE core family
move

WORKED SOLUTION

Question 10

WMA12/01 January 2021

WMA12-01_2021_Jan_Q10

1

Find the arithmetic difference

Plan A has 12 Sunday runs from 15 km to 37 km, so there are 11 equal increases. The common difference is $\frac{37-15}{11} = 2$.

2

Find week five for plan A

Week 5 is the fifth term, so the distance is $15 + 4(2) = 23$ km.

3

Find the geometric ratio

For plan B, $15r^{11} = 37$, so $r = \left(\frac{37}{15}\right)^{1/11}$.

4

Find week five for plan B

The fifth Sunday distance is $15r^4 = 15\left(\frac{37}{15}\right)^{4/11} = 20.829 \dots$, so it is 20.8 km.

5

Sum the Sunday runs for plan A

The arithmetic total is $\frac{12}{2}(15 + 37) = 312$ km.

6

Find the Wednesday limit for plan A

The total distance condition gives $312 + 12x \leq 360$, so $x \leq 4$. The maximum integer is 4.

7

Sum the Sunday runs for plan B

The geometric total is $15\frac{r^{12}-1}{r-1} = 294.185 \dots$ km.

8

Find the Wednesday limit for plan B

Now $294.185 \dots + 12x \leq 360$, so $x \leq 5.484 \dots$. The maximum integer is 5.

FINAL ANSWER

WMA12-01_2021_Jan_Q10

(a) 23 km. (b) 20.8 km. (c)(i) $x = 4$. (c)(ii) $x = 5$.

T08

08_LIMIT

08_LIMIT_CORE

CHAPTER

Modelling with Sequences &
Series

FAMILY

Limiting models

VARIANT

core family move

10 marks

WMA12/01 JANUARY 2026

Question 9

Past-paper question

WMA12-01_2026_Jan_Q09

9. A company mines lithium.

- in year 1, the company mined 12 400 tonnes of lithium
- in year 15, the company is predicted to mine 7 500 tonnes of lithium

Model *A* assumes that the mass of lithium the company mines will decrease by the same amount each year.

According to model *A*,

(a) find the mass of lithium the company will mine in year 5 (3)

(b) calculate the **total** mass of lithium the company will mine from year 1 to year 15 inclusive. (2)

Model *B* assumes that the mass of lithium the company mines will decrease by the same percentage each year.

(c) Find, according to model *B*, the mass of lithium the company will mine in year 10. Give the answer to the nearest 10 tonnes. (3)

According to model *B*, there is a limit to the **total** mass of lithium that can be mined.

(d) Calculate the value of this limit. Give the answer to the nearest 100 tonnes. (2)

T08 Modelling with
Sequences & Series

08_LIMIT Limiting models

08_LIMIT_CORE core family
move

WORKED SOLUTION

Question 9

WMA12/01 January 2026

WMA12-01_2026_Jan_Q09

1

Model A as an arithmetic sequence

For Model A, let the common difference be d . Since year 15 is the 15th term, $7500 = 12400 + 14d$.

2

Find year 5 for Model A

$d = \frac{7500-12400}{14} = -350$. Year 5 is $12400 + 4(-350) = 11000$ tonnes.

3

Find the total from year 1 to year 15

$S_{15} = \frac{15}{2}(12400 + 7500) = 149250$ tonnes.

4

Model B as a geometric sequence

Let the common ratio be r . Then $7500 = 12400r^{14}$, so $r = \left(\frac{7500}{12400}\right)^{1/14} \approx 0.964723$.

5

Find year 10 for Model B

Year 10 is $12400r^9 \approx 8975.26$, which is 8980 tonnes to the nearest 10 tonnes.

6

Find the limiting total

Since $0 < r < 1$, the sum to infinity is $S_{\infty} = \frac{12400}{1-r} \approx 351508.1$, so the limit is 351500 tonnes to the nearest 100 tonnes.

FINAL ANSWER

WMA12-01_2026_Jan_Q09

(a) 11000 tonnes. (b) 149250 tonnes. (c) 8980 tonnes. (d) 351500 tonnes.

TOPIC 08 | FAMILY 04

Model Setup

WHAT THIS FAMILY ASKS

Recognise the model setup demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

3 classified questions

21 total marks

T08

08_MODEL

08_MODEL_CORE

CHAPTER

Modelling with Sequences & Series

FAMILY

Model setup

VARIANT

core family move**6 marks**

WMA12/01 OCTOBER 2019

Question 2

Past-paper question

WMA12-01_2019_Oct_Q02

2. The adult population of a town at the start of 2019 is 25 000

A model predicts that the adult population will increase by 2% each year, so that the number of adults in the population at the start of each year following 2019 will form a geometric sequence.

- (a) Find, according to the model, the adult population of the town at the start of 2032 (3)

It is also modelled that every member of the adult population gives £5 to local charity at the start of each year.

- (b) Find, according to these models, the total amount of money that would be given to local charity by the adult population of the town from 2019 to 2032 inclusive. Give your answer to the nearest £1 000 (3)

T08 Modelling with
Sequences & Series

08_MODEL Model setup

08_MODEL_CORE core
family move

WORKED SOLUTION

Question 2

WMA12/01 October 2019

WMA12-01_2019_Oct_Q02

1

Identify the common ratio

An increase of 2% each year gives the geometric ratio $r = 1.02$.

2

Count the years to 2032

From the start of 2019 to the start of 2032 there are 13 yearly increases, so the population is $25000(1.02)^{13}$.

3

Calculate the 2032 population

$25000(1.02)^{13} \approx 32340.7$, so the model gives about 32341 adults.

4

Set up the total number of adult payments

From 2019 to 2032 inclusive there are 14 yearly populations, so the total number of adult payments is $25000 \frac{1.02^{14}-1}{1.02-1}$.

5

Multiply by the charity amount

The total money is $5 \times 25000 \frac{1.02^{14}-1}{0.02} \approx 1\,996\,766$, so this is about £1,996,766.

6

Round as requested

To the nearest £1000, this is £1,997,000.

FINAL ANSWER

(a) **32341 adults, approximately.** (b) **£1,997,000.**

WMA12-01_2019_Oct_Q02

T08

08_MODEL

08_MODEL_CORE

CHAPTER

Modelling with Sequences & Series

FAMILY

Model setup

VARIANT

core family move**9 marks****WMA12/01 JANUARY 2024****Question 7**

Past-paper question

WMA12-01_2024_Jan_Q07

7. Wheat is grown on a farm.

- In year 1, the farm produced 300 tonnes of wheat.
- In year 12, the farm is predicted to produce 4 000 tonnes of wheat.

Model *A* assumes that the amount of wheat produced on the farm will increase by the same amount each year.

- (a) Using model *A*, find the amount of wheat produced on the farm in year 4.
Give your answer to the nearest 10 tonnes.

(3)

Model *B* assumes that the amount of wheat produced on the farm will increase by the same percentage each year.

- (b) Using model *B*, find the amount of wheat produced on the farm in year 2.
Give your answer to the nearest 10 tonnes.

(3)

- (c) Calculate, according to the two models, the difference between the total amounts of wheat predicted to be produced on the farm from year 1 to year 12 inclusive.
Give your answer to the nearest 10 tonnes.

(3)

T08 Modelling with
Sequences & Series

08_MODEL Model setup

08_MODEL_CORE core
family move

WORKED SOLUTION

Question 7

WMA12/01 January 2024

WMA12-01_2024_Jan_Q07

1

Model A common difference

Model A is arithmetic. From year 1 to year 12 there are 11 equal increases, so $d = \frac{4000-300}{11} = \frac{3700}{11}$.

2

Find year 4 in model A

Year 4 is $300 + 3d = 300 + 3\left(\frac{3700}{11}\right) = 1309.09 \dots$, so the amount is 1310 tonnes to the nearest 10 tonnes.

3

Model B common ratio

Model B is geometric. If the ratio is r , then $300r^{11} = 4000$.

4

Find the ratio

$$r = \left(\frac{4000}{300}\right)^{1/11} = 1.2655 \dots$$

5

Find year 2 in model B

Year 2 is $300r = 379.65 \dots$, so the amount is 380 tonnes to the nearest 10 tonnes.

6

Total for model A

$$S_A = \frac{12}{2}(300 + 4000) = 25800 \text{ tonnes.}$$

7

Total for model B

$$S_B = 300 \frac{r^{12}-1}{r-1} = 17935.20 \dots \text{ tonnes.}$$

8

Find the difference

The difference is $25800 - 17935.20 \dots = 7864.79 \dots$, which is 7860 tonnes to the nearest 10 tonnes.

FINAL ANSWER

(a) 1310 tonnes. (b) 380 tonnes. (c) 7860 tonnes.

WMA12-01_2024_Jan_Q07

T08

08_MODEL

08_MODEL_CORE

CHAPTER

Modelling with Sequences & Series

FAMILY

Model setup

VARIANT

core family move**6 marks**

WMA12/01 OCTOBER 2024

Question 7

Past-paper question

WMA12-01_2024_Oct_Q07

7. Jem pays money into a savings scheme, A , over a period of 300 months.

Jem pays £20 into scheme A in month 1, £20.50 in month 2, £21 in month 3 and so on, so that the amounts Jem pays each month form an arithmetic sequence.

- (a) Show that Jem pays £69.50 into scheme A in month 100 (1)

- (b) Find the **total** amount that Jem pays into scheme A over the period of 300 months. (2)

Kim pays money into a different savings scheme, B , over the same period of 300 months.

In a model, the amounts Kim pays into scheme B increase by the same percentage each month, so that the amounts Kim pays each month form a geometric sequence.

Given that Kim pays

- £20 into scheme B in month 1
- £250 into scheme B in month 300

- (c) use the model to calculate, to the nearest £10, the difference between the total amount paid into scheme A and the total amount paid into scheme B over the period of 300 months. (3)

T08 Modelling with
Sequences & Series

08_MODEL Model setup

08_MODEL_CORE core
family move

WORKED SOLUTION

Question 7

WMA12/01 October 2024

WMA12-01_2024_Oct_Q07

1

Identify Jem's sequence

Jem's payments are arithmetic with first term 20 and common difference 0.50.

2

Show month 100

The month 100 payment is $20 + 99(0.50) = 69.50$, so Jem pays £69.50.

3

Sum Jem's payments

$$S_{300} = \frac{300}{2} \{2(20) + 299(0.50)\} = 150(189.5) = 28425.$$

4

Model Kim's payments

Let the common ratio be r . Since the first payment is 20 and the month 300 payment is 250, $20r^{299} = 250$.

5

Find the ratio

Thus $r^{299} = 12.5$ and $r = 12.5^{1/299}$.

6

Sum Kim's payments

$$S_B = 20 \frac{r^{300} - 1}{r - 1} \text{ with } r = 12.5^{1/299}, \text{ giving } S_B \approx 27362.95.$$

7

Find the difference

The difference is $28425 - 27362.95 \approx 1062.05$, which is £1,060 to the nearest £10.

FINAL ANSWER

(a) Proven. (b) £28,425. (c) £1,060.

WMA12-01_2024_Oct_Q07

PURE MATHEMATICS 2

ADAPTIVE ALL CLASSIFIED

Presentation Edition | Solutions

CLASSIFIED BY IDEA | ROOM TO THINK

Every demand appears clearly, with practical space for the mathematics.

Dr Eslam Ahmed

Elite Mathematics

ElitelGCSE.com | 011 2000 9622

PURE 2

PURE 2 | TOPIC 09

Laws of Logarithms

Your classified route

20

QUESTIONS

148

MARKS

4

FAMILIES

5

VARIANTS

DR ESLAM AHMED

CLASSIFIED BY IDEA

Move from recognition to method, then use the working space to complete the solution.

TOPIC 09 | FAMILY 01

Change Of Base

WHAT THIS FAMILY ASKS

Recognise the change of base demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE**V01** core family move**1 classified questions**

7 total marks

T09

09_BASE

09_BASE_CORE

CHAPTER
Laws of Logarithms

FAMILY
Change of base

VARIANT
core family move

7 marks

WMA12/01 OCTOBER 2022

Question 10

Past-paper question

WMA12-01_2022_Oct_Q10

10. Given $a = \log_2 3$

(i) write, in simplest form, in terms of a ,

(a) $\log_2 9$

(b) $\log_2 \left(\frac{\sqrt{3}}{16} \right)$

(3)

(ii) Solve

$$3^x \times 2^{x+4} = 6$$

giving your answer, in simplest form, in terms of a .

(4)

T09 Laws of Logarithms

09_BASE Change of base

09_BASE_CORE core family
move

WORKED SOLUTION

Question 10

WMA12/01 October 2022

WMA12-01_2022_Oct_Q10

1

Use the power rule for log 2 of 9

$$\log_2 9 = \log_2(3^2) = 2 \log_2 3 = 2a.$$

2

Split the second logarithm

$$\log_2 \left(\frac{\sqrt{3}}{16} \right) = \log_2(3^{1/2}) - \log_2(16).$$

3

Simplify in terms of a

$$\log_2(3^{1/2}) = \frac{1}{2} \log_2 3 = \frac{a}{2} \text{ and } \log_2(16) = 4, \text{ so the expression is } \frac{a}{2} - 4.$$

4

Take log base 2 of both sides

$$\text{For } 3^x \times 2^{x+4} = 6, \text{ take } \log_2 \text{ of both sides: } \log_2(3^x) + \log_2(2^{x+4}) = \log_2 6.$$

5

Rewrite each term

$$\log_2(3^x) = xa, \log_2(2^{x+4}) = x + 4, \text{ and } \log_2 6 = \log_2(2 \cdot 3) = 1 + a.$$

6

Solve for x

$$\text{So } xa + x + 4 = a + 1. \text{ Hence } x(a + 1) = a - 3, \text{ giving } x = \frac{a-3}{a+1}.$$

FINAL ANSWER

WMA12-01_2022_Oct_Q10

$$(i)(a) \ 2a. \quad (i)(b) \ \frac{a}{2} - 4. \quad (ii) \ x = \frac{a-3}{a+1}.$$

TOPIC 09 | FAMILY 02

Log Definition

WHAT THIS FAMILY ASKS

Recognise the log definition demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

1 classified questions

7 total marks

T09

09_EXP

09_EXP_CORE

CHAPTER

Laws of Logarithms

FAMILY

Log definition

VARIANT

core family move

7 marks

WMA12/01 JANUARY 2020

Question 9

Past-paper question

WMA12-01_2020_Jan_Q09

9. (a) Sketch the curve with equation

$$y = 3 \times 4^x$$

showing the coordinates of any points of intersection with the coordinate axes.

(2)

The curve with equation $y = 6^{1-x}$ meets the curve with equation $y = 3 \times 4^x$ at the point P .

- (b) Show that the x coordinate of P is $\frac{\log_{10} 2}{\log_{10} 24}$

(5)

T09 Laws of Logarithms

09_EXP Log definition

09_EXP_CORE core family
move

WORKED SOLUTION

Question 9

WMA12/01 January 2020

WMA12-01_2020_Jan_Q09

1

Describe the sketch

$y = 3 \times 4^x$ is an increasing exponential curve. At $x = 0$, $y = 3$, so the curve crosses the y -axis at $(0, 3)$ and has no x -axis intersection.

2

Equate the two curves

At P , $6^{1-x} = 3 \times 4^x$.

3

Take logarithms

$$(1-x) \log_{10} 6 = \log_{10} 3 + x \log_{10} 4.$$

4

Collect the x terms

$$\log_{10} 6 - \log_{10} 3 = x(\log_{10} 4 + \log_{10} 6).$$

5

Simplify the logarithms

$$\log_{10} 6 - \log_{10} 3 = \log_{10} 2 \text{ and } \log_{10} 4 + \log_{10} 6 = \log_{10} 24.$$

6

State the x coordinate

$$\text{Therefore } x = \frac{\log_{10} 2}{\log_{10} 24}.$$

FINAL ANSWER

WMA12-01_2020_Jan_Q09

(a) **Increasing exponential curve through $(0, 3)$, with no x -axis intersection.**

(b) $x = \frac{\log_{10} 2}{\log_{10} 24}.$

TOPIC 09 | FAMILY 03

Log Laws

WHAT THIS FAMILY ASKS

Recognise the log laws demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

3 classified questions

20 total marks

T09

09_LAWS

09_LAWS_CORE

CHAPTER
Laws of Logarithms

FAMILY
Log laws

VARIANT
core family move

7 marks

WMA12/01 OCTOBER 2019

Question 7

Past-paper question

WMA12-01_2019_Oct_Q07

7. Given $\log_a b = k$, find, in simplest form in terms of k ,

(i) $\log_a \left(\frac{\sqrt{a}}{b} \right)$ (2)

(ii) $\frac{\log_a a^2 b}{\log_a b^3}$ (2)

(iii) $\sum_{n=1}^{50} (k + \log_a b^n)$ (3)

T09 Laws of Logarithms

09_LAWS Log laws

09_LAWS_CORE core family
move

WORKED SOLUTION

Question 7

WMA12/01 October 2019

WMA12-01_2019_Oct_Q07

1

Use the given logarithmSince $\log_a b = k$, we can replace every $\log_a b$ by k .

2

Simplify part (i)

$$\log_a \left(\frac{\sqrt{a}}{b} \right) = \log_a (a^{1/2}) - \log_a b = \frac{1}{2} - k.$$

3

Simplify the numerator and denominator

$$\log_a (a^2 b) = \log_a a^2 + \log_a b = 2 + k, \text{ and } \log_a (b^3) = 3 \log_a b = 3k.$$

4

Write part (ii)

$$\text{Therefore } \frac{\log_a (a^2 b)}{\log_a (b^3)} = \frac{k+2}{3k}.$$

5

Simplify the summand

$$\log_a (b^n) = n \log_a b = nk, \text{ so } k + \log_a (b^n) = k + nk = (n+1)k.$$

6

Sum the series

$$\sum_{n=1}^{50} (n+1)k = k \left(\sum_{n=1}^{50} n + 50 \right) = k(1275 + 50) = 1325k.$$

FINAL ANSWER

WMA12-01_2019_Oct_Q07

$$(i) \frac{1}{2} - k. \quad (ii) \frac{k+2}{3k}. \quad (iii) 1325k.$$

T09

09_LAWS

09_LAWS_CORE

CHAPTER
Laws of Logarithms

FAMILY
Log laws

VARIANT
core family move

8 marks

WMA12/01 OCTOBER 2022

Question 4

Past-paper question

WMA12-01_2022_Oct_Q04

4. The weight of a baby mammal is monitored over a 16-month period.

The weight of the mammal, w kg, is given by

$$w = \log_a(t + 5) - \log_a 4 \quad 2 \leq t \leq 18$$

where t is the age of the mammal in months and a is a constant.

Given that the weight of the mammal was 10 kg when $t = 3$

- (a) show that $a = 1.072$ correct to 3 decimal places.

(3)

Using $a = 1.072$

- (b) find an equation for t in terms of w

(3)

- (c) find the value of t when $w = 15$, giving your answer to 3 significant figures.

(2)

T09 Laws of Logarithms

09_LAWS Log laws

09_LAWS_CORE core family
move

WORKED SOLUTION

Question 4

WMA12/01 October 2022

WMA12-01_2022_Oct_Q04

1

Substitute the given informationWhen $t = 3$, $w = 10$. So $10 = \log_a(3 + 5) - \log_a 4 = \log_a 8 - \log_a 4$.

2

Combine the logarithms $\log_a 8 - \log_a 4 = \log_a \left(\frac{8}{4}\right) = \log_a 2$, so $10 = \log_a 2$.

3

Find a $10 = \log_a 2$ means $a^{10} = 2$, hence $a = 2^{1/10} = 1.071773 \dots = 1.072$ to 3 decimal places.

4

Start from the modelUsing $a = 1.072$, $w = \log_{1.072}(t + 5) - \log_{1.072} 4$.

5

Combine the logs again $w = \log_{1.072} \left(\frac{t+5}{4}\right)$.

6

Make t the subjectChanging from logarithmic form to exponential form gives $(1.072)^w = \frac{t+5}{4}$, so $t = 4(1.072)^w - 5$.

7

Substitute w equals fifteenWhen $w = 15$, $t = 4(1.072)^{15} - 5 = 6.3496 \dots$, so $t = 6.35$ months to 3 significant figures.**FINAL ANSWER**

WMA12-01_2022_Oct_Q04

(a) $a = 1.072$ to 3 d.p. (b) $t = 4(1.072)^w - 5$. (c) $t = 6.35$ months to 3 s.f.

T09

09_LAWS

09_LAWS_CORE

CHAPTER
Laws of Logarithms

FAMILY
Log laws

VARIANT
core family move

5 marks

WMA12/01 OCTOBER 2024

Question 6

Past-paper question

WMA12-01_2024_Oct_Q06

6. In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.

Using the laws of logarithms, solve

$$\log_4(12 - 2x) = 2 + 2\log_4(x + 1)$$

(5)

T09 Laws of Logarithms

09_LAWS Log laws

09_LAWS_CORE core family
move

WORKED SOLUTION

Question 6

WMA12/01 October 2024

WMA12-01_2024_Oct_Q06

1

State the domain

The logarithms require $12 - 2x > 0$ and $x + 1 > 0$, so $-1 < x < 6$.

2

Combine the right hand side

$2 = \log_4(16)$, so $2 + 2\log_4(x + 1) = \log_4(16(x + 1)^2)$.

3

Equate the arguments

$\log_4(12 - 2x) = \log_4(16(x + 1)^2)$ gives $12 - 2x = 16(x + 1)^2$.

4

Form the quadratic

$12 - 2x = 16x^2 + 32x + 16$, so $8x^2 + 17x + 2 = 0$.

5

Factor and solve

$8x^2 + 17x + 2 = (8x + 1)(x + 2) = 0$, so $x = -\frac{1}{8}$ or $x = -2$.

6

Check the domain

The value $x = -2$ is outside $-1 < x < 6$, so the solution is $x = -\frac{1}{8}$.

FINAL ANSWER

$$x = -\frac{1}{8}.$$

WMA12-01_2024_Oct_Q06

TOPIC 09 | FAMILY 04

Log Equations

WHAT THIS FAMILY ASKS

Recognise the log equations demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

V02 Quadratics from logs can create invalid negative arguments.

15 classified questions

114 total marks

T09

09_SOLVE

09_SOLVE_CORE

CHAPTER

Laws of Logarithms

FAMILY

Log equations

VARIANT

core family move

8 marks

WMA12/01 JANUARY 2021

Question 3

Past-paper question

WMA12-01_2021_Jan_Q03

3. (i) Solve

$$7^{x+2} = 3$$

giving your answer in the form $x = \log_7 a$ where a is a rational number in its simplest form.

(3)

(ii) Using the laws of logarithms, solve

$$1 + \log_2 y + \log_2 (y + 4) = \log_2 (5 - y)$$

(5)

T09 Laws of Logarithms

09_SOLVE Log equations

09_SOLVE_CORE core family
move

WORKED SOLUTION

Question 3

WMA12/01 January 2021

WMA12-01_2021_Jan_Q03

1

Rewrite the exponential equationFrom $7^{x+2} = 3$, divide by 7^2 to get $7^x = \frac{3}{49}$.

2

Write x as a logarithmTherefore $x = \log_7 \left(\frac{3}{49} \right)$.

3

Convert one to a logarithmSince $1 = \log_2 2$, the equation becomes $\log_2 2 + \log_2 y + \log_2 (y + 4) = \log_2 (5 - y)$.

4

Combine the logarithms $\log_2 \{2y(y + 4)\} = \log_2 (5 - y)$.

5

Solve the quadraticThus $2y(y + 4) = 5 - y$, so $2y^2 + 9y - 5 = 0 = (2y - 1)(y + 5)$.

6

Check the domainThe original logarithms require $y > 0$ and $y < 5$, so $y = -5$ is invalid and $y = \frac{1}{2}$.**FINAL ANSWER**(i) $x = \log_7 \left(\frac{3}{49} \right)$. (ii) $y = \frac{1}{2}$.

WMA12-01_2021_Jan_Q03

T09

09_SOLVE

09_SOLVE_CORE

CHAPTER
Laws of Logarithms

FAMILY
Log equations

VARIANT
core family move

8 marks

WMA12/01 MAY/JUNE 2021
Question 2

Past-paper question

WMA12-01_2021_MayJune_Q02

2.




Figure 1

Figure 1 shows a sketch of the curve with equation $y = 4^x$.

A copy of Figure 1, labelled Diagram 1, is shown on the next page.

(a) On Diagram 1, sketch the curve with equation

(i) $y = 2^x$

(ii) $y = 4^x - 6$

Label clearly the coordinates of any points of intersection with the coordinate axes.

(4)

The curve with equation $y = 2^x$ meets the curve with equation $y = 4^x - 6$ at the point P .

(b) Using algebra, find the exact coordinates of P .

(4)

Question 2 continued




Diagram 1

T09 Laws of Logarithms

09_SOLVE Log equations

09_SOLVE_CORE core family
move

WORKED SOLUTION

Question 2

WMA12/01 May/June 2021

WMA12-01_2021_MayJune_Q02

1

Describe y equals two to the x

$y = 2^x$ is an increasing exponential curve. It crosses the y -axis at $(0, 1)$ and has no x -axis intercept.

2

Describe y equals four to the x minus six

$y = 4^x - 6$ is the graph of $y = 4^x$ translated 6 units down. It crosses the y -axis at $(0, -5)$.

3

Find the x -axis intercept

For $y = 4^x - 6$, set $4^x - 6 = 0$. Hence $4^x = 6$ and $x = \log_4 6$.

4

Set up the intersection equation

At P , $2^x = 4^x - 6$.

5

Use a substitution

Let $u = 2^x$. Then $4^x = (2^x)^2 = u^2$, so $u = u^2 - 6$.

6

Solve for u

$u^2 - u - 6 = 0$, so $(u - 3)(u + 2) = 0$. Since $u = 2^x > 0$, $u = 3$.

7

State P

$2^x = 3$, so $x = \log_2 3$, and $y = 3$. Therefore $P = (\log_2 3, 3)$.

FINAL ANSWER

WMA12-01_2021_MayJune_Q02

(a)(i) Sketch $y = 2^x$ through $(0, 1)$ with no x -axis intercept. (a)(ii) Sketch $y = 4^x - 6$ through $(0, -5)$ and $(\log_4 6, 0)$. (b) $P = (\log_2 3, 3)$.

T09

09_SOLVE

09_SOLVE_CORE

CHAPTER

Laws of Logarithms

FAMILY

Log equations

VARIANT

core family move

5 marks

WMA12/01 JANUARY 2022

Question 4

Past-paper question

WMA12-01_2022_Jan_Q04

4. Using the laws of logarithms, solve

$$\log_3(32 - 12x) = 2\log_3(1 - x) + 3$$

(5)

DR ESLAM AHMED

T09 Laws of Logarithms

09_SOLVE Log equations

09_SOLVE_CORE core family
move

WORKED SOLUTION

Question 4

WMA12/01 January 2022

WMA12-01_2022_Jan_Q04

1

Rewrite the right side $2 \log_3(1 - x) = \log_3(1 - x)^2$ and $3 = \log_3 27$.

2

Combine the logarithmsThe equation becomes $\log_3(32 - 12x) = \log_3\{27(1 - x)^2\}$.

3

Equate the arguments $32 - 12x = 27(1 - x)^2$.

4

Form the quadratic $32 - 12x = 27 - 54x + 27x^2$, so $27x^2 - 42x - 5 = 0$.

5

Solve $x = \frac{42 \pm 48}{54}$, giving $x = \frac{5}{3}$ or $x = -\frac{1}{9}$.

6

Check the domainThe original logarithms require $1 - x > 0$, so $x < 1$. Therefore $x = \frac{5}{3}$ is invalid and $x = -\frac{1}{9}$.**FINAL ANSWER**

$$x = -\frac{1}{9}.$$

WMA12-01_2022_Jan_Q04

T09

09_SOLVE

09_SOLVE_CORE

CHAPTER

Laws of Logarithms

FAMILY

Log equations

VARIANT

core family move

6 marks

WMA12/01 MAY/JUNE 2022

Question 4

Past-paper question

WMA12-01_2022_MayJune_Q04

4. In this question you must show all stages of your working.

Give your answers in fully simplified surd form.

Given that a and b are positive constants, solve the simultaneous equations

$$\begin{aligned}a - b &= 8 \\ \log_4 a + \log_4 b &= 3\end{aligned}$$

(6)

T09 Laws of Logarithms

09_SOLVE Log equations

09_SOLVE_CORE core family
move

WORKED SOLUTION

Question 4

WMA12/01 May/June 2022

WMA12-01_2022_MayJune_Q04

1

Combine the logarithms

$$\log_4 a + \log_4 b = \log_4(ab) = 3.$$

2

Find ab

$$\text{Therefore } ab = 4^3 = 64.$$

3

Use a minus b

$$\text{From } a - b = 8, b = a - 8.$$

4

Form a quadratic

$$a(a - 8) = 64, \text{ so } a^2 - 8a - 64 = 0.$$

5

Solve for a

$$a = \frac{8 \pm \sqrt{64 + 256}}{2} = 4 \pm 4\sqrt{5}.$$

6

Choose positive constants

$$\text{Since } b = a - 8 \text{ must also be positive, take } a = 4 + 4\sqrt{5}.$$

7

Find b

$$b = a - 8 = 4\sqrt{5} - 4.$$

FINAL ANSWER

$$a = 4 + 4\sqrt{5}, \quad b = 4\sqrt{5} - 4.$$

WMA12-01_2022_MayJune_Q04

T09

09_SOLVE

09_SOLVE_CORE

CHAPTER

Laws of Logarithms

FAMILY

Log equations

VARIANT

core family move

6 marks

WMA12/01 JANUARY 2023

Question 4

Past-paper question

WMA12-01_2023_Jan_Q04

4. (i) Using the laws of logarithms, solve

$$\log_3(4x) + 2 = \log_3(5x + 7) \quad (3)$$

- (ii) Given that

$$\sum_{r=1}^2 \log_a(y^r) = \sum_{r=1}^2 (\log_a y)^r \quad y > 1, a > 1, y \neq a$$

find y in terms of a , giving your answer in simplest form.

(3)

T09 Laws of Logarithms

09_SOLVE Log equations

09_SOLVE_CORE core family
move

WORKED SOLUTION

Question 4

WMA12/01 January 2023

WMA12-01_2023_Jan_Q04

1

Rewrite the constant

Since $2 = \log_3 9$, the equation becomes $\log_3(4x) + \log_3 9 = \log_3(5x + 7)$.

2

Combine the logarithms

$$\log_3(4x) + \log_3 9 = \log_3(36x).$$

3

Solve for x

Therefore $36x = 5x + 7$, so $31x = 7$ and $x = \frac{7}{31}$.

4

Introduce L

Let $L = \log_a y$. Then $\log_a(y^r) = r \log_a y = rL$.

5

Evaluate the left sum

$$\sum_{r=1}^2 \log_a(y^r) = L + 2L = 3L.$$

6

Evaluate the right sum

$$\sum_{r=1}^2 (\log_a y)^r = L + L^2.$$

7

Solve for L

So $3L = L + L^2$, which gives $L^2 - 2L = 0$ and $L(L - 2) = 0$.

8

Use the restrictions

Since $a > 1$ and $y > 1$, $\log_a y > 0$, so $L \neq 0$. Hence $L = 2$ and $y = a^2$.

FINAL ANSWER

WMA12-01_2023_Jan_Q04

(i) $x = \frac{7}{31}$. (ii) $y = a^2$.

T09

09_SOLVE

09_SOLVE_CORE

CHAPTER

Laws of Logarithms

FAMILY

Log equations

VARIANT

core family move

6 marks

WMA12/01 OCTOBER 2023

Question 5

Past-paper question

WMA12-01_2023_Oct_Q05

5. In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

(i) Solve

$$3^a = 70$$

giving the answer to 3 decimal places.

(2)

(ii) Find the exact value of b such that

$$4 + 3 \log_3 b = \log_3 5b$$

(4)

T09 Laws of Logarithms

09_SOLVE Log equations

09_SOLVE_CORE core family
move

WORKED SOLUTION

Question 5

WMA12/01 October 2023

WMA12-01_2023_Oct_Q05

1

Solve the exponential equation

$$3^a = 70 \text{ gives } a = \log_3 70 = \frac{\log 70}{\log 3}.$$

2

Round a $a = 3.867147 \dots$, so $a = 3.867$ to 3 decimal places.

3

Rewrite four as a logarithm

$$4 = \log_3(3^4) = \log_3 81.$$

4

Combine the left side

$$4 + 3 \log_3 b = \log_3 81 + \log_3(b^3) = \log_3(81b^3).$$

5

Equate argumentsThe equation becomes $\log_3(81b^3) = \log_3(5b)$, so $81b^3 = 5b$.

6

Use the log domainSince $\log_3 b$ is defined, $b > 0$. Divide by b to get $81b^2 = 5$.

7

Find b

$$b^2 = \frac{5}{81}, \text{ and } b > 0, \text{ so } b = \frac{\sqrt{5}}{9}.$$

FINAL ANSWER

WMA12-01_2023_Oct_Q05

$$(i) a = 3.867. \quad (ii) b = \frac{\sqrt{5}}{9}.$$

T09

09_SOLVE

09_SOLVE_CORE

CHAPTER

Laws of Logarithms

FAMILY

Log equations

VARIANT

core family move

8 marks

WMA12/01 JANUARY 2024

Question 6

Past-paper question

WMA12-01_2024_Jan_Q06

6. (a) Given that

$$2\log_4(x+3) + \log_4 x = \log_4(4x+2) + \frac{1}{2}$$

show that

$$x^3 + 6x^2 + x - 4 = 0 \quad (4)$$

(b) Given also that -1 is a root of the equation

$$x^3 + 6x^2 + x - 4 = 0$$

(i) use algebra to find the other two roots of the equation. (3)

(ii) Hence solve

$$2\log_4(x+3) + \log_4 x = \log_4(4x+2) + \frac{1}{2} \quad (1)$$

T09 Laws of Logarithms

09_SOLVE Log equations

09_SOLVE_CORE core family
move

WORKED SOLUTION

Question 6

WMA12/01 January 2024

WMA12-01_2024_Jan_Q06

1

Combine the logarithms $2 \log_4(x+3) = \log_4((x+3)^2)$, so the left side is $\log_4(x(x+3)^2)$.

2

Rewrite one half $\frac{1}{2} = \log_4 2$, so the right side is $\log_4(4x+2) + \log_4 2 = \log_4(8x+4)$.

3

Equate argumentsTherefore $x(x+3)^2 = 8x+4$.

4

Expand $x(x^2+6x+9) = 8x+4$, so $x^3+6x^2+9x = 8x+4$.

5

Show the cubicRearranging gives $x^3+6x^2+x-4=0$.

6

Use the given rootSince -1 is a root, $(x+1)$ is a factor of x^3+6x^2+x-4 .

7

Factor the cubic $x^3+6x^2+x-4 = (x+1)(x^2+5x-4)$.

8

Solve the quadratic $x^2+5x-4=0$ gives $x = \frac{-5 \pm \sqrt{25+16}}{2} = \frac{-5 \pm \sqrt{41}}{2}$.

9

Apply the logarithm domainThe original log equation requires $x > 0$. The only positive root is $x = \frac{-5+\sqrt{41}}{2}$.**FINAL ANSWER**

WMA12-01_2024_Jan_Q06

(a) **Proven.** (b)(i) $x = \frac{-5+\sqrt{41}}{2}$ and $x = \frac{-5-\sqrt{41}}{2}$. (b)(ii) $x = \frac{-5+\sqrt{41}}{2}$.

T09

09_SOLVE

09_SOLVE_CORE

CHAPTER

Laws of Logarithms

FAMILY

Log equations

VARIANT

core family move

9 marks

WMA12/01 MAY/JUNE (R) 2024

Question 9

Past-paper question

WMA12-01_2024_MayJuneR_Q09

9. Given that

$$3 \log_2(t+4) - 2 \log_2(t-2) = 7$$

(a) verify that $t = 4$ is a solution of the above equation,

(2)

(b) show that

$$t^3 - 116t^2 + 560t - 448 = 0$$

(3)

(c) Hence, using algebra and showing your working, solve

$$3 \log_2(t+4) - 2 \log_2(t-2) = 7$$

giving each answer in simplest form.

(Solutions based entirely on calculator technology are not acceptable.)

(4)

T09 Laws of Logarithms

09_SOLVE Log equations

09_SOLVE_CORE core family
move

WORKED SOLUTION

Question 9

WMA12/01 May/June (R) 2024

WMA12-01_2024_MayJuneR_Q09

1

Verify t equals fourSubstitute $t = 4$: $3 \log_2(8) - 2 \log_2(2) = 3(3) - 2(1) = 7$.

2

Combine the logarithms

$$3 \log_2(t+4) - 2 \log_2(t-2) = \log_2 \left(\frac{(t+4)^3}{(t-2)^2} \right).$$

3

Use seven as a logarithm

$$7 = \log_2(128), \text{ so } \frac{(t+4)^3}{(t-2)^2} = 128.$$

4

Clear the denominator

$$(t+4)^3 = 128(t-2)^2.$$

5

Expand

$$t^3 + 12t^2 + 48t + 64 = 128t^2 - 512t + 512.$$

6

Show the cubic

$$\text{Rearranging gives } t^3 - 116t^2 + 560t - 448 = 0.$$

7

Use the known rootSince $t = 4$ is a root, $(t - 4)$ is a factor.

8

Factor the cubic

$$t^3 - 116t^2 + 560t - 448 = (t - 4)(t^2 - 112t + 112).$$

9

Solve the quadratic

$$t^2 - 112t + 112 = 0 \text{ gives } t = \frac{112 \pm \sqrt{112^2 - 4(112)}}{2} = 56 \pm 12\sqrt{21}.$$

10

Use the log domain

The original equation requires $t > 2$. The value $56 - 12\sqrt{21}$ is less than 2, so it is rejected.
Hence $t = 4$ or $t = 56 + 12\sqrt{21}$.

FINAL ANSWER

WMA12-01_2024_MayJuneR_Q09

(a) **Verified.** (b) **Proven.** (c) $t = 4$ **or** $t = 56 + 12\sqrt{21}$.

DR ESLAM AHMED

T09

09_SOLVE

09_SOLVE_CORE

CHAPTER

Laws of Logarithms

FAMILY

Log equations

VARIANT

core family move

8 marks

WMA12/01 JANUARY 2025

Question 7

Past-paper question

WMA12-01_2025_Jan_Q07

7.

In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.

- (i) The table below shows values of x and y , where $y = \log_{10}(x + 5)$, for x values between -1 and 4

x	-1	0	1	2	3	4
$y = \log_{10}(x + 5)$	$\log_{10} 4$	$\log_{10} 5$	$\log_{10} 6$	$\log_{10} 7$	$\log_{10} 8$	$\log_{10} 9$

Using the trapezium rule with all the y values in the given table, show that

$$\int_{-1}^4 \log_{10}(x + 5) \, dx \approx \log_{10} k$$

where k is an integer to be found.

(3)

- (ii) Find the value of a such that

$$2\log_5(5 - a) - \log_5(a + 25) = 1$$

(5)

T09 Laws of Logarithms

09_SOLVE Log equations

09_SOLVE_CORE core family
move

WORKED SOLUTION

Question 7

WMA12/01 January 2025

WMA12-01_2025_Jan_Q07

1

Use the trapezium rule

The width is 1, so $\int_{-1}^4 \log_{10}(x+5) dx \approx \frac{1}{2} \{ \log_{10} 4 + \log_{10} 9 + 2(\log_{10} 5 + \log_{10} 6 + \log_{10} 7 + \log_{10} 8) \}$.

2

Combine the logarithms

This is $\log_{10} (\sqrt{4 \times 9} \times 5 \times 6 \times 7 \times 8)$.

3

Find k

$\sqrt{4 \times 9} \times 5 \times 6 \times 7 \times 8 = 6 \times 1680 = 10080$, so $k = 10080$.

4

State the domain

The logarithms require $5 - a > 0$ and $a + 25 > 0$, so $-25 < a < 5$.

5

Combine the logarithms

$2 \log_5 (5 - a) - \log_5 (a + 25) = \log_5 \left(\frac{(5-a)^2}{a+25} \right)$.

6

Use one as a logarithm

$1 = \log_5 5$, so $\frac{(5-a)^2}{a+25} = 5$.

7

Form the quadratic

$(5 - a)^2 = 5(a + 25)$, so $a^2 - 10a + 25 = 5a + 125$. Hence $a^2 - 15a - 100 = 0$.

8

Solve and check

$(a - 20)(a + 5) = 0$, so $a = 20$ or $a = -5$. Only $a = -5$ is in the domain, so $a = -5$.

FINAL ANSWER

(i) $k = 10080$. (ii) $a = -5$.

WMA12-01_2025_Jan_Q07

T09

09_SOLVE

09_SOLVE_CORE

CHAPTER

Laws of Logarithms

FAMILY

Log equations

VARIANT

core family move

10 marks

WMA12/01 MAY/JUNE 2025

Question 6

Past-paper question

WMA12-01_2025_MayJune_Q06

6:

**In this question you must show all stages of your working.
Solutions relying on calculator technology are not acceptable.**

(i) Solve

$$2\log_2(4-x) = 3 + \log_2\left(\frac{x+11}{2}\right) \quad (5)$$

(ii) The curves C_1 and C_2 with equations

$$y = 3^{2x+1} \quad \text{and} \quad y = 6 \times 3^x$$

meet at the point P .

Find the exact coordinates of P , writing your answer in the form $(\log_3 a, b)$ where a and b are integers.

(5)

T09 Laws of Logarithms

09_SOLVE Log equations

09_SOLVE_CORE core family
move

WORKED SOLUTION

Question 6

WMA12/01 May/June 2025

WMA12-01_2025_MayJune_Q06

1

State the domain

The logarithms require $4 - x > 0$ and $\frac{x+11}{2} > 0$, so $-11 < x < 4$.

2

Use log laws

$2 \log_2(4 - x) = \log_2((4 - x)^2)$ and $3 + \log_2\left(\frac{x+11}{2}\right) = \log_2\left(8 \cdot \frac{x+11}{2}\right)$.

3

Equate the arguments

Thus $(4 - x)^2 = 4(x + 11)$.

4

Solve the quadratic

$x^2 - 8x + 16 = 4x + 44$, so $x^2 - 12x - 28 = 0$. Hence $(x - 14)(x + 2) = 0$.

5

Check the domain

The root 14 is outside the domain, so $x = -2$.

6

Equate the curves

At P , $3^{2x+1} = 6 \times 3^x$.

7

Let u equal 3 to the x

Let $u = 3^x$. Then $3^{2x+1} = 3u^2$, so $3u^2 = 6u$.

8

Find u

Since $u > 0$, divide by $3u$ to get $u = 2$. Hence $3^x = 2$ and $x = \log_3 2$.

9

Find y

$y = 6 \times 3^x = 6 \times 2 = 12$, so $P = (\log_3 2, 12)$.

FINAL ANSWER

WMA12-01_2025_MayJune_Q06

(i) $x = -2$. (ii) $P = (\log_3 2, 12)$.

T09

09_SOLVE

09_SOLVE_CORE

CHAPTER

Laws of Logarithms

FAMILY

Log equations

VARIANT

core family move

10 marks

WMA12/01 OCTOBER 2025

Question 9

Past-paper question

WMA12-01_2025_Oct_Q09

9.

**In this question you must show detailed reasoning.
Solutions relying on calculator technology are not acceptable.**

(i) Solve

$$2\log_3(4x + 5) - \log_3(x + 3) = 2 \quad (5)$$

(ii) Given that $a > 0$, $b > 0$ and

$$\log_{10} a + \log_{10} b = \log_{10}(a + b)$$

(a) prove that $a = \frac{b}{b-1}$ (3)

(b) Hence write down the full restriction on the value of b , giving a reason for your answer. (2)

T09 Laws of Logarithms

09_SOLVE Log equations

09_SOLVE_CORE core family
move

WORKED SOLUTION

Question 9

WMA12/01 October 2025

WMA12-01_2025_Oct_Q09

1

State the domain

The logarithms require $4x + 5 > 0$ and $x + 3 > 0$, so $x > -\frac{5}{4}$.

2

Combine the logarithms

$$2\log_3(4x + 5) - \log_3(x + 3) = \log_3\left(\frac{(4x+5)^2}{x+3}\right).$$

3

Use two as a logarithm

$$2 = \log_3 9, \text{ so } \frac{(4x+5)^2}{x+3} = 9.$$

4

Form the quadratic

$$(4x + 5)^2 = 9(x + 3), \text{ so } 16x^2 + 40x + 25 = 9x + 27. \text{ Hence } 16x^2 + 31x - 2 = 0.$$

5

Solve the quadratic

$$16x^2 + 31x - 2 = (16x - 1)(x + 2) = 0, \text{ so } x = \frac{1}{16} \text{ or } x = -2.$$

6

Check the domain

$x = -2$ is outside the domain, so $x = \frac{1}{16}$.

7

Use the logarithm equation

$$\log_{10} a + \log_{10} b = \log_{10}(a + b) \text{ gives } \log_{10}(ab) = \log_{10}(a + b).$$

8

Equate the arguments

Since the logarithm function is one-to-one, $ab = a + b$.

9

Rearrange for a

$$ab - a = b, \text{ so } a(b - 1) = b \text{ and therefore } a = \frac{b}{b-1}.$$

10

Restrict b

Given $a > 0$ and $b > 0$, the fraction $\frac{b}{b-1}$ must be positive. Since the numerator is positive, the denominator must also be positive, so $b > 1$.

FINAL ANSWER

WMA12-01_2025_Oct_Q09

(i) $x = \frac{1}{16}$. (ii)(a) **Proven.** (ii)(b) $b > 1$.

DR ESLAM AHMED

T09

09_SOLVE

09_SOLVE_HV02

CHAPTER
Laws of Logarithms

FAMILY
Log equations

VARIANT
Quadratics from logs can
create invalid negative
arguments.

10 marks

WMA12/01 MAY/JUNE 2021

Question 7

Past-paper question

WMA12-01_2021_MayJune_Q07

7. (a) Given that

$$3 \log_3(2x - 1) = 2 + \log_3(14x - 25)$$

show that

$$2x^3 - 3x^2 - 30x + 56 = 0$$

(4)

- (b) Show that -4 is a root of this cubic equation.

(2)

- (c) Hence, using algebra and showing each step of your working, solve

$$3 \log_3(2x - 1) = 2 + \log_3(14x - 25)$$

(4)

T09 Laws of Logarithms

09_SOLVE Log equations

09_SOLVE_HV02 Quadratics from logs can create invalid negative arguments.

WORKED SOLUTION

Question 7

WMA12/01 May/June 2021

WMA12-01_2021_MayJune_Q07

1

Use the power law

$$3 \log_3(2x - 1) = \log_3(2x - 1)^3.$$

2

Rewrite two as a logarithm

Since $2 = \log_3 9$, the right side is $\log_3 9 + \log_3(14x - 25) = \log_3\{9(14x - 25)\}$.

3

Equate the arguments

So $(2x - 1)^3 = 9(14x - 25)$.

4

Expand and simplify

$8x^3 - 12x^2 + 6x - 1 = 126x - 225$, which simplifies to $2x^3 - 3x^2 - 30x + 56 = 0$.

5

Check the given root

Substituting $x = -4$ gives $2(-64) - 3(16) - 30(-4) + 56 = 0$, so -4 is a root.

6

Factorise the cubic

Dividing by $(x + 4)$ gives $2x^2 - 11x + 14 = (2x - 7)(x - 2)$.

7

Apply the logarithm domain

The possible roots are -4 , 2 , and $\frac{7}{2}$. The original logarithms require $2x - 1 > 0$ and $14x - 25 > 0$, so $x > \frac{25}{14}$.

8

State the valid solutions

Therefore the solutions are $x = 2$ and $x = \frac{7}{2}$.

FINAL ANSWER

WMA12-01_2021_MayJune_Q07

(a) Proven. (b) -4 is a root of the cubic. (c) $x = 2, \frac{7}{2}$.

T09

09_SOLVE

09_SOLVE_HV02

CHAPTER
Laws of Logarithms

FAMILY
Log equations

VARIANT
Quadratics from logs can
create invalid negative
arguments.

5 marks

WMA12/01 MAY/JUNE 2023

Question 5

Past-paper question

WMA12-01_2023_MayJune_Q05

5. Use the laws of logarithms to solve

$$\log_2(16x) + \log_2(x + 1) = 3 + \log_2(x + 6)$$

(5)

T09 Laws of Logarithms

09_SOLVE Log equations

09_SOLVE_HV02 Quadratics
from logs can create invalid
negative arguments.

WORKED SOLUTION

Question 5

WMA12/01 May/June 2023

WMA12-01_2023_MayJune_Q05

1

State the domain

The logarithms require $16x > 0$, $x + 1 > 0$ and $x + 6 > 0$, so $x > 0$.

2

Combine the left side

$$\log_2(16x) + \log_2(x + 1) = \log_2(16x(x + 1)).$$

3

Rewrite the right side

$$3 = \log_2 8, \text{ so } 3 + \log_2(x + 6) = \log_2(8(x + 6)).$$

4

Equate the arguments

$$\text{Therefore } 16x(x + 1) = 8(x + 6).$$

5

Solve the quadratic

$$16x^2 + 16x = 8x + 48, \text{ so } 16x^2 + 8x - 48 = 0. \text{ Dividing by 8 gives } 2x^2 + x - 6 = 0.$$

6

Factor and use the domain

$$(2x - 3)(x + 2) = 0, \text{ so } x = \frac{3}{2} \text{ or } x = -2. \text{ The domain } x > 0 \text{ rejects } -2, \text{ leaving } x = \frac{3}{2}.$$

FINAL ANSWER

$$x = \frac{3}{2}.$$

WMA12-01_2023_MayJune_Q05

T09

09_SOLVE

09_SOLVE_HV02

CHAPTER
Laws of Logarithms

FAMILY
Log equations

VARIANT
Quadratics from logs can
create invalid negative
arguments.

6 marks

WMA12/01 MAY/JUNE 2024

Question 3

Past-paper question

WMA12-01_2024_MayJune_Q03

3. **In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.**

- (i) Using the laws of logarithms, solve

$$2 \log_2 (2 - x) = 4 + \log_2 (x + 10) \quad (5)$$

- (ii) Find the value of

$$\log_{\sqrt{a}} a^6$$

where a is a positive constant greater than 1

(1)

T09 Laws of Logarithms

09_SOLVE Log equations

09_SOLVE_HV02 Quadratics
from logs can create invalid
negative arguments.

WORKED SOLUTION

Question 3

WMA12/01 May/June 2024

WMA12-01_2024_MayJune_Q03

1

State the domain

The logarithms require $2 - x > 0$ and $x + 10 > 0$, so $-10 < x < 2$.

2

Combine powers

$$2 \log_2(2 - x) = \log_2((2 - x)^2).$$

3

Rewrite four

$$4 = \log_2 16, \text{ so } 4 + \log_2(x + 10) = \log_2(16(x + 10)).$$

4

Equate arguments

$$(2 - x)^2 = 16(x + 10).$$

5

Solve the quadratic

$$x^2 - 4x + 4 = 16x + 160, \text{ so } x^2 - 20x - 156 = 0.$$

6

Use the domain

$x = 26$ or $x = -6$. The value 26 is outside the domain, so $x = -6$.

7

Evaluate the logarithm

$$\log_{\sqrt{a}}(a^6) = \log_{a^{1/2}}(a^6) = \frac{6}{1/2} = 12.$$

FINAL ANSWER

(i) $x = -6$. (ii) 12.

WMA12-01_2024_MayJune_Q03

T09

09_SOLVE

09_SOLVE_HV02

CHAPTER
Laws of Logarithms

FAMILY
Log equations

VARIANT
Quadratics from logs can
create invalid negative
arguments.

9 marks

WMA12/01 JANUARY 2026

Question 7

Past-paper question

WMA12-01_2026_Jan_Q07

7.

**In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.**

- (i) Find the exact solution of the equation

$$8^{2x-5} = 20$$

giving the answer in the form $a + b \log_2 5$ where a and b are rational constants.

(4)

- (ii) Using the laws of logarithms, solve the equation

$$\log_3 (13 + 2y) + 3 = 2 \log_3 (4 - y)$$

(5)

T09 Laws of Logarithms

09_SOLVE Log equations

09_SOLVE_HV02 Quadratics
from logs can create invalid
negative arguments.

WORKED SOLUTION

Question 7

WMA12/01 January 2026

WMA12-01_2026_Jan_Q07

1

Take logarithms base 2

From $8^{2x-5} = 20$, take $\log_2$ of both sides to get $(2x - 5) \log_2 8 = \log_2 20$.

2

Simplify the logarithms

Since $\log_2 8 = 3$ and $\log_2 20 = \log_2(4 \cdot 5) = 2 + \log_2 5$, we get $3(2x - 5) = 2 + \log_2 5$.

3

Solve for x

$6x - 15 = 2 + \log_2 5$, so $6x = 17 + \log_2 5$ and $x = \frac{17}{6} + \frac{1}{6} \log_2 5$.

4

Use log laws

$3 = \log_3 27$ and $2 \log_3(4 - y) = \log_3(4 - y)^2$.

5

Remove the logarithms

$\log_3(13 + 2y) + \log_3 27 = \log_3(4 - y)^2$, so $27(13 + 2y) = (4 - y)^2$.

6

Solve the quadratic

$351 + 54y = y^2 - 8y + 16$, hence $y^2 - 62y - 335 = 0 = (y - 67)(y + 5)$.

7

Check the logarithm domains

We need $13 + 2y > 0$ and $4 - y > 0$. Therefore $y = 67$ is not valid, and the solution is $y = -5$.

FINAL ANSWER

WMA12-01_2026_Jan_Q07

(i) $x = \frac{17}{6} + \frac{1}{6} \log_2 5$. (ii) $y = -5$.

PURE MATHEMATICS 2

ADAPTIVE ALL CLASSIFIED

Presentation Edition | Solutions

CLASSIFIED BY IDEA | ROOM TO THINK

Every demand appears clearly, with practical space for the mathematics.

Dr Eslam Ahmed

Elite Mathematics

ElitelGCSE.com | 011 2000 9622

PURE 2

PURE 2 | TOPIC 10

Trigonometric Equations

Your classified route

20

QUESTIONS

173

MARKS

4

FAMILIES

5

VARIANTS

DR ESLAM AHMED

CLASSIFIED BY IDEA

Move from recognition to method, then use the working space to complete the solution.

TOPIC 10 | FAMILY 01

Trig Equations

WHAT THIS FAMILY ASKS

Recognise the trig equations demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE**V01** core family move**9 classified questions**

79 total marks

T10

10_CAST

10_CAST_CORE

CHAPTER
Trigonometric Equations

FAMILY
Trig equations

VARIANT
core family move

12 marks

WMA12/01 OCTOBER 2019

Question 9

Past-paper question

WMA12-01_2019_Oct_Q09

9. Solutions based entirely on graphical or numerical methods are not acceptable in this question.

- (i) Solve, for $0 \leq \theta < 180^\circ$, the equation

$$3 \sin(2\theta - 10^\circ) = 1$$

giving your answers to one decimal place.

(4)

- (ii) The first three terms of an arithmetic sequence are

$$\sin \alpha, \quad \frac{1}{\tan \alpha} \quad \text{and} \quad 2 \sin \alpha$$

where α is a constant.

- (a) Show that $2 \cos \alpha = 3 \sin^2 \alpha$

(3)

Given that $\pi < \alpha < 2\pi$,

- (b) find, showing all working, the value of α to 3 decimal places.

(5)

T10 Trigonometric Equations 10_CAST Trig equations

10_CAST_CORE core family
move

WORKED SOLUTION

Question 9

WMA12/01 October 2019

WMA12-01_2019_Oct_Q09

1

Solve the sine equationLet $u = 2\theta - 10^\circ$. Then $\sin u = \frac{1}{3}$.

2

Find the valid angle valuesSince $0 \leq \theta < 180^\circ$, $-10^\circ \leq u < 350^\circ$. In this interval, $u = 19.471 \dots^\circ$ or $u = 160.529 \dots^\circ$.

3

Return to theta $\theta = \frac{u+10^\circ}{2}$, giving $\theta = 14.7^\circ$ and 85.3° to one decimal place.

4

Use the arithmetic sequence conditionFor three consecutive terms of an arithmetic sequence, second minus first equals third minus second. Hence $\frac{1}{\tan \alpha} - \sin \alpha = 2 \sin \alpha - \frac{1}{\tan \alpha}$.

5

Prove the required identityThis gives $\frac{2}{\tan \alpha} = 3 \sin \alpha$. Since $\frac{1}{\tan \alpha} = \frac{\cos \alpha}{\sin \alpha}$, we get $\frac{2 \cos \alpha}{\sin \alpha} = 3 \sin \alpha$, so $2 \cos \alpha = 3 \sin^2 \alpha$.

6

Form a quadratic in cos alphaUsing $\sin^2 \alpha = 1 - \cos^2 \alpha$, $2 \cos \alpha = 3(1 - \cos^2 \alpha)$, so $3 \cos^2 \alpha + 2 \cos \alpha - 3 = 0$.

7

Solve in the given interval $\cos \alpha = \frac{-1+\sqrt{10}}{3}$; the other root is less than -1 . Since $\pi < \alpha < 2\pi$ and cosine is positive, α is in the fourth quadrant.

8

State alpha $\alpha = 2\pi - \cos^{-1}\left(\frac{-1+\sqrt{10}}{3}\right) = 5.517$ radians to 3 decimal places.**FINAL ANSWER**

WMA12-01_2019_Oct_Q09

(i) $\theta = 14.7^\circ, 85.3^\circ$. (ii) (a) **Proven.** (ii) (b) $\alpha = 5.517$ radians.

T10

10_CAST

10_CAST_CORE

CHAPTER

Trigonometric Equations

FAMILY

Trig equations

VARIANT

core family move

8 marks

WMA12/01 JANUARY 2021

Question 6

Past-paper question

WMA12-01_2021_Jan_Q06

6. (a) Show that the equation

$$\frac{3 \sin \theta \cos \theta}{2 \sin \theta - 1} = 5 \tan \theta \quad \sin \theta \neq \frac{1}{2}$$

can be written in the form

$$3 \sin^3 \theta + 10 \sin^2 \theta - 8 \sin \theta = 0$$

(4)

- (b) Hence solve, for $-\frac{\pi}{4} < x < \frac{\pi}{4}$

$$\frac{3 \sin 2x \cos 2x}{2 \sin 2x - 1} = 5 \tan 2x$$

giving your answers to 3 decimal places where appropriate.

(4)

T10 Trigonometric Equations 10_CAST Trig equations

10_CAST_CORE core family
move

WORKED SOLUTION

Question 6

WMA12/01 January 2021

WMA12-01_2021_Jan_Q06

1

Clear the fractionStarting with $\frac{3 \sin \theta \cos \theta}{2 \sin \theta - 1} = 5 \tan \theta$, multiply by $(2 \sin \theta - 1) \cos \theta$.

2

Use tan thetaThis gives $3 \sin \theta \cos^2 \theta = 5 \sin \theta (2 \sin \theta - 1)$.

3

Use cos squaredSubstitute $\cos^2 \theta = 1 - \sin^2 \theta$: $3 \sin \theta (1 - \sin^2 \theta) = 10 \sin^2 \theta - 5 \sin \theta$.

4

Collect termsRearranging gives $3 \sin^3 \theta + 10 \sin^2 \theta - 8 \sin \theta = 0$.

5

Factor the sine equation $3s^3 + 10s^2 - 8s = 0$ gives $s(3s - 2)(s + 4) = 0$, where $s = \sin 2x$.

6

Use the valid sine valuesSince $-\frac{\pi}{2} < 2x < \frac{\pi}{2}$, the valid values are $\sin 2x = 0$ and $\sin 2x = \frac{2}{3}$.

7

Find x $2x = 0$ gives $x = 0$. Also $2x = \sin^{-1}\left(\frac{2}{3}\right)$, so $x = 0.364863... = 0.365$ to 3 decimal places.**FINAL ANSWER**

WMA12-01_2021_Jan_Q06

(a) Proven. (b) $x = 0, 0.365$.

T10

10_CAST

10_CAST_CORE

CHAPTER

Trigonometric Equations

FAMILY

Trig equations

VARIANT

core family move

8 marks

WMA12/01 JANUARY 2022

Question 7

Past-paper question

WMA12-01_2022_Jan_Q07

7. In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

- (i) Solve, for $-90^\circ < x < 90^\circ$, the equation

$$3 \sin(2x - 15^\circ) = \cos(2x - 15^\circ)$$

giving your answers to one decimal place.

(4)

- (ii) Solve, for $0 < \theta < 2\pi$, the equation

$$4 \sin^2 \theta + 8 \cos \theta = 3$$

giving your answers to 3 significant figures.

(4)

T10 Trigonometric Equations 10_CAST Trig equations

10_CAST_CORE core family
move

WORKED SOLUTION

Question 7

WMA12/01 January 2022

WMA12-01_2022_Jan_Q07

1

Solve the first tangent equationLet $u = 2x - 15^\circ$. Then $3 \sin u = \cos u$, so $\tan u = \frac{1}{3}$.

2

Use the interval for u Since $-90^\circ < x < 90^\circ$, $-195^\circ < u < 165^\circ$.

3

Find x $u = 18.434 \dots^\circ$ or $u = -161.565 \dots^\circ$, so $x = 16.7^\circ$ or $x = -73.3^\circ$.

4

Rewrite the second equation $4 \sin^2 \theta + 8 \cos \theta = 3$.

5

Use sin squaredSubstitute $\sin^2 \theta = 1 - \cos^2 \theta$ to get $4(1 - \cos^2 \theta) + 8 \cos \theta = 3$.

6

Solve for cos thetaThis gives $4 \cos^2 \theta - 8 \cos \theta - 1 = 0$, so $\cos \theta = 1 \pm \frac{\sqrt{5}}{2}$.

7

Choose the valid cosineOnly $\cos \theta = 1 - \frac{\sqrt{5}}{2}$ lies between -1 and 1 .

8

State thetaFor $0 < \theta < 2\pi$, $\theta = 1.69$ or 4.59 to 3 significant figures.**FINAL ANSWER**

WMA12-01_2022_Jan_Q07

(i) $x = -73.3^\circ, 16.7^\circ$. (ii) $\theta = 1.69, 4.59$.

T10

10_CAST

10_CAST_CORE

CHAPTER

Trigonometric Equations

FAMILY

Trig equations

VARIANT

core family move

6 marks

WMA12/01 MAY/JUNE 2022

Question 5

Past-paper question

WMA12-01_2022_MayJune_Q05

5. In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

Solve, for $-180^\circ < \theta \leq 180^\circ$, the equation

$$3 \tan(\theta + 43^\circ) = 2 \cos(\theta + 43^\circ)$$

(6)

T10 Trigonometric Equations 10_CAST Trig equations

10_CAST_CORE core family
move

WORKED SOLUTION

Question 5

WMA12/01 May/June 2022

WMA12-01_2022_MayJune_Q05

1

Substitute u Let $u = \theta + 43^\circ$. The equation becomes $3 \tan u = 2 \cos u$.

2

Clear the cosine denominator $3 \frac{\sin u}{\cos u} = 2 \cos u$, so $3 \sin u = 2 \cos^2 u$.

3

Use cos squared $3 \sin u = 2(1 - \sin^2 u)$, so $2 \sin^2 u + 3 \sin u - 2 = 0$.

4

Factor the quadratic $(2 \sin u - 1)(\sin u + 2) = 0$.

5

Use the valid sine value $\sin u = -2$ is impossible, so $\sin u = \frac{1}{2}$.

6

Find u in the intervalSince $-180^\circ < \theta \leq 180^\circ$, $-137^\circ < u \leq 223^\circ$. Thus $u = 30^\circ$ or 150° .

7

Return to theta $\theta = u - 43^\circ$, so $\theta = -13^\circ$ or 107° .**FINAL ANSWER** $\theta = -13^\circ, 107^\circ$.

WMA12-01_2022_MayJune_Q05

T10

10_CAST

10_CAST_CORE

CHAPTER
Trigonometric Equations

FAMILY
Trig equations

VARIANT
core family move

7 marks

WMA12/01 OCTOBER 2023

Question 3

Past-paper question

WMA12-01_2023_Oct_Q03

3. In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

(a) Solve, for $0 < \theta \leq 360^\circ$ the equation

$$2 \tan \theta + 3 \sin \theta = 0$$

giving your answers, as appropriate, to one decimal place.

(5)

(b) Hence, or otherwise, find the smallest positive solution of

$$2 \tan(2x + 40^\circ) + 3 \sin(2x + 40^\circ) = 0$$

giving your answer to one decimal place.

(2)

T10 Trigonometric Equations 10_CAST Trig equations

10_CAST_CORE core family
move

WORKED SOLUTION

Question 3

WMA12/01 October 2023

WMA12-01_2023_Oct_Q03

1

Factor the equation $2 \tan \theta + 3 \sin \theta = 0$ becomes $\frac{2 \sin \theta}{\cos \theta} + 3 \sin \theta = 0$.

2

Multiply by cos thetaSince $\tan \theta$ is defined, $\cos \theta \neq 0$. Multiplying by $\cos \theta$ gives $2 \sin \theta + 3 \sin \theta \cos \theta = 0$.

3

Factor $\sin \theta(2 + 3 \cos \theta) = 0$.

4

Solve the sine caseFor $0 < \theta \leq 360^\circ$, $\sin \theta = 0$ gives $\theta = 180^\circ$ or 360° .

5

Solve the cosine case $2 + 3 \cos \theta = 0$ gives $\cos \theta = -\frac{2}{3}$, so $\theta = 131.8^\circ$ or 228.2° to one decimal place.

6

Apply to the new angleFor $2 \tan(2x + 40^\circ) + 3 \sin(2x + 40^\circ) = 0$, the angle $2x + 40^\circ$ must be one of the solution angles, modulo 360° .

7

Find the smallest positive xThe smallest positive value comes from $2x + 40^\circ = 131.8 \dots^\circ$, so $x = \frac{131.8 \dots - 40}{2} = 45.9^\circ$ to one decimal place.**FINAL ANSWER**

WMA12-01_2023_Oct_Q03

(a) $\theta = 131.8^\circ, 180^\circ, 228.2^\circ, 360^\circ$. (b) $x = 45.9^\circ$.

T10

10_CAST

10_CAST_CORE

CHAPTER
Trigonometric Equations

FAMILY
Trig equations

VARIANT
core family move

14 marks

WMA12/01 JANUARY 2024

Question 9

Past-paper question

WMA12-01_2024_Jan_Q09

9. In this question you must show detailed reasoning.

Solutions relying entirely on calculator technology are not acceptable.

- (i) Solve, for $0 \leq x < 360^\circ$, the equation

$$\sin x \tan x = 5$$

giving your answers to one decimal place.

(6)

- (ii)




Figure 1

Figure 1 shows a sketch of part of the curve with equation

$$y = A \sin\left(2\theta - \frac{3\pi}{8}\right) + 2$$

where A is a constant and θ is measured in radians.

The points P , Q and R lie on the curve and are shown in Figure 1.

Given that the y coordinate of P is 7

- (a) state the value of A ,

(1)

- (b) find the exact coordinates of Q ,

(3)

- (c) find the value of θ at R , giving your answer to 3 significant figures.

(4)

T10 Trigonometric Equations 10_CAST Trig equations

10_CAST_CORE core family
move

WORKED SOLUTION

Question 9

WMA12/01 January 2024

WMA12-01_2024_Jan_Q09

1

Rewrite the equation $\sin x \tan x = 5$ becomes $\frac{\sin^2 x}{\cos x} = 5$.

2

Use sine squaredMultiplying by $\cos x$ gives $\sin^2 x = 5 \cos x$. Since $\sin^2 x = 1 - \cos^2 x$, $1 - \cos^2 x = 5 \cos x$.

3

Solve for $\cos x$ $\cos^2 x + 5 \cos x - 1 = 0$, so $\cos x = \frac{-5 \pm \sqrt{29}}{2}$.

4

Choose the possible cosine value $\frac{-5 - \sqrt{29}}{2} < -1$, so the valid value is $\cos x = \frac{-5 + \sqrt{29}}{2}$.

5

Find x For $0 \leq x < 360^\circ$, this gives $x = 78.9^\circ$ or 281.1° to one decimal place.

6

Find A from P The point P is a maximum point on the sine curve. Its y coordinate is 7, so $A + 2 = 7$ and $A = 5$.

7

Locate Q Point Q is the next minimum after P , so $\sin\left(2\theta - \frac{3\pi}{8}\right) = -1$.

8

Find θ at Q For this minimum, $2\theta - \frac{3\pi}{8} = \frac{3\pi}{2}$, so $2\theta = \frac{15\pi}{8}$ and $\theta = \frac{15\pi}{16}$.

9

Find y at Q At the minimum, $y = 2 - 5 = -3$, so $Q = \left(\frac{15\pi}{16}, -3\right)$.

10

Set up R At R , $y = 0$, so $5 \sin\left(2\theta - \frac{3\pi}{8}\right) + 2 = 0$.

11

Choose the shown crossing

Thus $\sin\left(2\theta - \frac{3\pi}{8}\right) = -\frac{2}{5}$. The crossing R shown after the second maximum has $2\theta - \frac{3\pi}{8} = 3\pi + \sin^{-1}\left(\frac{2}{5}\right)$.

12

Find theta at R

$\theta = \frac{1}{2}\left(3\pi + \sin^{-1}\left(\frac{2}{5}\right) + \frac{3\pi}{8}\right) = 5.507\dots$, so $\theta = 5.51$ to 3 significant figures.

FINAL ANSWER

WMA12-01_2024_Jan_Q09

(i) $x = 78.9^\circ, 281.1^\circ$. (ii)(a) $A = 5$. (ii)(b) $Q = \left(\frac{15\pi}{16}, -3\right)$. (ii)(c) $\theta = 5.51$.

DR ESLAM AHMED

T10

10_CAST

10_CAST_CORE

CHAPTER

Trigonometric Equations

FAMILY

Trig equations

VARIANT

core family move

7 marks

WMA12/01 MAY/JUNE (R) 2024

Question 7

Past-paper question

WMA12-01_2024_MayJuneR_Q07

7.

In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.

- (i) Solve, for $0 \leq x < 2\pi$, the equation

$$3 \sin x \tan x = 11 + \cos x$$

giving the answers in radians to 3 decimal places.

(5)

- (ii) Given that

- $0 < \theta < 90^\circ$
- $\cos \theta = \frac{1}{3}$

find, in simplest form, the exact value of $\tan \theta$

(2)

T10 Trigonometric Equations 10_CAST Trig equations

10_CAST_CORE core family
move

WORKED SOLUTION

Question 7

WMA12/01 May/June (R) 2024

WMA12-01_2024_MayJuneR_Q07

1

Rewrite $\tan x$ $3 \sin x \tan x = 11 + \cos x$ becomes $\frac{3 \sin^2 x}{\cos x} = 11 + \cos x$.

2

Multiply by $\cos x$ $3 \sin^2 x = 11 \cos x + \cos^2 x$.

3

Use sine squared $3(1 - \cos^2 x) = 11 \cos x + \cos^2 x$, so $4 \cos^2 x + 11 \cos x - 3 = 0$.

4

Solve for cosine $(4 \cos x - 1)(\cos x + 3) = 0$, so $\cos x = \frac{1}{4}$ or $\cos x = -3$.

5

Reject impossible value $\cos x = -3$ is impossible, so $\cos x = \frac{1}{4}$.

6

Find x valuesFor $0 \leq x < 2\pi$, this gives $x = 1.318 \dots$ or $x = 4.965 \dots$, so $x = 1.318, 4.965$ radians.

7

Use the right triangle identityGiven $0 < \theta < 90^\circ$ and $\cos \theta = \frac{1}{3}$, $\sin \theta = \sqrt{1 - \frac{1}{9}} = \frac{2\sqrt{2}}{3}$.

8

Find $\tan \theta$ $\tan \theta = \frac{\sin \theta}{\cos \theta} = \frac{2\sqrt{2}/3}{1/3} = 2\sqrt{2}$.**FINAL ANSWER**

WMA12-01_2024_MayJuneR_Q07

(i) $x = 1.318, 4.965$. (ii) $2\sqrt{2}$.

T10

10_CAST

10_CAST_CORE

CHAPTER
Trigonometric Equations

FAMILY
Trig equations

VARIANT
core family move

8 marks

WMA12/01 MAY/JUNE 2025

Question 8

Past-paper question

WMA12-01_2025_MayJune_Q08

8: **In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.**

(i) Solve, for $0 \leq x < \pi$, the equation

$$3 \tan \left(2x + \frac{\pi}{5} \right) = \sqrt{3}$$

giving the answers in radians in the form $k\pi$, where k is a rational constant to be found.

(3)

(ii) Solve, for $0 \leq \theta < 360^\circ$, the equation

$$5 \sin \theta \tan \theta = \cos \theta + 4$$

giving your answers, in degrees, to one decimal place.

(5)

T10 Trigonometric Equations 10_CAST Trig equations

10_CAST_CORE core family
move

WORKED SOLUTION

Question 8

WMA12/01 May/June 2025

WMA12-01_2025_MayJune_Q08

1

Solve the tangent equation

$$3 \tan \left(2x + \frac{\pi}{5} \right) = \sqrt{3} \text{ gives } \tan \left(2x + \frac{\pi}{5} \right) = \frac{1}{\sqrt{3}}.$$

2

Write the general angle

$$\text{Therefore } 2x + \frac{\pi}{5} = \frac{\pi}{6} + n\pi.$$

3

Find x

$$2x = -\frac{\pi}{30} + n\pi, \text{ so } x = -\frac{\pi}{60} + \frac{n\pi}{2}.$$

4

Use the interval

$$\text{For } 0 \leq x < \pi, \text{ the possible values are } x = \frac{29\pi}{60} \text{ and } x = \frac{59\pi}{60}.$$

5

Rewrite the second equation

$$\text{Using } \tan \theta = \frac{\sin \theta}{\cos \theta}, 5 \sin \theta \tan \theta = \cos \theta + 4 \text{ becomes } \frac{5 \sin^2 \theta}{\cos \theta} = \cos \theta + 4.$$

6

Form an equation in cos theta

$$\text{Multiplying by } \cos \theta \text{ gives } 5 \sin^2 \theta = \cos^2 \theta + 4 \cos \theta. \text{ Using } \sin^2 \theta = 1 - \cos^2 \theta \text{ gives } 6 \cos^2 \theta + 4 \cos \theta - 5 = 0.$$

7

Solve for cos theta

$$\cos \theta = \frac{-2 \pm \sqrt{34}}{6}. \text{ The negative option is less than } -1, \text{ so reject it.}$$

8

Find theta

$$\cos \theta = \frac{-2 + \sqrt{34}}{6} \text{ gives } \theta = 50.320 \dots^\circ \text{ or } 309.679 \dots^\circ.$$

9

Round the answers

$$\text{To one decimal place, } \theta = 50.3^\circ \text{ or } 309.7^\circ.$$

FINAL ANSWER

$$(i) x = \frac{29\pi}{60}, \frac{59\pi}{60}. \quad (ii) \theta = 50.3^\circ, 309.7^\circ.$$

WMA12-01_2025_MayJune_Q08

T10

10_CAST

10_CAST_CORE

CHAPTER

Trigonometric Equations

FAMILY

Trig equations

VARIANT

core family move

9 marks

WMA12/01 OCTOBER 2025

Question 5

Past-paper question

WMA12-01_2025_Oct_Q05

5. In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.

- (i) Solve, for $0 < \theta \leq 360^\circ$, the equation

$$4 \tan \theta + 5 \sin \theta = 0$$

giving any non-exact answers to one decimal place.

(5)

- (ii) Solve, for $0 < x < \pi$, the equation

$$\frac{\sin x}{\cos x} + \frac{\cos x}{\sin x} = \frac{5}{\cos x}$$

giving the answers, in radians, to 3 significant figures.

(4)

T10 Trigonometric Equations 10_CAST Trig equations

10_CAST_CORE core family
move

WORKED SOLUTION

Question 5

WMA12/01 October 2025

WMA12-01_2025_Oct_Q05

1

Rewrite the first equation $4 \tan \theta + 5 \sin \theta = 0$ gives $4 \frac{\sin \theta}{\cos \theta} + 5 \sin \theta = 0$.

2

Factor sin theta $\sin \theta \left(\frac{4}{\cos \theta} + 5 \right) = 0$.

3

Use sin theta equals zeroIn $0 < \theta \leq 360^\circ$, $\sin \theta = 0$ gives $\theta = 180^\circ$ or 360° .

4

Use the other factor $\frac{4}{\cos \theta} + 5 = 0$ gives $\cos \theta = -\frac{4}{5}$.

5

Find the remaining anglesFor $0 < \theta \leq 360^\circ$, $\cos \theta = -\frac{4}{5}$ gives $\theta = 143.130 \dots^\circ$ or $216.869 \dots^\circ$.

6

Round part iThe non-exact angles are 143.1° and 216.9° to one decimal place.

7

Clear the fractions in part ii $\frac{\sin x}{\cos x} + \frac{\cos x}{\sin x} = \frac{5}{\cos x}$. Multiplying by $\sin x \cos x$ gives $\sin^2 x + \cos^2 x = 5 \sin x$.

8

Use the identity $\sin^2 x + \cos^2 x = 1$, so $1 = 5 \sin x$ and $\sin x = \frac{1}{5}$.

9

Solve in the intervalFor $0 < x < \pi$, $x = \sin^{-1} \left(\frac{1}{5} \right)$ or $x = \pi - \sin^{-1} \left(\frac{1}{5} \right)$.

10

Round part iiThus $x = 0.201357 \dots$ or $2.94023 \dots$, so $x = 0.201$ or 2.94 to 3 significant figures.

FINAL ANSWER

WMA12-01_2025_Oct_Q05

(i) $\theta = 143.1^\circ, 180^\circ, 216.9^\circ, 360^\circ$. (ii) $x = 0.201, 2.94$.

DR ESLAM AHMED

TOPIC 10 | FAMILY 02

Trig Identities

WHAT THIS FAMILY ASKS

Recognise the trig identities demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

2 classified questions

15 total marks

T10

10_IDENT

10_IDENT_CORE

CHAPTER

Trigonometric Equations

FAMILY

Trig identities

VARIANT

core family move

7 marks

WMA12/01 JANUARY 2020

Question 7

Past-paper question

WMA12-01_2020_Jan_Q07

7. (a) Show that the equation

$$8 \tan \theta = 3 \cos \theta$$

may be rewritten in the form

$$3 \sin^2 \theta + 8 \sin \theta - 3 = 0$$

(3)

- (b) Hence solve, for $0 \leq x \leq 90^\circ$, the equation

$$8 \tan 2x = 3 \cos 2x$$

giving your answers to 2 decimal places.

(4)

T10 Trigonometric Equations 10_IDENT Trig identities

10_IDENT_CORE core family
move

WORKED SOLUTION

Question 7

WMA12/01 January 2020

WMA12-01_2020_Jan_Q07

1

Rewrite tan theta $8 \tan \theta = 3 \cos \theta$ becomes $8 \frac{\sin \theta}{\cos \theta} = 3 \cos \theta$.

2

Clear the denominatorMultiplying by $\cos \theta$ gives $8 \sin \theta = 3 \cos^2 \theta$.

3

Use the identitySince $\cos^2 \theta = 1 - \sin^2 \theta$, $8 \sin \theta = 3(1 - \sin^2 \theta)$, so $3 \sin^2 \theta + 8 \sin \theta - 3 = 0$.

4

Apply the result to 2xFor $8 \tan 2x = 3 \cos 2x$, we get $3 \sin^2(2x) + 8 \sin(2x) - 3 = 0$.

5

Solve for sin 2x $(3 \sin 2x - 1)(\sin 2x + 3) = 0$. Since $\sin 2x$ cannot be -3 , $\sin 2x = \frac{1}{3}$.

6

Find xFor $0 \leq x \leq 90^\circ$, $0 \leq 2x \leq 180^\circ$. Thus $2x = 19.471 \dots^\circ$ or $160.529 \dots^\circ$, so $x = 9.74^\circ$ or 80.26° .**FINAL ANSWER**

WMA12-01_2020_Jan_Q07

(a) **Proven.** (b) $x = 9.74^\circ, 80.26^\circ$.

T10

10_IDENT

10_IDENT_CORE

CHAPTER

Trigonometric Equations

FAMILY

Trig identities

VARIANT

core family move

8 marks

WMA12/01 JANUARY 2025

Question 10

Past-paper question

WMA12-01_2025_Jan_Q10

10. In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.

(a) Show that

$$\cos \theta \left(3 \tan \theta + \frac{2}{\tan \theta} \right) \equiv \sin \theta + \frac{2}{\sin \theta} \quad \theta \neq \frac{n\pi}{2} \quad (4)$$

(b) Hence solve, for $0 < x < 2\pi$, the equation

$$\cos x \left(3 \tan x + \frac{2}{\tan x} \right) = 4 \sin x - 5$$

giving your answers to 3 significant figures.

(4)

T10 Trigonometric Equations 10_IDENT Trig identities

10_IDENT_CORE core family
move

WORKED SOLUTION

Question 10

WMA12/01 January 2025

WMA12-01_2025_Jan_Q10

1

Expand the left side

$$\cos \theta \left(3 \tan \theta + \frac{2}{\tan \theta} \right) = 3 \sin \theta + \frac{2 \cos \theta}{\tan \theta}.$$

2

Rewrite the second term

$$\text{Since } \tan \theta = \frac{\sin \theta}{\cos \theta}, \frac{2 \cos \theta}{\tan \theta} = \frac{2 \cos^2 \theta}{\sin \theta}.$$

3

Use the identity

$$3 \sin \theta + \frac{2 \cos^2 \theta}{\sin \theta} = 3 \sin \theta + \frac{2(1 - \sin^2 \theta)}{\sin \theta}.$$

4

Simplify

$$3 \sin \theta + \frac{2}{\sin \theta} - 2 \sin \theta = \sin \theta + \frac{2}{\sin \theta}, \text{ as required.}$$

5

Use part a

$$\text{The equation becomes } \sin x + \frac{2}{\sin x} = 4 \sin x - 5.$$

6

Form a quadratic in $\sin x$

$$\text{Multiplying by } \sin x \text{ gives } \sin^2 x + 2 = 4 \sin^2 x - 5 \sin x, \text{ so } 3 \sin^2 x - 5 \sin x - 2 = 0.$$

7

Solve for $\sin x$

$$(3 \sin x + 1)(\sin x - 2) = 0. \text{ Since } \sin x = 2 \text{ is impossible, } \sin x = -\frac{1}{3}.$$

8

Find x in the interval

$$\text{For } 0 < x < 2\pi, \sin x = -\frac{1}{3} \text{ gives } x = \pi + \sin^{-1}\left(\frac{1}{3}\right) \text{ or } x = 2\pi - \sin^{-1}\left(\frac{1}{3}\right).$$

9

Round the answers

$$\text{Thus } x = 3.481 \dots \text{ or } x = 5.943 \dots, \text{ so to 3 significant figures } x = 3.48 \text{ or } x = 5.94.$$

FINAL ANSWER

WMA12-01_2025_Jan_Q10

(a) Proven. (b) $x = 3.48, 5.94$.

TOPIC 10 | FAMILY 03

Periodicity

WHAT THIS FAMILY ASKS

Recognise the periodicity demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

1 classified questions

10 total marks

T10

10_PERIOD

10_PERIOD_CORE

CHAPTER

Trigonometric Equations

FAMILY

Periodicity

VARIANT

core family move

10 marks

WMA12/01 OCTOBER 2021

Question 10

Past-paper question

WMA12-01_2021_Oct_Q10

10. In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

- (i) Solve, for $-\frac{\pi}{2} < x < \frac{\pi}{2}$

$$\tan^2\left(2x + \frac{\pi}{4}\right) = 3 \quad (5)$$

- (ii) Solve, for $0 < \theta < 360^\circ$

$$(2 \sin \theta - \cos \theta)^2 = 1$$

giving your answers, as appropriate, to one decimal place.

(5)

(Total 10 marks)

T10 Trigonometric Equations 10_PERIOD Periodicity

10_PERIOD_CORE core
family move

WORKED SOLUTION

Question 10

WMA12/01 October 2021

WMA12-01_2021_Oct_Q10

1

Take the square rootLet $u = 2x + \frac{\pi}{4}$. From $\tan^2 u = 3$, $\tan u = \pm\sqrt{3}$.

2

Write the angle familiesThus $u = \frac{\pi}{3} + n\pi$ or $u = -\frac{\pi}{3} + n\pi$, where n is an integer.

3

Return to x So $x = \frac{\pi}{24} + \frac{n\pi}{2}$ or $x = -\frac{7\pi}{24} + \frac{n\pi}{2}$.

4

Use the intervalFor $-\frac{\pi}{2} < x < \frac{\pi}{2}$, the solutions are $x = -\frac{11\pi}{24}, -\frac{7\pi}{24}, \frac{\pi}{24}, \frac{5\pi}{24}$.

5

Expand the squared equation $(2 \sin \theta - \cos \theta)^2 = 1$ gives $4 \sin^2 \theta - 4 \sin \theta \cos \theta + \cos^2 \theta = 1$.

6

Use the identitySince $1 = \sin^2 \theta + \cos^2 \theta$, this simplifies to $3 \sin^2 \theta - 4 \sin \theta \cos \theta = 0$.

7

Factor $\sin \theta (3 \sin \theta - 4 \cos \theta) = 0$.

8

Solve the two cases $\sin \theta = 0$ gives $\theta = 180^\circ$ in the interval. Also $3 \sin \theta = 4 \cos \theta$ gives $\tan \theta = \frac{4}{3}$.

9

State thetaFor $0 < \theta < 360^\circ$, $\tan \theta = \frac{4}{3}$ gives $\theta = 53.1^\circ$ and 233.1° . Therefore $\theta = 53.1^\circ, 180^\circ, 233.1^\circ$.**FINAL ANSWER**

WMA12-01_2021_Oct_Q10

(i) $x = -\frac{11\pi}{24}, -\frac{7\pi}{24}, \frac{\pi}{24}, \frac{5\pi}{24}$. (ii) $\theta = 53.1^\circ, 180^\circ, 233.1^\circ$.

TOPIC 10 | FAMILY 04

Transformed Trig

WHAT THIS FAMILY ASKS

Recognise the transformed trig demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

V02 Expand the interval for the inside expression before finding angles.

8 classified questions

69 total marks

T10

10_TRANS

10_TRANS_CORE

CHAPTER

Trigonometric Equations

FAMILY

Transformed trig

VARIANT

core family move

10 marks

WMA12/01 MAY/JUNE 2021

Question 8

Past-paper question

WMA12-01_2021_MayJune_Q08

8. In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

- (i) Solve, for $0 < \theta < 360^\circ$, the equation

$$3 \sin(\theta + 30^\circ) = 7 \cos(\theta + 30^\circ)$$

giving your answers to one decimal place.

(4)

- (ii) (a) Show that the equation

$$3 \sin^3 x = 5 \sin x - 7 \sin x \cos x$$

can be written in the form

$$\sin x (a \cos^2 x + b \cos x + c) = 0$$

where a , b and c are constants to be found.

- (b) Hence solve for $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ the equation

$$3 \sin^3 x = 5 \sin x - 7 \sin x \cos x$$

(6)

T10 Trigonometric Equations

10_TRANS

Transformed trig

10_TRANS_CORE core family
move

WORKED SOLUTION

Question 8

WMA12/01 May/June 2021

WMA12-01_2021_MayJune_Q08

1

Solve for theta plus thirtyLet $u = \theta + 30^\circ$. Then $3 \sin u = 7 \cos u$, so $\tan u = \frac{7}{3}$.

2

Find u in the intervalSince $0 < \theta < 360^\circ$, $30^\circ < u < 390^\circ$. Thus $u = 66.801 \dots^\circ$ or $246.801 \dots^\circ$.

3

Return to theta $\theta = u - 30^\circ$, giving $\theta = 36.8^\circ$ and 216.8° to one decimal place.

4

Rearrange the second equation $3 \sin^3 x = 5 \sin x - 7 \sin x \cos x$ gives $3 \sin^3 x - 5 \sin x + 7 \sin x \cos x = 0$.

5

Factor out sine $\sin x(3 \sin^2 x - 5 + 7 \cos x) = 0$.

6

Use sin squaredSince $\sin^2 x = 1 - \cos^2 x$, this becomes $\sin x(-3 \cos^2 x + 7 \cos x - 2) = 0$, or $\sin x(3 \cos^2 x - 7 \cos x + 2) = 0$.

7

Solve in the intervalFor $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$, $\sin x = 0$ gives $x = 0$. Also $3 \cos^2 x - 7 \cos x + 2 = 0$ gives $(3 \cos x - 1)(\cos x - 2) = 0$.

8

State the solutions $\cos x = 2$ is impossible, so $\cos x = \frac{1}{3}$. The solutions are $x = -\cos^{-1}\left(\frac{1}{3}\right)$, 0, and $\cos^{-1}\left(\frac{1}{3}\right)$.**FINAL ANSWER**

WMA12-01_2021_MayJune_Q08

(i) $\theta = 36.8^\circ, 216.8^\circ$. (ii) (a) $a = 3, b = -7, c = 2$. (ii) (b) $x = -\cos^{-1}\left(\frac{1}{3}\right), 0, \cos^{-1}\left(\frac{1}{3}\right)$.

T10

10_TRANS

10_TRANS_CORE

CHAPTER

Trigonometric Equations

FAMILY

Transformed trig

VARIANT

core family move

9 marks

WMA12/01 JANUARY 2023

Question 8

Past-paper question

WMA12-01_2023_Jan_Q08

8. In this question you must show all stages of your working.

Solutions based entirely on calculator technology are not acceptable.

- (i) Solve, for $-\frac{\pi}{2} < x < \pi$, the equation

$$5 \sin(3x + 0.1) + 2 = 0$$

giving your answers, **in radians**, to 2 decimal places.

(4)

- (ii) Solve, for $0 < \theta < 360^\circ$, the equation

$$2 \tan \theta \sin \theta = 5 + \cos \theta$$

giving your answers, **in degrees**, to one decimal place.

(5)

T10 Trigonometric Equations

10_TRANS

Transformed trig

10_TRANS_CORE

core family

move

WORKED SOLUTION

Question 8

WMA12/01 January 2023

WMA12-01_2023_Jan_Q08

1

Isolate the sine term $5 \sin(3x + 0.1) + 2 = 0$ gives $\sin(3x + 0.1) = -0.4$.

2

Find the angle intervalSince $-\frac{\pi}{2} < x < \pi$, multiplying by 3 and adding 0.1 gives $-\frac{3\pi}{2} + 0.1 < 3x + 0.1 < 3\pi + 0.1$.

3

Solve for the inner angleIn this interval, $3x + 0.1 = -2.7301 \dots, -0.4115 \dots, 3.5531 \dots, 5.8717 \dots$

4

Find x valuesSubtract 0.1 and divide by 3: $x = -0.9433 \dots, -0.1705 \dots, 1.1510 \dots, 1.9239 \dots$

5

Round the radian answersTo 2 decimal places, $x = -0.94, -0.17, 1.15, 1.92$.

6

Rewrite the second equation $2 \tan \theta \sin \theta = 5 + \cos \theta$. Using $\tan \theta = \frac{\sin \theta}{\cos \theta}$ gives $\frac{2 \sin^2 \theta}{\cos \theta} = 5 + \cos \theta$.

7

Use sine squaredMultiplying by $\cos \theta$ gives $2 \sin^2 \theta = 5 \cos \theta + \cos^2 \theta$. Substitute $\sin^2 \theta = 1 - \cos^2 \theta$.

8

Solve for cos theta $2(1 - \cos^2 \theta) = 5 \cos \theta + \cos^2 \theta$, so $3 \cos^2 \theta + 5 \cos \theta - 2 = 0$.

9

Factor and reject impossible value $(3 \cos \theta - 1)(\cos \theta + 2) = 0$. Since $\cos \theta = -2$ is impossible, $\cos \theta = \frac{1}{3}$.

10

Find thetaFor $0 < \theta < 360^\circ$, $\cos \theta = \frac{1}{3}$ gives $\theta = 70.5^\circ$ or 289.5° to one decimal place.

FINAL ANSWER

WMA12-01_2023_Jan_Q08

(i) $x = -0.94, -0.17, 1.15, 1.92$. (ii) $\theta = 70.5^\circ, 289.5^\circ$.

DR ESLAM AHMED

T10

10_TRANS

10_TRANS_CORE

CHAPTER

Trigonometric Equations

FAMILY

Transformed trig

VARIANT

core family move

12 marks

WMA12/01 MAY/JUNE 2024

Question 8

Past-paper question

WMA12-01_2024_MayJune_Q08

8. In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.

- (i) Solve, for $0 < x \leq \pi$, the equation

$$5 \sin x \tan x + 13 = \cos x$$

giving your answer in radians to 3 significant figures.

(5)

- (ii) The temperature inside a greenhouse is monitored on one particular day.

The temperature, $H^\circ\text{C}$, inside the greenhouse, t hours after midnight, is modelled by the equation

$$H = 10 + 12 \sin(kt + 18)^\circ \quad 0 \leq t < 24$$

where k is a constant.

Use the equation of the model to answer parts (a) to (c).

Given that

- the temperature inside the greenhouse was 20°C at 6 am
- $0 < k < 20$

- (a) find all possible values for k , giving each answer to 2 decimal places.

(4)

Given further that $0 < k < 10$

- (b) find the maximum temperature inside the greenhouse,
- (c) find the time of day at which this maximum temperature occurs.

(1)

Give your answer to the nearest minute.

(2)

T10 Trigonometric Equations

10_TRANS

Transformed trig

10_TRANS_CORE

core family

move

WORKED SOLUTION

Question 8

WMA12/01 May/June 2024

WMA12-01_2024_MayJune_Q08

1

Rewrite the trig equation
 $5 \sin x \tan x + 13 = \cos x$ becomes $\frac{5 \sin^2 x}{\cos x} + 13 = \cos x$.

2

Multiply by $\cos x$
 $5 \sin^2 x + 13 \cos x = \cos^2 x$.

3

Use sine squared
 $5(1 - \cos^2 x) + 13 \cos x = \cos^2 x$, so $6 \cos^2 x - 13 \cos x - 5 = 0$.

4

Solve for $\cos x$
 $(2 \cos x - 5)(3 \cos x + 1) = 0$, so $\cos x = \frac{5}{2}$ or $\cos x = -\frac{1}{3}$.

5

Choose the valid root
 $\cos x = \frac{5}{2}$ is impossible, so $\cos x = -\frac{1}{3}$. For $0 < x \leq \pi$, $x = 1.9106 \dots$, so $x = 1.91$ radians to 3 significant figures.

6

Use the temperature at 6 am
At $t = 6$, $20 = 10 + 12 \sin(6k + 18)^\circ$, so $\sin(6k + 18)^\circ = \frac{5}{6}$.

7

Solve for k
 $\sin^{-1}\left(\frac{5}{6}\right) = 56.442 \dots^\circ$. Since $0 < k < 20$, $18 < 6k + 18 < 138$, so $6k + 18 = 56.442 \dots$ or $123.557 \dots$

8

Find possible k values
This gives $k = 6.407 \dots$ or $k = 17.592 \dots$, so $k = 6.41$ or 17.59 to 2 decimal places.

9

Use the tighter condition
Given further that $0 < k < 10$, use $k = 6.407 \dots$

10

Find the maximum temperature
The maximum value of $\sin$ is 1, so the maximum temperature is $10 + 12 = 22^\circ\text{C}$.

11

Find the time of maximum

The maximum occurs when $kt + 18 = 90$, so $t = \frac{72}{6.407...} = 11.2375 \dots$ hours after midnight.

12

Convert to time of day

0.2375 ... hours is 14.25 ... minutes, so the maximum occurs at about 11:14 am.

FINAL ANSWER

WMA12-01_2024_MayJune_Q08

(i) $x = 1.91$ **radians.** (ii)(a) $k = 6.41$ **or** 17.59. (ii)(b) 22° **C.** (ii)(c) 11:14 **am.**

DR ESLAM AHMED

T10

10_TRANS

10_TRANS_CORE

CHAPTER

Trigonometric Equations

FAMILY

Transformed trig

VARIANT

core family move

6 marks

WMA12/01 MAY/JUNE (R) 2024

Question 5

Past-paper question

WMA12-01_2024_MayJuneR_Q05

5.

**In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.**

The depth of water, D metres, in a harbour on a particular day is given by the equation

$$D = 8 + 5 \sin\left(\frac{\pi t}{6} + 3\right) \quad 0 \leq t < 24$$

where t is the number of hours after **midnight**.

- (a) Show that the depth of water in the harbour at 2 am is just over 4 metres. (1)
- (b) Find, to the nearest minute, the first time after **midday** when the depth of water in the harbour is exactly 6 metres. (5)

T10 Trigonometric Equations

10_TRANS

Transformed trig

10_TRANS_CORE core family
move

WORKED SOLUTION

Question 5

WMA12/01 May/June (R) 2024

WMA12-01_2024_MayJuneR_Q05

1

Substitute t equals two

$$D = 8 + 5 \sin \left(\frac{\pi(2)}{6} + 3 \right) = 8 + 5 \sin \left(\frac{\pi}{3} + 3 \right).$$

2

Evaluate

$D = 4.064 \dots$, so the depth at 2 am is just over 4 metres.

3

Set depth equal to six

$$6 = 8 + 5 \sin \left(\frac{\pi t}{6} + 3 \right), \text{ so } \sin \left(\frac{\pi t}{6} + 3 \right) = -\frac{2}{5}.$$

4

Use the time interval

After midday means $t > 12$. Then $\frac{\pi t}{6} + 3$ is greater than $2\pi + 3$.

5

Choose the first suitable angle

The first angle after $2\pi + 3$ with sine $-\frac{2}{5}$ is $3\pi + \sin^{-1} \left(\frac{2}{5} \right)$.

6

Solve for t

$$\frac{\pi t}{6} + 3 = 3\pi + \sin^{-1} \left(\frac{2}{5} \right), \text{ so } t = 13.056 \dots \text{ hours after midnight.}$$

7

Convert to clock time

$0.056 \dots$ hours is about 3.38 minutes, so the time to the nearest minute is 13:03, or 1:03 pm.

FINAL ANSWER

WMA12-01_2024_MayJuneR_Q05

(a) $D(2) = 4.06 \dots$ m, just over 4 m. (b) 13:03, or 1:03 pm.

T10

10_TRANS

10_TRANS_CORE

CHAPTER

Trigonometric Equations

FAMILY

Transformed trig

VARIANT

core family move

7 marks

WMA12/01 OCTOBER 2024

Question 9

Past-paper question

WMA12-01_2024_Oct_Q09

9. In this question you must show detailed reasoning.

Solutions relying entirely on calculator technology are not acceptable.

(a) Show that the equation

$$2 \tan \theta = 3 \cos \theta$$

can be written as

$$3 \sin^2 \theta + 2 \sin \theta - 3 = 0$$

(3)

(b) Hence solve, for $-\pi < x < \pi$, the equation

$$2 \tan \left(2x + \frac{\pi}{3} \right) = 3 \cos \left(2x + \frac{\pi}{3} \right)$$

giving your answers to 3 significant figures.

(4)

T10 Trigonometric Equations

10_TRANS

Transformed trig

10_TRANS_CORE core family
move

WORKED SOLUTION

Question 9

WMA12/01 October 2024

WMA12-01_2024_Oct_Q09

1

Write tan in terms of sin and cos $2 \tan \theta = 3 \cos \theta$ gives $2 \frac{\sin \theta}{\cos \theta} = 3 \cos \theta$.

2

Multiply by cos thetaSince $\tan \theta$ is defined, $\cos \theta \neq 0$. Hence $2 \sin \theta = 3 \cos^2 \theta$.

3

Use the identity $2 \sin \theta = 3(1 - \sin^2 \theta)$, so $3 \sin^2 \theta + 2 \sin \theta - 3 = 0$.

4

Substitute the new angleLet $\theta = 2x + \frac{\pi}{3}$. Then $3 \sin^2 \theta + 2 \sin \theta - 3 = 0$.

5

Solve for sin theta $\sin \theta = \frac{-2 \pm \sqrt{40}}{6} = \frac{-1 \pm \sqrt{10}}{3}$. The root $\frac{-1 - \sqrt{10}}{3}$ is less than -1 , so reject it.

6

Find the angle rangeSince $-\pi < x < \pi$, $-\frac{5\pi}{3} < \theta < \frac{7\pi}{3}$.

7

Find the theta valuesFor $\sin \theta = \frac{-1 + \sqrt{10}}{3}$ in this range, $\theta \approx -3.94649, 0.804897, 2.33670$, and 7.08808 .

8

Convert back to xUsing $x = \frac{\theta - \pi/3}{2}$ gives $x \approx -2.49684, -0.121150, 0.644749$, and 3.02044 .

9

Round the answersTo 3 significant figures, $x = -2.50, -0.121, 0.645$, and 3.02 .**FINAL ANSWER**

WMA12-01_2024_Oct_Q09

(a) **Proven.** (b) $x = -2.50, -0.121, 0.645, 3.02$.

T10

10_TRANS

10_TRANS_HV01

CHAPTER

Trigonometric Equations

FAMILY

Transformed trig

VARIANT

Expand the interval for the
inside expression before
finding angles.

8 marks

WMA12/01 OCTOBER 2022

Question 5

Past-paper question

WMA12-01_2022_Oct_Q05

5.**In this question you must show detailed reasoning.****Solutions relying entirely on calculator technology are not acceptable.**

(a) Show that the equation

$$(3 \cos \theta - \tan \theta) \cos \theta = 2$$

can be written as

$$3 \sin^2 \theta + \sin \theta - 1 = 0$$

(3)(b) Hence solve for $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$

$$(3 \cos 2x - \tan 2x) \cos 2x = 2$$

(5)

T10 Trigonometric Equations 10_TRANS Transformed trig

10_TRANS_HV01 Expand the interval for the inside expression before finding angles.

WORKED SOLUTION

Question 5

WMA12/01 October 2022

WMA12-01_2022_Oct_Q05

1

Rewrite the left side $(3 \cos \theta - \tan \theta) \cos \theta = 3 \cos^2 \theta - \sin \theta$ because $\tan \theta \cos \theta = \sin \theta$.

2

Move everything to one sideThe equation becomes $3 \cos^2 \theta - \sin \theta = 2$.

3

Use cos squaredSince $\cos^2 \theta = 1 - \sin^2 \theta$, $3(1 - \sin^2 \theta) - \sin \theta = 2$.

4

Show the required formExpanding and rearranging gives $3 \sin^2 \theta + \sin \theta - 1 = 0$.

5

Apply the result to 2xFor $(3 \cos 2x - \tan 2x) \cos 2x = 2$, take $\theta = 2x$. Then $3 \sin^2(2x) + \sin(2x) - 1 = 0$.

6

Solve the quadraticLet $u = \sin(2x)$. Then $3u^2 + u - 1 = 0$, so $u = \frac{-1 \pm \sqrt{13}}{6}$.

7

Use the intervalSince $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$, we have $-\pi \leq 2x \leq \pi$. Solving $\sin(2x) = \frac{-1 + \sqrt{13}}{6}$ and $\sin(2x) = \frac{-1 - \sqrt{13}}{6}$ in this interval gives $2x = -2.2665 \dots, -0.8751 \dots, 0.4492 \dots, 2.6924 \dots$

8

Divide by twoTherefore $x = -1.133 \dots, -0.4375 \dots, 0.2246 \dots, 1.346 \dots$, giving $x = -1.13, -0.438, 0.225, 1.35$ radians.**FINAL ANSWER**

WMA12-01_2022_Oct_Q05

(a) Proven. (b) $x = -1.13, -0.438, 0.225, 1.35$ radians.

T10

10_TRANS

10_TRANS_HV01

CHAPTER

Trigonometric Equations

FAMILY

Transformed trig

VARIANT

Expand the interval for the
inside expression before
finding angles.

7 marks

WMA12/01 MAY/JUNE 2023

Question 9

Past-paper question

WMA12-01_2023_MayJune_Q09

9.

In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

(a) Show that

$$3 \cos \theta (\tan \theta \sin \theta + 3) = 11 - 5 \cos \theta$$

may be written as

$$3 \cos^2 \theta - 14 \cos \theta + 8 = 0 \quad (3)$$

(b) Hence solve, for $0 < x < 360^\circ$

$$3 \cos 2x (\tan 2x \sin 2x + 3) = 11 - 5 \cos 2x$$

giving your answers to one decimal place.

(4)

T10 Trigonometric Equations 10_TRANS Transformed trig

10_TRANS_HV01 Expand the interval for the inside expression before finding angles.

WORKED SOLUTION

Question 9

WMA12/01 May/June 2023

WMA12-01_2023_MayJune_Q09

1

Rewrite $\tan \theta \sin \theta$

$$\tan \theta \sin \theta = \frac{\sin^2 \theta}{\cos \theta}.$$

2

Simplify the left side

$$3 \cos \theta (\tan \theta \sin \theta + 3) = 3 \sin^2 \theta + 9 \cos \theta.$$

3

Use sine squared

$$\text{The equation becomes } 3(1 - \cos^2 \theta) + 9 \cos \theta = 11 - 5 \cos \theta.$$

4

Rearrange

$$3 - 3 \cos^2 \theta + 9 \cos \theta = 11 - 5 \cos \theta, \text{ so } 3 \cos^2 \theta - 14 \cos \theta + 8 = 0.$$

5

Apply the result to $2x$

$$\text{For the equation in part (b), take } \theta = 2x. \text{ Then } 3 \cos^2(2x) - 14 \cos(2x) + 8 = 0.$$

6

Solve the quadratic

$$\text{Let } u = \cos(2x). \text{ Then } 3u^2 - 14u + 8 = 0, \text{ so } (3u - 2)(u - 4) = 0.$$

7

Reject the impossible value

$$u = 4 \text{ is impossible for cosine, so } \cos(2x) = \frac{2}{3}.$$

8

Use the interval

$$\text{Since } 0 < x < 360^\circ, \text{ we have } 0 < 2x < 720^\circ.$$

9

Find x

In this interval, $2x = 48.189 \dots^\circ, 311.810 \dots^\circ, 408.189 \dots^\circ, 671.810 \dots^\circ$. Dividing by 2 gives $x = 24.1^\circ, 155.9^\circ, 204.1^\circ, 335.9^\circ$ to one decimal place.

FINAL ANSWER

WMA12-01_2023_MayJune_Q09

(a) **Proven.** (b) $x = 24.1^\circ, 155.9^\circ, 204.1^\circ, 335.9^\circ$.

T10

10_TRANS

10_TRANS_HV01

CHAPTER

Trigonometric Equations

FAMILY

Transformed trig

VARIANT

Expand the interval for the
inside expression before
finding angles.

10 marks

WMA12/01 JANUARY 2026

Question 8

Past-paper question

WMA12-01_2026_Jan_Q08

8.

**In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.**

- (i) Solve, for $0 \leq \theta < 180$

$$4 \tan^2(2\theta - 30)^\circ + 1 = 49$$

giving the answers to one decimal place.

(5)

- (ii) Solve, for $0 \leq x < 2\pi$

$$2 \tan x \sin x + 3 = 0$$

giving the answers, in radians, in the form $k\pi$ where k is a rational constant.

(5)

T10 Trigonometric Equations 10_TRANS Transformed trig

10_TRANS_HV01 Expand the interval for the inside expression before finding angles.

WORKED SOLUTION

Question 8

WMA12/01 January 2026

WMA12-01_2026_Jan_Q08

1

Make the tangent expression the subject $4 \tan^2(2\theta - 30^\circ) + 1 = 49$ gives $\tan^2(2\theta - 30^\circ) = 12$.

2

Find the possible angle valuesLet $\alpha = 2\theta - 30^\circ$. Since $0 \leq \theta < 180^\circ$, $-30^\circ \leq \alpha < 330^\circ$. The solutions of $\tan \alpha = \pm\sqrt{12}$ in this range are approximately $73.9^\circ, 106.1^\circ, 253.9^\circ, 286.1^\circ$.

3

Return to theta $\theta = \frac{\alpha + 30^\circ}{2}$, giving $\theta = 51.9^\circ, 68.1^\circ, 141.9^\circ, 158.1^\circ$ to one decimal place.

4

Rewrite tan x $2 \tan x \sin x + 3 = 0$ becomes $2 \left(\frac{\sin x}{\cos x} \right) \sin x + 3 = 0$.

5

Form a quadratic in cos xMultiplying by $\cos x$ gives $2 \sin^2 x + 3 \cos x = 0$. Using $\sin^2 x = 1 - \cos^2 x$, $2(1 - \cos^2 x) + 3 \cos x = 0$, so $2 \cos^2 x - 3 \cos x - 2 = 0$.

6

Solve for x $(2 \cos x + 1)(\cos x - 2) = 0$. Since $\cos x = 2$ is impossible, $\cos x = -\frac{1}{2}$. In $0 \leq x < 2\pi$, $x = \frac{2\pi}{3}$ or $x = \frac{4\pi}{3}$.**FINAL ANSWER**

WMA12-01_2026_Jan_Q08

(i) $\theta = 51.9^\circ, 68.1^\circ, 141.9^\circ, 158.1^\circ$. (ii) $x = \frac{2\pi}{3}, \frac{4\pi}{3}$.

PURE MATHEMATICS 2

ADAPTIVE ALL CLASSIFIED

Presentation Edition | Solutions

CLASSIFIED BY IDEA | ROOM TO THINK

Every demand appears clearly, with practical space for the mathematics.

Dr Eslam Ahmed

Elite Mathematics

ElitelGCSE.com | 011 2000 9622

PURE 2

PURE 2 | TOPIC 11

Applications of Differentiation

Your classified route

20

QUESTIONS

175

MARKS

4

FAMILIES

5

VARIANTS

DR ESLAM AHMED

CLASSIFIED BY IDEA

Move from recognition to method, then use the working space to complete the solution.

TOPIC 11 | FAMILY 01

Gradient/Rate

WHAT THIS FAMILY ASKS

Recognise the gradient/rate demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

2 classified questions

19 total marks

T11

11_GRAD

11_GRAD_CORE

CHAPTER
Applications of Differentiation

FAMILY
Gradient/rate

VARIANT
core family move

7 marks

WMA12/01 OCTOBER 2019
Question 10

Past-paper question

WMA12-01_2019_Oct_Q10

10. The curve C has equation

$$y = ax^3 - 3x^2 + 3x + b$$

where a and b are constants.

Given that

- the point $(2, 5)$ lies on C
- the gradient of the curve at $(2, 5)$ is 7

(a) find the value of a and the value of b .

(4)

(b) Prove that C has no turning points.

(3)

T11 Applications of
Differentiation

11_GRAD Gradient/rate

11_GRAD_CORE core family
move

WORKED SOLUTION

Question 10

WMA12/01 October 2019

WMA12-01_2019_Oct_Q10

1

Use the point on the curveSubstitute $(2, 5)$ into $y = ax^3 - 3x^2 + 3x + b$: $5 = 8a - 12 + 6 + b$, so $8a + b = 11$.

2

Differentiate

$$\frac{dy}{dx} = 3ax^2 - 6x + 3.$$

3

Use the given gradientAt $x = 2$, the gradient is 7, so $12a - 12 + 3 = 7$. Hence $12a = 16$ and $a = \frac{4}{3}$.

4

Find bUsing $8a + b = 11$, $b = 11 - 8\left(\frac{4}{3}\right) = \frac{1}{3}$.

5

Check for turning pointsWith $a = \frac{4}{3}$, $\frac{dy}{dx} = 4x^2 - 6x + 3$.

6

Use the discriminantFor $4x^2 - 6x + 3 = 0$, the discriminant is $(-6)^2 - 4(4)(3) = 36 - 48 = -12 < 0$.

7

Conclude

The derivative is never zero, so the curve has no turning points.

FINAL ANSWER(a) $a = \frac{4}{3}$, $b = \frac{1}{3}$. (b) **Proven: C has no turning points.**

WMA12-01_2019_Oct_Q10

T11

11_GRAD

11_GRAD_CORE

CHAPTER
Applications of Differentiation

FAMILY
Gradient/rate

VARIANT
core family move

12 marks

WMA12/01 OCTOBER 2023

Question 9

Past-paper question

WMA12-01_2023_Oct_Q09

9.




Figure 3

In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

Figure 3 shows a sketch of part of the curve C with equation

$$y = \frac{2}{3}x^2 - 9\sqrt{x} + 13 \quad x \geq 0$$

- (a) Find, using calculus, the range of values of x for which y is increasing.

(4)

The point P lies on C and has coordinates $(9, 40)$.

The line l is the tangent to C at the point P .

The finite region R , shown shaded in Figure 3, is bounded by the curve C , the line l , the x -axis and the y -axis.

- (b) Find, using calculus, the exact area of R .

(8)

T11 Applications of
Differentiation

11_GRAD Gradient/rate

11_GRAD_CORE core family
move

WORKED SOLUTION

Question 9

WMA12/01 October 2023

WMA12-01_2023_Oct_Q09

1

DifferentiateFor $y = \frac{2}{3}x^2 - 9\sqrt{x} + 13$, $\frac{dy}{dx} = \frac{4}{3}x - \frac{9}{2\sqrt{x}}$.

2

Set up increasing condition y is increasing when $\frac{dy}{dx} > 0$.

3

Multiply by a positive quantityFor $x > 0$, multiply by $6\sqrt{x}$: $8x\sqrt{x} - 27 > 0$.

4

Solve the inequality $x^{3/2} > \frac{27}{8} = \left(\frac{3}{2}\right)^3$, so $\sqrt{x} > \frac{3}{2}$ and $x > \frac{9}{4}$.

5

Find the tangent gradientAt $P(9, 40)$, $\frac{dy}{dx} = \frac{4}{3}(9) - \frac{9}{2 \cdot 3} = 12 - \frac{3}{2} = \frac{21}{2}$.

6

Write the tangent line $y - 40 = \frac{21}{2}(x - 9)$, so $y = \frac{21}{2}x - \frac{109}{2}$.

7

Find where the tangent meets the x-axisSet $y = 0$: $\frac{21}{2}x - \frac{109}{2} = 0$, so $x = \frac{109}{21}$.

8

Split the areaFrom $x = 0$ to $x = \frac{109}{21}$, the lower boundary is the x -axis. From $x = \frac{109}{21}$ to $x = 9$, the lower boundary is the tangent.

9

Set up the integral

$$R = \int_0^{109/21} \left(\frac{2}{3}x^2 - 9\sqrt{x} + 13 \right) dx + \int_{109/21}^9 \left(\frac{2}{3}x^2 - 9\sqrt{x} + 13 - \left(\frac{21}{2}x - \frac{109}{2} \right) \right) dx.$$

10

Use antiderivatives

$$\int \left(\frac{2}{3}x^2 - 9\sqrt{x} + 13 \right) dx = \frac{2}{9}x^3 - 6x^{3/2} + 13x, \text{ and } \int \left(\frac{21}{2}x - \frac{109}{2} \right) dx = \frac{21}{4}x^2 - \frac{109}{2}x.$$

11

Evaluate

Substituting the limits and simplifying gives $R = \frac{857}{21}$.

FINAL ANSWER

WMA12-01_2023_Oct_Q09

(a) y is increasing for $x > \frac{9}{4}$. (b) **Area** = $\frac{857}{21}$.

DR ESLAM AHMED

TOPIC 11 | FAMILY 02

Optimisation

WHAT THIS FAMILY ASKS

Recognise the optimisation demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 Use constraints to remove extra variables first.

V02 State the maximum/minimum value in context, not just x .

4 classified questions

40 total marks

T11

11_OPTIM

11_OPTIM_HV01

CHAPTER

Applications of Differentiation

FAMILY

Optimisation

VARIANT

Use constraints to remove
extra variables first.

10 marks

WMA12/01 MAY/JUNE 2021

Question 9

Past-paper question

WMA12-01_2021_MayJune_Q09

9.




Figure 3

Figure 3 shows a sketch of a square based, open top box.

The height of the box is h cm, and the base edges each have length l cm.

Given that the volume of the box is $250\,000\text{ cm}^3$

(a) show that the external surface area, $S\text{ cm}^2$, of the box is given by

$$S = \frac{250\,000}{h} + 2000\sqrt{h} \quad (3)$$

(b) Use algebraic differentiation to show that S has a stationary point when $h = 250^k$ where k is a rational constant to be found. (5)

(c) Justify by further differentiation that this value of h gives the minimum external surface area of the box. (2)

(Total 10 marks)

**T11 Applications of
Differentiation**

11_OPTIM Optimisation

**11_OPTIM_HV01 Use
constraints to remove extra
variables first.**

WORKED SOLUTION

Question 9

WMA12/01 May/June 2021

WMA12-01_2021_MayJune_Q09

1

Use the volume

The volume is $l^2h = 250000$, so $l^2 = \frac{250000}{h}$.

2

Write the surface area

The open box has one square base and four side faces, so $S = l^2 + 4lh$.

3

Substitute l

Since $l = \sqrt{\frac{250000}{h}} = \frac{500}{\sqrt{h}}$, $4lh = 4 \cdot \frac{500}{\sqrt{h}} \cdot h = 2000\sqrt{h}$.

4

Show S

Therefore $S = \frac{250000}{h} + 2000\sqrt{h}$.

5

Differentiate S

$S = 250000h^{-1} + 2000h^{1/2}$, so $\frac{dS}{dh} = -250000h^{-2} + 1000h^{-1/2}$.

6

Set the derivative to zero

$-250000h^{-2} + 1000h^{-1/2} = 0$, so $1000h^{-1/2} = 250000h^{-2}$.

7

Solve for h

Multiplying by h^2 gives $1000h^{3/2} = 250000$, so $h^{3/2} = 250$ and $h = 250^{2/3}$.

8

Find k

Since $h = 250^k$, we have $k = \frac{2}{3}$.

9

Use the second derivative

$\frac{d^2S}{dh^2} = 500000h^{-3} - 500h^{-3/2}$.

10

Check the sign

At $h = 250^{2/3}$, $h^{3/2} = 250$, so the second derivative is $\frac{500000}{250^2} - \frac{500}{250} = 8 - 2 = 6 > 0$.

11

Conclude

The stationary point is therefore a minimum.

FINAL ANSWER

WMA12-01_2021_MayJune_Q09

(a) **Proven:** $S = \frac{250000}{h} + 2000\sqrt{h}$. (b) $h = 250^{2/3}$, so $k = \frac{2}{3}$. (c) **Minimum.**

DR ESLAM AHMED

T11

11_OPTIM

11_OPTIM_HV01

CHAPTER
Applications of Differentiation

FAMILY
Optimisation

VARIANT
Use constraints to remove
extra variables first.

10 marks

WMA12/01 JANUARY 2025
Question 3

Past-paper question

WMA12-01_2025_Jan_Q03

3.




Figure 1

Figure 1 shows an open-topped container used for holding water.

The container is in the shape of a cuboid and is made of sheet metal.

The base of the container is a rectangle $3x$ metres by x metres.

The height of the container is y metres as shown in Figure 1.

Given that the capacity of the container is 120 m^3

(a) show that the area $A \text{ m}^2$ of the sheet metal used to make the container is given by

$$A = Px^2 + \frac{Q}{x}$$

where P and Q are positive constants to be found.

(4)

(b) Use calculus to find the value of x for which A has a stationary value, giving your answer to 3 significant figures.

(4)

(c) Find $\frac{d^2A}{dx^2}$ and hence show that the value of x found in part (b) gives the minimum value of A .

(2)

T11 Applications of
Differentiation

11_OPTIM Optimisation

11_OPTIM_HV01 Use
constraints to remove extra
variables first.

WORKED SOLUTION

Question 3

WMA12/01 January 2025

WMA12-01_2025_Jan_Q03

1

Use the volume

The volume is $(3x)(x)(y) = 3x^2y$. Since the capacity is 120, $3x^2y = 120$, so $y = \frac{40}{x^2}$.

2

Write the metal area

The open top means the metal area is the base plus four sides: $A = 3x^2 + 2(3xy) + 2(xy)$.

3

Substitute for y

$A = 3x^2 + 8xy = 3x^2 + 8x\left(\frac{40}{x^2}\right) = 3x^2 + \frac{320}{x}$. Hence $P = 3$ and $Q = 320$.

4

Differentiate

$$\frac{dA}{dx} = 6x - 320x^{-2} = 6x - \frac{320}{x^2}.$$

5

Use the stationary condition

Set $\frac{dA}{dx} = 0$: $6x = \frac{320}{x^2}$, so $6x^3 = 320$.

6

Find x

$x^3 = \frac{160}{3}$, so $x = \sqrt[3]{\frac{160}{3}} = 3.764\dots$. To 3 significant figures, $x = 3.76$.

7

Differentiate again

$$\frac{d^2A}{dx^2} = 6 + 640x^{-3} = 6 + \frac{640}{x^3}.$$

8

Show the minimum

For $x > 0$, $6 + \frac{640}{x^3} > 0$, so the stationary value found in part (b) is a minimum.

FINAL ANSWER

(a) $A = 3x^2 + \frac{320}{x}$, so $P = 3$ and $Q = 320$. (b) $x = 3.76$. (c) $\frac{d^2A}{dx^2} = 6 + \frac{640}{x^3}$, minimum.

WMA12-01_2025_Jan_Q03

T11

11_OPTIM

11_OPTIM_HV03

CHAPTER
Applications of Differentiation

FAMILY
Optimisation

VARIANT
State the maximum/minimum
value in context, not just $/(x/)$.

9 marks

WMA12/01 JANUARY 2023
Question 2

Past-paper question

WMA12-01_2023_Jan_Q02

2. In this question you must show all stages of your working.

Solutions based entirely on calculator technology are not acceptable.




Figure 2

A brick is in the shape of a cuboid with width x cm, length $3x$ cm and height h cm, as shown in Figure 2.

The volume of the brick is 972 cm^3

(a) Show that the surface area of the brick, $S \text{ cm}^2$, is given by

$$S = 6x^2 + \frac{2592}{x}$$

(3)

(b) Find $\frac{dS}{dx}$

(1)

(c) Hence find the value of x for which S is stationary.

(2)

(d) Find $\frac{d^2S}{dx^2}$ and hence show that the value of x found in part (c) gives the minimum value of S .

(2)

(e) Hence find the minimum surface area of the brick.

(1)

T11 Applications of
Differentiation

11_OPTIM Optimisation

11_OPTIM_HV03 State the
maximum/minimum value in
context, not just $f(x)$.

WORKED SOLUTION

Question 2

WMA12/01 January 2023

WMA12-01_2023_Jan_Q02

1

Use the volumeThe cuboid has dimensions x , $3x$ and h , so its volume is $3x^2h = 972$.

2

Write h in terms of x Hence $h = \frac{972}{3x^2} = \frac{324}{x^2}$.

3

Build the surface areaThe surface area is $S = 2(3x^2 + 3xh + xh) = 6x^2 + 8xh$.

4

Substitute h $S = 6x^2 + 8x \left(\frac{324}{x^2} \right) = 6x^2 + \frac{2592}{x}$.

5

Differentiate $S = 6x^2 + 2592x^{-1}$, so $\frac{dS}{dx} = 12x - 2592x^{-2} = 12x - \frac{2592}{x^2}$.

6

Set the derivative to zeroFor a stationary value, $12x - \frac{2592}{x^2} = 0$.

7

Solve for x Multiplying by x^2 gives $12x^3 = 2592$, so $x^3 = 216$ and $x = 6$.

8

Use the second derivative $\frac{d^2S}{dx^2} = 12 + \frac{5184}{x^3}$.

9

Show it is a minimumAt $x = 6$, $\frac{d^2S}{dx^2} = 12 + \frac{5184}{216} = 36 > 0$, so the stationary value is a minimum.

10

Find the minimum surface area $S = 6(6)^2 + \frac{2592}{6} = 216 + 432 = 648$, so the minimum surface area is 648 cm^2 .

FINAL ANSWER

WMA12-01_2023_Jan_Q02

- (a) $S = 6x^2 + \frac{2592}{x}$. (b) $\frac{dS}{dx} = 12x - \frac{2592}{x^2}$. (c) $x = 6$. (d) **Minimum shown.**
(e) **648 cm².**

DR ESLAM AHMED

T11

11_OPTIM

11_OPTIM_HV03

CHAPTER
Applications of Differentiation

FAMILY
Optimisation

VARIANT
State the maximum/minimum
value in context, not just $/(x/)$.

11 marks

WMA12/01 MAY/JUNE 2025
Question 9

Past-paper question

WMA12-01_2025_MayJune_Q09

9:




Figure 2

Figure 2 shows a sketch of an open container.

The sides $CDEF$ and $ABFE$ are rectangles.

The ends ADE and BCF are congruent (identical) right-angled triangles.

The container is made from metal of negligible thickness.

Given that

- $AE = BF = 3x$ metres
- $DE = CF = 2x$ metres
- $AB = DC = EF = L$ metres

and the capacity of the container is 12 m^3

(a) show that the area of metal used to make the container, $S \text{ m}^2$, is given by

$$S = Px^2 + \frac{Q}{x}$$

where P and Q are positive integers to be found.

(4)

Given that x can vary,

(b) use algebraic calculus to find the minimum value of S , giving your answer to one decimal place.

(5)

(c) Justify that the value of S found in part (b) is a minimum.

(2)

T11 Applications of
Differentiation

11_OPTIM Optimisation

11_OPTIM_HV03 State the
maximum/minimum value in
context, not just $f(x)$.

WORKED SOLUTION

Question 9

WMA12/01 May/June 2025

WMA12-01_2025_MayJune_Q09

1

Find the triangular end area

Each end is a right-angled triangle with perpendicular sides $3x$ and $2x$, so its area is $\frac{1}{2}(3x)(2x) = 3x^2$.

2

Use the capacity

The volume is cross-section area times length, so $3x^2L = 12$ and $L = \frac{4}{x^2}$.

3

Write the metal area

There are two triangular ends with total area $6x^2$, and two rectangular sides with total area $3xL + 2xL = 5xL$.

4

Substitute for L

$S = 6x^2 + 5x\left(\frac{4}{x^2}\right) = 6x^2 + \frac{20}{x}$. Hence $P = 6$ and $Q = 20$.

5

Differentiate

$$\frac{dS}{dx} = 12x - 20x^{-2} = 12x - \frac{20}{x^2}.$$

6

Find the stationary value of x

Set $\frac{dS}{dx} = 0$: $12x = \frac{20}{x^2}$, so $x^3 = \frac{5}{3}$.

7

Evaluate S

At $x = \sqrt[3]{\frac{5}{3}}$, $S = 6x^2 + \frac{20}{x} = 25.3029\dots$, so the minimum value is 25.3 to one decimal place.

8

Differentiate again

$$\frac{d^2S}{dx^2} = 12 + 40x^{-3} = 12 + \frac{40}{x^3}.$$

9

Justify the minimum

Since $x > 0$, $12 + \frac{40}{x^3} > 0$, so the stationary value found is a minimum.

FINAL ANSWER

WMA12-01_2025_MayJune_Q09

(a) $S = 6x^2 + \frac{20}{x}$, so $P = 6$ and $Q = 20$. (b) 25.3. (c) **Minimum justified.**

DR ESLAM AHMED

TOPIC 11 | FAMILY 03

Curve Behaviour

WHAT THIS FAMILY ASKS

Recognise the curve behaviour demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

3 classified questions

25 total marks

T11

11_SKETCH

11_SKETCH_CORE

CHAPTER

Applications of Differentiation

FAMILY

Curve behaviour

VARIANT

core family move

6 marks

WMA12/01 OCTOBER 2019

Question 1

Past-paper question

WMA12-01_2019_Oct_Q01

1. A curve C has equation $y = 2x^2(x - 5)$

(a) Find, using calculus, the x coordinates of the stationary points of C .

(4)

(b) Hence find the values of x for which y is increasing.

(2)

DR ESLAM AHMED

**T11 Applications of
Differentiation****11_SKETCH Curve behaviour****11_SKETCH_CORE core
family move****WORKED SOLUTION****Question 1**

WMA12/01 October 2019

WMA12-01_2019_Oct_Q01

1**Expand the curve**

$$y = 2x^2(x - 5) = 2x^3 - 10x^2.$$

2**Differentiate**

$$\frac{dy}{dx} = 6x^2 - 20x = 2x(3x - 10).$$

3**Find the stationary values of x**

At a stationary point, $\frac{dy}{dx} = 0$. Hence $2x(3x - 10) = 0$, so $x = 0$ or $x = \frac{10}{3}$.

4**Use the sign of the derivative**

Since $2x(3x - 10)$ is positive outside its two roots and negative between them, y is increasing for $x < 0$ and for $x > \frac{10}{3}$.

FINAL ANSWER

WMA12-01_2019_Oct_Q01

(a) $x = 0, \frac{10}{3}$. (b) $x < 0$ or $x > \frac{10}{3}$.

T11

11_SKETCH

11_SKETCH_CORE

CHAPTER

Applications of Differentiation

FAMILY

Curve behaviour

VARIANT

core family move

10 marks

WMA12/01 OCTOBER 2021

Question 8

Past-paper question

WMA12-01_2021_Oct_Q08

8. In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.




Figure 2

Figure 2 shows a sketch of part of the curve C with equation

$$y = \frac{4}{3}x^3 - 11x^2 + kx \quad \text{where } k \text{ is a constant}$$

The point M is the maximum turning point of C and is shown in Figure 2.

Given that the x coordinate of M is 2

- (a) show that $k = 28$ (3)

- (b) Determine the range of values of x for which y is increasing. (2)

The line l passes through M and is parallel to the x -axis.

The region R , shown shaded in Figure 2, is bounded by the curve C , the line l and the y -axis.

- (c) Find, by algebraic integration, the exact area of R . (5)

T11 Applications of
Differentiation

11_SKETCH Curve behaviour

11_SKETCH_CORE core
family move

WORKED SOLUTION

Question 8

WMA12/01 October 2021

WMA12-01_2021_Oct_Q08

1

Differentiate the curveFor $y = \frac{4}{3}x^3 - 11x^2 + kx$, $\frac{dy}{dx} = 4x^2 - 22x + k$.

2

Use the turning pointAt the turning point M , $x = 2$ and $\frac{dy}{dx} = 0$.

3

Show k $4(2)^2 - 22(2) + k = 0$, so $16 - 44 + k = 0$ and $k = 28$.

4

Factor the derivativeWith $k = 28$, $\frac{dy}{dx} = 4x^2 - 22x + 28 = 2(2x - 7)(x - 2)$.

5

Find where y is increasingThe derivative is positive outside its roots, so y is increasing for $x < 2$ or $x > \frac{7}{2}$.

6

Find the height of the lineAt $x = 2$, $y = \frac{4}{3}(8) - 11(4) + 28(2) = \frac{68}{3}$. Hence line l has equation $y = \frac{68}{3}$.

7

Set up the shaded areaThe region is between the line and the curve from $x = 0$ to $x = 2$, so $A = \int_0^2 \left(\frac{68}{3} - \left(\frac{4}{3}x^3 - 11x^2 + 28x \right) \right) dx$.

8

Integrate

$$A = \left[\frac{68}{3}x - \frac{1}{3}x^4 + \frac{11}{3}x^3 - 14x^2 \right]_0^2.$$

9

Evaluate exactlySubstituting $x = 2$ gives $\frac{136}{3} - \frac{16}{3} + \frac{88}{3} - 56 = \frac{40}{3}$.**FINAL ANSWER**

WMA12-01_2021_Oct_Q08

(a) **Proven:** $k = 28$. (b) $x < 2$ or $x > \frac{7}{2}$. (c) $\frac{40}{3}$.

T11

11_SKETCH

11_SKETCH_CORE

CHAPTER

Applications of Differentiation

FAMILY

Curve behaviour

VARIANT

core family move

9 marks

WMA12/01 MAY/JUNE (R) 2024

Question 6

Past-paper question

WMA12-01_2024_MayJuneR_Q06

6.

$$f(x) = 4x^3 + px^2 + 8x + q$$

where p and q are constants.

Given that

- $(2x + 3)$ is a factor of $f(x)$
- $f(x)$ has a remainder of -5 when divided by $(x + 2)$

(a) (i) show that $p = 10$

(ii) find the value of q .

(5)

(b) Hence find the range of values of x for which $f(x)$ is decreasing.

(Solutions based entirely on calculator technology are not acceptable.)

(4)

T11 Applications of
Differentiation

11_SKETCH Curve behaviour

11_SKETCH_CORE core
family move

WORKED SOLUTION

Question 6

WMA12/01 May/June (R) 2024

WMA12-01_2024_MayJuneR_Q06

1

Use the factor theoremSince $(2x + 3)$ is a factor, $x = -\frac{3}{2}$ is a root.

2

Substitute x equals negative three halves

$$4\left(-\frac{3}{2}\right)^3 + p\left(-\frac{3}{2}\right)^2 + 8\left(-\frac{3}{2}\right) + q = 0.$$

3

Simplify the first condition

$$-\frac{27}{2} + \frac{9}{4}p - 12 + q = 0, \text{ so } \frac{9}{4}p + q = \frac{51}{2}.$$

4

Use the remainder theoremThe remainder on division by $(x + 2)$ is $f(-2)$.

5

Substitute x equals negative two

$$f(-2) = 4(-8) + 4p - 16 + q = 4p + q - 48.$$

6

Use the given remainder

$$4p + q - 48 = -5, \text{ so } 4p + q = 43.$$

7

Solve for p and q

Subtract $\frac{9}{4}p + q = \frac{51}{2}$ from $4p + q = 43$: $\frac{7}{4}p = \frac{35}{2}$, so $p = 10$. Then $4(10) + q = 43$, so $q = 3$.

8

Differentiate

With $p = 10$, $f(x) = 4x^3 + 10x^2 + 8x + 3$, so $f'(x) = 12x^2 + 20x + 8$.

9

Factor the derivative

$$f'(x) = 4(3x + 2)(x + 1).$$

10

Find where decreasing

The quadratic derivative is negative between its roots, so $f(x)$ is decreasing for $-1 < x < -\frac{2}{3}$.

FINAL ANSWER

WMA12-01_2024_MayJuneR_Q06

$$(a)(i) p = 10. \quad (a)(ii) q = 3. \quad (b) -1 < x < -\frac{2}{3}.$$

DR ESLAM AHMED

TOPIC 11 | FAMILY 04

Stationary Points

WHAT THIS FAMILY ASKS

Recognise the stationary points demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

11 classified questions

91 total marks

T11

11_STAT

11_STAT_CORE

CHAPTER

Applications of Differentiation

FAMILY

Stationary points

VARIANT

core family move

10 marks

WMA12/01 JANUARY 2020

Question 10

Past-paper question

WMA12-01_2020_Jan_Q10

10. A curve C has equation

$$y = 4x^3 - 9x + \frac{k}{x} \quad x > 0$$

where k is a constant.

The point P with x coordinate $\frac{1}{2}$ lies on C .

Given that P is a stationary point of C ,

(a) show that $k = -\frac{3}{2}$ (4)

(b) Determine the nature of the stationary point at P , justifying your answer. (2)

The curve C has a second stationary point.

(c) Using algebra, find the x coordinate of this second stationary point. (4)

(Total 10 marks)

T11 Applications of
Differentiation

11_STAT Stationary points

11_STAT_CORE core family
move

WORKED SOLUTION

Question 10

WMA12/01 January 2020

WMA12-01_2020_Jan_Q10

1

Differentiate the curveWrite $y = 4x^3 - 9x + kx^{-1}$. Then $\frac{dy}{dx} = 12x^2 - 9 - kx^{-2}$.

2

Use the stationary pointAt $x = \frac{1}{2}$, $\frac{dy}{dx} = 0$, so $0 = 12\left(\frac{1}{2}\right)^2 - 9 - k\left(\frac{1}{2}\right)^{-2} = 3 - 9 - 4k$.

3

Find k $-6 - 4k = 0$, hence $k = -\frac{3}{2}$.

4

Use the second derivativeWith $k = -\frac{3}{2}$, $\frac{dy}{dx} = 12x^2 - 9 + \frac{3}{2x^2}$ and $\frac{d^2y}{dx^2} = 24x - 3x^{-3}$.

5

Determine the nature of PAt $x = \frac{1}{2}$, $\frac{d^2y}{dx^2} = 12 - 24 = -12 < 0$, so P is a local maximum.

6

Find the other stationary pointSet $12x^2 - 9 + \frac{3}{2x^2} = 0$. Multiplying by $2x^2$ gives $24x^4 - 18x^2 + 3 = 0$, or $8x^4 - 6x^2 + 1 = 0$.

7

Solve the quadratic in x squared $(4x^2 - 1)(2x^2 - 1) = 0$. The given point has $x = \frac{1}{2}$, so the second stationary point has $x^2 = \frac{1}{2}$.

8

State the positive x coordinateSince $x > 0$, the second stationary point has $x = \frac{1}{\sqrt{2}}$.**FINAL ANSWER**

WMA12-01_2020_Jan_Q10

(a) $k = -\frac{3}{2}$. (b) local maximum. (c) $x = \frac{1}{\sqrt{2}}$.

T11

11_STAT

11_STAT_CORE

CHAPTER

Applications of Differentiation

FAMILY

Stationary points

VARIANT

core family move

7 marks

WMA12/01 JANUARY 2021

Question 2

Past-paper question

WMA12-01_2021_Jan_Q02

2. A curve has equation

$$y = x^3 - x^2 - 16x + 2$$

- (a) Using calculus, find the x coordinates of the stationary points of the curve. (4)
- (b) Justify, by further calculus, the nature of all of the stationary points of the curve. (3)

T11 Applications of
Differentiation

11_STAT Stationary points

11_STAT_CORE core family
move

WORKED SOLUTION

Question 2

WMA12/01 January 2021

WMA12-01_2021_Jan_Q02

1

Differentiate the curveFor $y = x^3 - x^2 - 16x + 2$, $\frac{dy}{dx} = 3x^2 - 2x - 16$.

2

Set the gradient to zeroStationary points occur when $3x^2 - 2x - 16 = 0$.

3

Factorise $3x^2 - 2x - 16 = (3x - 8)(x + 2)$, so $x = \frac{8}{3}$ or $x = -2$.

4

Differentiate again $\frac{d^2y}{dx^2} = 6x - 2$.

5

Classify x equals negative twoAt $x = -2$, $6x - 2 = -14 < 0$, so the stationary point is a local maximum.

6

Classify x equals eight over threeAt $x = \frac{8}{3}$, $6x - 2 = 14 > 0$, so the stationary point is a local minimum.**FINAL ANSWER**

WMA12-01_2021_Jan_Q02

(a) $x = -2$, $\frac{8}{3}$. (b) local maximum at $x = -2$ and local minimum at $x = \frac{8}{3}$.

T11

11_STAT

11_STAT_CORE

CHAPTER

Applications of Differentiation

FAMILY

Stationary points

VARIANT

core family move

8 marks

WMA12/01 JANUARY 2022

Question 2

Past-paper question

WMA12-01_2022_Jan_Q02

2. In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

The curve C has equation

$$y = 27x^{\frac{1}{2}} - x^{\frac{3}{2}} - 20 \quad x > 0$$

(a) Find $\frac{dy}{dx}$, giving each term in simplest form.

(2)

(b) Hence find the coordinates of the stationary point of C .

(4)

(c) Find $\frac{d^2y}{dx^2}$ and hence determine the nature of the stationary point of C .

(2)

T11 Applications of
Differentiation

11_STAT Stationary points

11_STAT_CORE core family
move

WORKED SOLUTION

Question 2

WMA12/01 January 2022

WMA12-01_2022_Jan_Q02

1

DifferentiateFor $y = 27x^{1/2} - x^{3/2} - 20$, $\frac{dy}{dx} = \frac{27}{2}x^{-1/2} - \frac{3}{2}x^{1/2}$.

2

Set the gradient to zero

$$\frac{27}{2}x^{-1/2} - \frac{3}{2}x^{1/2} = 0.$$

3

Clear the powersMultiplying by $2x^{1/2}$ gives $27 - 3x = 0$, so $x = 9$.

4

Find yAt $x = 9$, $y = 27\sqrt{9} - 9^{3/2} - 20 = 81 - 27 - 20 = 34$.

5

State the stationary pointThe stationary point is $(9, 34)$.

6

Differentiate again

$$\frac{d^2y}{dx^2} = -\frac{27}{4}x^{-3/2} - \frac{3}{4}x^{-1/2}.$$

7

Determine the natureAt $x = 9$, both terms in $\frac{d^2y}{dx^2}$ are negative, so the stationary point is a local maximum.**FINAL ANSWER**

WMA12-01_2022_Jan_Q02

(a) $\frac{dy}{dx} = \frac{27}{2}x^{-1/2} - \frac{3}{2}x^{1/2}$. (b) $(9, 34)$. (c) **local maximum.**

T11

11_STAT

11_STAT_CORE

CHAPTER

Applications of Differentiation

FAMILY

Stationary points

VARIANT

core family move

8 marks

WMA12/01 MAY/JUNE 2022

Question 8

Past-paper question

WMA12-01_2022_MayJune_Q08

8. In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

A curve has equation

$$y = 256x^4 - 304x - 35 + \frac{27}{x^2} \quad x \neq 0$$

- (a) Find $\frac{dy}{dx}$ (3)
- (b) Hence find the coordinates of the stationary points of the curve. (5)

T11 Applications of
Differentiation

11_STAT Stationary points

11_STAT_CORE core family
move

WORKED SOLUTION

Question 8

WMA12/01 May/June 2022

WMA12-01_2022_MayJune_Q08

1

Rewrite with powers

$$y = 256x^4 - 304x - 35 + 27x^{-2}.$$

2

Differentiate

$$\frac{dy}{dx} = 1024x^3 - 304 - 54x^{-3} = 1024x^3 - 304 - \frac{54}{x^3}.$$

3

Set the derivative to zero

$$1024x^3 - 304 - \frac{54}{x^3} = 0.$$

4

Use a substitution

Multiplying by x^3 gives $1024x^6 - 304x^3 - 54 = 0$. Let $u = x^3$.

5

Solve for u

$$1024u^2 - 304u - 54 = 0, \text{ or } 512u^2 - 152u - 27 = 0 = (8u + 1)(64u - 27).$$

6

Find x

Thus $x^3 = -\frac{1}{8}$ or $x^3 = \frac{27}{64}$, giving $x = -\frac{1}{2}$ or $x = \frac{3}{4}$.

7

Find the y values

Substitution into $y = 256x^4 - 304x - 35 + \frac{27}{x^2}$ gives $y = 241$ at $x = -\frac{1}{2}$ and $y = -134$ at $x = \frac{3}{4}$.

8

State the stationary points

The stationary points are $(-\frac{1}{2}, 241)$ and $(\frac{3}{4}, -134)$.

FINAL ANSWER

WMA12-01_2022_MayJune_Q08

$$(a) \frac{dy}{dx} = 1024x^3 - 304 - \frac{54}{x^3}. \quad (b) \left(-\frac{1}{2}, 241\right), \left(\frac{3}{4}, -134\right).$$

T11

11_STAT

11_STAT_CORE

CHAPTER

Applications of Differentiation

FAMILY

Stationary points

VARIANT

core family move

9 marks

WMA12/01 OCTOBER 2022

Question 7

Past-paper question

WMA12-01_2022_Oct_Q07

7. The curve C has equation

$$y = \frac{12x^3(x-7) + 14x(13x-15)}{21\sqrt{x}} \quad x > 0$$

(a) Write the equation of C in the form

$$y = ax^{\frac{7}{2}} + bx^{\frac{5}{2}} + cx^{\frac{3}{2}} + dx^{\frac{1}{2}}$$

where a , b , c and d are fully simplified constants.

(3)

The curve C has three turning points.

Using calculus,

(b) show that the x coordinates of the three turning points satisfy the equation

$$2x^3 - 10x^2 + 13x - 5 = 0$$

(3)

Given that the x coordinate of one of the turning points is 1

(c) find, using algebra, the exact x coordinates of the other two turning points.

(Solutions based entirely on calculator technology are not acceptable.)

(3)

T11 Applications of
Differentiation

11_STAT Stationary points

11_STAT_CORE core family
move

WORKED SOLUTION

Question 7

WMA12/01 October 2022

WMA12-01_2022_Oct_Q07

1

Expand the numerator

$$12x^3(x-7) + 14x(13x-15) = 12x^4 - 84x^3 + 182x^2 - 210x.$$

2

Divide by 21 root x

Since $\sqrt{x} = x^{1/2}$, divide each term by $21x^{1/2}$.

3

Write in the required form

$$y = \frac{12}{21}x^{7/2} - \frac{84}{21}x^{5/2} + \frac{182}{21}x^{3/2} - \frac{210}{21}x^{1/2} = \frac{4}{7}x^{7/2} - 4x^{5/2} + \frac{26}{3}x^{3/2} - 10x^{1/2}.$$

4

Differentiate

$$\frac{dy}{dx} = 2x^{5/2} - 10x^{3/2} + 13x^{1/2} - 5x^{-1/2}.$$

5

Factor the derivative

$$\frac{dy}{dx} = x^{-1/2}(2x^3 - 10x^2 + 13x - 5).$$

6

Use the turning point condition

At a turning point, $\frac{dy}{dx} = 0$. Since $x > 0$, $x^{-1/2} \neq 0$, so the x coordinates satisfy $2x^3 - 10x^2 + 13x - 5 = 0$.

7

Use the known root

One root is $x = 1$, so divide $2x^3 - 10x^2 + 13x - 5$ by $(x - 1)$.

8

Factor the cubic

$$2x^3 - 10x^2 + 13x - 5 = (x - 1)(2x^2 - 8x + 5).$$

9

Solve the quadratic

$$2x^2 - 8x + 5 = 0 \text{ gives } x = \frac{8 \pm \sqrt{64 - 40}}{4} = \frac{8 \pm 2\sqrt{6}}{4} = 2 \pm \frac{\sqrt{6}}{2}.$$

FINAL ANSWER

WMA12-01_2022_Oct_Q07

$$(a) y = \frac{4}{7}x^{7/2} - 4x^{5/2} + \frac{26}{3}x^{3/2} - 10x^{1/2}. \quad (b) \text{ Proven.} \quad (c) x = 2 - \frac{\sqrt{6}}{2} \text{ and } x = 2 + \frac{\sqrt{6}}{2}.$$

T11

11_STAT

11_STAT_CORE

CHAPTER

Applications of Differentiation

FAMILY

Stationary points

VARIANT

core family move

8 marks

WMA12/01 MAY/JUNE 2023

Question 7

Past-paper question

WMA12-01_2023_MayJune_Q07

7. The height of a river above a fixed point on the riverbed was monitored over a 7-day period.

The height of the river, H metres, t days after monitoring began, was given by

$$H = \frac{\sqrt{t}}{20}(20 + 6t - t^2) + 17 \quad 0 \leq t \leq 7$$

Given that H has a stationary value at $t = \alpha$

- (a) use calculus to show that α satisfies the equation

$$5\alpha^2 - 18\alpha - 20 = 0 \quad (5)$$

- (b) Hence find the value of α , giving your answer to 3 decimal places. (1)

- (c) Use further calculus to prove that H is a maximum at this value of α . (2)

T11 Applications of
Differentiation

11_STAT Stationary points

11_STAT_CORE core family
move

WORKED SOLUTION

Question 7

WMA12/01 May/June 2023

WMA12-01_2023_MayJune_Q07

1

Rewrite H with powers

$$H = \frac{1}{20}t^{1/2}(20 + 6t - t^2) + 17.$$

2

Expand before differentiating

$$H = \frac{1}{20}(20t^{1/2} + 6t^{3/2} - t^{5/2}) + 17.$$

3

Differentiate

$$\frac{dH}{dt} = \frac{1}{20} \left(10t^{-1/2} + 9t^{1/2} - \frac{5}{2}t^{3/2} \right).$$

4

Put over a common factor

$$\frac{dH}{dt} = \frac{-5t^2 + 18t + 20}{40\sqrt{t}}.$$

5

Use the stationary condition

At $t = \alpha$, $\frac{dH}{dt} = 0$. Since $\alpha > 0$, the denominator is not zero, so $-5\alpha^2 + 18\alpha + 20 = 0$.

6

Show the required equation

Multiplying by -1 gives $5\alpha^2 - 18\alpha - 20 = 0$.

7

Find alpha

$$\alpha = \frac{18 \pm \sqrt{18^2 + 4(5)(20)}}{10} = \frac{9 \pm \sqrt{181}}{5}.$$

8

Use the time interval

Only the positive value lies in $0 \leq t \leq 7$, so $\alpha = \frac{9 + \sqrt{181}}{5} = 4.491$ to 3 decimal places.

9

Differentiate again

$$\frac{d^2H}{dt^2} = \frac{-15t^2 + 18t - 20}{80t^{3/2}}.$$

10

Use the stationary equation

At $t = \alpha$, $5\alpha^2 = 18\alpha + 20$. Therefore the numerator of $\frac{d^2H}{dt^2}$ is $-15\alpha^2 + 18\alpha - 20 = -3(18\alpha + 20) + 18\alpha - 20 = -36\alpha - 80$.

11

Show it is negative

Since $\alpha > 0$, $-36\alpha - 80 < 0$, and the denominator $80\alpha^{3/2}$ is positive. Thus $\frac{d^2H}{dt^2} < 0$, so H is a maximum.

FINAL ANSWER

WMA12-01_2023_MayJune_Q07

(a) **Proven.** (b) $\alpha = 4.491$. (c) H is a maximum at this value.

DR ESLAM AHMED

T11

11_STAT

11_STAT_CORE

CHAPTER

Applications of Differentiation

FAMILY

Stationary points

VARIANT

core family move

9 marks

WMA12/01 JANUARY 2024

Question 10

Past-paper question

WMA12-01_2024_Jan_Q10

10.

In this question you must show detailed reasoning.

Solutions relying entirely on calculator technology are not acceptable.




Figure 2

Figure 2 shows a sketch of the curve with equation

$$y = \frac{1}{2}x^2 + \frac{1458}{\sqrt{x^3}} - 74 \quad x > 0$$

The point P is the only stationary point on the curve.(a) Use calculus to show that the x coordinate of P is 9

(4)

The line l passes through the point P and is parallel to the x -axis.The region R , shown shaded in Figure 2, is bounded by the curve, the line l and the line with equation $x = 4$ (b) Use algebraic integration to find the exact area of R .

(5)

T11 Applications of
Differentiation

11_STAT Stationary points

11_STAT_CORE core family
move

WORKED SOLUTION

Question 10

WMA12/01 January 2024

WMA12-01_2024_Jan_Q10

1

Rewrite the curve

$$y = \frac{1}{2}x^2 + 1458x^{-3/2} - 74.$$

2

Differentiate

$$\frac{dy}{dx} = x + 1458 \left(-\frac{3}{2}\right) x^{-5/2} = x - 2187x^{-5/2}.$$

3

Use the stationary condition

$$\text{At a stationary point, } x - 2187x^{-5/2} = 0.$$

4

Solve for x

$$x = 2187x^{-5/2}, \text{ so } x^{7/2} = 2187 = 3^7.$$

5

Show x coordinate

Raising both sides to the power $\frac{2}{7}$ gives $x = 3^2 = 9$. Since $x > 0$, this is the only stationary point.

6

Find the horizontal line

$$\text{At } x = 9, y = \frac{1}{2}(9)^2 + \frac{1458}{\sqrt{9^3}} - 74 = \frac{81}{2} + 54 - 74 = \frac{41}{2}.$$

7

Set up the area

The region lies from $x = 4$ to $x = 9$ between the curve and the horizontal line $y = \frac{41}{2}$.

8

Integrate the difference

$$R = \int_4^9 \left(\frac{1}{2}x^2 + 1458x^{-3/2} - 74 - \frac{41}{2} \right) dx.$$

9

Find an antiderivative

$$\int \left(\frac{1}{2}x^2 + 1458x^{-3/2} - \frac{189}{2} \right) dx = \frac{x^3}{6} - 2916x^{-1/2} - \frac{189}{2}x.$$

10

Evaluate exactly

$$R = \left[\frac{x^3}{6} - 2916x^{-1/2} - \frac{189}{2}x \right]_4^9 = \frac{373}{3}.$$

FINAL ANSWER

WMA12-01_2024_Jan_Q10

(a) **Proven.** (b) $\frac{373}{3}$.

DR ESLAM AHMED

T11

11_STAT

11_STAT_CORE

CHAPTER

Applications of Differentiation

FAMILY

Stationary points

VARIANT

core family move

8 marks

WMA12/01 MAY/JUNE 2024

Question 9

Past-paper question

WMA12-01_2024_MayJune_Q09

9.




Figure 1

Figure 1 is a sketch of the curve C with equation

$$y = 2x^{\frac{3}{2}}(4 - x) \quad x \geq 0$$

The point P is the stationary point of C .

(a) Find, using calculus, the x coordinate of P .

(4)

The region R_1 , shown shaded in Figure 1, is bounded by C and the x -axis.

The region R_2 , also shown shaded in Figure 1, is bounded by C , the x -axis and the line with equation $x = k$, where k is a constant.

Given that the area of R_1 is equal to the area of R_2

(b) find, using calculus, the exact value of k .

(4)

T11 Applications of
Differentiation

11_STAT Stationary points

11_STAT_CORE core family
move

WORKED SOLUTION

Question 9

WMA12/01 May/June 2024

WMA12-01_2024_MayJune_Q09

1

Rewrite the curve

$$y = 2x^{3/2}(4 - x) = 8x^{3/2} - 2x^{5/2}.$$

2

Differentiate

$$\frac{dy}{dx} = 12x^{1/2} - 5x^{3/2} = \sqrt{x}(12 - 5x).$$

3

Find the stationary point

For the interior stationary point P , $12 - 5x = 0$, so $x = \frac{12}{5}$.

4

Set up the area condition

R_1 is the area from $x = 0$ to $x = 4$ above the x -axis. R_2 is the area from $x = 4$ to $x = k$ below the x -axis.

5

Use equal areas

Equal areas mean $\int_0^4 y \, dx = -\int_4^k y \, dx$, so $\int_0^k y \, dx = 0$.

6

Integrate y

$$\int 2x^{3/2}(4 - x) \, dx = \int (8x^{3/2} - 2x^{5/2}) \, dx = \frac{16}{5}x^{5/2} - \frac{4}{7}x^{7/2}.$$

7

Solve for k

$$\frac{16}{5}k^{5/2} - \frac{4}{7}k^{7/2} = 0.$$

8

Factor

$$4k^{5/2} \left(\frac{4}{5} - \frac{k}{7} \right) = 0.$$

9

Choose the value beyond four

Since $k > 4$, $\frac{4}{5} - \frac{k}{7} = 0$, so $k = \frac{28}{5}$.

FINAL ANSWER

$$(a) \, x = \frac{12}{5}. \quad (b) \, k = \frac{28}{5}.$$

WMA12-01_2024_MayJune_Q09

T11

11_STAT

11_STAT_CORE

CHAPTER

Applications of Differentiation

FAMILY

Stationary points

VARIANT

core family move

9 marks

WMA12/01 MAY/JUNE (R) 2024

Question 10

Past-paper question

WMA12-01_2024_MayJuneR_Q10

10.




Figure 1

In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.

Figure 1 shows a sketch of part of the curve C with equation

$$y = \frac{9x - x^2}{2\sqrt{x}} \quad x > 0$$

The point P is the stationary point on C .

(a) Find, using calculus, the x coordinate of P .

(4)

The finite region R , shown shaded in Figure 1, is bounded by the curve C , the x -axis and the line with equation $x = 1$

(b) Using calculus, calculate the exact area of R .

(5)

T11 Applications of
Differentiation

11_STAT Stationary points

11_STAT_CORE core family
move

WORKED SOLUTION

Question 10

WMA12/01 May/June (R) 2024

WMA12-01_2024_MayJuneR_Q10

1

Rewrite the curve

$$y = \frac{9x-x^2}{2\sqrt{x}} = \frac{1}{2}(9x^{1/2} - x^{3/2}).$$

2

Differentiate

$$\frac{dy}{dx} = \frac{1}{2} \left(\frac{9}{2}x^{-1/2} - \frac{3}{2}x^{1/2} \right) = \frac{3(3-x)}{4\sqrt{x}}.$$

3

Use the stationary condition

For $x > 0$, the denominator is not zero. So $\frac{dy}{dx} = 0$ when $3 - x = 0$.

4

Find P

Therefore the x coordinate of P is 3.

5

Find where the curve meets the x-axis

$y = 0$ when $9x - x^2 = 0$, so $x = 9$ for the positive intercept.

6

Set up the area

The region is under the curve from $x = 1$ to $x = 9$, so $R = \int_1^9 \frac{9x-x^2}{2\sqrt{x}} dx$.

7

Integrate

$$R = \int_1^9 \frac{1}{2}(9x^{1/2} - x^{3/2})dx = \left[3x^{3/2} - \frac{1}{5}x^{5/2} \right]_1^9.$$

8

Evaluate

$$R = \left(3(27) - \frac{1}{5}(243) \right) - \left(3 - \frac{1}{5} \right) = \frac{148}{5}.$$

FINAL ANSWER

$$(a) x = 3. \quad (b) \frac{148}{5}.$$

WMA12-01_2024_MayJuneR_Q10

T11

11_STAT

11_STAT_CORE

CHAPTER

Applications of Differentiation

FAMILY

Stationary points

VARIANT

core family move

7 marks

WMA12/01 OCTOBER 2024

Question 4

Past-paper question

WMA12-01_2024_Oct_Q04

4. In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

The curve C has equation

$$y = 4x^{\frac{1}{2}} + 9x^{-\frac{1}{2}} + 3 \quad x > 0$$

- (a) Find $\frac{dy}{dx}$ giving each term in simplest form. (2)
- (b) Hence find the x coordinate of the stationary point of C . (2)
- (c) (i) Find $\frac{d^2y}{dx^2}$ giving each term in simplest form.
- (ii) Hence determine the nature of the stationary point of C , giving a reason for your answer. (2)
- (d) State the range of values of x for which y is decreasing. (1)

T11 Applications of
Differentiation

11_STAT Stationary points

11_STAT_CORE core family
move

WORKED SOLUTION

Question 4

WMA12/01 October 2024

WMA12-01_2024_Oct_Q04

1

Rewrite using powers

$$y = 4x^{1/2} + 9x^{-1/2} + 3.$$

2

Differentiate

$$\frac{dy}{dx} = 2x^{-1/2} - \frac{9}{2}x^{-3/2} = \frac{2}{\sqrt{x}} - \frac{9}{2x^{3/2}}.$$

3

Set the derivative to zero

$$\frac{dy}{dx} = \frac{4x-9}{2x^{3/2}}. \text{ Since } x > 0, \text{ the stationary point occurs when } 4x - 9 = 0.$$

4

Find the x coordinate

$$\text{Therefore } x = \frac{9}{4}.$$

5

Differentiate again

$$\frac{d^2y}{dx^2} = -x^{-3/2} + \frac{27}{4}x^{-5/2} = -\frac{1}{x^{3/2}} + \frac{27}{4x^{5/2}}.$$

6

Test the stationary point

$$\frac{d^2y}{dx^2} = \frac{27-4x}{4x^{5/2}}. \text{ At } x = \frac{9}{4}, \text{ the numerator is } 27 - 9 = 18 > 0, \text{ so the stationary point is a minimum.}$$

7

Find where y is decreasing

$$\text{For } x > 0, 2x^{3/2} > 0, \text{ so } \frac{dy}{dx} < 0 \text{ when } 4x - 9 < 0. \text{ Hence } y \text{ is decreasing for } 0 < x < \frac{9}{4}.$$

FINAL ANSWER

WMA12-01_2024_Oct_Q04

$$(a) \frac{2}{\sqrt{x}} - \frac{9}{2x^{3/2}}. \quad (b) x = \frac{9}{4}. \quad (c)(i) -\frac{1}{x^{3/2}} + \frac{27}{4x^{5/2}}. \quad (c)(ii) \text{ Minimum.} \quad (d) 0 < x < \frac{9}{4}.$$

T11

11_STAT

11_STAT_CORE

CHAPTER

Applications of Differentiation

FAMILY

Stationary points

VARIANT

core family move

8 marks

WMA12/01 JANUARY 2026

Question 10

Past-paper question

WMA12-01_2026_Jan_Q10

10.




Figure 2

In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.

Figure 2 shows a sketch of the curve C with equation

$$y = \frac{\sqrt{x}(100 - x^2)}{40} \quad x \geq 0$$

The point P is a stationary point on C .

(a) Use algebraic differentiation to find the exact x coordinate of P .

(4)

The region R_1 , shown shaded in Figure 2, is bounded by C and the x -axis.

The region R_2 , also shown shaded in Figure 2, is bounded by C , the x -axis and the line with equation $x = k$, where k is a constant.

Given that the area of R_1 is equal to the area of R_2

(b) use algebraic integration to find the exact value of k .

(4)

T11 Applications of
Differentiation

11_STAT Stationary points

11_STAT_CORE core family
move

WORKED SOLUTION

Question 10

WMA12/01 January 2026

WMA12-01_2026_Jan_Q10

1

Rewrite the curve

$$y = \frac{\sqrt{x}(100-x^2)}{40} = \frac{5}{2}x^{1/2} - \frac{1}{40}x^{5/2}.$$

2

Differentiate

$$\frac{dy}{dx} = \frac{5}{4}x^{-1/2} - \frac{1}{16}x^{3/2}.$$

3

Use the stationary point

At P , $\frac{dy}{dx} = 0$. Thus $\frac{5}{4}x^{-1/2} = \frac{1}{16}x^{3/2}$. Multiplying by $16x^{1/2}$ gives $20 = x^2$.

4

Find the x-coordinate

Since $x \geq 0$, $x = \sqrt{20} = 2\sqrt{5}$.

5

Integrate the curve

$$\int y \, dx = \int \left(\frac{5}{2}x^{1/2} - \frac{1}{40}x^{5/2} \right) dx = \frac{5}{3}x^{3/2} - \frac{1}{140}x^{7/2}.$$

6

Use equal areas

The signed area from 0 to k is zero because the positive area R_1 equals the negative area R_2 in magnitude.

7

Solve for k

$\frac{5}{3}k^{3/2} - \frac{1}{140}k^{7/2} = 0$. Since $k > 0$, divide by $k^{3/2}$ to get $\frac{5}{3} - \frac{k^2}{140} = 0$, so $k^2 = \frac{700}{3}$.

8

Write the exact value

$$k = \sqrt{\frac{700}{3}} = \frac{10\sqrt{21}}{3}.$$

FINAL ANSWER

WMA12-01_2026_Jan_Q10

$$(a) \, x = 2\sqrt{5}. \quad (b) \, k = \frac{10\sqrt{21}}{3}.$$

PURE MATHEMATICS 2

ADAPTIVE ALL CLASSIFIED

Presentation Edition | Solutions

CLASSIFIED BY IDEA | ROOM TO THINK

Every demand appears clearly, with practical space for the mathematics.

Dr Eslam Ahmed

Elite Mathematics

ElitelGCSE.com | 011 2000 9622

PURE 2

PURE 2 | TOPIC 12

Integration

Your classified route

29

QUESTIONS

212

MARKS

4

FAMILIES

5

VARIANTS

DR ESLAM AHMED

CLASSIFIED BY IDEA

Move from recognition to method, then use the working space to complete the solution.

TOPIC 12 | FAMILY 01

Exact Area

WHAT THIS FAMILY ASKS

Recognise the exact area demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

9 classified questions

78 total marks

T12

12_AREA

12_AREA_CORE

CHAPTER
Integration

FAMILY
Exact area

VARIANT
core family move

9 marks

WMA12/01 OCTOBER 2019

Question 8

Past-paper question

WMA12-01_2019_Oct_Q08

8. Solutions relying on calculator technology are not acceptable in this question.

(i)




Figure 2

Figure 2 shows a sketch of part of a curve with equation

$$y = \frac{8\sqrt{x} - 5}{2x^2} \quad x > 0$$

The region R , shown shaded in Figure 2, is bounded by the curve, the line with equation $x = 2$, the x -axis and the line with equation $x = 4$

Find the exact area of R .

(5)

(ii) Find the value of the constant k such that

$$\int_{-3}^6 \left(\frac{1}{2}x^2 + k \right) dx = 55$$

(4)

T12 Integration

12_AREA Exact area

12_AREA_CORE core family
move

WORKED SOLUTION

Question 8

WMA12/01 October 2019

WMA12-01_2019_Oct_Q08

1

Rewrite the curve

$$y = \frac{8\sqrt{x}-5}{2x^2} = 4x^{-3/2} - \frac{5}{2}x^{-2}.$$

2

Integrate the curve

$$\int \left(4x^{-3/2} - \frac{5}{2}x^{-2}\right) dx = -8x^{-1/2} + \frac{5}{2x}.$$

3

Apply the limits for the area

$$\left[-8x^{-1/2} + \frac{5}{2x}\right]_2^4 = \left(-4 + \frac{5}{8}\right) - \left(-4\sqrt{2} + \frac{5}{4}\right).$$

4

Simplify the exact area

$$\text{This gives } 4\sqrt{2} - \frac{37}{8}.$$

5

Integrate the second expression

$$\int \left(\frac{1}{2}x^2 + k\right) dx = \frac{x^3}{6} + kx.$$

6

Use the limits

$$\left[\frac{x^3}{6} + kx\right]_{-3}^6 = (36 + 6k) - \left(-\frac{9}{2} - 3k\right) = \frac{81}{2} + 9k.$$

7

Solve for k

$$\frac{81}{2} + 9k = 55, \text{ so } 9k = \frac{29}{2} \text{ and } k = \frac{29}{18}.$$

FINAL ANSWER

$$(i) 4\sqrt{2} - \frac{37}{8}. \quad (ii) k = \frac{29}{18}.$$

WMA12-01_2019_Oct_Q08

T12

12_AREA

12_AREA_CORE

CHAPTER
Integration

FAMILY
Exact area

VARIANT
core family move

6 marks

WMA12/01 JANUARY 2020

Question 4

Past-paper question

WMA12-01_2020_Jan_Q04

4.




Figure 1

Figure 1 shows a sketch of the curve with equation

$$y = 2x^2 + 7 \quad x \geq 0$$

The finite region R , shown shaded in Figure 1, is bounded by the curve, the y -axis and the line with equation $y = 17$

Find the exact area of R .

(6)

T12 Integration

12_AREA Exact area

12_AREA_CORE core family
move

WORKED SOLUTION

Question 4

WMA12/01 January 2020

WMA12-01_2020_Jan_Q04

1

Find the intersection with y equals 17

The curve is $y = 2x^2 + 7$. Set $2x^2 + 7 = 17$ to get $x^2 = 5$, so the right-hand limit is $x = \sqrt{5}$.

2

Set up the area

The shaded area is between $y = 17$ and $y = 2x^2 + 7$, from $x = 0$ to $x = \sqrt{5}$.

3

Integrate

$$A = \int_0^{\sqrt{5}} \{17 - (2x^2 + 7)\} dx = \int_0^{\sqrt{5}} (10 - 2x^2) dx.$$

4

Evaluate the exact area

$$A = \left[10x - \frac{2}{3}x^3 \right]_0^{\sqrt{5}} = 10\sqrt{5} - \frac{2}{3}(5\sqrt{5}) = \frac{20\sqrt{5}}{3}.$$

FINAL ANSWER

$$\frac{20\sqrt{5}}{3}.$$

WMA12-01_2020_Jan_Q04

T12

12_AREA

12_AREA_CORE

CHAPTER
Integration

FAMILY
Exact area

VARIANT
core family move

10 marks

WMA12/01 JANUARY 2022

Question 9

Past-paper question

WMA12-01_2022_Jan_Q09

9.




Figure 2

Figure 2 shows

- the curve C with equation $y = x - x^2$
- the line l with equation $y = mx$, where m is a constant and $0 < m < 1$

The line and the curve intersect at the origin O and at the point P .

- (a) Find, in terms of m , the coordinates of P .

(2)

The region R_1 , shown shaded in Figure 2, is bounded by C and l .

- (b) Show that the area of R_1 is

$$\frac{(1-m)^3}{6}$$

(5)

The region R_2 , also shown shaded in Figure 2, is bounded by C , the x -axis and l .

Given that the area of R_1 is equal to the area of R_2

- (c) find the exact value of m .

(3)

T12 Integration

12_AREA Exact area

12_AREA_CORE core family
move

WORKED SOLUTION

Question 9

WMA12/01 January 2022

WMA12-01_2022_Jan_Q09

1

Find the second intersectionAt an intersection, $x - x^2 = mx$, so $x(1 - m - x) = 0$.

2

State PThe non-zero intersection has $x = 1 - m$, and then $y = m(1 - m)$, so $P = (1 - m, m(1 - m))$.

3

Set up area R oneRegion R_1 is between the curve and the line from $x = 0$ to $x = 1 - m$.

4

Integrate the difference

$$A_{R_1} = \int_0^{1-m} \{(x - x^2) - mx\} dx = \int_0^{1-m} \{(1 - m)x - x^2\} dx.$$

5

Evaluate

$$A_{R_1} = \left[\frac{1-m}{2} x^2 - \frac{1}{3} x^3 \right]_0^{1-m} = \frac{(1-m)^3}{2} - \frac{(1-m)^3}{3} = \frac{(1-m)^3}{6}.$$

6

Use the total curve areaThe total area under $y = x - x^2$ from $x = 0$ to $x = 1$ is $\int_0^1 (x - x^2) dx = \frac{1}{6}$.

7

Use equal regionsSince R_1 and R_2 are equal and together make this total area, each has area $\frac{1}{12}$.

8

Solve for m

$$\frac{(1-m)^3}{6} = \frac{1}{12}, \text{ so } (1 - m)^3 = \frac{1}{2} \text{ and } m = 1 - \sqrt[3]{\frac{1}{2}}.$$

FINAL ANSWER

WMA12-01_2022_Jan_Q09

(a) $P = (1 - m, m(1 - m))$. (b) **Proven:** $\frac{(1-m)^3}{6}$. (c) $m = 1 - \sqrt[3]{\frac{1}{2}}$.

T12

12_AREA

12_AREA_CORE

CHAPTER
Integration

FAMILY
Exact area

VARIANT
core family move

8 marks

WMA12/01 JANUARY 2023
Question 9

Past-paper question

WMA12-01_2023_Jan_Q09

9. In this question you must show all stages of your working.

Solutions based entirely on calculator technology are not acceptable.




Figure 3

Figure 3 shows

- the curve C with equation $y = x^2 - 4x + 5$
- the line l with equation $y = 2$

The curve C intersects the y -axis at the point D .

(a) Write down the coordinates of D .

(1)

The curve C intersects the line l at the points E and F , as shown in Figure 3.

(b) Find the x coordinate of E and the x coordinate of F .

(2)

Shown shaded in Figure 3 is

- the region R_1 which is bounded by C , l and the y -axis
- the region R_2 which is bounded by C and the line segments EF and DF

Given that $\frac{\text{area of } R_1}{\text{area of } R_2} = k$, where k is a constant,

(c) use algebraic integration to find the exact value of k , giving your answer as a simplified fraction.

(5)

T12 Integration

12_AREA Exact area

12_AREA_CORE core family
move

WORKED SOLUTION

Question 9

WMA12/01 January 2023

WMA12-01_2023_Jan_Q09

1

Find DAt the y -axis, $x = 0$. On C , $y = 0^2 - 4(0) + 5 = 5$, so $D = (0, 5)$.

2

Find intersections with the lineThe line is $y = 2$, so solve $x^2 - 4x + 5 = 2$.

3

Solve the quadratic $x^2 - 4x + 3 = 0$, so $(x - 1)(x - 3) = 0$. Hence $x_E = 1$ and $x_F = 3$.

4

Area of R1 R_1 lies between the curve and the line $y = 2$ from $x = 0$ to $x = 1$.

5

Integrate R1

$$R_1 = \int_0^1 (x^2 - 4x + 5 - 2) dx = \int_0^1 (x^2 - 4x + 3) dx.$$

6

Evaluate R1

$$R_1 = \left[\frac{x^3}{3} - 2x^2 + 3x \right]_0^1 = \frac{1}{3} - 2 + 3 = \frac{4}{3}.$$

7

Find the line DFThe line through $D = (0, 5)$ and $F = (3, 2)$ has gradient -1 , so its equation is $y = 5 - x$.

8

Split R2For $0 \leq x \leq 1$, R_2 is between $y = 5 - x$ and the curve. For $1 \leq x \leq 3$, it is between $y = 5 - x$ and $y = 2$.

9

Integrate R2

$$R_2 = \int_0^1 [(5 - x) - (x^2 - 4x + 5)] dx + \int_1^3 [(5 - x) - 2] dx.$$

10

Evaluate R2

$$R_2 = \int_0^1 (-x^2 + 3x) dx + \int_1^3 (3 - x) dx = \frac{7}{6} + 2 = \frac{19}{6}.$$

11

Find the ratio

$$k = \frac{\text{area of } R_1}{\text{area of } R_2} = \frac{4/3}{19/6} = \frac{8}{19}.$$

FINAL ANSWER

WMA12-01_2023_Jan_Q09

$$(a) D = (0, 5). \quad (b) x_E = 1, x_F = 3. \quad (c) k = \frac{8}{19}.$$

DR ESLAM AHMED

T12

12_AREA

12_AREA_CORE

CHAPTER
Integration

FAMILY
Exact area

VARIANT
core family move

10 marks

WMA12/01 MAY/JUNE 2023
Question 10

Past-paper question

WMA12-01_2023_MayJune_Q10

10. The curve C has equation

$$y = \frac{(x-k)^2}{\sqrt{x}} \quad x > 0$$

where k is a **positive** constant.

(a) Show that

$$\int_1^{16} \frac{(x-k)^2}{\sqrt{x}} \, dx = ak^2 + bk + \frac{2046}{5}$$

where a and b are integers to be found.

(5)




Figure 1

Figure 1 shows a sketch of the curve C and the line l .

Given that l intersects C at the point $A(1, 9)$ and at the point $B(16, q)$ where q is a constant,

(b) show that $k = 4$

(2)

The region R , shown shaded in Figure 1, is bounded by C and l

Using the answers to parts (a) and (b),

(c) find the area of region R

(3)

T12 Integration

12_AREA Exact area

12_AREA_CORE core family
move

WORKED SOLUTION

Question 10

WMA12/01 May/June 2023

WMA12-01_2023_MayJune_Q10

1

Expand the integrand

$$\frac{(x-k)^2}{\sqrt{x}} = \frac{x^2-2kx+k^2}{x^{1/2}} = x^{3/2} - 2kx^{1/2} + k^2x^{-1/2}.$$

2

Integrate

$$\int (x^{3/2} - 2kx^{1/2} + k^2x^{-1/2}) dx = \frac{2}{5}x^{5/2} - \frac{4k}{3}x^{3/2} + 2k^2x^{1/2}.$$

3

Substitute the limits

$$\text{From 1 to 16, this gives } \left(\frac{2048}{5} - \frac{256k}{3} + 8k^2\right) - \left(\frac{2}{5} - \frac{4k}{3} + 2k^2\right).$$

4

Simplify the integral

$$\text{The result is } 6k^2 - 84k + \frac{2046}{5}, \text{ so } a = 6 \text{ and } b = -84.$$

5

Use point A

$$\text{Since } A(1,9) \text{ lies on } C, 9 = \frac{(1-k)^2}{\sqrt{1}} = (1-k)^2.$$

6

Find k

$$\text{Thus } 1 - k = \pm 3, \text{ so } k = 4 \text{ or } k = -2. \text{ Since } k \text{ is positive, } k = 4.$$

7

Find point B

$$\text{When } k = 4 \text{ and } x = 16, q = \frac{(16-4)^2}{\sqrt{16}} = \frac{144}{4} = 36, \text{ so } B = (16, 36).$$

8

Find the line

$$\text{The line through } A(1,9) \text{ and } B(16,36) \text{ has gradient } \frac{36-9}{16-1} = \frac{9}{5}, \text{ so } l \text{ has equation } y = \frac{9}{5}x + \frac{36}{5}.$$

9

Area under the line

$$\int_1^{16} \left(\frac{9}{5}x + \frac{36}{5}\right) dx = \left[\frac{9}{10}x^2 + \frac{36}{5}x\right]_1^{16} = \frac{675}{2}.$$

10

Area under the curve

From part (a), with $k = 4$, $\int_1^{16} \frac{(x-4)^2}{\sqrt{x}} dx = 6(4)^2 - 84(4) + \frac{2046}{5} = \frac{846}{5}$.

11

Subtract curve from line

The shaded area is $\frac{675}{2} - \frac{846}{5} = \frac{1683}{10}$.

FINAL ANSWER

WMA12-01_2023_MayJune_Q10

(a) $a = 6$, $b = -84$. (b) $k = 4$. (c) **Area** = $\frac{1683}{10}$.

DR ESLAM AHMED

T12

12_AREA

12_AREA_CORE

CHAPTER
Integration

FAMILY
Exact area

VARIANT
core family move

8 marks

WMA12/01 OCTOBER 2024

Question 8

Past-paper question

WMA12-01_2024_Oct_Q08

8. In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.




Figure 1

Figure 1 shows a sketch of part of the curve C_1 with equation

$$y = x^2 + 3 \quad x > 0$$

and part of the curve C_2 with equation

$$y = 13 - \frac{9}{x^2} \quad x > 0$$

The curves C_1 and C_2 intersect at the points P and Q as shown in Figure 1.

- (a) Use algebra to find the x coordinate of P and the x coordinate of Q .

(4)

The finite region R , shown shaded in Figure 1, is bounded by C_1 and C_2

- (b) Use algebraic integration to find the exact area of R .

(4)

T12 Integration

12_AREA Exact area

12_AREA_CORE core family
move

WORKED SOLUTION

Question 8

WMA12/01 October 2024

WMA12-01_2024_Oct_Q08

1

Equate the curves

$$x^2 + 3 = 13 - \frac{9}{x^2}.$$

2

Clear the denominator

Multiplying by x^2 gives $x^4 + 3x^2 = 13x^2 - 9$, so $x^4 - 10x^2 + 9 = 0$.

3

Factor

Let $u = x^2$. Then $u^2 - 10u + 9 = (u - 1)(u - 9) = 0$, so $x^2 = 1$ or $x^2 = 9$.

4

Use x greater than zero

Since $x > 0$, the intersections have $x = 1$ and $x = 3$. Thus $x_P = 1$ and $x_Q = 3$.

5

Set up the area

Between $x = 1$ and $x = 3$, curve C_2 is above curve C_1 , so $R = \int_1^3 \left(13 - \frac{9}{x^2} - (x^2 + 3)\right) dx$.

6

Simplify the integrand

$$R = \int_1^3 (10 - x^2 - 9x^{-2}) dx.$$

7

Integrate

$$R = \left[10x - \frac{x^3}{3} + \frac{9}{x}\right]_1^3.$$

8

Evaluate

$$R = (30 - 9 + 3) - \left(10 - \frac{1}{3} + 9\right) = 24 - \frac{56}{3} = \frac{16}{3}.$$

FINAL ANSWER

WMA12-01_2024_Oct_Q08

(a) $x_P = 1, x_Q = 3$. (b) $\frac{16}{3}$.

T12

12_AREA

12_AREA_CORE

CHAPTER
Integration

FAMILY
Exact area

VARIANT
core family move

12 marks

WMA12/01 JANUARY 2025

Question 9

Past-paper question

WMA12-01_2025_Jan_Q09

9.




Figure 3

In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.

Figure 3 shows a sketch of part of the curve with equation

$$y = \frac{9x^2(5 - \sqrt{x})}{5} \quad x \geq 0$$

The curve has a turning point at the point M , as shown in Figure 3.

(a) Using calculus, find the coordinates of M .

(5)

The curve crosses the x -axis at the point P , as shown in Figure 3.

(b) Use algebra to find the x coordinate of P .

(2)

The finite region R , shown shaded in Figure 3, is bounded by the curve, the line through M parallel to the x -axis and the line through P parallel to the y -axis.

(c) Use algebraic integration to find the area of R , giving your answer to one decimal place.

(5)

T12 Integration

12_AREA Exact area

12_AREA_CORE core family
move

WORKED SOLUTION

Question 9

WMA12/01 January 2025

WMA12-01_2025_Jan_Q09

1

Rewrite the curve

$$y = \frac{9x^2(5-\sqrt{x})}{5} = 9x^2 - \frac{9}{5}x^{5/2}.$$

2

Differentiate

$$\frac{dy}{dx} = 18x - \frac{9}{2}x^{3/2} = \frac{9}{2}x(4 - \sqrt{x}).$$

3

Find the turning point x coordinate

For the turning point shown away from the origin, set $4 - \sqrt{x} = 0$. Hence $x = 16$.

4

Find the y coordinate

$$\text{At } x = 16, y = \frac{9(16)^2(5-4)}{5} = \frac{2304}{5}.$$

5

State M

Therefore $M\left(16, \frac{2304}{5}\right)$.

6

Find the positive x intercept

For the positive intercept, $\frac{9x^2(5-\sqrt{x})}{5} = 0$ gives $5 - \sqrt{x} = 0$, so $x = 25$.

7

Set up the area

The top boundary is $y = \frac{2304}{5}$ and the curve is below it from $x = 16$ to $x = 25$. So

$$R = \int_{16}^{25} \left(\frac{2304}{5} - 9x^2 + \frac{9}{5}x^{5/2} \right) dx.$$

8

Integrate

$$R = \left[\frac{2304}{5}x - 3x^3 + \frac{18}{35}x^{7/2} \right]_{16}^{25}.$$

9

Evaluate

$R = \frac{9189}{7} = 1312.714 \dots$, so the area is 1312.7 to one decimal place.

FINAL ANSWER

(a) $M\left(16, \frac{2304}{5}\right)$. (b) $x_P = 25$. (c) 1312.7.

WMA12-01_2025_Jan_Q09

T12

12_AREA

12_AREA_CORE

CHAPTER
Integration

FAMILY
Exact area

VARIANT
core family move

5 marks

WMA12/01 MAY/JUNE 2025

Question 3

Past-paper question

WMA12-01_2025_MayJune_Q03

3:




Figure 1

In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.

Figure 1 shows a sketch of the curve C with equation

$$y = 3x + \frac{16}{x^2} - 8 \quad x > 0$$

The points $P(1, 11)$ and $Q(4, 5)$ lie on C and are shown in Figure 1.

The region R , shown shaded in Figure 1, is bounded by C and line segment PQ .

Use algebraic integration to find the area of R .

(5)

T12 Integration

12_AREA Exact area

12_AREA_CORE core family
move

WORKED SOLUTION

Question 3

WMA12/01 May/June 2025

WMA12-01_2025_MayJune_Q03

1

Find the line PQ

The gradient of PQ is $\frac{5-11}{4-1} = -2$, so the line through $P(1, 11)$ is $y - 11 = -2(x - 1)$.

2

Write the line equation

Thus PQ has equation $y = -2x + 13$.

3

Set up the area

From $x = 1$ to $x = 4$, the line segment is above the curve. Hence $R = \int_1^4 \left[(-2x + 13) - \left(3x + \frac{16}{x^2} - 8 \right) \right] dx$.

4

Simplify the integrand

$$R = \int_1^4 (21 - 5x - 16x^{-2}) dx.$$

5

Integrate

$$R = \left[21x - \frac{5}{2}x^2 + \frac{16}{x} \right]_1^4.$$

6

Evaluate

$$R = (84 - 40 + 4) - \left(21 - \frac{5}{2} + 16 \right) = 48 - \frac{69}{2} = \frac{27}{2}.$$

FINAL ANSWER

$$\frac{27}{2}.$$

WMA12-01_2025_MayJune_Q03

T12

12_AREA

12_AREA_CORE

CHAPTER
Integration

FAMILY
Exact area

VARIANT
core family move

10 marks

WMA12/01 OCTOBER 2025

Question 7

Past-paper question

WMA12-01_2025_Oct_Q07

7.




Figure 3

In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.

Figure 3 shows a sketch of part of the curve C with equation

$$y = x^3 - 4x^{\frac{5}{2}} - kx^{\frac{1}{2}} + 28x - 44 \quad x \geq 0$$

where k is a positive constant.

- (a) Find $\frac{dy}{dx}$ in simplest form. (2)

The point T , shown in Figure 3, is a minimum stationary point on C .

Given that the x coordinate of T is 9

- (b) show that $k = 6$ (2)

The line through T parallel to the x -axis meets the y -axis at the point N .

The finite region R , shown shaded in Figure 3, is bounded by C , the y -axis and the line segment NT .

- (c) Use algebraic integration to find the area of R , giving the answer to 3 significant figures. (6)

T12 Integration

12_AREA Exact area

12_AREA_CORE core family
move

WORKED SOLUTION

Question 7

WMA12/01 October 2025

WMA12-01_2025_Oct_Q07

1

Differentiate

For $y = x^3 - 4x^{5/2} - kx^{1/2} + 28x - 44$, $\frac{dy}{dx} = 3x^2 - 10x^{3/2} - \frac{k}{2}x^{-1/2} + 28$.

2

Write in simplest form

$$\frac{dy}{dx} = 3x^2 - 10x^{3/2} - \frac{k}{2\sqrt{x}} + 28.$$

3

Use the stationary point

At T , $x = 9$ and $\frac{dy}{dx} = 0$.

4

Substitute x equals nine

$$0 = 3(9)^2 - 10(9)^{3/2} - \frac{k}{2\sqrt{9}} + 28 = 243 - 270 - \frac{k}{6} + 28.$$

5

Find k

This simplifies to $0 = 1 - \frac{k}{6}$, so $k = 6$.

6

Find the horizontal line

With $k = 6$, the y coordinate of T is $9^3 - 4(9)^{5/2} - 6\sqrt{9} + 28(9) - 44 = -53$.

7

Set up the area

The line NT is $y = -53$. The region is between the curve and this line from $x = 0$ to $x = 9$.

8

Write the integral

$$R = \int_0^9 (x^3 - 4x^{5/2} - 6x^{1/2} + 28x - 44 + 53) dx.$$

9

Integrate

$$R = \left[\frac{x^4}{4} - \frac{8}{7}x^{7/2} - 4x^{3/2} + 14x^2 + 9x \right]_0^9.$$

10

Evaluate

$R = \frac{6939}{28} = 247.821 \dots$, so the area is 248 to 3 significant figures.

FINAL ANSWER

WMA12-01_2025_Oct_Q07

(a) $3x^2 - 10x^{3/2} - \frac{k}{2\sqrt{x}} + 28$. (b) **Proven.** (c) 248.

DR ESLAM AHMED

TOPIC 12 | FAMILY 02

Power Integration

WHAT THIS FAMILY ASKS

Recognise the power integration demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

2 classified questions

12 total marks

T12

12_POWER

12_POWER_CORE

CHAPTER
Integration

FAMILY
Power integration

VARIANT
core family move

7 marks

WMA12/01 MAY/JUNE 2022

Question 7

Past-paper question

WMA12-01_2022_MayJune_Q07

7. $f(x) = Ax^3 + 6x^2 - 4x + B$

where A and B are constants.

Given that

- $(x + 2)$ is a factor of $f(x)$

- $\int_3^5 f(x) dx = 176$

find the value of A and the value of B .

(7)

T12 Integration

12_POWER Power integration

12_POWER_CORE core
family move

WORKED SOLUTION

Question 7

WMA12/01 May/June 2022

WMA12-01_2022_MayJune_Q07

1

Use the factor conditionSince $(x + 2)$ is a factor, $f(-2) = 0$.

2

Form the first equation $A(-2)^3 + 6(-2)^2 - 4(-2) + B = 0$, so $-8A + 32 + B = 0$ and $B = 8A - 32$.

3

Set up the integral

$$\int_3^5 (Ax^3 + 6x^2 - 4x + B) dx = 176.$$

4

Integrate

$$\left[\frac{A}{4}x^4 + 2x^3 - 2x^2 + Bx \right]_3^5 = 176.$$

5

Simplify the second equationThis gives $136A + 164 + 2B = 176$, so $68A + B = 6$.

6

Solve simultaneouslyUsing $B = 8A - 32$ in $68A + B = 6$ gives $76A = 38$, so $A = \frac{1}{2}$.

7

Find B

$$B = 8\left(\frac{1}{2}\right) - 32 = -28.$$

FINAL ANSWER

$$A = \frac{1}{2}, \quad B = -28.$$

WMA12-01_2022_MayJune_Q07

T12

12_POWER

12_POWER_CORE

CHAPTER
Integration

FAMILY
Power integration

VARIANT
core family move

5 marks

WMA12/01 JANUARY 2026

Question 4

Past-paper question

WMA12-01_2026_Jan_Q04

4.

**In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.**

Given that k is a positive constant and

$$\int_k^{2k} \left(\frac{12}{x^2} + 4 \right) dx = 14$$

find the possible values of k .

(5)

T12 Integration

12_POWER Power integration

12_POWER_CORE core
family move

WORKED SOLUTION

Question 4

WMA12/01 January 2026

WMA12-01_2026_Jan_Q04

1

Integrate first

$$\int \left(\frac{12}{x^2} + 4 \right) dx = \int (12x^{-2} + 4) dx = -\frac{12}{x} + 4x.$$

2

Apply the limits

$$\left[-\frac{12}{x} + 4x \right]_k^{2k} = \left(-\frac{6}{k} + 8k \right) - \left(-\frac{12}{k} + 4k \right) = \frac{6}{k} + 4k.$$

3

Form a quadratic

Given the integral is 14, $\frac{6}{k} + 4k = 14$. Multiplying by k gives $4k^2 - 14k + 6 = 0$, so $2k^2 - 7k + 3 = 0$.

4

Solve for k

$(2k - 1)(k - 3) = 0$, hence $k = \frac{1}{2}$ or $k = 3$. Both are positive, so both are valid.

FINAL ANSWER

$$k = \frac{1}{2} \text{ or } k = 3.$$

WMA12-01_2026_Jan_Q04

TOPIC 12 | FAMILY 03

Transformed Estimates

WHAT THIS FAMILY ASKS

Recognise the transformed estimates demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 For $f(x+c)$ or $f(-x)$, check the limits by substitution.

V02 Do not round the known estimate too early.

8 classified questions

53 total marks

T12

12_SHIFT

12_SHIFT_HV01

CHAPTER
Integration

FAMILY
Transformed estimates

VARIANT
For $\int f(x+c)$ or $\int f(-x)$, check
the limits by substitution.

6 marks

WMA12/01 JANUARY 2023

Question 1

Past-paper question

WMA12-01_2023_Jan_Q01

1.




Figure 1

Figure 1 shows a sketch of part of the curve with equation $y = f(x)$

The table below shows some corresponding values of x and y for this curve.

The values of y are given to 3 decimal places.

x	-1	-0.5	0	0.5	1
y	2.287	4.470	6.719	7.291	2.834

Using the trapezium rule with all the values of y in the given table,

(a) obtain an estimate for

$$\int_{-1}^1 f(x) \, dx$$

giving your answer to 2 decimal places.

(3)

(b) Use your answer to part (a) to estimate

(i) $\int_{-1}^1 (f(x) - 2) \, dx$

(ii) $\int_1^3 f(x-2) \, dx$

(3)

T12 Integration

12_SHIFT Transformed
estimates12_SHIFT_HV01 For $f(x+c)/$
or $f(-x)/$, check the limits by
substitution.

WORKED SOLUTION

Question 1

WMA12/01 January 2023

WMA12-01_2023_Jan_Q01

1

Use the trapezium widthThe x values run from -1 to 1 in steps of 0.5 , so $h = 0.5$.

2

Apply the trapezium rule

$$\int_{-1}^1 f(x) dx \approx \frac{0.5}{2} [2.287 + 2.834 + 2(4.470 + 6.719 + 7.291)].$$

3

Evaluate the estimateThis gives 10.52025 , so the estimate is 10.52 to 2 decimal places.

4

Estimate the shifted-down area

$$\int_{-1}^1 (f(x) - 2) dx = \int_{-1}^1 f(x) dx - \int_{-1}^1 2 dx.$$

5

Subtract the rectangleThe interval has width 2 , so $\int_{-1}^1 2 dx = 4$. Hence the estimate is $10.52 - 4 = 6.52$.

6

Use a substitutionFor $\int_1^3 f(x-2) dx$, let $u = x - 2$. When $x = 1$, $u = -1$, and when $x = 3$, $u = 1$.

7

Recognise the same integralSo $\int_1^3 f(x-2) dx = \int_{-1}^1 f(u) du$, which has the same estimate 10.52 .**FINAL ANSWER**

WMA12-01_2023_Jan_Q01

(a) 10.52 . (b)(i) 6.52 . (b)(ii) 10.52 .

T12

12_SHIFT

12_SHIFT_HV01

CHAPTER
Integration

FAMILY
Transformed estimates

VARIANT
For $f(x+c)$ or $f(-x)$, check
the limits by substitution.

9 marks

WMA12/01 JANUARY 2024
Question 4

Past-paper question

WMA12-01_2024_Jan_Q04

4. (a) Sketch the curve with equation

$$y = a^{-x} + 4$$

where a is a constant and $a > 1$

On your sketch show

- the coordinates of the point of intersection of the curve with the y -axis
- the equation of the asymptote to the curve.

(3)

x	-4	-1.5	1	3.5	6	8.5
y	13	6.280	4.577	4.146	4.037	4.009

The table above shows corresponding values of x and y for $y = 3^{-\frac{1}{2}x} + 4$

The values of y are given to four significant figures, as appropriate.

Using the trapezium rule with all the values of y in the table,

- (b) find an approximate value for

$$\int_{-4}^{8.5} \left(3^{-\frac{1}{2}x} + 4 \right) dx$$

giving your answer to two significant figures.

(3)

- (c) Using the answer to part (b), find an approximate value for

(i) $\int_{-4}^{8.5} \left(3^{-\frac{1}{2}x} \right) dx$

(ii) $\int_{-4}^{8.5} \left(3^{-\frac{1}{2}x} + 4 \right) dx + \int_{-8.5}^4 \left(3^{\frac{1}{2}x} + 4 \right) dx$

(3)

T12 Integration

12_SHIFT Transformed
estimates12_SHIFT_HV01 For $f(x+c)/$
or $f(x-c)/$, check the limits by
substitution.

WORKED SOLUTION

Question 4

WMA12/01 January 2024

WMA12-01_2024_Jan_Q04

1

Find the y-interceptFor $y = a^{-x} + 4$, set $x = 0$: $y = a^0 + 4 = 5$, so the curve meets the y -axis at $(0, 5)$.

2

Find the asymptoteSince $a > 1$, $a^{-x} \rightarrow 0$ as $x \rightarrow \infty$, so the horizontal asymptote is $y = 4$.

3

Describe the sketchThe curve is decreasing, passes through $(0, 5)$, and approaches the line $y = 4$ from above.

4

Use the trapezium widthThe x values increase by 2.5, so $h = 2.5$.

5

Apply the trapezium rule

$$\int_{-4}^{8.5} (3^{-x/2} + 4) dx \approx \frac{2.5}{2} [13 + 4.009 + 2(6.280 + 4.577 + 4.146 + 4.037)].$$

6

Round part b

This gives 68.86125, so to two significant figures the estimate is 69.

7

Remove the constant part

$$\int_{-4}^{8.5} 4 dx = 4(12.5) = 50.$$

8

Estimate part c iUsing the answer to part (b), $\int_{-4}^{8.5} 3^{-x/2} dx \approx 69 - 50 = 19$.

9

Use symmetry by substitutionFor $\int_{-8.5}^4 (3^{x/2} + 4) dx$, let $u = -x$. This changes it to the same integral as part (b).

10

Estimate part c ii

Therefore the two integrals in part (c)(ii) are approximately equal, so the total is $69 + 69 = 138$.

FINAL ANSWER

WMA12-01_2024_Jan_Q04

(a) **y-intercept** $(0, 5)$ and **asymptote** $y = 4$. (b) 69. (c)(i) 19. (c)(ii) 138.

DR ESLAM AHMED

T12

12_SHIFT

12_SHIFT_HV01

CHAPTER
Integration

FAMILY
Transformed estimates

VARIANT
For $\int f(x+c)$ or $\int f(-x)$, check
the limits by substitution.

5 marks

WMA12/01 MAY/JUNE (R) 2024

Question 2

Past-paper question

WMA12-01_2024_MayJuneR_Q02

2. The table shows corresponding values of x and y for a continuous curve with equation $y = f(x)$ between $x = -4$ and $x = 5$, where a is a constant.

x	-4	-2.5	-1	0.5	2	3.5	5
y	4.16	2.91	a	1.73	1.37	1.43	2.28

The trapezium rule is used with all the y values in the table to find an approximation for

$$\int_{-4}^5 f(x) \, dx$$

Given that the value of this approximation is 19.3

- (a) find the value of the constant a to 3 significant figures.

(3)

- (b) Use the given answer of 19.3 to find an approximate value for

$$\int_{-4}^5 (2f(x) - 3) \, dx$$

(2)

T12 Integration

12_SHIFT Transformed
estimates12_SHIFT_HV01 For $\int (f(x+c))$
or $\int (f(-x))$, check the limits by
substitution.

WORKED SOLUTION

Question 2

WMA12/01 May/June (R) 2024

WMA12-01_2024_MayJuneR_Q02

1

Use the trapezium widthThe x values increase by 1.5, so $h = 1.5$.

2

Set up the trapezium estimate

$$19.3 = \frac{1.5}{2} [4.16 + 2.28 + 2(2.91 + a + 1.73 + 1.37 + 1.43)].$$

3

Solve for a This simplifies to $19.3 = 15.99 + 1.5a$, so $a = \frac{3.31}{1.5} = 2.206 \dots$

4

Round a Therefore $a = 2.21$ to 3 significant figures.

5

Use the transformed integral

$$\int_{-4}^5 (2f(x) - 3) dx = 2 \int_{-4}^5 f(x) dx - \int_{-4}^5 3 dx.$$

6

Substitute the given estimateThis is $2(19.3) - 3(9) = 38.6 - 27 = 11.6$.**FINAL ANSWER**(a) $a = 2.21$ to 3 s.f. (b) 11.6.

WMA12-01_2024_MayJuneR_Q02

T12

12_SHIFT

12_SHIFT_HV01

CHAPTER
Integration

FAMILY
Transformed estimates

VARIANT
For $\int f(x+c)$ or $\int f(-x)$, check
the limits by substitution.

5 marks

WMA12/01 OCTOBER 2024

Question 1

Past-paper question

WMA12-01_2024_Oct_Q01

1. A continuous curve has equation $y = f(x)$.

A table of values of x and y for $y = f(x)$ is shown below.

x	0.5	1.75	3	4.25	5.5
y	3.479	6.101	7.448	6.823	5.182

Using the trapezium rule with all the values of y in the given table,

- (a) find an estimate for

$$\int_{0.5}^{5.5} f(x) \, dx$$

giving your answer to one decimal place.

(3)

- (b) Using your answer to part (a) and making your method clear, estimate

$$\int_{0.5}^{5.5} (f(x) + 4x) \, dx$$

(2)

T12 Integration

12_SHIFT Transformed
estimates12_SHIFT_HV01 For $f(x+c)$ or $f(-x)$, check the limits by substitution.

WORKED SOLUTION

Question 1

WMA12/01 October 2024

WMA12-01_2024_Oct_Q01

1

Find the strip widthThe x values increase by 1.25, so the trapezium width is $h = 1.25$.

2

Apply the trapezium rule

$$\int_{0.5}^{5.5} f(x) dx \approx \frac{1.25}{2} \{3.479 + 5.182 + 2(6.101 + 7.448 + 6.823)\}.$$

3

Evaluate the estimate

This gives 30.878125, which is 30.9 to one decimal place.

4

Split the second integral

$$\int_{0.5}^{5.5} (f(x) + 4x) dx = \int_{0.5}^{5.5} f(x) dx + \int_{0.5}^{5.5} 4x dx.$$

5

Integrate $4x$

$$\int_{0.5}^{5.5} 4x dx = [2x^2]_{0.5}^{5.5} = 2(5.5)^2 - 2(0.5)^2 = 60.$$

6

Use part aUsing the estimate from part (a), the required value is $30.9 + 60 = 90.9$.**FINAL ANSWER**

(a) 30.9. (b) 90.9.

WMA12-01_2024_Oct_Q01

T12

12_SHIFT

12_SHIFT_HV02

CHAPTER
Integration

FAMILY
Transformed estimates

VARIANT
Do not round the known
estimate too early.

9 marks

WMA12/01 MAY/JUNE 2024

Question 6

Past-paper question

WMA12-01_2024_MayJune_Q06

6. (a) Sketch the curve with equation

$$y = a^x + 4$$

where a is a positive constant greater than 1

On your sketch, show

- the coordinates of the point of intersection of the curve with the y -axis
- the equation of the asymptote of the curve

(3)

x	2	2.3	2.6	2.9	3.2	3.5
y	0	0.3246	0.8629	1.6643	2.7896	4.3137

The table shows corresponding values of x and y for

$$y = 2^x - 2x$$

with the values of y given to 4 decimal places as appropriate.

Using the trapezium rule with all the values of y in the given table,

- (b) obtain an estimate for $\int_2^{3.5} (2^x - 2x) \, dx$, giving your answer to 2 decimal places.

(3)

- (c) Using your answer to part (b) and making your method clear, estimate

(i) $\int_2^{3.5} (2^x + 2x) \, dx$

(ii) $\int_2^{3.5} (2^{x+1} - 4x) \, dx$

(3)

T12 Integration

12_SHIFT Transformed
estimates12_SHIFT_HV02 Do not round
the known estimate too early.

WORKED SOLUTION

Question 6

WMA12/01 May/June 2024

WMA12-01_2024_MayJune_Q06

1

Sketch key featuresFor $y = a^x + 4$ with $a > 1$, the curve is increasing.

2

Find the y-interceptAt $x = 0$, $y = a^0 + 4 = 5$, so the y -intercept is $(0, 5)$.

3

Find the asymptoteAs $x \rightarrow -\infty$, $a^x \rightarrow 0$, so the horizontal asymptote is $y = 4$.

4

Use the trapezium widthThe x values increase by 0.3, so $h = 0.3$.

5

Apply the trapezium rule

$$\int_2^{3.5} (2^x - 2x) dx \approx \frac{0.3}{2} [0 + 4.3137 + 2(0.3246 + 0.8629 + 1.6643 + 2.7896)].$$

6

Round part b

This gives 2.339475, so the estimate is 2.34 to 2 decimal places.

7

Estimate part c i

$$2^x + 2x = (2^x - 2x) + 4x, \text{ so } \int_2^{3.5} (2^x + 2x) dx \approx 2.34 + \int_2^{3.5} 4x dx.$$

8

Add the extra integral

$$\int_2^{3.5} 4x dx = [2x^2]_2^{3.5} = 16.5, \text{ so the estimate is } 2.34 + 16.5 = 18.84.$$

9

Estimate part c ii

$$2^{x+1} - 4x = 2(2^x - 2x), \text{ so the estimate is } 2(2.34) = 4.68.$$

FINAL ANSWER

WMA12-01_2024_MayJune_Q06

(a) y -intercept $(0, 5)$ and asymptote $y = 4$. (b) 2.34. (c) (i) 18.84. (c) (ii) 4.68.

T12

12_SHIFT

12_SHIFT_HV02

CHAPTER
Integration

FAMILY
Transformed estimates

VARIANT
Do not round the known
estimate too early.

6 marks

WMA12/01 MAY/JUNE 2025

Question 5

Past-paper question

WMA12-01_2025_MayJune_Q05

x	-2	-0.5	1	2.5	4	5.5	7
y	12	4.243	1.5	0.530	0.188	0.066	0.023

The table above shows corresponding values of x and y for

$$y = 3\left(\frac{1}{2}\right)^x$$

The values of y are given to 3 decimal places as appropriate.

- (a) Using the trapezium rule with all the values of y in the given table, obtain an estimate for

$$\int_{-2}^7 3\left(\frac{1}{2}\right)^x dx$$

giving the answer to one decimal place.

(3)

Using the answer to part (a) and making your method clear, estimate

(b) (i) $\int_{-2}^7 3\left(\frac{1}{2}\right)^{x+2} dx$

(ii) $\int_{-2}^7 (2^{-x} + 2x) dx$

(3)

T12 Integration

12_SHIFT Transformed
estimates12_SHIFT_HV02 Do not round
the known estimate too early.

WORKED SOLUTION

Question 5

WMA12/01 May/June 2025

WMA12-01_2025_MayJune_Q05

1

Find the strip widthThe x values increase by 1.5, so $h = 1.5$.

2

Apply the trapezium rule

$$\int_{-2}^7 3 \left(\frac{1}{2}\right)^x dx \approx \frac{1.5}{2} \{12 + 0.023 + 2(4.243 + 1.5 + 0.530 + 0.188 + 0.066)\}.$$

3

Evaluate

This gives 18.80775, which is 18.8 to one decimal place.

4

Use the power shift

$$3 \left(\frac{1}{2}\right)^{x+2} = \frac{1}{4} \cdot 3 \left(\frac{1}{2}\right)^x.$$

5

Estimate part b i

$$\text{So } \int_{-2}^7 3 \left(\frac{1}{2}\right)^{x+2} dx \approx \frac{1}{4} (18.80775) = 4.7019375, \text{ which is } 4.7.$$

6

Rewrite part b ii

$$\text{Since } 3 \left(\frac{1}{2}\right)^x = 3 \cdot 2^{-x}, \text{ we have } 2^{-x} = \frac{1}{3} \cdot 3 \left(\frac{1}{2}\right)^x.$$

7

Estimate part b ii

$$\int_{-2}^7 (2^{-x} + 2x) dx \approx \frac{18.80775}{3} + [x^2]_{-2}^7 = 6.26925 + 45 = 51.26925, \text{ which is } 51.3.$$

FINAL ANSWER

WMA12-01_2025_MayJune_Q05

(a) 18.8. (b)(i) 4.7. (b)(ii) 51.3.

T12

12_SHIFT

12_SHIFT_HV02

CHAPTER
Integration

FAMILY
Transformed estimates

VARIANT
Do not round the known
estimate too early.

7 marks

WMA12/01 OCTOBER 2025
Question 3

Past-paper question

WMA12-01_2025_Oct_Q03

3.




Figure 1

In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.

Figure 1 shows a sketch of the curve with equation $y = 3 \times 2^{-x}$

The point $P(k, 300\,000)$ lies on the curve.

(a) Use logarithms to find the value of k to 2 decimal places.

(2)

x	-0.5	1	2.5	4.0	5.5	7
y	4.243	1.5	0.530	0.188	0.066	0.023

The table shows corresponding values of x and y for $y = 3 \times 2^{-x}$

The values of y are given to 3 decimal places where appropriate.

(b) (i) Use the trapezium rule, with all the values of y from the table, to find an approximate value, to 2 decimal places, for

$$\int_{-0.5}^7 3 \times 2^{-x} \, dx$$

(3)

(ii) Use your answer to part (b)(i) to estimate

$$\int_{-0.5}^7 2^{-x} \, dx + \int_{-7}^{0.5} 2^x \, dx$$

(2)

T12 Integration

12_SHIFT Transformed
estimates12_SHIFT_HV02 Do not round
the known estimate too early.

WORKED SOLUTION

Question 3

WMA12/01 October 2025

WMA12-01_2025_Oct_Q03

1

Use the point on the curve

Since $P(k, 300000)$ lies on $y = 3 \times 2^{-x}$, $300000 = 3 \times 2^{-k}$.

2

Solve with logarithms

$100000 = 2^{-k}$, so $-k = \log_2(100000)$ and $k = -\log_2(100000) = -16.609 \dots$

3

Round k

To 2 decimal places, $k = -16.61$.

4

Find the strip width

The x values increase by 1.5, so $h = 1.5$.

5

Apply the trapezium rule

$$\int_{-0.5}^7 3 \times 2^{-x} dx \approx \frac{1.5}{2} \{4.243 + 0.023 + 2(1.5 + 0.530 + 0.188 + 0.066)\}.$$

6

Evaluate the estimate

This gives 6.6255, which is 6.63 to 2 decimal places.

7

Relate the first new integral

$$\int_{-0.5}^7 2^{-x} dx = \frac{1}{3} \int_{-0.5}^7 3 \times 2^{-x} dx.$$

8

Relate the second new integral

Using the substitution $u = -x$, $\int_{-7}^{0.5} 2^x dx = \int_{-0.5}^7 2^{-u} du$.

9

Estimate the total

The required estimate is therefore $\frac{2}{3}(6.6255) = 4.417$, which is 4.42 to 2 decimal places.

FINAL ANSWER

WMA12-01_2025_Oct_Q03

(a) $k = -16.61$. (b)(i) 6.63. (b)(ii) 4.42.

T12

12_SHIFT

12_SHIFT_HV02

CHAPTER
Integration

FAMILY
Transformed estimates

VARIANT
Do not round the known
estimate too early.

6 marks

WMA12/01 JANUARY 2026
Question 2

Past-paper question

WMA12-01_2026_Jan_Q02

2.




Figure 1

Figure 1 shows a sketch of part of the curve C with equation $y = 2^{-0.1x}$

The table below shows corresponding values of x and y for $y = 2^{-0.1x}$

The y values are given to 4 decimal places.

x	-4	-1	2	5	8	11
y	1.3195	1.0718	0.8706	0.7071	0.5743	0.4665

The region R , shown shaded in Figure 1, is bounded by C , the x -axis and the lines with equations $x = -4$ and $x = 11$

- (a) Use the trapezium rule with all the values of y in the table to find an estimate for the area of R . Give the answer to 2 decimal places.

(3)

- (b) State how you would use the trapezium rule to get a more accurate estimate for the true area of R .

(1)

Using the answer to part (a) and showing your working,

- (c) estimate the value of

$$\int_{-4}^{11} 2^{3-0.1x} dx$$

(2)

T12 Integration

12_SHIFT Transformed
estimates12_SHIFT_HV02 Do not round
the known estimate too early.

WORKED SOLUTION

Question 2

WMA12/01 January 2026

WMA12-01_2026_Jan_Q02

1

Find the strip widthThe x values increase by 3, so $h = 3$.

2

Apply the trapezium rule

$$A \approx \frac{3}{2} \{1.3195 + 0.4665 + 2(1.0718 + 0.8706 + 0.7071 + 0.5743)\}.$$

3

Calculate the estimated area

$$A \approx \frac{3}{2} (8.2336) = 12.3504, \text{ so the area is } 12.35 \text{ to 2 decimal places.}$$

4

Improve the estimate

A more accurate estimate would use more trapezia, so the strip width is smaller.

5

Use the scale factor in the integralSince $2^{3-0.1x} = 8 \cdot 2^{-0.1x}$, the required estimate is $8(12.35) = 98.8$.**FINAL ANSWER**

WMA12-01_2026_Jan_Q02

(a) 12.35. (b) Use narrower strips, or more trapezia. (c) 98.8.

TOPIC 12 | FAMILY 04

Numerical Integration

WHAT THIS FAMILY ASKS

Recognise the numerical integration demand and complete the required reasoning in a clear, exam-ready order.

VARIANT ROUTE

V01 core family move

10 classified questions

69 total marks

T12

12_TRAP

12_TRAP_CORE

CHAPTER
Integration

FAMILY
Numerical integration

VARIANT
core family move

7 marks

WMA12/01 OCTOBER 2019

Question 5

Past-paper question

WMA12-01_2019_Oct_Q05

5. (a) Given $0 < a < 1$, sketch the curve with equation

$$y = a^x$$

showing the coordinates of the point at which the curve crosses the y-axis.

(2)

x	2	2.5	3	3.5	4
y	4.25	6.427	9.125	12.34	16.06

The table above shows corresponding values of x and y for $y = x^2 + \left(\frac{1}{2}\right)^x$

The values of y are given to 4 significant figures as appropriate.

Using the trapezium rule with all the values of y in the given table,

- (b) obtain an estimate for $\int_2^4 \left(x^2 + \left(\frac{1}{2}\right)^x \right) dx$

(3)

Using your answer to part (b) and making your method clear, estimate

- (c) $\int_2^4 \left(x(x-3) + \left(\frac{1}{2}\right)^x \right) dx$

(2)

T12 Integration

12_TRAP Numerical
integration12_TRAP_CORE core family
move

WORKED SOLUTION

Question 5

WMA12/01 October 2019

WMA12-01_2019_Oct_Q05

1

Describe the sketch

For $0 < a < 1$, $y = a^x$ is a decreasing exponential curve. At $x = 0$, $y = a^0 = 1$, so it crosses the y -axis at $(0, 1)$.

2

Find the strip width

The x values increase by 0.5, so $h = 0.5$.

3

Apply the trapezium rule

$$A \approx \frac{0.5}{2} \{4.25 + 16.06 + 2(6.427 + 9.125 + 12.34)\}.$$

4

Calculate the estimate

$A \approx 0.25(76.094) = 19.0235$, so the estimate is 19.0 to 3 significant figures.

5

Relate the two integrands

$$x(x - 3) + \left(\frac{1}{2}\right)^x = x^2 + \left(\frac{1}{2}\right)^x - 3x.$$

6

Subtract the exact integral of $3x$

$$\int_2^4 3x \, dx = \left[\frac{3}{2}x^2\right]_2^4 = 24 - 6 = 18.$$

7

Estimate the new integral

Using the estimate from part (b), the required value is $19.0235 - 18 = 1.0235$, so the estimate is 1.0.

FINAL ANSWER

WMA12-01_2019_Oct_Q05

(a) A decreasing exponential curve crossing the y -axis at $(0, 1)$.

(b) 19.0. (c) 1.0.

T12

12_TRAP

12_TRAP_CORE

CHAPTER
Integration

FAMILY
Numerical integration

VARIANT
core family move

7 marks

WMA12/01 JANUARY 2020

Question 1

Past-paper question

WMA12-01_2020_Jan_Q01

1. The table below shows corresponding values of x and y for $y = \log_2(2x)$

The values of y are given to 2 decimal places as appropriate.

x	2	5	8	11	14
y	2	3.32	4	4.46	4.81

Using the trapezium rule with all the values of y in the given table,

- (a) obtain an estimate for $\int_2^{14} \log_2(2x) dx$, giving your answer to one decimal place. (3)

Using your answer to part (a) and making your method clear, estimate

- (b) (i) $\int_2^{14} \frac{\log_2(4x^2)}{5} dx$
 (ii) $\int_2^{14} \log_2\left(\frac{2}{x}\right) dx$ (4)

T12 Integration

12_TRAP Numerical
integration12_TRAP_CORE core family
move

WORKED SOLUTION

Question 1

WMA12/01 January 2020

WMA12-01_2020_Jan_Q01

1

Use the trapezium rule

The strip width is $h = 3$. Therefore $\int_2^{14} \log_2(2x) dx \approx \frac{3}{2}\{2 + 4.81 + 2(3.32 + 4 + 4.46)\}$.

2

Calculate part (a)

The estimate is $\frac{3}{2}(30.37) = 45.555$, so to one decimal place the answer is 45.6.

3

Rewrite part (b)(i)

$\log_2(4x^2) = \log_2\{(2x)^2\} = 2\log_2(2x)$, so $\int_2^{14} \frac{\log_2(4x^2)}{5} dx = \frac{2}{5} \int_2^{14} \log_2(2x) dx$.

4

Estimate part (b)(i)

Using part (a), the estimate is $\frac{2}{5}(45.555) = 18.222$, so 18.2 to one decimal place.

5

Rewrite part (b)(ii)

$\log_2\left(\frac{2}{x}\right) = 2 - \log_2(2x)$.

6

Estimate part (b)(ii)

$\int_2^{14} \log_2\left(\frac{2}{x}\right) dx \approx \int_2^{14} 2 dx - 45.555 = 24 - 45.555 = -21.555$, so -21.6.

FINAL ANSWER

WMA12-01_2020_Jan_Q01

(a) 45.6. (b)(i) 18.2. (b)(ii) -21.6.

T12

12_TRAP

12_TRAP_CORE

CHAPTER
Integration

FAMILY
Numerical integration

VARIANT
core family move

6 marks

WMA12/01 JANUARY 2021

Question 7

Past-paper question

WMA12-01_2021_Jan_Q07

7.




Figure 1

Solar panels are installed on the roof of a building.

The power, P , produced on a particular day, in kW, can be modelled by the equation

$$P = 0.95 + 2^{t-12} + 2^{12-t} - (t-12)^2 \quad 8.5 \leq t \leq 15.2$$

where t is the time in hours after midnight. The graph of P against t is shown in Figure 1.

A table of values of t and P is shown below, with the values of P given to 4 significant figures where appropriate.

Time, t (hours)	10	10.5	11	11.5	12
Power, P (kW)		1.882	2.45		2.95

- (a) Use the given equation to complete the table, giving the values of P to 4 significant figures where appropriate.

(2)

The amount of energy, in kWh, produced between 10:00 and 12:00 can be found by calculating the area of region R , shown shaded in Figure 1.

- (b) Use the trapezium rule, with all the values of P in the completed table, to find an estimate for the amount of energy produced between 10:00 and 12:00. Give your answer to 2 decimal places.

(4)

T12 Integration

12_TRAP Numerical
integration12_TRAP_CORE core family
move

WORKED SOLUTION

Question 7

WMA12/01 January 2021

WMA12-01_2021_Jan_Q07

1

Use the model at t equals ten

$$P(10) = 0.95 + 2^{-2} + 2^2 - (10 - 12)^2 = 0.95 + 0.25 + 4 - 4 = 1.20.$$

2

Use the model at t equals eleven point five

$$P(11.5) = 0.95 + 2^{-0.5} + 2^{0.5} - (11.5 - 12)^2 = 2.821 \dots$$

3

Identify the trapezium width

The time values increase by 0.5, so $h = 0.5$.

4

Apply the trapezium rule

The estimate is $\frac{0.5}{2} \{1.20 + 2.95 + 2(1.882 + 2.45 + 2.821 \dots)\}$.

5

Calculate the energy

This gives 4.614 ..., so the estimated energy is 4.61 kWh to 2 decimal places.

FINAL ANSWER

WMA12-01_2021_Jan_Q07

(a) $P(10) = 1.20$, $P(11.5) = 2.821$. (b) 4.61 kWh.

T12

12_TRAP

12_TRAP_CORE

CHAPTER
Integration

FAMILY
Numerical integration

VARIANT
core family move

10 marks

WMA12/01 MAY/JUNE 2021

Question 5

Past-paper question

WMA12-01_2021_MayJune_Q05

5.




Figure 2

Figure 2 shows a sketch of part of the graph of the curves C_1 and C_2

The curves intersect when $x = 2.5$ and when $x = 4$

A table of values for some points on the curve C_1 is shown below, with y values given to 3 decimal places as appropriate.

x	2.5	2.75	3	3.25	3.5	3.75	4
y	5.453	7.764	9.375	9.964	9.367	7.626	5

Using the trapezium rule with all the values of y in the table,

- (a) find, to 2 decimal places, an estimate for the area bounded by the curve C_1 , the line with equation $x = 2.5$, the x -axis and the line with equation $x = 4$

(4)

The curve C_2 has equation

$$y = x^{\frac{3}{2}} - 3x + 9 \quad x > 0$$

- (b) Find $\int \left(x^{\frac{3}{2}} - 3x + 9 \right) dx$

(3)

The region R , shown shaded in Figure 2, is bounded by the curves C_1 and C_2

- (c) Use the answers to part (a) and part (b) to find, to one decimal place, an estimate for the area of the region R .

(3)

T12 Integration

12_TRAP Numerical
integration12_TRAP_CORE core family
move

WORKED SOLUTION

Question 5

WMA12/01 May/June 2021

WMA12-01_2021_MayJune_Q05

1

Identify the strip widthThe x values increase by 0.25, so $h = 0.25$.

2

Apply the trapezium ruleThe area under C_1 is approximately $\frac{0.25}{2}\{5.453 + 5 + 2(7.764 + 9.375 + 9.964 + 9.367 + 7.626)\}$.

3

Calculate the estimate

This gives 12.330625, so the estimate is 12.33 to 2 decimal places.

4

Integrate C two

$$\int (x^{3/2} - 3x + 9) dx = \frac{2}{5}x^{5/2} - \frac{3}{2}x^2 + 9x + c.$$

5

Find the area under C twoBetween $x = 2.5$ and $x = 4$, the exact integral of C_2 is $\left[\frac{2}{5}x^{5/2} - \frac{3}{2}x^2 + 9x\right]_{2.5}^4 = 7.722 \dots$

6

Find the shaded areaThe shaded region is between C_1 and C_2 , so its area is approximately $12.330625 - 7.722 \dots = 4.608 \dots$

7

Round the resultTo one decimal place, the area of region R is 4.6.**FINAL ANSWER**

WMA12-01_2021_MayJune_Q05

(a) 12.33. (b) $\frac{2}{5}x^{5/2} - \frac{3}{2}x^2 + 9x + c$. (c) 4.6.

T12

12_TRAP

12_TRAP_CORE

CHAPTER
Integration

FAMILY
Numerical integration

VARIANT
core family move

8 marks

WMA12/01 OCTOBER 2021
Question 3

Past-paper question

WMA12-01_2021_Oct_Q03

3.




Figure 1

Figure 1 shows a sketch of the curve with equation $y = \log_{10} x$

The region R , shown shaded in Figure 1, is bounded by the curve, the line with equation $x = 2$, the x -axis and the line with equation $x = 14$

Using the trapezium rule with four strips of equal width,

(a) show that the area of R is approximately 10.10 (3)

(b) Explain how the trapezium rule could be used to obtain a more accurate estimate for the area of R . (1)

(c) Using the answer to part (a) and making your method clear, estimate the value of

(i) $\int_2^{14} \log_{10} \sqrt{x} \, dx$

(ii) $\int_2^{14} \log_{10} 100x^3 \, dx$

(4)

T12 Integration

12_TRAP Numerical
integration12_TRAP_CORE core family
move

WORKED SOLUTION

Question 3

WMA12/01 October 2021

WMA12-01_2021_Oct_Q03

1

Find the strip widthFour equal strips from $x = 2$ to $x = 14$ have width $h = \frac{14-2}{4} = 3$.

2

List the ordinatesUse $x = 2, 5, 8, 11, 14$ and $y = \log_{10} x$.

3

Apply the trapezium rule

$$A \approx \frac{3}{2} \{ \log_{10} 2 + \log_{10} 14 + 2(\log_{10} 5 + \log_{10} 8 + \log_{10} 11) \} = 10.10 \dots$$

4

Improve the estimate

Use more strips of smaller width to make the trapezium rule estimate more accurate.

5

Estimate the square-root integral

$$\log_{10} \sqrt{x} = \frac{1}{2} \log_{10} x, \text{ so } \int_2^{14} \log_{10} \sqrt{x} dx \approx \frac{1}{2} (10.10) = 5.05.$$

6

Rewrite the second logarithm

$$\log_{10}(100x^3) = \log_{10} 100 + \log_{10} x^3 = 2 + 3 \log_{10} x.$$

7

Estimate the second integral

$$\int_2^{14} (2 + 3 \log_{10} x) dx \approx 2(12) + 3(10.10) = 54.30.$$

FINAL ANSWER

WMA12-01_2021_Oct_Q03

(a) 10.10. (b) Use more strips. (c)(i) 5.05. (c)(ii) 54.30.

T12

12_TRAP

12_TRAP_CORE

CHAPTER
Integration

FAMILY
Numerical integration

VARIANT
core family move

7 marks

WMA12/01 JANUARY 2022

Question 1

Past-paper question

WMA12-01_2022_Jan_Q01

1. The table below shows corresponding values of x and y for

$$y = 2^{5-\sqrt{x}}$$

The values of y are given to 3 decimal places.

x	5	5.5	6	6.5	7
y	6.792	6.298	5.858	5.466	5.113

Using the trapezium rule with all the values of y in the given table,

- (a) obtain an estimate for

$$\int_5^7 2^{5-\sqrt{x}} dx$$

giving your answer to 2 decimal places.

(3)

- (b) Using your answer to part (a) and making your method clear, estimate

(i) $\int_5^7 2^{6-\sqrt{x}} dx$

(ii) $\int_5^7 (3 + 2^{5-\sqrt{x}}) dx$

(4)

T12 Integration

12_TRAP Numerical
integration12_TRAP_CORE core family
move

WORKED SOLUTION

Question 1

WMA12/01 January 2022

WMA12-01_2022_Jan_Q01

1

Identify the strip widthThe x values increase by 0.5, so $h = 0.5$.

2

Apply the trapezium rule

$$\int_5^7 2^{5-\sqrt{x}} dx \approx \frac{0.5}{2} \{6.792 + 5.113 + 2(6.298 + 5.858 + 5.466)\}.$$

3

Calculate the estimate

This gives 11.78725, so the estimate is 11.79 to 2 decimal places.

4

Use the factor of two

$$2^{6-\sqrt{x}} = 2 \cdot 2^{5-\sqrt{x}}, \text{ so } \int_5^7 2^{6-\sqrt{x}} dx \approx 2(11.78725) = 23.5745.$$

5

Estimate part i

This is 23.57 to 2 decimal places.

6

Add the constant integral

$$\int_5^7 (3 + 2^{5-\sqrt{x}}) dx = \int_5^7 3 dx + \int_5^7 2^{5-\sqrt{x}} dx.$$

7

Estimate part iiThe estimate is $3(7 - 5) + 11.78725 = 17.78725$, so 17.79 to 2 decimal places.**FINAL ANSWER**

WMA12-01_2022_Jan_Q01

(a) 11.79. (b)(i) 23.57. (b)(ii) 17.79.

T12

12_TRAP

12_TRAP_CORE

CHAPTER
Integration

FAMILY
Numerical integration

VARIANT
core family move

8 marks

WMA12/01 MAY/JUNE 2022

Question 2

Past-paper question

WMA12-01_2022_MayJune_Q02

2.




Figure 1

Figure 1 shows the graph of

$$y = 1 - \log_{10}(\sin x) \quad 0 < x < \pi$$

where x is in radians.

The table below shows some values of x and y for this graph, with values of y given to 3 decimal places.

x	0.5	1	1.5	2	2.5	3
y	1.319		1.001		1.223	1.850

(a) Complete the table above, giving values of y to 3 decimal places. (2)

(b) Use the trapezium rule with all the y values in the completed table to find, to 2 decimal places, an estimate for

$$\int_{0.5}^3 (1 - \log_{10}(\sin x)) dx \quad (3)$$

(c) Use your answer to part (b) to find an estimate for

$$\int_{0.5}^3 (3 + \log_{10}(\sin x)) dx \quad (3)$$

T12 Integration

12_TRAP Numerical
integration12_TRAP_CORE core family
move

WORKED SOLUTION

Question 2

WMA12/01 May/June 2022

WMA12-01_2022_MayJune_Q02

1

Complete the missing table valuesUsing $y = 1 - \log_{10}(\sin x)$ gives $y(1) = 1.075$ and $y(2) = 1.041$ to 3 decimal places.

2

Identify the strip widthThe x values increase by 0.5, so $h = 0.5$.

3

Apply the trapezium rule

$$A \approx \frac{0.5}{2} \{1.319 + 1.850 + 2(1.075 + 1.001 + 1.041 + 1.223)\}.$$

4

Calculate part b

This gives 2.9625 ..., so the estimate is 2.96 to 2 decimal places.

5

Rewrite the new integrandSince $y = 1 - \log_{10}(\sin x)$, we have $\log_{10}(\sin x) = 1 - y$.

6

Use the interval length $3 + \log_{10}(\sin x) = 4 - y$, and the interval from 0.5 to 3 has length 2.5.

7

Estimate part c

$$\int_{0.5}^3 (3 + \log_{10}(\sin x)) dx \approx 4(2.5) - 2.9625 \dots = 7.0374 \dots, \text{ so } 7.04.$$

FINAL ANSWER

WMA12-01_2022_MayJune_Q02

(a) $y(1) = 1.075$, $y(2) = 1.041$. (b) 2.96. (c) 7.04.

T12

12_TRAP

12_TRAP_CORE

CHAPTER
IntegrationFAMILY
Numerical integrationVARIANT
core family move

7 marks

WMA12/01 OCTOBER 2022

Question 6

Past-paper question

WMA12-01_2022_Oct_Q06

6. The curve C_1 has equation $y = f(x)$.

A table of values of x and y for $y = f(x)$ is shown below, with the y values rounded to 4 decimal places where appropriate.

x	0	0.5	1	1.5	2
y	3	2.6833	2.4	2.1466	1.92

- (a) Use the trapezium rule with all the values of y in the table to find an approximation for

$$\int_0^2 f(x) \, dx$$

giving your answer to 3 decimal places.

(3)




Figure 1




Figure 2

The region R , shown shaded in Figure 1, is bounded by

- the curve C_1
- the curve C_2 with equation $y = 2 - \frac{1}{4}x^2$
- the line with equation $x = 2$
- the y -axis

The region R forms part of the design for a logo shown in Figure 2.

The design consists of the shaded region R inside a rectangle of width 2 and height 3

Using calculus and the answer to part (a),

- (b) calculate an estimate for the percentage of the logo which is shaded.

(4)

T12 Integration

12_TRAP Numerical
integration12_TRAP_CORE core family
move

WORKED SOLUTION

Question 6

WMA12/01 October 2022

WMA12-01_2022_Oct_Q06

1

Identify the strip width

The x values increase by 0.5, so the trapezium-rule width is $h = 0.5$.

2

Apply the trapezium rule

$$\int_0^2 f(x) dx \approx \frac{0.5}{2} [3 + 1.92 + 2(2.6833 + 2.4 + 2.1466)].$$

3

Evaluate the estimate

This gives 4.84495, so the approximation is 4.845 to 3 decimal places.

4

Find the area under C2

$$\int_0^2 \left(2 - \frac{1}{4}x^2\right) dx = \left[2x - \frac{x^3}{12}\right]_0^2 = 4 - \frac{2}{3} = \frac{10}{3}.$$

5

Estimate the shaded area

The shaded region lies between C_1 and C_2 , so its estimated area is $4.845 - \frac{10}{3} = 1.5116\dots$

6

Compare with the rectangle

The rectangle has area $2 \times 3 = 6$.

7

Convert to a percentage

Percentage shaded = $\frac{1.5116\dots}{6} \times 100 = 25.19\dots\%$, so the estimate is 25.2%.

FINAL ANSWER

(a) 4.845. (b) 25.2%.

WMA12-01_2022_Oct_Q06

T12

12_TRAP

12_TRAP_CORE

CHAPTER
Integration

FAMILY
Numerical integration

VARIANT
core family move

3 marks

WMA12/01 MAY/JUNE 2023

Question 1

Past-paper question

WMA12-01_2023_MayJune_Q01

1. The continuous curve C has equation $y = f(x)$.

A table of values of x and y for $y = f(x)$ is shown below.

x	4.0	4.2	4.4	4.6	4.8	5.0
y	9.2	8.4556	3.8512	5.0342	7.8297	8.6

Use the trapezium rule with all the values of y in the table to find an approximation for

$$\int_4^5 f(x) \, dx$$

giving your answer to 3 decimal places.

(3)

T12 Integration

12_TRAP Numerical
integration12_TRAP_CORE core family
move

WORKED SOLUTION

Question 1

WMA12/01 May/June 2023

WMA12-01_2023_MayJune_Q01

1

Identify the interval width

The x values go from 4.0 to 5.0 in steps of 0.2, so $h = 0.2$.

2

Apply the trapezium rule

$$\int_4^5 f(x) dx \approx \frac{0.2}{2} [9.2 + 8.6 + 2(8.4556 + 3.8512 + 5.0342 + 7.8297)].$$

3

Evaluate

The value is 6.81414, so the approximation is 6.814 to 3 decimal places.

FINAL ANSWER

6.814.

WMA12-01_2023_MayJune_Q01

T12

12_TRAP

12_TRAP_CORE

CHAPTER
Integration

FAMILY
Numerical integration

VARIANT
core family move

6 marks

WMA12/01 OCTOBER 2023

Question 6

Past-paper question

WMA12-01_2023_Oct_Q06




Figure 1

A river is being studied.

At one particular place, the river is 15 m wide.

The depth, y metres, of the river is measured at a point x metres from one side of the river.

Figure 1 shows a plot of the cross-section of the river and the coordinate values (x, y)

- (a) Use the trapezium rule with all the y values given in Figure 1 to estimate the cross-sectional area of the river.

(3)

The water in the river is modelled as flowing at a constant speed of 1.5 m s^{-1} across the whole of the cross-section.

- (b) Use the model and the answer to part (a) to estimate the volume of water flowing through this section of the river each minute, giving your answer in m^3 to 2 significant figures.

(2)

Assuming the model,

- (c) state, giving a reason for your answer, whether your answer for part (b) is an overestimate or an underestimate of the true volume of water flowing through this section of the river each minute.

(1)

T12 Integration

12_TRAP Numerical
integration12_TRAP_CORE core family
move

WORKED SOLUTION

Question 6

WMA12/01 October 2023

WMA12-01_2023_Oct_Q06

1

Read the strip width

The x values are 0, 3, 6, 9, 12, 15, so the trapezium width is $h = 3$.

2

Apply the trapezium rule

Area $\approx \frac{3}{2} [0 + 0 + 2(1.52 + 2.74 + 3.12 + 3.08)]$.

3

Find the cross-sectional area

This gives 31.38 m^2 .

4

Convert flow speed to one minute

The speed is 1.5 m s^{-1} , so in one minute the water travels $1.5 \times 60 = 90 \text{ m}$.

5

Estimate volume

Volume per minute $\approx 31.38 \times 90 = 2824.2 \text{ m}^3$.

6

Round the volume

To 2 significant figures, this is $2.8 \times 10^3 \text{ m}^3$.

7

State over or under

It is an underestimate, because the curve between the plotted points lies below the straight-line trapezium chords in the diagram, so the true cross-sectional area is larger.

FINAL ANSWER

WMA12-01_2023_Oct_Q06

(a) 31.38 m^2 . (b) $2.8 \times 10^3 \text{ m}^3$. (c) **Underestimate.**