

Concept 4 | Solving Equations and Inequalities Involving Trigonometric Functions

English Student Edition • Focused formulas and guided practice

Formula opener

Sine

$$\sin x = \sin \alpha$$

Cosine

$$\cos x = \cos \alpha$$

Tangent

$$\tan x = \tan \alpha$$

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Q001 Written

Solve (a) $-2 \cos \theta = \sqrt{2}$ and (b) $\tan \theta + \sqrt{3} = 0$, giving the general solution for (b).

Q002 Written

Solve (a) $2 \cos \theta - 1 = 0$ and (b) $\tan \theta - 1 = 0$ for $0 \leq \theta \leq 2\pi$.

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Q003 Written

Solve all given equations on $0 \leq \theta \leq 2\pi$: (a) $2 \sin \theta + 1 = 0$, (b) $2 \sin \theta = \sqrt{3}$, (c) $3 \sin \theta + 3 = 0$, (d) $\sqrt{2} \cos \theta + 1 = 0$, (e) $2 \cos \theta = \sqrt{3}$, (f) $2 \cos \theta - \sqrt{2} = 0$, (g) $\tan \theta + 1 = 0$, (h) $\sqrt{3} \tan \theta = 1$, (i) $\sqrt{3} \tan \theta = 3$, **and give general solutions for** (j) $2 \sin \theta + \sqrt{3} = 0$, (k) $\cos \theta + 1 = 0$, (l) $\sqrt{3} \tan \theta + 3 = 0$.

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Q004 **Written**

Solve $\tan^2 \theta - (2 + \sqrt{3}) \tan \theta + 2\sqrt{3} = 0$ and give the general solution.

Q005 Written

Solve for $0 \leq \theta < 2\pi$: (a) $2 \cos \theta > -\sqrt{3}$, (b) $\tan \theta \leq \frac{1}{\sqrt{3}}$.

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Q006 Written

Solve for $0 \leq \theta < 2\pi$: (a) $2 \sin \theta \leq -1$, (b) $\cos \theta > \frac{1}{2}$, (c) $\tan \theta > 1$.

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Q007 Written

Solve all given inequalities for $0 \leq \theta < 2\pi$: (a) $\sin \theta > \frac{\sqrt{3}}{2}$, (b) $\sin \theta - \frac{\sqrt{2}}{2} > 0$, (c) $2 \sin \theta + \sqrt{3} \leq 0$, (d) $\cos \theta > 0$, (e) $\cos \theta - \frac{1}{2} \leq 0$, (f) $2 \cos \theta + \sqrt{2} \leq 0$, (g) $\tan \theta \leq -\frac{1}{\sqrt{3}}$, (h) $\tan \theta + \sqrt{3} > 0$, (i) $\tan \theta + 1 \geq 0$.

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Q008 Written

Solve $2 \sin^2 \theta - \cos \theta - 1 > 0$ **for** $0 \leq \theta < 2\pi$.

Q009 Written

Solve for $0 \leq \theta < 2\pi$: (a) $2 \cos^2 \theta - 3 \sin \theta = 0$, (b) $2 \sin^2 \theta - 3 \cos \theta - 3 \leq 0$.

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Q010 Written

Solve for $0 \leq \theta < 2\pi$: (a) $2 \sin^2 \theta - \sin \theta = 0$, (b) $2 \cos^2 \theta - \sin \theta = 2$, (c) $2 \sin^2 \theta - 4 < 5 \cos \theta$.

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Q011 Written

Solve all given equations and inequalities for $0 \leq \theta < 2\pi$: (a) $2 \sin^2 \theta + \sin \theta - 1 = 0$, (b) $2 \sin^2 \theta + 3 \cos \theta = 0$, (c) $2 \sin^2 \theta - \cos \theta = 2$, (d) $2 \cos^2 \theta - 5 \cos \theta - 3 = 0$, (e) $2 \cos^2 \theta + \sin \theta - 1 = 0$, (f) $2 \cos^2 \theta + 3 \sin \theta = 3$, (g) $2 \cos^2 \theta + 3 \sin \theta \leq 0$, (h) $2 \sin^2 \theta + \cos \theta > 1$, (i) $2 \cos^2 \theta - 4 \leq 5 \sin \theta$.

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Q012 Written

Solve $\frac{1-2\sin^2\theta+\sin\theta}{\cos^2\theta} \geq 1$ for $0 \leq \theta < 2\pi$.

Q013 Written

Solve for $0 \leq \theta < 2\pi$: (a) $\sin(\theta + \frac{\pi}{3}) = \frac{1}{2}$, (b) $\sin(\theta + \frac{\pi}{6}) > -\frac{1}{2}$.

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Q014 Written

Solve for $0 \leq \theta < 2\pi$: (a) $\sin(\theta + \frac{\pi}{6}) = \frac{1}{2}$, (b) $\tan(\theta + \frac{\pi}{4}) = \sqrt{3}$, (c) $\cos(\theta + \frac{\pi}{4}) < -\frac{\sqrt{2}}{2}$.

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Q015 Written

Solve for $0 \leq \theta < 2\pi$: (a) $\cos(\theta + \frac{\pi}{3}) = -\frac{\sqrt{3}}{2}$, (b) $\cos(\theta + \frac{\pi}{6}) = -\frac{1}{2}$, (c) $\sin(\theta + \frac{\pi}{4}) = \frac{\sqrt{3}}{2}$, (d) $\tan(\theta + \frac{\pi}{4}) = -1$, (e) $\tan(\theta + \frac{\pi}{3}) = \frac{1}{\sqrt{3}}$, (f) $\tan(\theta + \frac{\pi}{6}) = 1$, (g) $\sin(\theta + \frac{\pi}{6}) \leq \frac{\sqrt{3}}{2}$, (h) $\sin(\theta + \frac{\pi}{4}) \geq \frac{\sqrt{2}}{2}$, (i) $\cos(\theta + \frac{\pi}{3}) > \frac{1}{2}$.

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Q016 Written

Solve $4 \sin^2(\theta + \frac{\pi}{6}) \geq 1$ for $0 \leq \theta < 2\pi$.

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Q017 Written

A student divides $\tan x \times \cos x = \frac{1}{2}$ by $\cos x$, obtaining $\tan x = \frac{1}{2 \cos x}$. Evaluate this strategy and propose a better substitution on $[0^\circ, 360^\circ[$.

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Q018 MCQ

The period used for a general tangent solution is:**Choose the correct option.**A 2π B π C $\pi/2$ D 4π

Q019 MCQ**The solutions of $2 \cos \theta = 1$ on $[0, 2\pi]$ are:****Choose the correct option.****A $\pi/6, 11\pi/6$** **B $\pi/3, 5\pi/3$** **C $2\pi/3, 4\pi/3$** **D $\pi/4, 7\pi/4$**

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Q020 MCQ

Which root is impossible when solving $2 \sin^2 \theta + 3 \sin \theta - 2 = 0$?

Choose the correct option.

A $\sin \theta = 1/2$

B $\sin \theta = -2$

C $\sin \theta = 0$

D $\sin \theta = -1/2$

Q021 MCQ

The quadratic challenge factors into roots:**Choose the correct option.**

- A 1 and $\sqrt{3}$
- B 2 and $\sqrt{3}$
- C 2 and 3
- D $\pi/3$ and 2

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Q022 MCQ

For $2 \cos \theta > -\sqrt{3}$, the excluded arc is:

Choose the correct option.

A $[0, 5\pi/6]$

B $[5\pi/6, 7\pi/6]$

C $[7\pi/6, 2\pi)$

D $(\pi/6, 5\pi/6)$

Q023 MCQ**The inequality $\tan \theta > 1$ on $[0, 2\pi)$ holds on:****Choose the correct option.****A** $(\pi/4, \pi/2) \cup (5\pi/4, 3\pi/2)$ **B** $[0, \pi/4]$ **C** $(\pi/2, \pi)$ **D** $[\pi, 2\pi)$

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Q024 MCQ

A closed endpoint in a trig inequality is used when the sign is:

Choose the correct option.

- A strictly greater
- B strictly less
- C greater than or equal to
- D never

Q025 MCQ**The 5-14 challenge reduces to which cosine condition?****Choose the correct option.**

A $\cos \theta > 1/2$

B $-1 < \cos \theta < 1/2$

C $\cos \theta \leq -1$

D $\cos \theta \geq 1$

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Q026 MCQ

When a quadratic gives $\sin \theta = -2$, we:**Choose the correct option.**

A keep it

B reject it

C square it

D divide by 2

Q027 MCQ

The first step for $2 \cos^2 \theta - 3 \sin \theta = 0$ is to use:**Choose the correct option.**

A $\sin^2 \theta + \cos^2 \theta = 1$

B $\tan \theta = \sin \theta$

C $\cos \theta = 1$

D the tangent period

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Q028 MCQ

The solution of $2 \sin^2 \theta + \sin \theta - 1 = 0$ includes:

Choose the correct option.

A $\theta = \pi/6$

B $\theta = \pi/4$

C $\theta = \pi/3$

D $\theta = 2\pi/3$

Q029 MCQ**In the rational challenge, which angle must be excluded?****Choose the correct option.****A** 0**B** $\pi/4$ **C** $\pi/2$ **D** π

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Q030 MCQ

After setting $t = \theta + \pi/3$ with $0 \leq \theta < 2\pi$, the t-range is:

Choose the correct option.

A $[0, 2\pi)$

B $[\pi/3, 7\pi/3)$

C $[-\pi/3, 5\pi/3)$

D $[\pi, 3\pi)$

Q031 MCQ**The solutions of $\tan(\theta + \pi/4) = \sqrt{3}$ include:****Choose the correct option.**A $\pi/12$ B $\pi/6$ C $\pi/3$ D $\pi/2$

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Q032 MCQ

For $\sin(\theta + \pi/4) \geq \sqrt{2}/2$, the θ -solution is:

Choose the correct option.

A $[0, \pi/2]$

B $(0, \pi/2)$

C $[\pi/2, \pi]$

D $[0, 2\pi)$

Q033 MCQ**The 5-16 challenge is equivalent to:****Choose the correct option.**

A $|\sin(t)| \geq 1/2$

B $\sin(t) = 0$

C $\cos(t) > 1/2$

D $\tan(t) = 1$

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Q034 MCQ

In the Reflect problem, $\tan x \cos x$ simplifies to:**Choose the correct option.**A $\cos(x)$ B $\sin(x)$ C $\tan^2 x$

D 1

Quiz MCQ 1 **Concept Quiz · MCQ**

The general solution of $\tan \theta = -1$ is:

Choose the correct option.

A $-\pi/4 + 2n\pi$

B $-\pi/4 + n\pi$

C $\pi/4 + 2n\pi$

D $\pi/2 + n\pi$

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Quiz MCQ 2 **Concept Quiz · MCQ**

The solution of $\cos \theta > 0$ on $[0, 2\pi)$ is:

Choose the correct option.

A $(\pi/2, 3\pi/2)$

B $[0, \pi/2) \cup (3\pi/2, 2\pi)$

C $[0, \pi)$

D $(\pi, 2\pi)$

Quiz MCQ 3 Concept Quiz · MCQ

The valid sine root of $2 \sin^2 \theta + 3 \sin \theta - 2 = 0$ is:

Choose the correct option.

A $\sin \theta = -2$

B $\sin \theta = 1/2$

C $\sin \theta = 2$

D $\sin \theta = -1/2$

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Quiz MCQ 4 [Concept Quiz · MCQ](#)

The first operation in a shifted-argument problem is to:

Choose the correct option.

- A divide by $\cos \theta$
- B set t equal to the whole argument and find its range
- C change radians to degrees
- D square both sides

Quiz WRITTEN 5 Concept Quiz · Written

Solve $2 \cos \theta = \sqrt{3}$ for $0 \leq \theta < 2\pi$.

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Quiz WRITTEN 6 Concept Quiz · Written**Solve** $\tan \theta \leq -1$ for $0 \leq \theta < 2\pi$.

Quiz WRITTEN 7 Concept Quiz · Written

Solve $2 \sin^2 \theta + \sin \theta - 1 = 0$ **for** $0 \leq \theta < 2\pi$.

Classified Practice

Solving Equations and Inequalities Involving Trigonometric Functions

Quiz WRITTEN 8 Concept Quiz · Written**Solve** $\cos(\theta + \frac{\pi}{3}) > \frac{1}{2}$ **for** $0 \leq \theta < 2\pi$.

Answer key

Use the question number shown on each page.

- Q001** (a) $\theta = \frac{3\pi}{4}, \frac{5\pi}{4}$ (b) $\theta = \frac{2\pi}{3} + n\pi, n \in \mathbb{Z}$ **Q012** $0 \leq \theta < \frac{\pi}{2}$ or $\frac{\pi}{2} < \theta \leq \pi$ **Q020** B
- Q002** (a) $\theta = \frac{\pi}{3}, \frac{5\pi}{3}$ (b) $\theta = \frac{\pi}{4}, \frac{5\pi}{4}$ **Q013** (a) $\theta = \frac{\pi}{2}, \frac{11\pi}{6}$ (b) $0 \leq \theta < \pi$ or $\frac{5\pi}{3} < \theta < 2\pi$ **Q021** B
- Q003** (a) $\frac{3\pi}{4}, \frac{11\pi}{4}$ (b) $\frac{\pi}{2}, \frac{3\pi}{2}$ (c) $\frac{3\pi}{4}, \frac{5\pi}{4}$ (d) $\frac{3\pi}{4}, \frac{5\pi}{4}$ (e) $\frac{\pi}{2}, \frac{3\pi}{2}$ (f) $\frac{\pi}{2}, \frac{3\pi}{2}$ (g) $\frac{\pi}{2}, \frac{3\pi}{2}$ (h) $\frac{\pi}{2}, \frac{3\pi}{2}$ (i) $\frac{\pi}{2}, \frac{3\pi}{2}$ (j) $\frac{\pi}{2}, \frac{3\pi}{2}$ (k) $\frac{\pi}{2}, \frac{3\pi}{2}$ (l) $\frac{\pi}{2}, \frac{3\pi}{2}$ (m) $\frac{\pi}{2}, \frac{3\pi}{2}$ (n) $\frac{\pi}{2}, \frac{3\pi}{2}$ (o) $\frac{\pi}{2}, \frac{3\pi}{2}$ (p) $\frac{\pi}{2}, \frac{3\pi}{2}$ (q) $\frac{\pi}{2}, \frac{3\pi}{2}$ (r) $\frac{\pi}{2}, \frac{3\pi}{2}$ (s) $\frac{\pi}{2}, \frac{3\pi}{2}$ (t) $\frac{\pi}{2}, \frac{3\pi}{2}$ (u) $\frac{\pi}{2}, \frac{3\pi}{2}$ (v) $\frac{\pi}{2}, \frac{3\pi}{2}$ (w) $\frac{\pi}{2}, \frac{3\pi}{2}$ (x) $\frac{\pi}{2}, \frac{3\pi}{2}$ (y) $\frac{\pi}{2}, \frac{3\pi}{2}$ (z) $\frac{\pi}{2}, \frac{3\pi}{2}$ **Q014** (a) $\theta = 0, \frac{2\pi}{3}$ (b) $\theta = \pi, \frac{13\pi}{12}$ (c) $\frac{\pi}{2} < \theta < \pi$ **Q022** B
- Q004** $\tan \theta = 2$ or $\tan \theta = \sqrt{3}$; $\theta = \arctan(2) + n\pi$ or $\theta = \frac{\pi}{3} + n\pi$ **Q015** (a) $\frac{\pi}{2}, \frac{5\pi}{6}$ (b) $\frac{\pi}{2}, \frac{7\pi}{6}$ (c) $\pi, \frac{5\pi}{12}$ (d) $\frac{\pi}{2}, \frac{3\pi}{2}$ (e) $\frac{\pi}{2}, \frac{3\pi}{2}$ (f) $\frac{\pi}{2}, \frac{3\pi}{2}$ (g) $\frac{\pi}{2}, \frac{3\pi}{2}$ (h) $\frac{\pi}{2}, \frac{3\pi}{2}$ (i) $\frac{\pi}{2}, \frac{3\pi}{2}$ (j) $\frac{\pi}{2}, \frac{3\pi}{2}$ (k) $\frac{\pi}{2}, \frac{3\pi}{2}$ (l) $\frac{\pi}{2}, \frac{3\pi}{2}$ (m) $\frac{\pi}{2}, \frac{3\pi}{2}$ (n) $\frac{\pi}{2}, \frac{3\pi}{2}$ (o) $\frac{\pi}{2}, \frac{3\pi}{2}$ (p) $\frac{\pi}{2}, \frac{3\pi}{2}$ (q) $\frac{\pi}{2}, \frac{3\pi}{2}$ (r) $\frac{\pi}{2}, \frac{3\pi}{2}$ (s) $\frac{\pi}{2}, \frac{3\pi}{2}$ (t) $\frac{\pi}{2}, \frac{3\pi}{2}$ (u) $\frac{\pi}{2}, \frac{3\pi}{2}$ (v) $\frac{\pi}{2}, \frac{3\pi}{2}$ (w) $\frac{\pi}{2}, \frac{3\pi}{2}$ (x) $\frac{\pi}{2}, \frac{3\pi}{2}$ (y) $\frac{\pi}{2}, \frac{3\pi}{2}$ (z) $\frac{\pi}{2}, \frac{3\pi}{2}$ **Q023** A
- Q005** (a) $0 \leq \theta < \frac{5\pi}{6}$ or $\frac{7\pi}{6} < \theta < 2\pi$ (b) $0 \leq \theta \leq \frac{\pi}{6}$ or $\frac{\pi}{2} < \theta \leq \frac{7\pi}{6}$ or $\frac{3\pi}{2} < \theta < 2\pi$ **Q016** $0 \leq \theta \leq \frac{2\pi}{3}$ or $\pi \leq \theta \leq \frac{5\pi}{3}$ **Q024** C
- Q006** (a) $\frac{7\pi}{6} \leq \theta \leq \frac{11\pi}{6}$ (b) $0 \leq \theta < \frac{\pi}{3}$ or $\frac{5\pi}{3} < \theta < 2\pi$ (c) $\frac{\pi}{4} < \theta < \frac{\pi}{2}$ or $\frac{5\pi}{4} < \theta < \frac{3\pi}{2}$ **Q017** Dividing by $\cos(x)$ separates the interacting factors and is not helpful, use $\tan(x) \cos(x) = \sin(x)$ (with $\cos(x) \neq 0$), so $\sin(x) = \frac{1}{2}$ and $x = 30^\circ, 150^\circ$. **Q025** B
- Q007** (a) $\frac{\pi}{2}, \frac{3\pi}{2}$ (b) $\frac{\pi}{2}, \frac{3\pi}{2}$ (c) $\frac{\pi}{2}, \frac{3\pi}{2}$ (d) $\frac{\pi}{2}, \frac{3\pi}{2}$ (e) $\frac{\pi}{2}, \frac{3\pi}{2}$ (f) $\frac{\pi}{2}, \frac{3\pi}{2}$ (g) $\frac{\pi}{2}, \frac{3\pi}{2}$ (h) $\frac{\pi}{2}, \frac{3\pi}{2}$ (i) $\frac{\pi}{2}, \frac{3\pi}{2}$ (j) $\frac{\pi}{2}, \frac{3\pi}{2}$ (k) $\frac{\pi}{2}, \frac{3\pi}{2}$ (l) $\frac{\pi}{2}, \frac{3\pi}{2}$ (m) $\frac{\pi}{2}, \frac{3\pi}{2}$ (n) $\frac{\pi}{2}, \frac{3\pi}{2}$ (o) $\frac{\pi}{2}, \frac{3\pi}{2}$ (p) $\frac{\pi}{2}, \frac{3\pi}{2}$ (q) $\frac{\pi}{2}, \frac{3\pi}{2}$ (r) $\frac{\pi}{2}, \frac{3\pi}{2}$ (s) $\frac{\pi}{2}, \frac{3\pi}{2}$ (t) $\frac{\pi}{2}, \frac{3\pi}{2}$ (u) $\frac{\pi}{2}, \frac{3\pi}{2}$ (v) $\frac{\pi}{2}, \frac{3\pi}{2}$ (w) $\frac{\pi}{2}, \frac{3\pi}{2}$ (x) $\frac{\pi}{2}, \frac{3\pi}{2}$ (y) $\frac{\pi}{2}, \frac{3\pi}{2}$ (z) $\frac{\pi}{2}, \frac{3\pi}{2}$ **Q026** B
- Q008** $\frac{\pi}{3} < \theta < \pi$ or $\pi < \theta < \frac{5\pi}{3}$ **Q027** A
- Q009** (a) $\theta = \frac{\pi}{6}, \frac{5\pi}{6}$ (b) $0 \leq \theta \leq \frac{2\pi}{3}$ or $\theta = \pi$ or $\frac{4\pi}{3} \leq \theta < 2\pi$ **Q028** A
- Q010** (a) $\theta = 0, \frac{\pi}{6}, \frac{5\pi}{6}, \pi$ (b) $\theta = 0, \pi, \frac{7\pi}{6}, \frac{11\pi}{6}$ (c) $0 \leq \theta < \frac{2\pi}{3}$ or $\frac{4\pi}{3} < \theta < 2\pi$ **Q029** C
- Q011** (a) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (b) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (c) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (d) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (e) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (f) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (g) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (h) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (i) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (j) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (k) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (l) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (m) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (n) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (o) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (p) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (q) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (r) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (s) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (t) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (u) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (v) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (w) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (x) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (y) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (z) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ **Q018** B
- Q019** B

Q032 A

Q033 A

Q034 B

Quiz MCQ 1 B

Quiz MCQ 2 B

Quiz MCQ 3 B

Quiz MCQ 4 B

Quiz WRITTEN 5 $\theta = \frac{\pi}{6}, \frac{11\pi}{6}$

Quiz WRITTEN 6 $\frac{\pi}{2} < \theta \leq \frac{3\pi}{4}$ or $\frac{3\pi}{2} < \theta \leq \frac{7\pi}{4}$

Quiz WRITTEN 7 $\theta = \frac{\pi}{6}, \frac{5\pi}{6}, \frac{3\pi}{2}$

Quiz WRITTEN 8 $\frac{4\pi}{3} < \theta < 2\pi$