

Concept 4 | Solving Equations and Inequalities Involving Trigonometric Functions

English Student Edition • Focused formulas and guided practice

Formula opener

Sine

$$\sin x = \sin \alpha$$

Cosine

$$\cos x = \cos \alpha$$

Tangent

$$\tan x = \tan \alpha$$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Q001 Written

Solve (a) $-2 \cos \theta = \sqrt{2}$ **and (b)** $\tan \theta + \sqrt{3} = 0$, giving the general solution for (b).

Solution

1. Rewrite as $\cos \theta = -\frac{\sqrt{2}}{2}$ and use QII/QIII.
2. For tangent, the reference angle is $\pi/3$ and the period is π .

Final answer

$$(a) \theta = \frac{3\pi}{4}, \frac{5\pi}{4} \quad (b) \theta = \frac{2\pi}{3} + n\pi, n \in \mathbb{Z}$$

Q002 Written

Solve (a) $2 \cos \theta - 1 = 0$ **and (b)** $\tan \theta - 1 = 0$ for $0 \leq \theta \leq 2\pi$.

Solution

1. Use the unit circle for cosine and include both turns in the stated interval.
2. For tangent, repeat the $\pi/4$ reference angle after one period π .

Final answer

$$(a) \theta = \frac{\pi}{3}, \frac{5\pi}{3} \quad (b) \theta = \frac{\pi}{4}, \frac{5\pi}{4}$$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Q003 Written

Solve all given equations on $0 \leq \theta \leq 2\pi$: (a) $2 \sin \theta + 1 = 0$, (b) $2 \sin \theta = \sqrt{3}$, (c) $3 \sin \theta + 3 = 0$, (d) $\sqrt{2} \cos \theta + 1 = 0$, (e) $2 \cos \theta = \sqrt{3}$, (f) $2 \cos \theta - \sqrt{2} = 0$, (g) $\tan \theta + 1 = 0$, (h) $\sqrt{3} \tan \theta = 1$, (i) $\sqrt{3} \tan \theta = 3$, **and give general solutions for** (j) $2 \sin \theta + \sqrt{3} = 0$, (k) $\cos \theta + 1 = 0$, (l) $\sqrt{3} \tan \theta + 3 = 0$.

Solution

1. Put each equation into a single-function form, find the reference angle, then select the quadrants from the sign.
2. Use period 2π for sine/cosine and π for tangent; reject the endpoint 2π when the interval is half-open.

Final answer

(a) $\frac{7\pi}{6}, \frac{11\pi}{6}$ (b) $\frac{\pi}{3}, \frac{2\pi}{3}$ (c) $\frac{3\pi}{2}$ (d) $\frac{3\pi}{4}, \frac{5\pi}{4}$ (e) $\frac{\pi}{6}, \frac{11\pi}{6}$ (f) $\frac{\pi}{4}, \frac{7\pi}{4}$ (g) $\frac{3\pi}{4}, \frac{7\pi}{4}$ (h) $\frac{\pi}{6}, \frac{7\pi}{6}$ (i) $\frac{\pi}{3}, \frac{4\pi}{3}$ (j) $\frac{4\pi}{3} + 2n\pi, \frac{5\pi}{3} + 2n\pi$ (k) $\pi + 2n\pi$ (l) $\frac{2\pi}{3} + n\pi$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Q004 Written

Solve $\tan^2 \theta - (2 + \sqrt{3}) \tan \theta + 2\sqrt{3} = 0$ **and give the general solution.**

Solution

1. Factor the quadratic as $(\tan \theta - 2)(\tan \theta - \sqrt{3}) = 0$.
2. Apply the tangent period π to both roots.

Final answer

$$\tan \theta = 2 \text{ or } \tan \theta = \sqrt{3}; \quad \theta = \arctan(2) + n\pi \text{ or } \theta = \frac{\pi}{3} + n\pi$$

Q005 Written

Solve for $0 \leq \theta < 2\pi$: **(a)** $2 \cos \theta > -\sqrt{3}$, **(b)** $\tan \theta \leq \frac{1}{\sqrt{3}}$.

Solution

1. Solve the boundary equation first, then shade the correct arcs on the unit circle.
2. Tangent is undefined at odd multiples of $\pi/2$; those points are never included.

Final answer

$$(a) 0 \leq \theta < \frac{5\pi}{6} \text{ or } \frac{7\pi}{6} < \theta < 2\pi \quad (b) 0 \leq \theta \leq \frac{\pi}{6} \text{ or } \frac{\pi}{2} < \theta \leq \frac{7\pi}{6} \text{ or } \frac{3\pi}{2} < \theta < 2\pi$$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Q006 Written

Solve for $0 \leq \theta < 2\pi$: (a) $2 \sin \theta \leq -1$, (b) $\cos \theta > \frac{1}{2}$, (c) $\tan \theta > 1$.

Solution

1. Mark the equality angles and choose the arcs where the function has the required sign.
2. Open endpoints correspond to strict inequalities and to tangent asymptotes.

Final answer

$$(a) \frac{7\pi}{6} \leq \theta \leq \frac{11\pi}{6} \quad (b) 0 \leq \theta < \frac{\pi}{3} \text{ or } \frac{5\pi}{3} < \theta < 2\pi \quad (c) \frac{\pi}{4} < \theta < \frac{\pi}{2} \text{ or } \frac{5\pi}{4} < \theta < \frac{3\pi}{2}$$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Q007 Written

Solve all given inequalities for $0 \leq \theta < 2\pi$: (a) $\sin \theta > \frac{\sqrt{3}}{2}$, (b) $\sin \theta - \frac{\sqrt{2}}{2} > 0$, (c) $2 \sin \theta + \sqrt{3} \leq 0$, (d) $\cos \theta > 0$, (e) $\cos \theta - \frac{1}{2} \leq 0$, (f) $2 \cos \theta + \sqrt{2} \leq 0$, (g) $\tan \theta \leq -\frac{1}{\sqrt{3}}$, (h) $\tan \theta + \sqrt{3} > 0$, (i) $\tan \theta + 1 \geq 0$.

Solution

- For each item, solve the equality, draw the relevant horizontal line, and select the arcs satisfying the sign.
- Keep the distinction between open and closed endpoints and remove all tangent asymptotes.

Final answer

$$(a) \left(\frac{\pi}{3}, \frac{2\pi}{3}\right) (b) \left(\frac{\pi}{4}, \frac{3\pi}{4}\right) (c) \left[\frac{4\pi}{3}, \frac{5\pi}{3}\right] (d) \left[0, \frac{\pi}{2}\right) \cup \left(\frac{3\pi}{2}, 2\pi\right) (e) \left[\frac{\pi}{3}, \frac{5\pi}{3}\right] (f) \left[\frac{3\pi}{4}, \frac{5\pi}{4}\right] (g) \left(\frac{\pi}{2}, \frac{5\pi}{6}\right] \cup \left(\frac{3\pi}{2}, \frac{11\pi}{6}\right] (h) \left[0, \frac{\pi}{2}\right) \cup \left(\frac{2\pi}{3}, \frac{3\pi}{2}\right) \cup \left(\frac{5\pi}{3}, 2\pi\right) (i) \left[0, \frac{\pi}{2}\right) \cup \left[\frac{3\pi}{4}, \frac{3\pi}{2}\right) \cup \left[\frac{7\pi}{4}, 2\pi\right)$$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Q008 Written

Solve $2 \sin^2 \theta - \cos \theta - 1 > 0$ for $0 \leq \theta < 2\pi$.**Solution**

1. Use $\sin^2 \theta = 1 - \cos^2 \theta$ to obtain $(1 - 2 \cos \theta)(1 + \cos \theta) > 0$.
2. Since $-1 \leq \cos \theta \leq 1$, the solution is $-1 < \cos \theta < 1/2$; exclude $\theta = \pi$ where the product is zero.

Final answer

$$\frac{\pi}{3} < \theta < \pi \text{ or } \pi < \theta < \frac{5\pi}{3}$$

Q009 Written

Solve for $0 \leq \theta < 2\pi$: **(a)** $2 \cos^2 \theta - 3 \sin \theta = 0$, **(b)** $2 \sin^2 \theta - 3 \cos \theta - 3 \leq 0$.**Solution**

1. For (a), replace $\cos^2 \theta$ and solve $2s^2 + 3s - 2 = 0$; reject $s = -2$.
2. For (b), replace $\sin^2 \theta$, solve $(2c + 1)(c + 1) \geq 0$, and intersect with $[-1, 1]$.

Final answer

$$(a) \theta = \frac{\pi}{6}, \frac{5\pi}{6} \quad (b) 0 \leq \theta \leq \frac{2\pi}{3} \text{ or } \theta = \pi \text{ or } \frac{4\pi}{3} \leq \theta < 2\pi$$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Q010 Written

Solve for $0 \leq \theta < 2\pi$: (a) $2 \sin^2 \theta - \sin \theta = 0$, (b) $2 \cos^2 \theta - \sin \theta = 2$, (c) $2 \sin^2 \theta - 4 < 5 \cos \theta$.

Solution

1. Set $s = \sin \theta$ or $c = \cos \theta$ after using $s^2 + c^2 = 1$.
2. Solve the resulting quadratic, reject impossible trig values, then read the unit-circle intervals.

Final answer

$$(a) \theta = 0, \frac{\pi}{6}, \frac{5\pi}{6}, \pi \quad (b) \theta = 0, \pi, \frac{7\pi}{6}, \frac{11\pi}{6} \quad (c) 0 \leq \theta < \frac{2\pi}{3} \text{ or } \frac{4\pi}{3} < \theta < 2\pi$$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Q011 Written

Solve all given equations and inequalities for $0 \leq \theta < 2\pi$: (a) $2 \sin^2 \theta + \sin \theta - 1 = 0$, (b) $2 \sin^2 \theta + 3 \cos \theta = 0$, (c) $2 \sin^2 \theta - \cos \theta = 2$, (d) $2 \cos^2 \theta - 5 \cos \theta - 3 = 0$, (e) $2 \cos^2 \theta + \sin \theta - 1 = 0$, (f) $2 \cos^2 \theta + 3 \sin \theta = 3$, (g) $2 \cos^2 \theta + 3 \sin \theta \leq 0$, (h) $2 \sin^2 \theta + \cos \theta > 1$, (i) $2 \cos^2 \theta - 4 \leq 5 \sin \theta$.

Solution

1. Convert to one trig function, solve the quadratic, and apply $-1 \leq \sin \theta, \cos \theta \leq 1$.
2. For inequalities, use the sign chart in the trig variable and then translate the accepted interval(s) to θ .

Final answer

(a) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{3\pi}{2}$ (b) $\frac{2\pi}{3}, \frac{4\pi}{3}$ (c) $\frac{2\pi}{3}, \frac{\pi}{2}, \frac{4\pi}{3}, \frac{3\pi}{2}$ (d) $\frac{2\pi}{3}, \frac{4\pi}{3}$ (e) $\frac{\pi}{2}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (f) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{\pi}{2}$ (g) $\frac{7\pi}{6} \leq \theta \leq \frac{11\pi}{6}$ (h) $0 < \theta < \frac{2\pi}{3}$ or $\frac{4\pi}{3} < \theta < 2\pi$ (i) $0 \leq \theta \leq \frac{7\pi}{6}$ or $\frac{11\pi}{6} \leq \theta < 2\pi$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Q012 Written

Solve $\frac{1-2\sin^2\theta+\sin\theta}{\cos^2\theta} \geq 1$ for $0 \leq \theta < 2\pi$.

Solution

1. Subtract 1 and simplify the numerator to $\sin\theta(1 - \sin\theta)$.
2. The denominator is positive but undefined at $\theta = \pi/2$; hence $0 \leq \sin\theta \leq 1$ with that point removed.

Final answer

$$0 \leq \theta < \frac{\pi}{2} \text{ or } \frac{\pi}{2} < \theta \leq \pi$$

Q013 Written

Solve for $0 \leq \theta < 2\pi$: **(a)** $\sin(\theta + \frac{\pi}{3}) = \frac{1}{2}$, **(b)** $\sin(\theta + \frac{\pi}{6}) > -\frac{1}{2}$.

Solution

1. Let $t = \theta + \alpha$, write the range of t , solve there, then subtract α .
2. For the inequality, shade one complete revolution in the shifted t-range and preserve strict endpoints.

Final answer

$$(a) \theta = \frac{\pi}{2}, \frac{11\pi}{6} \quad (b) 0 \leq \theta < \pi \text{ or } \frac{5\pi}{3} < \theta < 2\pi$$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Q014 Written

Solve for $0 \leq \theta < 2\pi$: (a) $\sin(\theta + \frac{\pi}{6}) = \frac{1}{2}$, (b) $\tan(\theta + \frac{\pi}{4}) = \sqrt{3}$, (c) $\cos(\theta + \frac{\pi}{4}) < -\frac{\sqrt{2}}{2}$.

Solution

1. Find each shifted argument's exact range before selecting the unit-circle arcs.
2. For tangent, exclude asymptotes automatically; for strict cosine inequality, endpoints are open.

Final answer

$$(a) \theta = 0, \frac{2\pi}{3} \quad (b) \theta = \pi, 2\pi, \frac{13\pi}{12} \quad (c) \frac{\pi}{2} < \theta < \pi$$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Q015 Written

Solve for $0 \leq \theta < 2\pi$: (a) $\cos(\theta + \frac{\pi}{3}) = -\frac{\sqrt{3}}{2}$, (b) $\cos(\theta + \frac{\pi}{6}) = -\frac{1}{2}$, (c) $\sin(\theta + \frac{\pi}{4}) = \frac{\sqrt{3}}{2}$, (d) $\tan(\theta + \frac{\pi}{4}) = -1$, (e) $\tan(\theta + \frac{\pi}{3}) = \frac{1}{\sqrt{3}}$, (f) $\tan(\theta + \frac{\pi}{6}) = 1$, (g) $\sin(\theta + \frac{\pi}{6}) \leq \frac{\sqrt{3}}{2}$, (h) $\sin(\theta + \frac{\pi}{4}) \geq \frac{\sqrt{2}}{2}$, (i) $\cos(\theta + \frac{\pi}{3}) > \frac{1}{2}$.

Solution

1. Set $t = \theta + \alpha$, record its interval, solve the standard trig relation in that interval, then subtract α .
2. Check every endpoint against the original strict/non-strict sign and the domain $0 \leq \theta < 2\pi$.

Final answer

(a) $\frac{\pi}{2}, \frac{5\pi}{6}$ (b) $\frac{\pi}{2}, \frac{7\pi}{6}$ (c) $\pi, 2\pi$ (d) $\frac{\pi}{2}, \frac{3\pi}{2}$ (e) $\frac{5\pi}{6}, \frac{11\pi}{6}$ (f) $\pi, 2\pi$ (g) $[0, \frac{\pi}{6}] \cup [\frac{\pi}{2}, 2\pi)$ (h) $[0, \frac{\pi}{2}]$ (i) $\frac{4\pi}{3} < \theta < 2\pi$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Q016 Written

Solve $4 \sin^2(\theta + \frac{\pi}{6}) \geq 1$ for $0 \leq \theta < 2\pi$.**Solution**

1. Let $t = \theta + \pi/6$; then $|\sin(t)| \geq 1/2$.
2. On the shifted range $[\pi/6, 13\pi/6)$, the accepted arcs give the stated closed θ -intervals.

Final answer

$$0 \leq \theta \leq \frac{2\pi}{3} \text{ or } \pi \leq \theta \leq \frac{5\pi}{3}$$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Q017 Written

A student divides $\tan x \times \cos x = \frac{1}{2}$ by $\cos x$, obtaining $\tan x = \frac{1}{2 \cos x}$. Evaluate this strategy and propose a better substitution on $[0^\circ, 360^\circ[$.

Solution

1. Replace $\tan(x)$ by $\sin(x)/\cos(x)$ before cancelling; the original tangent already excludes $\cos(x) = 0$.
2. Solve the resulting sine equation on the stated degree interval.

Final answer

Dividing by $\cos(x)$ separates the interacting factors and is not helpful; use $\tan(x) \cos(x) = \sin(x)$ (with $\cos(x) \neq 0$), so $\sin(x) = \frac{1}{2}$ and $x = 30^\circ, 150^\circ$.

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Q018 MCQ

The period used for a general tangent solution is:**Solution**

1. Tangent repeats after π .

Correct option: B**Final answer** π

Q019 MCQ

The solutions of $2 \cos \theta = 1$ on $[0, 2\pi]$ are:**Solution**

1. Cosine is $1/2$ in QI and QIV.

Correct option: B**Final answer** $\pi/3, 5\pi/3$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Q020 MCQ

Which root is impossible when solving $2 \sin^2 \theta + 3 \sin \theta - 2 = 0$?

Solution

1. A sine value must lie in $[-1, 1]$.

Correct option: B

Final answer

$$\sin \theta = -2$$

Q021 MCQ

The quadratic challenge factors into roots:

Solution

1. Use $(u - 2)(u - \sqrt{3})$ with $u = \tan \theta$.

Correct option: B

Final answer

$$2, \sqrt{3}$$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Q022 MCQ

For $2 \cos \theta > -\sqrt{3}$, the excluded arc is:

Solution

1. Cosine is at most $-\sqrt{3}/2$ on the central QII arc.

Correct option: B

Final answer

$[5\pi/6, 7\pi/6]$

Q023 MCQ

The inequality $\tan \theta > 1$ on $[0, 2\pi)$ holds on:

Solution

1. Use the increasing tangent branches and omit asymptotes.

Correct option: A

Final answer

$(\pi/4, \pi/2) \cup (5\pi/4, 3\pi/2)$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Q024 MCQ

A closed endpoint in a trig inequality is used when the sign is:**Solution**

1. Equality is included for $\leq$ or $\geq$.

Correct option: C**Final answer**

greater than or equal to

Q025 MCQ

The 5-14 challenge reduces to which cosine condition?**Solution**

1. Factor the expression after replacing $\sin^2\theta$.

Correct option: B**Final answer** $-1 < \cos \theta < 1/2$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Q026 MCQ

When a quadratic gives $\sin \theta = -2$, we:**Solution**

1. Sine cannot be outside $[-1, 1]$.

Correct option: B

Final answer

reject it

Q027 MCQ

The first step for $2 \cos^2 \theta - 3 \sin \theta = 0$ is to use:**Solution**

1. Rewrite in one trig function.

Correct option: A

Final answer

$$\sin^2 \theta + \cos^2 \theta = 1$$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Q028 MCQ

The solution of $2 \sin^2 \theta + \sin \theta - 1 = 0$ includes:**Solution**

1. The valid root $\sin \theta = 1/2$ gives $\pi/6, 5\pi/6$.

Correct option: A**Final answer** $\pi/6$

Q029 MCQ

In the rational challenge, which angle must be excluded?**Solution**

1. The denominator $\cos^2 \theta$ is zero there.

Correct option: C**Final answer** $\pi/2$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Q030 MCQ

After setting $t = \theta + \pi/3$ with $0 \leq \theta < 2\pi$, the t-range is:

Solution

1. Add $\pi/3$ to both domain endpoints.

Correct option: B

Final answer

$[\pi/3, 7\pi/3)$

Q031 MCQ

The solutions of $\tan(\theta + \pi/4) = \sqrt{3}$ include:

Solution

1. Set the shifted argument to $\pi/3$.

Correct option: A

Final answer

$\pi/12$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Q032 MCQ

For $\sin(\theta + \pi/4) \geq \sqrt{2}/2$, the θ -solution is:

Solution

1. The shifted sine is at least $\sqrt{2}/2$ from $\pi/4$ to $3\pi/4$.

Correct option: A

Final answer

$$[0, \pi/2]$$

Q033 MCQ

The 5-16 challenge is equivalent to:

Solution

1. Divide by 4 and take the square-root inequality.

Correct option: A

Final answer

$$|\sin t| \geq 1/2$$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Q034 MCQ

In the Reflect problem, $\tan x \cos x$ simplifies to:**Solution**1. Use $\tan(x) = \sin(x)/\cos(x)$ where defined.

Correct option: B

Final answer $\sin x$

Quiz MCQ 1 Concept Quiz · MCQ

The general solution of $\tan \theta = -1$ is:**Solution**1. Tangent has period π .

Correct option: B

Final answer

$$\theta = -\frac{\pi}{4} + n\pi$$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Quiz MCQ 2 Concept Quiz · MCQ

The solution of $\cos \theta > 0$ on $[0, 2\pi)$ is:

Solution

1. Cosine is positive in QI and QIV.

Correct option: B

Final answer

$$[0, \pi/2) \cup (3\pi/2, 2\pi)$$

Quiz MCQ 3 Concept Quiz · MCQ

The valid sine root of $2 \sin^2 \theta + 3 \sin \theta - 2 = 0$ is:

Solution

1. The other algebraic root is outside the sine range.

Correct option: B

Final answer

$$\sin \theta = \frac{1}{2}$$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Quiz MCQ 4 **Concept Quiz · MCQ**

The first operation in a shifted-argument problem is to:

Solution

1. The t-range prevents losing or adding solutions.

Correct option: B

Final answer

set t equal to the whole argument and find its range

Quiz WRITTEN 5 **Concept Quiz · Written**

Solve $2 \cos \theta = \sqrt{3}$ **for** $0 \leq \theta < 2\pi$.

Solution

1. Cosine is $\sqrt{3}/2$ in QI and QIV.

Final answer

$$\theta = \frac{\pi}{6}, \frac{11\pi}{6}$$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Quiz WRITTEN 6 Concept Quiz · Written**Solve** $\tan \theta \leq -1$ for $0 \leq \theta < 2\pi$.**Solution**

1. Use the two tangent branches and include the equality angles.

Final answer

$$\frac{\pi}{2} < \theta \leq \frac{3\pi}{4} \text{ or } \frac{3\pi}{2} < \theta \leq \frac{7\pi}{4}$$

Quiz WRITTEN 7 Concept Quiz · Written**Solve** $2 \sin^2 \theta + \sin \theta - 1 = 0$ for $0 \leq \theta < 2\pi$.**Solution**

1. Factor to $(2 \sin \theta - 1)(\sin \theta + 1) = 0$, then use the unit circle.

Final answer

$$\theta = \frac{\pi}{6}, \frac{5\pi}{6}, \frac{3\pi}{2}$$

Worked Solutions

Solving Equations and Inequalities Involving Trigonometric Functions

Quiz WRITTEN 8 Concept Quiz · Written

Solve $\cos(\theta + \frac{\pi}{3}) > \frac{1}{2}$ **for** $0 \leq \theta < 2\pi$.

Solution

1. Set $t = \theta + \pi/3$, solve $\cos(t) > 1/2$ in $[\pi/3, 7\pi/3)$, then subtract.

Final answer

$$\frac{4\pi}{3} < \theta < 2\pi$$

Answer key

Use the question number shown on each page.

Q001 (a) $\theta = \frac{3\pi}{4}, \frac{5\pi}{4}$ (b) $\theta = \frac{2\pi}{3} + n\pi, n \in \mathbb{Z}$ **Q012** $0 \leq \theta < \frac{\pi}{2}$ or $\frac{\pi}{2} < \theta \leq \pi$ **Q020** B

Q002 (a) $\theta = \frac{\pi}{3}, \frac{5\pi}{3}$ (b) $\theta = \frac{\pi}{4}, \frac{5\pi}{4}$ **Q013** (a) $\theta = \frac{\pi}{2}, \frac{11\pi}{6}$ (b) $0 \leq \theta < \pi$ or $\frac{5\pi}{3} < \theta < \frac{7\pi}{3}$ **Q021** B

Q003 (a) $\frac{3\pi}{4}, \frac{11\pi}{4}$ (b) $\frac{\pi}{2}, \frac{5\pi}{2}$ (c) $\frac{3\pi}{4}, \frac{5\pi}{4}$ (d) $\frac{3\pi}{4}, \frac{5\pi}{4}$ (e) $\frac{\pi}{2}, \frac{3\pi}{2}$ (f) $\frac{\pi}{2}, \frac{3\pi}{2}$ (g) $\frac{\pi}{2}, \frac{3\pi}{2}$ (h) $\frac{\pi}{2}, \frac{3\pi}{2}$ (i) $\frac{\pi}{2}, \frac{3\pi}{2}$ (j) $\frac{\pi}{2}, \frac{3\pi}{2}$ (k) $\frac{\pi}{2}, \frac{3\pi}{2}$ (l) $\frac{\pi}{2}, \frac{3\pi}{2}$ (m) $\frac{\pi}{2}, \frac{3\pi}{2}$ (n) $\frac{\pi}{2}, \frac{3\pi}{2}$ (o) $\frac{\pi}{2}, \frac{3\pi}{2}$ (p) $\frac{\pi}{2}, \frac{3\pi}{2}$ (q) $\frac{\pi}{2}, \frac{3\pi}{2}$ (r) $\frac{\pi}{2}, \frac{3\pi}{2}$ (s) $\frac{\pi}{2}, \frac{3\pi}{2}$ (t) $\frac{\pi}{2}, \frac{3\pi}{2}$ (u) $\frac{\pi}{2}, \frac{3\pi}{2}$ (v) $\frac{\pi}{2}, \frac{3\pi}{2}$ (w) $\frac{\pi}{2}, \frac{3\pi}{2}$ (x) $\frac{\pi}{2}, \frac{3\pi}{2}$ (y) $\frac{\pi}{2}, \frac{3\pi}{2}$ (z) $\frac{\pi}{2}, \frac{3\pi}{2}$ **Q014** (a) $\theta = 0, \frac{2\pi}{3}$ (b) $\theta = \pi, \frac{13\pi}{12}$ (c) $\frac{\pi}{2} < \theta < \frac{3\pi}{2}$ **Q022** B

Q004 $\tan \theta = 2$ or $\tan \theta = \sqrt{3}$; $\theta = \arctan(2) + n\pi$ or $\theta = \frac{\pi}{3} + n\pi$ **Q015** (a) $\frac{\pi}{2}, \frac{5\pi}{6}$ (b) $\frac{\pi}{2}, \frac{7\pi}{6}$ (c) $\pi, \frac{5\pi}{12}$ (d) $\frac{\pi}{2}, \frac{3\pi}{2}$ (e) $\frac{\pi}{2}, \frac{3\pi}{2}$ (f) $\frac{\pi}{2}, \frac{3\pi}{2}$ (g) $\frac{\pi}{2}, \frac{3\pi}{2}$ (h) $\frac{\pi}{2}, \frac{3\pi}{2}$ (i) $\frac{\pi}{2}, \frac{3\pi}{2}$ (j) $\frac{\pi}{2}, \frac{3\pi}{2}$ (k) $\frac{\pi}{2}, \frac{3\pi}{2}$ (l) $\frac{\pi}{2}, \frac{3\pi}{2}$ (m) $\frac{\pi}{2}, \frac{3\pi}{2}$ (n) $\frac{\pi}{2}, \frac{3\pi}{2}$ (o) $\frac{\pi}{2}, \frac{3\pi}{2}$ (p) $\frac{\pi}{2}, \frac{3\pi}{2}$ (q) $\frac{\pi}{2}, \frac{3\pi}{2}$ (r) $\frac{\pi}{2}, \frac{3\pi}{2}$ (s) $\frac{\pi}{2}, \frac{3\pi}{2}$ (t) $\frac{\pi}{2}, \frac{3\pi}{2}$ (u) $\frac{\pi}{2}, \frac{3\pi}{2}$ (v) $\frac{\pi}{2}, \frac{3\pi}{2}$ (w) $\frac{\pi}{2}, \frac{3\pi}{2}$ (x) $\frac{\pi}{2}, \frac{3\pi}{2}$ (y) $\frac{\pi}{2}, \frac{3\pi}{2}$ (z) $\frac{\pi}{2}, \frac{3\pi}{2}$ **Q023** A

Q005 (a) $0 \leq \theta < \frac{5\pi}{6}$ or $\frac{7\pi}{6} < \theta < 2\pi$ (b) $0 \leq \theta \leq \frac{\pi}{6}$ or $\frac{\pi}{2} < \theta \leq \frac{7\pi}{6}$ or $\frac{3\pi}{2} < \theta < 2\pi$ **Q016** $0 \leq \theta \leq \frac{2\pi}{3}$ or $\pi \leq \theta \leq \frac{5\pi}{3}$ **Q024** C

Q006 (a) $\frac{7\pi}{6} \leq \theta \leq \frac{11\pi}{6}$ (b) $0 \leq \theta < \frac{\pi}{3}$ or $\frac{5\pi}{3} < \theta < 2\pi$ (c) $\frac{\pi}{4} < \theta < \frac{\pi}{2}$ or $\frac{5\pi}{4} < \theta < \frac{3\pi}{2}$ **Q017** Dividing by $\cos(x)$ separates the **Q025** B

Q007 (a) $\frac{\pi}{2}, \frac{3\pi}{2}$ (b) $\frac{\pi}{2}, \frac{3\pi}{2}$ (c) $\frac{\pi}{2}, \frac{3\pi}{2}$ (d) $\frac{\pi}{2}, \frac{3\pi}{2}$ (e) $\frac{\pi}{2}, \frac{3\pi}{2}$ (f) $\frac{\pi}{2}, \frac{3\pi}{2}$ (g) $\frac{\pi}{2}, \frac{3\pi}{2}$ (h) $\frac{\pi}{2}, \frac{3\pi}{2}$ (i) $\frac{\pi}{2}, \frac{3\pi}{2}$ (j) $\frac{\pi}{2}, \frac{3\pi}{2}$ (k) $\frac{\pi}{2}, \frac{3\pi}{2}$ (l) $\frac{\pi}{2}, \frac{3\pi}{2}$ (m) $\frac{\pi}{2}, \frac{3\pi}{2}$ (n) $\frac{\pi}{2}, \frac{3\pi}{2}$ (o) $\frac{\pi}{2}, \frac{3\pi}{2}$ (p) $\frac{\pi}{2}, \frac{3\pi}{2}$ (q) $\frac{\pi}{2}, \frac{3\pi}{2}$ (r) $\frac{\pi}{2}, \frac{3\pi}{2}$ (s) $\frac{\pi}{2}, \frac{3\pi}{2}$ (t) $\frac{\pi}{2}, \frac{3\pi}{2}$ (u) $\frac{\pi}{2}, \frac{3\pi}{2}$ (v) $\frac{\pi}{2}, \frac{3\pi}{2}$ (w) $\frac{\pi}{2}, \frac{3\pi}{2}$ (x) $\frac{\pi}{2}, \frac{3\pi}{2}$ (y) $\frac{\pi}{2}, \frac{3\pi}{2}$ (z) $\frac{\pi}{2}, \frac{3\pi}{2}$ **Q018** B

Q008 $\frac{\pi}{3} < \theta < \pi$ or $\pi < \theta < \frac{5\pi}{3}$ **Q019** B

Q009 (a) $\theta = \frac{\pi}{6}, \frac{5\pi}{6}$ (b) $0 \leq \theta \leq \frac{2\pi}{3}$ or $\theta = \pi$ or $\frac{4\pi}{3} \leq \theta < 2\pi$ **Q020** B

Q010 (a) $\theta = 0, \frac{\pi}{6}, \frac{5\pi}{6}, \pi$ (b) $\theta = 0, \pi, \frac{7\pi}{6}, \frac{11\pi}{6}$ (c) $0 \leq \theta < \frac{2\pi}{3}$ or $\frac{4\pi}{3} < \theta < 2\pi$ **Q021** B

Q011 (a) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (b) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (c) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (d) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (e) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (f) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (g) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (h) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (i) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (j) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (k) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (l) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (m) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (n) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (o) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (p) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (q) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (r) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (s) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (t) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (u) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (v) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (w) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (x) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (y) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ (z) $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$ **Q022** B

Q012 $0 \leq \theta < \frac{\pi}{2}$ or $\frac{\pi}{2} < \theta \leq \pi$ **Q023** A

Q013 (a) $\theta = \frac{\pi}{2}, \frac{11\pi}{6}$ (b) $0 \leq \theta < \pi$ or $\frac{5\pi}{3} < \theta < \frac{7\pi}{3}$ **Q024** C

Q014 (a) $\theta = 0, \frac{2\pi}{3}$ (b) $\theta = \pi, \frac{13\pi}{12}$ (c) $\frac{\pi}{2} < \theta < \frac{3\pi}{2}$ **Q025** B

Q015 (a) $\frac{\pi}{2}, \frac{5\pi}{6}$ (b) $\frac{\pi}{2}, \frac{7\pi}{6}$ (c) $\pi, \frac{5\pi}{12}$ (d) $\frac{\pi}{2}, \frac{3\pi}{2}$ (e) $\frac{\pi}{2}, \frac{3\pi}{2}$ (f) $\frac{\pi}{2}, \frac{3\pi}{2}$ (g) $\frac{\pi}{2}, \frac{3\pi}{2}$ (h) $\frac{\pi}{2}, \frac{3\pi}{2}$ (i) $\frac{\pi}{2}, \frac{3\pi}{2}$ (j) $\frac{\pi}{2}, \frac{3\pi}{2}$ (k) $\frac{\pi}{2}, \frac{3\pi}{2}$ (l) $\frac{\pi}{2}, \frac{3\pi}{2}$ (m) $\frac{\pi}{2}, \frac{3\pi}{2}$ (n) $\frac{\pi}{2}, \frac{3\pi}{2}$ (o) $\frac{\pi}{2}, \frac{3\pi}{2}$ (p) $\frac{\pi}{2}, \frac{3\pi}{2}$ (q) $\frac{\pi}{2}, \frac{3\pi}{2}$ (r) $\frac{\pi}{2}, \frac{3\pi}{2}$ (s) $\frac{\pi}{2}, \frac{3\pi}{2}$ (t) $\frac{\pi}{2}, \frac{3\pi}{2}$ (u) $\frac{\pi}{2}, \frac{3\pi}{2}$ (v) $\frac{\pi}{2}, \frac{3\pi}{2}$ (w) $\frac{\pi}{2}, \frac{3\pi}{2}$ (x) $\frac{\pi}{2}, \frac{3\pi}{2}$ (y) $\frac{\pi}{2}, \frac{3\pi}{2}$ (z) $\frac{\pi}{2}, \frac{3\pi}{2}$ **Q026** B

Q016 $0 \leq \theta \leq \frac{2\pi}{3}$ or $\pi \leq \theta \leq \frac{5\pi}{3}$ **Q027** A

Q017 Dividing by $\cos(x)$ separates the **Q028** A

Q018 B

Q019 B

Q020 B

Q021 B

Q022 B

Q023 A

Q024 C

Q025 B

Q026 B

Q027 A

Q028 A

Q029 C

Q030 B

Q031 A

Q032 A

Q033 A

Q034 B

Quiz MCQ 1 B

Quiz MCQ 2 B

Quiz MCQ 3 B

Quiz MCQ 4 B

Quiz WRITTEN 5 $\theta = \frac{\pi}{6}, \frac{11\pi}{6}$

Quiz WRITTEN 6 $\frac{\pi}{2} < \theta \leq \frac{3\pi}{4}$ or $\frac{3\pi}{2} < \theta \leq \frac{7\pi}{4}$

Quiz WRITTEN 7 $\theta = \frac{\pi}{6}, \frac{5\pi}{6}, \frac{3\pi}{2}$

Quiz WRITTEN 8 $\frac{4\pi}{3} < \theta < 2\pi$