

Definition: Multi-index

For $\alpha_1, \dots, \alpha_n \geq 0$ we call $\alpha = (\alpha_1, \dots, \alpha_n)$ a multi-index. One then defines the following:

1. $\alpha! = \alpha_1! \cdots \alpha_n!$;
2. for $x \in \mathbb{R}^n$ one defines $x^\alpha = x_1^{\alpha_1} \cdots x_n^{\alpha_n}$ if $x_i \neq 0$ for all i , otherwise $x^\alpha = 0$ if there exists a j such that $x_j = 0$;
3. let $|\alpha| = \alpha_1 + \cdots + \alpha_n$ be the order of α ;
4. let $D^\alpha f(x) = \frac{\partial^{|\alpha|} f(x)}{\partial x_1^{\alpha_1} \cdots \partial x_n^{\alpha_n}}$.

Theorem: Taylor's Formula over \mathbb{R}^n

Let $E \subseteq \mathbb{R}^n$ be open and let $\overline{B_r(a)} \subseteq E$. Assume $f \in \mathcal{C}^k(E)$ then

$$f(x) = \sum_{|\alpha| \leq k} \frac{D^\alpha f(a)}{\alpha!} (x-a)^\alpha + \sum_{|\alpha|=k} h_\alpha(x) (x-a)^\alpha$$

for any $x \in B_r(a)$ where α is a multi-index and $h_\alpha : B_r(a) \rightarrow \mathbb{R}$ such that $h_\alpha(x) \rightarrow 0$ when $x \rightarrow a$ for all $|\alpha| = k$.

Proof:

Remark

Note

$$|(x-a)^\alpha| = \left| \prod_{i=1}^n (x_i - a_i)^{\alpha_i} \right| = \prod_{i=1}^n |x_i - a_i|^{\alpha_i} \leq |x-a|^{|\alpha|}.$$

Therefore we have that

$$\left| \sum_{|\alpha|=k} h_\alpha(x) (x-a)^\alpha \right| \leq |x-a|^k \sum_{|\alpha|=k} |h_\alpha(x)|$$

such that $\sum_{|\alpha|=k} h_\alpha(x) (x-a)^\alpha = o(|x-a|^k)$.

Let $x \in B_r(a)$ and define $[a, x] = \{\varphi(t) \mid t \in [0, 1]\}$ where $\varphi : [-1, 1] \rightarrow \mathbb{R}^n$ is defined by $\varphi(t) = a + t(x-a)$. Then let $g(t) = f(\varphi(t))$ such that $g(0) = f(a)$ and $g(1) = f(x)$. Then note $g \in \mathcal{C}^k([-1, 1])$ such that

$$g(1) = g(0) + \frac{1}{1!} g'(0) + \cdots + \frac{1}{(k-1)!} g^{(k-1)}(0) + \frac{1}{k!} g^{(k)}(c)$$

for some $c \in [0, 1]$. As shown previously we have

$$g^{(m)}(t) = \sum_{|\alpha|=m} D^\alpha f(\varphi(t)) \cdot (x-a)^\alpha$$

for $m = 1, \dots, k$ by mixed derivative theorem. Therefore

$$f(x) = \sum_{|\alpha| \leq k-1} \frac{D^\alpha f(a)}{\alpha!} (x-a)^\alpha + \sum_{|\alpha|=k} \frac{D^\alpha f(y)}{\alpha!} (x-a)^\alpha$$

for some $y \in [a, x]$. Now note that $D^\alpha f(y) = D^\alpha f(a) + (D^\alpha f(y) - D^\alpha f(a))$ where $D^\alpha f(y) - D^\alpha f(a) \rightarrow 0$ as $y \rightarrow a$. Then define $h_\alpha(x) = \frac{D^\alpha f(y) - D^\alpha f(a)}{\alpha!} \rightarrow 0$ as $x \rightarrow a$ where y is a function of x . \square